



**HAL**  
open science

## Root traits of crop species contributing to soil shear strength

Matthieu Forster, Carolina Ugarte, Mathieu Lamandé, Michel-Pierre Faucon

► **To cite this version:**

Matthieu Forster, Carolina Ugarte, Mathieu Lamandé, Michel-Pierre Faucon. Root traits of crop species contributing to soil shear strength. *Geoderma*, 2022, 409, pp.115642. 10.1016/j.geoderma.2021.115642 . hal-03470170

**HAL Id: hal-03470170**

**<https://hal.science/hal-03470170>**

Submitted on 8 Dec 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 **Root traits of crop species contributing to soil shear strength**

2 Matthieu Forster<sup>a</sup>, Carolina Ugarte<sup>a</sup>, Mathieu Lamandé<sup>bc†</sup>, Michel-Pierre Faucon<sup>\*a†</sup>

3

4 <sup>a</sup> AGHYLE (SFR Condorcet FR CNRS 3417), Chair in Agricultural Machinery & New Technologies,  
5 UniLaSalle, 19 Rue Pierre Waguet, 60026 Beauvais, France

6 <sup>b</sup> Department of Agroecology, Aarhus University, Research Centre Foulum, Blichers Allé 20, P.O.  
7 Box 50, DK-8830 Tjele, Denmark

8 <sup>c</sup> Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of  
9 Life Sciences, P.O. Box 5003, NO-1432 Ås, Norway

10 \*Corresponding author: michel-pierre.faucon@unilasalle.fr

11 †These authors contributed equally to this work.

12

## 13 **Abstract**

14 Roots of annual species would be able to conserve soil properties during traffic induced compaction.  
15 The objective of this study was to determine and compare root traits and soil shear strength for three  
16 crop species with contrasted root types and morphological traits to see if roots of annual species are  
17 able to increase soil shear strength and thus explain the soil properties conservation. The experiment  
18 was performed under controlled conditions in steel cylinders containing a repacked loamy sand, with  
19 initial dry bulk density of 1.2 g cm<sup>-3</sup> and that was kept moist. Soil shear strength parameters, i.e.  
20 cohesion and angle of internal friction, were determined from direct shear tests for soil cores at a soil  
21 matric potential of -10 kPa (i.e. water content at field capacity). The direct shear tests were performed  
22 with six external normal stress levels (25, 34, 44, 63, 83 and 93 kPa) and at a constant shear rate of 3  
23 mm.min<sup>-1</sup>, and applying Mohr Coulomb equation. The difference of root type, root length density, root  
24 density, specific root length and root dry matter content among crop species was not related to a  
25 difference in soil shear strength. The root volume density was the main trait involved in both soil  
26 cohesion and the angle of internal friction. This study highlights the effect of roots of annual crop  
27 species on soil shear strength by comparing their root traits to the apparent cohesion of soil core where  
28 they grow. *Vicia faba* would be a good candidate to improve soil shear strength for soil conservation.  
29 This study constitutes a step towards improving the understanding of plants' effects on soil shear  
30 strength with regards to selecting species and designing cropping systems for soil conservation.

31 **Keywords:** Soil cohesion, angle of internal friction, root traits, soil shear strength, crop species,  
32 compaction

33

## 34 **Abbreviations**

35 ARD: Average Root Diameter (mm)

36 ATRD: Average Taproot Diameter (mm)

37 DMC: Dry Matter Content ( $\text{mg g}^{-1}$ , root dry weight divided by the root fresh weight)

38 RD: Root Density ( $\text{g.m}^{-3}$ )

39 RLD: Root Length Density ( $\text{cm.cm}^{-3}$ )

40 RVD: Root Volume Density ( $\text{cm}^3.\text{cm}^{-3}$ )

41 SRL: Specific Root Length ( $\text{m.g}^{-1}$ )

42 TRVD: Taproot Volume Density ( $\text{cm}^3.\text{cm}^{-3}$ )

43

## 44 **Introduction**

45 Soil compaction is known as a major threat to soil quality in Europe (Berge et al., 2017; Stolte et al.,  
46 2015). Soil compaction in agroecosystems results from three-dimensional stress propagation in the soil  
47 (Horn et al., 2003), which is mainly caused by traffic during agricultural operations in the field. The  
48 soil compaction observed in agrosystems results from the combination of two processes: compression  
49 and shearing (Horn et al., 2003). Both processes negatively impact the soil pore system. Compression  
50 mainly reduces pore volume, while shearing affects pore system continuity (Berisso et al., 2013;  
51 Huang et al., 2022). These modifications to the pore system reduce the capacity of air exchange  
52 between the soil and the atmosphere, and impair water transport through the soil profile. Compression  
53 and shearing present heavy impacts on soil functions given as they make root penetration more  
54 difficult, decrease soil permeability, reduce the availability of nutrients and, therefore, reduce yields  
55 (Soane et al., 1980; Schjønning et al., 2016). These soil functions are essential for plant growth (Unger  
56 and Kaspar, 1994). Technological and agronomical solutions that can mitigate the detrimental effects  
57 of compaction on soil functions are needed. This mitigation can be achieved by reducing the  
58 mechanical stress applied during traffic or by improving soil mechanical resistance to deformation  
59 (Batey, 2009). Reinforcement of soil shear strength by roots has been investigated in the prevention of  
60 landslides and improvement of riverbank stability (Pollen and Simon, 2005; Stokes et al., 2009).

61 Potentially, plant roots can help enhance the soil's mechanical strength, and this protective function of  
62 plants could be included in the design of cropping systems (Bodner et al., 2021).

63 Soil shear strength parameters, cohesion and angle of internal friction, are classically determined from  
64 the Coulomb (1776) equation (Eq. 1):

$$\tau = c + \sigma \tan \varphi$$

65 where  $\tau$  is the shear strength (kPa),  $\sigma$  is the normal stress acting on the failure surface (kPa),  $C$  is the  
66 cohesion (kPa) and  $\varphi$  is the angle of internal friction ( $^{\circ}$ ).

67 Plant roots could potentially influence both of these two shear strength parameters. Roots increase soil  
68 shear strength directly through their mechanical resistance, and their anchoring in the soil, and  
69 indirectly through water removal from the soil by transpiration (Pollen and Simon, 2005). A decrease  
70 in the soil water potential increases both the soil resistance to compression (Schjønning and Lamandé,  
71 2018) and to shearing (Fredlund Delwyn, 2006; Schjønning et al., 2020). Several studies examining the  
72 effects of roots on soil shear strength have highlighted the direct effect of the root tensile strength (the  
73 root's capacity to resist tension) and in most cases, effects of a range of root properties on cohesion:  
74 root area ratio (the surface proportion of a plane occupied by roots) (Operstein and Frydman, 2000);  
75 root density (the dry weight of roots per volume of soil) (Tengbeh, 1989; Ali and Osman, 2008); and  
76 root length density (the length of roots per volume of soil) (Osman and Barakbah, 2006).  
77 Physiological root traits (root exudates) and plant–soil microorganisms interactions involved in soil  
78 aggregation can contribute also to the increase of soil cohesion and soil shear strength (Rillig et al.,  
79 2015; Poirier et al. 2018; Garcia et al., 2019). Effects of the root architecture or root type (i.e. taproot  
80 or fibrous) on cohesion have also been highlighted (Ghestem et al., 2014; Li et al., 2017).

81 Only scarce evidence of root effects on soil internal friction exist (Graf et al., 2009). However, more  
82 research is needed to identify the specific root traits involved in the increase of soil friction (Yildiz et  
83 al., 2018). In our context, soil friction increase by root traits could be due to the impact of high normal  
84 forces applied during compaction by traffic which will amplify the role of friction. In our context,

85 shear is also not restricted to a single shear plane as it could be in landslide. Traffic indeed induce  
86 shear stress gradually in the whole soil profile. Root density and distribution in the soil could be key  
87 factors as they could be related to a homogeneous soil reinforcement.

88 Fundamental differences in soil properties, root traits, and the rate and magnitude of deformation exist  
89 between the contexts of landslide prevention or riverbank stability on the one hand and traffic-induced  
90 compaction of agricultural soils on the other hand. In agroecosystems, there is a high frequency and  
91 abundance of annual species, presenting a specific gradient of root traits. In addition, transient  
92 mechanical stresses are applied at very high rates during traffic to agricultural soils (Keller and  
93 Lamandé 2010). Vertical stresses applied by agricultural machinery can be much larger than  
94 overburden stress (e.g. Lamandé and Schjønning, 2011b). Shearing during traffic occurs in the shallow  
95 soil layer (<0.3 m depth) (Berisso et al., 2013), where the soil is not necessarily saturated with water,  
96 but where roots of annual species are the most densely present (Osman and Barakbah, 2006).

97 Direct shear tests with increasing normal stress are needed to determine the cohesion and angle of  
98 internal friction using the classical Coulomb approach (Eq. 1). It is challenging to specifically identify  
99 the effects of crop roots on soil cohesion and on the angle of internal friction. This requires the use of  
100 a direct shear apparatus adapted to reproduce the compression and shearing of the magnitude of what  
101 is observed beneath a running tyre (Giadrossich et al., 2017), and allowing measurements on soil cores  
102 large enough to include roots systems of annual crops. In addition, effects of crop roots on soil shear  
103 strength could be quite small as compared to the soil shear strength (Amiri Khaboushan et al., 2018).  
104 This can be overcome by performing tests using a repacked soil, presenting therefore a low cohesion  
105 in the range of 4 to 30 kPa for degrees of saturation from 20 to 70% (Hemmat et al., 2010). The aim of  
106 this study was to investigate the effects of root traits on soil shear strength parameters for three crop  
107 species with contrasting root types and morphological traits. Root morphological traits, cohesion and  
108 angle of internal friction were compared between crop species and soil without plants to identify the  
109 root traits involved in soil reinforcement.

## 111 **Materials and methods**

### 112 **Experimental design and plant materials**

113 A greenhouse experiment was conducted over two and a half months, from October to December  
114 2019. The greenhouse light was turned on between 8:00 AM and 10:00 PM, and the temperature was  
115 regulated at 24°C during the simulated daylight and 18°C during the simulated night. The experiment  
116 included four treatments: soil without plants (reference treatment); soil with *Brassica napus* L. which  
117 is the second oil crop in the world and an important source of edible oil, animal fodder, vegetables,  
118 condiments and biodiesel, and presents a taproot; soil with *Secale cereale* L. which is a grass cereal  
119 grown extensively as a grain, a cover crop and a forage crop, and presents a fibrous root; and soil with  
120 *Vicia faba* L. which is a major food and feed legume because of the high nutritional value of its seeds  
121 (rich in protein and starch) and presents a taproot system with larger fine roots than *Brassica napus*  
122 (Forster et al., 2020). These three species could be used as cash or cover crops. The soil consisted of a  
123 2:1 ratio mix of Luvisol (IUSS Working Group WRB, 2015: 20.5 g 100g<sup>-1</sup> of clay, 69.4 g 100g<sup>-1</sup> of silt  
124 and 10.1 g 100g<sup>-1</sup> of sand) and of river sand previously sieved at 5 mm and dried (at 105°C during 24  
125 h) and manually mixed.

126 Each treatment included 12 replicates divided into three blocks, with each block containing 16  
127 randomly placed pots. Pots were made of two stacked steel cylinders 0.1 m in height and 0.16 m in  
128 inner diameter with the wall thickness of 4 mm (Fig. 1.b). Pots were packed with soil to reach a dry  
129 bulk density of 1.2 g cm<sup>-3</sup>. To do so, pots were filled and compact three consecutive times. Soil was  
130 added and compact at 7 cm, 14 cm and 20 cm. Each time, the right quantity was added to reach the  
131 aimed bulk density and compact by hand using a handmade wooden disk. Pots size were selected to be  
132 large enough to allow roots of young crop to develop normally without too much space restriction and  
133 small enough to reduce the amount of dead weight needed to reach a maximum normal stress 100 kPa  
134 during the shear tests (See next paragraph for details). For the crop species treatment, three seeds were  
135 planted closed to the centre per pot. After two weeks, the healthiest plant in each pot was selected and  
136 others were removed. To avoid possible heterogeneous light exposure in the greenhouse, the  
137 placement of the pots was changed on the second, fourth and sixth weeks. Each time, pots were placed  
138 close to the light, each replacing its neighbouring pot, while the pot closest to the light was moved to

139 the farthest space. Each pot was irrigated once a week during the first month and then twice a week  
140 until the end of the experiment. After two and a half months of growth, root of these crop species  
141 occupied the bottom of the pot (i.e. 20 cm of deep) below the shear plane located at 10 cm.

142

### 143 **Soil shear strength**

144 A direct shear apparatus was built especially for this study to perform shear tests directly on the pots,  
145 thus avoiding unnecessary manipulation of the repacked soil cores (Fig. 1). The size of the soil cores  
146 to be used for the shear tests followed the ASTM recommendation of ratio between the size of the  
147 sample and the maximum particle size of the material to be tested (ASTM Standard D 420–D 4914,  
148 Soil and Rock, Annual Book of ASTM Standards Vol. 04.08, ASTM International, West  
149 Conshohocken, PA). Two pots per treatment were subjected to a direct shear test at six external  
150 normal stress levels,  $\sigma_n$  (25, 34, 44, 63, 83 and 93 kPa), resulting in twelve shear tests per treatment,  
151 each test being performed on a new pot (i.e. no consecutive shear test on the same pot). The external  
152 normal stress was applied using a dead weight system. The direct shear tests were performed two and  
153 a half months after sowing. Two days before the tests, each pots were saturated with tap water. Twice  
154 a day, the soil matric potential was measured at 0.08 m depth for each treatment using tensiometers  
155 (SMS 2030S1 type, SDEC, Reignac-sur-Indre, France) placed in three pots chosen randomly for each  
156 treatment. Shear tests were performed when the soil matric potential reached approximately -10 kPa,  
157 which correspond to a water content at field capacity, i.e. where most of macro- and mesopores (i.e.  
158 eq. pore diameter  $> 30 \mu\text{m}$ , Luxmoore, 1981) are air-filled (Table 1). The upper part of the soil core  
159 was moved horizontally at a constant rate of  $3 \text{ mm}\cdot\text{min}^{-1}$ , and the test was stopped when a total  
160 horizontal displacement of 25 mm was reached. Horizontal displacement rate was controlled using an  
161 actuator monitored by a software called “Kollmorgen WorkBench”. During the test, the force required  
162 to displace the upper part of the soil core was recorded by a H3-C3 500 kg load cell every 0.095  
163 seconds. The shear stress was calculated from the load cell readings (kg) and the contact area between  
164 the upper and the lower part of the soil core. This contact area was considered as a lens formed by the  
165 intersection of section area of the two parts of the soil core. The horizontal displacement (mm) was

166 derived from the duration of the test (s) and the rate of horizontal displacement ( $\text{mm}\cdot\text{s}^{-1}$ ). We  
167 quantified the steel-steel friction forces between the upper- and lower cylinders when empty. The  
168 mean friction measured for fourteen tests was 0.66 kPa (standard deviation of 0.15 kPa), which was  
169 low as compared to the level of maximum shear stress recorded for the tests with cylinders repacked  
170 with soil.

171 The maximum shear stress,  $\tau_{\max}$  (kPa), was determined graphically on each shear stress-displacement  
172 curve according to the shape of the curve, following the methodology of Ghestem et al., (2014). When  
173 no clear yielding point was identifiable, the maximum shear stress was defined as the inflexion point  
174 of the curve. For unsaturated soils, the negative pore water pressure will contribute to the cohesive  
175 forces. During the test, soil will deform until shear failure, and the pore water pressure will change  
176 accordingly. We chose a high shearing rate to get a good representation of the shearing during traffic  
177 in the field (Schjønning et al., 2020). The high shearing rate prevented measuring the pore water  
178 pressure at failure, and we are thus not able to distinguish between the effective cohesion, from  
179 physicochemical interaction between material particles, and the cohesive forces from the negative pore  
180 water pressure at failure. Therefore, the cohesive forces detected in our study will be labelled the  
181 “apparent cohesion”,  $c_{\text{app}}$  (kPa), as also suggested by Schjønning et al., (2020). Similarly, we will not  
182 be able to identify the effects of the negative pore water pressure at failure on the friction forces. The  
183 friction forces determined in our study will then be labelled  $\phi_{\text{app}}$  ( $^{\circ}$ ). Consequently,  $c_{\text{app}}$  and  $\phi_{\text{app}}$  were  
184 derived using the Coulomb equation (Eq.1) assuming a linear relationship between the maximum  
185 shear stress measured and the normal load applied during the direct shear test (Eq. 2):

$$\tau_{max} = c_{app} + \sigma_n \tan\phi_{app}$$

## 186 **Root trait characterisation**

187 After the shear tests, all pots were emptied, and roots were manually separated from the soil. Soil was  
188 then placed two consecutive times in a 5 mm sieves to extract the remaining roots. Six root traits were  
189 then measured: the average root diameter (ARD, mm), which represents the average diameter of the  
190 fibrous system; the root length density (RLD,  $\text{cm}\cdot\text{cm}^{-3}$ ), which is the root length divided by the soil  
191 volume; the root density (RD,  $\text{g}\cdot\text{m}^{-3}$ ), which is the roots' dry weight divided by the soil volume; the



192 root volume density (RVD,  $\text{cm}^3.\text{cm}^{-3}$ ), which is the roots' volume divided by the soil volume; the  
193 specific root length (SRL,  $\text{m.g}^{-1}$ ), which is the root length divided by the root dry weight and the dry  
194 matter content (DMC) which is the root dry weight divided by the root fresh weight. Trait  
195 measurements were performed following the method described by Ristova and Barbez (2018).

196 All roots found in each pot were weighed and scanned in a film of water using the Epson Perfection  
197 V850 Pro at a resolution of 600 dpi. The software WinRhizo 2016 (Regent Instrument Inc.,  
198 Instruments, Québec, Canada) was used to quantify the root length and root volume. The taproot  
199 volume (TRV) and the average taproot diameter (ATRD) were determined separately using the ImageJ  
200 freeware (Schneider et al. 2012). TRV was calculated using the sum of homogeneous volume sections  
201 in terms of diameter. The diameter and the length of each sections were used to calculate ATRD with  
202 the length of each section used to proportionally calculate the mean diameter. Each volume section  
203 was determined by its length and mean diameter. TRV was then divided by the soil volume to obtain  
204 the taproot volume density (TRVD) which could be compared to the RVD. TRV and TRVD were not  
205 used as a trait but were used to compare the root type of the species. Fresh roots were weighed and  
206 then dried at  $105^\circ\text{C}$  for 24 h and then weighed again to obtain the DMC and SRL. The dry weight,  
207 length and volume were then divided by the total volume of the pot to obtain the RD, RLD and RVD.

208

## 209 **Statistical analysis**

210 Root traits values did not follow a normal distribution. A nonparametric Kruskal–Wallis rank sum test  
211 followed by a pairwise Wilcoxon test were thus performed to identify differences between species in  
212 each root trait. Spearman's correlation was then used to identify correlation between root traits. Shear  
213 strength values followed a Gamma distribution and were not subjected to any transformation. A  
214 variance analysis was performed using the Coulomb linear regression model (Eq. 2) to test the  
215 differences in intercept and slope of the shear strength vs. normal stress relationship between the four  
216 treatments.

217

## 218 **Results**

### 219 **Root traits and root type comparison among species**

220 All root traits were found to be significantly different between species: ARD (Chi square = 23.36, p-  
221 value = <0.001, df = 2), DMC (Chi square = 29.12, p-value = <0.001, df = 2), RD (Chi square = 26.08,  
222 p-value = <0.001, df = 2), RLD (Chi square = 8.06, p-value = 0.018, df = 2), RVD (Chi square =  
223 19.92, p-value = <0.001, df = 2) (Table 2 and Fig. 2). For *Vicia faba*, significant differences from the  
224 other two species were observed in five of the six measured traits. *Vicia faba* had the highest ARD,  
225 RD and RVD, and had the lowest SRL and DMC. *Secale cereale* had the highest SRL and the lowest  
226 RD. *Brassica napus* had the highest DMC. RLD did not differ between *Brassica napus* and *Vicia faba*  
227 (Table 2 and Fig. 2).

228 The correlation matrix showed that RLD was the only trait not correlated with another. RD, RVD and  
229 ARD were positively correlated, and SRL and DMC were positively correlated. RD, RVD and ARD  
230 were negatively correlated with SRL and DMC (Table 3).

231 No differences in TRVD nor in ATRD were found between *Vicia faba* and *Brassica napus*. *Vicia faba*  
232 did not consistently grow a taproot. A taproot was observed in six pots out of twelve, generating a  
233 higher standard error for the TRVD and ATRD of *Vicia faba*. *Vicia faba* combined the taproot and  
234 fibrous systems.

235

### 236 **Species effects on soil shear strength**

237 Apparent cohesion and angle of internal friction for each species and for the control without plants are  
238 reported in Table 4. *Vicia faba* reached an angle of internal friction of 17.9° and a cohesion of 20.2  
239 kPa, while *Brassica napus* reached an angle of internal friction of 15.6° and a cohesion of 4.9 kPa and  
240 *Secale cereale* reached an angle of internal friction of 12.6° and a cohesion of 8.8 kPa. For the  
241 reference treatment, we measured an angle of internal friction of 10.5° and a cohesion of 9.6 kPa  
242 (Figure 3). The variance analysis of the regressions showed that only *Vicia faba* had a higher angle of

243 internal friction and cohesion than the reference treatment, while both *Secale cereale* and *Brassica*  
244 *napus* did not have significantly different angle of internal friction or cohesion than the reference  
245 treatment.

246

## 247 **Discussion**

248 The main hypothesis of the present study was that roots of annual crop species would increase soil  
249 cohesion. The effect of roots on the soil shear strength was determined for three annual crop species  
250 with contrasting traits: *Brassica napus*, *Secale cereale* and *Vicia faba* (Table 4 and Fig. 3). However,  
251 only the presence of *Vicia faba* lead to an increase in apparent cohesion. Surprisingly, *Vicia faba* also  
252 present a significant higher angle of internal friction, as indicated by the increase compared to the  
253 reference treatment (Fig. 3). Our results suggest that some annual crop species could have a positive  
254 effect on soil shear strength.

255 No increase in soil shear strength was observed for *Brassica napus* or *Secale cereale* (Table 4, Fig. 4),  
256 even though the two species showed contrasting root types: a taproot in *Brassica napus* and a fibrous  
257 root system for in *Secale cereale* (Table 2, Fig. 2). This indicate that a rooted soil did not present  
258 systemically a higher shear strength, and that the potential soil reinforcement did not rely on the root  
259 type. These results suggest that the effects of roots on shear strength would be controlled by root traits.  
260 In that case, characterisation of root traits would be necessary to improve our understanding of root–  
261 soil interactions with regards to soil conservation in agroecosystems.

262 Previous studies examining the relationships between root traits and mechanical soil properties have  
263 shown that the RLD (Osman and Barakbah, 2006, Stokes et al., 2009), the RD (Ali and Osman, 2008),  
264 the root area ratio, the root tensile strength, which is negatively correlated to root diameter (De Baets  
265 et al., 2008, Ali and Osman, 2008, Operstein and Frydman, 2000), and the root type (Ghestem et al.,  
266 2014, Li et al., 2017) were influencing soil shear strength, with regard to the soil's protection against  
267 landslides. Gravity is the most important force component acting on soil that cause shear stresses  
268 susceptible to trigger landslides. Slow shearing of sloppy soil masses at constant volume may be

269 accompanied by a decrease in pore water pressure, i.e. a decrease in apparent cohesion that can lead to  
270 failure (landslide). In that context, the plants with high root length density and root area ratio are  
271 efficient for landslide prevention. The direct shear test method used in the present study reproduce  
272 shearing processes similar to the ones acting for shearing of agricultural soils during traffic: high shear  
273 rate and large normal load. In comparing root traits and soil shear strength with and without three plant  
274 species, the present study has allowed the identification of the main traits involved in the increase in  
275 soil shear strength by annual crop roots. Out of the five traits that distinguish *Vicia faba* from the two  
276 other species, three root traits (RD, DMC and SRL) differed between *Brassica napus* and *Secale  
277 cereale* (Fig. 2) while the cohesion and the angle of internal friction remained similar. It indicates that  
278 these three root traits do not directly influence the soil shear strength under these controlled conditions  
279 (plants in pots, remoulded soil). In moist sands, the shear strength is due to frictional resistance  
280 between grains, which is a combination of rolling and sliding friction, and to interlocking (the stress  
281 required to supply the energy for sands being expended in volume), and to apparent cohesion due to  
282 the pore water pressure. In the present study, in addition to that, only a limited true cohesion is  
283 expected from the silt and clay particles as the soil was remoulded. In contrary to the observations  
284 from studies about protection against landslides (Osman and Barakbah, 2006; Stokes et al., 2009), our  
285 results showed that RLD of the three annual crop species did not influence shear strength as it did not  
286 discriminate *Vicia faba* from the other two species. The low RLD of the three crop species  
287 investigated here ( $2.1-2,8 \text{ cm.cm}^{-3}$ ) compared to the RLD measured for perennial herbaceous, shrub  
288 and tree species (more  $300 \text{ cm.cm}^{-3}$ ), might explain the absence of relationships between RLD and the  
289 soil shear strength observed in the present study. Interestingly, annual crop species with low RLD can  
290 positively influence soil shear strength. Our results showed that soil shear strength was largest when  
291 the ARD and RVD were high. RVD in *Vicia faba* reflects the number of roots and was expected to  
292 participate in the increase of the soil shear strength. The divergence of some root traits effects on soil  
293 shear strength in the literature could be explained by the various methods (Freschet et al., 2021):  
294 different soil conditions (e.g. saturated or unsaturated), different soil core sizes, different order of  
295 magnitude of the root traits between annual crops and perennial plants (Roumet et al., 2006). Perennial  
296 plants (shrubs, trees) are usually investigated in studies related to prevention of landslides (Freschet et

297 al., 2021). These differences may prevent the direct transfer of the findings from these studies to  
298 agroecosystems. However, other root traits not measured in this study could also be related to  
299 differences in soil shear strength, such as the insertion angle of the roots or the roots' vertical and  
300 horizontal distribution in the soil (Stokes et al., 2009). Future studies examining the effects of root  
301 traits on soil shear strength should include a larger range of root traits in crop species.

302 In our study, *Vicia faba* was found to be related to a larger cohesion and angle of internal friction.  
303 RVD appeared to be the root trait involved in increase of both soil cohesion and the angle of internal  
304 friction (figures 2 and 3). Here, higher RVD value for *Vicia faba* was mainly due to its fibrous system  
305 and not due to a large taproot as observed in *B. napus*. It was thus related to the root capacity to spread  
306 and colonize the soil. An explanation could be that the replacement of soil air by roots (i.e. roots  
307 filling up the macropores) helped to reinforce the soil, without influence of the roots capacity to resist  
308 tension. Species with large taproot present a low RLD, RVD and root tensile strength, and would not  
309 be thus good candidates to increase soil shear strength.

310 The positive roots effects on the angle of internal friction is in accordance with the study by Graf et al.,  
311 (2009). They applied a maximum normal stress of 100 kPa during the shear tests, which was larger  
312 than the maximum normal stress used in other studies showing no evidence of roots' effect on the soil  
313 angle of internal friction. A low maximum normal stress applied during shear tests might prevent  
314 identifying this effect (Giadrossich et al., 2017). Our results indicated an increase of  $0.14 \text{ kPa kPa}^{-1}$  in  
315 the slope of Eq. 2 for *Vicia faba* compared to the reference treatment without plants (figure 3). In the  
316 contexts of landslide protection and riverbank stability, normal stress is applied by overburden soil and  
317 is therefore lower (Yildiz et al, 2018). A small increase in soil shear strength by friction could thus  
318 easily be outcompeted by a significant increase in cohesion due to the large perennial plants roots. It  
319 could explain why roots effects on the angle of internal friction were not identified in this study. From  
320 our results, the angle of internal friction increase by annual plant roots could participate to soil  
321 shearing prevention during traffic in agricultural fields. Our results provide a step toward a better  
322 understanding of the effects of crop roots on soil shear strength in controlled conditions through the

323 increase of both soil cohesion and angle of internal friction. There is now an urgent need to evaluate  
324 the potential of annual plant roots to reinforce the shear strength of structured agricultural soils.

325

## 326 **Conclusion**

327 The present trait-based study allowed the identification of the effects of annual crop roots on soil shear  
328 strength and provided information about which root traits might be relevant for the mitigation of  
329 traffic-related soil compaction. Our results showed crop species with low root length density could  
330 improve the soil shear strength, and that the root volume density may be the main root trait correlated  
331 to a larger apparent cohesion and angle of internal friction. These results provide an evidence of the  
332 possible effect of roots on soil shear strength in agroecosystems and demonstrate the potential of  
333 designing cropping systems with improved soil structure protection. To do so, further experiment  
334 should study roots effects on soil shear strength by using larger root traits gradient and for various soil  
335 conditions, including using undisturbed soil sample from agricultural fields at a range of soil water  
336 potentials.

337

## 338 **Acknowledgements**

339 We gratefully thank the technical support of Céline Roisin, Aurore Coutelier and Vincent Hervé. We  
340 also thanks all the Agrilab team for the huge support provided to build the shear box. This work has  
341 been supported by the Chair in Agricultural Machinery and New Technologies, backed by Institut  
342 Polytechnique UniLaSalle with financial support from the Michelin Corporate Foundation, AGCO  
343 Massey-Ferguson, Kuhn, the Hauts-de-France Regional Council and the European Regional  
344 Development Fund (ERDF).

345

## 346 **References**

347

348 Ali, F.H., Osman, N., 2008. Shear strength of a soil containing vegetation roots. *Soils Found*  
349 48, 587–596. <https://doi.org/10.3208/sandf.48.587>  
350

351 American Society for Testing and Materials (ASTM), 1996. Annual Book of ASTM  
352 Standards, vol. 04-08. Soil and Rock (I), D420-D4914  
353

354 Amiri Khaboushan, E., Emami, H., Mosaddeghi, M.R., Astaraei, A.R., 2018. Estimation of  
355 unsaturated shear strength parameters using easily-available soil properties. *Soil*  
356 *Tillage Res* 184, 118–127. <https://doi.org/10.1016/j.still.2018.07.006>  
357

358 Batey, T. 2009. Soil compaction and soil management – a review. *Soil Use Manag* 25, 335–  
359 345. <https://doi.org/10.1111/j.1475-2743.2009.00236.x>  
360

361 Berge, H.F.M., Schroder, J.J., Olesen, J.E., Giraldez Cervera, J.V., 2017. Research  
362 for AGRI Committee – Preserving agricultural soils in the EU, European Parliament,  
363 Policy, Department for Structural and Cohesion Policies, Brussels  
364

365 Berisso, F.E., Schjønning, P., Lamandé, M., Weisskopf, P., Stettler, M., Keller, T., 2013.  
366 Effects of the stress field induced by a running tyre on the soil pore system. *Soil*  
367 *Tillage Res* 131, 36–46. <https://doi.org/10.1016/j.still.2013.03.005>  
368

369 Bodner, G., Mentler, A., Keiblinger, K., 2021. Plant Roots for Sustainable Soil Structure  
370 Management in Cropping Systems. *The Root Systems in Sustainable Agricultural*  
371 *Intensification*, 45-90.  
372

373 Coulomb, C.A., 1776. Essai sur une application des règles des maximis et minimis à quelques  
374 problèmes de statique relatifs à l'architecture. *Mém. acad. roy. pres. divers savants*,  
375 Vol. 7, Paris.

376 De Baets, S., Poesen, J., Reubens, B., Wemans, K., De Baerdemaeker, J., Muys, B., 2008.  
377 Root tensile strength and root distribution of typical Mediterranean plant species and  
378 their contribution to soil shear strength. *Plant Soil* 305, 207-226.  
379

380 Forster, M., Ugarte, C., Lamandé, M., & Faucon, M. P., 2020. Relationships between Root  
381 Traits and Soil Physical Properties after Field Traffic from the Perspective of Soil  
382 Compaction Mitigation. *Agronomy* 10, 1697.  
383

384 Fredlund Delwyn G., 2006. Unsaturated Soil Mechanics in Engineering Practice. *J. Geotech.*  
385 *Geoenvironmental Eng* 132, 286–321. [https://doi.org/10.1061/\(ASCE\)1090-](https://doi.org/10.1061/(ASCE)1090-0241(2006)132:3(286))  
386 [0241\(2006\)132:3\(286\)](https://doi.org/10.1061/(ASCE)1090-0241(2006)132:3(286))  
387

388 Freschet, G.T., Roumet, C., Comas, L.H., Weemstra, M., Bengough, A.G., Rewald, B.,  
389 Bardgett, R.D., De Deyn, G.B., Johnson, D., Klimešová, J., Lukac, M., McCormack,  
390 M.L., Meier, I.C., Pagès, L., Poorter, H., Prieto, I., Wurzbürger, N., Zadworny, M.,  
391 Bagniewska- Zadworna, A., Blancaflor, E.B., Brunner, I., Gessler, A., Hobbie, S.E.,  
392 Iversen, C.M., Mommer, L., Picon- Cochard, C., Postma, J.A., Rose, L., Ryser, P.,  
393 Scherer- Lorenzen, M., Soudzilovskaia, N.A., Sun, T., Valverde- Barrantes, O.J.,

394 Weigelt, A., York, L.M., Stokes, A., 2021. Root traits as drivers of plant and  
395 ecosystem functioning: current understanding, pitfalls and future research needs. *New*  
396 *Phytologist*  
397

398 Ghestem, M., Veylon, G., Bernard, A., Vanel, Q., Stokes, A., 2014. Influence of plant root  
399 system morphology and architectural traits on soil shear resistance. *Plant Soil* 377, 43–  
400 61. <https://doi.org/10.1007/s11104-012-1572-1>  
401

402 Giadrossich, F., Schwarz, M., Cohen, D., Cislighi, A., Vergani, C., Hubble, T., Phillips, C.,  
403 Stokes, A., 2017. Methods to measure the mechanical behaviour of tree roots: A  
404 review. *Ecol. Eng., Soil Bio- and Eco-Engineering: The Use of Vegetation to Improve*  
405 *Slope Stability, Proceedings of the Fourth International Conference* 109, 256–271.  
406 <https://doi.org/10.1016/j.ecoleng.2017.08.032>  
407

408 Graf, F., Frei, M., Böll, A., 2009. Effects of vegetation on the angle of internal friction of a  
409 moraine. *Snow Lands Res* 17  
410

411 Hemmat, A., Aghilinategh, N., Sadeghi, M., 2010. Shear strength of repacked remoulded  
412 samples of a calcareous soil as affected by long-term incorporation of three organic  
413 manures in central Iran. *Biosystems Engineering* 107, 251-261.  
414

415 Horn, R., Way, T., Rostek, J., 2003. Effect of repeated tractor wheeling on stress/strain  
416 properties and consequences on physical properties in structured arable soils. *Soil*  
417 *Tillage Res* 73, 101–106. [https://doi.org/10.1016/S0167-1987\(03\)00103-X](https://doi.org/10.1016/S0167-1987(03)00103-X)  
418

419 Huang, X., Horn, R., Ren, T., 2022. Soil structure effects on deformation, pore water  
420 pressure, and consequences for air permeability during compaction and subsequent  
421 shearing. *Geoderma* 406, 115452.  
422

423 IUSS Working Group WRB, 2015. World Reference Base for Soil Resources 2014, update  
424 2015 International soil classification system for naming soils and creating legends for  
425 soil maps. *World Soil Resources Reports No. 106*. FAO, Rome. E-ISBN 978-92-5-  
426 108370-3  
427

428 Lamandé, M., Schjønning, P., 2011. Transmission of vertical stress in a real soil profile. Part  
429 I: Site description, evaluation of the Söhne model, and the effect of topsoil tillage. *Soil*  
430 *Tillage Res.* 114, 57–70. <https://doi.org/10.1016/j.still.2011.05.004>  
431

432 Luxmoore, R.J., 1981. Micro-, meso-, and macroporosity of soil. *Soil Sci. Soc. Am. J.* 45,  
433 671.

434 Li, Y., Wang, Yunqi, Wang, Yujie, Song, S., 2017. Effects of Root Architecture  
435 Characteristics on Soil Reinforcement in Undisturbed Soil. *Curr. Sci* 113, 1993.  
436 <https://doi.org/10.18520/cs/v113/i10/1993-2003>  
437



- 438 Operstein, V., Frydman, S., 2000. The influence of vegetation on soil strength. Proceedings of  
439 the Institution of Civil Engineers - Ground Improvement 2000 4:2, 81-89 .  
440 <https://doi.org/10.1680/grim.2000.4.2.81>  
441
- 442 Osman, N., Barakbah, S.S., 2006. Parameters to predict slope stability—Soil water and root  
443 profiles. *Ecol. Eng* 28:90–95. <https://doi.org/10.1016/j.ecoleng.2006.04.004>  
444
- 445 Pollen, N., Simon, A., 2005. Estimating the mechanical effects of riparian vegetation on  
446 stream bank stability using a fiber bundle model: MODELING ROOT  
447 REINFORCEMENT OF STREAM BANKS. *Water Resour Res* 41.  
448 <https://doi.org/10.1029/2004WR003801>  
449
- 450 Ristova, D., Barbez, E. (Eds), 2018. *Root Development: Methods and Protocols, Methods in*  
451 *Molecular Biology*. Humana Press. <https://doi.org/10.1007/978-1-4939-7747-5>  
452
- 453 Roumet, C., Urcelay, C., Díaz, S., 2006. Suites of root traits differ between annual and  
454 perennial species growing in the field. *New Phytol* 170, 357–368.  
455 <https://doi.org/10.1111/j.1469-8137.2006.01667.x>  
456
- 457 Schjøning, P., Lamandé, M., Munkholm, L.J., Lyngvig, H.S., Nielsen, J.A., 2016. Soil  
458 precompression stress, penetration resistance and crop yields in relation to differently-  
459 trafficked, temperate-region sandy loam soils. *Soil Tillage Res* 163, 298–308.  
460
- 461 Schjøning, P., Lamandé, M., 2018. Models for prediction of soil precompression stress from  
462 readily available soil properties. *Geoderma* 320, 115–125.  
463 <https://doi.org/10.1016/j.geoderma.2018.01.028>  
464
- 465 Schjøning, P., Lamandé, M., Keller, T., Labouriau, R., 2020. Subsoil shear strength –  
466 Measurements and prediction models based on readily available soil properties. *Soil*  
467 *Tillage Res* 200, 104638. <https://doi.org/10.1016/j.still.2020.104638>  
468
- 469 Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of  
470 image analysis. *Nat. Methods* 9, 671–675. <https://doi.org/10.1038/nmeth.2089>  
471
- 472 Stokes, A., Atger, C., Bengough, A.G., Fourcaud, T., Sidle, R.C., 2009. Desirable plant root  
473 traits for protecting natural and engineered slopes against landslides. *Plant Soil* 324,  
474 1–30. <https://doi.org/10.1007/s11104-009-0159-y>  
475
- 476 Stolte, J., Tesfai, M., Øygarden, L., Kværnø, S., Keizer, J., Verheijen, F., Panagos, P.,  
477 Ballabio, C., Hessel, R., European Commission, Joint Research Centre, Institute for  
478 Environment and Sustainability, 2015. *Soil threats in Europe*. Publications Office,  
479 Luxembourg  
480
- 481 Soane, B.D., Blackwell, P.S., Dickson, J.W., Painter, D.J., 1980. Compaction by agricultural  
482 vehicles: A review I. Soil and wheel characteristics. *Soil Tillage Res.* 1, 207–237.

483

484 Tengbeh, G.T., 1989. The effect of grass cover on bank erosion. PhD Thesis, Cranfield  
485 Institute of Technolgy, Silsoe College

486

487 Unger, P.W., Kaspar, T.C., 1994. Soil Compaction and Root Growth: A Review. Agron. J 86,  
488 759–766. <https://doi.org/10.2134/agronj1994.00021962008600050004x>

489

490 Yildiz, A., Graf, F., Rickli, C., Springman, S.M., 2018. Determination of the shearing  
491 behaviour of root-permeated soils with a large-scale direct shear apparatus. CATENA  
492 166, 98–113. <https://doi.org/10.1016/j.catena.2018.03.022>

493

494

495

496 *Table 1 Mean soil water content and soil matric potential for each of the three species and for the reference treatment. The*  
497 *letters showed differences between treatments.*

	<b>Soil water content (g.g-1)</b>	<b>Soil matric potential (kPa)</b>
<b><i>Brassica napus</i></b>	0,11 ± 0,02 <sup>a</sup>	-20.8 ± 6.07 <sup>a</sup>
<b><i>Vicia faba</i></b>	0,07 ± 0,01 <sup>b</sup>	-6.73 ± 0.04 <sup>a</sup>
<b><i>Secale cereale</i></b>	0,11 ± 0,03 <sup>a</sup>	-7.93 ± 0.31 <sup>a</sup>
<b>Bare soil</b>	0,12 ± 0,01 <sup>a</sup>	-6.2 ± 1.7 <sup>a</sup>

498 Soil water content values are the mean and standard error of 12 replicates.

499 Soil matric potential values are the mean and standard error of 3 replicates.

500 Means with the same letter within the same column were not significantly different at a 5% level based on the Pairwise  
501 Wilcoxon test.

502 Table 2: Root traits' mean and standard error per species. The letters showed differences between species. ARD = Average root diameter, ATRD = Average Taproot Diameter, DMC = Dry  
 503 Matter Content, RD = Root Density, RLD = Root Length Density, RVD = Root Volume Density, SRL = Specific Root Length, TRVD= Tap Root Volume Density

504

	<b>ARD (mm)</b>	<b>DMC (g.g<sup>-1</sup>)</b>	<b>SRL (m.g<sup>-1</sup>)</b>	<b>RLD (cm.cm<sup>-3</sup>)</b>	<b>RVD (cm<sup>3</sup>.cm<sup>-3</sup>)</b>	<b>RD (g.m<sup>-3</sup>)</b>	<b>TRVD (cm<sup>3</sup>.cm<sup>-3</sup>)</b>	<b>ATRD (cm)</b>
<i>Brassica</i>					0.0063±0.005		0.001±0.0005	
<i>napus</i>	0.35±0.06 <sup>b</sup>	0.18±0.03 <sup>a</sup>	38.6±9.1 <sup>b</sup>	2.05±0.85 <sup>b</sup>	4 <sup>b</sup>	525.1±133.8 <sup>b</sup>	<sup>a</sup>	0.53±0.08 <sup>a</sup>
<i>Secale</i>					0.0055±0.003			
<i>cereale</i>	0.34±0.05 <sup>b</sup>	0.13±0.01 <sup>b</sup>	77.4±14.2 <sup>a</sup>	2.82±0.91 <sup>a</sup>	<sup>b</sup>	373.6±138.3 <sup>c</sup>	0	
<i>Vicia</i>					0.0203±0.007	1397.6±309.3	0.0008±0.001	
<i>faba</i>	0.95±0.12 <sup>a</sup>	0.09±0.01 <sup>c</sup>	15.7±3.7 <sup>c</sup>	2.14±0.44 <sup>ab</sup>	6 <sup>a</sup>	<sup>a</sup>	<sup>a</sup>	0.64±0.16 <sup>a</sup>
Kruskal– Wallis test results	Chi square = 23.36, p-value = <0.001, df = 2	Chi square = 29.12, p-value = <0.001, df = 2	Chi square = 29.32, p-value = <0.001, df = 2	Chi square = 8.06, p-value = 0.018, df = 2	Chi square = 19.92, p-value = <0.001, df = 2	Chi square = 29.12, p-value = <0.001, df = 2	Chi square = 1.35, NS	Chi square = 2.54, NS

505 Values are the mean and standard error of 12 replicates\*

506 \**Vicia faba*'s TRVD and ATRD values are the mean and standard error of 6 replicates

507 Means with the same letter within the same column were not significantly different at a 5% level based on the Pairwise Wilcoxon test. NS = non-significant

508

509

510 *Table 3: Correlation matrix of root traits. Multiplying factors are shown using the significance of each correlation with p-*  
 511 *value > 0.05 = NS, p-value <0.05 = \*, p-value<0.01 = \*\* and p-value<0.001 = \*\*\*. ARD = Average root diameter, DMC =*  
 512 *Dry Matter Content, RD = Root Density, RLD = Root Length Density, RVD = Root Volume Density, SRL = Specific Root*  
 513 *Length*

	<b>ARD</b>	<b>DMC</b>	<b>RD</b>	<b>RLD</b>	<b>RVD</b>	<b>SRL</b>
<b>ARD</b>	1.00					
<b>DMC</b>	-0.76***	1.00				
<b>RD</b>	0.81***	-0.62***	1.00			
<b>RLD</b>	0.02NS	-0.32NS	0.02NS	1.00		
<b>RVD</b>	0.93***	-0.77***	0.85***	0.30NS	1.00	
<b>SRL</b>	-0.73***	0.42*	-0.86***	0.46**	-0.63***	1.00


514

515 *Table 4: Apparent cohesion and angle of internal friction, and linear regression coefficient R<sup>2</sup> for the fit with Eq. 2 for each*  
 516 *species and the reference treatment without plants.*

	<b>Angle of internal angle</b>	<b>Apparent cohesion (kPa)</b>	<b>R<sup>2</sup></b>
<i>Vicia faba</i>	17.9 <sup>oa</sup>	20.19 <sup>a</sup>	0.63
<i>Brassica napus</i>	15.6 <sup>ob</sup>	4.87 <sup>b</sup>	0.71
<i>Secale cereale</i>	12.6 <sup>ob</sup>	8.78 <sup>b</sup>	0.44
<i>Control without plant</i>	10.5 <sup>ob</sup>	9.58 <sup>b</sup>	0.68

517

518 **Figures caption**




519

520 *Figure 1: Components and schema of the shear box: a) Homemade system built to apply normal stress on the soil cylinder b)*  
 521 *Soil cylinders filled with remoulded soil used in the experiment. c) Shear box schema. 1) The computer-controlled motor and*  
 522 *actuator 2) The H3-C3 500kg load cell 3) The soil cylinders*

523

524



525

526

527


528

529

530

531

Figure 2: Boxplots of each root traits for each species: a) the average diameter (mm), b) the specific root length (cm.g<sup>-1</sup>), the root density (g.m<sup>-3</sup>), the dry matter content (g.g<sup>-1</sup>), the root length density (cm.cm<sup>-3</sup>), the root volume density (cm<sup>3</sup>.cm<sup>-3</sup>). Stars indicate significant differences between species according to the Pairwise Wilcoxon test: \*\*\* = p-value < 0.001, \*\* = p-value < 0.01, \* = p-value < 0.05. Means with the same letter were not significantly different at a 5% level based on the Pairwise Wilcoxon test.



532

533 *Figure 3: Linear regression  $\tau = c + \sigma \tan \phi$  for each species and the control without plants.*