



HAL
open science

Mechanobiology of muscle and myofibril morphogenesis

Nuno Miguel Luis, Frank Schnorrer

► **To cite this version:**

Nuno Miguel Luis, Frank Schnorrer. Mechanobiology of muscle and myofibril morphogenesis. *Cells & Development*, 2021, pp.203760. 10.1016/j.cdev.2021.203760 . hal-03471272

HAL Id: hal-03471272

<https://hal.science/hal-03471272v1>

Submitted on 8 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mechanobiology of muscle and myofibril morphogenesis

Nuno Miguel Luis¹ & Frank Schnorrer¹

¹ Aix Marseille University, CNRS, IBDM, Turing Centre for Living Systems, 13288 Marseille,
France

Correspondence should be addressed to:

nuno.luis@univ-amu.fr

frank.schnorrer@univ-amu.fr

keywords: muscle; sarcomere; biomechanics; mitochondria; *Drosophila*; titin;

Abstract

Muscles generate forces for animal locomotion. The contractile apparatus of muscles is the sarcomere, a highly regular array of large actin and myosin filaments linked by gigantic titin springs. During muscle development many sarcomeres assemble in series into long periodic myofibrils that mechanically connect the attached skeleton elements. Thus, ATP-driven myosin forces can power movement of the skeleton. Here we review muscle and myofibril morphogenesis, with a particular focus on their mechanobiology. We describe recent progress on the molecular structure of sarcomeres and their mechanical connections to the skeleton. We discuss current models predicting how tension coordinates the assembly of key sarcomeric components to periodic myofibrils that then further mature during development. This requires transcriptional feedback mechanisms that may help to coordinate myofibril assembly and maturation states with the transcriptional program. To fuel the varying energy demands of muscles we also discuss the close mechanical interactions of myofibrils with mitochondria and nuclei to optimally support powerful or enduring muscle fibers.

Highlights

- Update on the pseudo-crystalline structure of the sarcomere
- Molecularly detailing the development of force resistant muscle-tendon attachments
- Model for tension-driven myofibril self-assembly and myofibril maturation
- Mechanical coordination of muscle morphogenesis by transcriptional feedbacks
- Muscle-type specific mitochondrial and myofibril architectures

Contents

1. The musculoskeletal system

1.1 The sarcomere – a pseudo-crystalline force producing machine

1.2 Titin – a ruling and force sensing giant spring protein

2. Muscle attachments

2.1 Force-resistant muscle attachments: how to link a sarcomere to the skeleton

2.2 Quantifying molecular forces at muscle attachments

3. Muscle and sarcomere morphogenesis

3.1 Muscle-tendon attachment

3.2 Myofibril and sarcomere assembly

3.3 Myofibril and sarcomere maturation

3.4 Muscle fiber growth

3.5 Transcriptional dynamics recapitulates morphogenesis phases

4. Morphogenesis coordination within a myofiber

5. Conclusions and perspectives

1. The musculoskeletal system

An average human individual synthesises per day about half its body weight in form of ATP to power basic body functions. Each ATP/ADP molecule is turned over about 1000 times per day^{1,2}. An exercising athlete, such as a Tour de France cyclist, can consume up to 32 Mega Joule during a five-hour stage, more than five times the energy used during a regular day³. This enormous amount of energy is largely needed to fuel the myosin motor proteins of the muscles that produce the mechanical forces to power cyclists up the climbs. Even more impressive, flying animals power themselves into the air, with mosquitos beating their wings up to 1000 times per second giving rise to their well appreciated tone⁴.

For the efficient conversion of chemical energy into body movements it is critical that first, the energy is produced close to the consuming motors; second, the force generation of the many individual motor proteins is efficiently coordinated; and third, the force is converted into effective movements of the skeleton. If any of these conditions were not met, the muscles would run out of energy or largely produce heat instead of movement. Hence, the musculoskeletal system is a highly optimised biochemical apparatus that needs to form during animal development to match specific needs. In this review, we aim to cover the morphogenesis of muscles with a particular focus on the mechanobiology of its contractile machine, the sarcomere, the force transduction to the skeleton and lastly muscles' intracellular compartmentalisation optimised to fuel the molecular motors.

Skeletal muscles in vertebrates or body muscles in insects are large multinucleated cells, called muscle fibers, which can be tens of centimetres long in large vertebrates and are about 1 mm long in small fruit flies^{5,6}. Both muscle fibers ends are connected via tendon cells to the skeleton, a bony endoskeleton in vertebrates or a chitin-based exoskeleton in insects⁵⁻⁷. The precise insertion sites of the tendons at the skeletal elements together with hinges placed between different skeletal elements ensure that shortening of the muscle fibers causes effective

1 movement of legs or wings (Figure 1A). Hence, the morphogenesis of skeleton, tendons and
2 muscles must be precisely coordinated during animal development⁸⁻¹⁰.
3
4
5

6 7 *1.1 The sarcomere – a pseudo-crystalline force producing machine*

8
9 At the ‘heart’ of every muscle fiber is the muscles’ contractile machine, the sarcomere. We
10 know since pioneering work in the 1950ies that sarcomeres contract by a sliding mechanism
11
12
13
14^{11,12}: thick bipolar myosin filaments, which are anchored at the sarcomeric M-band, pull on
15 precisely organised parallel actin filaments, which are cross-linked at the sarcomeric Z-discs
16 bordering the sarcomere (Figure 1B). Hence, by pulling on the actin filaments the myosin
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

How is sarcomere length controlled? Actin and myosin filaments are stably connected
by the largest human protein, titin, whose N-terminus is bound to α -actinin at the Z-disc,
whereas its C-terminus is located in the middle of the sarcomere embedded into the M-band
(Figure 1B)¹⁴⁻¹⁷. Thus, titin is thought to serve as the ruler of the sarcomere determining its
resting length, usually between 2 and 3 μm in vertebrate muscle, depending on species and
muscle type¹⁸⁻²⁰.

Mature sarcomeres display a pseudo-crystalline order of their components. In
particular, insect flight muscles are so regular that structural changes of contracting sarcomeres
can be observed in the flying animal (during tethered flight) using X-ray diffraction^{21,22}. This
provided molecular evidence for the long-proposed stretch activation mechanism of insect
flight²³⁻²⁵: stretch-induced myosin and possibly troponin deformations are needed for efficient
binding of myosin motors to actin and thus to trigger fast oscillations of the antagonistic insect
flight muscle pairs. This showed that sarcomeres are excellently suited for structural studies.
Understanding the mature structure is the first step to understand how a sarcomere is built

1 during muscle development. Hence, we briefly summarise our current structural knowledge of
2 the sarcomeric parts.
3

4 **Z-discs:** despite its high regularity, resolving the native molecular structure of a
5 sarcomere in muscles is still an ambitious goal, especially for protein dense and complex
6 regions such as the Z-disc. However, significant progress has been made using X-ray
7 crystallography of purified sarcomeric proteins that are present in the Z-disc, in particular its
8 core member α -actinin. α -actinin contains an N-terminal actin binding domain, four central
9 spectrin repeats and a C-terminal calmodulin-like domain. The latter binds to the N-terminal
10 Z-repeats of titin and thus anchors titin directly at the Z-disc^{26,27}. The high-resolution structure
11 of the purified 200 kDa α -actinin dimer provides a first basis for how this interaction may be
12 regulated^{28,29}: the structure revealed that PIP2 binding to α -actinin's actin binding domain can
13 release the calmodulin-like domain and hence make it available for titin binding²⁸, as has been
14 suggested by biochemical experiments^{30,31}. Interestingly, mutations in α -actinin that alter the
15 titin interaction dynamics show defects in sarcomere formation in cell culture models²⁸.
16 Furthermore, the structural model of the anti-parallel α -actinin dimer places the two actin
17 filaments bound to it 23 nm apart, which is in almost perfect agreement with the 24 nm spacing
18 of tetragonal Z-disc lattice found in native mammalian muscle by electron microscopy³². The
19 structure further suggested that the bound actin filaments have rotational freedom with respect
20 to the stiff α -actinin rod, in analogy to a 'pivot and rod' structure, resulting in some structural
21 flexibility of Z-disc²⁹.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48

49 The flexibility of the Z-disc can also be affected by its thickness, which varies
50 depending on mammalian muscle type from 30 nm in fast fibers up to 100 nm in slow and
51 cardiac fibers³³ and about 80 nm in insect flight muscles³⁴ strongly suggesting that multiple
52 columns of crosslinked α -actinin exist at the Z-disc (Figure 1B). Differences might be
53 regulated by alternatively spliced titin isoforms containing variable number of Z-repeats^{29,35}
54
55
56
57
58
59
60
61
62
63
64
65

1 and thus more α -actinin binding sites or by presence of alternatively spliced Zasp (Z-disc
2 Alternatively Spliced Protein) isoforms, part of the Alp/Enigma conserved family of
3 scaffolding proteins, which can also interact with α -actinin via its PDZ domain and were
4 recently proposed to regulate the diameter of the Z-disc in insect muscles by an oligomerisation
5 mechanism³⁶. The native sarcomeric Z-disc also houses a proposed titin cross-linker telethonin
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Thin and thick filaments: to investigate the architecture of thin and thick filaments as well as their regulated interaction structural studies using purified *in vitro* assembled subcomplexes have been very insightful. Cryo-electron microscopy has revealed the architecture of a F-actin–tropomyosin complex with and without a myosin head demonstrating how a tropomyosin shift enables myosin binding and how the myosin motor head interacts with the actin filament at the molecular level⁴¹⁻⁴³. This provided a molecular model for how myosin forces are transduced to actin filaments in a regulated way in the sarcomere⁴³. Thus, *in vitro* assembled structures of sarcomeric components can teach us about the likely *in vivo* arrangement of these components and will be key structural models to resolve the native sarcomere structure *in situ*⁴⁴.

Taking advantage of the stability of sarcomere subcomplexes, *Drosophila* or *Lethocerus* flight muscle thick filaments have been isolated and analysed with cryo-electron tomography. Although not yet at atomic resolution, these studies visualised the long α -helices of the myosin rods in their native environment and revealed how their parallel arranged tails are packed together: interestingly, the individual myosin tails run in an angle from the main direction of the thick filament, forming a layered or ‘ribbon-like’ super-structure slowly

1 wrapping around the thick filament axis ^{45,46}, similar to what has been proposed in the 1970ies

2 47

3
4
5 **M-band:** thick filament associated proteins, such as myosin binding protein C in
6 vertebrates and myofilin or paramyosin in insect muscles, have not yet been fully resolved
7 structurally. Similarly, the native structure of the key M-band components myomesin and
8 obscurin are unknown. Myomesin forms C-terminal dimers with a suggested role to bridge
9 between neighbouring thick filaments (Figure 1B) ^{48,49}. Interestingly, these myomesin dimers
10 can be elastically stretched to more than double their length at forces around 30 pN that might
11 be physiologically relevant during muscle contractions ⁵⁰. This is consistent with a proposed
12 strain absorber function of myomesin at the M-band ⁵¹. Myomesin is not present in insect
13 muscles, however the large M-band protein obscurin (Unc-89) is conserved and required for
14 M-band alignment in *Drosophila* flight muscles and *C. elegans* body muscles ⁵²⁻⁵⁵. In
15 mammalian muscle, obscurin binds to the C-terminus of titin as well as to myomesin and hence
16 also localises to the M-band ⁵⁶⁻⁵⁸. However, its role in sarcomerogenesis is unclear ⁵⁹.

17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34 **Towards a first native sarcomere structure:** one recent approach towards resolving
35 the native sarcomere nanostructure with all its components included was to perform cryo-
36 focused ion beam-milling electron tomography on intact myofibrils isolated from mouse
37 muscles. This revealed that α -actinin dimers indeed cross-link anti-parallel actin filaments at
38 native Z-discs ⁶⁰. It also provided a structural explanation of how actin filaments, helped by
39 stabilising myosin cross-bridges, transition from their square-like symmetry at Z-discs to their
40 hexagonal pattern in the A-band ⁶⁰. Future cryo-electron tomography studies should reveal
41 further details of the native sarcomere nanostructure.

52 53 54 55 56 *1.2 Titin – a ruling and force sensing giant spring protein*

1 Titin is a key sarcomeric protein that in vertebrate muscle spans across half the sarcomere, with
2 its N-terminus anchored at the Z-disc and its C-terminus embedded into the M-band. Its
3
4 function in muscle architecture as well as muscle mechanics has been excellently reviewed
5
6 recently ^{61,62}, thus, we focus here on some recent highlights providing overwhelming evidence
7
8 that titin does rule sarcomere length and that titin can be a force sensing spring.
9
10

11
12 In mammalian skeletal or cardiac muscle, the myosin filament is 1.6 μm long (30 nm
13 diameter) ⁶³, consisting of about 300 muscle myosin II hexamers ⁶⁴⁻⁶⁶. The A-band part of titin
14
15 consists of immunoglobulin (Ig) and fibronectin type 3 (Fn3) domain super-repeats that are
16
17 constitutively present in the classical long titin isoforms in all muscle types. Hence, the length
18
19 of these titin super-repeats was long suggested as a blueprint, or ruler, for the observed
20
21 stereotypic length of the thick filaments. Indeed, genetic reduction of the number of A-band
22
23 Ig-Fn3 titin super-repeats caused the expected reduction of thick filament length and as
24
25 consequence, these muscle fibers produce lower forces ⁶⁷. This provided direct evidence for
26
27 titin's role regulating thick filament length and thus A-band length of mammalian sarcomeres
28
29 (Figure 1C).
30
31
32
33
34
35

36
37 If titin determines A-band length, which is constant during the sarcomere contraction
38
39 cycle, how about the I-band, whose length changes during contraction? The I-band part of titin
40
41 consists of a long series of Ig domains followed by an unstructured region rich in amino acids
42
43 proline, glutamic acid, valine and lysine (PEVK), which are known to function as an elastic
44
45 spring, extending and relaxing during sarcomere contraction cycles (Figure 1D) ^{61,62,68}.
46
47
48

49
50 The length of the I-band part of titin is muscle-type specific and regulated largely by
51
52 alternative splicing, with the general correlation that long Ig series and flexible PEVK
53
54 containing titin isoforms result in long I-bands present in mammalian skeletal and insect body
55
56 muscles, whereas short titin I-band versions, containing few PEVK and less Ig domains, result
57
58 in short I-bands present in cardiac muscle or insect flight muscle ^{61,69-71}. As a result, cardiac or
59
60
61
62
63
64
65

1 flight muscles are much stiffer compared to skeletal muscle. This has important functional
2 consequences, particularly prominent during mammalian heart beat or insect flight, both of
3 which are regulated with a large stretch-activation component ^{72,73}. It is hypothesized that this
4 stretch-activation (or length-dependent activation) requires the shorter and stiffer titin isoform
5 to communicate stretch effectively to fully activate actin and myosin filaments for the next
6 contraction cycle ^{22,61,69,74}.
7
8
9
10
11
12
13

14 Nevertheless, these findings provided only indirect evidence that titin rules I-band
15 length in muscles. Direct evidence was recently obtained by deleting a large part of titin's
16 PEVK spring using mouse genetics. Interestingly, the resulting mice are viable, grow to normal
17 size and have muscles with normal lengths, however, their sarcomeric I-bands are shorter,
18 resulting in shorter sarcomeres (Figure 1C). As a consequence, the muscle fiber assembles
19 more sarcomeres, likely by a tension-driven assembly mechanism (see section 3), to
20 mechanically connect over the same length, forming a muscle fiber that is stiffer compared to
21 wild type ⁷⁵. Surprisingly, these mice are performing equally well, at least in a simple running
22 wheel assay ⁷⁵. This provided direct evidence that titin's flexible PEVK spring determines the
23 length of the I-band and largely contributes to the elasticity and thus passive stiffness of the
24 muscle fiber.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 The elastic properties of the PEVK spring have been investigated using single molecule
42 atomic force microscopy or optical tweezers on native titin or recombinantly expressed protein
43 domains. This established that titin's PEVK sequence acts as a perfectly elastic spring at low
44 pico-newton forces ⁷⁶⁻⁷⁸, which are likely present on endogenous titin during sarcomere
45 contraction-relaxation cycles *in vivo* (Figure 1D) ^{17,79}. However, initial pulling experiments
46 suggested that much larger forces are needed to unfold titin's Ig domains ^{77,80,81}, questioning
47 whether or not Ig domain unfolding may also happen during normal muscle contraction cycles.
48 Interestingly, recent experiments pulling with magnetic tweezers on recombinantly expressed
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 titin fragments showed that at least some titin Ig domains can already unfold and refold at
2 forces around 10 pN⁸²⁻⁸⁴, which are likely present across individual titin molecules *in vivo*.
3
4 Refolding of Ig and PEVK domains can produce work and hence can contribute to force
5 produced by a contracting sarcomere, in addition to the active myosin force (Figure 1D),
6
7 similarly to how a stretched spring stores elastic energy released when recoiling back to normal
8
9 its normal length⁸²⁻⁸⁴. It has been estimated that the refolding of titin's domains under load can
10
11 produce a work of about 40zJ per titin molecule⁸⁴, which is comparable to the work produced
12
13 by one myosin motor cycle⁸⁵. On the other hand, during muscle contraction cycles an
14
15 antagonistic muscle is usually stretched (not shown in Figure 1A), hence work is needed to
16
17 unfold PEVK and potentially Ig domains in the antagonistic muscle. Thus, the net contribution
18
19 of titin refolding to the active contractile force *in vivo* remains unclear.
20
21
22
23
24
25

26
27 The range of forces present across a titin molecule in a native sarcomere has not yet
28
29 been determined. However, the recent introduction of a TEV protease cleavage site at the end
30
31 of the I-band part of mouse titin provided very strong evidence that titin does transduce forces
32
33 *in vivo*: cleaving titin in the I-band in permeabilised muscles *ex vivo* resulted in quick
34
35 dislocation of the myosin filaments and a splitting of the sarcomeric A-band, when muscle
36
37 contractions were induced⁸⁶. Passive tension in these titin cleaved muscles is reduced by more
38
39 than 50%^{84,86}. This demonstrated the key role of titin as an elastic buffer for the forces exerted
40
41 by myosin during muscle contraction by stably linking the Z-disc to the sarcomeric M-band.
42
43 Hence, titin is in the perfect position to sense the molecular forces present in a native sarcomere.
44
45
46
47
48
49
50

51 **2. Muscle attachments**

52 *2.1 Force-resistant muscle attachments: how to link a sarcomere to the skeleton*

53
54 As discussed above, effective body movements require not only effective force production in
55
56 the muscle fibers but also an efficient force transduction to the skeleton (Figure 1A). As bone
57
58
59
60
61
62
63
64
65

1 or chitinous skeleton is generally hard, whereas muscle tissue is soft, this generates high
2 demands for the material interfaces connecting the two tissues⁸⁷. Nature solved this challenge
3
4 by placing tendon cells between muscles and skeleton, hence allowing for a more graded
5
6 transition between these different cell types with such different stiffness. In mammals, tendons
7
8 cells are surrounded by a graded extracellular matrix (ECM) containing triple helical collagen
9
10 I, as well as collagen II, together with increasing amounts of mineralised calcium phosphate
11
12 (apatite) closer to the bone surface^{5,88}. In *Drosophila*, which only possess a single tendon cell
13
14 layer between the cuticle and the flight muscles (Figure 2A), the ECM facing the apical surface
15
16 of the tendons is formed by the gigantic, fiber forming protein Dumpy⁸⁹⁻⁹¹, whereas its basal
17
18 side connecting the tendons with the muscle fibers is an integrin-based adhesion with tigrin,
19
20 laminins and possibly collagen IV as ECM components⁹². Thus, a material with a graded
21
22 stiffness is used to stably anchor soft muscles to the stiff skeleton.
23
24
25
26
27
28

29 The terminal sarcomere of each myofibril faces a similar challenge as it needs to stably
30
31 attach to the ECM, which links muscle cells to tendons. One strategy to increase the anchoring
32
33 stability is to increase the surface contact. Hence *Drosophila* flight muscles and also the
34
35 intercalated discs of cardiomyocytes linking neighbouring sarcomeres, contain extensively
36
37 folded membrane indentations that largely increase the interaction surface (Figure 2A)^{93,94}.
38
39 Recently, super-resolution microscopy of flight muscle-tendon attachments revealed an
40
41 interesting layered organisation between the ECM and the terminal Z-disc of a myofibril⁹⁴:
42
43 proximal to the membrane is the integrin signalling layer with an extended integrin adaptor
44
45 talin. Talin, in its extended conformation, recruits vinculin in a force dependent manner^{95,96}.
46
47 Additionally, focal adhesion kinase (Fak) is present in the signalling layer (Figure 2A). This is
48
49 followed by an actin-rich layer with Arp2/3, filamin, Zasp and α -actinin nucleating and cross-
50
51 linking actin into a dense but not well-ordered actin network that links to the terminal Z-disc
52
53 of the myofibril (Figure 2A)⁹⁴. This complex organisation can enable various feedback
54
55
56
57
58
59
60
61
62
63
64
65

1 mechanisms to optimally adjust adhesion in response to the high force produced by muscle
2 fibers.
3
4
5
6

7 2.2 Quantifying molecular forces at muscle attachments 8

9 Integrin adhesions are very well characterised force transducing macro-molecular assemblies
10
11
12 ⁹⁷. One key component linking integrin tails to the actin cytoskeleton and hence mediating the
13 force transduction between integrins and the cytoskeleton is the integrin adaptor talin ^{98,99}.
14 Talin binds with its head to the beta-integrin tail and with its rod to the actin cytoskeleton,
15 which upon the application of force opens to expose vinculin binding sites that in turn enable
16 more actin filaments to pull on talin and induce focal adhesion growth ^{100,101}. Hence, talin is at
17 a key location to sense the range of forces present at muscle attachment sites.
18
19
20
21
22
23
24
25

26 Recent developments of genetically encodable FRET-based molecular force sensors
27 made it possible to quantify molecular forces across proteins in cell culture or *in vivo* ¹⁰². These
28 sensors contain a small mechanosensitive peptide between the FRET pair with a calibrated
29 force range between 2 to 11 pN ¹⁰². The application of high forces unfolds the elastic peptide,
30 resulting in low FRET values (Figure 2B).
31
32
33
34
35
36
37
38

39 Such molecular tension sensors have been inserted into the endogenous talin gene in
40 *Drosophila* to quantify forces across talin at flight muscle attachments ¹⁰³. Surprisingly, it was
41 found that in mature adult flight muscles at rest, only a small amount of the talin molecules is
42 under force. However, if talin amounts are reduced to about half, the flight muscle-tendon
43 junction mechanically ruptures. This suggests that recruiting a large pool of talin molecules to
44 mature attachments is an efficient way to share peak forces likely present during active flight
45 muscle contractions ¹⁰³. We can wonder how recruitment of such a large pool of talin
46 molecules, and all other sarcomeric components, is coordinated temporally and spatially during
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 muscle development in a regulated way? We are addressing this question in the following
2 section.
3
4
5
6

7 **3. Muscle and sarcomere morphogenesis**

8
9 Muscle development with a focus on muscle patterning or myoblast fusion has been excellently
10 reviewed recently ¹⁰⁴⁻¹⁰⁹. Here we focus on recent advances in mechanical aspects of muscle
11 morphogenesis and sarcomere development in particular. We use the well-studied *Drosophila*
12 indirect flight muscles as one model system to elaborate common concepts of muscle and
13 sarcomere morphogenesis as well as to highlight specific differences to mammalian skeletal
14 muscle development.
15
16
17
18
19
20
21
22
23
24
25

26 *3.1 Muscle-tendon attachment*

27
28 Each of the *Drosophila* indirect flight muscles is built during pupal stages by fusion of several
29 hundred myoblasts to large syncytial myotubes ^{110,111}. These myotubes elongate at both ends
30 towards their future attachment sites to which they establish a force-resistant integrin
31 dependent attachment at 24 h after puparium formation (24 h APF, Figure 3A) ^{112,113}. During
32 myotube elongation, myoblast fusion is largely restricted to the lateral membranes and absent
33 from the dynamic leading edges, which are extending towards the tendons ^{113,114}. A similar
34 myotube elongation process towards developing tendon attachments initiates myotube
35 development in other *Drosophila* muscle types and also in a wide range of vertebrate muscles
36 ^{8,115-118}. Hence, elongation and attachment are common principles of early muscle
37 morphogenesis that define the long axis of each muscle fiber. This axis also determines the
38 future muscle contraction axis, which will not be changed anymore during later steps of muscle
39 morphogenesis.
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 During myotube-tendon attachment, mechanical tension is built-up within the
2 myotendinous system: initially, tissue tension is low and only small amounts of integrin, talin
3 and other integrin associated proteins are present at myotube attachment sites. When tissue
4 tension increases, large amounts of integrin and talin are recruited and attachments mature over
5 the next 6 h (until 30 h APF) ^{103,113}. This tension increase causes the compaction of the flight
6 muscle myotube and its transition into an immature myofiber at 30 h APF (Figure 3A),
7 coinciding with the self-assembling of immature myofibrils that connect both attachments (see
8 section 3.2) ^{103,113,119}. At this stage, attachments have matured enough to be able to resist the
9 first active muscle twitchings ¹²⁰. This indicates that attachment maturation and myofibril
10 assembly must be coordinated during development.
11
12
13
14
15
16
17
18
19
20
21
22
23

24 How is attachment maturation triggered molecularly and how is it coordinated with
25 tissue tension increase? A first step to understand this coordination was undertaken by
26 quantifying molecular forces across talin at developing muscle attachments (see Figure 2B).
27 Interestingly, molecular tension across each talin molecule is larger at the early stages when
28 few talin molecules are present and gradually reduces when more molecules are recruited ¹⁰³.
29 This suggests that force dependent recruitment of talin is one important mechanism of
30 attachment maturation. Integrin associated proteins are also recruited in a force dependent
31 manner in *Drosophila* embryonic muscles. In particular the more peripheral members vinculin
32 and paxillin strongly depend on muscle myosin activity, and thus muscle twitching, for
33 effective recruitment and maturation of embryonic muscle attachments ¹²¹. This fits with the
34 interpretation that force-induced stretching of the talin rod will enable vinculin recruitment ⁹⁵.
35 This is also well established in mammalian cells ^{98,101,102} and in beating cardiomyocytes, in the
36 latter the oscillating forces across talin have even been directly quantified *in vitro* ¹²². Together,
37 this suggests that the tension dependent recruitment of large amounts of integrin complex
38 proteins provides a buffer for peak forces produced by muscle contractions. This mechanical
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 feedback mechanism coordinates muscle-tendon attachment morphogenesis with formation of
2 contractile sarcomeres and prevents a fatal rupture of attachments.
3
4
5
6

7 *3.2 Myofibril and sarcomere assembly*

8
9

10 Developing myofibers are very long cells connecting two skeletal elements. However, each
11 contractile sarcomere is only a few micrometres long (see Figure 1). How do myofibers manage
12 to assemble the correct number of sarcomeres into long myofibrils to bridge the long distance,
13 maintaining but not exceeding the correct tension level in each of its sarcomeric components?
14
15 One possible mechanism to solve this ‘dimension problem’ is suggested by the tension-driven
16 myofibril self-organisation hypothesis^{68,113}. This hypothesis, postulated in *Drosophila*, is
17 likely to also apply to mammalian muscles¹²³⁻¹²⁶. It suggests that mechanical tension is
18 coordinating the assembly of many smaller molecular complexes into higher order periodic
19 structures throughout the entire large muscle fiber. These long continuous and periodic
20 myofibrils mechanically bridge both ends of the entire muscle (Figure 3B). Live imaging has
21 shown that a periodic pattern in myofibrils indeed appears simultaneously over long distances
22 in *Drosophila* muscles as well as in cultured cardiomyocytes, suggesting a self-organisation
23 mechanism^{113,117,123,124}. How could such a phenomenon work molecularly?
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 In early stage myotubes, actomyosin is organised in a polar network along the long
42 muscle axis possessing a rather low molecular order without distinct myofibrils^{127,128}. As most
43 major sarcomeric components such as titin, muscle myosin or troponin are absent at this stage
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

120, the main force-producer is non-muscle myosin (Figure 3B). Non-muscle myosin
molecularly orders actin filaments into parallel arrays and is needed for efficient assembly of
immature periodic myofibrils in flight muscles and mammalian cardiomyocytes^{124,125,128}.
However, non-muscle myosin assembles into rather short, up to 300 nm long, bipolar filaments¹²⁹,
too short to span the entire A-band of a mature sarcomere. On the other hand, muscle

1 myosin assembles several hundred hexamers (composed of 2 heavy chains, 2 essential and 2
2 regulatory light chains) into 1.6 μm long filaments in mammalian muscle or into more than
3
4 3.0 μm long bipolar filaments in insect muscles⁶⁵. Hence with the onset of muscle myosin
5
6 expression, non-muscle myosin filaments are either replaced with muscle myosin ones, as
7
8 suggested by the pre-myofibril model^{130,131}, or the simple increase in muscle myosin levels
9
10 results in the dominant incorporation of muscle myosin into periodic immature myofibrils that
11
12 span the entire muscle fiber length (Figure 3B). These myofibrils are chains of immature
13
14 sarcomeres, containing periodic cross-linked actin filaments of increasing molecular order as
15
16 development progresses¹²⁸, as well as titin, α -actinin, obscurin, Zasp and likely other
17
18 components^{54,113,132,133}. Thus, not only actin and myosin but additional, possibly preassembled,
19
20 protein complexes are organising into these continuous immature myofibrils¹³⁴⁻¹³⁶. They may
21
22 use mechanical tension as a compass to orient themselves properly to assemble the correct
23
24 number of immature sarcomeres within each myofibril. The better the assembly, the higher the
25
26 active force that myosin can produce by pulling on ordered actin filaments (Figure 3A, B)⁶⁸.
27
28
29
30
31
32
33
34
35

36 3.3 Myofibril and sarcomere maturation

37 Immature sarcomeres are contractile, but they are not yet fully adapted to the specific needs of
38
39 the respective muscle type, be it a slow or fast mammalian fiber or an insect flight muscle fiber.
40
41 Sarcomere maturation is particularly well illustrated in *Drosophila* flight muscles, which
42
43 simultaneously assemble all their 2000 myofibrils per muscle fiber that mature synchronously
44
45 without addition of new myofibrils until the adult stage (Figure 3A, B)¹²⁰. During these
46
47 maturation steps sarcomere length grows from about 2 μm to 3.4 μm and myofibril diameter
48
49 progressively increases to reach about 1.5 μm ^{120,133,137,138}. How is such a defined sarcomere
50
51 growth achieved?
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Although not understood in molecular detail, it is likely that the relative amounts and the specific sarcomeric protein isoforms present at any time point determine the precise sarcomere architecture during sarcomere maturation. Expression of muscle myosin heavy chain isoforms changes via a switch in alternative splicing in flight muscles^{137,139} or via a transcriptional switch during mammalian muscle development from embryonic to fetal and finally to adult isoforms (Figure 3B)¹⁴⁰. Actin filament length growth is controlled by the expression of capping proteins like Tmod and Sals¹⁴¹⁻¹⁴³, as well as actin nucleators, including the formin family member Fhos or the short nebulin homolog Lasp during flight muscle sarcomere maturation^{144,145}. Similar proteins are needed to control actin filament length in mammalian muscle development, such as the formin homolog FHOD3¹⁴⁶, and in particular the gigantic protein nebulin whose proposed function as actin filament ruler has been reviewed in detail^{62,147-149}.

A particularly impressive way of how myofibril diameter can be regulated was recently proposed for flight muscles: the three distinct Zasp genes (*Zasp52*, *Zasp66* and *Zasp67*) encode either short Zasp isoforms only containing a PDZ-domain and the Zasp-motif (ZM) domain or long isoforms with additional 1 to 4 LIM domains^{132,138}. Interestingly, these Zasp proteins are core components of the Z-discs and can form aggregates by an interaction of their ZM domain with one of the LIM domains, whereas their PDZ domain interacts with α -actinin^{132,150}. Consequently, Zasp isoforms with one or more LIM domains promote ZASP complex growth and hence Z-disc growth, whereas short isoforms, lacking any LIM domain, block further growth. Strikingly, expression of ‘growing’ isoforms is high during myofibril maturation and diameter growth, whereas ‘blocking’ isoforms are upregulated only very late to terminate growth. Consistently, ‘growing’ isoforms are found located more centrally in the Z-discs, compared to blocking isoforms¹³⁸. These findings underscore the importance to coordinate

transcriptional regulation with sarcomere morphogenesis to form functional sarcomeres (see sections 3.4 and 3.5).

3.4 Muscle fiber growth

In mammalian embryonic and adult muscles, muscle fiber growth is largely driven by continuous myoblast fusion to existing fibers^{151,152} resulting in a fiber volume per nucleus correlation termed the myonuclear domain size¹⁵³. Careful volume measurements in *Drosophila* larval muscles also revealed a striking correlation of muscle fiber volume with nuclear volume¹⁵⁴. This correlation is maintained during the rapid insulin-induced growth of the *Drosophila* larval muscles: instead of additional myoblasts fusing, the larval muscle nuclei undergo several rounds of DNA endo-replication to produce very large nuclei, with up to 64 copies of each chromosome^{154,155}. Interestingly, the number of endo-replication cycles appears to be regulated mechanically via the LINC complex, which links the lamin meshwork below the nuclear membrane to the sarcomere and microtubule cytoskeleton¹⁵⁶. This provides a possible mechanical feedback mechanism to control nuclear endo-replication cycles during larval muscle fiber growth.

Further support for a significant mechanical input to the muscle nuclei is given by their specific positions. In *Drosophila* larval muscles, nuclei are initially transported by microtubules to clusters located close to the fiber ends^{157,158}. When sarcomeres have assembled and the fibers grow, the nuclei are redistributed and equally spaced along the fiber length, suggesting that a force-balance mechanism equally positions them per muscle volume^{154,159}. Furthermore, the larval nuclei are pushed to the surface of the larval muscles by the growing myofibrils and microtubules¹⁶⁰. Very similarly, nuclei in mature mammalian muscle fibers are also equally spaced along the fiber axis and are pushed to the fiber surface by the maturing and contracting myofibrils^{161,162}. In developing flight muscles, most nuclei are located centrally at

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

24 h APF before myofibrils are forming (Figure 3A). Upon myofibril assembly, they relocate and fill the space between the growing myofibril bundles¹⁶³. As a consequence, all nuclei adopt an elongated shape along the myofibril axis. Together, this strongly suggests that myofibrils impact nuclear shape and mechanics during muscle fiber growth.

Drosophila flight muscles grow tremendously after myoblast fusion has ceased, increasing their volume more than ten-fold in three days (Figure 3A)¹⁶³. Since their final number of myofibrils is already determined at 30 h APF, flight muscle length growth coincides with the addition of new sarcomeres to each existing myofibril, while their increase in diameter causes muscle diameter growth (Figure 3A, see section 3.3). Additionally, the mitochondrial content increases enormously (see section 4)¹⁶⁴. It is unknown if flight muscle nuclei undergo endoreplication, however a classical tissue growth pathway was recently shown to be essential during postmitotic flight muscle growth. Loss of the transcriptional regulator Yorkie (YAP, TAZ in mammals) or gain of the Hippo or Warts kinases, which are negatively regulating Yorkie, result in a block of flight muscle fiber growth¹⁶³. While these fibers finish myoblast fusion normally, they assemble fewer myofibrils, which then fail to mature, resulting in a dramatic muscle growth defect¹⁶³. The Hippo pathway has a well described key mechanosensory function at the epithelial cell cortex regulating stretch-dependent epithelial cell growth¹⁶⁵⁻¹⁶⁸. However, how can the Hippo pathway sense stretch in muscle fibers to induce muscle growth?

Interestingly, the classical upstream Hippo regulators Expanded, Merlin or Kibra that are involved in sensing stretch at the epithelial cell cortex do not seem to be required in flight muscles. Instead, the STRIPAK phosphatase complex controls Hippo activity. Slmap, one member of the STRIPAK complex binds to Dlg5, both of which are present in internal membranes forming the future T-tubules in proximity to the assembling myofibrils¹⁶³. Thus, it was speculated that in analogy to sensing stretch at the epithelial cell membrane, mechanical

1 stretch might be sensed at these internal muscle membranes to control Hippo kinase activity
2 before and after myofibril assembly. Consequently, expression of key sarcomeric components
3
4 is reduced upon loss of Yorkie, suggesting that mechanics and possibly myofibril assembly
5
6 status may feedback on the transcriptional state of the developing muscle fiber to induce its
7
8 further growth ¹⁶³.
9
10

11 *3.5 Transcriptional dynamics recapitulates morphogenesis phases*

12
13
14 As highlighted above, muscle fiber development can be divided into multiple phases such as
15
16 early muscle fate choice, myoblast fusion, myotube elongation and attachment, myofibril
17
18 assembly and maturation, as well as muscle fiber growth. Flight muscles are a suitable model
19
20 to study how transcriptional dynamics may impact the different phases, as their development
21
22 is largely homogenous. A systematic transcriptomic time-course covering flight muscle
23
24 development from myoblast stage in third instar larvae to mature adult flight muscles revealed
25
26 that the different morphogenesis phases are indeed preceded by important transcriptional
27
28 transitions ¹²⁰. As expected, genes implicated in patterning and cell division dominate the early
29
30 phase when myoblasts are dividing and their future muscle fate is determined (Figure 3C).
31
32 Then cell adhesion and muscle attachment genes become upregulated, fitting with the next
33
34 morphogenesis step, the attachment of the myotubes to tendons. At this stage, the expression
35
36 of sarcomeric genes is still very low, showing that only after successful myotube attachment
37
38 the expression of the key sarcomeric genes is initiated enabling myofibril assembly (Figure
39
40 3C) ¹²⁰. As discussed above, one signalling pathway that may react to mechanical input is the
41
42 Hippo pathway needed for effective sarcomere gene expression ¹⁶³. However, as Hippo
43
44 signalling appears to be particularly important only in flight muscles it remains to be seen how
45
46 other muscles regulate this important transition.
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 A subset of sarcomeric genes or gene isoforms, including the titin-like protein
2 Stretchin-MLCK, are only expressed at late stages during sarcomere maturation, once the
3 general sarcomere scaffold is built ¹²⁰. How could this late boost of sarcomere expression be
4 controlled? General muscle-specific regulators such as Mef2 are needed throughout flight
5 muscle development ¹⁶⁹ and also the flight muscle identity transcription factor Spalt is needed
6 both at the beginning to set up muscle fate and for the efficient boost at late stages ¹²⁰.
7 Interestingly, E2F, known for its key role during the cell cycle in many cell types ¹⁷⁰, is
8 dispensable for myoblast proliferation or fusion in *Drosophila* muscles and only needed at late
9 stages for myofibril assembly and, in particular, for myofibril maturation and fiber growth ¹⁷¹.
10 Together with the results from the Hippo pathway, this demonstrates that key transcriptional
11 regulators controlling cell cycle and growth in dividing tissue can be repurposed allowing
12 postmitotic muscle cell growth.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

28
29 It is important to note that flight muscle morphogenesis is not only controlled at the
30 transcriptional level, but alternative splicing is also critical in many different muscle types ^{70,172}.
31 In flight muscles, the splicing regulator Bruno (arrest), a CELF family member, is
32 transcriptionally induced by Spalt at the stage of myofibril assembly. However, Bruno is
33 largely needed to instruct flight muscle specific splicing during sarcomere maturation, for
34 example enabling splicing of the correct Stretchin-MLCK isoform. Consequently, *bruno*
35 mutant flight muscles show severe myofibril maturation defects ^{139,173}. This further illustrates
36 that transcriptional dynamics must be well synchronised with the morphogenesis phases of the
37 muscle fibers.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 **4. Morphogenesis coordination within a myofiber**

54 As illustrated above, maturing muscle fibers are very densely packed cells with a defined
55 tension axis. While it is well established that the mechanical properties of the extracellular
56
57
58
59
60
61
62
63
64
65

1 environment can directly impact cell fate, including muscle fate choice ^{174,175}, it is less clear
2 how tension and mechanical interactions between various intracellular organelles and
3 molecular machines are sensed and coordinated ¹⁷⁶. For muscle fibers, it particularly matters to
4 correctly partition their myofibrils, their mitochondria and their nuclei in a fiber-type specific
5 way to efficiently support insect flight or long-distance running (Figure 4).
6
7
8
9
10

11 The increasing crowdedness of differentiating muscle fibers strongly restricts diffusion
12 of mRNAs ¹⁷⁷. Thus, in mature muscle fibers most mRNAs undergo microtubule-based
13 transport when exiting the nucleus before they are translated ¹⁷⁷. Interestingly, mature
14 cardiomyocytes and skeletal muscle fibers show a strong enrichment of ribosomes close to
15 their sarcomeric Z-discs, at which not only mRNAs coding for sarcomeric proteins, but most
16 mRNAs appear to be translated. ¹⁷⁷⁻¹⁷⁹. If proteins are mostly synthesised at these ‘translation
17 hubs’ ¹⁷⁷, how do they efficiently reach their destinations in the nuclei, mitochondria or muscle
18 attachments, as protein diffusion in dense myofibers is restricted ¹⁸⁰? This is a particular
19 challenge for the most abundant organelle inside muscle fibers, the mitochondria, which import
20 most of their proteins from the cytoplasm ¹⁸¹. The answer might be resolved by live imaging
21 of individual protein dynamics in muscle in the future.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 Three-dimensional electron microscopy techniques ¹⁸² are now allowing to reconstruct
40 the detailed 3D morphology of muscle fibers revealing precise mitochondria shapes and their
41 relation to the myofibrils. Interestingly, in mammalian muscles mitochondria form elongated
42 networks whose locations and shapes depend on the muscle fiber type. In glycolytic fibers,
43 elongated mitochondria are largely oriented perpendicular to the contraction axis, whereas in
44 oxidative slow fibers mitochondria form a grid like network and in cardiomyocytes the network
45 is exclusively oriented in the direction of the myofibrils ¹⁸³. This leads to an increased
46 proximity between mitochondria and myosin filaments in myofibrils of oxidative and cardiac
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 myofibers compared to glycolytic ones and hence increase the efficiency of ATP delivery for
2 their function ¹⁸³.
3

4
5 Oxidative fibers and in particular cardiomyocytes contain up to 35% of mitochondria
6 per total muscle volume ^{183,184}, making it likely that mitochondria are not only providing the
7 myofibers with energy but also influence them mechanically. Indeed, in mammalian skeletal
8 muscles these mitochondrial networks form thin extensions or ‘nanotunnels’ close to the
9 aligned myofibrils at their sarcomeric I-bands, in proximity to the sarcoplasmic reticulum and
10 the T-tubule network ^{183,185,186}. Some of these tunnels are connecting two distant larger
11 mitochondria generating an extensive network (Figure 4, right panel) ¹⁸⁶. Interestingly, also in
12 *Drosophila* muscles the mitochondrial shapes and networks strongly depend on muscle fiber-
13 types: cross-striated leg muscles contain complex shaped mitochondrial networks with similar
14 mitochondrial nanochannels along their sarcomeric I-bands as in mammalian muscle fibers ¹⁶⁴.
15
16 In contrast, flight muscle mitochondria are elongated along each of the individual myofibrils
17 adopting ellipsoid-like shapes that ensheath myofibrils and isolate them from their
18 neighbouring myofibrils (Figure 4, left panel) ¹⁶⁴. This creates a large mitochondria-myofibril
19 interface that will enable an optimised ATP/ADP exchange to feed the high energy demands
20 during flight. Mitochondria-myofibril proximity is emphasized by the finding that myofibrils
21 are pushing the mitochondria into the elongated shapes since releasing the pressure or
22 disassembling the myofibrils results in a rounding of the mitochondria ¹⁶⁴. How can such close
23 ties arise in a coordinated way during muscle development?
24
25

26
27 Recent studies of developing *Drosophila* flight muscles showed that in myotubes,
28 before myofibrils assembly, mitochondria are thin elongated tubes and assemble to local
29 networks largely excluded from the actin filament bundles ¹⁶⁴. Concomitantly with the
30 assembly of the immature myofibrils (see Figure 3B), mitochondria intercalate between them
31 and insulate each myofibril from its neighbours ^{164,187}. When myofibrils start to mature and
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 grow in diameter, mitochondria also dramatically increase in volume and grow until they are
2 pushed from the myofibrils into ellipse shapes (Figure 4, left panel) ¹⁶⁴. At the same time,
3 expression of mitochondrial genes is strongly boosted (Figure 3C) ^{120,188} and mitochondria
4 dramatically increase their cristae density ^{128,187}, which are invaginations of the internal
5 mitochondria membranes that house the electron transport chain components ¹⁸⁹. The higher
6 the cristae surface, the higher the respiration capacity of the mitochondria, a feature particularly
7 associated with oxidative fibers ¹⁹⁰. Similarly dynamic mitochondria remodelling has been
8 observed in mammalian cardiomyocytes shortly after birth when myofibrils grow in size ¹⁹¹.
9 This demonstrates that mitochondria morphogenesis is intimately coordinated with myofibril
10 morphogenesis.
11
12
13
14
15
16
17
18
19
20
21
22
23

24 One possible mechanism how to coordinate morphogenesis of both organelles has
25 recently been suggested from studies of developing flight muscles. Changing the mitochondrial
26 fusion and fission dynamics during development in the direction of increased fusion and hence
27 larger mitochondrial clusters prevents the intercalation of the mitochondria between the
28 developing myofibrils ¹⁶⁴. Interestingly, this intercalation block causes the alignment of the
29 otherwise individualised flight muscle myofibrils changing them to a cross-striated
30 morphology normally found in leg muscles. Strikingly, the transcriptional status of these flight
31 muscles also changes and sarcomeric protein isoforms that are normally flight muscle specific
32 are reduced, whereas leg muscle specific proteins are boosted. This suggested a mechanical
33 feedback mechanism from the mitochondria and myofibrils back to the nuclei adapting their
34 transcriptional program ¹⁶⁴. As the nuclei are in close proximity to and pushed by the myofibrils
35 to their fiber-type specific positions (Figure 4 and see section 3), it is conceivable that the nuclei
36 themselves, possibly using the LINC complex in the nuclear membrane, sense the mechanical
37 changes and adapt their transcriptional program accordingly ^{156,192}. Appropriate mitochondrial
38 fusion and fission dynamics are also needed during mammalian muscle and heart development
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

193-195 leading us to speculate that similar mechanical feedback mechanisms may coordinate mitochondria with myofibril development in mammalian muscle.

5. Concluding remarks

Cellular mechanics controls the function and the development of most eukaryotic cell types and tissues, with a prime role in muscle and heart. The development of the muscle contractile machinery directly impacts the mechanical properties of muscle cells and hence feedbacks to coordinate its development with its surrounding tissues as well as with the other muscle cell organelles. We are just beginning to understand how such feedback mechanisms work molecularly to coordinate the formation of functional muscle cells. In the future, muscle and heart biology will continue to profit from novel high-resolution microscopy techniques, in particular electron tomography, to investigate the native muscle nanostructure and similarly from the *in vivo* application of live imaging tools that will allow to quantify individual protein dynamics and turnover rates as well as mechanical forces in developing muscle cells. Combining these technologies with genetic models and automated analysis tools will hopefully create the required synergism to reveal the nanostructure of the muscle and understand how biomechanical coordination between its components can assemble such beautiful contractile machines.

Competing interests

The authors declare that no competing interests exist.

Acknowledgements

1 We are grateful to Jerome Avellaneda for providing the electron microscopy image shown in
2 Figure 2A and for Sandra Lemke for integrin attachment and talin tensor sensor schemes
3 modified for Figure 2A and B. We thank Holger Knaut, Pierre Mangeol and Qiyao Mao for
4 insightful comments on this review.
5
6
7
8
9

10 **Funding**

11 This work was supported by the Centre National de la Recherche Scientifique (CNRS, N.M.L.,
12 F.S.), the European Research Council under the European Union's Horizon 2020 Programme
13 (ERC-2019-SyG 856118, F.S.), the excellence initiative Aix-Marseille University A*MIDEX
14 (ANR-11-IDEX-0001-02, F.S.), the French National Research Agency with ANR-ACHN
15 MUSCLE-FORCES and ANR-18-CE45-0016-01 MITO-DYNAMICS (F.S.), the Human
16 Frontiers Science Program (HFSP, RGP0052/2018, F.S.), the Bettencourt Foundation (F.S.),
17 the France-BioImaging national research infrastructure (ANR-10-INBS-04-01) and the
18 Investissements d'Avenir, French Government program managed by the French National
19 Research Agency (ANR-16-CONV-0001) and from Excellence Initiative of Aix-Marseille
20 University - A*MIDEX (Turing Centre for Living Systems).
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

39 The funders had no role in study design, data collection and analysis, decision to publish, or
40 preparation of the manuscript.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

1. Neupane, P., Bhujju, S., Thapa, N. & Bhattarai, H. K. ATP Synthase: Structure, Function and Inhibition. *Biomol Concepts* **10**, 1–10 (2019).
2. Capaldi, R. A. & Aggeler, R. Mechanism of the F(1)F(0)-type ATP synthase, a biological rotary motor. *Trends in Biochemical Sciences* **27**, 154–160 (2002).
3. Saris, W. H., van Erp-Baart, M. A., Brouns, F., Westerterp, K. R. & Hoor, ten, F. Study on food intake and energy expenditure during extreme sustained exercise: the Tour de France. *Int J Sports Med* **10 Suppl 1**, S26–31 (1989).
4. Arthur, B. J., Emr, K. S., Wyttenbach, R. A. & Hoy, R. R. Mosquito (*Aedes aegypti*) flight tones: frequency, harmonicity, spherical spreading, and phase relationships. *J Acoust Soc Am* **135**, 933–941 (2014).
5. Zelzer, E., Blitz, E., Killian, M. L. & Thomopoulos, S. Tendon-to-bone attachment: from development to maturity. *Birth Defects Res C Embryo Today* **102**, 101–112 (2014).
6. Moussian, B. Chitin: Structure, Chemistry and Biology. *Adv. Exp. Med. Biol.* **1142**, 5–18 (2019).
7. Sidor, C. & Schnorrer, F. Mechanobiology: Forging a strong matrix at tendons. *Curr Biol* **31**, R347–R350 (2021).
8. Schnorrer, F. & Dickson, B. J. Muscle building; mechanisms of myotube guidance and attachment site selection. *Developmental Cell* **7**, 9–20 (2004).
9. Schweitzer, R., Zelzer, E. & Volk, T. Connecting muscles to tendons: tendons and musculoskeletal development in flies and vertebrates. *Development* **137**, 2807–2817 (2010).
10. Felsenthal, N. & Zelzer, E. Mechanical regulation of musculoskeletal system development. *Development* **144**, 4271–4283 (2017).
11. Huxley, A. F. & Niedergerke, R. Structural changes in muscle during contraction; interference microscopy of living muscle fibres. **173**, 971–973 (1954).
12. Huxley, H. & Hanson, J. Changes in the cross-striations of muscle during contraction and stretch and their structural interpretation. *Nature* **173**, 973–976 (1954).
13. Huxley, H. E. Memories of early work on muscle contraction and regulation in the 1950's and 1960's. *Biochemical and Biophysical Research Communications* **369**, 34–42 (2008).
14. Fürst, D. O., Osborn, M., Nave, R. & Weber, K. The organization of titin filaments in the half-sarcomere revealed by monoclonal antibodies in immunoelectron microscopy: a map of ten nonrepetitive epitopes starting at the Z line extends close to the M line. *Journal of Cell Biology* **106**, 1563–1572 (1988).
15. Linke, W. A. *et al.* Towards a molecular understanding of the elasticity of titin. *Journal of Molecular Biology* **261**, 62–71 (1996).
16. Maruyama, K., Natori, R. & Nonomura, Y. New elastic protein from muscle. *Nature* **262**, 58–60 (1976).
17. Gautel, M. & Goulding, D. A molecular map of titin/connectin elasticity reveals two different mechanisms acting in series. *FEBS letters* **385**, 11–14 (1996).
18. Tskhovrebova, L. & Trinick, J. Titin: properties and family relationships. *Nature Reviews Molecular Cell Biology* **4**, 679–689 (2003).
19. Llewellyn, M. E., Barretto, R. P. J., Delp, S. L. & Schnitzer, M. J. Minimally invasive high-speed imaging of sarcomere contractile dynamics in mice and humans. *Nature* **454**, 784–788 (2008).

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
20. Ehler, E. & Gautel, M. The sarcomere and sarcomerogenesis. *Adv. Exp. Med. Biol.* **642**, 1–14 (2008).
 21. Dickinson, M., Bekyarova, T., Gore, D. & Maughan, D. Molecular dynamics of cyclically contracting insect flight muscle in vivo. *Nature* **433**, 330–334 (2005).
 22. Iwamoto, H. & Yagi, N. The molecular trigger for high-speed wing beats in a bee. *Science* **341**, 1243–1246 (2013).
 23. Pringle, J. W. The Bidder Lecture - The evolution of fibrillar muscle in insects. *J Exp Biol* **94**, 1–14 (1981).
 24. Josephson, R. K., Malamud, J. G. & Stokes, D. R. Asynchronous muscle: a primer. *J Exp Biol* **203**, 2713–2722 (2000).
 25. Agianian, B. *et al.* A troponin switch that regulates muscle contraction by stretch instead of calcium. *The EMBO Journal* **23**, 772–779 (2004).
 26. Young, P., Ferguson, C., Bañuelos, S. & Gautel, M. Molecular structure of the sarcomeric Z-disk: two types of titin interactions lead to an asymmetrical sorting of alpha-actinin. *The EMBO Journal* **17**, 1614–1624 (1998).
 27. Atkinson, R. A. *et al.* Ca²⁺-independent binding of an EF-hand domain to a novel motif in the alpha-actinin-titin complex. *Nat Struct Biol* **8**, 853–857 (2001).
 28. Ribeiro, E. de A. *et al.* The structure and regulation of human muscle α -actinin. *CELL* **159**, 1447–1460 (2014).
 29. Gautel, M. & Djinović-Carugo, K. The sarcomeric cytoskeleton: from molecules to motion. *Journal of Experimental Biology* **219**, 135–145 (2016).
 30. Fukami, K. *et al.* Requirement of phosphatidylinositol 4,5-bisphosphate for alpha-actinin function. *Nature* **359**, 150–152 (1992).
 31. Young, P. & Gautel, M. The interaction of titin and alpha-actinin is controlled by a phospholipid-regulated intramolecular pseudoligand mechanism. *The EMBO Journal* **19**, 6331–6340 (2000).
 32. Goldstein, M. A., Schroeter, J. P. & Sass, R. L. The Z lattice in canine cardiac muscle. *Journal of Cell Biology* **83**, 187–204 (1979).
 33. Luther, P. K. The vertebrate muscle Z-disc: sarcomere anchor for structure and signalling. *J Muscle Res Cell Motil* **30**, 171–185 (2009).
 34. Szikora, S. *et al.* Nanoscopy reveals the layered organization of the sarcomeric H-zone and I-band complexes. *The Journal of Cell Biology* **219**, (2020).
 35. Gautel, M., Goulding, D., Bullard, B., Weber, K. & Fürst, D. O. The central Z-disk region of titin is assembled from a novel repeat in variable copy numbers. *Journal of Cell Science* **109 (Pt 11)**, 2747–2754 (1996).
 36. González-Morales, N., Holenka, T. K. & Schöck, F. Filamin actin-binding and titin-binding fulfill distinct functions in Z-disc cohesion. *PLoS Genetics* **13**, e1006880 (2017).
 37. Zou, P. *et al.* Palindromic assembly of the giant muscle protein titin in the sarcomeric Z-disk. *Nature* **439**, 229–233 (2006).
 38. Narita, A., Takeda, S., Yamashita, A. & Maéda, Y. Structural basis of actin filament capping at the barbed-end: a cryo-electron microscopy study. *The EMBO Journal* **25**, 5626–5633 (2006).
 39. van der Ven, P. F. *et al.* Characterization of muscle filamin isoforms suggests a possible role of gamma-filamin/ABP-L in sarcomeric Z-disc formation. *Cell Motil. Cytoskeleton* **45**, 149–162 (2000).
 40. Liao, K. A., González-Morales, N. & Schöck, F. Zasp52, a Core Z-disc Protein in Drosophila Indirect Flight Muscles, Interacts with α -Actinin via an Extended PDZ Domain. *PLoS Genetics* **12**, e1006400 (2016).

- 1 41. Behrmann, E. *et al.* Structure of the rigor actin-tropomyosin-myosin complex. *CELL*
2 **150**, 327–338 (2012).
- 3 42. Ecken, von der, J. *et al.* Structure of the F-actin--tropomyosin complex. *Nature* **519**,
4 114–117 (2015).
- 5 43. Ecken, von der, J., Heissler, S. M., Pathan-Chhatbar, S., Manstein, D. J. & Raunser,
6 S. Cryo-EM structure of a human cytoplasmic actomyosin complex at near-atomic
7 resolution. *Nature* **534**, 724–728 (2016).
- 8 44. Taylor, K. A., Rahmani, H., Edwards, R. J. & Reedy, M. K. Insights into Actin-
9 Myosin Interactions within Muscle from 3D Electron Microscopy. *Int J Mol Sci* **20**,
10 (2019).
- 11 45. Hu, Z., Taylor, D. W., Reedy, M. K., Edwards, R. J. & Taylor, K. A. Structure of
12 myosin filaments from relaxed *Lethocerus* flight muscle by cryo-EM at 6 Å
13 resolution. *Sci Adv* **2**, e1600058 (2016).
- 14 46. Daneshparvar, N. *et al.* CryoEM structure of *Drosophila* flight muscle thick filaments
15 at 7 Å resolution. *Life Sci. Alliance* **3**, (2020).
- 16 47. Squire, J. M. General model of myosin filament structure. II. Myosin filaments and
17 cross-bridge interactions in vertebrate striated and insect flight muscles. *Journal of*
18 *Molecular Biology* **72**, 125–138 (1972).
- 19 48. Lange, S. *et al.* Dimerisation of myomesin: implications for the structure of the
20 sarcomeric M-band. *Journal of Molecular Biology* **345**, 289–298 (2005).
- 21 49. Pinotsis, N., Lange, S., Perriard, J.-C., Svergun, D. I. & Wilmanns, M. Molecular
22 basis of the C-terminal tail-to-tail assembly of the sarcomeric filament protein
23 myomesin. *The EMBO Journal* **27**, 253–264 (2008).
- 24 50. Pinotsis, N. *et al.* Superhelical architecture of the myosin filament-linking protein
25 myomesin with unusual elastic properties. *PLoS Biol* **10**, e1001261 (2012).
- 26 51. Berkemeier, F. *et al.* Fast-folding alpha-helices as reversible strain absorbers in the
27 muscle protein myomesin. *Proceedings of the National Academy of Sciences* **108**,
28 14139–14144 (2011).
- 29 52. Qadota, H., Blangy, A., Xiong, G. & Benian, G. M. The DH-PH region of the giant
30 protein UNC-89 activates RHO-1 GTPase in *Caenorhabditis elegans* body wall
31 muscle. *Journal of Molecular Biology* **383**, 747–752 (2008).
- 32 53. Benian, G. M. & Mayans, O. Titin and obscurin: giants holding hands and discovery
33 of a new Ig domain subset. *Journal of Molecular Biology* **427**, 707–714 (2015).
- 34 54. Katzemich, A. *et al.* The function of the M-line protein obscurin in controlling the
35 symmetry of the sarcomere in the flight muscle of *Drosophila*. *Journal of Cell*
36 *Science* **125**, 3367–3379 (2012).
- 37 55. Schnorrer, F. *et al.* Systematic genetic analysis of muscle morphogenesis and
38 function in *Drosophila*. *Nature* **464**, 287–291 (2010).
- 39 56. Fukuzawa, A. *et al.* Interactions with titin and myomesin target obscurin and
40 obscurin-like 1 to the M-band: implications for hereditary myopathies. *Journal of*
41 *Cell Science* **121**, 1841–1851 (2008).
- 42 57. Sauer, F., Vahokoski, J., Song, Y.-H. & Wilmanns, M. Molecular basis of the head-
43 to-tail assembly of giant muscle proteins obscurin-like 1 and titin. *EMBO Rep* **11**,
44 534–540 (2010).
- 45 58. Pernigo, S. *et al.* The crystal structure of the human titin:obscurin complex reveals a
46 conserved yet specific muscle M-band zipper module. *Journal of Molecular Biology*
47 **427**, 718–736 (2015).
- 48 59. Blondelle, J. *et al.* Murine obscurin and Obsl1 have functionally redundant roles in
49 sarcolemmal integrity, sarcoplasmic reticulum organization, and muscle metabolism.
50 *Communications Biology* **2**, 178–13 (2019).
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
60. Wang, Z. *et al.* The molecular basis for sarcomere organization in vertebrate skeletal muscle. *CELL* **184**, 2135–2150.e13 (2021).
 61. Linke, W. A. Titin Gene and Protein Functions in Passive and Active Muscle. *Annu. Rev. Physiol.* **80**, 389–411 (2018).
 62. Tskhovrebova, L. & Trinick, J. in *Fibrous Proteins: Structures and Mechanisms* **82**, 285–318 (Springer International Publishing, 2017).
 63. Tskhovrebova, L., Bennett, P., Gautel, M. & Trinick, J. Titin ruler hypothesis not refuted. *Proceedings of the National Academy of Sciences* **112**, E1172 (2015).
 64. Sosa, H., Popp, D., Ouyang, G. & Huxley, H. E. Ultrastructure of skeletal muscle fibers studied by a plunge quick freezing method: myofilament lengths. *Biophysj* **67**, 283–292 (1994).
 65. Dasbiswas, K., Hu, S., Schnorrer, F., Safran, S. A. & Bershadsky, A. D. Ordering of myosin II filaments driven by mechanical forces: experiments and theory. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* **373**, 20170114 (2018).
 66. Gokhin, D. S. & Fowler, V. M. A two-segment model for thin filament architecture in skeletal muscle. *Nature Reviews Molecular Cell Biology* **14**, 113–119 (2013).
 67. Tonino, P. *et al.* The giant protein titin regulates the length of the striated muscle thick filament. *Nature Communications* **8**, 1041 (2017).
 68. Lemke, S. B. & Schnorrer, F. Mechanical forces during muscle development. *Mechanisms of Development* **144**, 92–101 (2017).
 69. Burkart, C. *et al.* Modular proteins from the *Drosophila* sallimus (sls) gene and their expression in muscles with different extensibility. *Journal of Molecular Biology* **367**, 953–969 (2007).
 70. Spletter, M. L. & Schnorrer, F. Transcriptional regulation and alternative splicing cooperate in muscle fiber-type specification in flies and mammals. *Experimental Cell Research* **321**, 90–98 (2014).
 71. Granzier, H. & Labeit, S. Structure-function relations of the giant elastic protein titin in striated and smooth muscle cells. *Muscle Nerve* **36**, 740–755 (2007).
 72. Bullard, B. & Pastore, A. Regulating the contraction of insect flight muscle. *J Muscle Res Cell Motil* **32**, 303–313 (2011).
 73. Shiels, H. A. & White, E. The Frank-Starling mechanism in vertebrate cardiac myocytes. *J Exp Biol* **211**, 2005–2013 (2008).
 74. de Tombe, P. P. *et al.* Myofilament length dependent activation. *J. Mol. Cell. Cardiol.* **48**, 851–858 (2010).
 75. Brynnel, A. *et al.* Downsizing the molecular spring of the giant protein titin reveals that skeletal muscle titin determines passive stiffness and drives longitudinal hypertrophy. *eLife* **7**, 1065 (2018).
 76. Watanabe, K. *et al.* Molecular mechanics of cardiac titin's PEVK and N2B spring elements. *Journal of Biological Chemistry* **277**, 11549–11558 (2002).
 77. Linke, W. A. *et al.* PEVK domain of titin: an entropic spring with actin-binding properties. *J. Struct. Biol.* **137**, 194–205 (2002).
 78. Leake, M. C., Wilson, D., Gautel, M. & Simmons, R. M. The elasticity of single titin molecules using a two-bead optical tweezers assay. *Biophysj* **87**, 1112–1135 (2004).
 79. Linke, W. A., Ivemeyer, M., Mundel, P., Stockmeier, M. R. & Kolmerer, B. Nature of PEVK-titin elasticity in skeletal muscle. *Proceedings of the National Academy of Sciences of the United States of America* **95**, 8052–8057 (1998).
 80. Rief, M., Gautel, M., Oesterhelt, F., Fernandez, J. M. & Gaub, H. E. Reversible unfolding of individual titin immunoglobulin domains by AFM. *Science* **276**, 1109–1112 (1997).

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
81. Kellermayer, M. S., Smith, S. B., Granzier, H. L. & Bustamante, C. Folding-unfolding transitions in single titin molecules characterized with laser tweezers. *Science* **276**, 1112–1116 (1997).
 82. Rivas-Pardo, J. A. *et al.* Work Done by Titin Protein Folding Assists Muscle Contraction. *CellReports* **14**, 1339–1347 (2016).
 83. Eckels, E. C., Haldar, S., Tapia-Rojo, R., Rivas-Pardo, J. A. & Fernandez, J. M. The Mechanical Power of Titin Folding. *CellReports* **27**, 1836–1847.e4 (2019).
 84. Rivas-Pardo, J. A. *et al.* A HaloTag-TEV genetic cassette for mechanical phenotyping of proteins from tissues. *Nature Communications* **11**, 2060–13 (2020).
 85. Piazzesi, G. *et al.* Skeletal muscle performance determined by modulation of number of myosin motors rather than motor force or stroke size. *CELL* **131**, 784–795 (2007).
 86. Li, Y. *et al.* Graded titin cleavage progressively reduces tension and uncovers the source of A-band stability in contracting muscle. *eLife* **9**, (2020).
 87. Paxton, J. Z. & Baar, K. *Current progress in enthesis repair: strategies for interfacial tissue engineering.* (2012). doi:10.4172/2161-0533.S1-003
 88. Genin, G. M. *et al.* Functional grading of mineral and collagen in the attachment of tendon to bone. *Biophysical journal* **97**, 976–985 (2009).
 89. Bökel, C., Prokop, A. & Brown, N. H. Papillote and Piopio: Drosophila ZP-domain proteins required for cell adhesion to the apical extracellular matrix and microtubule organization. *Journal of Cell Science* **118**, 633–642 (2005).
 90. Chu, W.-C. & Hayashi, S. Mechano-chemical enforcement of tendon apical ECM into nano-filaments during Drosophila flight muscle development. *Curr Biol* **31**, 1366–1378.e7 (2021).
 91. Sidor, C. & Schnorrer, F. Mechanobiology: Forging a strong matrix at tendons. *Curr Biol* **31**, R347–R350 (2021).
 92. Brown, N. H. *et al.* Integrins as mediators of morphogenesis in Drosophila. **223**, 1–16 (2000).
 93. Reedy, M. C. & Beall, C. Ultrastructure of Developing Flight Muscle in Drosophila. II. Formation of the Myotendon Junction. *Dev. Biol.* **160**, 466–479 (1993).
 94. Green, H. J., Griffiths, A. G., Yläne, J. & Brown, N. H. Novel functions for integrin-associated proteins revealed by analysis of myofibril attachment in Drosophila. *eLife* **7**, 197 (2018).
 95. Klapholz, B. & Brown, N. H. Talin - the master of integrin adhesions. *Journal of Cell Science* jcs.190991–12 (2017). doi:10.1242/jcs.190991
 96. Klapholz, B. *et al.* Alternative mechanisms for talin to mediate integrin function. *Curr Biol* **25**, 847–857 (2015).
 97. Sun, Z., Guo, S. S. & Fässler, R. Integrin-mediated mechanotransduction. *The Journal of Cell Biology* **215**, 445–456 (2016).
 98. Sun, Z., Costell, M. & Fässler, R. Integrin activation by talin, kindlin and mechanical forces. *Nature cell biology* **21**, 25–31 (2019).
 99. Moser, M., Legate, K. R., Zent, R. & Fassler, R. The Tail of Integrins, Talin, and Kindlins. *Science* **324**, 895–899 (2009).
 100. Grashoff, C. *et al.* Measuring mechanical tension across vinculin reveals regulation of focal adhesion dynamics. *Nature* **466**, 263–266 (2010).
 101. Elosegui-Artola, A. *et al.* Mechanical regulation of a molecular clutch defines force transmission and transduction in response to matrix rigidity. *Nature cell biology* **18**, 540–548 (2016).
 102. Fischer, L. S., Rangarajan, S., Sadhanasatish, T. & Grashoff, C. Molecular Force Measurement with Tension Sensors. *Annu Rev Biophys* (2021). doi:10.1146/annurev-biophys-101920-064756

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
103. Lemke, S. B., Weidemann, T., Cost, A.-L., Grashoff, C. & Schnorrer, F. A small proportion of Talin molecules transmit forces at developing muscle attachments in vivo. *PLoS Biol* **17**, e3000057 (2019).
 104. Schiaffino, S. & Reggiani, C. Fiber types in mammalian skeletal muscles. *Physiological Reviews* **91**, 1447–1531 (2011).
 105. Bothe, I. & Baylies, M. K. Drosophila myogenesis. *Curr Biol* **26**, R786–91 (2016).
 106. Gunage, R. D., Dhanyasi, N., Reichert, H. & VijayRaghavan, K. Drosophila adult muscle development and regeneration. *Semin. Cell Dev. Biol.* **72**, 56–66 (2017).
 107. Sampath, S. C., Sampath, S. C. & Millay, D. P. Myoblast fusion confusion: the resolution begins. *Skeletal Muscle* **8**, 3–10 (2018).
 108. Lee, D. M. & Chen, E. H. Drosophila Myoblast Fusion: Invasion and Resistance for the Ultimate Union. *Annu Rev Genet* **53**, 67–91 (2019).
 109. Poovathumkadavil, P. & Jagla, K. Genetic Control of Muscle Diversification and Homeostasis: Insights from Drosophila. *Cells* **9**, 1543 (2020).
 110. Fernandes, J., Bate, M. & VijayRaghavan, K. Development of the indirect flight muscles of Drosophila. *Development* **113**, 67–77 (1991).
 111. Mukherjee, P., Gildor, B., Shilo, B.-Z., VijayRaghavan, K. & Schejter, E. D. The actin nucleator WASp is required for myoblast fusion during adult Drosophila myogenesis. *Development* **138**, 2347–2357 (2011).
 112. Fernandes, J., Celniker, S. & VijayRaghavan, K. Development of the Indirect Flight Muscle Attachment Sites in Drosophila: Role of the PS Integrins and the stripe Gene. **176**, 166–184 (1996).
 113. Weitkunat, M., Kaya-Copur, A., Grill, S. W. & Schnorrer, F. Tension and force-resistant attachment are essential for myofibrillogenesis in Drosophila flight muscle. *Curr Biol* **24**, 705–716 (2014).
 114. Sarov, M. *et al.* A genome-wide resource for the analysis of protein localisation in Drosophila. *eLife* **5**, e12068 (2016).
 115. Kardon, G., Harfe, B. D. & Tabin, C. J. A Tcf4-positive mesodermal population provides a prepattern for vertebrate limb muscle patterning. *Developmental Cell* **5**, 937–944 (2003).
 116. Gros, J., Serralbo, O. & Marcelle, C. WNT11 acts as a directional cue to organize the elongation of early muscle fibres. *Nature* **457**, 589–593 (2009).
 117. Weitkunat, M., Brasse, M., Bausch, A. R. & Schnorrer, F. Mechanical tension and spontaneous muscle twitching precede the formation of cross-striated muscle in vivo. *Development* **144**, 1261–1272 (2017).
 118. Wood, A. J. & Currie, P. D. Development Aspects of Zebrafish Myotendinous Junction: a Model System for Understanding Muscle Basement Membrane Formation and Failure. 1–9 (2017). doi:10.1007/s40139-017-0140-z
 119. Vega-Macaya, F., Manieu, C., Valdivia, M., Mlodzik, M. & Olguin, P. Establishment of the Muscle-Tendon Junction During Thorax Morphogenesis in Drosophila Requires the Rho-Kinase. **204**, 1139–1149 (2016).
 120. Spletter, M. L. *et al.* A transcriptomics resource reveals a transcriptional transition during ordered sarcomere morphogenesis in flight muscle. *eLife* **7**, 1361 (2018).
 121. Bulgakova, N. A., Wellmann, J. & Brown, N. H. Diverse integrin adhesion stoichiometries caused by varied actomyosin activity. *Open Biol* **7**, (2017).
 122. Pandey, P. *et al.* Cardiomyocytes Sense Matrix Rigidity through a Combination of Muscle and Non-muscle Myosin Contractions. *Developmental Cell* **44**, 326–336.e3 (2018).

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
123. Chopra, A. *et al.* Force Generation via β -Cardiac Myosin, Titin, and α -Actinin Drives Cardiac Sarcomere Assembly from Cell-Matrix Adhesions. *Developmental Cell* **44**, 87–96.e5 (2018).
 124. Fenix, A. M. *et al.* Muscle-specific stress fibers give rise to sarcomeres in cardiomyocytes. *eLife* **7**, 1111 (2018).
 125. Taneja, N., Neininger, A. C. & Burnette, D. T. Coupling to substrate adhesions drives the maturation of muscle stress fibers into myofibrils within cardiomyocytes. *Molecular Biology of the Cell* **31**, 1273–1288 (2020).
 126. Mao, Q. *et al.* Tension-driven multi-scale self-organisation in human iPSC-derived muscle fibers. *bioRxiv* 2021.10.24.465614 (2021).
 127. Reedy, M. C. & Beall, C. Ultrastructure of developing flight muscle in *Drosophila*. I. Assembly of myofibrils. *Dev. Biol.* **160**, 443–465 (1993).
 128. Loison, O. *et al.* Polarization-resolved microscopy reveals a muscle myosin motor-independent mechanism of molecular actin ordering during sarcomere maturation. *PLoS Biol* **16**, e2004718 (2018).
 129. Hu, S. *et al.* Long-range self-organization of cytoskeletal myosin II filament stacks. *Nature cell biology* **19**, 133–141 (2017).
 130. Rhee, D., Sanger, J. M. & Sanger, J. W. The premyofibril: evidence for its role in myofibrillogenesis. *Cell Motil. Cytoskeleton* **28**, 1–24 (1994).
 131. Sanger, J. W., Wang, J., Fan, Y., White, J. & Sanger, J. M. Assembly and dynamics of myofibrils. *Journal of Biomedicine and Biotechnology* **2010**, 858606 (2010).
 132. Katzemich, A., Liao, K. A., Czerniecki, S. & Schöck, F. Alp/Enigma family proteins cooperate in Z-disc formation and myofibril assembly. *PLoS Genetics* **9**, e1003342 (2013).
 133. Orfanos, Z. *et al.* Sallimus and the dynamics of sarcomere assembly in *Drosophila* flight muscles. *Journal of Molecular Biology* **427**, 2151–2158 (2015).
 134. Holtzer, H. *et al.* Independent assembly of 1.6 microns long bipolar MHC filaments and I-Z-I bodies. *Cell Struct Funct* **22**, 83–93 (1997).
 135. Ehler, E., Rothen, B. M., Hämmerle, S. P., Komiyama, M. & Perriard, J. C. Myofibrillogenesis in the developing chicken heart: assembly of Z-disk, M-line and the thick filaments. *Journal of Cell Science* **112 (Pt 10)**, 1529–1539 (1999).
 136. Rui, Y., Rui, Y., Bai, J. & Perrimon, N. Sarcomere formation occurs by the assembly of multiple latent protein complexes. *PLoS Genetics* **6**, e1001208 (2010).
 137. Orfanos, Z. & Sparrow, J. C. Myosin isoform switching during assembly of the *Drosophila* flight muscle thick filament lattice. *Journal of Cell Science* **126**, 139–148 (2013).
 138. González-Morales, N. *et al.* Myofibril diameter is set by a finely tuned mechanism of protein oligomerization in *Drosophila*. *eLife* **8**, (2019).
 139. Spletter, M. L. *et al.* The RNA-binding protein Arrest (Bruno) regulates alternative splicing to enable myofibril maturation in *Drosophila* flight muscle. *EMBO Rep* **16**, 178–191 (2015).
 140. Schiaffino, S., Rossi, A. C., Smerdu, V., Leinwand, L. A. & Reggiani, C. Developmental myosins: expression patterns and functional significance. *Skeletal Muscle* **5**, 22–14 (2015).
 141. Mardahl-Dumesnil, M. & Fowler, V. M. Thin filaments elongate from their pointed ends during myofibril assembly in *Drosophila* indirect flight muscle. *Journal of Cell Biology* **155**, 1043–1053 (2001).
 142. Bai, J., Hartwig, J. H. & Perrimon, N. SALS, a WH2-domain-containing protein, promotes sarcomeric actin filament elongation from pointed ends during *Drosophila* muscle growth. *Developmental Cell* **13**, 828–842 (2007).

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
143. Gokhin, D. S., Ochala, J., Domenighetti, A. A. & Fowler, V. M. Tropomodulin 1 directly controls thin filament length in both wild-type and tropomodulin 4-deficient skeletal muscle. *Development* **142**, 4351–4362 (2015).
 144. Fernandes, I. & Schöck, F. The nebulin repeat protein Lasp regulates I-band architecture and filament spacing in myofibrils. *The Journal of Cell Biology* **206**, 559–572 (2014).
 145. Shwartz, A., Dhanyasi, N., Schejter, E. D. & Shilo, B.-Z. The Drosophila formin Fhos is a primary mediator of sarcomeric thin-filament array assembly. *eLife* **5**, D786 (2016).
 146. Iskratsch, T. *et al.* Formin follows function: a muscle-specific isoform of FHOD3 is regulated by CK2 phosphorylation and promotes myofibril maintenance. *The Journal of Cell Biology* **191**, 1159–1172 (2010).
 147. Bang, M.-L. *et al.* Nebulin-deficient mice exhibit shorter thin filament lengths and reduced contractile function in skeletal muscle. *Journal of Cell Biology* **173**, 905–916 (2006).
 148. Pappas, C. T., Krieg, P. A. & Gregorio, C. C. Nebulin regulates actin filament lengths by a stabilization mechanism. *The Journal of Cell Biology* **189**, 859–870 (2010).
 149. Chu, M., Gregorio, C. C. & Pappas, C. T. Nebulin, a multi-functional giant. *Journal of Experimental Biology* **219**, 146–152 (2016).
 150. Zhou, Q. *et al.* Ablation of Cypher, a PDZ-LIM domain Z-line protein, causes a severe form of congenital myopathy. *Journal of Cell Biology* **155**, 605–612 (2001).
 151. White, R. B., Biérinx, A.-S., Gnocchi, V. F. & Zammit, P. S. Dynamics of muscle fibre growth during postnatal mouse development. *BMC Dev Biol* **10**, 21–11 (2010).
 152. Abmayr, S. M. & Pavlath, G. K. Myoblast fusion: lessons from flies and mice. *Development* **139**, 641–656 (2012).
 153. Snijders, T. *et al.* Satellite cells in human skeletal muscle plasticity. *Front Physiol* **6**, 283 (2015).
 154. Windner, S. E., Manhart, A., Brown, A., Mogilner, A. & Baylies, M. K. Nuclear Scaling Is Coordinated among Individual Nuclei in Multinucleated Muscle Fibers. *Developmental Cell* **49**, 48–62.e3 (2019).
 155. Demontis, F. & Perrimon, N. Integration of Insulin receptor/Foxo signaling and dMyc activity during muscle growth regulates body size in Drosophila. *Development* **136**, 983–993 (2009).
 156. Wang, S. *et al.* Mechanotransduction via the LINC complex regulates DNA replication in myonuclei. *The Journal of Cell Biology* **217**, 2005–2018 (2018).
 157. Metzger, T. *et al.* MAP and kinesin-dependent nuclear positioning is required for skeletal muscle function. *Nature* **484**, 120–124 (2012).
 158. Folker, E. S., Schulman, V. K. & Baylies, M. K. Muscle length and myonuclear position are independently regulated by distinct Dynein pathways. *Development* **139**, 3827–3837 (2012).
 159. Elhanany-Tamir, H. *et al.* Organelle positioning in muscles requires cooperation between two KASH proteins and microtubules. *The Journal of Cell Biology* **198**, 833–846 (2012).
 160. Wang, S., Reuveny, A. & Volk, T. Nesprin provides elastic properties to muscle nuclei by cooperating with spectraplakins and EB1. *The Journal of Cell Biology* **209**, 529–538 (2015).
 161. Roman, W. *et al.* Myofibril contraction and crosslinking drive nuclear movement to the periphery of skeletal muscle. *Nature cell biology* **19**, 1189–1201 (2017).

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
162. Roman, W. & Gomes, E. R. Nuclear positioning in skeletal muscle. *Semin. Cell Dev. Biol.* **82**, 51–56 (2018).
 163. Kaya-Copur, A. *et al.* The Hippo pathway controls myofibril assembly and muscle fiber growth by regulating sarcomeric gene expression. *eLife* **10**, (2021).
 164. Avellaneda, J. *et al.* Myofibril and mitochondria morphogenesis are coordinated by a mechanical feedback mechanism in muscle. *Nature Communications* 1–18 (2021). doi:10.1038/s41467-021-22058-7
 165. Pan, D. The hippo signaling pathway in development and cancer. *Developmental Cell* **19**, 491–505 (2010).
 166. Deng, H. *et al.* Spectrin regulates Hippo signaling by modulating cortical actomyosin activity. *eLife* **4**, e06567 (2015).
 167. Fletcher, G. C. *et al.* The Spectrin cytoskeleton regulates the Hippo signalling pathway. *The EMBO Journal* **34**, 940–954 (2015).
 168. Fletcher, G. C. *et al.* Mechanical strain regulates the Hippo pathway in Drosophila. *Development* **145**, dev159467–10 (2018).
 169. Soler, C., Han, J. & Taylor, M. V. The conserved transcription factor Mef2 has multiple roles in adult Drosophila musculature formation. *Development* **139**, 1270–1275 (2012).
 170. van den Heuvel, S. & Dyson, N. J. Conserved functions of the pRB and E2F families. *Nature Reviews Molecular Cell Biology* **9**, 713–724 (2008).
 171. Zappia, M. P. & Frolov, M. V. E2F function in muscle growth is necessary and sufficient for viability in Drosophila. *Nature Communications* **7**, 10509 (2016).
 172. Nikonova, E., Kao, S.-Y. & Spletter, M. L. Contributions of alternative splicing to muscle type development and function. 1–0 (2020). doi:10.1016/j.semcdb.2020.02.003
 173. Oas, S. T., Bryantsev, A. L. & Cripps, R. M. Arrest is a regulator of fiber-specific alternative splicing in the indirect flight muscles of Drosophila. *The Journal of Cell Biology* **206**, 895–908 (2014).
 174. Swift, J. *et al.* Nuclear lamin-A scales with tissue stiffness and enhances matrix-directed differentiation. *Science* **341**, 1240104 (2013).
 175. Engler, A. J., Sen, S., Sweeney, H. L. & Discher, D. E. Matrix elasticity directs stem cell lineage specification. *CELL* **126**, 677–689 (2006).
 176. Shamipour, S., Caballero-Mancebo, S. & Heisenberg, C.-P. Cytoplasm's Got Moves. *Developmental Cell* **56**, 213–226 (2021).
 177. Denes, L. T., Kelley, C. P. & Wang, E. T. Microtubule-based Transport is Essential to Distribute RNA and Nascent Protein in Skeletal Muscle. *Nature Communications* **12**, 6079-19 (2021).
 178. Rudolph, F. *et al.* Resolving titin's lifecycle and the spatial organization of protein turnover in mouse cardiomyocytes. *Proceedings of the National Academy of Sciences* **112**, 201904385 (2019).
 179. Lewis, Y. E. *et al.* Localization of transcripts, translation, and degradation for spatiotemporal sarcomere maintenance. *J. Mol. Cell. Cardiol.* **116**, 16–28 (2018).
 180. Papadopoulos, S., Jürgens, K. D. & Gros, G. Protein diffusion in living skeletal muscle fibers: dependence on protein size, fiber type, and contraction. *Biophysj* **79**, 2084–2094 (2000).
 181. Wiedemann, N. & Pfanner, N. Mitochondrial Machineries for Protein Import and Assembly. *Annu. Rev. Biochem.* **86**, 685–714 (2017).
 182. Heinrich, L. *et al.* Automatic whole cell organelle segmentation in volumetric electron microscopy. *bioRxiv* 2020.11.14.382143 (2020). doi:10.1101/2020.11.14.382143

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
183. Bleck, C. K. E., Kim, Y., Willingham, T. B. & Glancy, B. Subcellular connectomic analyses of energy networks in striated muscle. *Nature Communications* **9**, 1–11 (2018).
 184. Mishra, P., Varuzhanyan, G., Pham, A. H. & Chan, D. C. Mitochondrial Dynamics is a Distinguishing Feature of Skeletal Muscle Fiber Types and Regulates Organellar Compartmentalization. *Cell Metabolism* **22**, 1033–1044 (2015).
 185. Glancy, B. *et al.* Mitochondrial reticulum for cellular energy distribution in muscle. *Nature* **523**, 617–620 (2015).
 186. Vincent, A. E. *et al.* Quantitative 3D Mapping of the Human Skeletal Muscle Mitochondrial Network. *CellReports* **26**, 996–1009.e4 (2019).
 187. Sauerwald, J., Backer, W., Matzat, T., Schnorrer, F. & Luschnig, S. Matrix metalloproteinase 1 modulates invasive behavior of tracheal branches during entry into *Drosophila* flight muscles. *eLife* **8**, 1079 (2019).
 188. Yim, A. *et al.* mitoXplorer, a visual data mining platform to systematically analyze and visualize mitochondrial expression dynamics and mutations. *Nucleic Acids Res.* **48**, 605–632 (2020).
 189. Zick, M., Rabl, R. & Reichert, A. S. Cristae formation-linking ultrastructure and function of mitochondria. *Biochim. Biophys. Acta* **1793**, 5–19 (2009).
 190. Cogliati, S. *et al.* Mitochondrial cristae shape determines respiratory chain supercomplexes assembly and respiratory efficiency. *CELL* **155**, 160–171 (2013).
 191. Porter, G. A. *et al.* Bioenergetics, mitochondria, and cardiac myocyte differentiation. *Prog Pediatr Cardiol* **31**, 75–81 (2011).
 192. Janota, C. S., Calero-Cuenca, F. J. & Gomes, E. R. The role of the cell nucleus in mechanotransduction. *Current opinion in cell biology* **63**, 204–211 (2020).
 193. Dorn, G. Mitochondrial fission/fusion and cardiomyopathy. *Current Opinion in Genetics & Development* **38**, 38–44 (2016).
 194. Favaro, G. *et al.* DRP1-mediated mitochondrial shape controls calcium homeostasis and muscle mass. *Nature Communications* **10**, 2576–17 (2019).
 195. Silva Ramos, E. *et al.* Mitochondrial fusion is required for regulation of mitochondrial DNA replication. *PLoS Genetics* **15**, e1008085 (2019).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Box 1.

Sarcomere - defined as the region flanked by two Z-discs with a M-band at its center. It is composed of bipolar myosin filaments intercalated with polar actin filaments. Sliding of actin filaments by pulling myosin motors results in sarcomere shortening.

Myofibril - a concatenation of sarcomeres that spans the length of a muscle fiber. Myofibrils form bundles which laterally align in cross-striated mammalian skeletal muscle or insect leg muscle. In fibrillar insect flight muscle, each myofibril is isolated from its neighbours by surrounding mitochondria.

I-band - or 'isotropic' band region, is the zone of the sarcomere without any myosin filaments. It spans across one Z-disc into to the adjacent sarcomere. When the muscle fiber contracts the I-band shortens.

A-band - or 'anisotropic' band region, is the region encompassing myosin bipolar filaments. Myosin filaments are cross-linked at the central M-band by M-band proteins, such as myomesin. A-band length is constant during sarcomere contraction.

M-band - the central part of the sarcomere composed of proteins that laterally cross-link the myosin filaments, likely providing rigidity and stability.

Z-disc - the connection between two sarcomeres. It is composed of several actin cross-linking proteins that stably anchor actin filaments at their plus ends. It also anchors the N-terminus of the gigantic titin protein.

Titin - the gigantic spring protein of the sarcomere. It connects actin to myosin filaments, by spanning in mammals from the Z-disc (titin N-terminus) to the M-band (titin C-terminus). It rules the length of the sarcomere, thus is shortens during sarcomere contraction.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38

Figure legends.

Fig. 1. (A) The musculoskeletal system: a biomechanical apparatus. The muscle fibers attach via tendons to two bones. Muscle contraction generates force to move the two skeletal elements closer together and thus powers body movement. **(B) The sarcomere: a force producing machine.** Scheme of the basic architecture of a vertebrate sarcomere with myosin filaments (blue) and actin filaments (red), which are stably linked by titin spanning from the Z-disc (dark green) to the M-band (light blue). Note that the myosin containing region is also called the A-band, whereas the myosin free actin region of the sarcomere is called the I-band. The scheme was modified from ⁶⁸. **(C) Titin: a molecular ruler.** Titin spans half the length of a vertebrate sarcomere providing a direct template for sarcomere length: PEVK spring domain deletions result in shorter sarcomeres with shorter I-bands, whereas deletions of titin A-band parts lead to shorter myosin filaments and thus shorter A-bands. **(D) Titin: a molecular spring and possible force sensor.** The elasticity of titin is generated by its Ig domain series and PEVK rich region, which extend and possibly unfold in relaxed muscle to store elastic energy. Upon contraction the domains compact or refold again elastically.

39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig. 2. (A) Muscle-tendon attachment: a force resistant connection at molecular resolution. Top: transmission electronic micrograph of a *Drosophila* flight muscle tendon cell connecting one flight muscle myofibril to the chitin exoskeleton (cuticle). Note the zig-zag pattern of the interface between tendon and muscle. Scale bar is 500 nm. **Bottom:** molecular scheme of a terminal Z-disc connecting to a tendon cell. A terminal integrin signalling mechanically connects via an actin-rich layer to the terminal Z-disc of a flight muscle myofibril. Scheme was inspired by ^{94,103}. **(B) Tension sensors: measuring molecular forces at muscle attachments.** A talin FRET-based tension sensor allows to quantify molecular forces across talin. The sensor is composed of a mechanosensitive peptide inserted between FRET

1 donor (D) and acceptor (A). Low tension results in a closed sensor and high FRET, while high
2 force stretches the peptide and reduces FRET. Scheme was inspired by ¹⁰³ and modified from
3
4 102.
5
6
7
8
9


10 **Fig. 3. (A) *Drosophila* flight muscle morphogenesis.** Myotubes have attached to tendons at
11 24 h APF and contain a cortical actomyosin network under lower tension. At 30 h APF tension
12 has increased, myotubes have converted to myofibers and have assembled all their immature
13 myofibrils into myofibril bundles. High tension produced long tendon extensions. After 30 h
14 APF each myofibril and hence the entire myofiber grow in length and diameter resulting in a
15 large muscle volume increase. Tendon extensions shrink. **(B) Myofibrillogenesis: sarcomere**
16 **assembly and sarcomere maturation.** Schemes of actomyosin filament network without
17 periodic patterns which assembles to long immature myofibrils containing periodic muscle
18 myosin and titin. Each myofibril matures by isoform exchange and recruitment of more
19 sarcomeric components to a mature sarcomere. **(C) Transcriptional dynamics recapitulates**
20 **morphogenesis phases.** Complex transcriptional regulation patterns generate the necessary
21 proteins at precise stages and sufficient quantities to the various muscle morphogenesis phases.
22 Scheme was inspired by ¹²⁰.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 **Fig. 4. Muscle fibers: coordinating morphogenesis in a crowded space. Two examples of**
45 **different mitochondria-myofibril architectures. Left:** Adult *Drosophila* fibrillar flight
46 muscles contain round individual myofibrils (red) isolated from each other by large ellipsoid
47 shaped mitochondria (green). Nuclei (blue) are located between the myofibril bundles. Note
48 that myofibrils are squeezing mitochondria into their elongated shapes. **Right:** Mammalian
49 skeletal muscle contain bundles of vertically aligned cross-striated myofibrils with
50 mitochondria concentrated in groups squeezed in between. These mitochondria form long thin
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 extensions along the sarcomeric I-bands that build networks. In both muscle types the crowded
2 cellular environment generates mechanical pressure that positions the nuclei at the periphery
3
4 of myofibril bundles adopting an elongated shape.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65


A

the musculoskeletal system: a biomechanical apparatus




B

the sarcomere: a force producing machine



C

titin: a molecular ruler



D

titin: a molecular spring and possible force sensor






Figure 1

A muscle-tendon attachment: a force resistant connection at molecular resolution



B tension sensors: measuring molecular forces at muscle attachments






Figure 2

A *Drosophila* flight muscle morphogenesis:



B myofibrillogenesis: sarcomere assembly and sarcomere maturation



C transcriptional dynamics recapitulates morphogenesis phases




Figure 3

muscle fibers: coordinating morphogenesis in a crowded space




Figure 4

Nuno M Luis and Frank Schnorrer conceptualised the content, prepared the figures and drafted and revised the main text for this review together.