



HAL
open science

Uniform bound for the volume of the solutions to Liouville type equations on the annulus

Samy Skander Bahoura

► **To cite this version:**

Samy Skander Bahoura. Uniform bound for the volume of the solutions to Liouville type equations on the annulus. 2021. hal-03476580

HAL Id: hal-03476580

<https://hal.science/hal-03476580>

Preprint submitted on 13 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Uniform bound for the volume of the solutions to Liouville type equations on the annulus.

Samy Skander Bahoura*

Equipe d'Analyse Complexe et Géométrie.

Université Pierre et Marie Curie, 75005 Paris, France.

Abstract

We consider variational problems with regular Hölderian weight or with weight and boundary singularity and, Dirichlet condition. We prove the boundedness of the volume of the solutions to these equations on the annulus.

Keywords: Regular Hölderian weight, boundary singularity, a priori estimate, annulus, Lipschitz condition.

MSC: 35J60, 35B45.

1 Introduction and Main Results

We set $\Delta = \partial_{11} + \partial_{22}$ on the annulus $\Omega = C(1, 1/2, 0)$ of \mathbb{R}^2 of radii 1 and 1/2 centered at the origin.

We consider the following equation:

$$(P) \begin{cases} -\Delta u = (1 + |x - x_0|^{2\beta})Ve^u & \text{in } \Omega \subset \mathbb{R}^2, \\ u = 0 & \text{in } \partial\Omega. \end{cases}$$

Here: $C(1)$ the unit circle and $C(1/2)$ the circle of radius 1/2 centered at the origin.

$$\beta \in (0, 1/2), \quad x_0 \in C(1/2),$$

and,

$$u \in W_0^{1,1}(\Omega), \quad e^u \in L^1(\Omega) \text{ and } 0 < a \leq V \leq b.$$

This is an equation with regular Hölderian weight not Lipschitz in x_0 but have a weak derivative.

We have in [10] a nice formulation of this problem (P) in the sense of the distributions. This Problem arises in various geometrical and physical situations, see for example [1, 5, 23, 27]. The above equation was studied by many authors, with and without boundary conditions, also for Riemannian surfaces, see [1-27], where one can find some existence and compactness results. In [9] we have the following important Theorem,

*e-mails: samybahoura@yahoo.fr, samybahoura@gmail.com

Theorem A (Brezis-Merle [9]). For u and V two functions relative to (P) with,

$$0 < a \leq V \leq b < +\infty$$

then it holds, for all compact set K of Ω :

$$\sup_K u \leq c,$$

with c depending on a, b, β, x_0, K and Ω .

We deduce from Theorem A and from the elliptic estimates that, u is uniformly bounded in $C_{loc}^2(\Omega)$.

In this paper we try to prove that we have on all Ω the boundedness of the volume of the solutions of (P) if we add the assumption that V is uniformly Lipschitz with particular Lipschitz number.

Here we have:

Theorem 1.1 Assume that u is a solution of (P) relative to V with the following conditions:

$$x_0 \in C(1/2) \subset \partial\Omega, \quad \beta \in (0, 1/2),$$

and,

$$0 < a \leq V \leq b, \quad \|\nabla V\|_{L^\infty} \leq A = \frac{a}{2(1 + 2^{2\beta})},$$

we have,

$$\int_{\Omega} e^u \leq c(a, b, \beta, x_0, \Omega).$$

A consequence of this theorem is a compactness result of the solutions to this Liouville type equation, see [4].

We have the same result if we consider Liouville equation with boundary singularity on the annulus $\Omega = C(1, 1/2, 0)$ for particular Lipschitz number of the potential V :

We consider the following equation in the sense of distributions:

$$(P_\beta) \begin{cases} -\Delta u = |x - x_0|^{2\beta} V e^u & \text{in } \Omega \subset \mathbb{R}^2, \\ u = 0 & \text{in } \partial\Omega. \end{cases}$$

Here: $C(1)$ the unit circle and $C(1/2)$ the circle of radius 1/2 centered at the origin.

$$\beta \in (-1/2, +\infty), \quad x_0 \in C(1/2),$$

and,

$$u \in W_0^{1,1}(\Omega), \quad |x - x_0|^{2\beta} e^u \in L^1(\Omega) \text{ and } 0 < a \leq V \leq b.$$

This is an equation with boundary singularity and Dirichlet condition on the annulus.

Here we have:

Theorem 1.2 Assume that u is a solution of (P_β) relative to V with the following conditions:

$$x_0 \in C(1/2) \subset \partial\Omega, \quad \beta \in (-1/2, +\infty),$$

and,

$$0 < a \leq V \leq b, \quad \|\nabla V\|_{L^\infty} \leq A = \frac{(\beta + 1)a}{2},$$

we have,

$$\int_{\Omega} |x - x_0|^{2\beta} e^u \leq c(a, b, \beta, x_0, \Omega).$$

For the previous theorem, the condition $\beta > -1/2$ is in relation with the regularity of the solutions to apply the Pohozaev identity.

A consequence of the previous theorem is a compactness result for the solutions to a Liouville type equation with boundary singularity on the annulus, see [3].

2 Proof of the Theorems:

Proof of the theorems:

By corollary 1 of the paper of Brezis-Merle, we have: $e^{ku} \in L^1(\Omega)$ for all $k > 2$ and the elliptic estimates and the Sobolev embedding imply that: $u \in W^{2,k}(\Omega) \cap C^{1,\epsilon}(\bar{\Omega})$, $\epsilon > 0$. By the maximum principle $u \geq 0$.

Step 1: By using the first eigenvalue and the first eigenfunction, with Dirichlet boundary condition, the volume is locally uniformly bounded, and thus the solutions are locally uniformly bounded by Brezis-Merle result. The solutions $u > 0$ are locally uniformly bounded in $C^{1,\epsilon}(\Omega)$ for ϵ small.

Step 2: Let's consider $C_1 = C(1, 3/4, 0)$ and $C_2 = C(3/4, 1/2, 0)$ the two annulus wich are the neighborhood of the two components of the boundary.

We multiply the equation by $(x - x_0) \cdot \nabla u$ on C_1 and C_2 and use the Pohozaev identity. We use the fact that u is uniformly bounded around the circle $C(3/4)$. We obtain:

1) We have on C_1 :

$$\int_{C_1} (\Delta u)[(x - x_0) \cdot \nabla u] dx = \int_{C_1} -[(1 + |x - x_0|^{2\beta})V(x - x_0) \cdot \nabla(e^u)] dx,$$

Thus,

$$\begin{aligned} & \int_{\partial C_1} [(x - x_0) \cdot \nabla u](\nabla u \cdot \nu) - \frac{1}{2}[(x - x_0) \cdot \nu]|\nabla u|^2 = \\ & = \int_{C_1} (2 + 2(\beta + 1)|x - x_0|^{2\beta})V e^u dx + \int_{C_1} (1 + |x - x_0|^{2\beta})(x - x_0) \cdot \nabla V e^u dx + \\ & \quad - \int_{\partial C_1} (1 + |x - x_0|^{2\beta})[(x - x_0) \cdot \nu]V e^u d\sigma \end{aligned}$$

We can write, ($u = 0$ on $C(1)$):

$$\begin{aligned} & \int_{C(1)} \frac{1}{2} [(x - x_0) \cdot \nu] (\partial_\nu u)^2 d\sigma + O(1) = \\ & = \int_{C_1} (2 + 2(\beta + 1)|x - x_0|^{2\beta}) V e^u dx + \int_{C_1} (1 + |x - x_0|^{2\beta})(x - x_0) \cdot \nabla V e^u dx + O(1) = \\ & = C \int_{C(1)} \partial_\nu u d\sigma + O(1), \end{aligned}$$

with $C > 0$ not depends on u .

Be cause $\nu = x, \|x\| = 1, \|x_0\| = 1/2$ and then by Cauchy-Schwarz, $(x - x_0) \cdot x = \|x\|^2 - x_0 \cdot x \geq 1/2$, we obtain:

$$\int_{C(1)} (\partial_\nu u)^2 d\sigma \leq 4C \int_{C(1)} \partial_\nu u d\sigma + O(1),$$

and, by the Cauchy-Schwarz inequality applied to the integral of the right hand side we obtain:

$$\int_{C(1)} (\partial_\nu u)^2 d\sigma = O(1), \text{ and } \int_{C(1)} (\partial_\nu u) d\sigma = O(1).$$

2) We have on C_2 , we use again, the uniform boundedness of u in C^1 norm around $C(3/4)$:

$$\int_{C_2} (\Delta u)[(x - x_0) \cdot \nabla u] dx = \int_{C_2} -[(1 + |x - x_0|^{2\beta})V(x - x_0) \cdot \nabla(e^u)] dx,$$

Thus,

$$\begin{aligned} & \int_{\partial C_2} [(x - x_0) \cdot \nabla u](\nabla u \cdot \nu) - \frac{1}{2} [(x - x_0) \cdot \nu] |\nabla u|^2 = \\ & = \int_{C_2} (2 + 2(\beta + 1)|x - x_0|^{2\beta}) V e^u dx + \int_{C_2} (1 + |x - x_0|^{2\beta})(x - x_0) \cdot \nabla V e^u dx + \\ & \quad - \int_{\partial C_2} (1 + |x - x_0|^{2\beta}) [(x - x_0) \cdot \nu] V e^u d\sigma \end{aligned}$$

We can write, ($u = 0$ on $C(1/2)$):

$$\begin{aligned} & \int_{C(1/2)} \frac{1}{2} [(x - x_0) \cdot \nu] (\partial_\nu u)^2 d\sigma + O(1) = \\ & = \int_{C_2} (2 + 2(\beta + 1)|x - x_0|^{2\beta}) V e^u dx + \int_{C_2} (1 + |x - x_0|^{2\beta})(x - x_0) \cdot \nabla V e^u dx + O(1), \end{aligned}$$

But here, $\nu = -2x$ and $(x - x_0) \cdot \nu = -(x - x_0) \cdot x \leq 0$ and thus:

$$\begin{aligned} & \int_{C_2} (2 + 2(\beta + 1)|x - x_0|^{2\beta})Ve^u dx + \int_{C_2} (1 + |x - x_0|^{2\beta})(x - x_0) \cdot \nabla Ve^u dx + \\ & + \int_{C(1/2)} \frac{1}{2}[-(x - x_0) \cdot \nu](\partial_\nu u)^2 d\sigma = O(1), \end{aligned}$$

If we choose:

$$\frac{|(x - x_0) \cdot \nabla V|}{V} \leq \frac{1}{2} \inf_{x \in \Omega} \frac{(2 + 2(\beta + 1)|x - x_0|^{2\beta})}{1 + |x - x_0|^{2\beta}}, \quad (1)$$

We obtain:

$$\begin{aligned} & \int_{C_2} (2 + 2(\beta + 1)|x - x_0|^{2\beta})Ve^u dx + \int_{C_2} (1 + |x - x_0|^{2\beta})(x - x_0) \cdot \nabla Ve^u dx \geq \\ & \geq \frac{1}{2} \int_{C_2} (2 + 2(\beta + 1)|x - x_0|^{2\beta})Ve^u dx \geq 0 \end{aligned}$$

thus,

$$\int_{C_2} (2 + 2(\beta + 1)|x - x_0|^{2\beta})Ve^u dx = O(1),$$

(This condition is true for the second theorem for $x \in \Omega$, we have: $\frac{|(x - x_0) \cdot \nabla V|}{V} \leq \frac{1}{2} \inf_{x \in \Omega} \frac{2(\beta + 1)|x - x_0|^{2\beta}}{|x - x_0|^{2\beta}} = \beta + 1$, this is true if $A \leq \frac{a(\beta + 1)}{2}$.)
and thus,

$$\int_{C(1/2)} \partial_\nu u d\sigma = O(1),$$

Thus, if: $2 \frac{\text{diam}(\Omega) \|\nabla V\|_\infty}{a} \leq \frac{2}{1 + [\text{diam}(\Omega)]^{2\beta}} \Rightarrow$ if $2A \leq \frac{2a}{2(1 + [\text{diam}(\Omega)]^{2\beta})}$,

we obtain a uniform bound for $\int_{C(1/2)} \partial_\nu u d\sigma$. But $\text{diam}(\Omega) = \text{diameter}(\Omega) = 2$, because it is the annulus.

If we use 1) and 2), we obtain: if $A \leq \frac{a}{2(1 + 2^{2\beta})}$ for the first theorem, then:

$$\int_{C(1)} (\partial_\nu u) d\sigma = O(1), \text{ and } \int_{C(1/2)} \partial_\nu u d\sigma = O(1),$$

and, thus:

$$\int_{\Omega} (1 + |x - x_0|^{2\beta})Ve^u dx = \int_{\partial\Omega} \partial_\nu u d\sigma = O(1).$$

We have the same conclusion for the second theorem if $A = \frac{a(\beta + 1)}{2}$.

References

- [1] T. Aubin. Some Nonlinear Problems in Riemannian Geometry. Springer-Verlag, 1998.
- [2] Ambrosio. L, Fusco. N, Pallara, D. Functions of Bounded variations and Free discontinuity Problems, Oxford Press. 2000.
- [3] Bahoura.S.S. A uniform boundedness result for solutions to the Liouville type equation with boundary singularity. J. Math. Sci. Univ. Tokyo, 23, no 2, 487-497. 2016.
- [4] Bahoura.S.S. A compactness result for an equation with Holderian condition. Commun. Math. Anal. Vol 21, no 1, 23-34, 2018.
- [5] C. Bandle. Isoperimetric Inequalities and Applications. Pitman, 1980.
- [6] L. Boccardo, T. Gallouet. Nonlinear elliptic and parabolic equations involving measure data. J. Funct. Anal. 87 no 1, (1989), 149-169.
- [7] H. Brezis, YY. Li and I. Shafrir. A sup+inf inequality for some nonlinear elliptic equations involving exponential nonlinearities. J.Funct.Anal.115 (1993) 344-358.
- [8] Brezis. H, Marcus. M, Ponce. A. C. Nonlinear elliptic equations with measures revisited. Mathematical aspects of nonlinear dispersive equations, 55-109, Ann. of Math. Stud., 163, Princeton Univ. Press, Princeton, NJ, 2007.
- [9] H. Brezis, F. Merle. Uniform estimates and Blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimension. Commun. in Partial Differential Equations, 16 (8 and 9), 1223-1253(1991).
- [10] H. Brezis, W. A. Strauss. Semi-linear second-order elliptic equations in L1. J. Math. Soc. Japan 25 (1973), 565-590.
- [11] Chang, Sun-Yung A, Gursky, Matthew J, Yang, Paul C. Scalar curvature equation on 2- and 3-spheres. Calc. Var. Partial Differential Equations 1 (1993), no. 2, 205-229.
- [12] W. Chen, C. Li. A priori estimates for solutions to nonlinear elliptic equations. Arch. Rational. Mech. Anal. 122 (1993) 145-157.
- [13] C-C. Chen, C-S. Lin. A sharp sup+inf inequality for a nonlinear elliptic equation in \mathbb{R}^2 . Commun. Anal. Geom. 6, No.1, 1-19 (1998).
- [14] D.G. De Figueiredo, P.L. Lions, R.D. Nussbaum, A priori Estimates and Existence of Positive Solutions of Semilinear Elliptic Equations, J. Math. Pures et Appl., vol 61, 1982, pp.41-63.
- [15] Droniou. J. Quelques resultats sur les espaces de Sobolev. Hal 2001.
- [16] Ding.W, Jost. J, Li. J, Wang. G. The differential equation $\Delta u = 8\pi - 8\pi h e^u$ on a compact Riemann surface. Asian J. Math. 1 (1997), no. 2, 230-248.
- [17] D. Gilbarg, N. S, Trudinger. Elliptic Partial Differential Equations of Second order, Berlin Springer-Verlag.

- [18] Hofmann, S. Mitrea, M. Taylor, M. Geometric and transformational properties of Lipschitz domains, Semmes-Kenig-Toro domains, and other classes of finite perimeter domains. *J. Geom. Anal.* 17 (2007), no. 4, 593-647.
- [19] Krantz, S. Geometric functions theory. Birkhauser.
- [20] YY. Li. Harnack Type Inequality: the method of moving planes. *Commun. Math. Phys.* 200,421-444 (1999).
- [21] YY. Li, I. Shafrir. Blow-up analysis for solutions of $-\Delta u = Ve^u$ in dimension two. *Indiana. Math. J.* Vol 3, no 4. (1994). 1255-1270.
- [22] L. Ma, J-C. Wei. Convergence for a Liouville equation. *Comment. Math. Helv.* 76 (2001) 506-514.
- [23] Nagasaki, K, Suzuki,T. Asymptotic analysis for two-dimensional elliptic eigenvalue problems with exponentially dominated nonlinearities. *Asymptotic Anal.* 3 (1990), no. 2, 173-188.
- [24] Necas, J. Direct Methods in the Theory of Elliptic Equations. Springer.
- [25] I. Shafrir. A sup+inf inequality for the equation $-\Delta u = Ve^u$. *C. R. Acad.Sci. Paris Sér. I Math.* 315 (1992), no. 2, 159-164.
- [26] Stoker, J. Differential Geometry.
- [27] Tarantello, G. Multiple condensate solutions for the Chern-Simons-Higgs theory. *J. Math. Phys.* 37 (1996), no. 8, 3769-3796.