

Snails in depth: Integrative taxonomy of Famelica, Glaciotomella and Rimosodaphnella (Conoidea: Raphitomidae) from the deep sea of temperate Australia

Francesco Criscione, Anders Hallan, Nicolas Puillandre, Alexander Fedosov

To cite this version:

Francesco Criscione, Anders Hallan, Nicolas Puillandre, Alexander Fedosov. Snails in depth: Integrative taxonomy of Famelica, Glaciotomella and Rimosodaphnella (Conoidea: Raphitomidae) from the deep sea of temperate Australia. Invertebrate Systematics, 2021, 35 (8), pp.940-962. $10.1071/I$ S21008. hal-03477619

HAL Id: hal-03477619 <https://hal.science/hal-03477619v1>

Submitted on 13 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

[Distributed under a Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

Invertebrate Systematics

Snails in depth: Integrative taxonomy of Famelica, Glaciotomella and Rimosodaphnella (Conoidea: Raphitomidae) from the deep sea of temperate Australia

650x400mm (96 x 96 DPI)

Short summary

Normalist Contraction Distribution patterns of deep-sea benthic gastropods remain poorly known, yet such knowledge is crucial to approaching their systematics and understanding their diversity. With focus on the raphitomid genera *Famelica*, *Glaciotomella* and *Rimosodaphnella*, we combine genetic, shell, anatomy and distribution data to identify four new species from Australia and beyond, which we here describe. We identify the set of morphological characters shared by congeneric species and highlight patterns of rarity and endemicity in some species. Our findings suggest that radula and protoconch features are good proxies for genus-level phylogenetic relationships. We discuss the significance of ecological factors (such as choice and availability of preys) as potential drivers in the evolution of this hyperdiverse group of gastropods.

http://www.publish.csiro.au/journals/is

Abstract

Mexical Solution Networkships and Mexican Solution Networkships and Mexical Solution Network and Alam Unjected a comprehensive mitochondrial DNA

Do Nara raphitomids to the species delimitation met

Ind 15 primary species The deep sea of temperate south-eastern Australia appears to be a 'hotspot' for diversity and endemism of conoidean neogastropods of the family Raphitomidae. Following a series of expeditions in the region, a considerable amount of relevant DNA-suitable material has become available. A molecular phylogeny based on this material has facilitated the identification of diagnostic morphological characters, allowing the circumscription of monophyletic genera and the introduction of a number of new genus-level taxa. Both named and new genera are presently being investigated through integrative taxonomy, with the discovery of a significant amount of undescribed species. As part of this ongoing investigation, this study focuses on the genera *Famelica* Bouchet & Warén, 1980, *Glaciotomella* Criscione, Hallan, Fedosov & Puillandre, 2020 and *Rimosodaphnella* Cossmann, 1914. We subjected a comprehensive mitochondrial DNA dataset of representative deep-sea raphitomids to the species delimitation methods ABGD and ASAP, which recognised 18 and 15 primary species hypotheses (PSHs) respectively. Following additional evaluation of shell and radular features, as well as examination of geographic and bathymetric ranges, nine of these PSHs were converted to secondary species hypotheses (SSHs). Four SSHs (two in *Famelica* and two in *Rimosodaphnella*) were recognised as new to science and their formal descriptions are provided herein.

Introduction

groups below family level are primarily based
35; Sysoev and Kantor 1986, 1987; Sysoev 198
97; Sysoev 1997; Bouchet and Sysoev 2001; S
fitto 2006; Kantor *et al.* 2012; Stahlschmidt ar
rassi and Bonfitto 2015; Kantor *et* Turriform Conoidea is the most diverse group of marine molluscs (Horton *et al.* 2019), with considerable species diversity occurring in the deep sea, where it is often coupled with low abundance (Sysoev 1997; Kantor *et al.* 2008; Bouchet *et al.* 2009). One of the total fifteen families of the group, Raphitomidae Bellardi, 1875, is among the most poorly studied (Bouchet *et al.* 2011), notably in the deep sea (Criscione *et al.* 2021). Because of their high diversity, low abundance and morphological complexity combined, the systematics of many deep-sea genera and species remains tentative, with a large proportion of taxa awaiting description (Bouchet *et al.* 2009). To further complicate matters, most of the taxonomic accounts available for groups below family level are primarily based on shell morphology (Sysoev and Ivanov 1985; Sysoev and Kantor 1986, 1987; Sysoev 1988, 1990, 1996a, b; Bouchet and Sysoev 1997; Sysoev 1997; Bouchet and Sysoev 2001; Sysoev and Bouchet 2001; Morassi and Bonfitto 2006; Kantor *et al.* 2012; Stahlschmidt and Chino 2012; Bonfitto and Morassi 2013; Morassi and Bonfitto 2015; Kantor *et al.* 2016) and only three studies do include molecular data (Fassio *et al.* 2019; Russini *et al.* 2020; Criscione *et al.* 2021). This is partly due to a lack of study material, particularly that suitable for molecular analysis (Criscione *et al.* 2021). Due to widespread shell homoplasy, some of the earlier studies have incorrectly attributed a considerable number of unrelated species to very few raphitomid genera, colloquially termed 'dumpsters' (such as *Pleurotomella* Verril, 1872 and *Gymnobela* Verrill, 1884) (Criscione *et al.* 2021). Based on their support as clades and on diagnostic morphological characters, Criscione *et al.* (2021) demonstrated the non-monophyly of such genus-level groups and, in constraining them*,* introduced a number of new genus-level taxa and described their type species. The same study also revealed a multitude of putatively undescribed species among deep-sea raphitomids of a temperate region roughly corresponding to the central and eastern part of the South Australia realm (#26 of Costello *et al.* 2017) and the southern part of the Tropical Australia and Coral Sea realm (#16 of Costello *et al.* 2017) (Fig. 1). With nearly half of the currently accepted deep-sea genera (24 out of 50) and nearly a third (7) endemic, this area is a potential 'hot spot' for raphitomid diversity (Criscione *et al.* 2021). Thanks to recent sampling efforts in the area (MacIntosh *et al.* 2018; Williams 2018; O'Hara *et al.* 2020), relatively abundant raphitomid samples, suitable for genetic analysis, are available. This material has allowed studies whose methodological design for species delimitation transcends the mere conchological approach of earlier taxonomical accounts and capitalises on the combination of morpho-anatomical and molecular data (i.e. 'integrative taxonomy ', Dayrat 2005; Will *et al.* 2005). As a result of these studies on the Australian Raphitomidae, new species have been described for the genera *Gladiobela* Criscione, Hallan, Puillandre & Fedosov, 2020 (5 species) and *Pagodibela* Criscione, Hallan, Puillandre & Fedosov, 2020 (2) (Hallan *et al.* 2021) and of *Austrobela* Criscione, Hallan, Puillandre & Fedosov, 2020 (4), *Austrotheta* Criscione, Hallan, Puillandre & Fedosov 2020 (1), *Spergo* Dall, 1895 (4) and *Theta* Clarke, 1959 (2) (Criscione et al., in press). Based on a larger sampling size, including specimens from beyond Australian waters, this study aims to revisit species delimitation and taxonomy in three genera: *Famelica* Bouchet & Warén, 1980, *Glaciotomella* Criscione, Hallan, Puillandre & Fedosov 2020 and *Rimosodaphnella* Cossmann, 1916. In the analysis of Criscione *et al.* (2021), these genera (along with *Veprecula* Melvill, 1917) constituted a monophyletic group (clade 'A' of fig. 2 in Criscione *et al.* 2021). Formal descriptions are here presented for newly recognised species. Furthermore, revised genus diagnoses and new anatomical and morphological data are introduced for both established and new taxa, which are discussed in terms of their diagnostic utility at the genus level. Finally, geographic and bathymetric distributions are presented and discussed for the taxa treated herein.

> http://www.publish.csiro.au/journals/is

PHS CAR

Materials and methods

Taxon sampling

e tropical and temperate Indian and Pacific Oc

I part of the Tropical Deep-sea Benthos program

Eig. 1, Table S1).

Systematic research on the Conoidea at the AN

published) sequences of the mitochondrial geobtained (see The samples studied herein were selected among all Raphitomidae ethanol-preserved material from the malacological collections of the Australian Museum, Sydney (AMS), the South Australian Museum, Adelaide (SAMA) and the Muséum national d'Histoire naturelle, Paris (MNHN). All studied material has been collected off Australia during the expeditions 106 IN2015 C01, IN2015 C02 (GAB) and IN2017 V03 (Tasman and Coral Seas), targeting (among other sites) a number of Commonwealth Marine Reserves (CMR), and other localities (mainly of the tropical and temperate Indian and Pacific Oceans), during a number of voyages that formed part of the Tropical Deep-sea Benthos programme of MNHN (expeditions.mnhn.fr; Fig. 1, Table S1). As a result of ongoing systematic research on the Conoidea at the AMS and MNHN, several hundreds of (mostly unpublished) sequences of the mitochondrial gene cytochrome oxidase

subunit 1 (*cox1*) were obtained (see methodology below) from a considerable number of

largely undescribed raphitomid taxa. In order to assist with the selection of the study

material, a pilot analysis was performed on a dataset including all raphitomid *cox1*

sequences, using the neighbour-joining method (NJ) (Saitou and Nei 1987) implemented in

MEGA 7 (Kumar *et al.* 2016). In particular, the dataset included *cox1* sequences of the

holotypes for the type species of several deep-sea raphitomid genera, including

Glaciotomella investigator Criscione, Hallan, Puillandre and Fedosov *et al.* 2020 (GenBank

AN: MN983178). The dataset also encompassed *cox1* sequences of well-recognisable

species, such as *Famelica tasmanica* Sysoev & Kantor, 1987, as well as of several Australian

undescribed species included in the phylogeny of Criscione *et al.* (2021), such as those

placed in the genus *Rimosodaphnella* therein. All *cox1* sequences of *Glaciotomella, Famelica*

or *Rimosodaphnella* used in (Criscione *et al.* 2021) and all sequences that (in the results of

the NJ analysis; Fig. S1) were more closely related to the species of these three genera than

to any other raphitomid genus, were selected as ingroup and used in the molecular analysis

described below (Table S1). Sequences representing 13 raphitomid species from 13 genera

were selected as outgroups (Table S1). Their choice was based on the phylogeny of Criscione

et al. (2021), containing many southern and south-eastern Australian Raphitomidae .

 Among the ingroup specimens, morphological examination was only conducted on those collected in Australian waters. However, for samples outside Australia, examination of shell photographs was possible and was utilised when necessary and appropriate. Geographic and bathymetric data were available for all ingroup specimens. Geographic distributions were assessed with reference to marine biogeographic realms as delimited in Costello *et al.* (2017). According to Bouchet *et al.* (2008), when inferring bathymetric distributions of species from sampling depth intervals, only shallower depth values were considered, as there is no evidence that the species collected occurs beyond that value.

-
- Molecular methods

erformed in laboratories at two different institutive stated, the same methodology was follow
m small pieces of foot muscle by use of a Bioli
animal tissue, following the standard procedures
5075 robot (Eppendorf), follow Molecular work was performed in laboratories at two different institutions (AMS and MNHN). Unless otherwise stated, the same methodology was followed by both laboratories. DNA was extracted from small pieces of foot muscle by use of a Bioline Isolate II Genomic DNA extraction kit for animal tissue, following the standard procedure of the manual (AMS) or using the Epmotion 5075 robot (Eppendorf), following the recommendations by the manufacturer (MNHN). A fragment of *cox1* was amplified using the primer pairs LCO1490/HCO2198 for *cox1* (Folmer *et al.* 1994). PCR reactions were performed in volumes of 25 μl, containing 3 ng DNA, 1X Qiagen CoralLoad PCR Buffer, 2.5mM MgCl2, 0.25mM dNTP, 0.5mM of each primer, 0.5 μg/μl of BSA and 0.2 μl of Bioline MyTaq DNA polymerase. Amplification consisted of an initial denaturation step at 94°C for 4 min, followed by 37 cycles of denaturation at 94°C for 30 s, annealing at 50°C for 30 s, followed by extension at 151 72°C for 1 min. The final extension was at 72°C for 5 min. PCR products were purified and sequenced by the Macrogen (AMS) and Eurofins (MNHN) sequencing facilities. When necessary, chromatograms were manually corrected for misreads and forward and reverse strands were merged into one sequence file using CodonCode Aligner v. 9.0.1 (CodonCode Corporation, Dedham, MA). Sequence alignments were generated using MUSCLE as implemented in MEGA7 (Kumar *et al.* 2016). Sequences were deposited in GenBank (Table S1). Phylogenetic trees were generated using Maximum Likelihood (ML) and Bayesian inference (BI) methods. ML was performed using the program IQ-Tree v. 2.1.1 (Minh *et al.* 2020), including the implemented model-finder function (Kalyaanamoorthy *et al.* 2017) and ultrafast bootstrapping (BS, 1000 replicates) (Hoang *et al.* 2017). The BI analysis was 161 performed in MrBayes 3.2.6 (Ronquist and Huelsenbeck 2003) and included 2 runs of $10⁷$

Invertebrate Systematics

 generations, with 4 chains each and a sampling frequency of one tree per 1000 generations. Other parameters were set to default. After checking for convergence (ESS>200) with Tracer (Rambaut *et al.* 2018), a consensus tree was calculated after discarding the first 25% trees as burn-in. According to MrBayes manual (p. 94), a priori model testing was not performed, and the GTR+G+I model was applied to the BI analysis. Nodal support was assessed as Bayesian posterior clade probabilities (BPP). Pairwise genetic distances were calculated for *cox1* sequences using the Kimura-2 parameter model (K2p, Kimura 1980) as implemented in MEGA7 with the option 'pair-wise deletion of gaps'.

Morphological examinations

ations

sisted of animals and shells, from which they

logy described in Criscione *et al.* (2021). We s

ternal anatomy, including radular morphology

d to plasticine and positioned with their vertic

h shell was then pho All studied samples consisted of animals and shells, from which they had been extracted following the methodology described in Criscione *et al.* (2021). We studied shell morphology and (when possible) internal anatomy, including radular morphology. Shells of sequenced specimens were affixed to plasticine and positioned with their vertical axis parallel to the observation plane. Each shell was then photographed from above using a digital SLR camera. Maximum shell length (SL) and width (SW) were measured on digitised images using the calibrated ruler tool in Adobe Photoshop CC v.20.0.6. Measurements were rounded to the nearest 0.1 mm. The number of shell whorls was counted under a Leica MZ8 stereomicroscope, according to Bouchet and Kantor (2004). While it was possible to count the number of teleoconch whorls for almost all studied specimens, protoconch whorls could only be counted occasionally due to widespread erosion of the apex. Anatomical studies were conducted on animals removed from ethanol and briefly rehydrated in distilled water. Using standard dissection tools, the venom apparatus was excised and the radular sac isolated and placed on a glass slide; during this dissection process, head-foot, mantle, genital and (non-radula) foregut characters were examined where possible. After dissolution in dilute commercial bleach, clusters of hypodermic teeth were rinsed repeatedly in distilled water, then separated into individuals and ligament-connected pairs/smaller clusters. Subsequently, the glass stub was affixed to a carbon adhesive placed on a 12 mm diameter aluminium mount. All samples were imaged at Macquarie University, Sydney, using a 191 Phenom XL Scanning Electron Microscope.

Species delimitation

Firstly, the Automatic Barcode Gap Discovery (ABGD, Puillandre *et al.* 2012a;

https://bioinfo.mnhn.fr/abi/public/abgd/abgdweb.html) was used with a p-distance model,

the relative gap width (X) set to 1 and the other parameters left to default.

Secondly, the Assemble Species by Automatic Partitioning (ASAP, Puillandre *et al.* 2020;

https://bioinfo.mnhn.fr/abi/public/asap/) was applied. Similar to ABGD, ASAP proposes

species partitions based on genetic distances only; however, ASAP screens all the genetic

distances from the lowest to the highest and merges sequences into "groups" that are

successively further merged until all sequences form a single group. At each merging step,

the partition is evaluated and given a score. At the end of the analyses, the 10 partitions

with the lowest scores are provided (the lower the ASAP-score, the better the partition).

Resulting ABGD and ASAP groups were considered primary species hypotheses, henceforth

referred to as PSHs. Following Puillandre *et al.* (2012b), conversion of PSHs to secondary

species hypotheses (SSHs) was conducted through comparative examination of

are provided (the lower the ASAP-score, the b
AP groups were considered primary species h
llowing Puillandre *et al.* (2012b), conversion o
Hs) was conducted through comparative exan
ers as well as through evaluation of ge morphological characters as well as through evaluation of geographic and bathymetric data

and phylogenetic reconstruction based on the DNA sequences.

209 In particular, when converting an individual PSH to SSH, the occurrence of the following

conditions was assessed: (i) the PSH is a highly supported clade (BPP>0.98 and BS>90%), (ii)

211 all its constituent specimens share at least one distinctive morphological feature deemed

212 not to be polymorphic or ecophenotypic and without exhibiting intermediate forms, (iii) the

PSH maintains genetic or morphological divergence and/or bathymetric partitioning when

occurring in sympatry with another PSH. When available, species names were assigned to

215 SSHs based on the current taxonomy. New species names were introduced when no names

were available. Formal descriptions for these taxa are given in the systematics section

below.

Results

- e (F2) included samples from Australian seas a

FA, GA, R3-R5, RA-RE) encompassed sequence

1 (i.e. with the lowest ASAP-score) returned 1!

1 resulting from ABGD and the remaining three

1+G2 and R4+R5. The second-best AS Molecular studies Molecular analyses were based on a total of 46 *cox1* sequences (28 newly produced and 18 GenBank-sourced). Of the total sequences employed, 33 constituted the ingroup and the remaining 13 were used as outgroups. In the vicinity of the barcode gap (*i.e.* between 0.8 and 3.3%, the highest intra-PSH genetic distance and the lowest inter-PSH genetic, respectively), the ABGD analysis of the *cox1* ingroup dataset consistently returned a primary partition with eighteen groups of sequences (PSHs). Among all PSHs, six (F1, F3, G1, G2, R1, R2) contained exclusively Australian samples, one (F2) included samples from Australian seas and beyond, while the remaining eleven (F4, FA, GA, R3-R5, RA-RE) encompassed sequences from outside Australian waters. The best ASAP partition (i.e. with the lowest ASAP-score) returned 15 PSHs, 12 of which were identical to those resulting from ABGD and the remaining three corresponding to the combinations F1+F3, G1+G2 and R4+R5. The second-best ASAP partition included 14 PSHs,
- with RD and RE grouped (when compared to the 15-species ASAP partition). The ASAP-score of the alternative partitions were much higher and are thus ignored.
- The BI and ML analyses generated trees (Fig. 2, S2) with similar topologies and no supported
- incongruences. Clades representing PSH-level relationships and above were generally well-
- 237 supported, with very few exceptions. In both analyses, three major genus-level clades were
- retrieved, namely *Famelica* (BPP=1, BS=95%), *Glaciotomella* (BPP=1 and BS=95%) and
- *Rimosodaphnella* (BPP=0.98 and BS=94%). These three clades included five, three and ten
- ABGD PSHs respectively, all forming highly supported (in terms of nodal support) and well-
- 241 differentiated (in terms of branch lengths) clades. The only exception was R1, which
- received no support from the BI analysis and moderate support from the ML analysis
- (BS=95%). ASAP PSHs R4+R5 and G1+G2 were highly supported by both analyses (Fig. 2, S2),
- while F1+F3 received no support.
- Within the genus-level clades, there was no overlap between intra- and inter-ABGD-PSH
- distances. In the *Famelica* clade, the intra-PSH pairwise distance in *cox1* measured for the
- only PSH with more than one sample (F2) was 0.8%, with inter-PSH distances ranging from
- 5.2 to 18.2% (average=14.5%) (Table 1). The lowest inter-PSH distances were observed
- between F1 and F3 (the two ABGD-PSH grouped by ASAP). No intra-ABGD-PSH pairwise

 distances could be measured for the *Glaciotomella* clade and the inter-PSH distances ranged from 3.3 to 16.0 % (average=9.7 %), with the lowest value recorded between G1 and G2 (the two ABGD-PSH grouped by ASAP). In the *Rimosodaphnella* clade, the intra-PSH pairwise distances in *cox1* ranged from 0 to 0.3 % (average=0.2 %) with inter-PSH distances ranging from 4.6 to 15.6 % (average=11.7 %) (Table 2). The lowest inter-PSH distances were observed between R4 and R5 (the two ABGD-PSH grouped by ASAP) and the highest intra- PSH distances were found within R2. The genetic distance between RD and RE, the two ABGD-PSH grouped in the second-best ASAP partition, is 6.7%.

Morphological studies

examined for all sequenced material and inter
then possible) for at least two sequenced spec
omical observations refer to PSHs examined h
with a letter suffix (i.e., FA, GA, etc.). Protocor
ded and could therefore not be s Shell morphology was examined for all sequenced material and internal anatomy, including radular morphology (when possible) for at least two sequenced specimens per PSH. Detailed morpho-anatomical observations refer to PSHs examined herein, and do not 263 include PSHs assigned with a letter suffix (i.e., FA, GA, etc.). Protoconchs of many specimens were considerably eroded and could therefore not be studied. For the remaining specimens, some sculptural detail could be inferred from heavily eroded protoconchs by careful examination using a microscope. However, owing to their very poor quality, these protoconchs are not figured herein. When observed, protoconch sculpture was highly conserved within each genus-level clade and could not be used to differentiate congeneric PSHs. While *Rimosodaphnella* PSHs possessed a protoconch with a typical raphitomid diagonally cancellate sculpture (Fig. 3C), *Famelica* and *Glaciotomella* PSHs exhibited a protoconch with median keel as the only spiral element (Fig. 3A-B). In these two genera, axial elements ('pillars' in Bouchet *et al.* 2011, p. 283) were present, that were restricted to the area below the keel in *Famelica* (Fig. 3A) and extended also above the keel in *Glaciotomella* (Fig. 3B). All PSHs in the *Famelica* clade exhibited a gross shell morphology and whorl profile that were comparatively similar (Figs 4-5), with only one (F4) differing in its sculptural elements (prominent spiral keels). The radula was absent in all *Famelica* specimens studied. Very similar shells were exhibited by PSHs in the *Glaciotomella* clade (Fig. 6), with shells of G1 and G2 being remarkably larger than GA and sharing the same sculptural pattern. While the radula of GA was not studied, G1 and G2 shared nearly identical hypodermic teeth (Fig. 6).

 Rimosodaphnella PSHs varied notably in shell morphology (Figs 7-8), from comparatively large and broad (R2, Fig. 7D-F), to large and elongate (R1, Fig. 7A-C), to very small (*e.g.* R4 and R5, Fig. 8B-C). Generally, axial sculpture was more prominent than spiral sculpture in all shells of all PSHs, with two exceptions: RA, where axials were absent (Fig. 5C) and R2, where spiral and axial elements were equally prominent (Fig. 7G). The radula of both R1 and R2 (Fig. 10) consisted of similar straight, unbarbed hypodermic teeth that in R2 were shorter and broader.

Geographic and bathymetric distributions

The *Famelica* clade is recorded from at least four marine realms (sensu Costello *et al.* 2017)

(Fig. 1A), the *Glaciotomella* clade from three (Fig. 1A) and the *Rimosodaphnella* clade from

at least six marine realms (Fig. 1B). *Famelica* and *Rimosodaphnella* both exhibit a wide

ecorded from at least four marine realms (sen

nella clade from three (Fig. 1A) and the *Rimose*

ms (Fig. 1B). *Famelica* and *Rimosodaphnella* b

pectively 398-4144 m and 350-4750 m), while

74 m) is narrower (Fig. 11). bathymetric range (respectively 398-4144 m and 350-4750 m), while the range of

Glaciotomella (600-2474 m) is narrower (Fig. 11). In *Famelica*, the sister PSHs F1 and F3 are

known only from off Australia, where they are found respectively at bathyal and abyssal

depths of the east coast (Fig. 1A). F2 is known from two widely separated localities, both

geographically (Fig. 1A) and bathymetrically (Fig. 11), namely in the temperate GAB (3389

m) and off the tropical New Caledonia (815 m). The only samples of the remaining *Famelica*

PSHs, F4 (Papua New Guinea, PNG – 398 m; Figs 1A, 8) and FA (Taiwan – 999 m; Figs 1A, 11),

were collected outside the seas of Australia.

Two of the three ABGD PSHs of *Glaciotomella*, G1 and G2, are found only off temperate

Australia (albeit G1 at the temperate-tropical boundary) at comparable depths (2474 and

2007 m). The remaining PSH, GA is only known from a single record in shallower waters (600

m) off the Solomon Islands.

For *Rimosodaphnella*, virtually all records of the only two Australian PSHs, R1 and R2, are

limited to a relatively restricted area off the east coast of Tasmania (Fig. 1B), where they

occupy fairly disjunct bathymetric ranges (bathyal vs. abyssal; Fig. 11). The only sample of

RA is found off the south coast of Chile (Fig. 1B) in much shallower waters (766 m, Fig. 11).

All other *Rimosodaphnella* ABGD PSHs occur at lower latitudes in the warmer and shallower

waters of the Philippines (R4 and R5; Figs 1B, 11), Melanesia (R3, RB, RC; Figs 1B, 11) and

Polynesia (RD and RE; Figs 1B, 11).

PSH to SSH conversion

nd intra-PSH genetic distance were not available to their geographic and bathymetric distribution
The evidence for PSHs to SSHs conversion is a
simpared with each other according to their re
sis (Figs 2 and S2).
Famelica c Comparative examination of the morphological, geographic and bathymetric data available was employed to attempt the conversion of PSHs to SSHs. As generating morphological data for most species with distribution outside Australian waters was beyond the scope of this study, testing of seven PSHs (i.e. FA, GA, RA-RE), out of the total retrieved by ABGD and ASAP, was not attempted and these are pending further sampling and taxonomic investigation. As detailed below, nine of the PSHs retrieved by ABGD and ten of those retrieved by ASAP satisfied at least one of the conditions described in the methodological section. It must be noted that some of the PSHs consisted of one single sequence and, for those, support value and intra-PSH genetic distance were not available. Also, the information available on their geographic and bathymetric distribution was limited to a single collection event. The evidence for PSHs to SSHs conversion is detailed below, where congeneric PSHs are compared with each other according to their relationships as resolved by the molecular analysis (Figs 2 and S2). Two haplotypes of the *Famelica* clade represent two distinct ABGD PSHs (F1 and F3) but are combined into one (F1+F3) by ASAP. While the comparatively low genetic distance

separating them (5.2%) could be interpreted as an indication of genetic connectivity

between two allopatric conspecific populations (Fig. 1A), this is not sufficient to convert

F1+F3 to SSH, as criteria (i) and (ii) are not met. In fact, F1+F3 is not statistically supported

(Figs 2, S2) and there is considerable difference between F1 and F3 in shell features (with F1

possessing a much more slender and darker shell; Fig. 4A, D-F). For these reasons F1 and F3

(as delimited by ABGD) are converted into two separate SSHs.

The only *Famelica* PSH including more than one sample, F2, exhibited a value of intra-PSH

genetic distance (0.8%) considerably lower than the distance value (13.8%) separating it

from its closest relative, F4. In addition, F2 and F4 differ significantly in shell features (Fig.

5C, F-G) and occur at different depths in distinct geographic areas (Figs 1, 11), which clearly

suggests that they should be treated as separate SSHs.

Two sequences in the *Glaciotomella* clade are treated as distinct PSHs by ABGD (G1 and G2)

but are combined into one highly supported PSH (G1+G2; BPP=1, BS=100%; Figs 2, S2) by

ASAP. G1 and G2 are found in allopatry, they are separated by comparatively low genetic

distance (3.3%), have very similar shells (Fig. 6A-B), and in terms of observed anatomical

characters differ only in G2 exhibiting a peculiar squamose osphradium. Whether the latter

character is of any taxonomic value or a result of a developmental anomaly is not clear,

therefore the evidence available is interpreted as supporting combination of G1 and G2 (as

suggested by ASAP) and subsequent conversion to a single SSH.

In the *Rimosodaphnella* clade, R1 was separated from its most closely related PSH (RA) by

comparatively large genetic distance (6.4%; Table 2) and could be differentiated by its more

pronounced cancellate shell sculpture (Fig. 7A-C). R1 occurs in a very distant realm (Fig. 1B)

at a different depth (Fig. 11). For these reasons, and despite the comparatively low support

(BPP=0.71, BS=96%) received in the molecular analysis (Figs 2, S2), R1 is converted into SSH.

R2 fulfils all criteria necessary to be converted to SSH. This PSH is separated from its closest

relatives (R1 and RA) by high genetic distance (9.3 and 6.4%; Table 2), far higher than its

intra-PSH genetic distance (0.3%). It has a notably different shell (Fig. 7D-F) and it is found in

sympatry with R1 in the South Australian realm (Fig. 1B) at deeper sites (Fig. 11).

R3, sister to the R1+RA+R2 clade, is separated from each of these by high genetic distance

(Table 1) and differs considerably in shell shape and sculpture (Fig. 8E). It is allopatric to RA,

R1 and R2 (Fig. 1B) and is found at much shallower depth than the two latter PSHs (Fig. 11).

Due to its marked genetic and morphological distinctiveness and distinct bathymetric

distribution, R3 is converted to SSH.

y high genetic distance (9.3 and 6.4%; Table 2)

nce (0.3%). It has a notably different shell (Fig

e South Australian realm (Fig. 1B) at deeper si

+R2 clade, is separated from each of these by

nsiderably in shell shape R4 and R5 were recognised by ABGD only, while ASAP combined them into a single PSH

(R4+R5). Their members exhibit very similar shells (Fig. 8B-C) and are separated by the

lowest inter-PSHs distance (4.6%) measured for PSHs of *Rimosodaphnella*. They also occur

sympatrically (Fig. 1B) and at comparable depth (Fig. 11). As no genetic, morphological,

geographic or bathymetric criterion can be applied to reliably distinguish them, they are

here combined (as indicated by ASAP) into a single SSH.

Assigning names to SSHs

A search was conducted for all names available and potentially applicable to the nine SSHs

resulting from the conversion process described above. By consulting the relevant literature

on Raphitomidae (Dall 1881; Watson 1881; Verrill 1884; Dautzenberg and Fischer 1896;

- Schepman 1913; Barnard 1963; McLean and Poorman 1971; Bouchet and Warén 1980;
- Sysoev and Kantor 1987; Sysoev 1990, 1996b; Figueira and Absalão 2012; Bonfitto and
- Morassi 2013; Criscione *et al.* 2021) and by comparison of molecular and morphological
- data available on type specimens with the data generated on sequenced specimens, we

found five names applicable to five SSHs. One SSH, G1+G2, including the holotype of *G.*

- *investigator*, could be readily assigned to this species. As the constituent specimens of F3,
- F4, R3 and R4+R5 shared shells that closely resembled the holotypes of respectively
- *Famelica tasmanica* Sysoev & Kantor 1987 (Fig. 4D), *Veprecula polyacantha* Stahlschmidt,
- Chino & Kilburn 2012 (Fig. 5G), *Rimosodaphnella solomonensis* Bonfitto & Morassi 2013 (Fig.
- 8D) and *R. brunneolineata* Bonfitto & Morassi 2013 (Fig. 8A), these SSH were attributed to
- these taxa. This required the formal transfer (as hereby proposed) of the latter species to
- *Famelica* as *Famelica polyacantha* (Stahlschmidt, Chino & Kilburn, 2012) n. comb. As no
- available names could be found for the remaining four SSHs, these were assigned to new
- taxa: namely *Famelica turritelloides* n. sp. (F1), *F. acus* n. sp. (F2), *Rimosodaphnella*
- *guraradara* n. sp. (R1) and *R. truvana* n. sp. (R2). Formal taxonomic descriptions of these
- newly recognised species are provided below.

South Creek Contract Creek Creek

of shell length. Outer lip thin, finely sculptured; inner lip smooth, with or without very thin

callus. Anal sinus moderately deep, U-shaped. Radula, and venom apparatus absent. Animal

with broad to long, narrow funnel; cephalic tentacles moderately short, cylindrical to weakly

tapering, with small eyes situated at their lower portion not recorded. Rhynchocoel

capacious.

Remarks

lão 2012, fig. 4), (b) a very slender, thin-walled
Figs 4-5; Dall 1889, p. 117, pl. 10, fig. 12; Barr
80, figs 186-189; Sysoev and Kantor 1987, figs
9; Figueira and Absalão 2012, fig. 3) and (c) ab:
fland (Criscione *et al Famelica* is characterised by the combination of: (a) a multispiral keeled protoconch with axial pillars below the keel (this study, Fig. 3A; Bouchet and Warén 1980, p. 88-90, figs 277- 281; Figueira and Absalão 2012, fig. 4), (b) a very slender, thin-walled and spirally sculptured teleoconch (this study, Figs 4-5; Dall 1889, p. 117, pl. 10, fig. 12; Barnard 1963, fig. 1b; Bouchet and Warén 1980, figs 186-189; Sysoev and Kantor 1987, figs а-д; Sysoev 1990, pl. 2, fig. 8; 1996b, figs 28-29; Figueira and Absalão 2012, fig. 3) and (c) absence of radula, proboscis and venom gland (Criscione *et al.* 2021). This combination of morpho-anatomical characters is unique among the Raphitomidae (Criscione *et al.* 2021). The distinctive protoconch morphology of *Famelica* is shared by only one other deep-sea raphitomid genus, the monotypic *Aliceia* Dautzenberg and Fischer, 1897 (Dautzenberg and Fischer 1897, p. 182, pl. 4, fig. 16; Bouchet and Warén 1980, p. 90, fig. 230; Figueira and Absalão 2012, p. 5, fig. 2). Although its type species, *A. aenigmatica* Dautzenberg and Fischer, 1897, exhibits a very distinctive shell, with a labial process and a deep umbilicus (Dautzenberg and Fischer 1897, p. 182, pl. 4, fig. 15, 17-18; Bouchet and Warén 1980, p. 90, fig. 190; Figueira and Absalão 2012, p. 5, fig. 2), its overall shape and whorl profile are not too dissimilar from those of some species of *Famelica* (e.g. *F. bitrudis* (Barnard, 1963). No anatomical details are available for *Aliceia*, including information on the presence of a radula. Their remarkable similarity may suggest a close phylogenetic relationship between *Aliceia* and *Famelica*. However, testing of this hypothesis depends on the availability of live-collected material of *A. aenigmatica* for molecular and anatomical investigation. Prior to this study, *Famelica* included ten Recent species: the type species *F. catharinae* (Verrill & S. Smith [in Verrill], 1884) (Fig. 4B), *F. monoceros* (R. B. Watson, 1881), *F. mirmidina* (Dautzenberg & H. Fischer, 1896), *F. monotropis* (Dautzenberg & H. Fischer, 1896) (Bouchet and Warén 1980 figs 187, 188 and 189 respectively) and *F. scipio* (Dall, 1889) (Dall

1889 pl. 10, fig. 12) from the Atlantic; *F. bitrudis* (Barnard, 1963) (Barnard 1963, fig. 1b;

ZooBank registration

 http://zoobank.org/urn:lsid:zoobank.org:act:69A31671-72A6-45E1-9048-1A04352BB83E

Description

te whorl, 32 on last whorl). Axial sculpture of
inct on subsutural ramp, and below - in inters
x, with pronounced concavity at its transition
re moderately wide; aperture length less than
insculptured, inner lip smooth, wi Shell (SL=23.3, SW=6.4 mm) thin-walled, elongate-fusiform to turriform with strongly convex whorls and long, slender siphonal canal. Protoconch pagodiform, multispiral, of at least 3 whorls, all bearing a median keel, with axial pillars below keel and no sculpture above. Teleoconch of 6.5 strongly convex, uniformly dark cream whorls. Suture adpressed; subsutural ramp steep, moderately wide. Spiral sculpture of fine, regularly set rounded cords (14 on penultimate whorl, 32 on last whorl). Axial sculpture of very fine collabral growth lines, most distinct on subsutural ramp, and below - in interspaces between spiral cords. Shell base convex, with pronounced concavity at its transition to long, undulating siphonal canal. Aperture moderately wide; aperture length less than one third of shell 499 length. Outer lip thin, unsculptured, inner lip smooth, with very thin callus. Anal sinus moderately deep, U-shaped. Shell colouration light to dark orange, with dark orange protoconch. Animal with long, cylindrical rhynchostome funnel; cephalic tentacles moderately short, cylindrical, eyes not detected. Rhynchocoel long, capacious.

Remarks

 This species exhibits a very similar sculpture to that of four other *Famelica* species: *F. tasmanica* (Fig. 4D-F), *F. pacifica* (Fig. 4C), *F. catharinae* (Fig. 4B) and *F. monoceros*, from which it can be distinguished by possessing a notably higher spire. Among the high-spired *Famelica* species, *F. turritelloides* can be recognised by its strongly convex whorl profile and orange shell colour. While eyes have not been observed in studied material, it cannot be ruled out that they occur.

-
-

Famelica acus n. sp. (PSH F2)

Material examined

Holotype (Fig. 5C): New Caledonia, (-22.33, 167.37), EXBODI, CP3844, 815-970 m (MNHN

IM-2009-24922). COI: MW459351.

(Figs 3A, 5C-D)

- Paratype (Fig. 5D): Australia, GAB, (-34.77, 130.71), IN2017_C01_207, 3389 m, 1 wet (=
- ethanol-preserved specimen) (SAMA D49339).
-
- Distribution
- New Caledonia and GAB.
-
- Etymology
- In reference to its very elongate shell, derived from 'acus' (Latin = needle). Noun in
- apposition.
-
- ZooBank registration

 http://zoobank.org/urn:lsid:zoobank.org:act:03D9ECD1-37D1-439D-9DF9-4DF189D8F801

1:lsid:zoobank.org:act:03D9ECD1-37D1-439D-
19.1, SW=3.4 mm) very thin-walled, fragile, lar
pagodiform, multispiral, of four dark orange w
with axial pillars below the keel and no sculpt
i, uniformly white semitransparent w Description. Shell (SL=19.1, SW=3.4 mm) very thin-walled, fragile, lanceolate-fusiform to turriform. Protoconch pagodiform, multispiral, of four dark orange whorls, three of which bearing a median keel, with axial pillars below the keel and no sculpture above. Teleoconch of seven evenly convex, uniformly white semitransparent whorls. Subsutural ramp indistinct on early whorls; more clearly demarcated after onset of spiral sculpture, steep, moderately wide. Sculpture of very thin cords (about 36 on last whorl) and very fine collabral growth lines. Last adult whorl convex, clearly demarcated from long, tapering siphonal canal by smooth, shallow, concave transition. Aperture elongate, one third of shell length. Outer lip thin, unsculptured, attenuated towards tip of siphonal canal in its lower portion; inner lip smooth, with no callus. Anal sinus rather shallow, U-shaped. Animal with short, broad funnel (possibly contracted upon fixation); cephalic tentacles moderately short, somewhat tapering, small eyes situated at their lower portion, adjacent to base. Rhynchocoel capacious.

Remarks

The very elongate, weakly sculptured shell (Fig. 5C-D) differentiates this species from most

of its congeners. *F. bitrudis* possesses a very similar, but broader and shorter shell, which

- lacks spiral sculpture (Barnard 1963, fig. 1b; Sysoev 1996b, figs 28-29). Remains of setae of
- syllid polychaetes were found in the digestive tract of the paratype specimen. The

Invertebrate Systematics

 rhynchostomal sphincter extremely large; rhynchocoel short, rhynchostome funnel moderately developed. Radula of long, straight, cylindrical hypodermic teeth with no distinct barbs or blades.

Remarks

 $-$ not figured due to erosion) and on the PSH
rotoconch morphology is uncommon among r
) is possessed only by species of *Neopleurotor*
s *N. rufoapicatus* (Schepman, 1913) (Schepma
pl. 1, fig. 2), *N. distincta* (Bouche *Glaciotomella* is defined by the combination of: (a) a multispiral, axially ribbed and keeled protoconch (Fig. 3B), (b) strongly convex whorls, with cancellate sculpture (Fig. 6C-D, but also Criscione *et al.* 2021, fig. 3g) and (c) straight hypodermic teeth with no barbs or blade (Fig. 6). The protoconch description contained in the diagnosis (above) is based on the type species (SAMA D44120 – not figured due to erosion) and on the PSH GA (MNHN IM-2009- 19042; Fig. 3B). Such protoconch morphology is uncommon among raphitomids and (besides *Glaciotomella*) is possessed only by species of *Neopleurotomoides* Shuto, 1971, such as the type species *N. rufoapicatus* (Schepman, 1913) (Schepman 1913, p. 75, pl. 29, fig. 6c; Shuto 1971p. 6, pl. 1, fig. 2), *N. distincta* (Bouchet & Warén, 1980) (Bouchet and Warén 1980, p. 43, fig. 233), *N. callembryon* (Dautzenberg and Fischer, 1896) (Bouchet and Warén 1980, p. 43, fig. 232) and *N. aembe* (Figueira & Absalão, 2012) (Figueira and Absalão 2012, p. 8, fig. 7). General shell morphology and sculpture are very similar in species of *Glaciotomella* (this study, Fig. 6A-C; Criscione *et al.* 2021, fig. 3g) and *Neopleurotomoides* (Fig. 6D and Bouchet and Warén 1980, figs 100-103; Figueira and Absalão 2012, figs 5-6), with species of the former differing mainly by their considerably larger shells. In addition, species of the two genera share nearly identical, unbarbed hypodermic teeth (Fig. 6 for Glaciotomella; see Bouchet and Warén 1980, fig. 45 for Neopleurotomoides). The remarkable morphological similarity between *Glaciotomella* and *Neopleurotomoides* may be an indication of a close phylogenetic relationship. However, the extent of this relationship and its resulting taxonomic implications are still to be determined, pending availability of molecular data on the type species of *Neopleurotomoides*. *Glaciotomella* species are also similar to species of *Pleurotomella s.s.* (see Criscione *et al.* 2021) in possessing a shell with prominent sculpture, strongly convex whorls with an impressed suture and a long siphonal canal and a comparatively straight, awl-shaped radula. However, they differ from the latter genus in having a distinctly broader, more convex and less shouldered whorl profile. While further study is required on the radula in both genera,

differences based on examinations thus far are that *Glaciotomella* does not possess a barb

Diagnosis

re elongate, outer lip thin, inner lip with thin c
wide, deep, u-shaped.

I: head broad, blunt. Cephalic tentacles taperin

oboscis relatively long (broad and short when

scular bulb pearlescent, kidney-shaped to ova

iler Shell elongate-fusiform, moderately thick, with high spire. Protoconch multispiral, dark orange, with diagonally cancellate sculpture. Teleoconch white to cream to pale purple; whorl profile evenly convex (with angulation) to subcylindrical or pagodiform. Suture deeply impressed. Subsutural ramp wide, concave, rather steep, sculptured by dense, raised arcuate growth lines, indicating shape of the anal sinus. Below subsutural ramp, axial sculpture of regular, opisthocline ribs, rarely absent. Spiral sculpture of rather densely set cords, overriding axials, and typically forming nodules at intersections. Last adult whorl evenly convex below subsutural ramp, clearly demarcated from straight, moderately long siphonal canal. Aperture elongate, outer lip thin, inner lip with thin callus, gently recurved toward left. Anal sinus wide, deep, u-shaped. Animal uniform cream; head broad, blunt. Cephalic tentacles tapering evenly toward blunt tip; eyes very small. Proboscis relatively long (broad and short when retracted); venom

gland, convoluted; muscular bulb pearlescent, kidney-shaped to ovate.

Radular teeth of hypodermic type, straight, tightly rolled, often rugose; barbs absent;

adapical opening elongate; ventral blade sharp; base moderately broad; lateral process

present; basal opening subcircular. Ligament moderately large, broad.

Remarks

 Two main combined diagnostic features define *Rimosodaphnella*, namely: (a) a high-spired shell with cancellate sculpture (Brocchi 1814, p. 283, pl. 8, fig. 14; McLean and Poorman 1971, p. 111, fig. 50; Bonfitto and Morassi 2013, figs. 1A-C, G-I, P, 2A-B) and (b) straight, tightly rolled, unbarbed hypodermic teeth with a sharp dorsal blade (Fig. 10). An exhaustive comparison of the shell of *Rimosodaphnella* with the shell of other raphitomid genera can be found in Bonfitto and Morassi (2013). Bonfitto and Morassi (2013) critically reviewed the Recent and fossil species attributed to *Rimosodaphnella*, restricting this genus to its fossil type species *R. textilis* (Brocchi, 1814) (Fig. 7I) and to four additional Recent species: *R. morra* (Dall, 1881) (Fig. 7J, off Cuba, 822 m), *R. brunneolineata* Bonfitto & Morassi, 2013 (Fig. 8A) and *R. tenuipurpurata* Bonfitto & Morassi, 2013 (Fig. 8J) and (both from the Philippines, 180-250 m) as well as *R. solomonensis* Bonfitto & Morassi, 2013 (Fig. 8D) (from Solomon Islands and Vanuatu, 456-551 m). Bonfitto and Morassi (2013) removed from the genus *R. deroyae* McLean & Poorman 1971 (off Galapagos Islands, 200 m). However, the

Page 26 of 66

g evenly toward blunt tip; weak longitudinal fu

en; eyes very small. Proboscis relatively long (

d of medium length, convoluted; muscular bu

dium size. Radular tooth hypodermic, straight

y 175 µm in length; barb absent narrow striae. Ribs more elevated on early teleoconch whorls and becoming lower, with shallower interspaces on last adult whorl. Spiral cords forming weak, horizontally elongated nodules at intersections with axial ribs. Subsutural ramp sculptured by dense, raised arcuate growth lines, indicating shape of the anal sinus. Last adult whorl evenly convex below subsutural ramp, clearly demarcated from straight, moderately long siphonal canal. Last whorl periphery with predominant spiral sculpture; axial ribs becoming obsolete toward base. Aperture elongate, approximately 40% of shell length; outer lip thin, unsculptured; inner lip whitish, with thin callus, gently recurved toward left. Anal sinus wide, deep, U- shaped. Animal uniform cream; head broad, blunt. Cephalic tentacles of medium length, broad at base, tapering evenly toward blunt tip; weak longitudinal furrow observed in right tentacle of one specimen; eyes very small. Proboscis relatively long (broad and short when retracted); venom gland of medium length, convoluted; muscular bulb pearlescent, bean- shaped to ovate, of medium size. Radular tooth hypodermic, straight, tightly rolled, attaining approximately 175 µm in length; barb absent; adapical opening elongate; ventral blade sharp, approximately 1/5 of length of tooth; basal third of tooth somewhat rugose, in some teeth distinctly rugose throughout, in places with deep indentations; base moderately broad, rugose; lateral process present; basal opening subcircular, unrestricted. Ligament relatively large and broad.

Remarks

 The shell of this species (Fig. 7A-C) can be differentiated from most congeners by the combination of larger size and convex whorl profile with weakly-defined shoulder. *R. truvana* is comparable in size and sculpture but possesses distinctly shouldered whorls. Most other congeners, particularly *R. deroyae* (Fig. 7H) and the PSHs RC, RD and RE (Fig. 8G- I), possess fewer but more prominent sculptural elements. Finally, the PSH RA (Fig. 7G) can be differentiated from *R. guraradara* by a complete lack of axial elements in the former. The radula of *R. guraradara* is considerably longer than that of *R. truvana* based on the material examined.

http://www.publish.csiro.au/journals/is

- Material examined
- Holotype (Fig. 7D). Australia, TAS/VIC, Bass Strait, (-39.55, 149.55), IN2017_V03_030, 4197-
- 4133 m, (AMS C.571685). COI: MN983200.
- Paratypes. As per holotype, 1 wet (AMS C.519315), 1 wet (AMS C.519316), 1 wet (AMS
- C.571615), 1 wet (AMS C.571686), 1 wet (AMS C.571689); TAS, Flinders CMR, (-40.47,
- 747 149.40), IN2017 V03 015, 4114-4139 m, 1 wet (AMS C.571687).
-
- Etymology
- In reference to the closest land mass to the type locality, derived from 'Truvana' (Aboriginal
- Australian language Palawa Kani = Cape Barren Island), name in apposition.
-
- Zoobank registration number
- http://zoobank.org/urn:lsid:zoobank.org:act:614EE2C7-1A95-4049-9179-08DA467FD60D
-
- Distribution
- Tasman Sea.
-
- lawa Kani = Cape Barren Island), name in appo

umber

1:lsid:zoobank.org:act:614EE2C7-1A95-4049-9

1:lsid:zoobank.org:act:614EE2C7-1A95-4049-9

1:lsid:zoobank.org:act:614EE2C7-1A95-4049-9

1:lsid:zoobank.org:act:614EE2C7-1 Description. Shell solid, broadly fusiform, (SL=24.5 mm, SW=11.2 mm), protoconch eroded. Teleoconch uniformly whitish, of at least six whorls. Early teleoconch whorls convex, subsequent whorls distinctly shouldered with nearly subcylindrical profile, giving spire somewhat stepped appearance. Suture impressed. Subsutural ramp concave, sculptured with fine, arcuate collabral growth lines. Sculpture of rounded, slightly opisthocline axial ribs (about 22 on penultimate and last whorl) intersected by dense, uneven cords. Axials more elevated, with deeper interstices on early teleoconch whorls and lower, densely set on last adult whorl. Spiral cords rounded, becoming somewhat nodulose at intersections with axial elements. Last adult whorl strongly convex, sculptured predominantly by spiral elements, clearly demarcated from long and slender siphonal canal. Aperture wide, elongate-pyriform, approximately half of shell length; outer lip thin, simple; inner lip whitish, with well- developed callus, gently twisting toward left. Anal sinus wide, moderately deep, U-shaped. Anatomy (based on holotype and paratypes AMS C.571687, C.571686, C.571689): animal uniform whitish to cream; head broad, blunt. Penis large, somewhat flattened, muscular, coiling clockwise, ¼ distal end tapering. Cephalic tentacles broad, of medium length,

 cylindrical, tip blunt; eyes very small. Osphradium and ctenidium large. Venom apparatus: Proboscis far retracted into rhyncocoel in specimens examined; venom gland moderately long, convoluted; muscular bulb large, pearlescent, kidney shaped.

777 Radular teeth of hypodermic type, straight, tightly rolled, attaining approximately 125 µm in length; barb absent; terminal pore elongate; ventral blade sharp, approximately 1/5 of length of tooth; some teeth rugose, in places with deep indentations; base moderately broad, of coarse texture; lateral process present; basal opening subcircular. Ligament

- comparatively broad and large.
-

Remarks

 Rimosodaphnella truvana n. sp. can be distinguished from all other congeners by its much broader and larger shell with a subcylindrical rather than convex whorl outline (Fig. 7D-F). It differs from the shell of *R. guraradara* (Fig. 7A-C) by its far more angular shoulder at the whorl periphery, the length of its aperture which is approximately equal to the length of the spire (compared to the relatively shorter aperture in the former), and its more pronounced cancellate sculpture. The radula of *R. truvana* is shorter than that of *R. guraradara.*

is Not Note

Discussion

 The five-gene phylogeny of Criscione et al. (2020) established the phylogenetic framework upon which the new genus *Glaciotomella* was recognised and described, and it is shown herein that there is strong support in both BI and ML analyses for its monophyly, as is the case for *Famelica* and *Rimosodaphnella* (Figs 2, S2).

el. It remains unclear whether this is a reflecting
ling (most species of these genera have low ab
on rate in these lineages when compared with
ther than being supported by molecular evide
is also corroborated by morpho-an The level of genetic distinctiveness reported for *cox1* sequences of the three genera (Tables 796 1, 2) was generally comparable with those reported for the other deep-sea raphitomid genera (Hallan *et al.* 2021; Criscione et al., in press) at the intra-specific level, but overall higher (up to more than three times; *e.g.* average 4.1% in *Theta* vs. 14.2% in *Glaciotomella*) at the inter-specific level. It remains unclear whether this is a reflection of a comparatively reduced species sampling (most species of these genera have low abundance) or indication 801 of a higher diversification rate in these lineages when compared with other raphitomid genus-level lineages. Other than being supported by molecular evidence, the integrity of the genera studied herein is also corroborated by morpho-anatomical features diagnostic for each genus. Among those features, the sculpture of the protoconch and the radula were particularly conserved among congeneric species and therefore represented solid diagnostic genus-level characters.

 Rimosodaphnella possessed a multispiral larval shell, with protoconch II exhibiting a diagonally cancellate sculpture (Fig. 3C). This configuration is the most commonly recorded in Raphitomidae, where it may display a range of variations. Conversely, protoconchs with a keel and axial 'pillars', such as those observed here for *Famelica* and *Glaciotomella* (Fig. 3A- B) are considered a departure from the prevailing raphitomid pattern (Bouchet *et al.* 2011). Among deep-sea raphitomid genera, keeled protoconchs are shared by *Famelica*, *Aliceia*, *Neopleurotomoides* and *Glaciotomella*, which can be separated into pairs by the two former genera having no axial elements above the keel, as opposed to the two latter ones where axial elements persist above the keel. This, and other obvious morpho-anatomical affinities, exhibited in pairs by these genera (see above Remarks to genera), coupled with the close phylogenetic relationship between *Famelica* and *Glaciotomella* (this study, Figs 2, S2; Criscione *et al.* 2021, fig. 2), might suggest that all deep-sea raphitomid genera with keeled protoconchs are part of the same lineage (clade A of fig. 2 of Criscione *et al.* 2021), although this requires testing through molecular data. Nevertheless, owing to the presence in that 821 lineage of at least one genus with a typical diagonally cancellate protoconch

Invertebrate Systematics

 (Rimosodaphnella: this study, Figs 2, S1; Criscione *et al.* 2021, fig. 2), it is unlikely that a 'keeled-pillared' protoconch is the plesiomorphic state for this lineage (Criscione *et al.* 2021).

 Three main types of radular morphology have been reported for Australian deep-sea raphitomid genera (Criscione *et al.* 2021): (1) double-barbed, (2) awl-shaped with dorsal blade and (3) awl-shaped with no distinct blade. The latter two were recorded for two genera in this study: (2) in *Rimosodaphnella* (Fig. 10) and (3) in *Glaciotomella* (Fig. 9). The hypodermic teeth lacking barbs or blades, or reduced in size are commonly found in raphitomids, and are thought to be a result of secondary simplification of morphology (Kantor and Taylor 2002, see also below)(Kantor & Taylor 2002, also see discussion below). Unfortunately, diagnostic value of simplified morphology is questionable as it is virtually impossible to extract a phylogenetic signal associated with loss of a structure and to ascertain, whether loss of certain feature has happened once or several times in the evolution.

2, see also below)(Kantor & Taylor 2002, also
tic value of simplified morphology is question
phylogenetic signal associated with loss of a s
of certain feature has happened once or seve
Famelica studied here possess a r None of the species of *Famelica* studied here possess a radula or a venom apparatus, in agreement with that reported by Criscione *et al.* (2021) for the genus. As hypodermic teeth as well as a well-developed intraembolic proboscis and venom gland are plesiomorphic in Raphitomidae (Kantor and Sysoev 1989; Bouchet *et al.* 2011), the lack or reduction of these structures in some lineages has been interpreted as a derived state, which has occurred independently in multiple unrelated deep-sea lineages (Kantor and Taylor 2002; Criscione *et al.* 2021). The loss of such structures (involved in predation by stabbing and injecting venom) has in some studies been associated with adaptation to alternative feeding strategies (Kantor and Sysoev 1989; Taylor *et al.* 1993; Kantor and Taylor 2002). In some raphitomid groups, the loss of radula and venom apparatus has been accompanied by substantial anatomical modifications, such as the development of a prominent rhynchodeal introvert (*e.g.* some *Raphitoma* spp.), the expansion of the cavity between rhynchodeum and body walls (*e.g. Abyssobela* Kantor & Sysoev, 1986 and *Teretiopsis* Kantor & Sysoev, 1989) or the evolution of accessory rhynchodeal organs (e.g. Tritonoturris subrissoides (Hervier, 1897); Fedosov 2007). No obvious anatomical rearrangements have occurred in some other groups (*e.g. Taranis* Jeffreys, 1870). The well-developed rhynchodeal introvert was also found to be a common feature in Terebridae, another lineage of Conoidea remarkable for tendency toward complete loss of the venom apparatus (Taylor 1990; Taylor

http://www.publish.csiro.au/journals/is

Page 32 of 66

nomation (which has occurred in most living R
te reduction. It is possible that initially, the ro
prey during envenomation, as for cone-snails
in this group first engulf fish-preys into an enor
hem with a radular harpoon (*et al.* 1993; Castelin *et al.* 2012). Furthermore, terebrid species with a well-developed proboscis and hypodermic radulae appeared to have smaller introverts compared to those with reduced or lacking radula and/or proboscis (Taylor 1990). This observation had corroborated earlier observation on the feeding of terebrids, which demonstrated that the rhynchodeal introvert plays a primary role in feeding for some terebrids (Miller 1975). In Raphitomidae it was also showed that a well-developed introvert can occur in taxa possessing a prominent venom apparatus and radula: *Hemilienardia*, *Austrobela* and *Pueridaphne* (Kantor and Taylor 2002; Criscione *et al.* 2021). One could hypothesize that development of a functional rhynchodeal introvert is a prerequisite to simplification of the organs related to envenomation (which has occurred in most living Raphitomidae), often 864 leading to their complete reduction. It is possible that initially, the role of the introvert was limited to retention of prey during envenomation, as for cone-snails of the subgenus *Conus* (*Gastridium*). Conids of this group first engulf fish-preys into an enormous rhynchostome funnel, and then stab them with a radular harpoon (Olivera *et al.* 2015) Species of *Conus* (*Gastridium*) are remarkable for their simplified radular teeth, lacking elaborated barbs and serrations, that are structures that keep prey 'hooked' preventing its escape. Indeed, once prey is enclosed in a rhynchocoel, it cannot escape, and as the retention of prey is taken over by rhynchostome funnel, elaborated radular weaponry becomes unnecessary, and undergoes reduction. The widespread morphological simplification of the hypodermic teeth in Raphitomidae can possibly be explained by the same causal link (*i.e.* by partial transfer of 874 their functions to the rhynchodeal introvert). Progressively, the introvert might enable alternative feeding mechanisms, which do not require envenomation (similar to the mechanisms observed in Terebridae) and which eventually led to the complete loss of radula, venom gland and proboscis. Some occasional prey findings in radula-less raphitomids (*e.g.* polychaete setae identified as belonging to a member of the family Syllidae, recovered from *Famelica acus* n. sp. SAMA D49339) suggest that these species retain a predatory lifestyle. Shells are heterogeneous for most species of all three genera, but generally exhibit similar gross morphology, with very few exceptions. The overall strongly elongate shell shape of

Famelica (Fig. 4-5), the fusiform shell with strongly convex whorls of *Glaciotomella* (Fig. 6)

- and the cancellate teleoconch sculpture of *Rimosodaphnella* (Fig. 7-8) are the main
- recognisable characters in their respective composite taxa. However, while most congeneric

species (or PSHs) of shallower waters (*i.e. R. bi*
ach genus respectively). A size-depth clinal pa
(Rex and Etter 1998; Harasewych and Kantor
i conoideans in particular ('turrids' in Rex *et al*
erpreted as an adaptive re species are similar in shell morphology [*e.g. F. tasmanica* (Fig. 4D-F), *F. acus* (Fig. 5D-E) and *F. turritelloides* (Fig. 4A), *R. guraradara* (Fig. 7A-C) and *R. truvana* (Fig. 7D-F) or *R. brunneolineata* (Fig. 8A-C) and *R. solomonensis* (Fig. 8D-F)], other species or PSHs [*e.g. F. polyacantha* (Fig. 5F-G) or *Rimosodaphnella* RA (Fig. 7G)] exhibit shell features that are somewhat divergent from those shared by their congeners. Some species of *Rimosodaphnella* and *Glaciotomella* studied here have much larger shells than others. 892 Although the small sample size did not allow thorough statistical testing, the size of their shells (and bodies) appears to be positively correlated with depth, with species from the lower bathyal (*i.e. R. guraradara* and *R. truvana* for the former genus and *G. investigator* for the latter) larger than species (or PSHs) of shallower waters (*i.e. R. brunneolineata* and *Glaciotomella* GA for each genus respectively). A size-depth clinal pattern is well-known for bathyal neogastropods (Rex and Etter 1998; Harasewych and Kantor 2004; McClain *et al.* 2005) and for turriform conoideans in particular ('turrids' in Rex *et al.* 1999; McClain and Rex 899 2001) and has been interpreted as an adaptive response to an environmental gradient in the deep sea. The increase in maximum size attained with depth would provide these snails with increasing metabolic rates and competitive advantages as food resources diminish (Rex and Etter 1998; Rex *et al.* 1999). For instance, as their polychaete prey becomes scarcer with depth (Thistle *et al.* 1985), larger raphitomids may benefit from increased mobility as well as ability to feed on a wider range of prey sizes (Levinton 1982, 1987). As is often the case for deep-sea studies, it is difficult to establish the extent to which inevitably limited observations reflect such adaptive mechanisms, and to which they are confounded by sampling biases (see Criscione *et al.* 2021).

Biogeographic and bathymetric patterns

 Most species treated herein occur within an area corresponding approximately to the South Australia marine realm of Costello *et al.* (2017). Unsurprisingly, the records are concentrated in the areas of sampling: a relatively restricted portion of the GAB and on a section of the temperate eastern Australian coast between south-eastern Tasmania and Moreton Bay (Fig. 1). Samples included in this study that were collected outside this region tend to cluster around the areas targeted during MNHN-led voyages in the tropical Indo-Pacific and Pacific (Table S1).

al., in press) as well as *Pagodibela* (Hallan *et*
 e (pothetical *Glaciotomella* + *Neopleurotomoide*

species from three different Oceans, would als
 nosodaphnella is regarded in its entirety (*i.e.* in

molecular t Of the three genera studied here (restricted to the sequenced samples), one (*Glaciotomella*) seems to be relatively restricted, recorded only in Australia and Melanesia (Fig. 1A) whereas the other two have ranges that from South Australia extend respectively to the Central Pacific (*Famelica*, Fig 1A) and to the Central and Eastern Pacific (*Rimosodaphnella*, Fig. 1B). When *Famelica* is regarded in its entirety (*i.e.* including all species herein attributed to it, irrespective of the availability of molecular data for them – see Remarks under this genus), it displays a much broader, transoceanic range, encompassing all of the world's Oceans. Geographic ranges of comparable size, purportedly stretching over two or more Oceans, have been reported for other raphitomid genera present in Australia, such as *Austrobela* and *Theta* (Criscione et al., in press) as well as *Pagodibela* (Hallan *et al.* 2021). If confirmed by molecular data, a hypothetical *Glaciotomella* + *Neopleurotomoides* genus-level group (see above), including species from three different Oceans, would also feature a similarly broad range. When *Rimosodaphnella* is regarded in its entirety (*i.e.* including species whose attribution is pending molecular testing), it exhibits a prevalent South Pacific distribution. However, further sampling outside this area (such as the Atlantic, where *R. morra* is found) may reveal the presence of further species. Among the species reported here for Australia, none occur outside the two realms encompassing Australian waters (Fig. 1). However, future deep-sea sampling may reveal records beyond these realms, particularly for species occurring further north, such as *F. acus* or *F. turritelloides* n. spp. Species of other deep-sea raphitomid genera, recorded for the Tropical Australia & Coral Sea realm (#16 of Costello *et al.* 2017), commonly have ranges extending into neighbouring tropical realms [Hallan *et al.* (2021) for *Pagodibela baruna* (Sysoev, 1997); Criscione et al., in press for *Austrobela procera* (Sysoev & Bouchet, 2001) and *Spergo fusiformis* (Habe, 1962)]. The comparatively wide geographic distributions of *F. acus* and *G. investigator* are not entirely unexpected given the mounting evidence of wide distributions in some species of other deep-sea raphitomid genera, such as *Austrobela*, *Spergo* and *Theta* (Criscione et al., in press) as well as *Gladiobela* and *Pagodibela* (Hallan *et al.* 2021). As neighbouring regions with comparable environmental conditions remain virtually unexplored, it is plausible that disjunct distributions, like that observed for *F. acus* and *G. investigator* (Fig. 1A), are the result of sampling bias, rather than a reflection of any underlying biological process. Low mean *cox1* genetic distance between two populations, when compared with the range of distances calculated within one of the populations of the

 raphitomid *Gladiobela angulata* Criscione, Hallan, Puillandre & Fedosov, 2020 has been interpreted as indication of high genetic connectivity (Hallan *et al.* 2021). In order to perform similar comparisons for *F. acus* and *G. investigator*, more than one sequence from each of the two disjunct populations would be necessary. In addition, due to the aforementioned sampling bias, it remains unclear whether genetic connectivity in these species is realised through a stepping-stone process, involving additional geographically intervening populations, or through long-range dispersal of the planktotrophic larva (or both).

ere there is no overlap in their bathymetric raid of the state of the state of some sampling coverage within
thas been suggested that partitioning into sep
peciation of sister species of some other Aust
th as *Gladiobela Rimosodaphnella guraradara* and *R. truvana* are present in sympatry off the coast of East Tasmania (Fig. 1B), where there is no overlap in their bathymetric ranges, with a gap of more than 1000 m (Fig. 11), despite some sampling coverage within this gap at corresponding areas. It has been suggested that partitioning into separate bathymetric niches is the driver of speciation of sister species of some other Australian deep-sea raphitomid genera, such as *Gladiobela* (Hallan *et al.* 2021) and *Austrobela* (Criscione et al., in press), albeit the ecological factors involved remain unknown. As *R. guraradara* and *R. truvana* are not recovered as sister taxa based on our results, speciation via bathymetric partitioning appears unlikely; rather, a scenario of a secondary contact between the two species appears more plausible.

 Due to the limited material of *F. turritelloides*, we cannot infer much about its biogeography. While it cannot be ruled out that its rarity may be an artefact of sampling, it is possible that this species may indeed be relatively rare, as the regions in which it has been collected are comparatively well-sampled (MacIntosh *et al.* 2018; O'Hara *et al.* 2020) and have revealed the presence of a diverse raphitomid fauna (Criscione et al., 2020). For species of other raphitomid genera from the same regions, such as *Gladiobela* (Hallan *et al.* 2021)*, Theta and Austrotheta* (Criscione et al., in press), scarce records have been interpreted as evidence of their intrinsic rarity. Conversely, some *Rimosodaphnella* species studied here, notably *R. guraradara* and *R. truvana*, can be considered relatively common, as observed for most species of *Austrobela* (Criscione et al., in press) and some of *Gladiobela* (Hallan *et al.* 2021). Estimates of bathymetric ranges of deep-sea turriform conoideans genera have traditionally relied on morphology-based genus attribution (Bouchet 1990; Kantor *et al.* 2016). For raphitomid genera, Criscione *et al.* (2021) provided the first available depth range size, based on Australian species whose generic attribution was confirmed by molecular data. By

st oti.
Container
De Container
Container
Container
Container complementing molecular data on further species from beyond Australia, this and other studies suggest range size expansions for a number of these genera. Among these expanded depth ranges, those of *Spergo* Dall, 1895 (4432 m; Criscione et al., in press), *Famelica* (4352 m; this study), *Gladiobela* Criscione, Hallan, Puillandre and Fedosov, 2020 (4276 m; (Hallan *et al.* 2021) and *Rimosodaphnella* (3874 m; this study) are the widest four reported for the family to date. The bathymetric range of *F. acus* (at least 2574 m; Fig. 11) constitutes one of 987 the widest ever reported for a molecularly confirmed conoidean species, second only to that of *Gladiobela acris* Hallan, Criscione, Fedosov and Puillandre, 2021 (Hallan *et al.* 2021). Due to the scarcity of material for most other species studied here, little inference can be made with regards to their bathymetric zonation.

> http://www.publish.csiro.au/journals/is

- re part of: (a) the GAB Research Program [GA

e South Australian Research and Development

and Flinders University] and (b) the GAB Deep

d research program sponsored by Chevron Au

(IN2017_V03) was provided by the Marine Conflict of interest The authors declare no conflicts of interest. Acknowledgments This work has been made possible through financial support from the Australian Government (ABRS grant RF217-57, principal investigator FC). The participation of AF was also supported by the Russian Science Foundation (grant 19-74-10020). The participation of NP was also supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 865101). Voyages in the GAB were part of: (a) the GAB Research Program [GABRP – a collaboration between BP, CSIRO, the South Australian Research and Development Institute (SARDI), the University of Adelaide and Flinders University] and (b) the GAB Deepwater Marine Program (GABDMP – a CSIRO led research program sponsored by Chevron Australia]. Funding for the 'Eastern Abyss' voyage (IN2017_V03) was provided by the Marine Biodiversity Hub (MBH), supported through the Australian Government's National Environmental Science Program (NESP). The authors wish to thank the CSIRO MNF for its support in the form of sea time onboard, support personnel, scientific equipment and data management. We also thank the scientific staff and crew who participated in all voyages generating the samples studied herein. The MNHN samples used in this study originates from shore-based expeditions (PANGLAO 2004, PAPUA NIUGINI; PI Philippe Bouchet) and deep-sea cruises (AURORA 2007, BIOPAPUA, BOA 1, EXBODI, KANACONO, KAVIENG 2014, MAINBAZA, PANGLAO 2005, SALOMON 2, SALOMON BOA 3, TARASOC, ZHONGSHA 2015; PIs Philippe Bouchet, Tin-Yam Chan, Laure Corbari, Nicolas Puillandre, Sarah Samadi, Wei-Jen Chen, Bertrand Richer de Forges) conducted by MNHN, Pro-Natura International (PNI) and Institut de Recherche pour
	- le Développement as part of the Our Planet Reviewed and the Tropical Deep-Sea Benthos programs. Funders and sponsors included a bilateral cooperation research funding from the Taiwan Ministry of Science and Technology (MOST 102-2923-B-002-001-MY3, PI Wei-Jen Chen) and the French National Research Agency (ANR 12-ISV7-0005-01, PI Sarah Samadi), the Total Foundation, Prince Albert II of Monaco Foundation, Stavros Niarchos Foundation, and Richard Lounsbery Foundation. One sample was collected during the INSPIRE cruise (Chile) onboard R/V Melville (SCRIPPS institutions). All expeditions operated under the
	- regulations then in force in the countries in question and satisfy the conditions set by the

Vor Distriction Nagoya Protocol for access to genetic resources (expeditions.mnhn.fr). We would like to express our gratitude to Mandy Reid, Alison Miller and Jennifer Caiza (AMS) for assistance with registration and databasing of material, to Virginie Héros, Philippe Maestrati and Barbara Buge (MNHN) for the sample preparation and databasing and to Andrea Crowther (SAMA, Adelaide. Thanks are also due to Sue Lindsay and Chao Shen (Macquarie University, Sydney) for facilitating SEM work. We are indebted to several people providing photos of types and other relevant specimens: Adam Baldinger and Alana Rivera (MCZ, Cambridge, USA), Antonio Bonfitto (Bologna University, Italy), Jeroen Goud and Bram van der Bijl (NMNL, Leiden), Kazunori Hasegawa (NSMT, Tokyo, Japan), Lindsey Groves (LACM, Los Angeles), Peter Stahlschmidt (Universität Koblenz-Landau, Koblenz, Germany), Ellen Strong (NMNH, Washington), Alexander Sysoev (ZMMU, Moscow), Giorgio Teruzzi and Anna Alessandrello (MSNMi, Milan).

- Rex, M.A., Etter, R.J., Clain, A.J., and Hill, M.S. (1999) Bathymetric patterns of body size in deep-sea gastropods. *Evolution* **53**(4), 1298-1301.
- Ronquist, F., and Huelsenbeck, J.P. (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**(12), 1572-1574.
- Russini, V., Giannuzzi-Savelli, R., Pusateri, F., Prkic, J., Fassio, G., Modica, M.V., and Oliverio,
- M. (2020) Candidate cases of poecilogony in Neogastropoda: implications for the
- systematics of the genus *Raphitoma* Bellardi, 1847. *Invertebrate Systematics* **34**(3), 293-318.
- Saitou, N., and Nei, M. (1987) The neighbor-joining method: a new method for
- reconstructing phylogenetic trees. *Molecular Biology and Evolution* **4**(4), 406-425.
- Schepman, M.M. (1913) The Prosobranchia of the Siboga Expedition. Part V. Toxoglossa, with a supplement. *Siboga-Expeditie* **49**, 365-452.
- Shuto, T. (1971) Taxonomical notes on the turrids of the Siboga Collection originally described by M. M. Schepman, 1931 (Part 3). *Venus Kyoto* **30**, 5-22.
- Stahlschmidt, P., and Chino, M. (2012) A new species of *Gymnobela* (Gastropoda: Raphitomidae) from the Central Pacific. *Miscellanea Malacologica* **5**, 95-98.
- Sysoev, A.V. (1988) Ultra-abyssal findings of the family Turridae (Gastropoda, Toxoglossa) in the Pacific Ocean. *Zoologicheskii Zhurnal* **67**(7), 965-973.
- phylogenetic trees. *Molecular Biology and Evo*

2) The Prosobranchia of the Siboga Expedition

ent. *Siboga-Expeditie* **49**, 365-452.

20 mical notes on the turrids of the Siboga Colle

1. M. Schepman, 1931 (Part 3). *Ven* Sysoev, A.V. (1990) Gastropods of the family Turridae (Gastropoda: Toxoglossa) of the Nasca and Sala-y-Gomez underwater ridges. *Trudy Instituta Okeanologii Akademii Nauk SSSR* **124**, 245-260.
- Sysoev, A.V. (1996a) Deep-sea conoidean gastropods collected by the John Murray
- Expedition, 1933-34. *Bulletin of the Natural History Museum Zoology Series* **62**(1), 1- 30.
- Sysoev, A.V. (1996b) Taxonomic notes on South African deep-sea conoidean gastropods (Gastropoda: Conoidea) described by K.H. Barnard, 1963. *Nautilus* **110**(1), 22-29.
- Sysoev, A.V. (1997) Mollusca Gastropoda: new deep-water turrid gastropods (Conoidea)
- from eastern Indonesia. *Mémoires du Museum National d'Histoire Naturelle* **172**, 325-355.
- Sysoev, A.V., and Bouchet, P. (2001) New and uncommon turriform gastropods
- (Gastropoda: Conoidea) from the South-West Pacific. *Mémoires du Museum National d'Histoire Naturelle* **185**, 271-320.

Figure captions

ares: 1 – R1/R. guraradara, 2 - R2/R. truvana, 3

ta, A – RA, B – RB, C – RC, D – RD, E – RE). Thir

(numbered as in Costello et al., 2017). Main sc

(m.

Sometimes are also in Costello et al., 2017). Main sc

(m.

sensus Figure 1. Distribution of taxa studied herein with focus on the South West Pacific. (A) Records of sequenced specimens of *Famelica* (circles) and *Glaciotomella* (triangles). (B) Records of sequenced specimens of *Rimosodaphnella* (squares). The insets show records of *Rimosodaphnella* in off the coast of Chile (upper righolotype) and in the Tuamotu Archipelago (lower left). Numbers and letters in shapes indicate PSHs/species of: *Famelica* (circles: 1 – F1/F*. turritelloides*, 2 – F2/*F. acus*, 3 – F3/*F. tasmanica*, 4 – F4/*F. polyacantha*, A – FA); *Glaciotomella* (1 – G1/*G. investigator*, 2 – G2/*G. investigator*, A – GA) and *Rimosodaphnella* (squares: 1 – R1/*R. guraradara*, 2 - R2/*R. truvana,* 3 – R3/*R. solomonensis*, 4 – R4/*R. brunneolineata*, A – RA, B – RB, C – RC, D – RD, E – RE). Thin lines mark limits among marine realms (numbered as in Costello et al., 2017). Main scalebars = 1000 km, insets scalebars = 200 Km. Figure 2. Bayesian consensus phylogram (BI) based on analyses of *cox1* sequences. Numbers above branches indicate nodal support by Bayesian posterior probabilities (BPP). Numbers below PSH nodes indicate nodal support (%) by bootstrap (BS) resulting from the ML analysis of Fig. S1, with 100% values indicated by asterisks. Names of species described herein and sequences of Australian samples are in bold. Vertical bars mark distinct PSHs as delimited by ABGD on the corresponding *cox1* dataset. Dashed rectangles mark alternative PSH groupings as delimited by the ASAP method. Symbols and numbers next to taxon names correspond to those used in Figure 1. Samples whose shells are figured are underlined. Scale bar (right) = 7.5 mm (shells of R3-R5, RB-RE, GA), 10 mm (other shells). Figure 3. Protoconchs of species/PSHs studied here. (A) F2/*F. acus* n. sp. holotype MNHN IM- 2009-249221; (B) GA/*G.* sp. MNHN IM-2009-19042; (C) R5/*R. brunneolineata*, MNHN IM-2009-18977. Figure 4. Shells of *Famelica* PSHs/species. (A) F1/*F. turritelloides* n. sp. holotype AMS

C.482253; (B) *F. catharinae* (Verrill & S. Smith [in Verrill], 1884) holotype USMN 37871; (C) *F.*

pacifica Sysoev & Kantor, 1987 holotype ZMMU LC 5369; (D) *F. tasmanica* Sysoev & Kantor,

- Bonfitto & Morassi, 2013, holotype MZB 49758. Scale bar = 10 mm.
-
- Figure 9. Hypodermic radular teeth of *Glaciotomella* PSHs/species studied herein. (A)
- G1/*Glaciotomella investigator* Criscione et al., 2020 holotype AMS C.571621; (B)
- R2/*Glaciotomella investigator* Criscione et al. 2020, SAMA D44120. Scale bar = 100 µm.
-
- Figure 10. Hypodermic radular teeth of *Rimosodaphnella* PSHs/species studied herein. (A)
- R1/*Rimosodaphnella guraradara* n. sp., paratype AMS C.519274; (B) R2/*Rimosodaphnella*
- *truvana* n. sp., paratype AMS C.571689. Scale bar = 50 µm.
-
- PSHs/1.
Control Control Contro Figure 11. Bathymetric ranges of PSHs/taxa studied herein as inferred from records of
- sequenced specimens. Species represented by a single record are indicated by a circle.
-

 http://www.publish.csiro.au/journals/is Tables captions

Table 1. Intra- and inter-PSHs/specific genetic differentiation of *cox1* sequences in *Famelica*

by means of K2p distances. Intra-PSHs/specific distances shaded. Maximum and minimum

values of inter-PSHs/specific distance in bold. Inset: minimum, maximum and average intra-

and inter-PSHs/specific p-distances within *Famelica*. Species codes: acu, *acus*; pol,

polyacantha; tas, *tasmanica*; tur, *turritelloides*. Codes of species described herein in bold.

Table 2. Intra- and inter-PSHs/specific genetic differentiation of *cox1* sequences in

Rimosodaphnella by means of K2p distances. Intra-PSHs/specific distances shaded.

Maximum and minimum values of inter-PSHs/specific distance in bold. Inset: minimum,

maximum and average intra- and inter-PSHs/specific p-distances within *Rimosodaphnella*.

Species codes: bru, *brunneolineata*; gur, *guraradara*; sol, *solomonensis*; tru, *truvana*. Codes

Don Notice of species described herein in bold.

 http://www.publish.csiro.au/journals/is

Distribution of taxa studied herein with focus on the South West Pacific. (A) Records of sequenced specimens of Famelica (circles) and Glaciotomella (triangles). (B) Records of sequenced specimens of Rimosodaphnella (squares). The insets show records of Rimosodaphnella in off the coast of Chile (upper righolotype) and in the Tuamotu Archipelago (lower left). Numbers and letters in shapes indicate PSHs/species of: Famelica (circles: 1 – F1/F. turritelloides, 2 – F2/F. acus, 3 – F3/F. tasmanica, 4 – F4/F. polyacantha, A – FA); Glaciotomella (1 – G1/G. investigator, 2 – G2/G. investigator, A – GA) and Rimosodaphnella (squares: 1 – R1/R. guraradara, 2 - R2/R. truvana, 3 – R3/R. solomonensis, 4 – R4/R. brunneolineata, A – RA, B – RB, C – RC, D – RD, E – RE). Thin lines mark limits among marine realms (numbered as in Costello et al., 2017). Main scalebars = 1000 km, insets scalebars = 200 Km.

170x126mm (600 x 600 DPI)

Bayesian consensus phylogram (BI) based on analyses of cox1 sequences. Numbers above branches indicate nodal support by Bayesian posterior probabilities (BPP). Numbers below PSH nodes indicate nodal support (%) by bootstrap (BS) resulting from the ML analysis of Fig. S1, with 100% values indicated by asterisks. Names of species described herein and sequences of Australian samples are in bold. Vertical bars mark distinct PSHs as delimited by ABGD on the corresponding cox1 dataset. Dashed rectangles mark alternative PSH groupings as delimited by the ASAP method. Samples whose shells are figured are underlined. Scale bar (right) = 7.5 mm (shells of R3-R5, RB-RE, GA), 10 mm (other shells).

168x171mm (600 x 600 DPI)

Figure 3. Protoconchs of species/PSHs studied here. (A) F2/F. acus n. sp. holotype MNHN IM- 2009-249221; Protoconchs of species/PSHs studied here. (A) F2/F. acus n. sp. holotype MNHN IM-2009-18977.
(B) GA/G. sp. MNHN IM-2009-19042; (C) R5/R. brunneolineata, MNHN IM-2009-18977.
199x65mm (300 x 300 DPI)

199x65mm (300 x 300 DPI)

Figure 4. Shells of Famelica PSHs/species. (A) F1/F. turritelloides n. sp. holotype AMS C.482253; (B) F. catharinae (Verrill & S. Smith [in Verrill], 1884) holotype USMN 37871; (C) F. pacifica Sysoev & Kantor, 1987 holotype ZMMU LC 5369; (D) F. tasmanica Sysoev & Kantor, 1987 holotype ZMMU LC 5370; (E) F3/F. tasmanica AMS C.519370; (F) F3/F. tasmanica AMS C.571629. Scale bar = 10 mm.

Figure 5. Shells of Famelica PSHs/species. (A) F. nitida Sysoev, 1990 holotype ZMMU LC 5737; (B) F. tajourensis Sysoev & Kantor, 1987 holotype ZMMU LC 5371; (C) F2/F. acus n. sp. holotype MNHN IM- 2009- 249221; (D) F2/F. acus n. sp. paratype SAMA D49339; (E) FA/F. sp. MNHN IM-2013-61627; (F) F4/F. polyacantha, MNHN IM-2009-17104; (G) F. polyacantha (Stahlschmidt, Chino & Kilburn 2012) holotype NSMT Mo 78456. Scale bar = 10 mm (A-D, F-G); 12.5 mm (E).

Figure 6. Shells of Glaciotomella and Neopleurotomoides PSHs/species. (A) G1/G. investigator Criscione et al. 2020 holotype AMS C.571621; (B) G2/G. investigator Criscione et al. 2020, SAMA D44120; (C) GA/G. sp. MNHN IM-2009-19042, (D) N. rufoapicatus (Schepman, 1913) holotype NMNL MOLL.136877. Scale bar = 10 mm (A-B), 5 mm (C), 4 mm (D).

Figure 7. Shells of Rimosodaphnella PSHs/species. (A) R1/R. guraradara n. sp., holotype AMS C.571613; (B) R1/R. guraradara n. sp., paratype AMS C.571684; (C) R1/R. guraradara n. sp., paratype AMS C.519332; (D) R2/R. truvana n. sp., holotype AMS C.571685; (E) R2/R. truvana n. sp., paratype AMS C.571615; (F) R2/R. truvana n. sp., paratype AMS C.571686; (G) RA/R. sp., MNHN IM-2013-57410; (H) R. deroyae n. sp., holotype LACM 1543; (I) R. textilis (Brocchi, 1814), holotype MSNMi 5340; (J) R. morra (Dall, 1881) holotype MCZ 7107. Scale bar = 10 mm (A-H), 7.5 mm (I), 2.5 mm (J).

Figure 8. Shells of Rimosodaphnella PSHs/species. (A) R. brunneolineata Bonfitto & Morassi, 2013, holotype MZB 49760; (B) R5/R. brunneolineata, MNHN IM-2009-18977; (C) R4/ R. brunneolineata, MNHN IM-2007- 42460; (D) R. solomonensis Bonfitto & Morassi, 2013, holotype MNHN 25803; (E) R3/R. solomonensis Bonfitto & Morassi, 2013, MNHN IM-2007-42524; (F) RB/R. sp., MNHN IM-2007-17836; (G) RC/R. sp., MNHN IM-2013-58315; (H) RD/R. sp., MNHN IM-2007-38690; (I) RE/R. sp., MNHN IM-2007-39365; (J) R. tenuipurpurata Bonfitto & Morassi, 2013, holotype MZB 49758. Scale bar = 10 mm.

Figure 9. Hypodermic radular teeth of Glaciotomella PSHs/species studied herein. (A) G1/Glaciotomella investigator Criscione et al., 2020 holotype AMS C.571621; (B) R2/Glaciotomella investigator Criscione et al. 2020, SAMA D44120. Scale bar = 100 µm.

Figure 10. Hypodermic radular teeth of Rimosodaphnella PSHs/species studied herein. (A) R1/Rimosodaphnella guraradara n. sp., paratype AMS C.519274; (B) R2/Rimosodaphnella truvana n. sp., paratype AMS C.571689. Scale bar = 50 μ m.

Figure 11. Bathymetric ranges of PSHs/taxa studied herein as inferred from records of sequenced specimens. Species represented by a single record are indicated by a circle.

83x84mm (600 x 600 DPI)

OUNDROINSING

OUNDROINSING

Figure S1 - NJ tree generated based on a datased including all available AMS and MNHN COI sequences of Raphitomidae. In red/sequences selected as ingrioup for further analysis, based on the criteria described in the text.

method.

168x225mm (600 x 600 DPI)