

What evidence exists on how changes in marine ecosystem structure and functioning affect ecosystem services delivery? A systematic map protocol

C. Sylvie Sylvie Campagne, Joseph Langridge, Joachim Claudet, Rémi Mongruel, Eric Thiébaut

▶ To cite this version:

C. Sylvie Sylvie Campagne, Joseph Langridge, Joachim Claudet, Rémi Mongruel, Eric Thiébaut. What evidence exists on how changes in marine ecosystem structure and functioning affect ecosystem services delivery? A systematic map protocol. Environmental Evidence, 2021, 10 (1), pp.36. 10.1186/s13750-021-00251-x. hal-03479042

HAL Id: hal-03479042

https://hal.science/hal-03479042

Submitted on 14 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

How changes in marine ecosystem structure and functioning

affect ecosystem services delivery? A systematic map protocol

3 4

1

2

5

Authors' contacts

- 6 C. Sylvie Campagne^{1,2}, Joseph Langridge² Joachim Claudet³, Rémi Mongruel⁴, Eric Thiébaut¹
- 7 ¹ Sorbonne Université, CNRS, Station Biologique de Roscoff, UMR7144, Adaptation et Diversité en
- 8 Milieu Marin, Place Georges Teissier, F-29680 Roscoff, France
- 9 ² Fondation pour la Recherche sur la Biodiversité, Centre de Synthèse et d'Analyse sur la Biodiversité
- 10 (FRB-Cesab), 5 rue de l'école de médecine 34000 Montpellier, France
- 11 ³ National Center for Scientific Research, PSL Université Paris, CRIOBE, CNRS-EPHE-UPVD, Maison des
- 12 Océans, 195 rue Saint-Jacques, 75005 Paris, France
- 13 ⁴ Ifremer, Univ Brest, CNRS, UMR 6308, AMURE, Unité d'Economie Maritime, IUEM, F-29280, Plouzane,
- 14 France

15

16

17

18

19 20

21

22

23

24

25

26

27

Abstract

Background: The current biodiversity crisis calls for an urgent need to sustainably manage human uses of nature. The Ecosystem Services (ES) concept defined as « the benefits humans obtain from nature » support decisions aimed at promoting nature conservation. However, marine ecosystems, in particular, endure numerous direct pressures (e.g., habitat loss and degradation, overexploitation, pollution, climate change, and the introduction of non-indigenous species) all of which threaten ecosystem structure, functioning, and the very provision of ES. While marine ecosystems often receive less attention than terrestrial ecosystems in ES literature, it would also appear that there is a heterogeneity of knowledge within marine ecosystems and within the different ES provided. Hence, a systematic map on the existing literature will aim to highlight knowledge clusters and knowledge gaps on how changes in marine ecosystems influence the provision of marine ecosystem services. This will provide an evidence base for possible future reviews, and may help to inform eventual management and policy decision-making.

28 29 30

31

32 33

34 35

36

37

38

Method: We will search for all evidence documenting how changes in structure and functioning of marine ecosystems affect the delivery of ES, across scientific and grey literature sources. Two bibliographic databases, Scopus and Web of Science Core Collection, will be used with a supplementary search undertaken in Google scholar. Multiple organisational websites related to intergovernmental agencies, supra-national or national structures, and NGOs will also be searched. Searches will be performed with English terms only without any geographic or temporal limitations. Literature screening, against predefined inclusion criteria, will be undertaken on title, abstract, and then full texts. All qualifying literature will be subjected to coding and meta-data extraction. No formal validity appraisal will be undertaken. Indeed, the map will highlight how marine ecosystem changes impact the ES provided. Knowledge gaps will be identified in terms of which ecosystem types, biodiversity components, or ES types are most or least studied and how these categories are correlated. Finally, a database will be provided, we will narratively describe this evidence base with summary figures and tables of pertinent study characteristics.

Keywords: Ecosystem disservices; Coastal; Marine; Biodiversity; Nature's contribution to people; Spatio-temporal dynamics

Background

In the context of the current biodiversity erosion crisis, there is an increasingly urgent need to manage nature's contribution to people in a sustainable way and at a sustainable rate, thereby maintaining its potential to meet the needs of the present and future human generations [1]. Nature's Contribution to People (NCP) and Ecosystem Services (ES) concepts have gained interest in their ability to highlight our dependency on nature and all the services we extract from it [2–4]. The concept of ES is relatively recent—being introduced in the late 1970s—and has its roots in the recognition that ecosystems provide irreplaceable goods and services [5]. It has since been largely popularized by the Millennium Ecosystem Assessment as a way of thinking about the relationships between humans and nature [6]. Defined as « the benefits humans obtain from nature » [6], the ES concept helps in producing knowledge to support decisions aimed at promoting nature conservation. The related concept of NCP, popularized first by Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) regional assessments, goes beyond ES by integrating a wider range of values (e.g., rational values) and the consideration of negative contributions of nature (also called disservices [7,8]). NCP are defined as « all the contributions, both positive and negative, of living nature to people's quality of life » [1,2].

These concepts allow the studying of socio-ecological systems by proposing to analyse the interactions between living components (humans and non-humans), which concerns rigorous approaches across different scientific disciplines - ecology (e.g. [9,10]), economics (e.g. [11]), anthropology (e.g. [12,13]), philosophy (e.g.[14]) or geography (e.g. [15]) - in order to describe the numerous interactions and to understand their scope. ES concept allows for improving interactions between disciplines but also among scientists, managers, stakeholders and politicians by redefining the debates that exist around the conflicts between development and conservation [16]. The different ES can be divided into three main categories: (1) provisioning services which are products obtained from ecosystems (e.g., foods, raw materials for industry); (2) regulation and maintenance services which are benefits obtained from ecosystems (e.g., climate regulation, coastal protection) and (3) cultural services which are non-material benefits obtained from ecosystems (e.g., Recreation activities) [17–19].

Marine ecosystems provide a wide range of ES. Several lists of marine ES are available in the literature such as Bordt and Sander [20], Kermagoret *et al.* [21], Barbier [22] or Mongruel *et al.* [13], generally inspired by the classification proposed in Liquete *et al.* [19] and Beaumont *et al.*

[23]. For instance, based on the Common International Classification for Ecosystem Services (CICES) [24] and Liquete *et al.* [19], the French platform for the evaluation of ecosystems and ecosystem services listed the ES provided by marine ecosystems [13] as follows: food provision, raw materials from aquaculture, macro-algae production, molecules production, coastal protection, climate regulation, nutrient regulation, pest and disease control, symbolic, emblematic and aesthetic values, recreation and tourism, landscape amenity and knowledge production. They also considered "nursery function" and "maintenance of food webs" in their assessment even if these are sometime considered as functions [13] or as regulating services [19]. We also decided to include them. However, ecological functions like primary and secondary production provided by marine ecosystems and sometimes defined as support services were not included in this review [21,25].

Marine ecosystems endure numerous direct drivers of change: mainly habitat loss and degradation, overexploitation, pollution, introduction of non-indigenous species, and climate change. All of which threaten the future sustainability of marine and coastal areas [26]. The magnitude of these direct drivers may also depend on indirect drivers such as demographic pressure, socio-cultural context, economy, technological development, institutions and governance, and conflicts and epidemics. In 2008, Halpern et al. [27] showed in a multi-driver analysis that no area of the global ocean is unaffected by human influence and that more than 40 % of the ocean, mainly in coastal areas (e.g., NE of USA, NW Europe, East Asia, Eastern Caribbean) are strongly affected. From 2008 to 2013, « sixty-six per cent of the ocean experienced increases in cumulative human impact [...], especially in tropical, subtropical and coastal regions, while only 13% experienced decreases in response to management measures » [28]. Indeed, threats and pressures endured by marine ecosystems induce changes which have impacted marine ecosystem services and have negatively impacted human health and well-being, more specifically for Indigenous peoples and local communities dependant on fisheries [26]. For example Selim et al. (2016) highlighted pathways linking fishing and climate (drivers) to Spawning Stock Biomass and recruitment of three demersal fish species (ecosystem processes) and the consequences for delivery of the Fisheries and ultimately on food provision and economic livelihoods (ecosystem services) (Figure 1).

While marine ecosystems are increasingly included in new international legislation, the need to develop effective conservation and protection strategies remains. For instance, marine protected areas concern only 7% of the marine realm, only partly cover important sites for biodiversity and are not fully ecologically representative, well-connected, and effectively managed [29]. It is therefore crucial to apply rigorous sustainable management practices in order to help guarantee the delivery of ES and conserve the multiple benefits provided by marine ecosystems, that so many people rely on [29,30]. Hence, it first is particularly necessary to better understand such ecosystems and to highlight the related socio-ecological aspects.




Figure 1 Examples of how drivers of change can affect ecosystem services (adapted from Selim et al. 2016)

In the ES literature, marine ecosystems receive less attention than terrestrial ecosystems; less than 9% of the ES literature [31]. In the numerous ES provided by marine ecosystems, food provision (i.e., fisheries and offshore aquaculture) seems to be by far the most intensively studied marine ES [17,19]. It can be related to the fact that some marine species groups are more assessed and studied like commercial species and top predator fish stocks [29]. In addition, another focus seems to be on specific ecosystems such as coral reefs, mudflats, and seagrass beds [13]. Also, knowledge on marine ecosystems seems to decrease with distance from the coastline [13]. Only a few studies have explored ES in deep-sea ecosystems [31]. Also, the literature seems to focus on snapshot assessments instead on multi-time assessments in relation to the ecological dynamics.

The heterogeneity of knowledge in marine and coastal ecosystems and their services is a major obstacle to the effective use of scientific results by decision-makers. Thus, in order to structure existing knowledge and produce results that are useful for decision-making, a systematic map seems particularly relevant. A systematic map on the existing literature will highlight knowledge gaps and knowledge clusters on how changes in marine ecosystems influence the provision of marine ES. This will provide an evidence base for possible future reviews, and should help to inform eventual management and stakeholder decisions.

Stakeholder engagement

 This systematic map is part of the InDySEM project - Influence of ecological Dynamics on production and demand for marine Ecosystem Services, submitted to a call for research projects launched by The French Foundation for Research on Biodiversity - Center for Biodiversity Synthesis and Analysis –(FRB-CESAB-) and is supervised by a scientific team and a methodological team. The scientific team is composed of researchers with expertise from ecology, economy, and sociology, with expertise on the different main marine ecosystems. The scientific team developed and built the project for the call and follow the project leader and the project officer during

Commenté [s1]: Note pour Sylvie : à ajouter et à reprendre dans la biblio:

Selim, S.A., Blanchard, J.L., Bedford, J., Webb, T.J. (2016) Direct and indirect effects of climate change and fishing on changes in coastal ecosystem services: a historical perspective from the North Sea. Regional Environmental Change, 16, 341-351 recurring meetings. Also, they validated the adjustment of the research question, the PECO elements, the search strings as well as all the ROSES elements. The methodological team is composed of systematic review and data analysis experts and follow all CEE methodological steps for systematic maps. The FRB's CESAB is a research structure with an international scope whose objective is to implement innovative work on the synthesis and analysis of existing data sets in the field of biodiversity.

Objective of the review

The principal objective of this review is firstly to record and synthesise all evidence on how spatio-temporal changes in marine biodiversity impact ES provided by these ecological systems. We will focus on changes in biodiversity from species level to ecosystems, including functional and structural diversity, and how these changes influence the services provided (i.e., provisioning, regulation and cultural). The associated disservices—negative benefits from nature perceived by Human—will also be taken into account where it is studied.

Indeed, the proposed mapping exercise differs from the 'traditional' ES literature, which often relies heavily on expert scientific knowledge and judgement for interpretation. This often results in a process that is rather subjective. By systematically mapping the existing literature, and by using recognised ES classification methods [19], our current strategy is expected to overcome previous biases and provide an objective and transparent overview.

Thus, in order to highlight knowledge gaps on how changes in marine ecosystems influence marine ES, a systematic map is chosen following specific PECO components (Table 1). The main goal is to collate existing literature and configure a map on the resulting evidence base concerning our primary question: what are the impacts of spatio-temporal dynamics of marine biodiversity such as changes in species abundance, and of ecosystems structure and functioning on the ecosystem services they provide?

177 In add 178 second

In addition, the systematic map will summarise the evidence base in terms of the following secondary questions:

- 179 How do the spatio-temporal dynamics of marine biodiversity affect ecosystem disservices?
 - $\hbox{-} \ \ \hbox{How are marine ecosystem services and disservices linked to natural or anthropogenic drivers?}$

Table 1: The different components of the systematic map

	PECO elements	Definition(s)				
Population	Marine biodiversity	This will include all types of marine ecosystems and				
	(ecosystems and species)	the species that they contain.				
Exposure	Types of changes in marine	All changes at all levels, from species to the				
	biodiversity	ecosystems, functional and structural.				
Comparator	Spatial difference - temporal	Spatial difference or difference in time				
	difference					
Outcomes	Marine ecosystem services	All qualitative or quantitative values of marine				
	(and disservices)	ecosystem services and disservices				

Methods

This systematic map follows the methodological guidelines in accordance with the Collaboration for Environmental Evidence Guidelines and Standards for Evidence Synthesis [32] and conforms to the 'RepOrting standards for Systematic Evidence Syntheses' (ROSES) for Systematic Map Protocols presented by Haddaway *et al.* [33] (See Supplementary Information 1 for the ROSES systematic map template of this study).

Searching for articles

Search string

The research sub-strings are composed in accordance to the key elements of the question representing the Population, the Exposure, and the Outcomes (Table 2). The search terms used for the sub-string on ES types includes different synonyms for each ES in order to be as inclusive as possible and comes from the different lists of marine ES based on Mongruel et al. [13]; Global Ocean Accounts Partnership [18] and Liquete et al.[19]. The search terms for the sub-string on Exposure, which concerns changes in biodiversity (from species to marine ecosystems and ecosystems) are composed of key words synonymous to "change".

The search string was tested and constructed in Web of Science Core collection in order to have the highest efficiency and the best comprehensiveness related to the test list (See more in Supplementary Information 2 for information of the building process of the search string).

Table 2: Full search string grouped by sub-strings in relation to our concept categories

Sub-string	Search terms
Term 1	(marine OR coast* OR ocean OR sea OR littoral OR maritime) AND (species OR
POPULATIO	biodiversity OR ecosystem OR ecological)
N	
Ecosystem	
Term 2 OUTCOMES Ecosystem service	("ecosystem service\$" OR "contribution to people" OR "ecosystem function\$" OR "ecosystem process" OR "landscape service\$" OR disservice\$ OR "provisioning service\$" OR ((provision OR production OR exploitation) AND (food OR fisher* OR macro-algae\$ OR molecules)) OR "biomass for nutrition" OR "biomass for materials" OR "genetic materials" OR "raw materials" OR "maintain* food webs" OR "life cycle maintenance and habitat protection" OR "habitat provision" OR "nursery function" OR "regulation service\$" OR "climate regulation" OR "carbon sequestration" OR "weather regulation" OR "atmospheric composition and conditions" OR "air quality regulation" OR "coastal protection" OR "water retention" OR "nutrient regulation" OR "nutrient cycling" OR "pathogen regulation" OR "pest and disease control" OR "mediation of waste" OR "mediation of mass" OR "cultural service\$" OR "intellectual interaction" OR "physical interaction" OR "experiential interaction\$" OR tourism OR recreation OR amenity OR aesthetic OR heritage OR symbolic OR "cognitive effect\$" OR "knowledge production" OR education)
Term 3	(dynamic\$ OR impact\$ OR effect\$ OR variation\$ OR interaction\$ OR evolution
EXPOSURE	OR change\$)
Dynamic	

The asterisk (*) at the end of a search term/word is used to accept any variant of a base term. The dollar (\$) is used to accept single or no added characters; useful for acquiring plural and singular forms. Quotation marks were used to search the exact word order (or phrase).

Search language

Searches will be performed using English terms only. All relevant international literature published in English will be included in this systematic map. Indeed, this will include diverse bibliographic documents (e.g., books, journal articles, theses and technical reports).

Table 3: Databases and search engines used

Platform	Databases and search engines	Years	Platform or Provider
Scopus	Scopus	1788–2020	Elsevier
	Arts & Humanities Citation Index	1975 -present	
	Conference Proceedings Citation Index:	1998-present	
	- Science		
Web of Science	- Social Science & Humanities	1998-present	
Core Collection	Current Chemical Reactions	1985-present	Clarivate Analytics
Core Collection	Emerging Sources Citation Index	2015-present	
	Index Chemicus	1993-present	
	Science Citation Index Expanded	1900-present	
	Social Sciences Citation Index	1956-present	
Google scholar	Google scholar	N/A	Google

Bibliographic databases

The 3 sub-strings of the search equation will be combined with "AND" and searched in title, abstract and keywords of the Scopus and WOS databases using the search tags "TITLE-ABS-KEY" and "TS", respectively. In Web of science core collection, we will use exact search mode. All databases will be accessed with the subscription of the CNRS (Table 3).

Web-based search engines

Google Scholar, with the aid of the software Publish and Perish [34], will be used to retrieve scientific and grey literature (Table 3). Indeed, Google scholar's use of Boolean characters differs from WOS and Scopus and is limited in terms of number of characters, and thus search terms [35]. Therefore, we adapted the search string to correspond to what the review team deemed as the most important keywords (Table 4) and searched in the section "keywords" which searches in the title, abstract, and body text. We will export the first 300 search hits, in line with recommendations by Haddaway et al. [35].

Table 4: Search strings and Search hits

	Name	Section of search	Search string	Search hits	Date Search
LITERATURE	Web of science	TS	-	17329	20/07/2021
DATABASES	Scopus	TITLE-ABS-KEY	-	24051	20/07/2021

ONLINE SEARCH ENGINE	Google scholar	keywords	(marine OR coastal OR ocean) AND (species OR biodiversity OR ecosystem) AND "ecosystem services" AND change	300	22/07/2021
	FAO	language: "English"	fishery	50	27/08/2021
ORGANISA-	UNESCO	Filter: language: "English" - source: "UNESCO" - AuthoCorporate-en-s: "Intergovernmental Oceanographic Commission" - nature of content: "guide" AND "manuals and handbooks"	marine ecosystem service	50	19/08/2021
TIONAL WEBSITES	UNEP	Filters: "Reports and publications" AND "Publication" AND "Report", "Ecosystems and biodiversity" AND "oceans and seas"	marine ecosystem service	50	19/08/2021
	US NOAA		ecosystem service	15	19/08/2021
	EEA		marine ecosystem service	7	19/08/2021
	IUCN		ecosystem service	32	27/08/2021

231 Organisational websites

- The following specialist organisations will be searched in order to collect reports with primary data related to our question.
 - (1) Intergovernmental agencies under the aegis of the UN
 - Food and Agriculture Organization of the United Nations (FAO)
 - United Nations Educational, Scientific and Cultural Organization (UNESCO)
 - United Nations Environment Programme (UNEP)
 - (2) National or supra-national structures:
 - US National Oceanic and Atmospheric Administration (NOAA)
 - European Environment Agency (EEA)
 - (3) Non-governmental organization and international associations:
 - International Union for Conservation of Nature (IUCN)

For each organisation website, the search string is adapted with specific keywords with manual-systematic hand-searches and vary between the websites (Table 4). The main keywords used will be « marine ecosystem services » which contains the keywords of the population and the outcomes. Adaptation of the keywords used depends on the main topic of the organisational websites. For example, as the NOAA focuses on marine ecosystems, the search string will only be « ecosystem services ». For the FAO, the main keywords did not lead to any results, so we will focus on one ecosystem service : « fishery ». Again, the main keywords did not lead to any results in the IUCN publication websites, so we will focus only on « ecosystem service ». Other websites were tested such as the websites of the Intergovernmental Panel on Climate Change (IPCC) and of the IPBES. Nevertheless, the main keywords of our search string did not lead to any results. These Intergovernmental websites only proposed review reports and no records with primary results. A maximum of 50 references will be considered in each organisational website.

Estimating comprehensiveness of the search

The search terms were tested in Web of Science Core Collection. The review team compiled a list

of 30 articles that we considered as important and relevant for our respected fields and the research questions. These articles are listed in Supplementary Information 3. Search terms were modified and refined several times until the benchmark publications were retrieved. Indeed, words related to the population, the outcomes and the exposure were progressively added as described in Supplementary Information 3. Concerning Web of Science Core Collection, 25 out of the 30 articles in our test list were retrieved with the search terms while 3 out of the 30 articles were not found at all in Web of Science Core Collection. With all the results extracted (Web of Science Core Collection, Scopus and Google Scholar) 29 out of the 30 articles in our test list were retrieved, equating to a 96,7% comprehensiveness (Supplementary Information 3). The only article we did not retrieve was Roessig et al. (2004). We tried different search string, nevertheless the numbers of documents fund with other search string retriveing Roessig et al (2004) was either increased or other documents in the test list were not fund. The current search string at 96.7% is the best compromise.

Organisation of search hits

Once the extraction of records from each database and website is completed, we will first reassemble all bibliographic information from all the different databases in one file using Excel software and Mendeley. This will allow us to collate all records retrieved from each database. Then we will remove clear and partial duplicates based on similar DOI and similar titles.

Article screening and study inclusion criteria

Screening strategy

In order to select the relevant literature to be analysed, a three-stage filtering process will be undertaken in accordance with pre-defined screening and study eligibility criteria. Firstly, title screening will be performed followed by abstracts screening, then full-text screening. If we are unable to retrieve/find the full-text of a record where its abstract was accepted during abstract screening, then the article's full text will not be screened. The proportion of documents missing relative to accepted records will be provided. The search for full texts will be undertaken for all accepted abstracts with the use of journal subscriptions from the CNRS and the Sorbonne University. If the articles cannot be found, request for texts may be made via ResearchGate (www.researchgate.net) or the authors may be contacted through ResearchGate or directly by email. Following a conservative approach, if articles are not clearly in the inclusion criteria or do not clearly follow a reason for exclusion (details in the Eligibility criteria section), the articles will be kept for screening at the next eligibility stage.

Consistency checking

To test the consistency of the screening process, a Cohen's Kappa test [36] will be performed between two reviewers (C.S.C. and E.T.). Accordingly, 1000 titles, 10% of the abstracts selected, then 10% of retained full texts will be pre-screened by the two reviewers separately to check for

consistency in the interpretation of eligibility criteria. Due to resource limitations and the considerable number of records within all databases used (more than 40 000 in total - see Table 4), it will not be possible to do the Kappa test on 10% of the titles. If the Kappa score is less than 0.6, (which would mean a lack of consistency), the sources of disagreement will be analysed and discussed in order to resolve the reasons for disagreement. Another set of articles will then be screened and the Kappa scores re-calculated. This will be repeated until statistical agreement is reached *i.e.* the Kappa score is greater than 0.6 [37]. After which, all (if any) remaining disagreements will be discussed before beginning the full screening process.

Eligibility criteria

The selection of records will depend on the inclusion and exclusion criteria presented in Table 5. The inclusion/exclusion decisions at each screening stage will be documented. A list of records will be provided with reasons for exclusion during the full-text (see Supplementary Information 4). Concerning the title screening, only articles with a clear mention of "marine ecosystems" and "ecosystem services" with the wording of ES or ES-related concepts directly mentioning an ES will be accepted (see list Liquete *et al.* [19], or Préat [17] for ES list of marine ecosystems). In the abstract screening process, we will consider in addition to the previous criteria, the Exposure and Comparator. If an article respects the inclusion criteria of population and outcome but not of exposure – (i.e., article on marine ecosystem and ES but without consideration of spatial or temporal difference), the article will be excluded. As we are looking for ES values, we will not consider documents only focusing on methods or on policy analysis. The full-text screening will consider the previous criteria and will also consider a new criteria related to the outcomes with the need to contain ES values.

Each screening stage will be undertaken with a conservative approach in respect to our selection criteria. If qualifying information is unclear or unknown then records in question will be passed to the next eligible screening stage.

Table 5: Eligibility criteria

Criteria	Screening Steps	Inclusion criteria	Exclusion criteria
Population	Title	titles on any marine biodiversity, marine species, habitats and ecosystems	OUT: any title referring to ecosystem services provided by terrestrial and/or fresh water ecosystems.
Outcomes	Title	titles on any marine ecosystem service (as well as related terms of ES like "nature contributions to people" and all the relevant terms in the search string) no matter the types of values. titles on ecosystem service of food supply in terms of indicators of stock or population size of commercial species.	OUT: titles that address commercial species criteria with indicators other than the stock or the population size of the species
Exposure	Abstract	abstracts have to present a change (spatial or temporal change) in marine biodiversity, marine species, habitats and ecosystems or in ecosystem services	OUT: abstracts presenting an assessment - a one- time state - of the population or of the outcomes.
Comparator	Abstract	abstracts presenting temporal and spatial differences.	OUT: abstracts only assessing ES.
Temporal period	Abstract	abstract analysing data covering period at least part of the Twentieth century and/or the twenty-first century	OUT: full-texts analysing data covering period ended before 1900 (e.g., Palaeoecology analysis).

		full-text	s hav	e to conta	ain qualitative	or quantit	tative	OUT: ful	-tex	ts withou	t qualitative	or quantit	ative
Outcomes	Full-text	values	of	marine	ecosystem	services	and	values	of	marine	ecosystem	services	and
		disservi	ces					disservi	ces				

Data coding strategy

A thorough metadata extraction for the map will be performed by the same two members of the mapping team. The metadata form will be tested on 20 articles in order to test its applicability. Each selected article will be double coded. If, due to resource limitations, true double coding is not possible, an *a posteriori* cross-check will be carried out and potential disagreements will be discussed until a consensus is reached. Any missing information will be highlighted with an 'unknown'. For each retained article after the screening process, the following metadata will be extracted and coded (Supplementary Information 4):

Biblio

Bibliographic information

Authors, Article Title, Keywords, Journal, Abstract, Affiliation first author, Country affiliation, DOI, Type of document (peer-reviewed article, book, book chapter, conference object, PhD thesis, technical report, or other), Sources of funding, Year of publication

Information to be coded

All items which will be coded in the selected articles are presented in Table 6. Each option of each item will be coded « yes » or « no ». If several items are present options as well as a free space for the text present in the article.

For the services classification, we used an adaptation of Préat [17] and leave the option to right the name of the ES if it is not in the list of ES proposed.

Complementary information to be coded

In addition, on the bibliographic information and information related to the PECO elements, information on the type of documents, the type of bibliographic content and if it is primary data made in the document will be coded. Also, we will code information about the types of pressures and management in order to know more about the origin of the change of marine ecosystems.

Table 6: Meta-data to be extracted from the selected articles

-			
Item	Description	Referen	ces
Population			
Ecosystem type	A1 - Intertidal rock and other hard substrates; A2 - Intertidal sediment; A3 - Subtidal rock and other hard substrates; A4 - Circalittoral rock and other hard substrates; A5 - Subtidal sediment; A6 - Deep-sea habitats; A7 - Pelagic habitats; A8 - Ice-associated marine habitats (sub-categories of EUNIS would be classified in its higher categories) + free space for others ecosystem types	Classification EUNIS Nivea European Commission	au 2 -
Specific ecosystem	Tidal marsh; Seagrass; Coral reefs; Mangroves; Kelp forests + free space for others specific ecosystem types		
Level of biological organisation	Species; population; community	Lausch et al. Bruford et al	
Characteristics of biodiversity	Taxonomic; structural; functional		
Outcomes			
Number of ES per	Number of ES for the following ES categories: Provisioning services; Regulating services; Cultural services; Disservices		

Categories		
ES	Food provision; Raw materials; Genetic materials; Water provision; Water purification; Air quality regulation; Coastal protection; Climate regulation; Weather regulation; Nutrients cycling; Habitat provision; Pest and disease control; Symbolic and aesthetic values; Recreation and tourism; Cognitive effects; Educational opportunities (related ES terms would be considered in each ES type) + free space for other ES	Préat [17]
EDS	Free space to record all EDS assessed (e.g. release of greenhouse gases, wild animal attacks)	
ES-EDS components	Potential; capacity; use/flow; demand; preferences; desires	
ES -EDS values	Economic; social; biophysical	
Exposure / Comparator		
Scale of study area	L: local; N-: subnational; N: national; N+: supranational; C: continental; G: global; N: no case study	Liquete et al. [19]
Location of analysis	List of country names; no application; global	
Specific location	Free space	
Temporal scale of analysis	Interval (time elapsed between successive temporal replicates): free space Duration (time elapsed between first and last temporal replicates): free space	
Time frame	Past; present; future	
Spatial scale of analysis	Free space (e.g., « between a MPA and a no-MPA area»)	
Type of data	Observation; prediction	
Complementary inform	ation	
Bibliographic document type	Journal article; book; book chapter; conference object; PhD thesis; technical documentation	Following Langridge et al., [40]
Bibliographic content	Study; review; modelling; meta-analysis; other	
Primary data	Yes; No	
Pressure type	Land/sea use change; Direct exploitation; Climate change; Pollution; Invasive alien species + free space for other pressures	IPBES [41]
Management type	Status of protection; conservation; specific management + free space for other type of management	

Study validity assessment

The validity of evidence will not be assessed within this systematic map but we will code study design elements that may provide some preliminary indication of internal validity. Thus, two additional types of information will be coded. The 'bibliographic content' i.e., study, review, modelling, meta-analysis. In addition, the 'type of data' will be reported with an « yes » if the record produces primary data (Table 6).

Study Mapping and Presentation

The results will be synthesized using a narrative approach, graphs and summary tables to highlight the state of the evidence related to our primary and secondary questions. We will first depict the increase in the number of papers addressing the links between the changes of marine biodiversity and ecosystems and their provision of ES.

Summary tables and/or graphs will be created to identify the types of ecosystems, the biodiversity metrics, the ES category, and the temporal and spatial scale of each study. The types of ecosystems will be determined following the EUNIS (European Union Nature Information System) habitat classification which is a hierarchical comprehensive pan-European system for habitat classification. According to this typology at level 2, eight main categories of marine habitats will be considered: one category for pelagic habitats, one category for ice-associated

marine habitats and 6 categories for benthic habitats depending on depth and substrate type. Although this typology has been developed for European waters, its application to the world's oceans does not raise any major concerns. This list of eight habitats will be completed by specific charismatic habitats such as tidal marshes or seagrass meadows. To describe which facet of marine biodiversity is monitored to depict changes in marine populations or ecosystems, we will consider 3 classes of Essential Biodiversity Variables [42]: species populations, community composition, and ecosystem structure. For each class which corresponds to 3 different levels of biological organization (i.e., population, community, ecosystem), three essential characteristics of diversity will be used: (i) the taxonomic diversity, which is the number of different biotic entities; (ii) the structural biodiversity, which is the distribution of biological entities; and (iii) the functional diversity, which is the diversity of functions or functional traits [38]. The ES will be classified following the classification framework of Mongruel et al. [13] adapted from Liquete et al. [19] and Barbier [22]. This classification relies on the CICES which includes 3 groups of ES subdivided into different categories: provisioning services, regulation services and cultural services. In comparison with the framework proposed by the Millennium Ecosystem Assessment [43], this classification framework excludes the supporting services. As there is no classification of disservices commonly used the disservices would be reported as written in the records and then an analysis of the main concept or words used would be made.

392393394

395

396

397

398

399

400 401

402

403

404

405

406

407

375 376

377378

379

380

381

382

383

384

385 386

387

388

389

390

391

In addition, to depict the source of impact on marine biodiversity and the provision of ES, we will consider the 5 main direct drivers of change as identified by the IPBES [41]: (i) sea use change leading to habitat degradation, (ii) exploitation of marine resources, (iii) climate change, (iv) pollution, and (v) introduction of non-indigenous species. As management measures can counteract the negative effects of human stressors on marine biodiversity, we will also take into account the implementation of marine protected areas as a human intervention that may influence the ecosystem health and the provision of marine ES.

at bio

Heat maps will be used to identify knowledge clusters and knowledge gaps. A particular attention will be given to the links between habitat categories and ES types, and between biodiversity classes and levels of biological organization and ES. The results of the map will contribute to identifying the variability of the state of knowledge according to the ecosystems and their distance from the coast and the diversity of interactions between marine biodiversity and ecosystem services. We hypothesize a decrease in knowledge in offshore waters and deepsea habitats, a focus on the most emblematic or charismatic habitats as opposed to ordinary nature, while offshore habitats will be associated with global and common goods issues such as climate change.

408 409 410

The results will be used to identify key meta-data variables for a meta-analysis planned after the map review.

411 412

413

414

Declarations

· Ethics approval and consent to participate

415 Not applicable

• Consent for publication

417 Not applicable

416

418

421

426

· Availability of data and materials

- Data sharing is not applicable to the systematic map protocol in that no datasets were generated
- for this article. Datasets produced by the systematic map will be shared in open access.

Competing interests

422 Authors declare having no competing interests.

423 • Funding

- 424 This research is a product of the IndySEM project, funded by the synthesis center CESAB of the
- 425 French Foundation for Research on Biodiversity (FRB; www.fondationbiodiversite.fr).

Authors' contributions

- 427 All authors contributed to the conception and the design of the protocol. CSC was the major
- 428 contributor in writing the manuscript. ET contributed in writing the manuscript. JL contributed
- 429 more specifically to all elements of the methods and CEE guidelines. JC and RM provided
- 430 comments on the manuscript. All authors read and approved the final manuscript.

431 • Authors' information

- 432 ¹ Sorbonne Université, CNRS, Station Biologique de Roscoff, UMR7144, Adaptation et Diversité en
- 433 Milieu Marin, Place Georges Teissier, F-29680 Roscoff, France
- 434 ² Fondation pour la Recherche sur la Biodiversité, Centre de Synthèse et d'Analyse sur la Biodiversité
- 435 (FRB-Cesab), 5 rue de l'école de médecine 34000 Montpellier, France
- 436 ³ National Center for Scientific Research, PSL Université Paris, CRIOBE, CNRS-EPHE-UPVD, Maison des
- 437 Océans, 195 rue Saint-Jacques, 75005 Paris, France
- 438 ⁴ Ifremer, Univ Brest, CNRS, UMR 6308, AMURE, Unité d'Economie Maritime, IUEM, F-29280, Plouzane,
- 439 France

Acknowledgements

- 441 CSC thanks Sini Savilaakso for advice. CSC, JL and ET thank Nicolas Casajus, Data Scientist of
- 442 the Synthesis Center of the French Foundation for Research on Biodiversity (FRB-Cesab) for
- all his advice on the planned data extraction.

Supplementary information

1-10		Supplementary information 1. Excer mes Noses protocor med
449		
450	•	Supplementary Information 2: Search string evolution and estimating
451		comprehensiveness of the search
452		·
453		
454	•	Supplementary Information 3: Test list of articles for assessing comprehensiveness
455		
456		
457	•	Supplementary Information 4: Codebook outlining the meta-data extraction
458		methods
459		
		•
460	Ke	ferences
461	[1]	IPBES. Global assessment report on biodiversity and ecosystem services of the
462		Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. 2019.
463	[2]	Díaz S, Pascual U, Stenseke M, Martín-López B, Watson RT, Molnár Z, et al. Assessing nature's
464		contributions to people: Recognizing culture, and diverse sources of knowledge, can improve
465		assessments. Science (80-) 2018;359:270–2. https://doi.org/10.1126/science.aap8826.
466	[3]	Harrison PA, Harmáčková Z V., Karabulut AA, Brotons L, Cantele M, Claudet J, et al.
467		Synthesizing plausible futures for biodiversity and ecosystem services in europe and central
468		asia using scenario archetypes. Ecol Soc 2019;24. https://doi.org/10.5751/ES-10818-240227.
469	[4]	Hill R, Díaz S, Pascual U, Stenseke M, Molnár Z, Van Velden J. Nature's contributions to people:
470		Weaving plural perspectives. One Earth 2021;4:910–5.
471		https://doi.org/10.1016/j.oneear.2021.06.009.
472	[5]	Mongruel R, Méral P, Doussan I, Levrel H. L'institutionnalisation de l'approche par les services
473		écosystémiques : dimensions scientifiques, juridiques et politiques. Roche, P., Geijzendorffer,
474		I., Levrel, H., Maris, V. (Coord.), Val. la biodiversité Serv. écosystémiques Perspect. Interdiscip.
475		éditions QUAE, Paris, Fr., 2016, p. 191–216.
476	[6]	MEA. Millenium Ecosystem Assessment. Washington D.C.: Island Press; 2005.
477	[7]	Shackleton CM, Ruwanza S, Sinasson Sanni GK, Bennett S, De Lacy P, Modipa R, et al.
478		Unpacking Pandora's Box: Understanding and Categorising Ecosystem Disservices for
479		Environmental Management and Human Wellbeing. Ecosystems 2016;19:587–600.
480		https://doi.org/10.1007/s10021-015-9952-z.
481	[8]	Campagne CS, Roche PK, Salles J-M. Looking into Pandora's Box: Ecosystem disservices
482		assessment and correlations with ecosystem services. Ecosyst Serv 2018;30.

- 483 https://doi.org/10.1016/j.ecoser.2018.02.005.
- 484 [9] Vihervaara P, Mononen L, Nedkov S, Viinikka A. Biophysical Mapping and Assessment 485 Methods for Ecosystem Services 2018:72.
- 486 [10] Lavorel S, Locatelli B, Colloff MJ, Bruley E. Co-producing ecosystem services for adapting to 487 2020:375 climate change. Philos Trans R Soc В Biol Sci 488 https://doi.org/10.1098/rstb.2019.0119.
- 489 [11] Kubiszewski I, Costanza R, Anderson S, Sutton P. The future value of ecosystem services: Global 490 scenarios and national implications. Ecosyst Serv 2017;26:289–301. 491 https://doi.org/10.1016/j.ecoser.2017.05.004.
- 492 [12] Santos-martín F, Martín-lópez B, et al. Social Assessment Methods and Applications. 2016.
- 493 [13] Mongruel R, Kermagoret C, Carlier A, Scemama P, Le Mao P, Levain A, et al. Milieux marins et
 494 littoraux : évaluation des écosystèmes et des services rendus. Rapport de l'étude réalisée pour
 495 le compte du programme EFESE. 2019. Version finale du 31 octobre 2019.
- 496 https://archimer.ifremer.fr/doc/00600/71260/
- 497 [14] Maris V. Nature à vendre, les limites des services écosystémiques. 2014.
- 498 [15] Drakou EG, Virdin J, Pendleton L. Mapping the global distribution of locally-generated marine
 499 ecosystem services: The case of the West and Central Pacific Ocean tuna fisheries. Ecosyst Serv
 500 2018;31:278–88.
- 501 [16] Harrison PA, Dunford R, Barton DN, Kelemen E, Martin-Lopez B, Norton L, et al. Selecting 502 methods for ecosystem service assessment: A decision tree approach. Ecoser 503 2017;submitted:21–3. https://doi.org/10.1016/j.ecoser.2017.09.016.
- 504 [17] Preat N. Development of environmental sustainability impact assessment methods for marine 505 sourced products. PhD thesis, Ghent University, Belgium, 2021.
- 506 [18] Global Ocean Accounts Partnership. Technical Guidance on Ocean Accounting for Sustainable 507 Development. United Nat. 2019.
- 508 [19] Liquete C, Piroddi C, Drakou EG, Gurney L, Katsanevakis S, Charef A, et al. Current status and 509 future prospects for the assessment of marine and coastal ecosystem services: a systematic 510 review. PLoS One 2013;8:e67737. https://doi.org/10.1371/journal.pone.0067737.
- 511 [20] Bordt M, Saner MA. A critical review of ecosystem accounting and services frameworks 2018. 512 https://doi.org/10.3897/oneeco.3.e29306.
- 513 [21] Kermagoret C, Claudet J, Derolez V, Nugues MM, Ouisse V, Quillien N, et al. How does
 514 eutrophication impact bundles of ecosystem services in multiple coastal habitats using state515 and-transition models. Ocean Coast Manag 2019;174:144–53.
 516 https://doi.org/10.1016/j.ocecoaman.2019.03.028.
- 517 [22] Barbier EB. Marine ecosystem services. Curr Biol 2017;27:R507–10. 518 https://doi.org/10.1016/j.cub.2017.03.020.
- Beaumont NJ, Austen MC, Atkins JP, Burdon D, Degraer S, Dentinho TP, et al. Identification, definition and quantification of goods and services provided by marine biodiversity: implications for the ecosystem approach. Mar Pollut Bull 2007;54:253–65.
- 522 https://doi.org/10.1016/j.marpolbul.2006.12.003.
- 523 [24] Haines-Young R, Potschin M, Haines-Young, R., Potschin M. CICES V4.3 Report prepared 524 following consultation on CICES Version 4, August-December 2012. EEA Framework Contract 525 No EEA/IEA/09/003 2013.
- 526 [25] Bindoff NL, Cheung WWL, Kairo JG, Aristegui J, Guinder VA, Hallberg R, et al. Changing Ocean,

527	Marine Ecosystems, and Dependent Communities. IPCC Spec Rep Ocean Cryosph a Chang Clim
528	2019:447–588.

- 529 [26] IPCC, Portner HO, Roberts DC, Masson-Delmotte V, Zhai P, Tignor M, et al. IPCC The Ocean and 530 Cryosphere in a Changing Climate Summary for Policmakers. IPCC Spec Rep Ocean Cryosph a 531 Chang Clim 2019:SPM-1-SPM-42.
- 532 [27] Halpern BS, Walbridge S, Selkoe KA, Kappel CV, Micheli F, D'Agrosa C, et al. A global map of human impact on marine ecosystems. Science (80-) 2008:948–952.
- 534 [28] Halpern BS, et al. Spatial and temporal changes in cumulative human impacts on the world's ocean. Nat Commun 2015;6. https://doi.org/10.1038/ncomms8615.
- 536 Butchart SHM, Miloslavich P, Reyers B, Adams C, Bennett E, Czúcz B, et al. Chapter 3 Assessing 537 progress towards meeting major international objectives related to nature and nature's contri-538 butions to people. E. S. Brondízio, J. Settele, S. Díaz, H. Ngo (Eds.), Glob. Assess. Rep. Intergov. 539 Platf. Biodivers. Ecosyst. Serv. Bonn, Ger. IPBES, 2019. 540 https://doi.org/10.5281/zenodo.3832053.
- 541 [30] Schultz L, Folke C, Osterblom H, Olsson P. Adaptive governance, ecosystem management, and 542 natural capital. Proc Natl Acad Sci U S A 2015;112:7369–74.
- 543 [31] Townsend M, Davies K, Hanley N, Hewitt JE, Lundquist CJ, Lohrer AM. The challenge of 544 implementing the marine ecosystem service concept. Front Mar Sci 2018;5:1–13. 545 https://doi.org/10.3389/fmars.2018.00359.
- 546 [32] Collaboration for Environmental Evidence. Guidelines and Standards for Evidence synthesis in 547 Environmental Management. Version 5.0. 2018.
- 548 [33] Haddaway NR, Macura B, Whaley P, Pullin AS. ROSES for Systematic Map Protocols. Version 1.0. 2017. https://doi.org/10.6084/m9.figshare.5897284.
- 550 [34] Harzing A. Publish or Perish 2007. https://harzing.com/resources/publish-or-perish (accessed 551 July 22, 2021).
- Haddaway NR, Collins AM, Coughlin D, Kirk S. The role of google scholar in evidence reviews and its applicability to grey literature searching. PLoS One 2015;10:1–17. https://doi.org/10.1371/journal.pone.0138237.
- Cohen J. Weighted kappa: nominal scale agreement provision for scaled disagreement or partial credit. Psychol Bull 1968;70:213–20.
- [37] Landis JR, Koch GG. The Measurement of Observer Agreement for Categorical Data. Biometrics
 1977;33:159. https://doi.org/10.2307/2529310.
- 559 [38] Lausch A, Bannehr L, Beckmann M, Boehm C, Feilhauer H, Hacker JM, et al. Linking Earth 560 Observation and taxonomic, structural and functional biodiversity: Local to ecosystem 561 perspectives. Ecol Indic 2016;70:317–39. https://doi.org/10.1016/j.ecolind.2016.06.022.
- 562 [39] Bruford MW, Brummitt N, Butchart SHM, Cardoso AC, Coops NC, Dulloo E. Essential Biodiversity Variables. Science (80-) 2013;339:277–8.
- [40] Langridge J, Sordello R, Reyjol Y. Outcomes of wildlife translocations in protected areas: what is
 the type and extent of existing evidence? A systematic map protocol. Environ Evid 2020;9:1–
 11. https://doi.org/10.1186/s13750-020-00199-4.
- 567 [41] IPBES. Global assessment report of the Intergovernmental Science-Policy Platform on 568 Biodiversity and Ecosystem Services. IPBES secretariat, Bonn, Germany; 2019.
- For the pereira H. M., Ferrier S., Walters M., Geller G. N., Jongman R. H. G., Scholes R. J., et al. Essential Biodiversity Variables. Cience 2013;339:277–78.

https://doi.org/10.1126/science.1229931.

Millenium Ecosystem Service Assessment. Ecosystem Services and Human Well-Being 2005:1–

6.

74