

Stress responses in fish: From molecular to evolutionary processes

Quentin Petitjean, Séverine Jean, Allison Gandar, Jessica Côte, Pascal

Laffaille, Lisa Jacquin

► To cite this version:

Quentin Petitjean, Séverine Jean, Allison Gandar, Jessica Côte, Pascal Laffaille, et al.. Stress responses in fish: From molecular to evolutionary processes. Science of the Total Environment, 2019, 684, pp.371 - 380. 10.1016/j.scitotenv.2019.05.357 . hal-03484451

HAL Id: hal-03484451 https://hal.science/hal-03484451

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Version of Record: https://www.sciencedirect.com/science/article/pii/S0048969719324106 Manuscript_ef7e142cbbec216b89f559d49aa75a72

Stress responses in fishes: from molecular to evolutionary processes

PETITJEAN Quentin^{1,2}, JEAN Séverine², GANDAR Allison², CÔTE Jessica¹, LAFFAILLE Pascal², JACQUIN Lisa¹

 ¹ Laboratoire EDB Évolution & Diversité Biologique UMR 5174; Université de Toulouse, Université Toulouse 3 Paul Sabatier, UPS, CNRS, IRD; 118 route de Narbonne, 31062 Toulouse, France.
 ²Laboratoire ECOLAB UMR 5245; CNRS, INPT-ENSAT, Université Toulouse 3 Paul Sabatier ; avenue de l'Agrobiopole, 31326 Castanet-Tolosan, France.

Corresponding author:

Dr Lisa Jacquin Université de Toulouse Laboratoire Evolution et Diversité Biologique 118 route de Narbonne, 31062 Toulouse cedex 9 Tel: 033 5 61 55 64 74 Email: lisa.jacquin@univ-tlse3.fr

Running title: Evolution of fish stress responses

Word count: 5272 words

1 Abstract

In the context of global changes, fish are increasingly exposed to multiple stressors that have 2 cascading effects from molecules to the whole individual, thereby affecting wild fish populations 3 through selective processes. In this review, we synthetize recent advances in molecular biology and 4 evolutionary biology to outline some potentially important effects of stressors on fish across 5 6 biological levels. Given the burgeoning literature, we highlight four promising avenues of research. 7 First, (1) the exposure to multiple stressors can lead to unexpected synergistic or antagonistic effects, which should be better taken into account to improve our predictions of the effects of actual 8 9 and future human activities on aquatic organisms. Second, (2) we argue that such interactive effects might be due to switches in energy metabolism leading to threshold effects. Under multiple stress 10 exposure, fish could switch from a "compensation" strategy, i.e. a reallocation of energy to defenses 11 and repair to a "conservation" strategy, i.e. blocking of stress responses leading to strong 12 deleterious effects and high mortality. Third, (3) this could have cascading effects on fish survival 13 14 and population persistence but multiscale studies are still rare. We propose emerging tools merging 15 different levels of biological organization to better predict population resilience under multiple stressors. Fourth (4), there are strong variations in sensitivity among populations, which might arise 16 from transgenerational effects of stressors through plastic, genetic, and epigenetic mechanisms. This 17 can lead to local adaptation or maladaptation, with strong impacts on the evolutionary trajectories of 18 wild fish populations. With this review, we hope to encourage future research to bridge the gap 19 between molecular ecology, ecotoxicology and evolutionary biology to better understand the 20 evolution of responses of fishes to current and future multiple stressors in the context of global 21 22 changes.

23 Keywords: multiple stressors, metabolic strategy, adaptation, evolutionary ecotoxicology,
24 transgenerational effects

26 Introduction

27

Due to global changes, wild fish populations are exposed to multiple environmental stressors, such 28 29 as global warming, pollution or emerging pathogens, which have attracted a number of exciting 30 research questions in the last decades (Woodward et al. 2010; Crozier and Hutchings 2014; Schulte 2014; Matthaei and Lange 2016; Lange et al. 2018). For instance, what are the consequences of 31 multiple environmental stressors on fish at the molecular, physiological and whole organism levels? 32 How are stress responses integrated across levels of biological organization, especially under 33 34 multiple stressors? Why are some populations able to cope with environmental stressors while some others dwindle? What are the long-term consequences of stressors across generations and the 35 evolutionary mechanisms involved? 36

To address these questions, a large body of research has been dedicated to the molecular and 37 38 physiological pathways involved in the stress response, with a rise in the literature focusing on fish 39 in the last decades (Barton 2002; Killen et al. 2013; Schulte 2014; Schreck et al. 2016; Balasch and Tort 2019). In parallel, other studies have focused on the evolutionary mechanisms explaining 40 41 phenotypic and genetic divergence across fish populations exposed to different levels of perturbations (Crozier and Hutchings 2014; Reid et al. 2016; Whitehead et al. 2017), leading to the 42 emerging field of evolutionary ecotoxicology (Bickham 2011; Coutellec and Barata 2011; Oziolor 43 et al. 2016, 2017a; Brady et al. 2017a). However, much remains to be done to bridge the gap 44 between studies examining the impacts of multiple stressors on cells and organisms and studies 45 46 examining the evolutionary causes of variations in genotypes and sensitivity across populations. This is partly because different fields of biology (e.g. molecular ecology, ecotoxicology, 47 ecophysiology, evolutionary biology) have developed different approaches. In this review, we 48

49 attempt to build on recent advances in molecular and evolutionary ecology to give a synthetic view50 of the evolution of stress response in fish across levels of biological organization.

51

52 Definitions and objectives

53 The stress response *per se* (i.e. the physiological response triggered by stressors) is a complex series of biochemical, physiological and behavioral adjustments to environmental variations, that can 54 enable the maintenance of internal homeostasis and survival under some conditions (Bonga 1997; 55 56 Chrousos 1998; Barton 2002). Stress responses can also have negative long-term consequences depending on the level/duration/number of stressors (Schulte 2014; Balasch and Tort 2019). Some 57 stress responses are mediated by changes in stress hormones levels (catecholamines and 58 glucocorticoids, especially cortisol), while some responses are directly mediated by the stressor 59 itself, for instance temperature-inducible molecular changes in Heat Shock Proteins (HSP) or stress-60 61 induced epigenetic changes (Iwama et al. 1998; Feil and Fraga 2012; Schulte 2014; Rey et al. 2016; Schreck et al. 2016). We will here adopt a broad definition of stress responses and consider 62 responses that are initiated by stress hormones or directly by the stressor itself, following the 63 64 definition given in Schulte et al. (2014). We chose this broad definition of stress responses for two reasons. First, some mechanisms underpinning some stress-induced behavioral, reproductive or 65 morphological changes are still unknown. Second, there is now recent evidence that direct effects of 66 stressors play a larger role than previously thought, for instance through epigenetic changes (e.g. 67 Rey et al. 2016). 68

Interestingly, empirical studies bring contrasted results depending on the species, population, and stressor(s), making it difficult to predict the responses of aquatic organisms in their natural environment (Lavergne et al. 2015; Pédron et al. 2017b, 2017a; Attard et al. 2018). In this review, we synthetize recent advances in molecular biology (especially omics approaches) and 73 evolutionary ecotoxicology. We focus on abiotic thermal (changes in temperature) and chemical 74 stressors (contaminants and more generally, altered water quality), because they are of increasing concern in wild and farmed fish populations due to global changes (Schinegger et al. 2012, 2018). 75 76 We also include biotic stressors such as predators and parasites because they are also affected by global changes (e.g. invasive predators and emerging diseases) (Rahel and Olden 2008; Altizer et 77 al. 2013; Cable et al. 2017) and because they can interact with abiotic stressors (Thilakaratne et al. 78 2007; Blanar et al. 2009; Marcogliese and Pietrock 2011; Schmitz et al. 2016). However, the 79 combined effects of multiple biotic and abiotic stressors on fish across levels of biological 80 organization are still poorly known (Jackson et al. 2015; Schinegger et al. 2016a). This highlights 81 82 the need of multiscale studies from molecules to populations to better understand their effects on aquatic organisms. 83

Based on the existing literature, we propose four promising avenues of future research. We 84 first (1) propose that exposure to multiple stressors results in complex interactive effects (synergism 85 86 or antagonism) across biological scales, which need to be taken into account in future studies. Second, (2) we suggest that such effects might result from switches from a "metabolic 87 compensation strategy" to a "metabolic conservation strategy", when a certain threshold or 88 combination of stressor(s) is reached, resulting in contrasted deleterious effects at higher levels of 89 biological organization. Third, (3) we expose how this might differentially affect individual fitness 90 and population persistence, potentially explaining why some populations resist to perturbations 91 while others dwindle. Fourth, (4) we then review evidence of demonstrated interpopulation 92 variability in fish stress responses and expose how trans-generational effects of stressors could have 93 94 strong consequences for the evolutionary trajectories of wild populations, leading in some cases to extinction or to local adaptation. We believe that developing such promising avenues of future 95 research will significantly enhance our understanding of fish response to current and future stressors 96 97 in the context of global changes (Table 1).

98

99 (1) Interactive effects of multiple stressors

Most existing studies focus on the effect of one specific stress factor in isolation. Due to multiple 100 101 anthropogenic activities and global changes, aquatic organisms are however often exposed to 102 several stressors at once (Matthaei and Lange 2016; Schinegger et al. 2016b, 2018). Evaluating these multistress effects remains however a challenge: complex interactions between stressors can 103 be expected, because different kinds of stressors trigger different physiological pathways that can 104 interfere (Killen et al. 2013; Segner et al. 2014). Experimental studies using specific combinations 105 106 of stress factors have for instance unveiled complex interactive effects of thermal, toxic or biotic stressors (Marcogliese et al. 2005; Holmstrup et al. 2010; McBryan et al. 2013; Gandar et al. 2017b, 107 108 2017a; Pédron et al. 2017a), resulting in different types of interactions between stressors (Folt et al. 109 1999; Christensen et al. 2006; Côté et al. 2016; Liess et al. 2016).

Synergism occurs when individuals exposed to a stressor have a lower resistance or 110 tolerance to another stressor (Liess et al. 2016; Lange et al. 2018). For instance, an increase in 111 temperature can amplify the deleterious effects of contaminants, because increased temperature can 112 increase metabolic rates and assimilation of contaminants in fish (Noyes et al. 2009). In addition, 113 114 thermal stress can disrupt the ability of fish to set up an adequate stress response, thereby impairing their ability to limit the deleterious effects of contaminants (Kennedy and Ross 2012; Gandar et al. 115 2015, 2017b). Inversely, the metabolic costs of pollutant exposure may disturb the molecular and 116 117 physiological processes involved in thermal acclimation (Gordon 2005; López-Olmeda and 118 Sánchez-Vázquez 2011). In these cases, synergism between stressors can be expected. As a consequence, more severe deleterious effects at the cell and individual levels can be observed under 119 120 multiple stressors compared to single stressors with potential important negative consequences for fish survival and population persistence in the wild. For instance, pesticides have a higher negative 121

effect on goldfish *Carassius auratus* when they are combined with a thermal stress (Gandar et al.
2015, 2017b, 2017a; Jacquin et al. 2019).

On the opposite, antagonism occurs when exposure to one stressor limits or decreases the 124 effects of another stressor (Folt et al. 1999; Jackson et al. 2015), especially when physiological 125 adaptations to one stressor are beneficial against the exposure to additional stressors (e.g. co-126 tolerance) (Vinebrooke et al. 2004). For instance, some families of Atlantic salmon Salmo salar that 127 are tolerant to high-temperature are also more tolerant to hypoxia, because of increased heart 128 129 ventricle size and myoglobin levels (e.g. Anttila et al. 2013). Fish are also interacting with natural stressors and a number of other species (predators, parasites) in their environment, which may also 130 131 change the outcomes of stressor exposure (Holmstrup et al. 2010). For instance, a growing number of studies showed that fish exposed to contaminants are more susceptible to predation due to 132 neurotoxic effects and disruption of social interactions by pollution (e.g. Lurling and Scheffer 2007; 133 134 Tudorache et al. 2008). In addition, fish exposed to contaminants can be more susceptible to parasitism. For instance, Nile tilapia Oreochromis niloticus exposed to contamined sediments have 135 136 more gill monogean parasites due to immunosuppressive effects of toxicants (Sanchez-Ramirez et al. 2007). Conversely, parasitism can increase the sensitivity of hosts to pollutants in three-spined 137 sticklebacks Gasterosteus aculeatus (Pascoe and Cram 1977). Thermal stress can also have strong 138 effects on host-parasite interactions. For instance, the occurrence and negative effects of the 139 emerging Proliferative Kidney Disease (PKD) on brown trout Salmo trutta are increased by water 140 warming, with dramatic consequences for European salmonid populations (Okamura et al. 2011; 141 Bruneaux et al. 2017). 142

Experimental studies manipulating the composition of communities, and field studies evaluating these multiple stress effects with realistic biotic interactions should now be conducted to refine our ability to predict the effects of future stressors on aquatic wildlife (Hahn 2011).

146

Recent studies help unraveling the underpinning mechanisms explaining these complex multiple 148 stressor effects. During the last decade, a growing body of research used "omics" tools, providing 149 new insights into physiological pathways involved in fish stress responses (Gonzalez and Pierron 150 2015; Oziolor et al. 2017b). These "omics" methods have been applied at different levels of 151 152 biological organization (i.e. genomics, transcriptomics, proteomics, metabolomics) to evaluate the response of fish to temperature (e.g. Kullgren et al. 2013; Bermejo-Nogales et al. 2014a; Yang et al. 153 154 2016), altered water quality and pollution (e.g. Li et al. 2016; Oziolor et al. 2017b; Ortiz-Villanueva et al. 2018), pathogens (e.g. Lü et al. 2014), and recently to multiple stressors (Gandar et al. 2017b; 155 Pédron et al. 2017a). Although response patterns can vary according to the species and the nature, 156 intensity or duration of stressors, most studies highlight the central role of defense systems and 157 metabolism (Kassahn et al. 2009). More specifically, stress responses generally involve changes in 158 defense systems (e.g. chaperone proteins HSP, antioxidant systems, immunity), and more 159 160 importantly, energy metabolism (e.g. Evrard et al. 2013; Bermejo-Nogales et al. 2014a; Gandar et 161 al. 2015, 2017b; Naour et al. 2017). For instance, a transcriptomic approach showed that most transcript changes of delta smelt Hypomesus transpacificus exposed to salinity stress are related to 162 metabolic processes (Komoroske et al. 2016). Naour et al. (2017) also showed that stressed red 163 cusk-eel Genypterus chilensis have altered expression levels of energy metabolism pathways, 164 especially gluconeogenesis, triacylglyceride catabolism, cholesterol and fatty acid biosynthesis. 165

More generally, under single stressor exposure, energy and oxygen consumption is generally increased to face the cost of stress response, as shown in bluegill sunfish *Lepomis macrochirus* exposed to wastewater effluents (Du et al. 2018). However, responses to multiple stressors are often very different compared to single stressors. For instance, goldfish exposed to multiple thermal and toxic stressors show on the contrary a decrease of energy consumption, a decrease of amino acid metabolism, energy production and defense systems (Gandar et al. 2017a). Metabolism and the regulation of energy balance are thus central in the understanding of (multi)stress responses
(Kassahn et al. 2009). Below we propose a theoretical framework based on metabolic strategies
proposed by Sokolova et al. (2012; 2013) to better predict the impacts of single vs. multiple
stressors across levels of biological organization.

The tolerance to environmental stressors depends largely on the energy available, the 176 efficiency of acquisition and conversion of food, but also on the ability to mobilize energy resources 177 to compensate for the metabolic cost of defenses (Sokolova et al. 2012; Sokolova 2013). The 178 179 constraint of maintaining the aerobic metabolic capacity determines the stress tolerance of organisms and therefore their chances of survival (Kassahn et al. 2009). In addition, the 180 bioenergetic costs of maintaining homeostasis and defense systems while repairing damages 181 increase with the intensity and number of stressors, and this can lead to different metabolic 182 strategies and result in strong threshold effects (Sokolova et al. 2012). In other words, the effects of 183 184 stressors are very different depending on the duration, dose and combinations of stressors because alternative metabolic strategies can be displayed. Below, we describe the two main types of 185 186 metabolic responses and the associated physiological effects that can be expected under single vs. multiple stressors (Fig 1). 187

Specifically, when the dose, duration and/or number of stressors is low, fish might display a 188 "metabolic compensation" strategy (sensu Sokolova 2013). Fish implement a complete stress 189 190 response from molecules to the whole individual. In a nutshell, cortisol and catecholamine levels are increasing, leading to the reallocation of energetical resources to defence mechanisms and 191 maintenance (e.g. detoxification, immunity, damage repairs), which enables maintaining 192 193 homeostasis and survival. For instance, stressed gilthead sea bream Sparus aurata show increased mitochondrial metabolism (Bermejo-Nogales et al. 2014) and goldfish exposed to single stressor 194 (thermal or toxic stressor alone) show increased energy consumption and oxidative defenses 195 196 (Gandar et al. 2017a). Because the energy demand for maintenance increases, the energy available

for growth, reproduction, activity and storage of reserves decreases (Sokolova 2013) (Fig 1). This 197 can lead to a temporary reduction in body mass, growth and/or reproduction. For instance, fathead 198 minnows Pimephales promelas exposed to temporary anthropogenic pollution in urban streams 199 200 show reduced egg production (Crago et al. 2011). However, negative long-term consequences of single stressors for fitness might be reduced, if defence and repair mechanisms have been efficiently 201 202 mobilized, thereby reducing long-term irreversible cellular damage and mortality in the long term (Fig 1). Accordingly, goldfish exposed to single thermal or toxic stressor alone show limited 203 204 irreversible cellular damage in their gills and liver, which may ensure long-term survival (Jacquin et al. 2019). This is a "metabolic compensation" strategy where the energy is allocated to stress 205 206 resistance, allowing the organism to efficiently face stressors with limited long-term damage. In this case, strong short-term changes in metabolism, immune and anti-oxydant defenses might be 207 208 observed at the molecular and physiological level, but with low damage at the cellular level. At the 209 individual level, this could translate into low effects on mortality and fish fitness can be temporarily maintained (Fig 1). Long-term studies in wild populations are however needed to formally test these 210 211 fitness effects.

In contrast, when the dose, duration and/or number of stressors is high, fish could display a 212 "metabolic conservation" strategy (sensu Sokolova 2013) (Fig 1). In such conditions, the energy 213 demand for maintenance might be too high (i.e. beyond the aerobic metabolic capacity of the fish), 214 215 or there might be a disruption of aerobic metabolism, leading to a decrease in the energy available below the energy demand (Sokolova et al. 2012). In both cases, aerobic metabolism cannot cover 216 217 the energy demand. Metabolism can then switch to partial anaerobic to cover the costs of maintenance. Organisms can then enter into a state of "metabolic shutdown", i.e. a suppression of 218 all metabolic activity above basal metabolism, a decreased catabolism of carbohydrates and amino 219 acids. This phenomenon, called a "conservation strategy" (Sokolova et al. 2012, 2013), has been 220 221 observed under multiple stress exposure. For instance, gilthead seabream exposed to multiple

stressors (water level changes and chasing) show a blocked stress response (no cortisol increase) 222 223 and a switch from aerobic to anaerobic metabolism (Bermejo-Nogales et al. 2014). Because stress responses are blocked and defense systems are not triggered, no effects on routine behaviors and 224 225 growth and reproduction are expected (Fig 1). However, strong deleterious effects on cellular structure (high levels of cellular lesions and DNA damage) can be expected. For instance, goldfish 226 conjointly exposed to high temperature and pesticides show reduced metabolism associated with 227 increased cellular lesions of the gills and liver (Gandar et al. 2017a, 2017b; Jacquin et al. 2019). 228 229 This can then translate into high mortality and low fitness, with potential detrimental effects for population persistence (de Montaudouin et al. 2010). However, empirical studies testing the long-230 term effects of these "compensation" vs "conservation" metabolic strategies on fish fitness and 231 populations are now needed to formally test these hypotheses. 232

Taken together, the existing studies show that the stress tolerance and long-term effects of 233 stressors largely depends on the energy balance and metabolic capacity of the fish, especially in a 234 235 multistress context. However, the specific stimuli triggering a switch from a "compensation" to a "conservation" metabolic strategy remain to be determined. Identifying the thresholds and 236 combinations of stressors leading to each metabolic strategy is a major challenge for future 237 research. To tackle this question, we need more empirical studies in realistic ecological conditions 238 239 to better understand when and how the physiological endpoints of individuals are reached, and to help refining environmental risks posed by multiple anthropogenic stressors. 240

241

242 (3) Cascading effects of stressors from molecules to populations

As explained above, exposure to single vs multiple stressors thus have strong consequences across levels of biological organization, depending on the metabolic strategy adopted, with potentially strong consequences for population persistence. However, most existing studies focused on the molecular and cellular levels, and less work is devoted to the population level. Here we highlight
some potential effects of damage induced by single vs multiple stressors on individual fitness and
population persistence (Fig 2).

As seen above, exposure to moderate and/or single stressors is expected to lead to a 249 "compensation" metabolic strategy (Sokolova 2013), resulting in limited oxidative and cellular 250 251 damage (Fig 1). This is because Reactive Oxygen Species (ROS) induced by moderate/single stressors (Malek et al. 2004; Slaninova et al. 2009; Lushchak 2011; Cui et al. 2014), might be 252 rapidly neutralized via the anti-oxidant and repair machinery that is fully triggered under a 253 compensation metabolic strategy (Fig 1). This is expected to result in limited irreversible damage to 254 DNA, cells and organ functions and low effects on individual survival and fitness (e.g. Gandar et al. 255 256 2015, Jacquin et al. 2019). If individuals can maintain their long-term fitness under single or moderate stress conditions, one can expect a low risk of population extinction on the long term (e.g. 257 de Mautaudoin et al. 2010). 258

In contrast, under multiple stressors, individuals often display a "*conservation*" metabolic strategy which leads to a metabolic shutdown blocking the anti-oxydant and repair machinery (Fig 1). Excessive oxidative stress under multiple stress exposure can lead to irreversible DNA and cell damage (Valavanidis et al. 2006; Raphael et al. 2014; Santos et al. 2017), with potential impacts for organ function, fish survival and fitness (Costa et al. 2011; Xing et al. 2012; Birnie-Gauvin et al. 2017). Amplified negative effects of multiple stressors are thus expected on fish survival (Wang et al. 2017). This might have severe consequences for population persistence for at least two reasons.

First, alterations of cell structure and cycle (through activations of the tumor protein p53 for instance) can lead to enhanced cancer risk (Hagenaars et al. 2013; Topal et al. 2014), as shown in zebrafish *Danio rerio* exposed to multiple pesticides (Weber et al. 2013) or in goldfish exposed to increased temperature and pesticides levels (Gandar et al. 2017b). At the population level, this could lead to increased cancer prevalence and high extinction risk in fish populations exposed to
multiple environmental stressors (Baumann 1984, 1998; Black and Baumann 1991), although more
empirical studies are necessary to unravel the complex interactions between individual and
environmental determinants of cancer occurrence and extinction in wild populations (Hochberg and
Noble 2017; Vittecoq et al. 2018).

275 Second, excessive oxidative stress observed under multiple stressors can affect DNA structure and integrity (Cachot et al. 2006; Polard et al. 2011; Santos et al. 2017). In some cases, 276 this can lead for instance to accelerated attrition of telomeres that are non-coding, repetitive 277 sequences located at the ends of chromosomes, protecting them during replication. Individuals that 278 have stress-shortened telomeres have accelerated senescence, and diminished longevity (Bize et al. 279 280 2009; Herborn et al. 2014; Bernadotte et al. 2016; Reichert and Stier 2017). Interestingly, environmental stressors such as heat stress and pollutants have been shown to increase telomere 281 attrition in various vertebrates including fish (Boonekamp et al. 2014; Blévin et al. 2016; Debes et 282 283 al. 2016; Simide et al. 2016; McLennan et al. 2017; Angelier et al. 2018), and this has been associated with enhanced extinction risk at the population level (Dupoué et al. 2017). Integrative 284 markers such as telomeres could thus be highly relevant because they integrate stressors effects 285 from molecules to populations and help anticipating the extinction risks of wild fish populations 286 exposed to multiple stressors. 287

Ultimately, predicting the emerging properties of aquatic systems exposed to multiple stressors would need more integrative approaches to better anticipate extinction risk at the population level. Biological systems respond to multiple perturbations as multiple-level interactions, within networks of cellular traits, connected physiological processes and interspecies interactions (Geist 2011; Segner et al. 2014; Balasch and Tort 2019). Theoretical frameworks examining the dynamics of cellular to organismal and population extinction risk have been proposed (Connon et al. 2012). For instance, synthetic biology (Kitano 2002) or the Adverse

Outcome Pathways (AOP) assessment aim to establish associations between different levels of 295 296 biological organization and ultimately on population extinction risk (Ankley et al. 2010). This methodology has been applied to high-density transcription data in fathead minnows, to relate 297 298 chemical exposure to disruption of the hypothalamus-pituitary-gonadalendocrine pathway leading to reprotoxic effects and decreased reproduction in fish populations (Perkins et al. 2011). However, 299 empirical data on population dynamics and extinction risk are still lacking to fully implement such 300 301 methodologies under (multi)stress exposure and to better integrate the evolutionary responses of stressed wild populations. 302

303

304

(4) Population variability and evolutionary effects of stressors

Interestingly, a growing number of studies highlights a high variability in sensitivity to stressors 305 306 among wild fish populations exposed to contrasted environmental conditions. For instance, different populations of European flounder *Platichthys flesus* present different sensitivities to temperature, 307 hypoxia and contamination (Lavergne et al. 2015; Pédron et al. 2017b). Wild guppies Poecilia 308 309 reticulata evolving under contrasted PAHs pollution regimes in Trinidadian rivers have different morphology and behaviors (Rolshausen et al. 2015; Jacquin et al. 2017). Atlantic killifish Fundulus 310 311 *heteroclitus* living in industrially polluted sites show enhanced pollution tolerance, indicating rapid adaptation to extreme pollution (Reid et al. 2016; Whitehead et al. 2017). Such examples have led 312 to the rise of the emerging field of evolutionary toxicology, which is a promising route towards a 313 314 better understanding of long-term population responses to environmental stressors (Bickham 2011; Coutellec and Barata 2011; Brady et al. 2017b; Hamilton et al. 2017; Oziolor et al. 2017b), while 315 raising some exciting challenges for future research. For instance, the observed population 316 317 divergence can sometimes result in adaptation (enhanced fitness) or to maladaptation (decreased fitness) in stressful environments, and the conditions in which adaptive divergence occurs are not 318 clear yet (Loria et al. in press.; Ghalambor et al. 2007; Merilä and Hendry 2014). In addition, the 319

evolutionary mechanisms leading to divergence among populations are still debated, because
genetic, plastic and epigenetic processes often interact to shape their evolutionary trajectories
(Merilä and Hendry 2014; Merilä 2015; Beaman et al. 2016) (Fig 2). We discuss some of these
mechanisms below.

Under some conditions, genetic local adaptation can occur. Due to selective and neutral 324 processes, some specific resistant genotypes can be fixed in natural populations exposed to 325 stressors, leading to local adaptation, i.e. enhanced fitness in stressful environments. However, 326 empirical evidence of such genetic-based local adaptation in fish is still scarce. One of the best 327 documented example is the killifish (e.g. Oziolor et al. 2014; Reid et al. 2016). Fundulus 328 heteroclitus living in urban estuaries show strong genetic variations in toxicity mediating genes 329 330 (such as the aryl hydrocarbon receptors AHR) which confer them a high tolerance to normally lethal levels of pollution (Reid et al. 2016, Whitehead et al. 2017). Interestingly, different 331 populations having evolved in different sites show distinct molecular variations in genes connected 332 333 to signaling and detoxification pathways. This suggests that different genetic combinations might have been selected for to confer the same adaptive tolerance in different killifish populations (Reid 334 et al. 2016). High standing genetic diversity is thus crucial to determine the evolutionary potential 335 of fish populations exposed to stressors. 336

One major challenge is however to predict the evolutionary responses to multiple stressors. For instance some genetic adaptations might confer advantages against additional stressors, such as in some *Daphnia magna* populations co-adapted to warming and increased toxicity (Zhang et al. 2018). In other cases, adaptation to one stressor might impede adaptation to another stressor, such as in European flounder populations exposed to contamination that are less able to respond to thermal stress (Lavergne et al. 2015). Modelling approaches might help determining the effects of multiple environmental stressors on selective processes and population extinction risk. For instance, an individual-based demo-genetic model predicted the combined effects of river flow and ocean
warming on salmon population resilience and genetic evolution (Piou and Prévost 2013).

In addition, plasticity and transgenerational plasticity (parental effects) are major drivers of 346 rapid phenotypic divergence among populations under stress exposure, because they can enable 347 individuals with a given genotype to rapidly adjust their phenotypes to environmental conditions 348 349 (Hendry et al. 2008; Merilä and Hendry 2014; Beaman et al. 2016). For instance, thermal plasticity enable fish to rapidly adjust their metabolism to environmental temperature (Narum et al. 2013). 350 Transgenerational plasticity through parental effects is also an important evolutionary mechanism, 351 because it can directly translate the effects of parental stress environment on offspring phenotype 352 (Mousseau 1998). However, parental effects can have negative or positive effects on offspring 353 fitness. 354

355 On one hand, negative parental effects are expected for instance if there is a mismatch between parental and offspring environmental conditions, which is the case in stochastic 356 environments (Bateson et al. 2014). In addition, intrinsic costs of plasticity and stress response can 357 also reduce offspring fitness (Ghalambor et al. 2007; Räsänen and Kruuk 2007). For instance, 358 descendants of cortisol-treated females have a lower tolerance to thermal stress in Atlantic salmon 359 360 (Eriksen et al. 2006, 2007), and stressed parents of Atlantic cod Gadus morhua sire offspring with malformations and decreased survival (Morgan et al. 1999). Toxic stressors can also trigger the 361 induction of genes coding for oxidative and cellular stress across generations in zebrafish (Armant 362 363 et al. 2017). In addition, DNA alterations resulting from oxidative stress in parental gonadic cells 364 decrease the quality of gametes and offspring (Devaux et al. 2015; Herráez et al. 2017). In these cases, negative maladaptative effects of transgenerational stress are expected on offspring fitness. 365

366 On the other hand, some parental effects can be beneficial for offspring fitness, if they 367 preadapt the young to the same conditions encountered by parents (Mousseau 1998). For instance,

parents of three-spined sticklebacks Gasterosteus aculeatus exposed to parasites can improve 368 369 offspring response to the same parasites later on (Kaufmann et al. 2014). Sheepshead minnows Cyprinodon variegatus adaptively program their offspring growth to the temperature they will 370 experience, which improves their growth and survival (Salinas and Munch 2012). However, there is 371 often a trade-off between positive and negative effects generated by trans-generational stress 372 effects. For instance, positive trans-generational effects of UV radiation on offspring survival are 373 374 associated with increased costs on molecular damage and infection rate in guppies (Ghanizadeh Kazerouni et al. 2017). These examples suggest that transgenerational effects of stressors are more 375 important than previously thought, but that their associated costs can cause adaptive or maladaptive 376 377 plasticity depending on conditions and underpinning mechanisms.

378 Recent studies have unveiled some mechanisms underpinning transgenerational effects of stressors and have highlighted the crucial role of epigenetics in adaptive processes (Beldade et al. 379 2011; Rey et al. 2016; Danchin et al. 2019). Epigenetic changes in DNA in response to stressors 380 381 (such as changes in methylation and expression levels) are more frequent than previously thought, although their functional effects are not always clearly characterized (Nikinmaa and Rytkönen 382 2011). Stress exposure can cause strong epigenetic changes in some genes of parents that are 383 transmitted to their eggs and embryos (Feil and Fraga 2012; Mirbahai and Chipman 2014; Pierron 384 et al. 2014; Vandegehuchte and Janssen 2014; Cavalieri and Spinelli 2017). For instance, stress 385 386 induce epigenetic changes in genes regulating defense and metabolism pathways, cell formation and development, bones and muscles formation, as well as neurologic development (e.g. Mirbahai and 387 Chipman 2014; Vandegehuchte and Janssen 2014). Interestingly, these epigenetic changes are 388 389 transmitted to offspring, and thus impact the evolutionary trajectory of populations. For instance, offspring of stressed sticklebacks Gasterosteus aculeatus have divergent brain transcriptomes and 390 different levels of expression of DNA methylases, which is likely affecting their behavior and 391 392 fitness across generations (Metzger and Schulte 2016, 2017). These studies suggest that

transgenerational stress effects and stress tolerance depend partly on epigenetic effects, and that 393 394 adaptation (or maladaptation) to environmental stressors might well be faster than previously thought (Fig 2). Under global changes, the pace of environmental changes likely exceeds the rate of 395 genetic evolutionary changes, so that such rapid plastic and epigenetic responses to environmental 396 stressors might be crucial and result in local adaptation depending on the rate of environmental 397 changes (Hendry et al. 2008; Crozier and Hutchings 2014; Rey et al. 2016; Danchin et al. 2019). 398 However, more studies are needed to characterize the resulting effects on population adaptation and 399 persistence. 400

In addition, determining which evolutionary mechanisms are responsible for population 401 divergence is still a challenge, because a combination of all selective processes is likely to occur. 402 403 For instance, adaptive plasticity depends on genetic variations, and there is often a selection for different levels of plasticity, which complicates the evolutionary trajectories of fish evolving in 404 stressful environments (Kawecki and Ebert 2004; Ghalambor et al. 2007; Bateson et al. 2014; 405 406 Beaman et al. 2016). Consequences for population persistence will also strongly depend on standing intraspecific variations in existing populations and gene flow, and on spatio-temporal scale of 407 environmental perturbations (Fitzpatrick and Reid n.d.; Ghalambor et al. 2007; Merilä and Hendry 408 2014; Hendry 2016; Hamilton et al. 2017; Hendry et al. 2017; Whitehead et al. 2017). One of the 409 challenges for future studies will be to take into account these spatio-temporal scales of variations, 410 and to integrate the effects of multiple stressors to better predict the effects of human activities on 411 the evolutionary trajectories of wild fish populations. 412

413

414 Conclusion

To conclude, existing studies highlight the need to use multiple stress approaches across biologicalscales to better understand the effects of human-driven perturbations on fish populations. Recent

literature highlight the central role of metabolism in the stress response of fish from molecules to 417 the whole individual. Depending on the duration, level and combination of stressors, fish can set up 418 419 different metabolic strategies and energy allocation patterns, which could explain the high variability in deleterious effects on fish fitness and population persistence. Importantly, some of 420 421 these effects are transmitted across generations with potential impacts on the evolutionary trajectories of wild populations. With this review, we hope to encourage future studies to bridge the 422 gap between molecular and evolutionary approaches to better predict the impacts of present and 423 424 future stressors on aquatic wildlife under global changes.

425

426

427 Acknowledgements

- 428 We are grateful to A Perrault, C Veyssière, L Tudesque and M Cousseau for assistance. We thank S
- 429 Blanchet and G Loot for fruitful discussions. This work was supported by the Agence de l'Eau
- 430 Adour-Garonne (PHYPAT) and by the French national program CNRS EC2CO-ECODYN. QP and
- AG were supported by doctoral grants of the French Ministry of Higher Education and Research. JC
- 432 was supported by an AEAG postdoctoral grant. LJ was supported by an IDEX starting grant at
- 433 Toulouse University (IDEX-V5RJACQ). EDB is part of the French Laboratory of Excellence
- 434 project TULIP (ANR-10-LABX-41; ANR-11-IDEX-0002-02).

435

436 **References**

437

- Altizer S., R.S. Ostfeld, P.T.J. Johnson, S. Kutz, and C.D. Harvell. 2013. Climate Change and Infectious
 Diseases: From Evidence to a Predictive Framework. Science 341:514–519.
- Angelier F., D. Costantini, P. Blévin, and O. Chastel. 2018. Do glucocorticoids mediate the link between
 environmental conditions and telomere dynamics in wild vertebrates? A review. Gen Comp
 Endocrinol 256:99–111.
- Ankley G.T., R.S. Bennett, R.J. Erickson, D.J. Hoff, M.W. Hornung, R.D. Johnson, D.R. Mount, et al. 2010.
 Adverse outcome pathways: A conceptual framework to support ecotoxicology research and risk
 assessment. Environ Toxicol Chem 29:730–741.
- Anttila K., R.S. Dhillon, E.G. Boulding, A.P. Farrell, B.D. Glebe, J.A.K. Elliott, W.R. Wolters, et al. 2013.
 Variation in temperature tolerance among families of Atlantic salmon (Salmo salar) is associated
 with hypoxia tolerance, ventricle size and myoglobin level. J Exp Biol 216:1183–1190.
- Armant O., K. Gombeau, S.M.E. Houdigui, M. Floriani, V. Camilleri, I. Cavalie, and C. Adam-Guillermin. 2017.
 Zebrafish exposure to environmentally relevant concentration of depleted uranium impairs
 progeny development at the molecular and histological levels. PLOS ONE 12:e0177932.
- Attard C.R.M., C.J. Brauer, J. Sandoval-Castillo, L.K. Faulks, P.J. Unmack, D.M. Gilligan, and L.B. Beheregaray.
 2018. Ecological disturbance influences adaptive divergence despite high gene flow in golden perch (Macquaria ambigua): Implications for management and resilience to climate change. Mol Ecol 27:196–215.
- 456 Balasch J.C. and L. Tort. 2019. Netting the Stress Responses in Fish. Front Endocrinol 10.
- Barton B.A. 2002. Stress in fishes: a diversity of responses with particular reference to changes in circulating
 corticosteroids. Integr Comp Biol 42:517–525.
- Bateson P., P. Gluckman, and M. Hanson. 2014. The biology of developmental plasticity and the Predictive
 Adaptive Response hypothesis. J Physiol 592:2357–2368.
- Baumann P.C. 1984. Cancer in Wild Freshwater Fish Populations with Emphasis on the Great Lakes. J Gt
 Lakes Res 10:251–253.
- Baumann P.C. 1998. Epizootics of cancer in fish associated with genotoxins in sediment and water. Mutat
 Res Mutat Res 411:227–233.
- Beaman J.E., C.R. White, and F. Seebacher. 2016. Evolution of Plasticity: Mechanistic Link between
 Development and Reversible Acclimation. Trends Ecol Evol 31:237–249.
- Beldade P., A.R.A. Mateus, and R.A. Keller. 2011. Evolution and molecular mechanisms of adaptive
 developmental plasticity. Mol Ecol 20:1347–1363.

Bermejo-Nogales A., M. Nederlof, L. Benedito-Palos, G.F. Ballester-Lozano, O. Folkedal, R.E. Olsen, A. Sitjà Bobadilla, et al. 2014a. Metabolic and transcriptional responses of gilthead sea bream (Sparus

- 471 aurata L.) to environmental stress: new insights in fish mitochondrial phenotyping. Gen Comp
 472 Endocrinol 205:305–315.
- 473 Bernadotte A., V.M. Mikhelson, and I.M. Spivak. 2016. Markers of cellular senescence. Telomere shortening
 474 as a marker of cellular senescence. Aging 8:3–11.
- 475 Bickham J.W. 2011. The four cornerstones of Evolutionary Toxicology. Ecotoxicology 20:497–502.
- Birnie-Gauvin K., D. Costantini, S.J. Cooke, and W.G. Willmore. 2017. A comparative and evolutionary
 approach to oxidative stress in fish: A review. Fish Fish 18:928–942.
- Bize P., F. Criscuolo, N.B. Metcalfe, L. Nasir, and P. Monaghan. 2009. Telomere dynamics rather than age
 predict life expectancy in the wild. Proc R Soc Lond B Biol Sci rspb.2008.1817.
- Black J.J. and P.C. Baumann. 1991. Carcinogens and cancers in freshwater fishes. Environ Health Perspect
 90:27.
- Blanar C.A., K.R. Munkittrick, J. Houlahan, D.L. MacLatchy, and D.J. Marcogliese. 2009. Pollution and
 parasitism in aquatic animals: A meta-analysis of effect size. Aquat Toxicol 93:18–28.
- Blévin P., F. Angelier, S. Tartu, S. Ruault, P. Bustamante, D. Herzke, B. Moe, et al. 2016. Exposure to
 oxychlordane is associated with shorter telomeres in arctic breeding kittiwakes. Sci Total Environ
 563–564:125–130.
- 487 Bonga S.W. 1997. The stress response in fish. Physiol Rev 77:591–625.
- Boonekamp J.J., G.A. Mulder, H.M. Salomons, C. Dijkstra, and S. Verhulst. 2014. Nestling telomere
 shortening, but not telomere length, reflects developmental stress and predicts survival in wild
 birds. Proc Biol Sci 281:20133287.
- Brady S.P., E. Monosson, C.W. Matson, and J.W. Bickham. 2017a. Evolutionary toxicology: Toward a unified
 understanding of life's response to toxic chemicals. Evol Appl 10:745–751.
- Brady S.P., J.L. Richardson, and B.K. Kunz. 2017b. Incorporating evolutionary insights to improve
 ecotoxicology for freshwater species. Evol Appl 10:829–838.
- Bruneaux M., M. Visse, R. Gross, L. Pukk, L. Saks, and A. Vasemägi. 2017. Parasite infection and decreased
 thermal tolerance: impact of proliferative kidney disease on a wild salmonid fish in the context of
 climate change. (B. Tschirren, ed.)Funct Ecol 31:216–226.
- Cable J., I. Barber, B. Boag, A.R. Ellison, E.R. Morgan, K. Murray, E.L. Pascoe, et al. 2017. Global change,
 parasite transmission and disease control: lessons from ecology. Philos Trans R Soc Lond B Biol Sci
 372.
- Cachot J., O. Geffard, S. Augagneur, S. Lacroix, K. Le Menach, L. Peluhet, J. Couteau, et al. 2006. Evidence of
 genotoxicity related to high PAH content of sediments in the upper part of the Seine estuary
 (Normandy, France). Aquat Toxicol 79:257–267.
- 504 Cavalieri V. and G. Spinelli. 2017. Environmental epigenetics in zebrafish. Epigenetics Chromatin 10:46.
- 505 Christensen M.R., M.D. Graham, R.D. Vinebrooke, D.L. Findlay, M.J. Paterson, and M.A. Turner. 2006.
 506 Multiple anthropogenic stressors cause ecological surprises in boreal lakes. Glob Change Biol
 507 12:2316–2322.

- Chrousos G.P. 1998. Stressors, stress, and neuroendocrine integration of the adaptive response. The 1997
 Hans Selye Memorial Lecture. Ann N Y Acad Sci 851:311–335.
- Connon R.E., J. Geist, and I. Werner. 2012. Effect-Based Tools for Monitoring and Predicting the
 Ecotoxicological Effects of Chemicals in the Aquatic Environment. Sensors 12:12741–12771.
- Costa P.M., S. Caeiro, J. Lobo, M. Martins, A.M. Ferreira, M. Caetano, C. Vale, et al. 2011. Estuarine
 ecological risk based on hepatic histopathological indices from laboratory and in situ tested fish.
 Mar Pollut Bull 62:55–65.
- Côté I.M., E.S. Darling, and C.J. Brown. 2016. Interactions among ecosystem stressors and their importance
 in conservation. Proc R Soc B Biol Sci 283:20152592.
- 517 Coutellec M.-A. and C. Barata. 2011. An introduction to evolutionary processes in ecotoxicology. Ecotoxicol
 518 Lond Engl 20:493–496.
- Crago J., S.R. Corsi, D. Weber, R. Bannerman, and R. Klaper. 2011. Linking biomarkers to reproductive
 success of caged fathead minnows in streams with increasing urbanization. Chemosphere 82:1669–
 1674.
- 522 Crozier L.G. and J.A. Hutchings. 2014. Plastic and evolutionary responses to climate change in fish. Evol Appl
 523 7:68–87.
- Cui Y., B. Liu, J. Xie, P. Xu, H.-M. Habte-Tsion, and Y. Zhang. 2014. Effect of heat stress and recovery on
 viability, oxidative damage, and heat shock protein expression in hepatic cells of grass carp
 (Ctenopharyngodon idellus). Fish Physiol Biochem 40:721–729.
- D. Vinebrooke R., K. L. Cottingham, J. Norberg, Marten Scheffer, S. I. Dodson, S. C. Maberly, and U. Sommer.
 2004. Impacts of multiple stressors on biodiversity and ecosystem functioning: the role of species
 co-tolerance. Oikos 104:451–457.
- Danchin E., A. Pocheville, O. Rey, B. Pujol, and S. Blanchet. 2019. Epigenetically facilitated mutational
 assimilation: epigenetics as a hub within the inclusive evolutionary synthesis. Biol Rev 94:259–282.
- de Montaudouin X., I. Paul-Pont, C. Lambert, P. Gonzalez, N. Raymond, F. Jude, A. Legeay, et al. 2010.
 Bivalve population health: multistress to identify hot spots. Mar Pollut Bull 60:1307–1318.
- Debes P.V., M. Visse, B. Panda, P. Ilmonen, and A. Vasemägi. 2016. Is telomere length a molecular marker
 of past thermal stress in wild fish? Mol Ecol 25:5412–5424.
- Devaux A., S. Bony, S. Plénet, P. Sagnes, S. Segura, R. Suaire, Morgane Novak, et al. 2015. Field evidence of
 reproduction impairment through sperm DNA damage in the fish nase (Chondrostoma nasus) in
 anthropized hydrosystems. Aquat Toxicol 169:113–122.
- Du S.N.N., E.S. McCallum, M. Vaseghi-Shanjani, J.A. Choi, T.R. Warriner, S. Balshine, and G.R. Scott. 2018.
 Metabolic Costs of Exposure to Wastewater Effluent Lead to Compensatory Adjustments in
 Respiratory Physiology in Bluegill Sunfish. Environ Sci Technol 52:801–811.
- 542 Dupoué A., A. Rutschmann, J.F. Le Galliard, J. Clobert, F. Angelier, C. Marciau, S. Ruault, et al. 2017. Shorter
 543 telomeres precede population extinction in wild lizards. Sci Rep 7:16976.
- Eriksen M.S., M. Bakken, A. Espmark, B.O. Braastad, and R. Salte. 2006. Prespawning stress in farmed
 Atlantic salmon Salmo salar: maternal cortisol exposure and hyperthermia during embryonic

- 546 development affect offspring survival, growth and incidence of malformations. J Fish Biol 69:114– 547 129.
- Eriksen M.S., å. Espmark, B.O. Braastad, R. Salte, and M. Bakken. 2007. Long-term effects of maternal
 cortisol exposure and mild hyperthermia during embryogeny on survival, growth and morphological
 anomalies in farmed Atlantic salmon Salmo salar offspring. J Fish Biol 70:462–473.
- Evrard E., A. Devaux, S. Bony, J. Cachot, G. Charrier, L. Quiniou, and J. Laroche. 2013. Responses of juvenile
 European flounder (Platichthys flesus) to multistress in the Vilaine estuary, during a 6-month
 survey. Environ Sci Pollut Res 20:676–689.
- Feil R. and M.F. Fraga. 2012. Epigenetics and the environment: emerging patterns and implications. Nat Rev
 Genet 13:97–109.
- Fitzpatrick S.W. and B.N. Reid. Does gene flow aggravate or alleviate maladaptation to environmental stress
 in small populations? Evol Appl in press.
- Folt C.L., C.Y. Chen, M.V. Moore, and J.L. Burnaford. 1999. Synergism and antagonism among multiple
 stressors.
- Gandar A. 2015. Réponse aux stress multiples chez les poissons : effets croisés de la température et des
 cocktails de pesticides. Toulouse 3.
- Gandar A., S. Jean, J. Canal, N. Marty-Gasset, F. Gilbert, and P. Laffaille. 2015. Multistress effects on goldfish
 (Carassius auratus) behavior and metabolism. Environ Sci Pollut Res.
- Gandar A., P. Laffaille, C. Canlet, M. Tremblay-Franco, R. Gautier, A. Perrault, L. Gress, et al. 2017a. Adaptive
 response under multiple stress exposure in fish: From the molecular to individual level.
 Chemosphere 188:60–72.
- Gandar A., P. Laffaille, N. Marty-Gasset, D. Viala, C. Molette, and S. Jean. 2017b. Proteome response of fish
 under multiple stress exposure: Effects of pesticide mixtures and temperature increase. Aquat
 Toxicol 184:61–77.
- 570 Geist J. 2011. Integrative freshwater ecology and biodiversity conservation. Ecol Indic 11:1507–1516.
- Ghalambor C.K., J.K. McKAY, S.P. Carroll, and D.N. Reznick. 2007. Adaptive versus non-adaptive phenotypic
 plasticity and the potential for contemporary adaptation in new environments. Funct Ecol 21:394–
 407.
- 574 Ghanizadeh Kazerouni E., C.E. Franklin, and F. Seebacher. 2017. Parental exposure modulates the effects of 575 UV-B on offspring in guppies. Funct Ecol 31:1082–1090.
- Gonzalez P. and F. Pierron. 2015. Chapter 8 Omics in Aquatic Ecotoxicology: The Ultimate Response to
 Biological Questions? Pp. 183–203 in C. Amiard-Triquet, J.-C. Amiard, and C. Mouneyrac eds. Aquat
 Ecotoxicol. Academic Press.
- Gordon C. 2005. Temperature and Toxicology: An Integrative, Comparative, and Environmental Approach.
 CRC Press.
- Hagenaars A., L. Vergauwen, D. Benoot, K. Laukens, and D. Knapen. 2013. Mechanistic toxicity study of
 perfluorooctanoic acid in zebrafish suggests mitochondrial dysfunction to play a key role in PFOA
 toxicity. Chemosphere 91:844–856.

- Hahn M.E. 2011. Mechanistic Research in Aquatic Toxicology: Perspectives and Future Directions. Aquat
 Toxicol Amst Neth 105:67–71.
- Hamilton P.B., G. Rolshausen, T.M. Uren Webster, and C.R. Tyler. 2017. Adaptive capabilities and fitness
 consequences associated with pollution exposure in fish. Philos Trans R Soc B Biol Sci
 372:20160042.
- Hendry A.P. 2016. Key Questions on the Role of Phenotypic Plasticity in Eco-Evolutionary Dynamics. J Hered
 107:25–41.
- Hendry A.P., T.J. Farrugia, and M.T. Kinnison. 2008. Human influences on rates of phenotypic change in wild
 animal populations. Mol Ecol 17:20–29.
- Hendry A.P., K.M. Gotanda, and E.I. Svensson. 2017. Human influences on evolution, and the ecological and
 societal consequences. Philos Trans R Soc B Biol Sci 372:20160028.
- Herborn K.A., B.J. Heidinger, W. Boner, J.C. Noguera, A. Adam, F. Daunt, and P. Monaghan. 2014. Stress
 exposure in early post-natal life reduces telomere length: an experimental demonstration in a long lived seabird. Proc R Soc B Biol Sci 281:20133151–20133151.
- Herráez M.P., J. Ausió, A. Devaux, S. González-Rojo, C. Fernández-Díez, S. Bony, N. Saperas, et al. 2017.
 Paternal contribution to development: Sperm genetic damage and repair in fish. Aquaculture,
 Recent advances in fish gametes and embryo research 472:45–59.
- Hochberg M.E. and R.J. Noble. 2017. A framework for how environment contributes to cancer risk. (K.
 Lafferty, ed.)Ecol Lett 20:117–134.
- Holmstrup M., A.-M. Bindesbøl, G.J. Oostingh, A. Duschl, V. Scheil, H.-R. Köhler, S. Loureiro, et al. 2010.
 Interactions between effects of environmental chemicals and natural stressors: A review. Sci Total
 Environ, Cumulative Stressors Risk assessment of mixtures of chemicals and combinations of
 chemicals and natural stressors 408:3746–3762.
- Iwama G.K., P.T. Thomas, R.B. Forsyth, and M.M. Vijayan. 1998. Heat shock protein expression in fish. Rev
 Fish Biol Fish 8:35–56.
- Jackson M.C., C.J.G. Loewen, R.D. Vinebrooke, and C.T. Chimimba. 2015. Net effects of multiple stressors in
 freshwater ecosystems: a meta-analysis. Glob Change Biol.
- Jacquin L., C. Dybwad, G. Rolshausen, A.P. Hendry, and S.M. Reader. 2017. Evolutionary and immediate
 effects of crude-oil pollution: depression of exploratory behavior across populations of Trinidadian
 guppies. Anim Cogn 20:97–108.
- Jacquin L., A. Gandar, M. Aguirre-Smith, A. Perrault, M.L. Hénaff, L.D. Jong, S. Paris-Palacios, et al. 2019.
 High temperature aggravates the effects of pesticides in goldfish. Ecotoxicol Environ Saf 172:255–
 264.
- Kassahn K.S., R.H. Crozier, H.O. Pörtner, and M.J. Caley. 2009. Animal performance and stress: responses
 and tolerance limits at different levels of biological organisation. Biol Rev Camb Philos Soc 84:277–
 292.
- Kaufmann J., T.L. Lenz, M. Milinski, and C. Eizaguirre. 2014. Experimental parasite infection reveals costs
 and benefits of paternal effects. Ecol Lett 17:1409–1417.

- 622 Kawecki T.J. and D. Ebert. 2004. Conceptual issues in local adaptation. Ecol Lett 7:1225–1241.
- Kennedy C.J. and P.S. Ross. 2012. Stress syndromes: Heightened bioenergetic costs associated with
 contaminant exposure at warm temperatures in teleosts. Integr Environ Assess Manag 8:202–204.
- Killen S.S., S. Marras, N.B. Metcalfe, D.J. McKenzie, and P. Domenici. 2013. Environmental stressors alter
 relationships between physiology and behavior. Trends Ecol Evol 28:651–658.
- 627 Kitano H. 2002. Systems Biology: A Brief Overview. Science 295:1662–1664.
- Komoroske L.M., K.M. Jeffries, R.E. Connon, J. Dexter, M. Hasenbein, C. Verhille, and N.A. Fangue. 2016.
 Sublethal salinity stress contributes to habitat limitation in an endangered estuarine fish. Evol Appl 9:963–981.
- Kullgren A., F. Jutfelt, R. Fontanillas, K. Sundell, L. Samuelsson, K. Wiklander, P. Kling, et al. 2013. The impact
 of temperature on the metabolome and endocrine metabolic signals in Atlantic salmon (Salmo
 salar). Comp Biochem Physiol A Mol Integr Physiol 164:44–53.
- Lange K., A. Bruder, C.D. Matthaei, J. Brodersen, and R.A. Paterson. 2018. Multiple-stressor effects on
 freshwater fish: Importance of taxonomy and life stage. Fish Fish 19:974–983.
- Lavergne E., N. Pedron, I. Calves, G. Claireaux, D. Mazurais, J. Zambonino-Infante, N. Le Bayon, et al. 2015.
 Does the chronic chemical contamination of a European flounder population decrease its thermal
 tolerance? Mar Pollut Bull, The English Channel and it's catchments: Status and Responses to
 Contaminants 95:658–664.
- Li C., P. Li, Y.M. Tan, S.H. Lam, E.C.Y. Chan, and Z. Gong. 2016. Metabolomic Characterizations of Liver Injury
 Caused by Acute Arsenic Toxicity in Zebrafish. PLOS ONE 11:e0151225.
- Liess M., K. Foit, S. Knillmann, R.B. Schäfer, and H.-D. Liess. 2016. Predicting the synergy of multiple stress
 effects. Sci Rep 6.
- López-Olmeda J.F. and F.J. Sánchez-Vázquez. 2011. Thermal biology of zebrafish (Danio rerio). J Therm Biol
 36:91–104.
- Loria A., M.E. Cristescu, and A. Gonzalez. Mixed evidence for adaptation to environmental pollution. EvolAppl in press.
- Lü A., X. Hu, Y. Wang, X. Shen, X. Li, A. Zhu, J. Tian, et al. 2014. iTRAQ analysis of gill proteins from the
 zebrafish (Danio rerio) infected with Aeromonas hydrophila. Fish Shellfish Immunol 36:229–239.
- Lurling M. and M. Scheffer. 2007. Info-disruption: pollution and the transfer of chemical information
 between organisms. Trends Ecol Evol 22:374–379.
- 652 Lushchak V.I. 2011. Environmentally induced oxidative stress in aquatic animals. Aquat Toxicol 101:13–30.
- Malek R.L., H. Sajadi, J. Abraham, M.A. Grundy, and G.S. Gerhard. 2004. The effects of temperature
 reduction on gene expression and oxidative stress in skeletal muscle from adult zebrafish. Comp
 Biochem Physiol Toxicol Pharmacol CBP 138:363–373.

Marcogliese D., L. Brambilla, F. Gagné, and A. Gendron. 2005. Joint effects of parasitism and pollution on oxidative stress biomarkers in yellow perch Perca flavescens. Dis Aquat Organ 63:77–84.

- Marcogliese D.J. and M. Pietrock. 2011. Combined effects of parasites and contaminants on animal health:
 parasites do matter. Trends Parasitol 27:123–130.
- Matthaei C. and K. Lange. 2016. Multiple stressor effects on freshwater fish: a review and meta-analysis
 (Cambridge University Press.). Closs G.p., Krkodsek and Olden J.D.
- McBryan T.L., K. Anttila, T.M. Healy, and P.M. Schulte. 2013. Responses to temperature and hypoxia as
 interacting stressors in fish: implications for adaptation to environmental change. Integr Comp Biol
 53:648–659.
- McLennan D., J.D. Armstrong, D.C. Stewart, S. Mckelvey, W. Boner, P. Monaghan, and N.B. Metcalfe. 2017.
 Shorter juvenile telomere length is associated with higher survival to spawning in migratory Atlantic
 salmon. Funct Ecol 31:2070–2079.
- 668 Merilä J. 2015. Evolutionary biology: Perplexing effects of phenotypic plasticity. Nature.
- Merilä J. and A.P. Hendry. 2014. Climate change, adaptation, and phenotypic plasticity: the problem and
 the evidence. Evol Appl 7:1–14.
- Metzger D.C.H. and P.M. Schulte. 2016. Maternal stress has divergent effects on gene expression patterns
 in the brains of male and female threespine stickleback. Proc R Soc B Biol Sci 283:20161734.
- Metzger D.C.H. and P.M. Schulte. 2017. Persistent and plastic effects of temperature on DNA methylation
 across the genome of threespine stickleback (Gasterosteus aculeatus). Proc R Soc B 284:20171667.
- Mirbahai L. and J.K. Chipman. 2014. Epigenetic memory of environmental organisms: A reflection of
 lifetime stressor exposures. Mutat Res Toxicol Environ Mutagen, Epigenetics and chemical safety
 764–765:10–17.
- Morgan M.J., C.E. Wilson, and L.W. Crim. 1999. The effect of stress on reproduction in Atlantic cod. J Fish
 Biol 54:477–488.
- 680 Mousseau T. 1998. The adaptive significance of maternal effects. Trends Ecol Evol 13:403–407.
- Naour S., B.M. Espinoza, J.E. Aedo, R. Zuloaga, J. Maldonado, M. Bastias-Molina, H. Silva, et al. 2017.
 Transcriptomic analysis of the hepatic response to stress in the red cusk-eel (Genypterus chilensis):
 Insights into lipid metabolism, oxidative stress and liver steatosis. PloS One 12:e0176447.
- Narum S.R., N.R. Campbell, K.A. Meyer, M.R. Miller, and R.W. Hardy. 2013. Thermal adaptation and
 acclimation of ectotherms from differing aquatic climates. Mol Ecol 22:3090–3097.
- Nikinmaa M. and K.T. Rytkönen. 2011. Functional genomics in aquatic toxicology—Do not forget the
 function. Aquat Toxicol, Jubileum 105:16–24.
- Noyes P.D., M.K. McElwee, H.D. Miller, B.W. Clark, L.A. Van Tiem, K.C. Walcott, K.N. Erwin, et al. 2009. The
 toxicology of climate change: Environmental contaminants in a warming world. Environ Int 35:971–
 986.
- Okamura B., H. Hartikainen, H. Schmidt-Posthaus, and T. Wahli. 2011. Life cycle complexity, environmental
 change and the emerging status of salmonid proliferative kidney disease: PKD as an emerging
 disease of salmonid fish. Freshw Biol 56:735–753.

- Ortiz-Villanueva E., J. Jaumot, R. Martínez, L. Navarro-Martín, B. Piña, and R. Tauler. 2018. Assessment of
 endocrine disruptors effects on zebrafish (Danio rerio) embryos by untargeted LC-HRMS
 metabolomic analysis. Sci Total Environ 635:156–166.
- Oziolor E.M., J.W. Bickham, and C.W. Matson. 2017a. Evolutionary toxicology in an omics world. Evol Appl
 10:752–761.
- Oziolor E.M., E. Bigorgne, L. Aguilar, S. Usenko, and C.W. Matson. 2014. Evolved resistance to PCB- and
 PAH-induced cardiac teratogenesis, and reduced CYP1A activity in Gulf killifish (Fundulus grandis)
 populations from the Houston Ship Channel, Texas. Aquat Toxicol 150:210–219.
- Oziolor E.M., K.D. Schamphelaere, and C.W. Matson. 2016. Evolutionary toxicology: Meta-analysis of
 evolutionary events in response to chemical stressors. Ecotoxicology 25:1858–1866.
- Pascoe D. and P. Cram. 1977. The effect of parasitism on the toxicity of cadmium to the three-spined
 stickleback, Gasterosteus aculeatus L. J Fish Biol 10:467–472.
- Pédron N., S. Artigaud, J.-L.Z. Infante, N. Le Bayon, G. Charrier, V. Pichereau, and J. Laroche. 2017a.
 Proteomic responses of European flounder to temperature and hypoxia as interacting stressors:
 Differential sensitivities of populations. Sci Total Environ 586:890–899.
- Pédron N., J. Le Du, G. Charrier, J.-L. Zambonino-Infante, N. Le Bayon, R.P. Vasconcelos, V.F. Fonseca, et al.
 2017b. Contrasting patterns of energy metabolism in northern vs southern peripheral European
 flounder populations exposed to temperature rising and hypoxia. Mar Environ Res.
- Perkins E.J., J.K. Chipman, S. Edwards, T. Habib, F. Falciani, R. Taylor, G.V. Aggelen, et al. 2011. Reverse
 engineering adverse outcome pathways. Environ Toxicol Chem 30:22–38.
- Pierron F., S. Bureau du Colombier, A. Moffett, A. Caron, L. Peluhet, G. Daffe, P. Lambert, et al. 2014.
 Abnormal Ovarian DNA Methylation Programming during Gonad Maturation in Wild Contaminated
 Fish. Environ Sci Technol 48:11688–11695.
- Piou C. and E. Prévost. 2013. Contrasting effects of climate change in continental vs. oceanic environments
 on population persistence and microevolution of Atlantic salmon. Glob Change Biol 19:711–723.
- Polard T., S. Jean, L. Gauthier, C. Laplanche, G. Merlina, J.M. Sánchez-Pérez, and E. Pinelli. 2011. Mutagenic
 impact on fish of runoff events in agricultural areas in south-west France. Aquat Toxicol 101:126–
 134.
- Rahel F.J. and J.D. Olden. 2008. Assessing the Effects of Climate Change on Aquatic Invasive Species.
 Conserv Biol 22:521–533.
- Raphael S., P.-L. Mélissa, B. Aurélien, V. Emmanuelle, P. Jean Marc, B. Sylvie, D. Alain, et al. 2014. Kinetic
 response of a genotoxicity biomarker in the three-spined stickleback and implication for
 environmental monitoring. Ecotoxicol Environ Saf 102:6–11.
- Räsänen K. and L.E.B. Kruuk. 2007. Maternal effects and evolution at ecological time-scales. Funct Ecol
 21:408–421.
- Reichert S. and A. Stier. 2017. Does oxidative stress shorten telomeres in vivo? A review. Biol Lett 13.

- Reid N.M., D.A. Proestou, B.W. Clark, W.C. Warren, J.K. Colbourne, J.R. Shaw, S.I. Karchner, et al. 2016. The
 genomic landscape of rapid repeated evolutionary adaptation to toxic pollution in wild fish. Science
 354:1305–1308.
- Rey O., E. Danchin, M. Mirouze, C. Loot, and S. Blanchet. 2016. Adaptation to Global Change: A
 Transposable Element–Epigenetics Perspective. Trends Ecol Evol 31:514–526.
- Rolshausen G., S. Muttalib, R. Kaeuffer, K.B. Oke, D. Hanson, and A.P. Hendry. 2015. When maladaptive
 gene flow does not increase selection. Evol Int J Org Evol 69:2289–2302.
- Salinas S. and S.B. Munch. 2012. Thermal legacies: transgenerational effects of temperature on growth in a
 vertebrate. Ecol Lett 15:159–163.
- Sanchez-Ramirez C., V.M. Vidal-Martinez, M.L. Aguirre-Macedo, R.P. Rodriguez-Canul, G. Gold-Bouchot, and
 B. Sures. 2007. Cichlidogyrus sclerosus (Monogenea: Ancyrocephalinae) and Its Host, the Nile
 Tilapia (Oreochromis niloticus), as Bioindicators of Chemical Pollution. J Parasitol 93:1097–1106.
- Santos R., A. Joyeux, A. Besnard, C. Blanchard, C. Halkett, S. Bony, W. Sanchez, et al. 2017. An integrative
 approach to assess ecological risks of surface water contamination for fish populations. Environ
 Pollut 220:588–596.
- Schinegger R., M. Palt, P. Segurado, and S. Schmutz. 2016a. Untangling the effects of multiple human
 stressors and their impacts on fish assemblages in European running waters. Sci Total Environ
 573:1079–1088.
- Schinegger R., M. Pucher, C. Aschauer, and S. Schmutz. 2018. Configuration of multiple human stressors
 and their impacts on fish assemblages in Alpine river basins of Austria. Sci Total Environ 616–
 617:17–28.
- Schinegger R., C. Trautwein, A. Melcher, and S. Schmutz. 2012. Multiple human pressures and their spatial
 patterns in European running waters. Water Environ J 26:261–273.

Schmitz M., J. Douxfils, S.N.M. Mandiki, C. Morana, S. Baekelandt, and P. Kestemont. 2016. Chronic hyperosmotic stress interferes with immune homeostasis in striped catfish (Pangasianodon hypophthalmus, S.) and leads to excessive inflammatory response during bacterial infection. Fish Shellfish Immunol 55:550–558.

- Schreck C.B., L. Tort, A.P. Farrell, and C.J. Brauner, eds. 2016. Biology of stress in fish. Fish Physiol, Fish
 physiology.
- Schulte P.M. 2014. What is environmental stress? Insights from fish living in a variable environment. J Exp
 Biol 217:23–34.
- Segner H., M. Schmitt-Jansen, and S. Sabater. 2014. Assessing the impact of multiple stressors on aquatic
 biota: the receptor's side matters. Environ Sci Technol 48:7690–7696.
- Simide R., F. Angelier, S. Gaillard, and A. Stier. 2016. Age and Heat Stress as Determinants of Telomere
 Length in a Long-Lived Fish, the Siberian Sturgeon. Physiol Biochem Zool PBZ 89:441–447.
- Slaninova A., M. Smutna, H. Modra, and Z. Svobodova. 2009. A review: oxidative stress in fish induced by
 pesticides. Neuro Endocrinol Lett 30 Suppl 1:2–12.

- Sokolova I.M. 2013. Energy-Limited Tolerance to Stress as a Conceptual Framework to Integrate the Effects
 of Multiple Stressors. Integr Comp Biol 53:597–608.
- Sokolova I.M., M. Frederich, R. Bagwe, G. Lannig, and A.A. Sukhotin. 2012. Energy homeostasis as an
 integrative tool for assessing limits of environmental stress tolerance in aquatic invertebrates. Mar
 Environ Res 79:1–15.
- Thilakaratne I.D.S.I.P., J.D. McLaughlin, and D.J. Marcogliese. 2007. Effects of pollution and parasites on
 biomarkers of fish health in spottail shiners Notropis hudsonius (Clinton). J Fish Biol 71:519–538.
- Topal A., M. Atamanalp, E. Oruç, M. Kırıcı, and E.M. Kocaman. 2014. Apoptotic effects and glucose-6 phosphate dehydrogenase responses in liver and gill tissues of rainbow trout treated with
 chlorpyrifos. Tissue Cell 46:490–496.
- Tudorache C., R. Blust, and G. De Boeck. 2008. Social interactions, predation behavior and fast start
 performance are affected by ammonia exposure in brown trout (Salmo trutta L.). Aquat Toxicol
 Amst Neth 90:145–153.
- Valavanidis A., T. Vlahogianni, M. Dassenakis, and M. Scoullos. 2006. Molecular biomarkers of oxidative
 stress in aquatic organisms in relation to toxic environmental pollutants. Ecotoxicol Environ Saf
 64:178–189.
- Vandegehuchte M.B. and C.R. Janssen. 2014. Epigenetics in an ecotoxicological context. Mutat Res Genet
 Toxicol Environ Mutagen 764–765:36–45.
- Vittecoq M., M. Giraudeau, T. Sepp, D.J. Marcogliese, M. Klaassen, F. Renaud, B. Ujvari, et al. 2018. Turning
 natural adaptations to oncogenic factors into an ally in the war against cancer. Evol Appl 11:836–
 844.
- Wang Y., L. Lv, Y. Yu, G. Yang, Z. Xu, Q. Wang, and L. Cai. 2017. Single and joint toxic effects of five selected
 pesticides on the early life stages of zebrafish (Denio rerio). Chemosphere 170:61–67.
- Weber G.J., M.S. Sepúlveda, S.M. Peterson, S.S. Lewis, and J.L. Freeman. 2013. Transcriptome Alterations
 Following Developmental Atrazine Exposure in Zebrafish Are Associated with Disruption of
 Neuroendocrine and Reproductive System Function, Cell Cycle, and Carcinogenesis. Toxicol Sci
 132:458–466.
- Whitehead A., B.W. Clark, N.M. Reid, M.E. Hahn, and D. Nacci. 2017. When evolution is the solution to
 pollution: Key principles, and lessons from rapid repeated adaptation of killifish (Fundulus
 heteroclitus) populations. Evol Appl 10:762–783.
- Woodward G., D.M. Perkins, and L.E. Brown. 2010. Climate change and freshwater ecosystems: impacts
 across multiple levels of organization. Philos Trans R Soc Lond B Biol Sci 365:2093–2106.
- Xing H., S. Li, Z. Wang, X. Gao, S. Xu, and X. Wang. 2012. Oxidative stress response and histopathological
 changes due to atrazine and chlorpyrifos exposure in common carp. Pestic Biochem Physiol
 103:74–80.
- Yang Y., H. Yu, H. Li, A. Wang, and H. Yu. 2016. Effect of high temperature on immune response of grass
 carp (Ctenopharyngodon idellus) by transcriptome analysis. Fish Shellfish Immunol 58:89–95.

Zhang C., M. Jansen, L.D. Meester, and R. Stoks. 2018. Thermal evolution offsets the elevated toxicity of a contaminant under warming: A resurrection study in Daphnia magna. Evol Appl 11:1425–1436.

Fig 1. Schematic representation of theoretical allocation of energy following the concept of 806 807 metabolic capacity of Fry (1947) and metabolic strategies of Sokolova (2013). Under normal 808 conditions (no stress, on the left), the total energy available (represented by the circle) is first 809 allocated to maintenance and damage repair (standard energy power), then to routine behaviors (locomotion, foraging). The energy surplus is invested in somatic and/or gonadal growth (growth 810 811 and reproduction). Under exposure to a single stressor, a "compensation strategy" response is 812 observed: the energy allocated to maintenance increases, at the expense of routine behaviors, and growth. This stress response usually leads to limited irreversible damage to cells and tissues, and 813 814 low fitness effects. Under multiple stressors exposure, the energy available might be insufficient to 815 cover the costs of maintenance, leading to a "conservation strategy": the stress response is blocked 816 and no reallocation of energy is observed. In this case, high level of irreversible cellular damage is expected, with increased mortality and deleterious effects on fitness (adapted from Sokolova et al. 817 818 2013, Gandar et al. 2015).

819

820 Fig 2: Fish stress responses across levels of biological organisation, from molecules to 821 populations.

822

« Compensation strategy »



Multiple environmental stressors



Adaptation or maladaptation

- **Question 1:** What are the effects of multiple abiotic and biotic stressors across biological levels? Are synergic or antagonistic effects more frequent than previously thought?
- **Current limitations and future prospects:** Multiple stressors can have unexpected synergistic or antagonistic interaction effects on fish depending on the level of biological organization considered. Empirical studies in semi-controlled conditions from molecules to populations will help unravelling such interaction effects.
- **Question 2:** What can explain these interactions effects? Are different metabolic strategies set up under single vs. multiple stressors?
- **Current limitations and future prospects:** The metabolic strategy framework (compensation vs conservation strategy, sensu Sokolowa 2013) is a powerful framework for a better understanding of the high variability of stress responses observed in wild populations. Fish could switch from a "compensation" to a "conservation" metabolic strategy when exposed to multiple stressors, with contrasted effects across biological levels (Fig 1). More empirical studies are needed to test this promising theoretical framework.
- **Question 3**: What are the consequences on individual fitness and population persistence? Are there emerging tools to better link different levels of biological organization, from molecules to populations?
- **Current limitations and future prospects**: Most studies have focused on molecular pathways involved in the stress response while other studies compared individual fitness and population dynamics across environments. Integrated biomarkers and theories linking molecular and population effects could be powerful tools to bridge the gap between small- and large-scale effects of environmental stressors (Fig 2).
- **Question 4**: Why do different populations differ in their sensitivity to stressors? Can local adaptation occur and which evolutionary processes are involved (genetic, plastic, epigenetic effects)?
- **Current limitations and future prospects**: Molecular studies have enhanced our understanding of genetic and epigenetic processes underpinning adaptive processes, but they have been rarely applied to wild populations exposed to stressors. Future studies tracking molecular, genetic and epigenetic changes among stressed and unstressed fish populations while taking into account their evolutionary history will give insight into the evolutionary potential of wild fish populations exposed to multiple global changes.

Graphical abstract

Multiple environmental stressors



Adaptation or maladaptation