



HAL
open science

DYRK1A and cognition: A lifelong relationship

Maria L. Arbones, Aurore Thomazeau, Akiko Nakano-Kobayashi, Masatoshi Hagiwara, Jean M. Delabar

► **To cite this version:**

Maria L. Arbones, Aurore Thomazeau, Akiko Nakano-Kobayashi, Masatoshi Hagiwara, Jean M. Delabar. DYRK1A and cognition: A lifelong relationship. *Pharmacology & Therapeutics*, 2019, 194, pp.199 - 221. <10.1016/j.pharmthera.2018.09.010>. <hal-03484540>

HAL Id: hal-03484540

<https://hal.science/hal-03484540v1>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

P&T #23212

DYRK1A and cognition: A lifelong relationship

Maria L Arbones^a, Aurore Thomazeau^b, Akiko Nakano-Kobayashi^c, Masatoshi Hagiwara^c, Jean M Delabar^d

^a Department of Developmental Biology, Instituto de Biología Molecular de Barcelona, CSIC, and Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), 08028 Barcelona, Spain

^b Picower Institute for Learning and Memory, Massachusetts Institute of Technology, Cambridge, United States

^c Department of Anatomy and Developmental Biology, Graduate School of Medicine, Kyoto University, Kyoto 606-8501, Japan

^d INSERM U1127, CNRS UMR 7225, Sorbonne Universités, UPMC Univ Paris 06 UMRS 1127, Institut du Cerveau et de la Moelle épinière, ICM, Paris, France

Abstract:

The dosage of the serine threonine kinase DYRK1A is critical in the central nervous system (CNS) during development and aging. This review analyzes the functions of this kinase by considering its interacting partners and pathways. The role of DYRK1A in controlling the differentiation of prenatal newly formed neurons is presented separately from its role at the pre- and post-synaptic levels in the adult CNS; its effects on synaptic plasticity are also discussed. Because this kinase is positioned at the crossroads of many important processes, dosage medication errors in this protein produce devastating effects arising from DYRK1A deficiency, such as in MRD7, an autism spectrum disorder, or from DYRK1A excess, such as in Down syndrome. Effects of errors have been shown in various animal models including *Drosophila*, zebrafish, and mice. Dysregulation of DYRK1A levels also occurs in neurodegenerative diseases such as Alzheimer's and Parkinson's diseases. Finally, this review describes inhibitors that have been assessed *in vivo*. Accurate targeting of DYRK1A levels in the brain, with either inhibitors or activators, is a future research challenge.

Keywords:

Neurogenesis

Dendrites

MRD7

Alzheimer's disease

Parkinson's disease

Abbreviations:

APP, β -amyloid precursor protein

ASF1, alternative splicing factor 1

CGMC, group of kinases: CDK, CDKL, CK2, CLK, DYRK, GSK, MAPK, RCK, SRPK

CREB1, cAMP responsive element-binding protein 1

DREAM, dimerization partner, RB-like, E2F, and multivulval class B

DS, Down syndrome

DYRK1A, dual-specificity tyrosine phosphorylation regulated kinase 1A

EGCG, epigallocatechin gallate

EGFR, epidermal growth factor receptor

EPSP, excitatory postsynaptic potential

GAD1, glutamate decarboxylase 1

HSA21, human chromosome 21

HIP1, Huntingtin-interacting protein 1

IPSC, induced pluripotent stem cell

Mnb, minibrain (*Drosophila melanogaster*)

MRD7, mental retardation, autosomal dominant 7

PEST, peptide sequence rich in proline (P), glutamic acid (E), serine (S), and threonine (T);

Ser, serine

SGZ, subgranular zone

SNCA, α -synuclein

SPRED1, sprouty-related EVH1 domain-containing protein 1

STAT1, signal transducer and activator of transcription 1

SYNJ1, synaptojanin 1

Thr, threonine

Tyr, tyrosine

SVZ, subventricular zone

VGLUT1, vesicular glutamate transporter 1

VZ, ventricular zone

Table of Contents

1. Introduction		4
2. Protein structure and interactors	5	
3. Neural numbers and organ size		13
4. Synaptogenesis and neuronal functions		19
5. DYRK1A and neurotransmission		23
6. DYRK1A deficit, autism MRD7		27
7. DYRK1A and Down syndrome		30
8. Neurodegenerative diseases		36
9. Inhibitors		42
10. Conclusion		46
Disclosure/Conflict of interest		
References		55

1. Introduction

Protein kinases are vital for converting extracellular signals into biological responses. Functional alterations in kinases may directly contribute to age-dependent neuronal dysfunctions. Various kinases are positive regulators for brain development and memory function, and altered kinase signaling pathways can provoke memory disturbances. Many kinases associated with synaptic function are also age sensitive and are involved in neurodegenerative processes. Protein kinases are attractive drug targets because their ATP binding pocket is an excellent binding site for inhibitors, and many kinases are pivotal in pathological processes.

DYRK1A (dual-specificity tyrosine phosphorylation regulated kinase 1A), also designated as Minibrain (MNB) on the basis of its *Drosophila melanogaster* ortholog, is a protein kinase encoded on human chromosome 21 (HSA21). Compelling data implicate DYRK1A in the regulation of different cellular processes involved in brain development and function—ranging from early embryogenesis through late aging. Herein, we summarize existing data and highlight the roles of DYRK1A in healthy and affected brains. The molecular and physiological

functions of DYRK1A are described in parts 2-5. We present the phenotypic consequences of DYRK1A genetic alterations in mental retardation autosomal dominant 7 (MRD7) (Part 6) and trisomy 21 (Down syndrome [DS]) (Part 7), and we discuss the DYRK1A-mediated functions that are associated with neurodegenerative diseases including Alzheimer's disease (AD) and Parkinson's disease (PD) (Part 8). Finally, the role of DYRK1A in several human diseases makes DYRK1A a potential target for therapeutic drugs; the use of several inhibitors is described in part 9. Because of the functional and molecular complexity of the DYRK1A protein kinase, the advantages and drawbacks of its possible use as a drug target for DS, MRD7, and AD must be carefully considered.

2 Protein structure and interactors

2.1 Expression patterns

DYRK1A is encoded on HSA21 in 21q22.2 (Guimera, et al., 1996; Hattori, et al., 2000; Patil, et al., 1995). The gene comprises 151 kb and 15 exons (Ensemble release 90); it encodes two main protein isoforms of 763 and 754 amino acids. DYRK1A belongs to the DYRK family of dual-specificity protein kinases (CMGC group: DYRK family: DYRK subfamily). The kinase domain is located centrally in the primary structure of the protein. In mammals, the DYRK subfamily members (DYRK1A, DYRK1B, DYRK2, DYRK3, and DYRK4) share a conserved motif N-terminal to the kinase domain known as the DYRK homology (DH)-box. DYRK1A also harbors a functional bipartite nuclear localization signal (NLS) N-terminal to the DH-box, a second NLS between subdomains X and XI within the kinase domain, a C-terminal PEST motif, and a polyhistidine tract that acts as a nuclear speckle targeting signal (Alvarez, Estivill, & de la Luna, 2003; Aranda, Laguna, & de la Luna, 2011; W. Becker & Joost, 1999) (Figure 1A). DYRK1A and its closest family member DYRK1B are 73.9% identical (89.4% similar), and their kinase domains are 85.0% identical (95.6% similar). They share no sequence similarity within their C-terminal domains. *DYRK1A* is ubiquitously expressed in human and mouse tissues,

but *DYRK1B* exhibits a more restricted pattern, with the highest mRNA levels in the testes and muscles.

DYRK1A and other human kinases from the CMGC group share a similar fold, the kinase domain, comprising an N-terminal lobe (N-lobe) with five antiparallel β -strands and a conserved regulatory α C-helix and a larger C-terminal (C-lobe) consisting of α -helices. The N- and C-lobes are connected by a hinge region, an important part of the ATP-pocket. The “gatekeeper” is a vital residue in the ATP pocket responsible for the selectivity of inhibitors. DYRKs have phenylalanines as gatekeepers, F238 in DYRK1A and F190 in DYRK1B, located at the beginning of their hinge regions. These features have been described in crystallographic studies (Alexeeva, Aberg, Engh, & Rothweiler, 2015; Anderson, et al., 2013; Falke, et al., 2015; Ogawa, et al., 2010; Rothweiler, et al., 2016; Tahtouh, et al., 2012) mostly performed on complexes of the N-terminal fragment of DYRK1A, including the kinase domain, and competitive inhibitors targeting the ATP pocket (Figure 1B). The functional specificity of each DYRK family member likely arises from the distinct C-terminal domain of that member. These domains have only recently been considered in crystallographic studies.

DYRKs are an evolutionarily conserved family, and homologs have been found in yeast, for example, Yak1p from *Schizosaccharomyces pombe* (Garrett & Broach, 1989), and in nematodes, for example, the two kinases MBK-1 and MBK-2 (homologs of the MNB kinase) from *Caenorhabditis elegans* (Raich, et al., 2003). MNB dysfunction leads to reduced optic lobe size in the brain of adult flies, thus highlighting the role of MNB in neurogenesis (Tejedor, et al., 1995). In zebrafish, gene dosage modifications of *dyrk1aa* induce alterations in the brain development. (O. H. Kim, et al., 2017). Comparison of the modern human genome with Neanderthal and Denisovan genomes identified modern human *DYRK1A* variants, thus suggesting that this gene was influenced by late selective pressures (Mozzi, et al., 2017).

On the basis of its RNA expression profile, *Dyrk1a* is ubiquitous during rodent development but has stronger reactivity in central nervous system (CNS) regions (Okui, et al., 1999; Rahmani, Lopes, Rachidi, & Delabar, 1998). The *Dyrk1a* gene in chick and mouse embryos is

expressed in proliferative regions of the nervous system at a very early stage during development, before the onset of neurogenesis. Expression of the *Dyrk1a* gene in the developing CNS shows a dynamic spatio-temporal pattern, and the protein is detected in cycling neurogenic progenitors and in differentiated neurons (Hammerle, Elizalde, & Tejedor, 2008; Hammerle, et al., 2002). Immunofluorescence studies performed on the developing brain and retina of the mouse indicate that the location of DYRK1A in both neuronal progenitors and differentiated neurons is mainly cytoplasmic (Hammerle, et al., 2002; Laguna, et al., 2008). Expression of DYRK1A during brain development peaks around birth and is maintained at lower levels until adulthood (Okui, et al., 1999). In the adult mouse brain, varying levels of DYRK1A immunoreactivity are detected throughout the neuropil (Marti, et al., 2003). Biochemical fractionation of the mouse brain confirmed that DYRK1A partitions between the nucleus and the cytoplasm (Aranda, Alvarez, Turro, Laguna, & de la Luna, 2008), which correlates with histological studies showing various levels of DYRK1A in the soma and dendrites as well as in the nuclei of different adult brain neuronal types (Hammerle, et al., 2008). Consistently, the distribution of DYRK1A in primary cell cultures revealed the presence of this protein in the cytoplasm and the nucleus of neurons and astrocytes (Marti, et al., 2003). In the human brain, DYRK1A is also found in the cytoplasm and the nucleus of various neuronal types as well as in astrocytes, ependymal, and endothelial cells. The distribution of DYRK1A in these cells indicates cell type- and brain structure-specific patterns of trafficking and utilization of this kinase (Wegiel, et al., 2004).

Two NLSs contribute to DYRK1A nuclear translocation, and the histidine stretch is responsible for the accumulation of the protein in the splicing factor compartment. The C-terminal region of DYRK1A interacts with a brain-specific protein, phytanoyl-CoA hydroxylase-associated protein 1 (PAHX-AP1, also named PHYHIP), which may induce relocalization of nuclear DYRK1A to the cytosol (Bescond & Rahmani, 2005).

When overexpressed in mammalian cells, the DYRK1A protein mainly localizes to the nucleus with a punctate pattern of immunostaining consistent with accumulation in nuclear speckles

(Alvarez, et al., 2003; W. Becker, et al., 1998). Nuclear speckles containing DYRK1A can be detected in neural progenitors and differentiating neurons *in vivo* (Hammerle, et al., 2008; Laguna, et al., 2008). In differentiating neurons (pyramidal neocortical neurons), DYRK1A is mainly found in the growing dendrites and developing neuropil (Hammerle, Elizalde, Galceran, Becker, & Tejedor, 2003).

DYRK1A autophosphorylates on tyrosine, serine, and threonine residues but phosphorylates its substrates only on serine and threonine residues. Autophosphorylation of Tyr 312/321 (754/763 in variants) in the activation loop of DYRK1A is required for the full catalytic activity of the protein (Himpel, et al., 2001). This tyrosine autophosphorylation is an intramolecular reaction that takes place during or immediately after translation (Lochhead, Sibbet, Morrice, & Cleghon, 2005). This reaction is due to an intrinsic ability of the catalytic domain of DYRK1A and does not require chaperone proteins (Gockler, et al., 2009). Comparative analysis of inhibitors suggests that during DYRK1A autoactivation, tyrosine autophosphorylation in the activation loop stabilizes a conformation of the catalytic domain with enhanced serine/threonine kinase activity without disabling tyrosine phosphorylation (Walte, et al., 2013). Further, DYRK1A autophosphorylates serine residues such as Ser520, which modulates binding to 14-3-3 proteins and kinase activity (Alvarez, Altafaj, Aranda, & de la Luna, 2007). Mass spectrometry analysis disclosed different phosphorylation patterns according to the intracellular distribution of DYRK1A and supports the hypothesis that compartment-specific functions of DYRK1A may depend on its phosphorylation status (Kaczmariski, et al., 2014).

2.2 Regulation of DYRK1A transcription, translation, and enzymatic activity

Dyrk1a transcription is regulated by multiple proteins. P44 (a short isoform of the p53 start site in exon 4 of p53) binds to the mouse promoter of *Dyrk1a* and activates transcription; by contrast, p53 induces the expression of miR-1246, which, in turn, reduces the level of *Dyrk1a* mRNA (Pehar, Ko, Li, Scrabble, & Puglielli, 2014; Zhang, Liao, Zeng, & Lu, 2011). miR-199b, which is under the control of NFATc, is another microRNA that targets *DYRK1A* in cardiac hypertrophy but has not been investigated in the CNS (da Costa Martins, et al., 2010). The

transcription factor E2F1 modulates the choice between two *Dyrk1a* promoters, which differ strongly depending on reporter gene activity (Maenz, Hekerman, Vela, Galceran, & Becker, 2008). OLIG2, an HSA21 protein, may also regulate *Dyrk1a* gene expression (W. Liu, et al., 2015). Mena (ENAH), which is highly expressed in the developing and adult CNS and is a member of the Ena/VASP protein family, is part of a ribonucleoprotein complex that modulates DYRK1A translation and synthesis, both under steady state conditions and after brain derived neurotrophic factor (BDNF) stimulation (Vidaki, et al., 2017). REST can activate *DYRK1A* transcription through a neuron-restrictive silencer element in the human *DYRK1A* promoter (Lu, et al., 2011).

DYRK1A may also be regulated post-translationally. Dephosphorylation of its activation loop by a phosphatase, yet to be identified, may inactivate its kinase activity. In addition, phosphorylation events outside the activation loop may modulate DYRK1A activity. Phosphorylation of DYRK1A by LAST2, a component of the hippo pathway, enhances the ability of DYRK1A to phosphorylate the DREAM complex subunit LIN52 (Tschop, et al., 2011). DYRK1A also contains a PEST sequence that allows truncation of the protein by Calpain and may increase kinase activity (Jin, et al., 2015). Interacting proteins such as 14-3-3 and SPRED1 may also modulate DYRK1A kinase activity (Alvarez, et al., 2007; Li, Jackson, Yusoff, & Guy, 2010).

2.3 Interactants/substrates

A consensus phosphorylation sequence for DYRK1A serine/threonine kinase activity has been proposed (RPXS/TP) (Himpel, et al., 2000), although some phosphorylation sites do not fit with the consensus, for instance, LTAT(434)P, SF3B1/SAP155, or RPAS(640)V in glycogen synthase (de Graaf, et al., 2004; Skurat & Dietrich, 2004). Analysis of *in vitro* phosphorylated synthetic peptide substrates established the preference of DYRK1A for an arginine residue in the -2 or -3 position and for a proline at the +1 position (Himpel, et al., 2000). Substrates are related to the subcellular localization of DYRK1A (nuclear or cytosolic).

Chromatin regulators and transcription factors are an important group of DYRK1A substrates (Table I and Figure 2), thus suggesting that DYRK1A plays a role in regulating gene expression. DYRK1A is directly associated with transcriptional control: DYRK1A phosphorylates the C-terminal domain (CTD) of RNA polymerase II (RNAPII) at Ser2 and Ser5 (Di Vona, et al., 2015). Phosphorylation of histone H3 by DYRK1A antagonizes transcriptional repression mediated by heterochromatin protein 1 and participates in the activation of a subset of genes, including those encoding cytokines (Jang, Azebi, Soubigou, & Muchardt, 2014). When tested in gene reporter assays, DYRK1A acts as a transcriptional activator of cAMP responsive element-binding protein 1 (CREB1) (E. J. Yang, Ahn, & Chung, 2001), GLI1 (Ehe, et al., 2017; Mao, et al., 2002), forkhead box protein O1 (FOXO1/FKHR) (von Groote-Bidlingmaier, et al., 2003; Woods, Rena, et al., 2001), and androgen receptor-interacting protein 4 (ARIP4/RAD54L2) (Sitz, Tigges, Baumgartel, Khaspekov, & Lutz, 2004) and as an inhibitor of Notch-dependent transcription (Fernandez-Martinez, et al., 2009). DYRK1A is also a negative regulator of NFAT transcription factors in distinct cellular environments and acts by phosphorylating NFATs and retaining them in the cytoplasm (Arron, et al., 2006; Fernandez-Martinez, et al., 2009; Lee, et al., 2009).

DYRK1A is proposed as a regulator of splicing on the basis of the localization of kinase in nuclear speckles (Alvarez, et al., 2003) and the identification of several splicing factors as DYRK1A substrates (de Graaf, et al., 2006). DYRK1A phosphorylation of the alternative splicing factor (ASF) prevents ASF-mediated inclusion of the alternatively spliced exon 10 in tau mRNA (Shi, et al., 2008). There is evidence that DYRK1A phosphorylates other proteins involved in mRNA splicing, including the SR proteins SC35, SRp55, and 9G8, at several serine residues (Ding, et al., 2012; Qian, et al., 2011; Yin, et al., 2012).

DYRK1A inhibits apoptosis by phosphorylating the NAD-dependent deacetylase Sirtuin-1 (SIRT1), an inhibitor of p53 (Guo, Williams, Schug, & Li, 2010), and HIP1 (Kang, Choi, Park, & Chung, 2005), a protein normally associated with Huntingtin. SIRT1 together with HIPPI (HIP1 protein interactor) forms a complex with caspase-8 and induces apoptosis through

components of the extrinsic cell death pathway (Gervais, et al., 2002). Phosphorylation of p53 was demonstrated in neuronal cells in the context of proliferation; DYRK1A-mediated deacetylation did not affect proliferation but instead inhibited cell death. Moreover, DYRK1A restrains the activity of the intrinsic cell death pathway by phosphorylating Thr125 in caspase-9 (CASP9), a phosphorylation event that prevents CASP9 cleavage and subsequent activation of the pathway (Laguna, et al., 2008; Seifert, Allan, & Clarke, 2008).

Increasing evidence indicates that DYRK1A is a negative regulator of cell cycle progression. DYRK1A phosphorylation of LIN52 is required for the assembly of the DREAM complex (Litovchick, Florens, Swanson, Washburn, & DeCaprio, 2011). This complex mediates gene repression during the G0 phase and coordinates periodic gene expression with peaks during the G1/S and G2/M phases (Sadasivam & DeCaprio, 2013). DYRK1A regulates protein levels of regulators of the G1/S phase transition in different ways. It controls the nuclear levels of Cyclin D proteins by phosphorylating a particular threonine residue in Cyclin D1/D3 (CCND1, CCND3), and this promotes Cyclin D nuclear export and degradation of the protein through the ubiquitin-proteasome pathway (J. Y. Chen, Lin, Tsai, & Meyer, 2013; Thompson, et al., 2015). Knockdown of DYRK1A in fibroblast cells greatly increases Cyclin D1 levels and splits cells into two fates, with one subpopulation having an accelerated cell cycle with a significantly shortened G1 duration and the other entering an arrested state by costabilizing Cyclin D1 and the CDK inhibitor p21^{Cip1} (J. Y. Chen, et al., 2013). In addition, DYRK1A regulates p21^{Cip1} at the transcriptional level through phosphorylation of p53 (J. Park, et al., 2010).

2.4 Participation of DYRK1A in other cellular processes and signal transduction pathways

DYRK1A associated with the scaffold protein HAN11 (or DCAF7, WDR68) regulates glycogen synthase, a key enzyme in the regulation of glycogen synthesis by insulin (Skurat & Dietrich, 2004). DYRK1A is also a modifier of signaling mechanisms: it directly acts on receptor tyrosine kinase (RTK) signaling by phosphorylating Sprouty2. Both DYRK1A and Sprouty2 localize to growth cones of nerve terminals (Aranda, et al., 2008). DYRK1A also interacts with SPRED1

and SPRED2, which inhibits the ability of DYRK1A to phosphorylate its substrates by reducing access to its kinase domain (Li, et al., 2010).

DYRK1A contributes to the maintenance of normal circadian clock oscillation by phosphorylating Ser553 in cryptochrome 2 (CRY2), which serves as a priming step for glycogen synthase kinase 3 (GSK3)-mediated phosphorylation, thus leading to proteosomal degradation of CRY2 (Kurabayashi, Hirota, Sakai, Sanada, & Fukada, 2010).

DYRK1A interacts with proteins involved in protein processing and degradation: it phosphorylates presenilin, a component of the gamma secretase complex producing the A β peptide (Ryu, et al., 2010), and it modifies neprilysin (NEP), which cleaves peptides on the amino side of hydrophobic residues and degrades the A β peptide (Kawakubo, Mori, Shirotani, Iwata, & Asai, 2017). Results of a yeast two-hybrid assay showed that DYRK1A is a novel binding partner of parkin (PRKN): DYRK1A phosphorylates parkin at Ser131, a modification that inhibits the E3 ubiquitin ligase activity of parkin (Im & Chung, 2015).

There is strong experimental evidence positioning DYRK1A within the endocytic network. DYRK1A phosphorylates synaptojanin1 (SYNJ1) and modifies its binding with amphiphysin and intersectin (Adayev, Chen-Hwang, Murakami, Wang, & Hwang, 2006; Adayev, Chen-Hwang, Murakami, Wegiel, & Hwang, 2006). In *Drosophila*, MNB-dependent phosphorylation of Synj1 is required for complex endocytic protein interactions and to enhance Synj1 activity *in vivo* (C. K. Chen, et al., 2014). Through phosphorylation of Synj1 at Ser1029, MNB modulates the synaptic vesicle pool size (Geng, Wang, Lee, Chen, & Chang, 2016). DYRK1A was shown to phosphorylate Ser293 of amphiphysin in SY5Y cells (Murakami, et al., 2006). It also phosphorylates Ser857 of dynamin1, and this phosphorylation is dependent on neuronal activity (Xie, et al., 2012). DYRK1A can regulate neural development and synaptic plasticity through phosphorylation of Ser1048 in GluN2A, a subunit of the N-methyl-D-aspartate glutamate receptors (NMDARs) (Grau, et al., 2014). Another target of DYRK1A is α -synuclein (Ser87), which is involved in the control of presynaptic signaling and membrane trafficking (E. J. Kim, et al., 2006).

Cytoskeletal proteins involved in cell structure and trafficking, such as Septin4, a GTPase family member that serves as a scaffold for diverse molecules beneath the plasma membrane, are also DYRK1A targets (Sitz, et al., 2008). In addition, DYRK1A influences the interplay between neuronal Wiskott–Aldrich syndrome protein (N-WASP) and growth factor receptor-bound protein 2 (GRB2), which control the actin skeleton and cytoskeletal architecture. DYRK1A phosphorylation of N-WASP inhibits Arp2/3-complex-mediated actin polymerization (Abekhoukh, et al., 2013; J. Park, et al., 2012; Shin, Guedj, Delabar, & Lubec, 2007). Tau (MAPT), which promotes microtubule assembly and stability, is phosphorylated by DYRK1A (Frost, et al., 2011; Yin, et al., 2012) as well as two huntingtin-interacting proteins, HIP1 and HAP1a, associated with protein trafficking and the cytoskeleton, and these proteins also interact with DCAF7 (Kang, et al., 2005; Z. Xiang, Haroutunian, Ho, Purohit, & Pasinetti, 2006). β -Amyloid precursor protein (APP) is phosphorylated by DYRK1A *in vitro*, and overexpression of DYRK1A increases the phosphorylation of APP on Thr668, which has been reported to modify APP conformation and could alter the role of APP in axonal transport and its processing by γ -secretases (Ryoo, et al., 2008; Salehi, et al., 2006). A summary of DYRK1A substrates/interactors is given in Table I. The relevance of these interactions in brain development and function is discussed in parts 3, 4, and 5.

3 Neural numbers and organ size

The mammalian brain is formed of many neural types. During development, neurons arise from progenitors located in the proliferative regions within the anterior neural tube through a highly regulated process known as neurogenesis. Newborn neurons, which in most cases are produced prenatally, migrate and differentiate to form functional circuits during postnatal development. A proportion of differentiating neurons die by physiological apoptosis to balance the number of various neuron types with their corresponding target cells. Thus, alterations in neurogenesis and/or physiological cell death have an impact on the neuronal component of the brain circuits and, consequently, on its function. In this section, we summarize the current

knowledge of the mechanisms by which DYRK1A/MNB kinases may regulate neural homeostasis and ultimately organ size. Owing to the relevance of the neocortex in cognition, the DYRK1A-mediated activities involved in neurogenesis and astrogliogenesis in this structure are explained in more detail (see schematic representation in Figure 3).

3.1 Neurogenesis

The first evidence implicating DYRK1A/MNB in brain growth was published by Tejedor et al. in 1995. The authors showed that loss-of-function mutations in the *D. melanogaster mnb* gene caused a prominent size reduction in the optic lobes and central brain hemispheres in the adult fly; this reduction is due to the failure of *mnb* mutants to generate sufficient numbers of neurons during postembryonic neurogenesis. Recently, Shaikh et al. (2016) showed that *mnb* is weakly expressed in neuroblasts and that the expression of this gene is upregulated in newborn neurogenic precursors known as ganglion cells (Ceron, Gonzalez, & Tejedor, 2001). MNB in these precursors promotes the expression of the cyclin-dependent kinase inhibitor Dacapo, a homolog of vertebrate p27^{Kip1}, through a mechanism that involves the upregulation of the proneuronal transcription factor Asense and the homeodomain transcription factor Prospero. On the basis of these results and on the phenotype of *mnb* mutants, Shaikh et al. concluded that MNB is necessary for cell cycle exit and terminal differentiation in larval ganglion cells. In these mutants, many of these precursors continue to proliferate instead of exiting the cell cycle and they eventually die by apoptosis.

Drosophila MNB interacts with the adaptor protein Wings Apart (WAP, also known as Riquiqui), and this interaction promotes Yorkie-dependent tissue growth by stimulating phosphorylation-dependent inhibition of the Warts kinase (Degoutin, et al., 2013). Yorkie (YKI; YAP/TAZ in mammals) is the major effector of the Hippo pathway, a universal governor of organ size and tissue homeostasis (Yu, Daniels, Wu, & Wolf, 2015). In addition, WAP interaction with MNB can control organ growth through the downregulation of the transcriptional repressor Capicua (CIC) (L. Yang, et al., 2016), which is a key sensor of RTK signaling in *Drosophila* and mammals (Jimenez, Shvartsman, & Paroush, 2012). There is evidence that MNB, WAP, and

CIC are implicated in a common pathway that controls brain growth (Tejedor, et al., 1995; L. Yang, et al., 2016). Collectively, the studies performed in the fly indicate that in addition to the function of MNB in cell cycle regulation, other MNB-mediated activities may contribute to the brain size defects observed in *mnb* mutant flies.

The microcephaly displayed by human (see part 6), mouse (Fotaki, et al., 2002; Raveau, et al., 2018), and zebrafish (O. H. Kim, et al., 2017) models with loss-of-function mutations in *DYRK1A* indicates that the function of DYRK1A/MNB in brain growth is conserved across evolution. Morphometric studies performed in adult mutant mice with one (*Dyrk1a*^{+/-} mice) or three functional copies of *Dyrk1a* (mBACTgDyrk1a mice) revealed that the effect of DYRK1A on brain growth is dosage dependent and region specific. In these two *Dyrk1a* mutants, neuron numbers in the adult neocortex inversely correlate with *Dyrk1a* gene dosage, whereas the opposite correlation has been observed in other brain regions (Guedj, et al., 2012). This suggests that different DYRK1A-mediated activities are likely contributing to the cell number defects in mBACTgDyrk1a and *Dyrk1a*^{+/-} mice.

There is evidence that DYRK1A, like MNB, inhibits cell cycle progression in neural progenitors by controlling the expression of cell cycle regulatory proteins. In SH-SY5Y neuroblastoma cells, the overexpression of DYRK1A induces cell cycle exit and neuronal differentiation, an effect that correlates with increased phosphorylation of Cyclin D1 on Thr286, which promotes degradation of the protein, and of p27^{Kip1} in Ser10, an event that stabilizes the protein (Soppa, et al., 2014). Forced overexpression of DYRK1A in the chick spinal cord induces cell cycle exit and upregulates the expression of p27^{Kip1} at the transcriptional level (Hammerle, et al., 2011). In immortalized rat embryonic hippocampal progenitor cells, DYRK1A phosphorylates the transcription factor p53 at Ser15, thus leading to increased expression of several p53 target genes, including p21^{CIP1}, and attenuation of proliferation. Impaired proliferation and increased levels of phosphorylated p53 have also been observed in cortical neural precursors derived from a transgenic mouse line that expresses human *DYRK1A* (J. Park, et al., 2010).

Regulation of cell cycle, particularly G1 to S phase progression, is crucial for the generation of

appropriate number of neurons in the developing neocortex (Caviness, Takahashi, & Nowakowski, 1995; Dehay & Kennedy, 2007). These neurons are generated from pluripotent apical progenitors (also named as radial glial progenitors) in the ventricular zone (VZ) surrounding the ventricles or from neurogenic basal progenitors in the subventricular zone (SVZ) (Franco & Muller, 2013). In the mouse embryo, DYRK1A is expressed before the onset of neurogenesis, and the expression is maintained throughout corticogenesis (Franco & Muller, 2013; Hammerle, et al., 2008; Kurabayashi & Sanada, 2013; Najas, et al., 2015). The effect of an acute overexpression of DYRK1A on mouse cortical neurogenesis has been assessed by transfecting DYRK1A expression plasmids in apical progenitors through *in utero* or *ex vivo* electroporation techniques. In 2010, Yabut et al. showed that the overexpression of DYRK1A during mid-corticogenesis inhibits proliferation, thus leading to advanced differentiation (advanced production of basal progenitors and newborn neurons). Moreover, they provided the first evidence that DYRK1A overexpression reduces nuclear Cyclin D1 protein levels. In similar electroporation experiments, Hammerle et al. showed that overexpression of DYRK1A promotes cell cycle exit, but in these experiments, differentiation was abrogated (Hammerle, et al., 2011). To explain the failure of the progenitors that express DYRK1A under a heterologous promoter to differentiate and based on the dynamic expression of DYRK1A in precursor and differentiating neurons (Hammerle, et al., 2008), the authors hypothesized that *DYRK1A* transcripts must be downregulated in the progenitors that exit the cell cycle to allow these cells to differentiate. Nonetheless, differences in the levels of the DYRK1A protein in the electroporated progenitors may explain the apparent discrepancy between these two studies. Indeed, it has been noted that *in vivo* electroporation of DYRK1A expression plasmids at concentrations that produce only a moderate increase in the DYRK1A protein does not appear to alter cortical neurogenesis (Kurabayashi & Sanada, 2013). Interestingly, Kurabayashi and Sanada showed that moderate overexpression of DYRK1A and RCAN1 delays neuron production, thus leading to an alteration in the fate of the neuronal progeny. Moreover, the authors demonstrated that this delay in neurogenesis resulted from the cooperative action of DYRK1A and RCAN1 on NFAT transcriptional activity. Thus, on the basis of these various *in*

in vivo overexpression studies, it is likely that DYRK1A regulates cortical neurogenesis by impinging on multiple pathways (e.g., cell cycle progression and/or NFAT activity).

The effect of DYRK1A on cortical neurogenesis has also been studied using the mBACTgDyrk1a mouse model, which overexpresses DYRK1A in a space- and time-regulated manner (Guedj et al., 2012; (Najas, et al., 2015). In this model, neuronal production is delayed, thus leading to a deficit of neocortical neurons that extends until postnatal stages. This delay is caused by an augmented production of basal progenitors at the expense of neurons, which correlates with a lengthened G1 phase of the cell cycle in apical progenitors at the onset of neurogenesis. The levels of p27^{Kip1} and p21^{Cip1} in these progenitors are normal, but there is a significant deficit of nuclear CyclinD1. Conversely, *Dyrk1a*^{+/-} apical progenitors have excess nuclear Cyclin D1 and produce more neurons at the expense of basal progenitors (Najas et al., 2015). These data indicate that variations in the amount of the DYRK1A protein change the division mode of apical radial glial progenitors through a mechanism that likely involves DYRK1A-mediated phosphorylation of Cyclin D1.

Active neurogenesis in the adult brain is restricted to the SVZ of the lateral ventricle and to the subgranular zone (SGZ) of the hippocampus (Alvarez-Buylla & Lim, 2004). There is evidence that epidermal growth factor receptor (EGFR) distribution varies between the daughters of adult SVZ neural stem cells (NSCs) and that inheritance of unequal receptor amounts determines the stem cell potential of this progenitor type (Andreu-Agullo, Morante-Redolat, Delgado, & Farinas, 2009). In these NSCs, DYRK1A distributes either symmetrically or asymmetrically during mitosis, and EGFR level and distribution in NSC daughter cells depend on the amount of inherited DYRK1A. Mechanistically, this is explained by the inhibitory effect of DYRK1A on endocytosis-mediated degradation of EGFR signaling. In the loss-of function *Dyrk1a*^{+/-} mouse model, EGF-dependent cell fate decisions and long-term persistence of NSCs in the niche are disturbed (Ferron, et al., 2010). Thus, normal levels of DYRK1A are required to sustain neurogenesis in the adult SVZ. Evidence obtained in DS mouse models (see part 7) indicates that DYRK1A also plays a role in neurogenesis of the adult hippocampus, but the

underlying mechanism has not yet been described.

3.2 Astrogliogenesis

After the neurogenic period, the apical progenitors of the developing neocortex acquire the capacity to produce glial cells. In 2015, Kurabayashi et al. showed that the overexpression of DYRK1A upregulates the activity of the astroglial transcription factor STAT in wild-type progenitors as well as the levels of Ser727 phosphorylation in STAT3, a modification that enhances STAT3 activity. They provide evidence that STAT3 phosphorylation by DYRK1A contributes to the overproduction of astrocytes in the neocortex of a trisomic DS mouse model with three copies of *Dyrk1a*. This pro-astroglial role of DYRK1A is in apparent contradiction with the altered number of astrocytes observed in the adult hippocampus of the mBACTgDyrk1a and *Dyrk1a*^{+/-} mouse models, which is decreased in the gain-of-function model and increased in the loss-of-function model (Fotaki, et al., 2002; Guedj, et al., 2012). Further investigations are needed to understand the possible role of DYRK1A in astroglial homeostasis.

3.3 Developmental apoptosis

The retina, a part of the CNS, has been extensively used as a model system to study neurogenesis in vertebrates (Cepko, 2014). The effect of *Dyrk1a* dosage imbalance in retinal neurogenesis has been assessed in the loss-of-function *Dyrk1a*^{+/-} and gain-of-function tgYAC152f7 mice. In both models, retinal neurons are produced at normal rates and at the correct time. However, retinal thickness and cellularity at postnatal developmental stages are altered: decreased in the loss-of-function model and increased in the gain-of-function model. These alterations affect only the internal layers of the retina (internal nuclear layer and ganglion cell layer) and result from aberrant activity of the intrinsic apoptotic pathway in retinal differentiating neurons through a process involving CASP9 phosphorylation by DYRK1A (Laguna, et al., 2008). Consistent with the antiapoptotic role of DYRK1A in retinal differentiating cells, the retinas of mBACTgDyrk1a mice are thicker and have more neurons in their internal layers than those of wild-type mice (Laguna, et al., 2013).

Studies performed in the ventral brain mesencephalon of the *Dyrk1a*^{+/-} mouse model showed that there is a deficit of dopaminergic neurons in this brain region like in the retina and that this deficit is not due to an altered neurogenesis but rather due to an increased CASP9-mediated cell death (Barallobre, et al., 2014; Fotaki, et al., 2002; Guedj, et al., 2012). In the *dyrk1aa* zebrafish knockout model, brain size reduction seems to be caused by an increased neuronal cell death during development (Y. M. Kim, et al., 2017). Collectively, these studies indicate that DYRK1A regulation of programmed cell death may contribute to the brain size defects associated with *DYRK1A* haploinsufficiency.

4 Synaptogenesis and neuronal functions

There are several spatiotemporal functional profiles for DYRK1A/MNB. DYRK1A is expressed during development at later stages (Marti, et al., 2003; Song, et al., 1996) and has been detected on the apical side of dendrites, in growing axons (Hammerle, Carnicero, et al., 2003; Hammerle, et al., 2008), and in axonal growth cones (Vidaki, et al., 2017). The expression of DYRK1A in differentiated neurons suggests a possible role in neurite formation (Gockler, et al., 2009; Hammerle, Elizalde, et al., 2003; Hammerle, et al., 2008). This is supported by several studies that reported alterations in axon growth, dendritic arborization, or dendritic spines due to changes in DYRK1A dosage in gain-of-function or loss-of function models (transgenic mice, cell culture with siRNA, etc.). For instance, DYRK1A overexpression in primary mouse cortical neurons significantly reduced dendritic growth and complexity through disruption of the REST/NRSF-SWI/SNF chromatin remodeling complex (Lepagnol-Bestel, et al., 2009). In addition, a genetic knockdown of *Dyrk1a* in cultured cortical neurons resulted in neurons with shorter and more branched neurites and fewer axons (Scales, Lin, Kraus, Goold, & Gordon-Weeks, 2009). This result is in line with that reported in a previous study that showed that pyramidal neurons in the *Dyrk1a* haploinsufficient mouse cortex have considerably reduced dendritic arborization and dendritic branching as well as fewer dendritic spines than those of wild-type animals (Benavides-Piccione, et al., 2005; Fotaki, et al., 2002). Additionally,

although specific knockdown of DYRK1A in COS-7 cells promoted filopodia formation (J. Park, et al., 2012), DYRK1A overexpression caused a reduction in dendritic spine formation in cultured hippocampal neurons. An additional study reported that cortical neurons from *Dyrk1a* overexpressing mice exhibited reduced dendritic spine density, dendritic filopodia length, and synapse formation and a dendritic spine phenotype with thinner and more immature spines (Martinez de Lagran, et al., 2012). More recently, it was found that either knockdown or overexpression of *Dyrk1a* strongly represses dendritic spine formation in hippocampal neurons, thus indicating that *Dyrk1a* gene dosage is critical for proper dendritic spine development (Dang, et al., 2017). However, previous studies suggested that the overexpression of DYRK1A increases the spine density of cortical pyramidal neurons (Altafaj, et al., 2001; Thomazeau, et al., 2014), an effect that can be prevented by a green tea extract enriched in the DYRK1A inhibitor EGCG. A recent work found that mutations in MNB/DYRK1A kinases perturb the overall neuronal outgrowth and maintenance of terminal branch length (Ori-McKenney, et al., 2016). Finally, while DYRK1A overexpression reduces total neurite numbers and axon length in primary cultured mouse cortical neurons, a knockdown of *Dyrk1a* by RNA interference or missense mutations in *Dyrk1a* induced a significant decrease in neurite and axon length (Dang, et al., 2017). All these findings highlight the impact of *Dyrk1a/Mnb* gene dosage on dendrite and axon development, and consequently synaptogenesis, an active process of synapse formation and maintenance that affects neuronal phenotypes and circuit structure.

The mechanisms underlying the function of DYRK1A in neurite formation, dendritogenesis, or synaptogenesis are largely unknown. Cytoskeletal modifications, regulation of transcription factors, and membrane trafficking have been suggested as potential stakeholders (Figure 4).

4.1 Cytoskeletal arrangement

Axon growth and guidance is an important step in synapse formation. At the tip of outgrowing axons, motile growth cones sense guidance cues and translate this information into dynamic cytoskeletal reorganizations that orient growth in a specific direction (Sanes & Yamagata,

1999). Only one study has provided direct evidence that DYRK1A may control synaptogenesis through axon guidance. This study showed that DYRK1A and RTK, which are Sprouty antagonists, co-localize in the growth cone-like structures of cultured cortical neurons (Aranda, et al., 2008). The authors proposed that DYRK1A phosphorylates and inhibits Sprouty, thereby promoting FGF signaling (Aranda, et al., 2008), which in turn regulates axon guidance (Dabrowski, Terauchi, Strong, & Umemori, 2015; Shirasaki, Lewcock, Lettieri, & Pfaff, 2006). Two phases, synaptic assembly and synaptic formation, define synaptogenesis (Colon-Ramos, 2009). Cytoskeletal components, cytoplasmic scaffold assembly, and the trans-synaptic complex might collectively establish a checkpoint for the maturation of initial unstable assemblies into stable synapses. F-actin and its regulatory proteins are intriguing candidates for this task because they possess the combinatorial ability to interact with cell-surface proteins and with active zone proteins.

DYRK1A can regulate the cytoskeletal machinery (Colon-Ramos, 2009; K. Dowjat, Adayev, Kaczmarek, Wegiel, & Hwang, 2012; F. Liu, et al., 2008; Martinez de Lagran, et al., 2012; J. Park, et al., 2012; Scales, et al., 2009), thereby contributing to the development/establishment and maintenance of neurites and dendritic spines (F. Liu, et al., 2008; J. Park, et al., 2012) and ultimately neuronal function. As already mentioned, DYRK1A is both a nuclear and a cytoplasmic protein, with the latter existing in three pools: soluble, cytoskeletal associated, and membrane bound (Aranda, et al., 2008; Kaczmarek, et al., 2014; Marti, et al., 2003). DYRK1A phosphorylates cytoskeletal proteins, which then affect the outgrowth of microscopic fibers and actin assembly. Because of its enzymatic targets, DYRK1A is strongly implicated in the regulation of cytoskeletal protein assemblies such as actin (T. Liu, Sims, & Baum, 2009; J. Park, et al., 2012), tubulin in the form of microtubules (Ori-McKenney, et al., 2016; Scales, et al., 2009), and the microtubule-associated protein (MAP) tau (F. Liu, et al., 2008; Ryoo, et al., 2008; Woods, Cohen, et al., 2001).

4.2 Membrane trafficking

Synaptogenesis as well as spine morphology and function are also regulated by membrane trafficking pathways (Winkle & Gupton, 2016) in coordination with cytoskeletal dynamics (Hotulainen & Saarikangas, 2016; Newpher, Harris, Pringle, Hamilton, & Soderling, 2017; Spence & Soderling, 2015). DYRK1A is found in the presynaptic terminal of the neuromuscular junctions and in axons from the facial nucleus, thus suggesting a function for DYRK1A in these structures (Arque, Casanovas, & Dierssen, 2013; C. K. Chen, et al., 2014). DYRK1A phosphorylates and thus regulates the protein–protein interactions of various factors involved in membrane trafficking, for example, the core endocytic proteins (Adayev, Chen-Hwang, Murakami, Wegiel, et al., 2006; Chen-Hwang, Chen, Elzinga, & Hwang, 2002; C. K. Chen, et al., 2014; Geng, et al., 2016; Murakami, Bolton, & Hwang, 2009; Murakami, Bolton, Kida, Xie, & Hwang, 2012; Murakami, et al., 2006). DYRK1A-dependent regulation of clathrin-coated vesicle formation also contributes significantly to synaptic function. In both fibroblasts and neurons, overexpressed DYRK1A appears to inhibit endocytosis in transferrin internalization assays by causing defects in clathrin-mediated endocytosis (Y. Kim, Park, Song, & Chang, 2010). Moreover, DYRK1A phosphorylates dynamin, which is also critically linked to endocytosis and important for synaptogenesis (Fan, Funk, & Lou, 2016; Schmid, McNiven, & De Camilli, 1998), and amphiphysin, which decreases the binding of amphiphysin to endophilin (Murakami, et al., 2006). Endophilin is an endocytic protein, which like the clathrin heavy chain, can bind to DYRK1A (Murakami, et al., 2009). DYRK1A inhibits the onset of clathrin-mediated endocytosis in neurons by phosphorylating dynamin1, amphiphysin1, and Synj1 and can promote the uncoating process of endocytosed clathrin-coated vesicles. Phosphorylation of the clathrin-coated vesicle adaptor proteins MAP1A, MAP2, AP180, and α - and β -adaptilins by DYRK1A promotes the dissociation of these proteins from the clathrin-coated vesicle membranes (Murakami, et al., 2012).

4.3 Regulation of transcription factors

DYRK1A is localized in the nucleus in hippocampal neurons and has an NLS and a 13-histidine repeat for nuclear speckle targeting (Alvarez, et al., 2003; Kentrup, Joost, Heimann, & Becker,

2000). This localization is consistent with the findings reported in several studies showing that DYRK1A regulates the splicing machinery at post-transcriptional and post-translational levels (Toiber, et al., 2010; Wegiel, Gong, & Hwang, 2011; Wegiel, Kaczmarek, et al., 2011). DYRK1A overexpression (in either fetal DS brains, engineered mouse models, or cultured cells) alters the phosphorylation and subcellular location of splicing machinery components and modifies key synaptic transcripts such as neuroligin 1 (Toiber, et al., 2010). Such DYRK1A-regulated cell adhesion proteins have an important role in synaptogenesis and maintenance of adult synapse number (Kwon, et al., 2012). The neuroligin–neuroligin pair is an example of the synaptogenic membrane protein interactions needed for the establishment of functional trans-synaptic communication to guide synapse assembly. Moreover, overexpression of DYRK1A causes splicing aberrations resulting in changes in the expression levels of various mRNAs such as *TrkBT1*, *Bdnf*, *AchE-S*, and *AchE-R* (Toiber, et al., 2010) involved in synaptogenesis.

Additionally, local mRNA translation is required for axon guidance and synaptogenesis (Dabrowski, et al., 2015; Gao, et al., 2012; Jung, et al., 2011; Piper & Holt, 2004; Shirasaki, et al., 2006), and *Dyrk1a* was found to be locally translated in axons in a Mena-dependent manner (Vidaki, et al., 2017).

5. DYRK1A and neurotransmission

DYRK1A expression in mature brain neurons (Marti, et al., 2003; Wegiel, et al., 2004) and the possible role of two DYRK1A-regulated transcription factors, NFAT (Arron, et al., 2006) and CREB (E. J. Yang, et al., 2001), in synaptic function imply that DYRK1A may have a role in the maintenance of adult brain neural activity (Table II).

5.1 Basal synaptic activity

Monitoring spontaneous neural activity revealed that DYRK1A might be involved in synaptic transmission. Cortical primary cultures with transgenic overexpression of DYRK1A exhibit significantly reduced numbers of spikes, thus suggesting a reduction in spontaneous activity

(Martinez de Lagran, et al., 2012). However, electrophysiological recordings revealed an increased miniature excitatory postsynaptic potential (mEPSP) amplitude but normal-evoked EPSP amplitudes at the *Drosophila* neuromuscular junction in a classical *Dyrk1a/mnb* fly mutant, which has a mutation next to the critical ATP-binding residue in the kinase-active site previously shown to influence MNB function and protein level (C. K. Chen, et al., 2014). The authors observed no change in postsynaptic AMPAR in the *mnb* mutant, thus supporting the notion that the increased mEPSP amplitude was due to presynaptic changes. Indeed, they found that presynaptic overexpression of MNB in the *mnb* mutant protected the mEPSP phenotype. In addition, an increased synaptic vesicle size has been found in a variety of endocytic mutants (C. K. Chen, et al., 2014). This is consistent with the role of DYRK1A in endocytosis as discussed above. Synaptic activity increases DYRK1A/MNB mobilization to endocytic zones and efficiently promotes synaptic vesicle recycling by regulating synaptojanin function, which directly has an impact on synaptic transmission. A similar increase in the amplitude of miniature excitatory postsynaptic currents (mEPSCs) correlated with increased spine density was observed in prefrontal cortex pyramidal neurons of mBACTgDyrk1a mice (Thomazeau, et al., 2014).

5.2 Presynaptic changes: synaptic vesicle pool and dynamics

Changes in mEPSCs amplitude could formally occur through postsynaptic (receptor abundance or activity) or presynaptic (such as synaptic vesicle size) changes, but evidence discussed above favors a presynaptic mechanism. Consistent with this, DYRK1A/MNB is also located presynaptically (Arque, et al., 2013; C. K. Chen, et al., 2014) and is tightly coupled to the endocytic machinery, which can affect synaptic vesicle endocytosis and subsequently neurotransmitter release. Moreover, clathrin-mediated endocytosis is essential for the recycling of membrane after neurotransmitter release (Saheki & De Camilli, 2012).

In addition to regulating endocytosis, DYRK1A, through its interactors, could affect exocytosis, neurotransmitter release, or delivery of receptors at synaptic membranes. It was recently shown that the DYRK1A substrate dynamin 1 affects exocytosis in addition to its essential role

in vesicle endocytosis (Anantharam, et al., 2011; Chan, Doreian, & Smith, 2010; Gonzalez-Jamett, et al., 2010), but the role of phosphorylation of dynamin 1 by DYRK1A in this process remains unclear (J. H. Park, Jung, Kim, Song, & Chung, 2012). It is known, however, that DYRK1A-mediated phosphorylation of MUNC18-1, a central regulator of neurotransmitter release, controls the interaction of MUNC18-1 with syntaxin 1, which stimulates SNARE-mediated exocytosis (with the formation of the SNARE complex required for membrane fusion during synaptic vesicle exocytosis) (J. H. Park, et al., 2012). Finally, α -synuclein, a presynaptic protein involved in neurotransmitter release, is an additional substrate of DYRK1A (Burre, 2015) and has an important role in curvature stabilization, which is important for synaptic vesicle trafficking during both endocytosis and exocytosis (E. J. Kim, et al., 2006; Lautenschlager, Kaminski, & Kaminski Schierle, 2017).

5.3 Postsynaptic changes

In addition to its effect on neurotransmitter release, DYRK1A can regulate synaptic receptor trafficking, including membrane delivery and internalization of glutamate receptors, AMPARs or NMDARs. NMDARs are involved in neural development, survival, synaptic plasticity, and memory processes. DYRK1A phosphorylates the NMDAR subunit GRIN2A on its C-terminal domain. This modifies the biophysical properties of the receptor in favor of channel opening, thereby resulting in a longer NMDAR-induced Ca^{2+} transient decay (Altafaj, et al., 2008). This modification also decreases the internalization of NR2A-containing NMDAR, thus leading to an increase in the surface density of these receptors. The DYRK1A-dependent regulation of the amount and biophysical properties of synaptic membrane NMDARs indicates that DYRK1A participates in regulating Ca^{2+} signaling and ultimately contributes to synaptic transmission (Altafaj, et al., 2008; Grau, et al., 2014). DYRK1A might also contribute to the regulation of intracellular receptor trafficking by interacting with Sprouty, a protein associated with cytosolic vesicles involved in endocytic events controlling receptor trafficking (Aranda, et al., 2008). Unlike NMDARs, NR1 and NR2AR receptors appear to be negatively regulated by DYRK1A.

Mice overexpressing DYRK1A have lower levels of NR1 and NR2A receptors and CAMKII as well as a lower pCAMKII/CAMKII ratio (Souchet, et al., 2014).

5.4 Excitation/Inhibition balance

DYRK1A seems to have an important role in controlling the excitatory/inhibitory balance, with DYRK1A overexpression increasing the number of inhibitory synapses and leading to enhanced inhibition. Increased expression of *Dyrk1a* in the mouse (mBACtgDyrk1a) induced molecular alterations in synaptic plasticity pathways and expression changes in GABAergic- and glutamatergic-related proteins leading to an excitation/inhibition imbalance related to GABA synthesis (Souchet, et al., 2014). More precisely, this study showed that excess DYRK1A dosage induces the activation of the GABA pathway, thereby increasing the enzymes involved in decarboxylation of glutamate to produce GABA (GADs), a marker of GABAergic synapses, as well as vesicular transport of GABA (VGAT). By contrast, the protein involved in glutamate transport decreased, thereby resulting in elevated GABAergic neurotransmission and reduced glutamatergic transmission. Conversely, decreasing the level of DYRK1A induced a decrease in GABA production and an increase in glutamate transport-associated proteins (Souchet, et al., 2014). Similar results were found in an additional study where DYRK1A overexpression enhanced the expression of GADs and reduced the expression of VGLUT, a marker of glutamatergic synapses, and these effects could be reversed by removing one copy of the *Dyrk1a* gene (Garcia-Cerro, et al., 2014). It was recently reported that this DYRK1A-mediated increase in inhibition decreases the firing rate of cortical neurons. This perturbs the gamma range activity and thus affects the overall neuronal network (Ruiz-Mejias, et al., 2016).

5.5 Monoamine modulation

Dyrk1a overexpression induces major deficits in serotonergic, dopaminergic, and noradrenergic systems, thus revealing that DYRK1A modulates monoamine neurotransmission (London, et al., 2017). Serotonin is related to GABA synthesis in the CNS, and evidence suggests interactions between glutamatergic transmission and the monoamine systems (Yuen, et al., 2014). Thus, the dopamine and noradrenergic neurotransmitter systems

are also important for neural and synaptic functions, especially in the prefrontal cortex known to be involved in intellectual disabilities (IDs) (Xing, Li, & Gao, 2016).

5.6 Alterations in synaptic plasticity

Long-term potentiation (LTP) and long-term depression (LTD) are examples of plasticity associated with either an increase or decrease, respectively, in synaptic strength following high-frequency stimulation of a chemical synapse. Only a few studies have reported that DYRK1A is involved in synaptic plasticity. One study found that overexpression of DYRK1A impairs the long-term synaptic plasticity of hippocampal Schaeffer collateral-CA1 synapses without affecting normal basal synaptic transmission. Indeed, the authors found that although NMDAR-LTP is increased, a paired-pulse low-frequency stimulation-induced LTD (NMDAR and mGluR dependent) is reduced and classical low-frequency stimulation-induced NMDAR-LTD is abolished in hippocampi of mice overexpressing *Dyrk1a* (Ahn, et al., 2006). Moreover, *Dyrk1a* overexpression induces functional alterations in the prefrontal cortex of mBACtgDyrk1a mice with dendritic alterations and anomalous NMDAR-LTP and endocannabinoid-dependent LTD (Thomazeau, et al., 2014). Other studies used Ts65Dn mice, the most common rodent model for DS, to examine how normalization of the *Dyrk1a* gene dosage affects synaptic plasticity. Pharmacological inhibition of DYRK1A with EGCG protects the LTP deficit observed in the hippocampus of the Ts65Dn mice (Xie, Ramakrishna, Wieraszko, & Hwang, 2008). Moreover, injecting the hippocampus with an adeno-associated virus carrying an shRNA sequence specific for *Dyrk1a* normalized DYRK1A expression in Ts65Dn mice and restored hippocampal LTP (Altafaj, et al., 2013). Finally, crossing Ts65Dn mice with *Dyrk1a*^{+/-} mice protected hippocampal LTP in offspring with normalized *Dyrk1a* copy number (Garcia-Cerro, et al., 2014).

DYRK1A could also indirectly affect synaptic plasticity. As mentioned previously, DYRK1A overexpression turns the excitation/inhibition balance toward inhibition. LTP deficits in the hippocampus have been associated with enhanced GABA-mediated inhibition (Fernandez, et al., 2007; Yoshiike, et al., 2008). An increase in the inhibitory function mediated by GABAergic

synapses may interfere with processes required for learning and memory, as indicated by LTP deficits in the dentate gyrus (DG; Palop, et al., 2007). Consistent with this, the GABA_A receptor antagonist picrotoxin was found to prevent such LTP deficits observed in an animal model of AD (Kleschevnikov, et al., 2004). In addition, DYRK1A could indirectly affect synaptic plasticity by phosphorylating and inactivating GSK3 β (Song, et al., 2015), a protein that has a pivotal role in synaptic plasticity (Bradley, et al., 2012). Finally, by forming a multiprotein complex with Ras, Raf, and MEK1 (Kelly & Rahmani, 2005), DYRK1A might affect MAPK/ERK signaling, which is vital for synaptic plasticity (Impey, Obrietan, & Storm, 1999).

6. DYRK1A deficit, autism, and MRD7

Classical genotype–phenotype association studies of rare cases with partial monosomy 21 (Chettouh, et al., 1995; Matsumoto, et al., 1997) in combination with the physical map of the chromosomal region for DS (Osoegawa, et al., 1996) narrowed the loci of monosomy 21-associated microcephaly and intrauterine growth retardation to a 1.2 Mb segment at 21.q22.2 that contains several genes including *DYRK1A* (Matsumoto, et al., 1997). In 2008, Moller *et al.* identified two unrelated patients carrying a *de novo* balanced translocation that truncates *DYRK1A*. The overlapping phenotypic traits of these two patients, which include microcephaly, intrauterine growth retardation, and febrile seizures, indicated that *DYRK1A* haploinsufficiency alters brain development and function. Three years later, a study performed in a cohort of 3009 individuals with ID (intelligence quotient (IQ) below 70) identified one patient with a *de novo* microdeletion of 52 Kb affecting the last three exons of *DYRK1A* (van Bon, et al., 2016). Depending on the clinical features presented by this patient and by previously reported patients with either chromosomal rearrangements that truncate *DYRK1A* (Moller, et al., 2008) or with large deletions encompassing this gene (Fujita, et al., 2010; Lyle, et al., 2009; Matsumoto, et al., 1997), van Bon *et al.* proposed that the heterozygous disruption of *DYRK1A* causes a distinctive clinical syndrome, characterized by the presence of mild to severe ID, microcephaly, intrauterine growth retardation, facial dimorphisms, impaired motor functions, and behavioral

problems. The phenotype presented by haploinsufficient MNB/*Dyrk1a* flies and mice supported this hypothesis (Fotaki, et al., 2002; Tejedor, et al., 1995).

The sequencing of *DYRK1A* in a cohort of 105 patients with ID who exhibit two or more symptoms from the Angelman syndrome spectrum identified one patient with a frameshift *DYRK1A* variant affecting the N-terminal part of *DYRK1A* (Courcet, et al., 2012). This patient presented epileptic seizures, speech delay, and other clinical manifestations compared to the patient described in the study by van Bon *et al.* (2011). On the basis of this and the phenotype of patients with microcephaly and large deletions encompassing more than one chromosome 21 gene (Oegema, et al., 2010; Valetto, et al., 2012; T. Yamamoto, et al., 2011), Courcet *et al.*, proposed that MRD7 (OMIM #614104) is caused by heterozygous disruption of the *DYRK1A* gene.

Three additional *de novo* *DYRK1A* pathogenic variants, one splice site variant and two frameshift variants predicted to delete parts of the *DYRK1A* protein that are important for its function (O'Roak, et al., 2012), were identified by exome sequence analysis of a large number of families from the Simons Simplex Collection, which is a well-characterized cohort of patients with idiopathic autism spectrum disorder (ASD) (Fischbach & Lord, 2010). ASD is a complex and heterogeneous developmental disorder (DD) characterized by impairments in social interaction and communication and repetitive, restricted behaviors. ID and language delay are the most frequent comorbidities of the disorder (Lord and Bishop 2015; Geschwind and State 2015). Interestingly, the three ASD probands carrying pathogenic *DYRK1A* variants show a complex phenotype that includes mild to severe ID, microcephaly, and impaired speech (van Bon et al. 2016). Since 2013, high-throughput (whole-exome or gene target) sequencing applied to cohorts of patients with DD/ID, with or without ASD, identified 42 *de novo* truncating variants (frameshift, splice-acceptor, splice-donor, and nonsense) and missense mutations (Bronicki, et al., 2015; Dang, et al., 2017; Ji, et al., 2015; Luco, et al., 2016; Redin, et al., 2014; Ruaud, et al., 2015; van Bon, et al., 2016) in *DYRK1A*. Additional studies performed with larger cohorts of patients with DD, including the Deciphering Developmental Disorders Study

(Deciphering Developmental Disorders, 2015), substantially increased the number of *DYRK1A* point mutations and small insertions and deletions (INDELS) associated with these disorders (De Rubeis, et al., 2014; Kosmicki, et al., 2017; Stessman, et al., 2017). These studies found that *DYRK1A* is one of the most frequent *de novo*-mutated genes in ASD (Iossifov, et al., 2014; Kosmicki, et al., 2017; Stessman, et al., 2017; Yuen, et al., 2014), thus accounting for 0.1–0.5% of the ASD population (van Bon et al. 2015) and for approximately 0.5% of syndromic ID (Evers, et al., 2017).

DYRK1A-truncating variants are concentrated in the N-terminal region and the kinase domain of *DYRK1A*, whereas most missense mutations are within the kinase domain (Earl, et al., 2017; Ji, et al., 2015; Luco, et al., 2016; van Bon, et al., 2016) and are predicted to affect the enzymatic activity of the protein (Evers, et al., 2017). Functional studies showed that missense *DYRK1A* variants might impair enzymatic function by affecting catalytic residues or by compromising the structural integrity of the *DYRK1A* kinase domain (Widowati, Ernst, Hausmann, Muller-Newen, & Becker, 2018).

The clinical phenotypes of patients with *DYRK1A de novo* pathogenic variants or chromosomal mutations involving this gene defined a distinct ID syndrome. Core symptoms of the *DYRK1A*-related syndrome (also known as *DYRK1A*-haploinsufficiency syndrome or MRD7) are microcephaly, intrauterine growth retardation, developmental delay, seizures, speech problems, ASD or ASD-related deficits (e.g. stereotypies and anxious behaviors), neonatal feeding problems, hypertonia, gait disturbances, and a characteristic dysmorphic facies (Bronicki, et al., 2015; Earl, et al., 2017; Ji, et al., 2015; Luco, et al., 2016; van Bon, et al., 2016). Consistent with the effect of *Dyrk1a* haploinsufficiency on mouse brain development (Fotaki, et al., 2002), several structural brain alterations including cerebral atrophy, hypoplasia of the corpus callosum, and a thin optic chiasm have been detected by magnetic resonance imaging (MRI) in some patients with mutations in *DYRK1A* (Ji, et al., 2015; Y. M. Kim, et al., 2017). Clinical studies, especially brain imaging studies, in a larger group of patients, together with studies in cellular model systems and in animal models, like the *Dyrk1a*^{+/-} mouse model that displays many of the neurological traits in *DYRK1A*-related syndrome (Arque, et al., 2008;

Fotaki, et al., 2002; Fotaki, Martinez De Lagran, Estivill, Arbones, & Dierssen, 2004), are necessary to understand the pathogenesis of the syndrome.

7 DYRK1A and Down syndrome

DS, most commonly resulting from a complete trisomy of HSA21, is associated with several alterations in brain development and function. The trisomy underlying DS has provided particular insight into how deregulated gene expression in the brain leads to altered brain function, specifically cognitive impairment.

7.1 Overexpression in the trisomic context

The *DYRK1A* gene is located on chromosome 21, thus suggesting that this gene may be overexpressed in cells from patients with DS. In a microarray study of mRNA from DS lymphoblastoid cells, 29% of expressed chromosome 21 transcripts were elevated in DS. Of these 29%, 22% were increased proportionally with gene dosage and 7% were amplified. In this cell type, *DYRK1A* was 1.4-fold overexpressed (Ait Yahya-Graison, et al., 2007). In the brain of mice with three copies of the *Dyrk1a* gene, the *Dyrk1a* mRNA level was increased by 1.5-fold (hYACtgDyrk1a and mBACtgDyrk1a) (Guedj, et al., 2012; Guedj, et al., 2009) and the *DYRK1A* protein level was overexpressed by 1.6-fold (cortex), 1.9-fold (hippocampus), or 1.7-fold (cerebellum) fold in mBACtgDyrk1a mice. Similar overexpression levels were observed in models with partial trisomy (Ts65Dn and Dp(16)1Yey) (Souchet, et al., 2014). *DYRK1A* expression in the brain of patients with DS-associated trisomy was also, on average, 1.5-fold elevated, and this overexpression was preserved across a wide range of ages (W. K. Dowjat, et al., 2007).

7.2 Phenotypes in patients with DS

7.2.1 Genetics and analyses of partial duplications

DS occurs in 1 in every 750 live births and encompasses a constellation of features caused by partial or complete trisomy for chromosome 21 HSA. In particular, an altered copy number for segments of Hsa21 containing the *DYRK1A* gene can induce morphological defects and

cognitive impairments (Delabar, et al., 1993; Papoulidis, et al., 2014; Rahmani, et al., 1989; Ronan, et al., 2007). However, the genotype–phenotype correlation approach is limited by the small number of partial trisomies and the heterogeneity of clinical phenotypic descriptions; another group of partial trisomies produced discrepant results (Korbel, et al., 2009), which could not associate a specific set of genes with impairment of cognition. Linking phenotypes observed in trisomy 21 to the overexpression of *DYRK1A* can be done more efficiently by using two complementary strategies: i) analysis of mice models with different levels of *DYRK1A* and ii) assessing effects of treatments using *DYRK1A* inhibitors in mouse models and in patients with DS (see below).

Defects similar to those observed in DS have been reproduced in a number of different mouse models of DS (Ts1Rhr, Ts65Dn, Ts1Cje, Dp(16)1Yey) as well as in mice with altered copy numbers of *Dyrk1a* (hBACtgDyrk1a, hYACtgDyrk1a, mBACtgDyrk1a, *Dyrk1a*^{+/-}). Interestingly, a phenotype rescue experiment crossing Ts65Dn mice with mice monosomic for a 33-gene chromosomal segment containing *Dyrk1a* (Ms1Rhr) produced progeny with a normal learning phenotype, thus indicating that triplication of this 33-gene region produces the cognitive deficit (Belichenko, et al., 2009). A complete phenotypic assessment of Ts1Rhr mice, trisomic for the 33-gene segment, showed that trisomy of this region is sufficient to produce significant alterations in behavioral tests such as the open-field, novel object recognition, and T-maze tasks. In Ts65Dn, Ts1Cje, and Ts1Rhr mice, LTP in the fascia dentata, a brain region critical to learning and memory, could be induced only after blocking GABA(A)-dependent inhibitory neurotransmission. In addition, widespread enlargement of dendritic spines and decreased density of spines in the fascia dentata were preserved (Haas, et al., 2013). Thus, cognitive impairment in DS appears to derive from molecular and structural changes related to an altered copy number within this 33-gene region. Among the 33-genes in this region, *DYRK1A* is an attractive candidate for inducing brain structure alterations and cognitive impairment phenotypes.

7.2.2 Morphology

Brachycephaly and increased ventricle size have been reported in individuals with DS (Allanson, O'Hara, Farkas, & Nair, 1993; Pearlson, et al., 1998; Schimmel, Hammerman, Bromiker, & Berger, 2006). Prenatal EGCG treatment and *Dyrk1a* dosage reduction modified craniofacial features in Ts65Dn mice and brain morphology in YACtgDyrk1a mice (Guedj, et al., 2009; McElyea, et al., 2016), thus suggesting that DYRK1A overexpression alters brain morphology.

7.2.3 Dendritic alterations

Several studies have reported abnormal cortical lamination patterns, altered dendritic arbors and spines, aberrant membrane electrophysiological properties, reduced synaptic density, and abnormal synaptic morphology in patients with DS (L. E. Becker, Armstrong, & Chan, 1986; Dierssen & Ramakers, 2006; Marin-Padilla, 1976; Takashima, Iida, Mito, & Arima, 1994). Alterations in the dendritic structure should have a major impact on the processing of afferent information by single neurons. At the level of the neuronal network, even modest alterations in dendritic structure and organization of many neurons, as seen in other MR, will lead to considerable changes in overall information processing.

7.2.4 Cognition

Trisomy 21 reduces IQ to between 20 and 80 (Anneren & Edman, 1993). In contrast to normally developing children and other cases of MR, there is a progressive IQ decline in DS beginning in the first year of life; therefore, the ratio of mental age to chronological age is not constant. By adulthood, IQ is usually in the moderate-to-severe level of impairment, thus suggestive of ID (IQ 25–55) with an upper limit on mental age of approximately 7–8 years, although a few individuals have IQs in the lower normal range (IQ 70–80) (Pennington, Moon, Edgin, Stedron, & Nadel, 2003).

A randomized, double-blinded, placebo-controlled, pilot study has shown preliminary results on the safety and clinical effects of EGCG, a DYRK1A inhibitor, in young adults with DS. A visual recognition task that measures visuoperceptual processing, the weakest component of visual memory in individuals with DS, was used. This measure has proven to be sensitive to

hippocampal functioning, in particular of the perirhinal cortex, which is critically involved in object recognition memory and is reduced in size in DS. EGCG-treated individuals showed higher accuracy in visual memory recognition and spatial working memory, thus suggesting a positive effect of this compound on both the hippocampal and prefrontal system, in particular the ventromedial, ventrolateral, and dorsolateral cortices (De la Torre, et al., 2014).

Following up on these promising results, a phase II study compared an “EGCG plus cognitive training” group and a “placebo plus cognitive training” group. This study showed that patients in the EGCG group performed better than those in the placebo group in some cognitive tests and in adaptive behavior after the 12 months of treatment. The EGCG and cognitive training group had better preservation of recognition memory tasks and improvement in executive function than the placebo and cognitive training groups.

In both trials, the main effect of EGCG was the improvement of immediate recognition memory. This measure is sensitive to hippocampus, in particular, perirhinal cortex activity, in addition to regions such as the ventromedial cortices. In DS, the altered function of the hippocampus and prefrontal cortex contributes to memory and executive functioning deficits and to distinct connectivity disturbances in frontal and anterior temporal structures. It was proposed that the efficacy of EGCG depends, at least partly, on the inhibition of DYRK1A kinase activity. Total homocysteine plasma concentrations, used as a surrogate biomarker of DYRK1A kinase activity, were increased in the “EGCG and cognitive training” group to levels not observed in the “placebo and cognitive training” group and returned to baseline concentrations after discontinuation of treatment (de la Torre, et al., 2016). However, in this study, a reduction in total cholesterol and oxidized LDL concentrations was also noted. Thus, a lipid-lowering effect combined with a reduced lipid oxidation might contribute to the therapeutic effect. Other mechanisms of action of EGCG should not be disregarded, including epigenetic, protection from mitochondrial dysfunction, and antioxidant effects (Vacca & Valenti, 2015).

7.3 Phenotypes in murine models

7.3.1 Morphogenesis

Comparison of mice models with increased (hYACtgDyrk1a, hBACtgDyrk1a, and mBACtgDyrk1a) and decreased (*Dyrk1a*^{+/-}) gene copy number suggests that DYRK1A controls brain morphogenesis: global brain region-specific variations observed in gain-of-function models mirror their counterparts in the loss-of-function model (Ahn, et al., 2006; Guedj, et al., 2012; Sebrie, et al., 2008). Increased ventricle size has been observed in coronal sections from BACtgDyrk1a and through *in vivo* MRI from YACtgDyrk1a mice (Guedj, et al., 2012). This phenotype is also present in Ts65Dn, Ts1Cje, Ts2Cje mice, and additional trisomic mouse models that contain additional copies of orthologous regions of HSA21 (Ishihara, et al., 2010). In the above described trisomy models, the dorsal and lateral portions of the third ventricle are more enlarged than the ventral portion.

7.3.2 Neurogenesis

As in the TgBACDyrk1a mouse, cortical neurogenesis in Ts65Dn and Ts1Cje trisomic embryos is impaired (Chakrabarti, Galdzicki, & Haydar, 2007). Moreover, Ts65Dn apical progenitors have longer cell cycles and less nuclear Cyclin D1 than the euploid littermates (Chakrabarti, et al., 2007; Najas, et al., 2015). Remarkably, the genetic normalization of *Dyrk1a* gene dosage in Ts65Dn embryos increases the amount of Cyclin D1 to normal levels and normalizes the production of early-born cortical neurons (Najas, et al., 2015). The importance of DYRK1A overexpression in DS cortical neurogenesis is supported by the effect on the expansion of the cortical wall produced by the treatment of Ts1Cje embryos with a potent inhibitor of DYRK1A kinase activity (Nakano-Kobayashi, et al., 2017). These results indicate that alterations in the formation of the neocortex in DS start early in development and that triplication of the *DYRK1A* gene contributes to the neurogenic cortical defects associated with this syndrome.

Ts65Dn mice also exhibit postnatal neurogenic defects, but treatment of Ts65Dn pups from day 3 to day 15 with the DYRK1A-inhibiting phytochemical EGCG restored neurogenesis; total hippocampal granule cell number; and levels of pre- and postsynaptic proteins in the DG, hippocampus, and neocortex of these animals (Stagni, et al., 2016). Genetic normalization of *Dyrk1a* expression in Ts65Dn mice restored the proliferation and differentiation of hippocampal

cells in the adult DG and the density of GABAergic and glutamatergic synapse markers in the molecular layer of the hippocampus (Garcia-Cerro, et al., 2014). The increased dosage of *Dyrk1a* in the Ts1Cje mouse model of DS augments the propensity of progenitors to differentiate into astrocytes. This tendency is associated with enhanced astrogliogenesis in the developing neocortex, which is likely due to the potentiation of the DYRK1A-STAT pathway in progenitors (Kurabayashi, Nguyen, & Sanada, 2015). This work indicates that overexpression of DYRK1A likely contributes to the aberrant astrogliogenesis associated with DS.

7.3.3 Differentiation

The density of interneurons that control the level and type of inhibition is significantly increased in layers 3-6 of the somatosensory cortex of Ts65Dn mice (Perez-Cremades, et al., 2010). The density of GABAergic synapse markers is also increased in the inner molecular layer of the hippocampus (Martinez-Cue, et al., 2013). Similar density changes have been reported for the stratum radiatum of YACtgDyrk1a, BACtgDyrk1a, and Dp(16)1Yey (Souchet, et al., 2014) mice, thus suggesting that this phenotype is strongly linked with *Dyrk1a* overexpression.

7.3.4 Learning and plasticity

A functional screen using yeast artificial chromosomes covering 2 Mb of human 21q22.2 identified a 180 kb fragment containing *DYRK1A* that associated with learning defects in the Morris water maze (MWM) (Smith, et al., 1997). Additionally, animals overexpressing the full-length mouse *Dyrk1a* cDNA exhibit altered motor skill acquisition and impaired spatial learning and cognitive flexibility in the MWM (Altafaj, et al., 2001). Transgenic mice containing one extra copy of the human *DYRK1A* gene display significant impairment in hippocampal-dependent memory tasks in the MWM. Interestingly, shifts in both long-term potentiation and long-term depression were observed in these animals (Ahn, et al., 2006). In addition to alterations in LTP and LTD in the prefrontal cortex, the BACtgDyrk1a mouse also showed impairment in motor learning and novel object recognition task (Souchet, et al., 2014; Thomazeau, et al., 2014).

Mice with partial trisomy, which more closely mimics the alterations seen in DS, have also been used to study the role of *Dyrk1a* overexpression in specific phenotypes. In these animals,

genetic or pharmacological correction of DYRK1A levels not only corrected the defects in neurogenesis but also the excitation–inhibition imbalance, density of GABAergic and glutamatergic markers, and learning impairment (Catuara-Solarz, et al., 2016; De la Torre, et al., 2014; Garcia-Cerro, et al., 2014; Guedj, et al., 2009; Nakano-Kobayashi, et al., 2017; Souchet, et al., 2015).

8 Neurodegenerative diseases

8.1 AD and DS with AD

The cytoplasm and the nuclei of scattered neurons of the neocortex, entorhinal cortex, and hippocampus of patients with AD, DS, and Pick's Disease have increased DYRK1A immunoreactivity. DYRK1A is found in sarkosyl-insoluble fractions, which are enriched in phosphorylated tau in AD brains, thus suggesting a possible association of DYRK1A with neurofibrillary tangle pathology (Ferrer, et al., 2005). Compared to control levels, the *DYRK1A* mRNA level in the hippocampus is significantly elevated in patients with AD (Kimura, et al., 2007). In the AD brain, overactivation of calpain is associated with truncation of the C-terminus of DYRK1A. Quantification of the different forms of DYRK1A suggests that AD is associated with a decrease in the full-length DYRK1A protein and an increase in the truncated form, which is still active (Jin, et al., 2015). In parallel with proteomic studies, genetic studies have tried to establish a link between *DYRK1A* variants and AD. A study in a Japanese population reported a clear association between *DYRK1A* and AD risk, but the risk genotype (rs2835740) was not associated with increased expression of *DYRK1A*, thus suggesting that increased *DYRK1A* expression is induced by AD (Kimura, et al., 2007). In a case–control Spanish cohort, a study examined genetic variations in *DYRK1A* by genotyping and haplotype tagging SNPs and detected no difference between the patient and control groups either overall or after stratification by the APOE epsilon4 allele (Vazquez-Higuera, et al., 2009) (Figure 5).

8.1.1 Tau

Cellular shape is mainly determined by microtubules. MAPs are responsible for brain microtubule stabilization and maintenance of the assembled tubulin polymers. Tau is one of the most widely studied MAPs. Tau was originally found to enhance microtubule stability and polymerization but is currently known to have a number of other functions including controlling microtubule modifications, altering the stiffness and mechanical properties of the microtubule polymer, controlling the spacing of microtubules within axons, and regulating microtubule motor transport (Ramkumar, Jong, & Ori-McKenney, 2017, 2018). The DYRK1A protein can modify the functional properties of Tau by two different mechanisms that are altered in AD: i) *Tau* splicing and ii) Tau phosphorylation.

i) Tau splicing:

Six different isoforms of *Tau* generated through alternative splicing are expressed in the adult human brain (Goedert, Spillantini, Jakes, Rutherford, & Crowther, 1989). Inclusion or exclusion of *Tau* exon 10 (E10), which encodes the second microtubule-binding repeat, gives rise to Tau isoforms with either four (4R) or three (3R) microtubule-binding repeats. Almost equal levels of 3R-tau and 4R-tau are expressed in the normal adult human brain. Mutations in the *Tau* gene associated with frontotemporal dementia FTDP-17 cause dysregulation of *Tau* E10 splicing. An altered 3R-to-4R ratio in the brain also occurs in corticobasal degeneration, Pick's disease, and progressive supranuclear palsy. DYRK1A plays a very significant role in multidimensional control of tau isoforms through various splicing factors. Of note, expression of 3R-tau can be modulated by peptide amyloid β ($A\beta$), and 3R-tau levels increase with the progression of AD. This increase might also result from aberrant DYRK1A levels. Alternative splicing is controlled by both exonic and intronic enhancers and silencers, which are controlled upstream by splice factors. DYRK1A phosphorylates the splicing factor ASF at different residues (see part 2) and inhibits the ability of ASF to promote *tau* E10 inclusion (Shi, et al., 2008). DYRK1A phosphorylates ASF, thereby driving the splice factor into nuclear speckles. Moreover, phosphorylation of ASF by DYRK1A inhibits their association with nascent *tau* transcripts, thus increasing 3R-tau levels and causing an imbalance of the 3R-4R tau isoforms

(Wegiel, Kaczmarek, et al., 2011). DYRK1A regulation of 9G8 activity through phosphorylation modulates *tau* E10 splicing (Ding, et al., 2012). SC35 phosphorylation by DYRK1A suppresses the ability of SC35 to promote *Tau* E10 inclusion, whereas downregulation of DYRK1A promotes 4R-Tau expression (Qian, et al., 2011). Phosphorylation by DYRK1A also inhibits SRp55 ability to promote *Tau* exon 10 inclusion. Upregulation of *DYRK1A*, as in DS, could lead to neurofibrillary degeneration by altering the alternative splicing of *tau* exon 10 and increasing the 3R-tau-to-4R-tau ratio (Yin, et al., 2012). Treatment with EGCG from gestation to adulthood suppressed 3R-tau expression in mice (Yin, et al., 2017).

ii) Tau phosphorylation

The accumulation of paired-helical filaments (PHF) is the most characteristic neuropathological lesion of AD, including AD in DS. Abnormally phosphorylated Tau is a component of PHF (Banerjee, et al., 1991; Grundke-Iqbal, et al., 1986; Hanger, et al., 1991). DYRK1A phosphorylates human tau at Thr212 *in vitro*, a residue that is phosphorylated in fetal tau and hyperphosphorylated in filamentous tau from AD brain. Phosphorylation of Thr212 primes tau for phosphorylation by GSK3 at Ser208 *in vitro* (Woods, Cohen, et al., 2001). In a screen of 572 kinases in an AD model, Azorsa et al. found increased levels of *Dyrk1a* mRNA in the brain of Tg-PS1/APP mice and revealed that DYRK1A is involved in tau phosphorylation pathways (Azorsa, et al., 2010). Consistent with this, an elevated *DYRK1A* mRNA level was associated with tau phosphorylation at Thr212 (Kimura, et al., 2007). DYRK1A also phosphorylates other tau residues. These sites are phosphorylated in adult DS brains but not in age-matched controls. Increased expression of DYRK1A in Ts65Dn mice is also associated with increased phosphorylation of tau (F. Liu, et al., 2008). In cell cultures, DYRK1A inhibition by harmine reduced Tau phosphorylation at multiple AD-related sites (Frost, et al., 2011). Finally, DYRK1A inhibition reduced β -amyloid and tau pathology in mouse models of AD (Branca, et al., 2017; Naert, et al., 2015).

8.1.2 APP processing

DYRK1A phosphorylates APP on threonine 668 *in vitro* and in mammalian cells. The amounts of phospho-APP and A β are increased in the brains of transgenic mice overexpressing human DYRK1A, and the amounts of phospho-APP as well as those of APP are elevated in human DS brains (Ryoo, et al., 2008).

APP function may also be indirectly affected by changes in Tau, which, as discussed above, can be modulated by DYRK1A. Tau overexpression and/or mutations impair axonal transport. A study that used a trans-splicing strategy to modulate Tau exon 10 inclusion/exclusion in differentiated human-derived neurons found that 3R-tau favored anterograde movement of APP vesicles, thus suggesting that an alteration of the 3R-to-4R ratio may alter APP metabolism (Goldstein, 2012; Lacovich, et al., 2017).

DYRK1A might also affect the degradation of A β . Gene expression of the major A β -degrading enzyme NEP in sporadic AD brains is decreased beginning at the early stages of disease development, and there is an inverse relationship between *MME (NEP)* gene expression and A β accumulation, thus indicating that the downregulation of NEP is at least one cause of sporadic AD. NEP is downregulated in fibroblasts of patients with DS compared to fibroblasts from healthy controls. Treatment with the DYRK1A inhibitor harmine and knockdown of the *DYRK1A* gene upregulate NEP in fibroblasts (Kawakubo, et al., 2017). Recently, EGCG was shown to induce extracellular degradation of amyloid β -protein by increasing NEP secretion from astrocytes through the activation of ERK and PI3K pathways (N. Yamamoto, et al., 2017). Normalizing *Dyrk1A* gene dosage in aged Ts65Dn mice protected them from senescent cell density in the cingulate cortex, hippocampus, and septum; prevented cholinergic neuron degeneration; and reduced APP expression in the hippocampus, A β load in the cortex and hippocampus, the expression of phosphorylated Tau in Ser202 in the hippocampus and cerebellum, and the levels of total tau in the cortex, hippocampus, and cerebellum (Garcia-Cerro, Rueda, Vidal, Lantigua, & Martinez-Cue, 2017). Proposed consequences of increased DYRK1A level on β amyloidosis and neurofibrillary degeneration are schematized in Figure 5.

8.1.3 Peripheral DYRK1A

A search for novel peripheral biomarkers in plasma from two different cohorts consisting of patients with AD and control patients revealed that the DYRK1A level is decreased in AD and also in AD with mild cognitive impairment. This decrease is also present in lymphoblastoid cell lines derived from patients with AD. Combining the assessment of DYRK1A with BDNF and homocysteine, two DYRK1A-related markers, provides high sensitivity, specificity, and accuracy for detecting AD. Further investigations are needed to identify the cause of these variations (Janel, et al., 2017; Janel, et al., 2014) and to classify plasma DYRK1A as a diagnostic or prognostic marker.

8.2 α -Synuclein dementias

Synuclein is a soluble, natively unfolded protein that is highly enriched in the presynaptic terminals of neurons in the CNS. Dementia with Lewy bodies (LB) is characterized by α -synuclein (SNCA) accumulation and degeneration of dopaminergic and cholinergic pathways, and Ser129 in SNCA is selectively and extensively phosphorylated in synucleinopathic lesions (Fujiwara, et al., 2002). In a *Drosophila* model of PD, Ser129 phosphorylation is crucial for mediating α -synuclein neurotoxicity and inclusion formation (L. Chen & Feany, 2005). DYRK1A phosphorylates α -synuclein in transformed and primary neuronal cells on Ser87, which leads to α -synuclein aggregation and a decrease in cell viability (E. J. Kim, et al., 2006). Furthermore, phosphorylation of α -synuclein at Ser129 promoted fibril formation *in vitro*. *In vitro* kinase assays of anti-DYRK1A immunocomplexes demonstrated that DYRK1A can phosphorylate α -synuclein at Ser87. It is not yet clear whether DYRK1A can also phosphorylate Ser129. It was proposed that α -synuclein neurotoxicity in PD and related synucleinopathies may result from an imbalance between the detrimental, oligomer-promoting effect of Ser129 phosphorylation and the neuroprotective action of Tyr125 and Ser87 phosphorylations that inhibit toxic oligomer formation (L. Chen, et al., 2009). A pilot study identified the DYRK1A rs8126696 polymorphism as a risk factor for developing an α -synuclein-associated dementia (Jones, Aarsland, Londos, & Ballard, 2012). Collectively, the above data indicate that DYRK1A in pathological brain

regions may contribute to aggregations of SNCA peptides, which are a major component of amyloid plaques in the brains of patients with AD (Irwin, Lee, & Trojanowski, 2013).

8.3 Parkinson's disease

PD is a long-term degenerative disorder of the CNS that mainly affects the motor system. The motor symptoms of PD are the result of reduced dopamine production in the basal ganglia of the brain. Cognitive and behavioral problems such as dementia and depression often arise as the disorder develops.

The *DYRK1A* rs8126696 T allele was associated with early onset in a cohort 297 Chinese patients with PD (K. Fan, et al., 2016). An additional study in the Chinese Han population identified the TT genotype derived from SNP rs8126696 of the *DYRK1A* gene as a possible risk factor for developing sporadic PD, especially for men (Cen, et al., 2016). Increased *DYRK1A* expression in the BACtgDyrk1a mouse model is associated with increased survival of mesencephalic dopaminergic neurons (mDA) of animals injected with 1-methyl-4-phenyl-1,2,3,6 tetrahydropyridine (MPTP), a toxin that activates caspase-9-dependent apoptosis in mDA neurons (Barallobre, et al., 2014).

DYRK1A may influence the behavior of parkin, the protein product of the first gene known to cause autosomal recessive familial PD. Multiple kinases phosphorylate parkin at several distinct sites and regulate its ubiquitin E3 ligase activity. *DYRK1A* directly phosphorylates parkin at Ser131 *in vitro*, which inhibits the E3 ubiquitin ligase activity of parkin and, consequently, its neuroprotective function in dopaminergic SH-SY5Y cells exposed to 6-hydroxydopamine (Im & Chung, 2015). In PD, α -synuclein aggregates often contain and sequester Septin4 (Sept4), a polymerizing scaffold protein. In a yeast two-hybrid screen, Septin4 was identified as a *DYRK1A*-binding partner, which colocalizes with *DYRK1A* in mouse neurons. *DYRK1A* phosphorylation of septin4 is inhibited by harmine (Sitz, et al., 2008), thus suggesting a role for *DYRK1A* in health of dopaminergic neurons.

9 Inhibitors

9.1 *DYRK1A* inhibitors and their bioavailability

The elevated activity of DYRK1A in several human diseases makes DYRK1A an attractive potential target for therapeutic drugs, and much effort has been applied toward developing DYRK1A inhibitors. The chemical compounds shown to inhibit DYRK1A in living cells are listed in Table II. In this chapter, we focus on chemicals applicable to disease models and discuss issues relating to their future application.

As mentioned before, prenatal exposure to green tea polyphenols protects from brain defects induced by overexpression of DYRK1A (Guedj, et al., 2009), particularly EGCG, which is the most abundant catechin in green tea and a known DYRK1A inhibitor (Bain, McLauchlan, Elliott, & Cohen, 2003; Bain, et al., 2007). Of note, EGCG is a noncompetitive inhibitor, thus suggesting that regions outside the DYRK1A active site are important for its enzymatic activity. However, EGCG has profound effects on other signaling pathways. EGCG is an antioxidant/metal chelator and a known inhibitor of proteasomes, matrix metalloproteinase, dihydrofolate reductase, DNA methyltransferase, topoisomerase II, and telomerase (Khan, Afaq, Saleem, Ahmad, & Mukhtar, 2006; Mandel, Amit, Weinreb, Reznichenko, & Youdim, 2008), and it is difficult to attribute its pharmacological effect solely to the inhibition of DYRK1A. Harmine is a β -carboline alkaloid that inhibits the kinase activity of DYRK1A and interferes with neurite formation (Gockler, et al., 2009). Harmine has psychotropic effects owing to its inhibitory activity on monoamine oxidase A (MAO-A); therefore, groups have applied structure–activity relationship (SAR) analysis of harmine derivatives to design DYRK1A inhibitors without MAO-A inhibitory activity (Drung, et al., 2014; Ruben, et al., 2015). Meridianin is a natural alkaloid from the marine tunicate (Giraud, et al., 2011; Yadav, et al., 2015) shown to inhibit DYRK1A. Additionally, Leucettine L41 was extracted and optimized from marine sponges, thus showing potent inhibitory activity against DYRK1A in cell culture and in brain slices (Debdab, et al., 2011; Tahtouh, et al., 2012). L41 treatment rescued memory impairment in A β -injected mice (Naert, et al., 2015), thus indicating its bioavailability.

Based on the structural similarity of the ATP-binding pocket of DYRKs and CLKs, INDY (inhibitor of DYRKs) was developed by altering the CLK inhibitor TG003 to enhance its

specificity for DYRKs (Ogawa, et al., 2010). INDY has potent inhibitory effects on DYRK1A in cultured cells and prevented the abnormal development of *Xenopus laevis* embryos overexpressing *Dyrk1a*. On the basis of the crystal structure of the DYRK1A/INDY complex, the novel inhibitor BINDY was designed and synthesized by replacing the phenol group of INDY with dibenzofuran to produce the BINDY derivative. Treatment of 3T3-L1 pre-adipocytes with BINDY-hampered adipogenesis was by suppressing gene expression of the critical transcription factors PPAR γ and C/EBP α (Masaki, et al., 2015). GMF7156 (along with GMF4877) was identified by screening of the pancreatic β -cell proliferation-promoter for insulin secretion in diabetes (Shen, et al., 2015). Two new inhibitors, FINDY and CaNDY, targeting DYRK1A protein stability/maturation have been developed. These inhibitors suppress intramolecular autophosphorylation of Ser97 in DYRK1A and antagonize the interaction of DYRK1A and the kinase-specific co-chaperone CDC37, thus leading to the degradation of DYRK1A in living cells (Kii, et al., 2016; Sonamoto, et al., 2015). DANDY was also obtained from the search of the common structure supposedly inhibiting DYRK1A (Gourdain, et al., 2013). Recently, its effectiveness in animals has been reported in a DS model with 2 weeks of administration (Neumann, et al., 2018).

Additional inhibitors have been identified by screening for promoter activity of specific target genes. GMF7156 (along with GMF4877) was identified by screening of the pancreatic β -cell proliferation-promoter for insulin secretion in diabetes (Shen, et al., 2015). ALGERNON was first identified from the screening of proliferation-enhancer in NSCs, thereby aiming to correct the aberrant neurogenesis observed in individuals with DS/DS murine models (Nakano-Kobayashi, et al., 2017). ALGERNON is water soluble, distributes to the brain tissue after oral administration, and exhibits good pharmacokinetics. ALGERNON enhances the proliferation of NSCs in the DG of the mouse hippocampus. ALGERNON treatment of pregnant dams prevented morphological brain abnormalities, including the thinned cortical plate, typically observed in Ts1Cje embryonic mice. Remarkably, when compared to untreated trisomic offspring, ALGERNON-treated mice exhibited normal cognitive behavior.

Finally, a DYRK1A inhibitor was identified by a drug repurposing strategy. 5-Iodotubercidin (5-IT), an annotated adenosine kinase inhibitor previously reported to increase proliferation of islets, was found to be a potent and selective inhibitor of DYRK1A and proven to be effective in diabetes models (Dirice, et al., 2016).

9.2 Future of DYRK1A inhibitors: possible applications and safety

Considering the involvement of DYRK1A in psychiatric/developmental diseases such as DS and AD, clinicians and patients have long anticipated the development of practical DYRK1A inhibitors. Of note, a clinical trial of EGCG in patients with DS has been conducted as discussed in section 7 (de la Torre, et al., 2016). EGCG supplementation for 12 months significantly improved visual recognition memory and adaptive behavior after cognitive training compared to those parameters in untreated controls.

ALGERNON is a more specific inhibitor of DYRK1A than EGCG and, as discussed above, has been used prenatally to prevent cognitive deficits in the Ts1Cje mouse model (Nakano-Kobayashi, et al., 2017), thus suggesting the possibility of a prenatal therapy for DS. Prenatal treatment is not realistic at present, but ALGERNON also stimulates adult neurogenesis meaning that ALGERNON has therapeutic potential not only for the treatment of DS but also for a wide range of disorders involving progressive or permanent neuronal loss, including neurodegenerative diseases and traumatic brain injury. ALGERNON may also be applicable for treating the consequences of Zika virus infection. Zika is associated with an increased rate of microcephaly, and DYRK1A is reported to be upregulated in Zika-infected human neural progenitor cells that exhibit dysregulated cell cycle progression and attenuated growth (Tang, et al., 2016).

Before DYRK1A inhibitors can be applied for therapeutic purposes, we need to consider that these inhibitors might be double-edged swords. DYRK1A has crucial roles in brain development, synaptic maturation, and functions (as discussed in previous chapters), and excess inhibition of DYRK1A could lead to deleterious side effects. Truncating mutations that reduce *DYRK1A* gene dose cause microcephaly and autism in humans, and loss-of-function

mutations in *DYRK1A Drosophila* and mice orthologs produce defects in neurogenesis (Courcet, et al., 2012; O'Roak, et al., 2012). Of note, WT offspring of pregnant mice treated with ALGERNON trended toward impaired learning behaviors (Nakano-Kobayashi, et al., 2017), which is reminiscent of MDR7 patients with known *DYRK1A* disruptions (Bronicki, et al., 2015; Moller, et al., 2008; Oegema, et al., 2010; Ruaud, et al., 2015; van Bon, et al., 2016). However, there is information indicating the safety of ALGERNON; prenatal administration of ALGERNON in mice had no effect on the pup weight or other physical or behavioral factors, namely, including rectal temperature; righting reflex; whisker twitch; ear twitch; reaching behavior; wire hang performance; grip strength; reaction to key jangling (Nakano-Kobayashi, et al., 2017); and the pup's performance in several behavioral tests, including the open field test, light/dark transition test, elevated plus maze test, Crawly's social interaction test, tail suspension test, and forced swim test (Nakano-Kobayashi et al., unpublished data). Effective biomarkers that reflect *DYRK1A* activity are needed to establish the clinical therapeutic potential of these *DYRK1A* inhibitors.

10 Conclusion

The large number of substrates phosphorylated by *DYRK1A* and the wide range of interacting partners indicate that *DYRK1A* is capable of controlling a variety of molecular processes. These processes underlie several physiological functions at different stages of life: during neurogenesis at early development, in neuronal plasticity during brain functioning, and during aging. Analysis of chromosomal rearrangements and mutations have revealed that a deficit in the *DYRK1A* gene dosage is a likely cause of MRD7, a form of autism, and an excess in gene dosage is a likely cause of the cognitive alterations present in DS. These observations indicate that *DYRK1A* is a critical gene for synaptopathies. The role of *DYRK1A* in synaptic plasticity has been confirmed in models designed to reproduce alterations of *DYRK1A* levels in *Drosophila*, mice, and zebrafish. Many reports from humans and animal models also indicate a role for *DYRK1A* during neurodegenerative processes: with abnormal truncation events or

abnormal peripheral levels of DYRK1A in patients with AD or with altered phosphorylation of DYRK1A targets involved in α -synuclein dementias or in PD.

The molecular pathways controlling the DYRK1A level and the overall molecular and regulatory controls exerted by DYRK1A on cognitive processes remain to be completely established. Evidence for crosstalk between DYRK1A-related pathways and for regulatory loops is still emerging.

The role of DYRK1A in developmental and neurodegenerative diseases makes this protein kinase an attractive drug target to not only normalize prenatal development and improve cognition in youths and adults with DS but also treat neurodegenerative diseases. However, considering the complexity of DYRK1A targets, adverse effects could be a key issue in clinical trials. Therapeutic risks may arise from i) off-target drug effects, as it is known that other serine threonine kinases can be targeted by DYRK1A inhibitors or ii) excessive inhibition, as it has been shown that severe developmental anomalies can arise when DYRK1A levels fall below normal. In the case of AD, inhibition in the periphery might be associated with adverse effects, for example, inducing an increase in homocysteine. Therefore, the proper objective is to achieve a balanced level of DYRK1A, and this will depend on the controlled use of existing or novel inhibitors; conversely, in the case of MRD7 or other conditions with decreased DYRK1A levels, therapeutic perspectives will rely on the discovery of activators.

Conflict of interest

The authors do not have any conflict of interest to declare.

Table I. DYR1KA interactors, partners, and targets

Functions	Proteins	Subcellular localization	References
Chromatin regulator and Transcription	Arip4	nucleus	(Sitz, et al., 2004)
	Creb1	nucleus	(E. J. Yang, et al., 2001)
	CRY2	nucleus, cytosol	(Kurabayashi, et al., 2010)
	FOXO1	nucleus	(von Groote-Bidlingmaier, et al., 2003)
	Gli1	nucleus	(Ehe, et al., 2017)
	HIST1H3E	nucleus	(Jang, et al., 2014)
	NFAT	nucleus/cytosol	(Arron, et al., 2006; Lee, et al., 2009)
	NOTCH	nucleus	(Fernandez-Martinez, et al., 2009)
	P53	nucleus	(J. Park, et al., 2010)
	RNAPII	nucleus	(Di Vona, et al., 2015)
	SIRT1	nucleus	(Guo, et al., 2010)
	STAT3	nucleus/cytosol	(Wiechmann, et al., 2003)
Splicing	SC35	nucleus	(Qian, et al., 2011)
	SF2/ASF	nucleus	(Shi, et al., 2008)
	SRp55	nucleus	(Yin, et al., 2012)
	9G8	nucleus	(Ding, et al., 2012)
Cell cycle, cell signaling, and pathways	BRAF	cytosol	(Kelly & Rahmani, 2005)
	CASP9	nucleus/cytosol	(Laguna, et al., 2008)
	CCND1	nucleus/cytosol	(Soppa, et al., 2014)
	DCAF7	cytosol	(Skurat & Dietrich, 2004; J. Xiang, et al., 2017)
	GSK3B	nucleus/cytosol	(Song, et al., 2015)
	LIN52	nucleus	(Guiley, et al., 2015)
	MEK1	cytosol	(Kelly & Rahmani, 2005)
	NEP	membrane	(Kawakubo, et al., 2017)
	PAHX-AP1	cytosol	(Bescond & Rahmani, 2005)
	PSEN1	membrane	(Ryu, et al., 2010)
	PRKN	cytosol	(Im & Chung, 2015)
	PROX1, ASCL1	nucleus	(Shaikh, et al., 2016)
	SPRED1	cytosol	(Li, et al., 2010)
	SPRY2	cytosol	(Aranda, et al., 2008)
YWHAE	cytosol	(Alvarez, et al., 2007)	
Synaptic plasticity, endocytosis, and trafficking	SNCA	cytosol, membrane	(Murakami, et al., 2012)
	AMPH	cytosol, membrane	(Murakami, et al., 2006)









	APP	cytosol, membrane	(Ryoo, et al., 2008)
	CLTC	cytosol, membrane	(Murakami, et al., 2012)
	DNM1	cytosol	(Chen-Hwang, et al., 2002)
	SH3GLB1	Cytosol, membrane	(Murakami, et al., 2009)
	GRB2	nucleus, cytosol	(Abekhoukh, et al., 2013; Shin, et al., 2007)
	GRIN2A	membrane	(Grau, et al., 2014)
	HAP1	cytosol	(J. Xiang, et al., 2017)
	HIP1	cytosol, membrane	(Kang, et al., 2005)
	N-WASP	cytosol	(J. Park, et al., 2012)
	SEPT4	cytosol	(Sitz, et al., 2008)
	SYNJ1	cytosol	(Adayev, Chen-Hwang, Murakami, Wang, et al., 2006)
	MAPT (TAU)	cytosol	(Kimura, et al., 2007; Woods, Cohen, et al., 2001)





Table II. DYRK1A implication in synaptic activity

Model	Approach	Phenotype	References
Basal synaptic activity			
Dyrk1a overexpression	cell culture, cortex	↓ spike number	Martinez de Lagran et al., 2012
Mnb overexpression	Drosophila NMJ	↑ mEPSP amplitude ↑ synaptic vesicle size	Chen et al., 2014
Dyrk1a overexpression	Mice, PFC, slice	↑ mEPSC amplitude ↑ dendritic spine density	Thomazeau et al., 2014
Dyrk1a overexpression	Mice, cerebellum, neuron culture	↑ NR2A-containing NMDAR expression ↑ decay NMDA-induced calcium transient	Altafaj et al., 2008
Dyrk1a overexpression	Mice,	E/I imbalance toward inhibition	Souchet, et al., 2014
Ts65Dn crossed with Dyrk1a ^{+/-}	Mice, HPC	Normalization of the density of GABAergic and glutamatergic synapse markers	Garcia-Cerro, et al., 2014
Synaptic plasticity			
Dyrk1a overexpression	Mice, HPC, slice	↑ NMDAR-LTP ↓ PPF-LTD Ø NMDAR-LTD	Ahn et al., 2006
Dyrk1a overexpression	Mice, PFC, slice	Ø NMDAR-LTP Ø eCB-LTD	Thomazeau et al., 2014
Ts65Dn + EGCG	Mice, HPC, slice	NMDAR-LTP protected by EGCG treatment	Xie et al., 2008
Ts65Dn + shRNA Dyrk1a viral injection	Mice, HPC, slice	NMDAR-LTP protected by shRNA Dyrk1a	Altafaj et al., 2013
Ts65Dn crossed with Dyrk1a ^{+/-}	Mice, HPC, slice	NMDAR-LTP protected	Garcia-Cerro et al., 2014

NMJ: neuromuscular junction; PFC: prefrontal cortex; HPC: hippocampus; mEPSP: miniature excitatory postsynaptic potential; mEPSC: miniature excitatory postsynaptic current; NR2A: 2A subunit-containing N-methyl-D-aspartate receptor; NMDA: N-methyl-D-aspartate; E/I: excitation/inhibition; GABA: gamma-aminobutyric acid; NMDAR: N-methyl-D-aspartate receptor; PPF: paired-pulse facilitation; LTP: long-term potentiation; LTD: long-term depression; eCB: endocannabinoid; shRNA: short hairpin ribonucleic acid.

Table III: DYRK1A inhibitors

	Structure	Specificity (other targets except for Clks)	Bioavailability and disease models tested	References
From natural products and their derivatives				
EGCG		proteasomes matrix metalloproteinase dihydrofolate reductase DNA methyltransferase topoisomerase II telomerase COMT Vimentin	Yes, but low DS Cancer	(Bain, et al., 2003; Bain, et al., 2007)
Harmine (β -carboline)		MAO-A	Yes (plant alkaloid)	(Gockler, et al., 2009)
AnnH75 (β -carboline)		Haspin	ND	(Ruben, et al., 2015)
Leucettine L41		GSK3 α/β	Yes AD model	(Debdab, et al., 2011; Naert, et al., 2015; Tahtouh, et al., 2012)
Meridianin (31)		CDK5	ND	(Giraud, et al., 2011; Yadav, et al., 2015)
INDY type (TG003 derivatives)				
INDY		Pim1, CDK2, CK1, GSK3 α/β *	No INDY tested in <i>Xenopus</i> embryos with DYRK1A- overexpression	(Ogawa, et al., 2010)
BINDY		Pims, CHK2, Pims, CDK2, FLT3**	ND	(Masaki, et al., 2015)
Targeting DYRK1A protein stability				
CaNDY		Haspin, CK2, CDK9**	ND	(Sonamoto, et al., 2015)

FINDY		Gsk3 β , MARK4, Pims, PLK3*	ND	(Kii, et al., 2016)
From cell-based functional assay				
GNF7156		GSK3 β	ND Diabetes	(Shen, et al., 2015)
ALGERNON		CDK2, CDK5, Haspin, HGK, ROCKs**	Yes DS	(Nakano-Kobayashi, et al., 2017)
From the drug repurposing				
5-IT		GSK3 β	Yes Diabetes	(Dirice, et al., 2016)

* determined by kinase panel assay at 10 μ M (INDY and FINDY) and **1 μ M (BINDY, CaNDY, and ALGERNON) concentrations of compounds.

Figure legends:

Figure 1. (A) Representation of the domain structure of DYRK1A: NLS, nuclear localization signal; DH, DYRK homology domain; PEST, domain enriched in proline, glutamic acid, serine, and threonine residues; His, histidine repeats; S/T, serine and threonine-rich region. **(B)** Structure of the DYRK1A–harmine complex. The kinase domain (amino acids 126–490) of human DYRK1A was expressed in *E. coli* and co-crystallized with harmine (3ANR in ProteinDataBase) (NGL view with harmine in the spacefill mode) (from Ogawa, 2010).

Figure 2. Main functional roles of DYRK1A and proteins described as interactants or targets.

Figure 3. Scheme illustrating the DYRK1A-mediated functions/activities involved in neural cell differentiation in the developing neocortex.

Neurons and glial cells are produced sequentially from pluripotent radial glial progenitors (apical progenitors) or from progenitors with a more restricted fate (basal progenitors (IPs) and OPCs). During neurogenesis, DYRK1A regulates the neurogenic potential of radial glial progenitors regulating the expression of cell cycle regulator proteins (Cyclin D1, p27^{Kip1}, and p21^{Cip}) involved in the G1 to S phase transition. DYRK1A phosphorylation of NFATc transcription factors may regulate neuronal differentiation. Moreover, DYRK1A regulation of the pro-astroglial factor STAT may have a role in astrocyte production. CP: cortical plate; IP; intermediate progenitor; IZ: intermediate zone; NE: neuroepithelial cell; OPC: oligodendrocyte precursor; SVZ: subventricular zone; VZ: ventricular zone.

Figure 4: Schematic model illustrating the putative involvements of DYRK1A at the developing and the mature synapse.

DYRK1A regulates the exocytosis–endocytosis cycle of synaptic vesicles containing neurotransmitter in the presynaptic bouton and of receptors located at the postsynaptic membrane by phosphorylating core proteins such as dynamin 1, amphiphysin 1, synaptojanin 1, endophilin, or adaptor proteins such as AP180 (1).

The presynaptic vesicle cluster and the postsynaptic compartment throughout the spine head are surrounded by a dense scaffold, the cytoskeleton, mainly composed of actin filaments and microtubules. DYRK1A is involved in cytoskeletal rearrangement by direct or indirect interactions with actin filaments, microtubules, or the microtubule-associated protein tau (2).

DYRK1A can also impact the synapse formation and activity by regulating the transcription machinery, by its direct or indirect effects on transcription factors such as CREB, NFAT, or FKHR, or transcripts such as neuroligin, TrkBT1, BDNF, AchE-S, and AchE-R (3).

AP180: clathrin adaptor protein; CREB: cAMP responsive element-binding protein; NFAT: nuclear factor of activated T cells; FKHR: forkhead transcription factor; TrkBT1: truncated Tropomyosin receptor kinase B receptor; BDNF: Brain-derived neurotrophic factor; AchE-S: acetylcholinesterase-S; AchE-R: acetylcholinesterase-R.

Figure 5: Consequences of increased DYRK1A level on β -amyloidosis and neurofibrillary degeneration.

References

- Abekhoukh, S., Planque, C., Ripoll, C., Urbaniak, P., Paul, J. L., Delabar, J. M., & Janel, N. (2013). Dyrk1A, a serine/threonine kinase, is involved in ERK and Akt activation in the brain of hyperhomocysteinemic mice. *Mol Neurobiol*, *47*, 105-116.
- Adayev, T., Chen-Hwang, M. C., Murakami, N., Wang, R., & Hwang, Y. W. (2006). MNB/DYRK1A phosphorylation regulates the interactions of synaptojanin 1 with endocytic accessory proteins. *Biochem Biophys Res Commun*, *351*, 1060-1065.
- Adayev, T., Chen-Hwang, M. C., Murakami, N., Wegiel, J., & Hwang, Y. W. (2006). Kinetic properties of a MNB/DYRK1A mutant suitable for the elucidation of biochemical pathways. *Biochemistry*, *45*, 12011-12019.
- Ahn, K. J., Jeong, H. K., Choi, H. S., Ryoo, S. R., Kim, Y. J., Goo, J. S., Choi, S. Y., Han, J. S., Ha, I., & Song, W. J. (2006). DYRK1A BAC transgenic mice show altered synaptic plasticity with learning and memory defects. *Neurobiol Dis*, *22*, 463-472.
- Ait Yahya-Graison, E., Aubert, J., Dauphinot, L., Rivals, I., Prieur, M., Golfier, G., Rossier, J., Personnaz, L., Creau, N., Blehaut, H., Robin, S., Delabar, J. M., & Potier, M. C. (2007). Classification of human chromosome 21 gene-expression variations in Down syndrome: impact on disease phenotypes. *Am J Hum Genet*, *81*, 475-491.
- Alexeeva, M., Aberg, E., Engh, R. A., & Rothweiler, U. (2015). The structure of a dual-specificity tyrosine phosphorylation-regulated kinase 1A-PKC412 complex reveals disulfide-bridge formation with the anomalous catalytic loop HRD(HCD) cysteine. *Acta Crystallogr D Biol Crystallogr*, *71*, 1207-1215.
- Allanson, J. E., O'Hara, P., Farkas, L. G., & Nair, R. C. (1993). Anthropometric craniofacial pattern profiles in Down syndrome. *Am J Med Genet*, *47*, 748-752.
- Altafaj, X., Dierssen, M., Baamonde, C., Marti, E., Visa, J., Guimera, J., Oset, M., Gonzalez, J. R., Florez, J., Fillat, C., & Estivill, X. (2001). Neurodevelopmental delay, motor abnormalities and cognitive deficits in transgenic mice overexpressing Dyrk1A (minibrain), a murine model of Down's syndrome. *Hum Mol Genet*, *10*, 1915-1923.
- Altafaj, X., Martin, E. D., Ortiz-Abalia, J., Valderrama, A., Lao-Peregrin, C., Dierssen, M., & Fillat, C. (2013). Normalization of Dyrk1A expression by AAV2/1-shDyrk1A attenuates hippocampal-dependent defects in the Ts65Dn mouse model of Down syndrome. *Neurobiol Dis*, *52*, 117-127.
- Altafaj, X., Ortiz-Abalia, J., Fernandez, M., Potier, M. C., Laffaire, J., Andreu, N., Dierssen, M., Gonzalez-Garcia, C., Cena, V., Marti, E., & Fillat, C. (2008). Increased NR2A expression and prolonged decay of NMDA-induced calcium transient in cerebellum of TgDyrk1A mice, a mouse model of Down syndrome. *Neurobiol Dis*, *32*, 377-384.
- Alvarez-Buylla, A., & Lim, D. A. (2004). For the long run: maintaining germinal niches in the adult brain. *Neuron*, *41*, 683-686.
- Alvarez, M., Altafaj, X., Aranda, S., & de la Luna, S. (2007). DYRK1A autophosphorylation on serine residue 520 modulates its kinase activity via 14-3-3 binding. *Mol Biol Cell*, *18*, 1167-1178.
- Alvarez, M., Estivill, X., & de la Luna, S. (2003). DYRK1A accumulates in splicing speckles through a novel targeting signal and induces speckle disassembly. *J Cell Sci*, *116*, 3099-3107.
- Anantharam, A., Bittner, M. A., Aikman, R. L., Stuenkel, E. L., Schmid, S. L., Axelrod, D., & Holz, R. W. (2011). A new role for the dynamin GTPase in the regulation of fusion pore expansion. *Mol Biol Cell*, *22*, 1907-1918.
- Anderson, K., Chen, Y., Chen, Z., Dominique, R., Glenn, K., He, Y., Janson, C., Luk, K. C., Lukacs, C., Polonskaia, A., Qiao, Q., Railkar, A., Rossman, P., Sun, H., Xiang, Q., Vilenchik, M., Wovkulich, P., & Zhang, X. (2013). Pyrido[2,3-d]pyrimidines: discovery and preliminary SAR of a novel series of DYRK1B and DYRK1A inhibitors. *Bioorg Med Chem Lett*, *23*, 6610-6615.

- Andreu-Agullo, C., Morante-Redolat, J. M., Delgado, A. C., & Farinas, I. (2009). Vascular niche factor PEDF modulates Notch-dependent stemness in the adult subependymal zone. *Nat Neurosci*, *12*, 1514-1523.
- Anneren, G., & Edman, B. (1993). Down syndrome--a gene dosage disease caused by trisomy of genes within a small segment of the long arm of chromosome 21, exemplified by the study of effects from the superoxide-dismutase type 1 (SOD-1) gene. *APMIS Suppl*, *40*, 71-79.
- Aranda, S., Alvarez, M., Turro, S., Laguna, A., & de la Luna, S. (2008). Sprouty2-mediated inhibition of fibroblast growth factor signaling is modulated by the protein kinase DYRK1A. *Mol Cell Biol*, *28*, 5899-5911.
- Aranda, S., Laguna, A., & de la Luna, S. (2011). DYRK family of protein kinases: evolutionary relationships, biochemical properties, and functional roles. *FASEB J*, *25*, 449-462.
- Arque, G., Casanovas, A., & Dierssen, M. (2013). Dyrk1A is dynamically expressed on subsets of motor neurons and in the neuromuscular junction: possible role in Down syndrome. *PLoS One*, *8*, e54285.
- Arque, G., Fotaki, V., Fernandez, D., Martinez de Lagran, M., Arbones, M. L., & Dierssen, M. (2008). Impaired spatial learning strategies and novel object recognition in mice haploinsufficient for the dual specificity tyrosine-regulated kinase-1A (Dyrk1A). *PLoS One*, *3*, e2575.
- Arron, J. R., Winslow, M. M., Polleri, A., Chang, C. P., Wu, H., Gao, X., Neilson, J. R., Chen, L., Heit, J. J., Kim, S. K., Yamasaki, N., Miyakawa, T., Francke, U., Graef, I. A., & Crabtree, G. R. (2006). NFAT dysregulation by increased dosage of DSCR1 and DYRK1A on chromosome 21. *Nature*, *441*, 595-600.
- Azorsa, D. O., Robeson, R. H., Frost, D., Meec hoovet, B., Brautigam, G. R., Dickey, C., Beaudry, C., Basu, G. D., Holz, D. R., Hernandez, J. A., Bisanz, K. M., Gwinn, L., Grover, A., Rogers, J., Reiman, E. M., Hutton, M., Stephan, D. A., Mousses, S., & Dunckley, T. (2010). High-content siRNA screening of the kinome identifies kinases involved in Alzheimer's disease-related tau hyperphosphorylation. *BMC Genomics*, *11*, 25.
- Bain, J., McLauchlan, H., Elliott, M., & Cohen, P. (2003). The specificities of protein kinase inhibitors: an update. *Biochem J*, *371*, 199-204.
- Bain, J., Plater, L., Elliott, M., Shpiro, N., Hastie, C. J., McLauchlan, H., Klevernic, I., Arthur, J. S., Alessi, D. R., & Cohen, P. (2007). The selectivity of protein kinase inhibitors: a further update. *Biochem J*, *408*, 297-315.
- Bancher, C., Grundke-Iqbal, I., Iqbal, K., Fried, V. A., Smith, H. T., & Wisniewski, H. M. (1991). Abnormal phosphorylation of tau precedes ubiquitination in neurofibrillary pathology of Alzheimer disease. *Brain Res*, *539*, 11-18.
- Barallobre, M. J., Perier, C., Bove, J., Laguna, A., Delabar, J. M., Vila, M., & Arbones, M. L. (2014). DYRK1A promotes dopaminergic neuron survival in the developing brain and in a mouse model of Parkinson's disease. *Cell Death Dis*, *5*, e1289.
- Becker, L. E., Armstrong, D. L., & Chan, F. (1986). Dendritic atrophy in children with Down's syndrome. *Ann Neurol*, *20*, 520-526.
- Becker, W., & Joost, H. G. (1999). Structural and functional characteristics of Dyrk, a novel subfamily of protein kinases with dual specificity. *Prog Nucleic Acid Res Mol Biol*, *62*, 1-17.
- Becker, W., Weber, Y., Wetzels, K., Eirnbter, K., Tejedor, F. J., & Joost, H. G. (1998). Sequence characteristics, subcellular localization, and substrate specificity of DYRK-related kinases, a novel family of dual specificity protein kinases. *J Biol Chem*, *273*, 25893-25902.
- Belichenko, N. P., Belichenko, P. V., Kleschevnikov, A. M., Salehi, A., Reeves, R. H., & Mobley, W. C. (2009). The "Down syndrome critical region" is sufficient in the mouse model to confer behavioral, neurophysiological, and synaptic phenotypes characteristic of Down syndrome. *J Neurosci*, *29*, 5938-5948.
- Benavides-Piccione, R., Dierssen, M., Ballesteros-Yanez, I., Martinez de Lagran, M., Arbones, M. L., Fotaki, V., DeFelipe, J., & Elston, G. N. (2005). Alterations in the phenotype of neocortical pyramidal cells in the Dyrk1A^{+/-} mouse. *Neurobiol Dis*, *20*, 115-122.

- Bescond, M., & Rahmani, Z. (2005). Dual-specificity tyrosine-phosphorylated and regulated kinase 1A (DYRK1A) interacts with the phytanoyl-CoA alpha-hydroxylase associated protein 1 (PAHX-AP1), a brain specific protein. *Int J Biochem Cell Biol*, *37*, 775-783.
- Bradley, C. A., Peineau, S., Taghibiglou, C., Nicolas, C. S., Whitcomb, D. J., Bortolotto, Z. A., Kaang, B. K., Cho, K., Wang, Y. T., & Collingridge, G. L. (2012). A pivotal role of GSK-3 in synaptic plasticity. *Front Mol Neurosci*, *5*, 13.
- Branca, C., Shaw, D. M., Belfiore, R., Gokhale, V., Shaw, A. Y., Foley, C., Smith, B., Hulme, C., Dunckley, T., Meechoovet, B., Caccamo, A., & Oddo, S. (2017). Dyrk1 inhibition improves Alzheimer's disease-like pathology. *Aging Cell*, *16*, 1146-1154.
- Bronicki, L. M., Redin, C., Drunat, S., Piton, A., Lyons, M., Passemard, S., Baumann, C., Faivre, L., Thevenon, J., Riviere, J. B., Isidor, B., Gan, G., Francannet, C., Willems, M., Gunel, M., Jones, J. R., Gleeson, J. G., Mandel, J. L., Stevenson, R. E., Friez, M. J., & Aylsworth, A. S. (2015). Ten new cases further delineate the syndromic intellectual disability phenotype caused by mutations in DYRK1A. *Eur J Hum Genet*, *23*, 1482-1487.
- Burre, J. (2015). The Synaptic Function of alpha-Synuclein. *J Parkinsons Dis*, *5*, 699-713.
- Catuara-Solarz, S., Espinosa-Carrasco, J., Erb, I., Langohr, K., Gonzalez, J. R., Notredame, C., & Dierssen, M. (2016). Combined Treatment With Environmental Enrichment and (-)-Epigallocatechin-3-Gallate Ameliorates Learning Deficits and Hippocampal Alterations in a Mouse Model of Down Syndrome. *eNeuro*, *3*.
- Caviness, V. S., Jr., Takahashi, T., & Nowakowski, R. S. (1995). Numbers, time and neocortical neuronogenesis: a general developmental and evolutionary model. *Trends Neurosci*, *18*, 379-383.
- Cen, L., Xiao, Y., Wei, L., Mo, M., Chen, X., Li, S., Yang, X., Huang, Q., Qu, S., Pei, Z., & Xu, P. (2016). Association of DYRK1A polymorphisms with sporadic Parkinson's disease in Chinese Han population. *Neurosci Lett*, *632*, 39-43.
- Cepko, C. (2014). Intrinsically different retinal progenitor cells produce specific types of progeny. *Nat Rev Neurosci*, *15*, 615-627.
- Ceron, J., Gonzalez, C., & Tejedor, F. J. (2001). Patterns of cell division and expression of asymmetric cell fate determinants in postembryonic neuroblast lineages of *Drosophila*. *Dev Biol*, *230*, 125-138.
- Chakrabarti, L., Galdzicki, Z., & Haydar, T. F. (2007). Defects in embryonic neurogenesis and initial synapse formation in the forebrain of the Ts65Dn mouse model of Down syndrome. *J Neurosci*, *27*, 11483-11495.
- Chan, S. A., Doreian, B., & Smith, C. (2010). Dynamin and myosin regulate differential exocytosis from mouse adrenal chromaffin cells. *Cell Mol Neurobiol*, *30*, 1351-1357.
- Chen-Hwang, M. C., Chen, H. R., Elzinga, M., & Hwang, Y. W. (2002). Dynamin is a minibrain kinase/dual specificity Yak1-related kinase 1A substrate. *J Biol Chem*, *277*, 17597-17604.
- Chen, C. K., Bregere, C., Paluch, J., Lu, J. F., Dickman, D. K., & Chang, K. T. (2014). Activity-dependent facilitation of Synaptojanin and synaptic vesicle recycling by the Minibrain kinase. *Nat Commun*, *5*, 4246.
- Chen, J. Y., Lin, J. R., Tsai, F. C., & Meyer, T. (2013). Dosage of Dyrk1a shifts cells within a p21-cyclin D1 signaling map to control the decision to enter the cell cycle. *Mol Cell*, *52*, 87-100.
- Chen, L., & Feany, M. B. (2005). Alpha-synuclein phosphorylation controls neurotoxicity and inclusion formation in a *Drosophila* model of Parkinson disease. *Nat Neurosci*, *8*, 657-663.
- Chen, L., Periquet, M., Wang, X., Negro, A., McLean, P. J., Hyman, B. T., & Feany, M. B. (2009). Tyrosine and serine phosphorylation of alpha-synuclein have opposing effects on neurotoxicity and soluble oligomer formation. *J Clin Invest*, *119*, 3257-3265.
- Chettouh, Z., Croquette, M. F., Delobel, B., Gilgenkrants, S., Leonard, C., Maunoury, C., Prieur, M., Rethore, M. O., Sinet, P. M., Chery, M., & et al. (1995). Molecular mapping of 21 features associated with partial monosomy 21: involvement of the APP-SOD1 region. *Am J Hum Genet*, *57*, 62-71.

- Colon-Ramos, D. A. (2009). Synapse formation in developing neural circuits. *Curr Top Dev Biol*, *87*, 53-79.
- Courcet, J. B., Faivre, L., Malzac, P., Masurel-Paulet, A., Lopez, E., Callier, P., Lambert, L., Lemesle, M., Thevenon, J., Gigot, N., Duplomb, L., Ragon, C., Marle, N., Mosca-Boidron, A. L., Huet, F., Philippe, C., Moncla, A., & Thauvin-Robinet, C. (2012). The DYRK1A gene is a cause of syndromic intellectual disability with severe microcephaly and epilepsy. *J Med Genet*, *49*, 731-736.
- da Costa Martins, P. A., Salic, K., Gladka, M. M., Armand, A. S., Leptidis, S., el Azzouzi, H., Hansen, A., Coenen-de Roo, C. J., Bierhuizen, M. F., van der Nagel, R., van Kuik, J., de Weger, R., de Bruin, A., Condorelli, G., Arbones, M. L., Eschenhagen, T., & De Windt, L. J. (2010). MicroRNA-199b targets the nuclear kinase Dyrk1a in an auto-amplification loop promoting calcineurin/NFAT signalling. *Nat Cell Biol*, *12*, 1220-1227.
- Dabrowski, A., Terauchi, A., Strong, C., & Umemori, H. (2015). Distinct sets of FGF receptors sculpt excitatory and inhibitory synaptogenesis. *Development*, *142*, 1818-1830.
- Dang, T., Duan, W. Y., Yu, B., Tong, D. L., Cheng, C., Zhang, Y. F., Wu, W., Ye, K., Zhang, W. X., Wu, M., Wu, B. B., An, Y., Qiu, Z. L., & Wu, B. L. (2017). Autism-associated Dyrk1a truncation mutants impair neuronal dendritic and spine growth and interfere with postnatal cortical development. *Mol Psychiatry*.
- de Graaf, K., Czajkowska, H., Rottmann, S., Packman, L. C., Lilischkis, R., Luscher, B., & Becker, W. (2006). The protein kinase DYRK1A phosphorylates the splicing factor SF3b1/SAP155 at Thr434, a novel in vivo phosphorylation site. *BMC Biochem*, *7*, 7.
- de Graaf, K., Hekerman, P., Spelten, O., Herrmann, A., Packman, L. C., Bussow, K., Muller-Newen, G., & Becker, W. (2004). Characterization of cyclin L2, a novel cyclin with an arginine/serine-rich domain: phosphorylation by DYRK1A and colocalization with splicing factors. *J Biol Chem*, *279*, 4612-4624.
- de la Torre, R., de Sola, S., Hernandez, G., Farre, M., Pujol, J., Rodriguez, J., Espadaler, J. M., Langohr, K., Cuenca-Royo, A., Principe, A., Xicota, L., Janel, N., Catuara-Solarz, S., Sanchez-Benavides, G., Blehaut, H., Duenas-Espin, I., Del Hoyo, L., Benejam, B., Blanco-Hinojo, L., Videla, S., Fito, M., Delabar, J. M., Dierssen, M., & group, T. s. (2016). Safety and efficacy of cognitive training plus epigallocatechin-3-gallate in young adults with Down's syndrome (TESDAD): a double-blind, randomised, placebo-controlled, phase 2 trial. *Lancet Neurol*, *15*, 801-810.
- De la Torre, R., De Sola, S., Pons, M., Duchon, A., de Lagran, M. M., Farre, M., Fito, M., Benejam, B., Langohr, K., Rodriguez, J., Pujadas, M., Bizot, J. C., Cuenca, A., Janel, N., Catuara, S., Covas, M. I., Blehaut, H., Herault, Y., Delabar, J. M., & Dierssen, M. (2014). Epigallocatechin-3-gallate, a DYRK1A inhibitor, rescues cognitive deficits in Down syndrome mouse models and in humans. *Mol Nutr Food Res*, *58*, 278-288.
- De Rubeis, S., He, X., Goldberg, A. P., Poultney, C. S., Samocha, K., Cicek, A. E., Kou, Y., Liu, L., Fromer, M., Walker, S., Singh, T., Klei, L., Kosmicki, J., Shih-Chen, F., Aleksic, B., Biscaldi, M., Bolton, P. F., Brownfeld, J. M., Cai, J., Campbell, N. G., Carracedo, A., Chahrour, M. H., Chiocchetti, A. G., Coon, H., Crawford, E. L., Curran, S. R., Dawson, G., Duketis, E., Fernandez, B. A., Gallagher, L., Geller, E., Guter, S. J., Hill, R. S., Ionita-Laza, J., Jimenez Gonzalez, P., Kilpinen, H., Klauck, S. M., Klevzon, A., Lee, I., Lei, I., Lei, J., Lehtimaki, T., Lin, C. F., Ma'ayan, A., Marshall, C. R., McInnes, A. L., Neale, B., Owen, M. J., Ozaki, N., Parellada, M., Parr, J. R., Purcell, S., Puura, K., Rajagopalan, D., Rehnstrom, K., Reichenberg, A., Sabo, A., Sachse, M., Sanders, S. J., Schafer, C., Schulte-Ruther, M., Skuse, D., Stevens, C., Szatmari, P., Tammimies, K., Valladares, O., Voran, A., Li-San, W., Weiss, L. A., Willsey, A. J., Yu, T. W., Yuen, R. K., Study, D. D., Homozygosity Mapping Collaborative for, A., Consortium, U. K., Cook, E. H., Freitag, C. M., Gill, M., Hultman, C. M., Lehner, T., Palotie, A., Schellenberg, G. D., Sklar, P., State, M. W., Sutcliffe, J. S., Walsh, C. A., Scherer, S. W., Zwick, M. E., Barrett, J. C., Cutler, D. J., Roeder, K., Devlin, B., Daly, M. J., & Buxbaum, J. D. (2014). Synaptic, transcriptional and chromatin genes disrupted in autism. *Nature*, *515*, 209-215.

- Debdab, M., Carreaux, F., Renault, S., Soundararajan, M., Fedorov, O., Filippakopoulos, P., Lozach, O., Babault, L., Tahtouh, T., Baratte, B., Ogawa, Y., Hagiwara, M., Eisenreich, A., Rauch, U., Knapp, S., Meijer, L., & Bazureau, J. P. (2011). Leucettines, a class of potent inhibitors of cdc2-like kinases and dual specificity, tyrosine phosphorylation regulated kinases derived from the marine sponge leucettamine B: modulation of alternative pre-RNA splicing. *J Med Chem*, *54*, 4172-4186.
- Deciphering Developmental Disorders, S. (2015). Large-scale discovery of novel genetic causes of developmental disorders. *Nature*, *519*, 223-228.
- Degoutin, J. L., Milton, C. C., Yu, E., Tipping, M., Bosveld, F., Yang, L., Bellaiche, Y., Veraksa, A., & Harvey, K. F. (2013). Riquiqui and minibrain are regulators of the hippo pathway downstream of Dachshous. *Nat Cell Biol*, *15*, 1176-1185.
- Dehay, C., & Kennedy, H. (2007). Cell-cycle control and cortical development. *Nat Rev Neurosci*, *8*, 438-450.
- Delabar, J. M., Theophile, D., Rahmani, Z., Chettouh, Z., Blouin, J. L., Prieur, M., Noel, B., & Sinet, P. M. (1993). Molecular mapping of twenty-four features of Down syndrome on chromosome 21. *Eur J Hum Genet*, *1*, 114-124.
- Di Vona, C., Bezdán, D., Islam, A. B., Salichs, E., Lopez-Bigas, N., Ossowski, S., & de la Luna, S. (2015). Chromatin-wide profiling of DYRK1A reveals a role as a gene-specific RNA polymerase II CTD kinase. *Mol Cell*, *57*, 506-520.
- Dierssen, M., & Ramakers, G. J. (2006). Dendritic pathology in mental retardation: from molecular genetics to neurobiology. *Genes Brain Behav*, *5 Suppl 2*, 48-60.
- Ding, S., Shi, J., Qian, W., Iqbal, K., Grundke-Iqbal, I., Gong, C. X., & Liu, F. (2012). Regulation of alternative splicing of tau exon 10 by 9G8 and Dyrk1A. *Neurobiol Aging*, *33*, 1389-1399.
- Dirice, E., Walpita, D., Vetere, A., Meier, B. C., Kahraman, S., Hu, J., Dancik, V., Burns, S. M., Gilbert, T. J., Olson, D. E., Clemons, P. A., Kulkarni, R. N., & Wagner, B. K. (2016). Inhibition of DYRK1A Stimulates Human beta-Cell Proliferation. *Diabetes*, *65*, 1660-1671.
- Dowjat, K., Adayev, T., Kaczmarek, W., Wegiel, J., & Hwang, Y. W. (2012). Gene dosage-dependent association of DYRK1A with the cytoskeleton in the brain and lymphocytes of down syndrome patients. *J Neuropathol Exp Neurol*, *71*, 1100-1112.
- Dowjat, W. K., Adayev, T., Kuchna, I., Nowicki, K., Palmieriello, S., Hwang, Y. W., & Wegiel, J. (2007). Trisomy-driven overexpression of DYRK1A kinase in the brain of subjects with Down syndrome. *Neurosci Lett*, *413*, 77-81.
- Drung, B., Scholz, C., Barbosa, V. A., Nazari, A., Sarragiotto, M. H., & Schmidt, B. (2014). Computational & experimental evaluation of the structure/activity relationship of beta-carbolines as DYRK1A inhibitors. *Bioorg Med Chem Lett*, *24*, 4854-4860.
- Earl, R. K., Turner, T. N., Mefford, H. C., Hudac, C. M., Gerds, J., Eichler, E. E., & Bernier, R. A. (2017). Clinical phenotype of ASD-associated DYRK1A haploinsufficiency. *Mol Autism*, *8*, 54.
- Ehe, B. K., Lamson, D. R., Tarpley, M., Onyenwoke, R. U., Graves, L. M., & Williams, K. P. (2017). Identification of a DYRK1A-mediated phosphorylation site within the nuclear localization sequence of the hedgehog transcription factor GLI1. *Biochem Biophys Res Commun*, *491*, 767-772.
- Evers, J. M., Laskowski, R. A., Bertolli, M., Clayton-Smith, J., Deshpande, C., Eason, J., Elmslie, F., Flinter, F., Gardiner, C., Hurst, J. A., Kingston, H., Kini, U., Lampe, A. K., Lim, D., Male, A., Naik, S., Parker, M. J., Price, S., Robert, L., Sarkar, A., Straub, V., Woods, G., Thornton, J. M., Study, D. D. D., & Wright, C. F. (2017). Structural analysis of pathogenic mutations in the DYRK1A gene in patients with developmental disorders. *Hum Mol Genet*, *26*, 519-526.
- Falke, H., Chaikuad, A., Becker, A., Loaec, N., Lozach, O., Abu Jhaisha, S., Becker, W., Jones, P. G., Preu, L., Baumann, K., Knapp, S., Meijer, L., & Kunick, C. (2015). 10-iodo-11H-indolo[3,2-c]quinoline-6-carboxylic acids are selective inhibitors of DYRK1A. *J Med Chem*, *58*, 3131-3143.
- Fan, F., Funk, L., & Lou, X. (2016). Dynamin 1- and 3-Mediated Endocytosis Is Essential for the Development of a Large Central Synapse In Vivo. *J Neurosci*, *36*, 6097-6115.

- Fan, K., Tang, B. S., Wang, Y. Q., Kang, J. F., Li, K., Liu, Z. H., Sun, Q. Y., Xu, Q., Yan, X. X., & Guo, J. F. (2016). The GBA, DYRK1A and MS4A6A polymorphisms influence the age at onset of Chinese Parkinson patients. *Neurosci Lett*, *621*, 133-136.
- Fernandez-Martinez, J., Vela, E. M., Tora-Ponsioen, M., Ocana, O. H., Nieto, M. A., & Galceran, J. (2009). Attenuation of Notch signalling by the Down-syndrome-associated kinase DYRK1A. *J Cell Sci*, *122*, 1574-1583.
- Fernandez, F., Morishita, W., Zuniga, E., Nguyen, J., Blank, M., Malenka, R. C., & Garner, C. C. (2007). Pharmacotherapy for cognitive impairment in a mouse model of Down syndrome. *Nat Neurosci*, *10*, 411-413.
- Ferrer, I., Barrachina, M., Puig, B., Martinez de Lagran, M., Marti, E., Avila, J., & Dierssen, M. (2005). Constitutive Dyrk1A is abnormally expressed in Alzheimer disease, Down syndrome, Pick disease, and related transgenic models. *Neurobiol Dis*, *20*, 392-400.
- Ferron, S. R., Pozo, N., Laguna, A., Aranda, S., Porlan, E., Moreno, M., Fillat, C., de la Luna, S., Sanchez, P., Arbones, M. L., & Farinas, I. (2010). Regulated segregation of kinase Dyrk1A during asymmetric neural stem cell division is critical for EGFR-mediated biased signaling. *Cell Stem Cell*, *7*, 367-379.
- Fischbach, G. D., & Lord, C. (2010). The Simons Simplex Collection: a resource for identification of autism genetic risk factors. *Neuron*, *68*, 192-195.
- Fotaki, V., Dierssen, M., Alcantara, S., Martinez, S., Marti, E., Casas, C., Visa, J., Soriano, E., Estivill, X., & Arbones, M. L. (2002). Dyrk1A haploinsufficiency affects viability and causes developmental delay and abnormal brain morphology in mice. *Mol Cell Biol*, *22*, 6636-6647.
- Fotaki, V., Martinez De Lagran, M., Estivill, X., Arbones, M., & Dierssen, M. (2004). Haploinsufficiency of Dyrk1A in mice leads to specific alterations in the development and regulation of motor activity. *Behav Neurosci*, *118*, 815-821.
- Franco, S. J., & Muller, U. (2013). Shaping our minds: stem and progenitor cell diversity in the mammalian neocortex. *Neuron*, *77*, 19-34.
- Frost, D., Meechoovet, B., Wang, T., Gately, S., Giorgetti, M., Shcherbakova, I., & Dunckley, T. (2011). beta-carboline compounds, including harmine, inhibit DYRK1A and tau phosphorylation at multiple Alzheimer's disease-related sites. *PLoS One*, *6*, e19264.
- Fujita, H., Torii, C., Kosaki, R., Yamaguchi, S., Kudoh, J., Hayashi, K., Takahashi, T., & Kosaki, K. (2010). Microdeletion of the Down syndrome critical region at 21q22. *Am J Med Genet A*, *152A*, 950-953.
- Fujiwara, H., Hasegawa, M., Dohmae, N., Kawashima, A., Masliah, E., Goldberg, M. S., Shen, J., Takio, K., & Iwatsubo, T. (2002). alpha-Synuclein is phosphorylated in synucleinopathy lesions. *Nat Cell Biol*, *4*, 160-164.
- Gao, J., Yang, X., Yin, P., Hu, W., Liao, H., Miao, Z., Pan, C., & Li, N. (2012). The involvement of FoxO in cell survival and chemosensitivity mediated by Mirk/Dyrk1B in ovarian cancer. *Int J Oncol*, *40*, 1203-1209.
- Garcia-Cerro, S., Martinez, P., Vidal, V., Corrales, A., Florez, J., Vidal, R., Rueda, N., Arbones, M. L., & Martinez-Cue, C. (2014). Overexpression of Dyrk1A is implicated in several cognitive, electrophysiological and neuromorphological alterations found in a mouse model of Down syndrome. *PLoS One*, *9*, e106572.
- Garcia-Cerro, S., Rueda, N., Vidal, V., Lantigua, S., & Martinez-Cue, C. (2017). Normalizing the gene dosage of Dyrk1A in a mouse model of Down syndrome rescues several Alzheimer's disease phenotypes. *Neurobiol Dis*, *106*, 76-88.
- Garrett, S., & Broach, J. (1989). Loss of Ras activity in *Saccharomyces cerevisiae* is suppressed by disruptions of a new kinase gene, YAK1, whose product may act downstream of the cAMP-dependent protein kinase. *Genes Dev*, *3*, 1336-1348.
- Geng, J., Wang, L., Lee, J. Y., Chen, C. K., & Chang, K. T. (2016). Phosphorylation of Synaptojanin Differentially Regulates Endocytosis of Functionally Distinct Synaptic Vesicle Pools. *J Neurosci*, *36*, 8882-8894.

- Gervais, F. G., Singaraja, R., Xanthoudakis, S., Gutekunst, C. A., Leavitt, B. R., Metzler, M., Hackam, A. S., Tam, J., Vaillancourt, J. P., Houtzager, V., Rasper, D. M., Roy, S., Hayden, M. R., & Nicholson, D. W. (2002). Recruitment and activation of caspase-8 by the Huntingtin-interacting protein Hip-1 and a novel partner Hip1. *Nat Cell Biol*, *4*, 95-105.
- Giraud, F., Alves, G., Debiton, E., Nauton, L., Thery, V., Durieu, E., Ferandin, Y., Lozach, O., Meijer, L., Anizon, F., Pereira, E., & Moreau, P. (2011). Synthesis, protein kinase inhibitory potencies, and in vitro antiproliferative activities of meridianin derivatives. *J Med Chem*, *54*, 4474-4489.
- Gockler, N., Jofre, G., Papadopoulos, C., Soppa, U., Tejedor, F. J., & Becker, W. (2009). Harmine specifically inhibits protein kinase DYRK1A and interferes with neurite formation. *FEBS J*, *276*, 6324-6337.
- Goedert, M., Spillantini, M. G., Jakes, R., Rutherford, D., & Crowther, R. A. (1989). Multiple isoforms of human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of Alzheimer's disease. *Neuron*, *3*, 519-526.
- Goldstein, L. S. (2012). Axonal transport and neurodegenerative disease: can we see the elephant? *Prog Neurobiol*, *99*, 186-190.
- Gonzalez-Jamett, A. M., Baez-Matus, X., Hevia, M. A., Guerra, M. J., Olivares, M. J., Martinez, A. D., Neely, A., & Cardenas, A. M. (2010). The association of dynamin with synaptophysin regulates quantal size and duration of exocytotic events in chromaffin cells. *J Neurosci*, *30*, 10683-10691.
- Gourdain, S., Dairou, J., Denhez, C., Bui, L. C., Rodrigues-Lima, F., Janel, N., Delabar, J. M., Cariou, K., & Dodd, R. H. (2013). Development of DANDYs, new 3,5-diaryl-7-azaindoles demonstrating potent DYRK1A kinase inhibitory activity. *J Med Chem*, *56*, 9569-9585.
- Grau, C., Arato, K., Fernandez-Fernandez, J. M., Valderrama, A., Sindreu, C., Fillat, C., Ferrer, I., de la Luna, S., & Altafaj, X. (2014). DYRK1A-mediated phosphorylation of GluN2A at Ser(1048) regulates the surface expression and channel activity of GluN1/GluN2A receptors. *Front Cell Neurosci*, *8*, 331.
- Grundke-Iqbal, I., Iqbal, K., Tung, Y. C., Quinlan, M., Wisniewski, H. M., & Binder, L. I. (1986). Abnormal phosphorylation of the microtubule-associated protein tau (tau) in Alzheimer cytoskeletal pathology. *Proc Natl Acad Sci U S A*, *83*, 4913-4917.
- Guedj, F., Pereira, P. L., Najas, S., Barallobre, M. J., Chabert, C., Souchet, B., Sebric, C., Verney, C., Herault, Y., Arbones, M., & Delabar, J. M. (2012). DYRK1A: a master regulatory protein controlling brain growth. *Neurobiol Dis*, *46*, 190-203.
- Guedj, F., Sebric, C., Rivals, I., Ledru, A., Paly, E., Bizot, J. C., Smith, D., Rubin, E., Gillet, B., Arbones, M., & Delabar, J. M. (2009). Green tea polyphenols rescue of brain defects induced by overexpression of DYRK1A. *PLoS One*, *4*, e4606.
- Guiley, K. Z., Liban, T. J., Felthousen, J. G., Ramanan, P., Litovchick, L., & Rubin, S. M. (2015). Structural mechanisms of DREAM complex assembly and regulation. *Genes Dev*, *29*, 961-974.
- Guimera, J., Casas, C., Pucharcos, C., Solans, A., Domenech, A., Planas, A. M., Ashley, J., Lovett, M., Estivill, X., & Pritchard, M. A. (1996). A human homologue of Drosophila minibrain (MNB) is expressed in the neuronal regions affected in Down syndrome and maps to the critical region. *Hum Mol Genet*, *5*, 1305-1310.
- Guo, X., Williams, J. G., Schug, T. T., & Li, X. (2010). DYRK1A and DYRK3 promote cell survival through phosphorylation and activation of SIRT1. *J Biol Chem*, *285*, 13223-13232.
- Haas, M. A., Bell, D., Slender, A., Lana-Elola, E., Watson-Scales, S., Fisher, E. M., Tybulewicz, V. L., & Guillemot, F. (2013). Alterations to dendritic spine morphology, but not dendrite patterning, of cortical projection neurons in Tc1 and Ts1Rhr mouse models of Down syndrome. *PLoS One*, *8*, e78561.
- Hammerle, B., Carnicero, A., Elizalde, C., Ceron, J., Martinez, S., & Tejedor, F. J. (2003). Expression patterns and subcellular localization of the Down syndrome candidate protein MNB/DYRK1A suggest a role in late neuronal differentiation. *Eur J Neurosci*, *17*, 2277-2286.

- Hammerle, B., Elizalde, C., Galceran, J., Becker, W., & Tejedor, F. J. (2003). The MNB/DYRK1A protein kinase: neurobiological functions and Down syndrome implications. *J Neural Transm Suppl*, 129-137.
- Hammerle, B., Elizalde, C., & Tejedor, F. J. (2008). The spatio-temporal and subcellular expression of the candidate Down syndrome gene Mnb/Dyrk1A in the developing mouse brain suggests distinct sequential roles in neuronal development. *Eur J Neurosci*, 27, 1061-1074.
- Hammerle, B., Ulin, E., Guimera, J., Becker, W., Guillemot, F., & Tejedor, F. J. (2011). Transient expression of Mnb/Dyrk1a couples cell cycle exit and differentiation of neuronal precursors by inducing p27KIP1 expression and suppressing NOTCH signaling. *Development*, 138, 2543-2554.
- Hammerle, B., Vera-Samper, E., Speicher, S., Arencibia, R., Martinez, S., & Tejedor, F. J. (2002). Mnb/Dyrk1A is transiently expressed and asymmetrically segregated in neural progenitor cells at the transition to neurogenic divisions. *Dev Biol*, 246, 259-273.
- Hanger, D. P., Brion, J. P., Gallo, J. M., Cairns, N. J., Luthert, P. J., & Anderton, B. H. (1991). Tau in Alzheimer's disease and Down's syndrome is insoluble and abnormally phosphorylated. *Biochem J*, 275 (Pt 1), 99-104.
- Hattori, M., Fujiyama, A., Taylor, T. D., Watanabe, H., Yada, T., Park, H. S., Toyoda, A., Ishii, K., Totoki, Y., Choi, D. K., Groner, Y., Soeda, E., Ohki, M., Takagi, T., Sakaki, Y., Taudien, S., Blechschmidt, K., Polley, A., Menzel, U., Delabar, J., Kumpf, K., Lehmann, R., Patterson, D., Reichwald, K., Rump, A., Schillhabel, M., Schudy, A., Zimmermann, W., Rosenthal, A., Kudoh, J., Schibuya, K., Kawasaki, K., Asakawa, S., Shintani, A., Sasaki, T., Nagamine, K., Mitsuyama, S., Antonarakis, S. E., Minoshima, S., Shimizu, N., Nordsiek, G., Hornischer, K., Brant, P., Scharfe, M., Schon, O., Desario, A., Reichelt, J., Kauer, G., Blocker, H., Ramser, J., Beck, A., Klages, S., Hennig, S., Riesselmann, L., Dagand, E., Haaf, T., Wehrmeyer, S., Borzym, K., Gardiner, K., Nizetic, D., Francis, F., Lehrach, H., Reinhardt, R., Yaspo, M. L., Chromosome, m., & sequencing, c. (2000). The DNA sequence of human chromosome 21. *Nature*, 405, 311-319.
- Himpel, S., Panzer, P., Eirnbter, K., Czajkowska, H., Sayed, M., Packman, L. C., Blundell, T., Kentrup, H., Grotzinger, J., Joost, H. G., & Becker, W. (2001). Identification of the autophosphorylation sites and characterization of their effects in the protein kinase DYRK1A. *Biochem J*, 359, 497-505.
- Himpel, S., Tegge, W., Frank, R., Leder, S., Joost, H. G., & Becker, W. (2000). Specificity determinants of substrate recognition by the protein kinase DYRK1A. *J Biol Chem*, 275, 2431-2438.
- Hotulainen, P., & Saarikangas, J. (2016). The initiation of post-synaptic protrusions. *Commun Integr Biol*, 9, e1125053.
- Im, E., & Chung, K. C. (2015). Dyrk1A phosphorylates parkin at Ser-131 and negatively regulates its ubiquitin E3 ligase activity. *J Neurochem*, 134, 756-768.
- Impey, S., Obrietan, K., & Storm, D. R. (1999). Making new connections: role of ERK/MAP kinase signaling in neuronal plasticity. *Neuron*, 23, 11-14.
- Iossifov, I., O'Roak, B. J., Sanders, S. J., Ronemus, M., Krumm, N., Levy, D., Stessman, H. A., Witherspoon, K. T., Vives, L., Patterson, K. E., Smith, J. D., Paepers, B., Nickerson, D. A., Dea, J., Dong, S., Gonzalez, L. E., Mandell, J. D., Mane, S. M., Murtha, M. T., Sullivan, C. A., Walker, M. F., Waqar, Z., Wei, L., Willsey, A. J., Yamrom, B., Lee, Y. H., Grabowska, E., Dalkic, E., Wang, Z., Marks, S., Andrews, P., Leotta, A., Kendall, J., Hakker, I., Rosenbaum, J., Ma, B., Rodgers, L., Troge, J., Narzisi, G., Yoon, S., Schatz, M. C., Ye, K., McCombie, W. R., Shendure, J., Eichler, E. E., State, M. W., & Wigler, M. (2014). The contribution of de novo coding mutations to autism spectrum disorder. *Nature*, 515, 216-221.
- Irwin, D. J., Lee, V. M., & Trojanowski, J. Q. (2013). Parkinson's disease dementia: convergence of alpha-synuclein, tau and amyloid-beta pathologies. *Nat Rev Neurosci*, 14, 626-636.
- Ishihara, K., Amano, K., Takaki, E., Shimohata, A., Sago, H., Epstein, C. J., & Yamakawa, K. (2010). Enlarged brain ventricles and impaired neurogenesis in the Ts1Cje and Ts2Cje mouse models of Down syndrome. *Cereb Cortex*, 20, 1131-1143.

- Janel, N., Alexopoulos, P., Badel, A., Lamari, F., Camproux, A. C., Lagarde, J., Simon, S., Feraudet-Tarisse, C., Lamourette, P., Arbones, M., Paul, J. L., Dubois, B., Potier, M. C., Sarazin, M., & Delabar, J. M. (2017). Combined assessment of DYRK1A, BDNF and homocysteine levels as diagnostic marker for Alzheimer's disease. *Transl Psychiatry*, *7*, e1154.
- Janel, N., Sarazin, M., Corlier, F., Corne, H., de Souza, L. C., Hamelin, L., Aka, A., Lagarde, J., Blehaut, H., Hindie, V., Rain, J. C., Arbones, M. L., Dubois, B., Potier, M. C., Bottlaender, M., & Delabar, J. M. (2014). Plasma DYRK1A as a novel risk factor for Alzheimer's disease. *Transl Psychiatry*, *4*, e425.
- Jang, S. M., Azebi, S., Soubigou, G., & Muchardt, C. (2014). DYRK1A phosphorylates histone H3 to differentially regulate the binding of HP1 isoforms and antagonize HP1-mediated transcriptional repression. *EMBO Rep*, *15*, 686-694.
- Ji, J., Lee, H., Argiropoulos, B., Dorrani, N., Mann, J., Martinez-Agosto, J. A., Gomez-Ospina, N., Gallant, N., Bernstein, J. A., Hudgins, L., Slattery, L., Isidor, B., Le Caignec, C., David, A., Oberszty, E., Wisniewiecka-Kowalnik, B., Fox, M., Deignan, J. L., Vilain, E., Hendricks, E., Horton Harr, M., Noon, S. E., Jackson, J. R., Wilkens, A., Mirzaa, G., Salamon, N., Abramson, J., Zackai, E. H., Krantz, I., Innes, A. M., Nelson, S. F., Grody, W. W., & Quintero-Rivera, F. (2015). DYRK1A haploinsufficiency causes a new recognizable syndrome with microcephaly, intellectual disability, speech impairment, and distinct facies. *Eur J Hum Genet*, *23*, 1473-1481.
- Jimenez, G., Shvartsman, S. Y., & Paroush, Z. (2012). The Capicua repressor--a general sensor of RTK signaling in development and disease. *J Cell Sci*, *125*, 1383-1391.
- Jin, N., Yin, X., Gu, J., Zhang, X., Shi, J., Qian, W., Ji, Y., Cao, M., Gu, X., Ding, F., Iqbal, K., Gong, C. X., & Liu, F. (2015). Truncation and Activation of Dual Specificity Tyrosine Phosphorylation-regulated Kinase 1A by Calpain I: A MOLECULAR MECHANISM LINKED TO TAU PATHOLOGY IN ALZHEIMER DISEASE. *J Biol Chem*, *290*, 15219-15237.
- Jones, E. L., Aarsland, D., Londos, E., & Ballard, C. (2012). A pilot study examining associations between DYRK1A and alpha-synuclein dementias. *Neurodegener Dis*, *10*, 229-231.
- Jung, K. H., Chu, K., Lee, S. T., Park, K. I., Kim, J. H., Kang, K. M., Kim, S., Jeon, D., Kim, M., Lee, S. K., & Roh, J. K. (2011). Molecular alterations underlying epileptogenesis after prolonged febrile seizure and modulation by erythropoietin. *Epilepsia*, *52*, 541-550.
- Kaczmarek, W., Barua, M., Mazur-Kolecka, B., Frackowiak, J., Dowjat, W., Mehta, P., Bolton, D., Hwang, Y. W., Rabe, A., Albertini, G., & Wegiel, J. (2014). Intracellular distribution of differentially phosphorylated dual-specificity tyrosine phosphorylation-regulated kinase 1A (DYRK1A). *J Neurosci Res*, *92*, 162-173.
- Kang, J. E., Choi, S. A., Park, J. B., & Chung, K. C. (2005). Regulation of the proapoptotic activity of huntingtin interacting protein 1 by Dyrk1 and caspase-3 in hippocampal neuroprogenitor cells. *J Neurosci Res*, *81*, 62-72.
- Kawakubo, T., Mori, R., Shirogami, K., Iwata, N., & Asai, M. (2017). Nephilysin Is Suppressed by Dual-Specificity Tyrosine-Phosphorylation Regulated Kinase 1A (DYRK1A) in Down-Syndrome-Derived Fibroblasts. *Biol Pharm Bull*, *40*, 327-333.
- Kelly, P. A., & Rahmani, Z. (2005). DYRK1A enhances the mitogen-activated protein kinase cascade in PC12 cells by forming a complex with Ras, B-Raf, and MEK1. *Mol Biol Cell*, *16*, 3562-3573.
- Kentrup, H., Joost, H. G., Heimann, G., & Becker, W. (2000). [Minibrain/DYRK1A gene: candidate gene for mental retardation in Down's syndrome?]. *Klin Padiatr*, *212*, 60-63.
- Khan, N., Afaq, F., Saleem, M., Ahmad, N., & Mukhtar, H. (2006). Targeting multiple signaling pathways by green tea polyphenol (-)-epigallocatechin-3-gallate. *Cancer Res*, *66*, 2500-2505.
- Kii, I., Sumida, Y., Goto, T., Sonamoto, R., Okuno, Y., Yoshida, S., Kato-Sumida, T., Koike, Y., Abe, M., Nonaka, Y., Ikura, T., Ito, N., Shibuya, H., Hosoya, T., & Hagiwara, M. (2016). Selective inhibition of the kinase DYRK1A by targeting its folding process. *Nat Commun*, *7*, 11391.
- Kim, E. J., Sung, J. Y., Lee, H. J., Rhim, H., Hasegawa, M., Iwatsubo, T., Min do, S., Kim, J., Paik, S. R., & Chung, K. C. (2006). Dyrk1A phosphorylates alpha-synuclein and enhances intracellular inclusion formation. *J Biol Chem*, *281*, 33250-33257.

- Kim, O. H., Cho, H. J., Han, E., Hong, T. I., Ariyasiri, K., Choi, J. H., Hwang, K. S., Jeong, Y. M., Yang, S. Y., Yu, K., Park, D. S., Oh, H. W., Davis, E. E., Schwartz, C. E., Lee, J. S., Kim, H. G., & Kim, C. H. (2017). Zebrafish knockout of Down syndrome gene, DYRK1A, shows social impairments relevant to autism. *Mol Autism*, *8*, 50.
- Kim, Y., Park, J., Song, W. J., & Chang, S. (2010). Overexpression of Dyrk1A causes the defects in synaptic vesicle endocytosis. *Neurosignals*, *18*, 164-172.
- Kim, Y. M., Lee, Y. J., Park, J. H., Lee, H. D., Cheon, C. K., Kim, S. Y., Hwang, J. Y., Jang, J. H., & Yoo, H. W. (2017). High diagnostic yield of clinically unidentifiable syndromic growth disorders by targeted exome sequencing. *Clin Genet*, *92*, 594-605.
- Kimura, R., Kamino, K., Yamamoto, M., Nuripa, A., Kida, T., Kazui, H., Hashimoto, R., Tanaka, T., Kudo, T., Yamagata, H., Tabara, Y., Miki, T., Akatsu, H., Kosaka, K., Funakoshi, E., Nishitomi, K., Sakaguchi, G., Kato, A., Hattori, H., Uema, T., & Takeda, M. (2007). The DYRK1A gene, encoded in chromosome 21 Down syndrome critical region, bridges between beta-amyloid production and tau phosphorylation in Alzheimer disease. *Hum Mol Genet*, *16*, 15-23.
- Kleschevnikov, A. M., Belichenko, P. V., Villar, A. J., Epstein, C. J., Malenka, R. C., & Mobley, W. C. (2004). Hippocampal long-term potentiation suppressed by increased inhibition in the Ts65Dn mouse, a genetic model of Down syndrome. *J Neurosci*, *24*, 8153-8160.
- Korbel, J. O., Tirosh-Wagner, T., Urban, A. E., Chen, X. N., Kasowski, M., Dai, L., Grubert, F., Erdman, C., Gao, M. C., Lange, K., Sobel, E. M., Barlow, G. M., Aylsworth, A. S., Carpenter, N. J., Clark, R. D., Cohen, M. Y., Doran, E., Falik-Zaccari, T., Lewin, S. O., Lott, I. T., McGillivray, B. C., Moeschler, J. B., Pettenati, M. J., Puschel, S. M., Rao, K. W., Shaffer, L. G., Shohat, M., Van Riper, A. J., Warburton, D., Weissman, S., Gerstein, M. B., Snyder, M., & Korenberg, J. R. (2009). The genetic architecture of Down syndrome phenotypes revealed by high-resolution analysis of human segmental trisomies. *Proc Natl Acad Sci U S A*, *106*, 12031-12036.
- Kosmicki, J. A., Samocha, K. E., Howrigan, D. P., Sanders, S. J., Slowikowski, K., Lek, M., Karczewski, K. J., Cutler, D. J., Devlin, B., Roeder, K., Buxbaum, J. D., Neale, B. M., MacArthur, D. G., Wall, D. P., Robinson, E. B., & Daly, M. J. (2017). Refining the role of de novo protein-truncating variants in neurodevelopmental disorders by using population reference samples. *Nat Genet*, *49*, 504-510.
- Kurabayashi, N., Hirota, T., Sakai, M., Sanada, K., & Fukada, Y. (2010). DYRK1A and glycogen synthase kinase 3beta, a dual-kinase mechanism directing proteasomal degradation of CRY2 for circadian timekeeping. *Mol Cell Biol*, *30*, 1757-1768.
- Kurabayashi, N., Nguyen, M. D., & Sanada, K. (2015). DYRK1A overexpression enhances STAT activity and astroglialogenesis in a Down syndrome mouse model. *EMBO Rep*, *16*, 1548-1562.
- Kurabayashi, N., & Sanada, K. (2013). Increased dosage of DYRK1A and DSCR1 delays neuronal differentiation in neocortical progenitor cells. *Genes Dev*, *27*, 2708-2721.
- Kwon, H. B., Kozorovitskiy, Y., Oh, W. J., Peixoto, R. T., Akhtar, N., Saulnier, J. L., Gu, C., & Sabatini, B. L. (2012). Neuroligin-1-dependent competition regulates cortical synaptogenesis and synapse number. *Nat Neurosci*, *15*, 1667-1674.
- Lacovich, V., Espindola, S. L., Alloatti, M., Pozo Devoto, V., Cromberg, L. E., Carna, M. E., Forte, G., Gallo, J. M., Bruno, L., Stokin, G. B., Avale, M. E., & Falzone, T. L. (2017). Tau Isoforms Imbalance Impairs the Axonal Transport of the Amyloid Precursor Protein in Human Neurons. *J Neurosci*, *37*, 58-69.
- Laguna, A., Aranda, S., Barallobre, M. J., Barhoum, R., Fernandez, E., Fotaki, V., Delabar, J. M., de la Luna, S., de la Villa, P., & Arbones, M. L. (2008). The protein kinase DYRK1A regulates caspase-9-mediated apoptosis during retina development. *Dev Cell*, *15*, 841-853.
- Laguna, A., Barallobre, M. J., Marchena, M. A., Mateus, C., Ramirez, E., Martinez-Cue, C., Delabar, J. M., Castelo-Branco, M., de la Villa, P., & Arbones, M. L. (2013). Triplication of DYRK1A causes retinal structural and functional alterations in Down syndrome. *Hum Mol Genet*, *22*, 2775-2784.
- Lautenschlager, J., Kaminski, C. F., & Kaminski Schierle, G. S. (2017). alpha-Synuclein - Regulator of Exocytosis, Endocytosis, or Both? *Trends Cell Biol*, *27*, 468-479.

- Lee, Y., Ha, J., Kim, H. J., Kim, Y. S., Chang, E. J., Song, W. J., & Kim, H. H. (2009). Negative feedback Inhibition of NFATc1 by DYRK1A regulates bone homeostasis. *J Biol Chem*, *284*, 33343-33351.
- Lepagnol-Bestel, A. M., Zvara, A., Maussion, G., Quignon, F., Ngimbous, B., Ramoz, N., Imbeaud, S., Loe-Mie, Y., Benihoud, K., Agier, N., Salin, P. A., Cardona, A., Khung-Savatovsky, S., Kallunki, P., Delabar, J. M., Puskas, L. G., Delacroix, H., Aggerbeck, L., Delezoide, A. L., Delattre, O., Gorwood, P., Moalic, J. M., & Simonneau, M. (2009). DYRK1A interacts with the REST/NRSF-SWI/SNF chromatin remodelling complex to deregulate gene clusters involved in the neuronal phenotypic traits of Down syndrome. *Hum Mol Genet*, *18*, 1405-1414.
- Li, D., Jackson, R. A., Yusoff, P., & Guy, G. R. (2010). Direct association of Sprouty-related protein with an EVH1 domain (SPRED) 1 or SPRED2 with DYRK1A modifies substrate/kinase interactions. *J Biol Chem*, *285*, 35374-35385.
- Litovchick, L., Florens, L. A., Swanson, S. K., Washburn, M. P., & DeCaprio, J. A. (2011). DYRK1A protein kinase promotes quiescence and senescence through DREAM complex assembly. *Genes Dev*, *25*, 801-813.
- Liu, F., Liang, Z., Wegiel, J., Hwang, Y. W., Iqbal, K., Grundke-Iqbal, I., Ramakrishna, N., & Gong, C. X. (2008). Overexpression of Dyrk1A contributes to neurofibrillary degeneration in Down syndrome. *FASEB J*, *22*, 3224-3233.
- Liu, T., Sims, D., & Baum, B. (2009). Parallel RNAi screens across different cell lines identify generic and cell type-specific regulators of actin organization and cell morphology. *Genome Biol*, *10*, R26.
- Liu, W., Zhou, H., Liu, L., Zhao, C., Deng, Y., Chen, L., Wu, L., Mandrycky, N., McNabb, C. T., Peng, Y., Fuchs, P. N., Lu, J., Sheen, V., Qiu, M., Mao, M., & Lu, Q. R. (2015). Disruption of neurogenesis and cortical development in transgenic mice misexpressing Olig2, a gene in the Down syndrome critical region. *Neurobiol Dis*, *77*, 106-116.
- Lochhead, P. A., Sibbet, G., Morrice, N., & Cleghon, V. (2005). Activation-loop autophosphorylation is mediated by a novel transitional intermediate form of DYRKs. *Cell*, *121*, 925-936.
- London, J., Rouch, C., Bui, L. C., Assayag, E., Souchet, B., Daubigney, F., Medjaoui, H., Luquet, S., Magnan, C., Delabar, J. M., Dairou, J., & Janel, N. (2017). Overexpression of the DYRK1A Gene (Dual-Specificity Tyrosine Phosphorylation-Regulated Kinase 1A) Induces Alterations of the Serotonergic and Dopaminergic Processing in Murine Brain Tissues. *Mol Neurobiol*.
- Lu, M., Zheng, L., Han, B., Wang, L., Wang, P., Liu, H., & Sun, X. (2011). REST regulates DYRK1A transcription in a negative feedback loop. *J Biol Chem*, *286*, 10755-10763.
- Luco, S. M., Pohl, D., Sell, E., Wagner, J. D., Dymont, D. A., & Daoud, H. (2016). Case report of novel DYRK1A mutations in 2 individuals with syndromic intellectual disability and a review of the literature. *BMC Med Genet*, *17*, 15.
- Lyle, R., Bena, F., Gagos, S., Gehrig, C., Lopez, G., Schinzel, A., Lespinasse, J., Bottani, A., Dahoun, S., Taine, L., Doco-Fenzy, M., Cornillet-Lefebvre, P., Pelet, A., Lyonnet, S., Toutain, A., Colleaux, L., Horst, J., Kennerknecht, I., Wakamatsu, N., Descartes, M., Franklin, J. C., Florentin-Arar, L., Kitsiou, S., Ait Yahya-Graison, E., Costantine, M., Sinet, P. M., Delabar, J. M., & Antonarakis, S. E. (2009). Genotype-phenotype correlations in Down syndrome identified by array CGH in 30 cases of partial trisomy and partial monosomy chromosome 21. *Eur J Hum Genet*, *17*, 454-466.
- Maenz, B., Hekerman, P., Vela, E. M., Galceran, J., & Becker, W. (2008). Characterization of the human DYRK1A promoter and its regulation by the transcription factor E2F1. *BMC Mol Biol*, *9*, 30.
- Mandel, S. A., Amit, T., Weinreb, O., Reznichenko, L., & Youdim, M. B. (2008). Simultaneous manipulation of multiple brain targets by green tea catechins: a potential neuroprotective strategy for Alzheimer and Parkinson diseases. *CNS Neurosci Ther*, *14*, 352-365.
- Mao, J., Maye, P., Kogerman, P., Tejedor, F. J., Toftgard, R., Xie, W., Wu, G., & Wu, D. (2002). Regulation of Gli1 transcriptional activity in the nucleus by Dyrk1. *J Biol Chem*, *277*, 35156-35161.

- Marin-Padilla, M. (1976). Pyramidal cell abnormalities in the motor cortex of a child with Down's syndrome. A Golgi study. *J Comp Neurol*, *167*, 63-81.
- Marti, E., Altafaj, X., Dierssen, M., de la Luna, S., Fotaki, V., Alvarez, M., Perez-Riba, M., Ferrer, I., & Estivill, X. (2003). Dyrk1A expression pattern supports specific roles of this kinase in the adult central nervous system. *Brain Res*, *964*, 250-263.
- Martinez-Cue, C., Martinez, P., Rueda, N., Vidal, R., Garcia, S., Vidal, V., Corrales, A., Montero, J. A., Pazos, A., Florez, J., Gasser, R., Thomas, A. W., Honer, M., Knoflach, F., Trejo, J. L., Wettstein, J. G., & Hernandez, M. C. (2013). Reducing GABAA alpha5 receptor-mediated inhibition rescues functional and neuromorphological deficits in a mouse model of down syndrome. *J Neurosci*, *33*, 3953-3966.
- Martinez de Lagran, M., Benavides-Piccione, R., Ballesteros-Yanez, I., Calvo, M., Morales, M., Fillat, C., Defelipe, J., Ramakers, G. J., & Dierssen, M. (2012). Dyrk1A influences neuronal morphogenesis through regulation of cytoskeletal dynamics in mammalian cortical neurons. *Cereb Cortex*, *22*, 2867-2877.
- Masaki, S., Kii, I., Sumida, Y., Kato-Sumida, T., Ogawa, Y., Ito, N., Nakamura, M., Sonamoto, R., Kataoka, N., Hosoya, T., & Hagiwara, M. (2015). Design and synthesis of a potent inhibitor of class 1 DYRK kinases as a suppressor of adipogenesis. *Bioorg Med Chem*, *23*, 4434-4441.
- Matsumoto, N., Ohashi, H., Tsukahara, M., Kim, K. C., Soeda, E., & Niikawa, N. (1997). Possible narrowed assignment of the loci of monosomy 21-associated microcephaly and intrauterine growth retardation to a 1.2-Mb segment at 21q22.2. *Am J Hum Genet*, *60*, 997-999.
- McElyea, S. D., Starbuck, J. M., Tumbleson-Brink, D. M., Harrington, E., Blazek, J. D., Ghoneima, A., Kula, K., & Roper, R. J. (2016). Influence of prenatal EGCG treatment and Dyrk1a dosage reduction on craniofacial features associated with Down syndrome. *Hum Mol Genet*, *25*, 4856-4869.
- Moller, R. S., Kubart, S., Hoeltzenbein, M., Heye, B., Vogel, I., Hansen, C. P., Menzel, C., Ullmann, R., Tommerup, N., Ropers, H. H., Tumer, Z., & Kalscheuer, V. M. (2008). Truncation of the Down syndrome candidate gene DYRK1A in two unrelated patients with microcephaly. *Am J Hum Genet*, *82*, 1165-1170.
- Mozzi, A., Forni, D., Cagliani, R., Pozzoli, U., Clerici, M., & Sironi, M. (2017). Distinct selective forces and Neanderthal introgression shaped genetic diversity at genes involved in neurodevelopmental disorders. *Sci Rep*, *7*, 6116.
- Murakami, N., Bolton, D., & Hwang, Y. W. (2009). Dyrk1A binds to multiple endocytic proteins required for formation of clathrin-coated vesicles. *Biochemistry*, *48*, 9297-9305.
- Murakami, N., Bolton, D. C., Kida, E., Xie, W., & Hwang, Y. W. (2012). Phosphorylation by Dyrk1A of clathrin coated vesicle-associated proteins: identification of the substrate proteins and the effects of phosphorylation. *PLoS One*, *7*, e34845.
- Murakami, N., Xie, W., Lu, R. C., Chen-Hwang, M. C., Wieraszko, A., & Hwang, Y. W. (2006). Phosphorylation of amphiphysin I by minibrain kinase/dual-specificity tyrosine phosphorylation-regulated kinase, a kinase implicated in Down syndrome. *J Biol Chem*, *281*, 23712-23724.
- Naert, G., Ferre, V., Meunier, J., Keller, E., Malmstrom, S., Givalois, L., Carreaux, F., Bazureau, J. P., & Maurice, T. (2015). Leucettine L41, a DYRK1A-preferential DYRKs/CLKs inhibitor, prevents memory impairments and neurotoxicity induced by oligomeric Abeta25-35 peptide administration in mice. *Eur Neuropsychopharmacol*, *25*, 2170-2182.
- Najas, S., Arranz, J., Lochhead, P. A., Ashford, A. L., Oxley, D., Delabar, J. M., Cook, S. J., Barallobre, M. J., & Arbones, M. L. (2015). DYRK1A-mediated Cyclin D1 Degradation in Neural Stem Cells Contributes to the Neurogenic Cortical Defects in Down Syndrome. *EBioMedicine*, *2*, 120-134.
- Nakano-Kobayashi, A., Awaya, T., Kii, I., Sumida, Y., Okuno, Y., Yoshida, S., Sumida, T., Inoue, H., Hosoya, T., & Hagiwara, M. (2017). Prenatal neurogenesis induction therapy normalizes brain structure and function in Down syndrome mice. *Proc Natl Acad Sci U S A*, *114*, 10268-10273.
- Neumann, F., Gourdain, S., Albac, C., Dekker, A. D., Bui, L. C., Dairou, J., Schmitz-Afonso, I., Hue, N., Rodrigues-Lima, F., Delabar, J. M., Potier, M. C., Le Caer, J. P., Touboul, D., Delatour, B.,

- Cariou, K., & Dodd, R. H. (2018). DYRK1A inhibition and cognitive rescue in a Down syndrome mouse model are induced by new fluoro-DANDY derivatives. *Sci Rep*, *8*, 2859.
- Newpher, T. M., Harris, S., Pringle, J., Hamilton, C., & Soderling, S. (2017). Regulation of spine structural plasticity by Arc/Arg3.1. *Semin Cell Dev Biol*.
- O'Roak, B. J., Vives, L., Fu, W., Egertson, J. D., Stanaway, I. B., Phelps, I. G., Carvill, G., Kumar, A., Lee, C., Ankenman, K., Munson, J., Hiatt, J. B., Turner, E. H., Levy, R., O'Day, D. R., Krumm, N., Coe, B. P., Martin, B. K., Borenstein, E., Nickerson, D. A., Mefford, H. C., Doherty, D., Akey, J. M., Bernier, R., Eichler, E. E., & Shendure, J. (2012). Multiplex targeted sequencing identifies recurrently mutated genes in autism spectrum disorders. *Science*, *338*, 1619-1622.
- Oegema, R., de Klein, A., Verkerk, A. J., Schot, R., Dumeé, B., Douben, H., Eussen, B., Dubbel, L., Poddighe, P. J., van der Laar, I., Dobyns, W. B., van der Spek, P. J., Lequin, M. H., de Coo, I. F., de Wit, M. C., Wessels, M. W., & Mancini, G. M. (2010). Distinctive Phenotypic Abnormalities Associated with Submicroscopic 21q22 Deletion Including DYRK1A. *Mol Syndromol*, *1*, 113-120.
- Ogawa, Y., Nonaka, Y., Goto, T., Ohnishi, E., Hiramatsu, T., Kii, I., Yoshida, M., Ikura, T., Onogi, H., Shibuya, H., Hosoya, T., Ito, N., & Hagiwara, M. (2010). Development of a novel selective inhibitor of the Down syndrome-related kinase Dyrk1A. *Nat Commun*, *1*, 86.
- Okui, M., Ide, T., Morita, K., Funakoshi, E., Ito, F., Ogita, K., Yoneda, Y., Kudoh, J., & Shimizu, N. (1999). High-level expression of the Mnb/Dyrk1A gene in brain and heart during rat early development. *Genomics*, *62*, 165-171.
- Ori-McKenney, K. M., McKenney, R. J., Huang, H. H., Li, T., Meltzer, S., Jan, L. Y., Vale, R. D., Wiita, A. P., & Jan, Y. N. (2016). Phosphorylation of beta-Tubulin by the Down Syndrome Kinase, Minibrain/DYRK1a, Regulates Microtubule Dynamics and Dendrite Morphogenesis. *Neuron*, *90*, 551-563.
- Osoegawa, K., Susukida, R., Okano, S., Kudoh, J., Minoshima, S., Shimizu, N., de Jong, P. J., Groet, J., Ives, J., Lehrach, H., Nizetic, D., & Soeda, E. (1996). An integrated map with cosmid/PAC contigs of a 4-Mb Down syndrome critical region. *Genomics*, *32*, 375-387.
- Palop, J. J., Chin, J., Roberson, E. D., Wang, J., Thwin, M. T., Bien-Ly, N., Yoo, J., Ho, K. O., Yu, G. Q., Kreitzer, A., Finkbeiner, S., Noebels, J. L., & Mucke, L. (2007). Aberrant excitatory neuronal activity and compensatory remodeling of inhibitory hippocampal circuits in mouse models of Alzheimer's disease. *Neuron*, *55*, 697-711.
- Papoulidis, I., Papageorgiou, E., Siomou, E., Oikonomidou, E., Thomaidis, L., Vetro, A., Zuffardi, O., Liehr, T., Manolakos, E., & Vassilis, P. (2014). A patient with partial trisomy 21 and 7q deletion expresses mild Down syndrome phenotype. *Gene*, *536*, 441-443.
- Park, J., Oh, Y., Yoo, L., Jung, M. S., Song, W. J., Lee, S. H., Seo, H., & Chung, K. C. (2010). Dyrk1A phosphorylates p53 and inhibits proliferation of embryonic neuronal cells. *J Biol Chem*, *285*, 31895-31906.
- Park, J., Sung, J. Y., Park, J., Song, W. J., Chang, S., & Chung, K. C. (2012). Dyrk1A negatively regulates the actin cytoskeleton through threonine phosphorylation of N-WASP. *J Cell Sci*, *125*, 67-80.
- Park, J. H., Jung, M. S., Kim, Y. S., Song, W. J., & Chung, S. H. (2012). Phosphorylation of Munc18-1 by Dyrk1A regulates its interaction with Syntaxin 1 and X11alpha. *J Neurochem*, *122*, 1081-1091.
- Patil, N., Cox, D. R., Bhat, D., Faham, M., Myers, R. M., & Peterson, A. S. (1995). A potassium channel mutation in weaver mice implicates membrane excitability in granule cell differentiation. *Nat Genet*, *11*, 126-129.
- Pearlson, G. D., Breiter, S. N., Aylward, E. H., Warren, A. C., Grygorcewicz, M., Frangou, S., Barta, P. E., & Pulsifer, M. B. (1998). MRI brain changes in subjects with Down syndrome with and without dementia. *Dev Med Child Neurol*, *40*, 326-334.
- Pehar, M., Ko, M. H., Li, M., Scrabble, H., & Puglielli, L. (2014). P44, the 'longevity-assurance' isoform of P53, regulates tau phosphorylation and is activated in an age-dependent fashion. *Aging Cell*, *13*, 449-456.
- Pennington, B. F., Moon, J., Edgin, J., Stedron, J., & Nadel, L. (2003). The neuropsychology of Down syndrome: evidence for hippocampal dysfunction. *Child Dev*, *74*, 75-93.

- Perez-Cremades, D., Hernandez, S., Blasco-Ibanez, J. M., Crespo, C., Nacher, J., & Varea, E. (2010). Alteration of inhibitory circuits in the somatosensory cortex of Ts65Dn mice, a model for Down's syndrome. *J Neural Transm (Vienna)*, *117*, 445-455.
- Piper, M., & Holt, C. (2004). RNA translation in axons. *Annu Rev Cell Dev Biol*, *20*, 505-523.
- Qian, W., Liang, H., Shi, J., Jin, N., Grundke-Iqbal, I., Iqbal, K., Gong, C. X., & Liu, F. (2011). Regulation of the alternative splicing of tau exon 10 by SC35 and Dyrk1A. *Nucleic Acids Res*, *39*, 6161-6171.
- Rahmani, Z., Blouin, J. L., Creau-Goldberg, N., Watkins, P. C., Mattei, J. F., Poissonnier, M., Prieur, M., Chettouh, Z., Nicole, A., Aurias, A., & et al. (1989). Critical role of the D21S55 region on chromosome 21 in the pathogenesis of Down syndrome. *Proc Natl Acad Sci U S A*, *86*, 5958-5962.
- Rahmani, Z., Lopes, C., Rachidi, M., & Delabar, J. M. (1998). Expression of the mnb (dyrk) protein in adult and embryonic mouse tissues. *Biochem Biophys Res Commun*, *253*, 514-518.
- Raich, W. B., Moorman, C., Lacefield, C. O., Lehrer, J., Bartsch, D., Plasterk, R. H., Kandel, E. R., & Hobert, O. (2003). Characterization of *Caenorhabditis elegans* homologs of the Down syndrome candidate gene DYRK1A. *Genetics*, *163*, 571-580.
- Ramkumar, A., Jong, B. Y., & Ori-McKenney, K. M. (2017). ReMAPping the microtubule landscape: How phosphorylation dictates the activities of microtubule-associated proteins. *Dev Dyn*.
- Ramkumar, A., Jong, B. Y., & Ori-McKenney, K. M. (2018). ReMAPping the microtubule landscape: How phosphorylation dictates the activities of microtubule-associated proteins. *Dev Dyn*, *247*, 138-155.
- Raveau, M., Polygalov, D., Boehringer, R., Amano, K., Yamakawa, K., & McHugh, T. J. (2018). Alterations of in vivo CA1 network activity in Dp(16)1Yey Down syndrome model mice. *Elife*, *7*.
- Redin, C., Gerard, B., Lauer, J., Herenger, Y., Muller, J., Quartier, A., Masurel-Paulet, A., Willems, M., Lesca, G., El-Chehadeh, S., Le Gras, S., Vicaire, S., Philipps, M., Dumas, M., Geoffroy, V., Feger, C., Haumesser, N., Alembik, Y., Barth, M., Bonneau, D., Colin, E., Dollfus, H., Doray, B., Delrue, M. A., Drouin-Garraud, V., Flori, E., Fradin, M., Francannet, C., Goldenberg, A., Lumbroso, S., Mathieu-Dramard, M., Martin-Coignard, D., Lacombe, D., Morin, G., Polge, A., Sukno, S., Thauvin-Robinet, C., Thevenon, J., Doco-Fenzy, M., Genevieve, D., Sarda, P., Edery, P., Isidor, B., Jost, B., Olivier-Faivre, L., Mandel, J. L., & Piton, A. (2014). Efficient strategy for the molecular diagnosis of intellectual disability using targeted high-throughput sequencing. *J Med Genet*, *51*, 724-736.
- Ronan, A., Fagan, K., Christie, L., Conroy, J., Nowak, N. J., & Turner, G. (2007). Familial 4.3 Mb duplication of 21q22 sheds new light on the Down syndrome critical region. *J Med Genet*, *44*, 448-451.
- Rothweiler, U., Stensen, W., Brandsdal, B. O., Isaksson, J., Leeson, F. A., Engh, R. A., & Svendsen, J. S. (2016). Probing the ATP-Binding Pocket of Protein Kinase DYRK1A with Benzothiazole Fragment Molecules. *J Med Chem*, *59*, 9814-9824.
- Ruad, L., Mignot, C., Guet, A., Ohl, C., Nava, C., Heron, D., Keren, B., Depienne, C., Benoit, V., Maystadt, I., Lederer, D., Amsallem, D., & Piard, J. (2015). DYRK1A mutations in two unrelated patients. *Eur J Med Genet*, *58*, 168-174.
- Ruben, K., Wurzlbauer, A., Walte, A., Sippl, W., Bracher, F., & Becker, W. (2015). Selectivity Profiling and Biological Activity of Novel beta-Carbolines as Potent and Selective DYRK1 Kinase Inhibitors. *PLoS One*, *10*, e0132453.
- Ruiz-Mejias, M., Martinez de Lagran, M., Mattia, M., Castano-Prat, P., Perez-Mendez, L., Ciria-Suarez, L., Gener, T., Sancristobal, B., Garcia-Ojalvo, J., Gruart, A., Delgado-Garcia, J. M., Sanchez-Vives, M. V., & Dierssen, M. (2016). Overexpression of Dyrk1A, a Down Syndrome Candidate, Decreases Excitability and Impairs Gamma Oscillations in the Prefrontal Cortex. *J Neurosci*, *36*, 3648-3659.
- Ryoo, S. R., Cho, H. J., Lee, H. W., Jeong, H. K., Radnaabazar, C., Kim, Y. S., Kim, M. J., Son, M. Y., Seo, H., Chung, S. H., & Song, W. J. (2008). Dual-specificity tyrosine(Y)-phosphorylation regulated

- kinase 1A-mediated phosphorylation of amyloid precursor protein: evidence for a functional link between Down syndrome and Alzheimer's disease. *J Neurochem*, *104*, 1333-1344.
- Ryu, Y. S., Park, S. Y., Jung, M. S., Yoon, S. H., Kwen, M. Y., Lee, S. Y., Choi, S. H., Radnaabazar, C., Kim, M. K., Kim, H., Kim, K., Song, W. J., & Chung, S. H. (2010). Dyrk1A-mediated phosphorylation of Presenilin 1: a functional link between Down syndrome and Alzheimer's disease. *J Neurochem*, *115*, 574-584.
- Sadasivam, S., & DeCaprio, J. A. (2013). The DREAM complex: master coordinator of cell cycle-dependent gene expression. *Nat Rev Cancer*, *13*, 585-595.
- Saheki, Y., & De Camilli, P. (2012). Synaptic vesicle endocytosis. *Cold Spring Harb Perspect Biol*, *4*, a005645.
- Salehi, A., Delcroix, J. D., Belichenko, P. V., Zhan, K., Wu, C., Valletta, J. S., Takimoto-Kimura, R., Kleschevnikov, A. M., Sambamurti, K., Chung, P. P., Xia, W., Villar, A., Campbell, W. A., Kulnane, L. S., Nixon, R. A., Lamb, B. T., Epstein, C. J., Stokin, G. B., Goldstein, L. S., & Mobley, W. C. (2006). Increased App expression in a mouse model of Down's syndrome disrupts NGF transport and causes cholinergic neuron degeneration. *Neuron*, *51*, 29-42.
- Sanes, J. R., & Yamagata, M. (1999). Formation of lamina-specific synaptic connections. *Curr Opin Neurobiol*, *9*, 79-87.
- Scales, T. M., Lin, S., Kraus, M., Goold, R. G., & Gordon-Weeks, P. R. (2009). Nonprimed and DYRK1A-primed GSK3 beta-phosphorylation sites on MAP1B regulate microtubule dynamics in growing axons. *J Cell Sci*, *122*, 2424-2435.
- Schimmel, M. S., Hammerman, C., Bromiker, R., & Berger, I. (2006). Third ventricle enlargement among newborn infants with trisomy 21. *Pediatrics*, *117*, e928-931.
- Schmid, S. L., McNiven, M. A., & De Camilli, P. (1998). Dynamin and its partners: a progress report. *Curr Opin Cell Biol*, *10*, 504-512.
- Sebrie, C., Chabert, C., Ledru, A., Guedj, F., Po, C., Smith, D. J., Rubin, E., Rivals, I., Beloeil, J. C., Gillet, B., & Delabar, J. M. (2008). Increased dosage of DYRK1A and brain volumetric alterations in a YAC model of partial trisomy 21. *Anat Rec (Hoboken)*, *291*, 254-262.
- Seifert, A., Allan, L. A., & Clarke, P. R. (2008). DYRK1A phosphorylates caspase 9 at an inhibitory site and is potently inhibited in human cells by harmine. *FEBS J*, *275*, 6268-6280.
- Shaikh, M. N., Gutierrez-Avino, F., Colonques, J., Ceron, J., Hammerle, B., & Tejedor, F. J. (2016). Minibrain drives the Dacapo-dependent cell cycle exit of neurons in the Drosophila brain by promoting asense and prospero expression. *Development*, *143*, 3195-3205.
- Shen, W., Taylor, B., Jin, Q., Nguyen-Tran, V., Meeusen, S., Zhang, Y. Q., Kamireddy, A., Swafford, A., Powers, A. F., Walker, J., Lamb, J., Bursalaya, B., DiDonato, M., Harb, G., Qiu, M., Filippi, C. M., Deaton, L., Turk, C. N., Suarez-Pinzon, W. L., Liu, Y., Hao, X., Mo, T., Yan, S., Li, J., Herman, A. E., Hering, B. J., Wu, T., Martin Seidel, H., McNamara, P., Glynn, R., & Laffitte, B. (2015). Inhibition of DYRK1A and GSK3B induces human beta-cell proliferation. *Nat Commun*, *6*, 8372.
- Shi, J., Zhang, T., Zhou, C., Chohan, M. O., Gu, X., Wegiel, J., Zhou, J., Hwang, Y. W., Iqbal, K., Grundke-Iqbal, I., Gong, C. X., & Liu, F. (2008). Increased dosage of Dyrk1A alters alternative splicing factor (ASF)-regulated alternative splicing of tau in Down syndrome. *J Biol Chem*, *283*, 28660-28669.
- Shin, J. H., Guedj, F., Delabar, J. M., & Lubec, G. (2007). Dysregulation of growth factor receptor-bound protein 2 and fascin in hippocampus of mice polytransgenic for chromosome 21 structures. *Hippocampus*, *17*, 1180-1192.
- Shirasaki, R., Lewcock, J. W., Lettieri, K., & Pfaff, S. L. (2006). FGF as a target-derived chemoattractant for developing motor axons genetically programmed by the LIM code. *Neuron*, *50*, 841-853.
- Sitz, J. H., Baumgartel, K., Hammerle, B., Papadopoulos, C., Hekerman, P., Tejedor, F. J., Becker, W., & Lutz, B. (2008). The Down syndrome candidate dual-specificity tyrosine phosphorylation-regulated kinase 1A phosphorylates the neurodegeneration-related septin 4. *Neuroscience*, *157*, 596-605.


- Sitz, J. H., Tigges, M., Baumgartel, K., Khaspekov, L. G., & Lutz, B. (2004). Dyrk1A potentiates steroid hormone-induced transcription via the chromatin remodeling factor Arip4. *Mol Cell Biol*, *24*, 5821-5834.
- Skurat, A. V., & Dietrich, A. D. (2004). Phosphorylation of Ser640 in muscle glycogen synthase by DYRK family protein kinases. *J Biol Chem*, *279*, 2490-2498.
- Smith, D. J., Stevens, M. E., Sudanagunta, S. P., Bronson, R. T., Makhinson, M., Watabe, A. M., O'Dell, T. J., Fung, J., Weier, H. U., Cheng, J. F., & Rubin, E. M. (1997). Functional screening of 2 Mb of human chromosome 21q22.2 in transgenic mice implicates minibrain in learning defects associated with Down syndrome. *Nat Genet*, *16*, 28-36.
- Sonamoto, R., Kii, I., Koike, Y., Sumida, Y., Kato-Sumida, T., Okuno, Y., Hosoya, T., & Hagiwara, M. (2015). Identification of a DYRK1A Inhibitor that Induces Degradation of the Target Kinase using Co-chaperone CDC37 fused with Luciferase nanoKAZ. *Sci Rep*, *5*, 12728.
- Song, W. J., Song, E. A., Jung, M. S., Choi, S. H., Baik, H. H., Jin, B. K., Kim, J. H., & Chung, S. H. (2015). Phosphorylation and inactivation of glycogen synthase kinase 3beta (GSK3beta) by dual-specificity tyrosine phosphorylation-regulated kinase 1A (Dyrk1A). *J Biol Chem*, *290*, 2321-2333.
- Song, W. J., Sternberg, L. R., Kasten-Sportes, C., Keuren, M. L., Chung, S. H., Slack, A. C., Miller, D. E., Glover, T. W., Chiang, P. W., Lou, L., & Kurnit, D. M. (1996). Isolation of human and murine homologues of the Drosophila minibrain gene: human homologue maps to 21q22.2 in the Down syndrome "critical region". *Genomics*, *38*, 331-339.
- Soppa, U., Schumacher, J., Florencio Ortiz, V., Pasqualon, T., Tejedor, F. J., & Becker, W. (2014). The Down syndrome-related protein kinase DYRK1A phosphorylates p27(Kip1) and Cyclin D1 and induces cell cycle exit and neuronal differentiation. *Cell Cycle*, *13*, 2084-2100.
- Souchet, B., Guedj, F., Penke-Verdier, Z., Daubigney, F., Duchon, A., Herault, Y., Bizot, J. C., Janel, N., Creau, N., Delatour, B., & Delabar, J. M. (2015). Pharmacological correction of excitation/inhibition imbalance in Down syndrome mouse models. *Front Behav Neurosci*, *9*, 267.
- Souchet, B., Guedj, F., Sahun, I., Duchon, A., Daubigney, F., Badel, A., Yanagawa, Y., Barallobre, M. J., Dierssen, M., Yu, E., Herault, Y., Arbones, M., Janel, N., Creau, N., & Delabar, J. M. (2014). Excitation/inhibition balance and learning are modified by Dyrk1a gene dosage. *Neurobiol Dis*, *69*, 65-75.
- Spence, E. F., & Soderling, S. H. (2015). Actin Out: Regulation of the Synaptic Cytoskeleton. *J Biol Chem*, *290*, 28613-28622.
- Stagni, F., Giacomini, A., Emili, M., Trazzi, S., Guidi, S., Sassi, M., Ciani, E., Rimondini, R., & Bartesaghi, R. (2016). Short- and long-term effects of neonatal pharmacotherapy with epigallocatechin-3-gallate on hippocampal development in the Ts65Dn mouse model of Down syndrome. *Neuroscience*, *333*, 277-301.
- Stessman, H. A., Xiong, B., Coe, B. P., Wang, T., Hoekzema, K., Fenckova, M., Kvarnung, M., Gerds, J., Trinh, S., Cosemans, N., Vives, L., Lin, J., Turner, T. N., Santen, G., Ruivenkamp, C., Kriek, M., van Haeringen, A., Aten, E., Friend, K., Liebelt, J., Barnett, C., Haan, E., Shaw, M., Gecz, J., Anderlid, B. M., Nordgren, A., Lindstrand, A., Schwartz, C., Kooy, R. F., Vandeweyer, G., Helsmoortel, C., Romano, C., Alberti, A., Vinci, M., Avola, E., Giusto, S., Courchesne, E., Pramparo, T., Pierce, K., Nalabolu, S., Amaral, D. G., Scheffer, I. E., Delatycki, M. B., Lockhart, P. J., Hormozdiari, F., Harich, B., Castells-Nobau, A., Xia, K., Peeters, H., Nordenskjold, M., Schenck, A., Bernier, R. A., & Eichler, E. E. (2017). Targeted sequencing identifies 91 neurodevelopmental-disorder risk genes with autism and developmental-disability biases. *Nat Genet*, *49*, 515-526.
- Tahtouh, T., Elkins, J. M., Filippakopoulos, P., Soundararajan, M., Burgy, G., Durieu, E., Cochet, C., Schmid, R. S., Lo, D. C., Delhommel, F., Oberholzer, A. E., Pearl, L. H., Carreaux, F., Bazureau, J. P., Knapp, S., & Meijer, L. (2012). Selectivity, cocrystal structures, and neuroprotective properties of leucettines, a family of protein kinase inhibitors derived from the marine sponge alkaloid leucettamine B. *J Med Chem*, *55*, 9312-9330.

- Takashima, S., Iida, K., Mito, T., & Arima, M. (1994). Dendritic and histochemical development and ageing in patients with Down's syndrome. *J Intellect Disabil Res*, *38* (Pt 3), 265-273.
- Tang, H., Hammack, C., Ogden, S. C., Wen, Z., Qian, X., Li, Y., Yao, B., Shin, J., Zhang, F., Lee, E. M., Christian, K. M., Didier, R. A., Jin, P., Song, H., & Ming, G. L. (2016). Zika Virus Infects Human Cortical Neural Progenitors and Attenuates Their Growth. *Cell Stem Cell*, *18*, 587-590.
- Tejedor, F., Zhu, X. R., Kaltenbach, E., Ackermann, A., Baumann, A., Canal, I., Heisenberg, M., Fischbach, K. F., & Pongs, O. (1995). minibrain: a new protein kinase family involved in postembryonic neurogenesis in *Drosophila*. *Neuron*, *14*, 287-301.
- Thomazeau, A., Lassalle, O., Iafrafi, J., Souchet, B., Guedj, F., Janel, N., Chavis, P., Delabar, J., & Manzoni, O. J. (2014). Prefrontal deficits in a murine model overexpressing the down syndrome candidate gene *dyrk1a*. *J Neurosci*, *34*, 1138-1147.
- Thompson, B. J., Bhansali, R., Diebold, L., Cook, D. E., Stolzenburg, L., Casagrande, A. S., Besson, T., Leblond, B., Desire, L., Malinge, S., & Crispino, J. D. (2015). DYRK1A controls the transition from proliferation to quiescence during lymphoid development by destabilizing Cyclin D3. *J Exp Med*, *212*, 953-970.
- Toiber, D., Azkona, G., Ben-Ari, S., Toran, N., Soreq, H., & Dierssen, M. (2010). Engineering DYRK1A overdosage yields Down syndrome-characteristic cortical splicing aberrations. *Neurobiol Dis*, *40*, 348-359.
- Tschop, K., Conery, A. R., Litovchick, L., Decaprio, J. A., Settleman, J., Harlow, E., & Dyson, N. (2011). A kinase shRNA screen links LATS2 and the pRB tumor suppressor. *Genes Dev*, *25*, 814-830.
- Vacca, R. A., & Valenti, D. (2015). Green tea EGCG plus fish oil omega-3 dietary supplements rescue mitochondrial dysfunctions and are safe in a Down's syndrome child. *Clin Nutr*, *34*, 783-784.
- Valetto, A., Orsini, A., Bertini, V., Toschi, B., Bonuccelli, A., Simi, F., Sammartino, I., Taddeucci, G., Simi, P., & Saggese, G. (2012). Molecular cytogenetic characterization of an interstitial deletion of chromosome 21 (21q22.13q22.3) in a patient with dysmorphic features, intellectual disability and severe generalized epilepsy. *Eur J Med Genet*, *55*, 362-366.
- van Bon, B. W., Coe, B. P., Bernier, R., Green, C., Gerdts, J., Witherspoon, K., Kleefstra, T., Willemsen, M. H., Kumar, R., Bosco, P., Fichera, M., Li, D., Amaral, D., Cristofoli, F., Peeters, H., Haan, E., Romano, C., Mefford, H. C., Scheffer, I., Gecz, J., de Vries, B. B., & Eichler, E. E. (2016). Disruptive de novo mutations of DYRK1A lead to a syndromic form of autism and ID. *Mol Psychiatry*, *21*, 126-132.
- Vazquez-Higuera, J. L., Sanchez-Juan, P., Rodriguez-Rodriguez, E., Mateo, I., Pozueta, A., Frank, A., Sastre, I., Valdivieso, F., Berciano, J., Bullido, M. J., & Combarros, O. (2009). DYRK1A genetic variants are not linked to Alzheimer's disease in a Spanish case-control cohort. *BMC Med Genet*, *10*, 129.
- Vidaki, M., Drees, F., Saxena, T., Lanslots, E., Taliaferro, M. J., Tatarakis, A., Burge, C. B., Wang, E. T., & Gertler, F. B. (2017). A Requirement for Mena, an Actin Regulator, in Local mRNA Translation in Developing Neurons. *Neuron*, *95*, 608-622 e605.
- von Groote-Bidlingmaier, F., Schmoll, D., Orth, H. M., Joost, H. G., Becker, W., & Barthel, A. (2003). DYRK1 is a co-activator of FKHR (FOXO1a)-dependent glucose-6-phosphatase gene expression. *Biochem Biophys Res Commun*, *300*, 764-769.
- Walte, A., Ruben, K., Birner-Gruenberger, R., Preisinger, C., Bamberg-Lemper, S., Hilz, N., Bracher, F., & Becker, W. (2013). Mechanism of dual specificity kinase activity of DYRK1A. *FEBS J*, *280*, 4495-4511.
- Wegiel, J., Gong, C. X., & Hwang, Y. W. (2011). The role of DYRK1A in neurodegenerative diseases. *FEBS J*, *278*, 236-245.
- Wegiel, J., Kaczmarek, W., Barua, M., Kuchna, I., Nowicki, K., Wang, K. C., Wegiel, J., Yang, S. M., Frackowiak, J., Mazur-Kolecka, B., Silverman, W. P., Reisberg, B., Monteiro, I., de Leon, M., Wisniewski, T., Dalton, A., Lai, F., Hwang, Y. W., Adayev, T., Liu, F., Iqbal, K., Iqbal, I. G., & Gong, C. X. (2011). Link between DYRK1A overexpression and several-fold enhancement of neurofibrillary degeneration with 3-repeat tau protein in Down syndrome. *J Neuropathol Exp Neurol*, *70*, 36-50.

- Wegiel, J., Kuchna, I., Nowicki, K., Frackowiak, J., Dowjat, K., Silverman, W. P., Reisberg, B., DeLeon, M., Wisniewski, T., Adayev, T., Chen-Hwang, M. C., & Hwang, Y. W. (2004). Cell type- and brain structure-specific patterns of distribution of minibrain kinase in human brain. *Brain Res*, *1010*, 69-80.
- Widowati, E. W., Ernst, S., Hausmann, R., Muller-Newen, G., & Becker, W. (2018). Functional characterization of DYRK1A missense variants associated with a syndromic form of intellectual deficiency and autism. *Biol Open*, *7*.
- Wiechmann, S., Czajkowska, H., de Graaf, K., Grotzinger, J., Joost, H. G., & Becker, W. (2003). Unusual function of the activation loop in the protein kinase DYRK1A. *Biochem Biophys Res Commun*, *302*, 403-408.
- Winkle, C. C., & Gupton, S. L. (2016). Membrane Trafficking in Neuronal Development: Ins and Outs of Neural Connectivity. *Int Rev Cell Mol Biol*, *322*, 247-280.
- Woods, Y. L., Cohen, P., Becker, W., Jakes, R., Goedert, M., Wang, X., & Proud, C. G. (2001). The kinase DYRK phosphorylates protein-synthesis initiation factor eIF2Bepsilon at Ser539 and the microtubule-associated protein tau at Thr212: potential role for DYRK as a glycogen synthase kinase 3-priming kinase. *Biochem J*, *355*, 609-615.
- Woods, Y. L., Rena, G., Morrice, N., Barthel, A., Becker, W., Guo, S., Unterman, T. G., & Cohen, P. (2001). The kinase DYRK1A phosphorylates the transcription factor FKHR at Ser329 in vitro, a novel in vivo phosphorylation site. *Biochem J*, *355*, 597-607.
- Xiang, J., Yang, S., Xin, N., Gaertig, M. A., Reeves, R. H., Li, S., & Li, X. J. (2017). DYRK1A regulates Hap1-Dcaf7/WDR68 binding with implication for delayed growth in Down syndrome. *Proc Natl Acad Sci U S A*, *114*, E1224-E1233.
- Xiang, Z., Haroutunian, V., Ho, L., Purohit, D., & Pasinetti, G. M. (2006). Microglia activation in the brain as inflammatory biomarker of Alzheimer's disease neuropathology and clinical dementia. *Dis Markers*, *22*, 95-102.
- Xie, W., Adayev, T., Zhu, H., Wegiel, J., Wieraszko, A., & Hwang, Y. W. (2012). Activity-dependent phosphorylation of dynamin 1 at serine 857. *Biochemistry*, *51*, 6786-6796.
- Xie, W., Ramakrishna, N., Wieraszko, A., & Hwang, Y. W. (2008). Promotion of neuronal plasticity by (-)-epigallocatechin-3-gallate. *Neurochem Res*, *33*, 776-783.
- Xing, B., Li, Y. C., & Gao, W. J. (2016). Norepinephrine versus dopamine and their interaction in modulating synaptic function in the prefrontal cortex. *Brain Res*, *1641*, 217-233.
- Yadav, R. R., Sharma, S., Joshi, P., Wani, A., Vishwakarma, R. A., Kumar, A., & Bharate, S. B. (2015). Meridianin derivatives as potent Dyrk1A inhibitors and neuroprotective agents. *Bioorg Med Chem Lett*, *25*, 2948-2952.
- Yamamoto, N., Shibata, M., Ishikuro, R., Tanida, M., Taniguchi, Y., Ikeda-Matsuo, Y., & Sobue, K. (2017). Epigallocatechin gallate induces extracellular degradation of amyloid beta-protein by increasing neprilysin secretion from astrocytes through activation of ERK and PI3K pathways. *Neuroscience*, *362*, 70-78.
- Yamamoto, T., Shimojima, K., Nishizawa, T., Matsuo, M., Ito, M., & Imai, K. (2011). Clinical manifestations of the deletion of Down syndrome critical region including DYRK1A and KCNJ6. *Am J Med Genet A*, *155A*, 113-119.
- Yang, E. J., Ahn, Y. S., & Chung, K. C. (2001). Protein kinase Dyrk1 activates cAMP response element-binding protein during neuronal differentiation in hippocampal progenitor cells. *J Biol Chem*, *276*, 39819-39824.
- Yang, L., Paul, S., Trieu, K. G., Dent, L. G., Frolidi, F., Fores, M., Webster, K., Siegfried, K. R., Kondo, S., Harvey, K., Cheng, L., Jimenez, G., Shvartsman, S. Y., & Veraksa, A. (2016). Minibrain and Wings apart control organ growth and tissue patterning through down-regulation of Capicua. *Proc Natl Acad Sci U S A*, *113*, 10583-10588.
- Yin, X., Jin, N., Gu, J., Shi, J., Zhou, J., Gong, C. X., Iqbal, K., Grundke-Iqbal, I., & Liu, F. (2012). Dual-specificity tyrosine phosphorylation-regulated kinase 1A (Dyrk1A) modulates serine/arginine-rich protein 55 (SRp55)-promoted Tau exon 10 inclusion. *J Biol Chem*, *287*, 30497-30506.


- Yin, X., Jin, N., Shi, J., Zhang, Y., Wu, Y., Gong, C. X., Iqbal, K., & Liu, F. (2017). Dyrk1A overexpression leads to increase of 3R-tau expression and cognitive deficits in Ts65Dn Down syndrome mice. *Sci Rep*, *7*, 619.
- Yoshiike, Y., Kimura, T., Yamashita, S., Furudate, H., Mizoroki, T., Murayama, M., & Takashima, A. (2008). GABA(A) receptor-mediated acceleration of aging-associated memory decline in APP/PS1 mice and its pharmacological treatment by picrotoxin. *PLoS One*, *3*, e3029.
- Yu, L., Daniels, J. P., Wu, H., & Wolf, M. J. (2015). Cardiac hypertrophy induced by active Raf depends on Yorkie-mediated transcription. *Sci Signal*, *8*, ra13.
- Yuen, E. Y., Qin, L., Wei, J., Liu, W., Liu, A., & Yan, Z. (2014). Synergistic regulation of glutamatergic transmission by serotonin and norepinephrine reuptake inhibitors in prefrontal cortical neurons. *J Biol Chem*, *289*, 25177-25185.
- Zhang, Y., Liao, J. M., Zeng, S. X., & Lu, H. (2011). p53 downregulates Down syndrome-associated DYRK1A through miR-1246. *EMBO Rep*, *12*, 811-817.


A



B








Time →

Neurogenesis

Astroglialogenesis





3. Transcription factors and affected transcripts

- CREB
- NFAT
- FKHR
- Neuroligin
- TrkB1
- BDNF
- AchE-S
- AchE-R

1. Membrane trafficking

- Dynamin 1
- Amphiphysin 1
- Endophilin
- Synaptojanin 1
- AP180

Activity-dependent transcription



2. Cytoskeleton rearrangement

- Actin filament
- Microtubule
- MT-associated protein tau

- Synaptic receptor
- Transcription factor
- Actin filament
- Clathrin-coated vesicle
- Neurexin/neuroligin
- Microtubule
- Tau

Soma

