



Physical injury and depression in six low- and middle-income countries: A nationally representative study

Louis Jacob, Damiano Pizzol, Nicola Veronese, Brendon Stubbs, Ai Koyanagi

► To cite this version:

Louis Jacob, Damiano Pizzol, Nicola Veronese, Brendon Stubbs, Ai Koyanagi. Physical injury and depression in six low- and middle-income countries: A nationally representative study. *Journal of Affective Disorders*, 2019, 248, pp.99 - 107. <10.1016/j.jad.2019.01.023>. <hal-03485688>

HAL Id: hal-03485688

<https://hal.science/hal-03485688v1>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

**Physical injury and depression in six low- and middle-income countries:
a nationally representative study**

Louis Jacob PhD^{a,*}; Damiano Pizzol MD-PhD^b; Nicola Veronese MD-PhD^c;
Brendon Stubbs PhD^{d,e}; Ai Koyanagi MD-PhD^{f,g}

^a Faculty of Medicine, University of Versailles Saint-Quentin-en-Yvelines,
Montigny-le-Bretonneux 78180, France

^b Operational Research Unit, Doctors with Africa, Mozambique

^c Aging Branch National Research Council, Neuroscience Institute, Padova,
Italy

^d Department of Psychological Medicine, Institute of Psychiatry, Psychology
and Neuroscience, King's College London, London, United Kingdom

^e Physiotherapy Department, South London and Maudsley NHS Foundation
Trust, Denmark Hill, London SE5 8AZ, United Kingdom

^f Research and Development Unit, Parc Sanitari Sant Joan de Déu,
Universitat de Barcelona, Fundació Sant Joan de Déu, Dr. Antoni Pujadas,
42, Sant Boi de Llobregat, Barcelona 08830, Spain

^g Instituto de Salud Carlos III, Centro de Investigación Biomédica en Red de
Salud Mental, CIBERSAM, Monforte de Lemos 3-5 Pabellón 11, Madrid
28029, Spain

*** Correspondence:**

Dr. Louis Jacob

Faculty of Medicine, University of Versailles Saint-Quentin-en-Yvelines

2 avenue de la Source de la Bièvre, Montigny-le-Bretonneux 78180

France

Tel.: +33-(0)6-27-88-37-06

louis.jacob.contacts@gmail.com

Table count: 3

Figure count: 2

Appendix count: 3

Number of words in the main body of the article: 4150

Number of words in the abstract: 245

1. Introduction

Major depressive disorder accounted for 8.2% of years lived with disability in the world in 2010 (Ferrari et al., 2013), and this psychiatric condition is responsible for an important share of the total burden attributable to non-communicable diseases (Patel, 2007). Furthermore, depression is positively associated with low quality of life (Zeng et al., 2013), chronic diseases (Kilzieh et al., 2008), and mortality (Kozela et al., 2016). Depression is now a major problem in low- and middle-income countries (LMICs), as these countries bear more than 80% of the global years lost to disability due to depression (World Health Organization, 2017), while the majority of people with depression in LMICs do not receive appropriate treatment (Cuijpers et al., 2018). Therefore, there is an urgent need to better understand the risk factors for depression in LMICs to establish effective prevention measures for this disorder.

Numerous factors that can increase the risk for depression have been identified in LMICs. These factors include female gender (Tomlinson et al., 2009), age (Geldsetzer et al., 2018), low socioeconomic status (Fernández-Niño et al., 2014), lack of social support (Cheng et al., 2014), low self-esteem (Azizi et al., 2013), stressful life events (Tao et al., 2011), family history of psychiatric disorders (Chin et al., 2016), and chronic conditions (Das et al., 2013; Wong et al., 2013). However, physical injuries (e.g., traffic accident, fall, gunshot, burns) have rarely been investigated as a risk factor for depression in LMICs despite growing scientific evidence from high-income countries showing that depression is highly prevalent among people who have experienced injuries (O'Donnell et al., 2004; Mitchell et al.,

2008; Wiseman et al., 2015; Rayner et al., 2016; Agtarap et al., 2017; Ahl et al., 2017; Martin et al., 2017).

Investigating the physical injury-depression relationship in LMICs is of particular importance because the incidence of injuries is disproportionately high in this setting (Kalaiselvana et al., 2011; Bashah et al., 2015), with 90% of the global physical injury-related deaths and 94% of disability-adjusted life years resulting from injuries occurring in these countries (Hyder, 2013). In addition, physical injury is associated with major costs at the individual and national level in this setting (Arokiasamy and Krishnan, 1994). For example, the economic burden of motorcycle accidents was estimated to be approximately US\$ 1.2 million in Northern Ghana in 2008 (Kudebong et al., 2011). The high prevalence of physical injury in LMICs may be explained by factors such as insufficient law enforcement for the use of helmets or seat belts when driving (Routley et al., 2007; Wadhvaniya et al., 2017). In addition, there has been a rapid increase in the number of registered vehicles (Solanki et al., 2016), while traffic laws have remained relatively inadequate (Gopalakrishnan, 2012). Furthermore, some LMICs have very high rates of alcohol consumption (e.g., Eastern Europe) and this may be an important contributor to traffic injuries and other types of injury (e.g., falls, physical violence) (De Boni et al., 2011; Razvodovsky, 2012). Finally, it has been suggested that injuries may increase risk for depression via pain (Geerlings et al., 2002; Rivara et al., 2008; Stubbs et al., 2017b), impairments and disabilities in activities of daily living (Turner and Noh, 1988; Anke et al., 1997), stress (Wiseman et al., 2015; Yang et al., 2015), and job loss (Doctor et al., 2005; Andreeva et al., 2015). Thus, the impact of injuries on depression in LMICs may differ from that of high-income countries for factors such as differences in types

of injury, suboptimal treatment for injury/disability, lack of disability pensions, and high out-of-pocket costs associated with treatment which may lead to catastrophic expenditures (Wesson et al., 2014).

Although several studies have investigated the association between physical injury (or some conditions which may be attributable to injury such as fractures) and depression in LMICs (Rasmussen et al., 2007; Ukpong et al., 2007; Hawamdeh et al., 2008; de Moraes et al., 2010; Husain et al., 2010; Eldin et al., 2012; Dogu et al., 2014; Gandjalikhan-Nassab et al., 2016; Asuquo et al., 2017; Braimah et al., 2017; Chen et al., 2017; Wu et al., 2017), many of them were of small sample size (i.e., <200 participants) (Rasmussen et al., 2007; Ukpong et al., 2007; Hawamdeh et al., 2008; de Moraes et al., 2010; Eldin et al., 2012; Dogu et al., 2014; Gandjalikhan-Nassab et al., 2016; Asuquo et al., 2017; Braimah et al., 2017), conducted in either a single hospital or a limited area of a single country (Ukpong et al., 2007; Hawamdeh et al., 2008; de Moraes et al., 2010; Husain et al., 2010; Eldin et al., 2012; Dogu et al., 2014; Gandjalikhan-Nassab et al., 2016; Asuquo et al., 2017; Braimah et al., 2017; Wu et al., 2017), or lacked a control group (Rasmussen et al., 2007; Ukpong et al., 2007; Hawamdeh et al., 2008; Husain et al., 2010; Eldin et al., 2012; Dogu et al., 2014; Braimah et al., 2017; Wu et al., 2017). In addition, the few studies which did conduct an adjusted analysis failed to adjust for alcohol consumption despite the fact that alcohol consumption is known to strongly influence the likelihood of experiencing injury and also depression (Watt et al., 2004; Jorge et al., 2005). Finally, there is a lack of data on the differences in terms of risk for depression between physical injury with and without disability, despite the fact that previous

research has highlighted the major impact of disability on this psychiatric condition (Barry et al., 2013; Noh et al., 2016).

Therefore, the goal of the present study was to investigate the relationship between physical injury (with and without disability) and depression in six LMICs (China, Ghana, India, Mexico, Russia, South Africa) using nationally representative data of the Study on Global Ageing and Adult Health (SAGE). These countries broadly represent different geographical locations and levels of socio-economic and demographic transition.

2. Methods

2.1. The survey

Data from the Study on Global Ageing and Adult Health (SAGE) survey were analyzed. This dataset is publically available through the World Health Organization (WHO) website (<http://www.who.int/healthinfo/sage/en/>). The survey was undertaken in China, Ghana, India, Mexico, Russia, and South Africa between 2007 and 2010. Based on the World Bank classification at the time of the survey, Ghana was the only low-income country, and China and India were lower middle-income countries although China became an upper middle-income country in 2010. The remaining countries were upper middle-income countries. This classification is based on the gross national income (GNI) per capita (in US dollars), converted from local currency using the World Bank Atlas method. For example, the specific figures for year 2010 were: low income (\leq \$1005), lower middle income (\$1006-\$3975) and upper middle income (\$3976-\$12275).

Details of the survey methodology have been published elsewhere (Kowal et al., 2012). In brief, in order to obtain nationally representative samples, a multistage clustered sampling design method was used. The sample consisted of adults aged ≥ 18 years with oversampling of those aged ≥ 50 years. Following a common research protocol across countries, trained interviewers conducted face-to-face interviews using a standard questionnaire to collect information. The questionnaires were translated from English into the local languages, following the WHO translation guidelines. All interviews in Mexico were completed using a computer-assisted personal interview (CAPI), while a paper and pencil interview (PAPI) was used in the remaining countries with the exception of China where both CAPI and PAPI were used. The response rate for each country was: 93% (China); 81% (Ghana); 68% (India); 53% (Mexico); 83% (Russia); 75% (South Africa). Sampling weights were constructed to adjust for the population structure as reported by the United Nations Statistical Division. Ethical approval was obtained from the WHO Ethical Review Committee and local ethics research review boards. Written informed consent was obtained from all participants.

2.2. Variables

2.2.1. Depression

Questions based on the World Mental Health Survey version of the Composite International Diagnostic Interview (Kessler and Üstün, 2004) were used for the endorsement of past 12-month DSM-IV depression with the same algorithm as in previous studies using the same dataset (**eTable 1** of the supplementary material) (Koyanagi et al., 2014; Garin et al., 2016).

2.2.2. Physical injury

Physical injury was assessed with two main questions with yes/no answer options: “In the last 12 months, have you been involved in a road traffic accident where you suffered from bodily injury?” (traffic accident injury); and “In the last 12 months, have you had any other event where you suffered from bodily injury?” (other injury). For both questions, for those who responded affirmatively, follow-up questions were asked: (1) “How did the injury happen?” [answer options “it was an accident (unintentional)”, “someone else did it to me deliberately (intentional)”, “I did it to myself deliberately (self-inflicted)”]; (2) “Did you suffer a physical disability as a result of being injured?” (yes/no answer options). Any injury in our analysis referred to having experienced either or both traffic injury or other injury in the past 12 months. Variables on any injury, traffic accident injury, and other injury were used as dichotomous (yes or no) or three-category variables (no accident, accident without disability, accident with disability) in our analysis. Traffic injuries were distinguished from other types of injury in some analyses because they are associated with a major burden in LMICs (Dalal et al., 2013), and their public health implication (e.g., preventive strategies) may differ from other forms of injury. For those who suffered a physical disability, a follow-up question asked in which ways they were physically disabled. Information on the cause of “other injury” was also obtained. If there were multiple injuries, the participant was asked to refer to the most recent incident.

2.2.3. Control variables

The control variables were selected based on past literature (Clausen et al., 2015) and included sex, age, wealth (poorest, poorer, middle, richer, richest), highest level of education achieved (\leq primary, secondary, \geq tertiary), setting (rural or urban), and

alcohol consumption (never, non-heavy, heavy). Wealth reflected capital at the individual level, whereas country income reflected capital at the national level.

Wealth quintiles were derived from the household ownership of durable goods (e.g., chairs, television, access to electricity). Using a Bayesian post-estimation method, households were arranged in increasing order of assets. The raw continuous income estimates were classified into country-wise quintiles using the asset order. Alcohol consumption was assessed by the question “Have you ever consumed a drink that contains alcohol (such as beer, wine, spirits, etc)?” Those who answered “no” were categorized as “never” drinkers. For those answering “yes”, a separate question asked about how many drinks of any alcohol beverage they had consumed on each day of the past week. Consumers of at least 4 (females) or 5 drinks (males) of any alcoholic beverage per day on at least one day in the past week were considered “heavy” drinkers. Those who had ever consumed alcohol but were not heavy drinkers were categorized as “non-heavy” drinkers (Koyanagi et al., 2015).

2.3. Statistical analysis

The statistical analysis was performed with Stata 14.1 (Stata Corp LP, College station, Texas). We conducted multivariable logistic regression analysis to assess the association between the three-category physical injury variable (exposure; i.e., no injury, injury without disability, injury with disability) and depression (outcome) using the overall sample. As a sensitivity analysis, we also assessed whether the results remain similar after the omission of those who had a self-inflicted physical injury as self-inflicted injury may have a different effect on mental health.

Country-wise analysis was also conducted with the dichotomous any injury variable (yes/no) as the exposure variable. We did not use the three-category variable for this

part of the analysis due to the limited sample size in each country. In order to assess the between-country heterogeneity that may exist in the association between injury and depression, we calculated the Higgin's I^2 based on estimates for each country. The Higgin's I^2 represents the degree of heterogeneity that is not explained by sampling error with a value of 25% often considered as low, 50% as moderate, and 75% as high heterogeneity (Higgins et al., 2003). In line with previous studies using multi-country data (Prince et al., 2012; Koyanagi et al., 2016; Stubbs et al., 2017a; DeVlyder et al., 2018), a pooled estimate was obtained by random-effect meta-analysis based on country-wise estimates using the Stata *metan* command.

All models were adjusted for sex, age, wealth, education, setting, and alcohol consumption. The model using the overall sample was additionally adjusted for country by including dummy variables for each country (Koyanagi et al., 2014). All variables were included in the models as categorical variables with the exception of age. The sample weighting and the complex study design were taken into account in all analyses with the use of the Taylor linearization method. Results from the logistic regression models are presented as odds ratios (OR) with 95% confidence intervals (CI). The level of statistical significance was set at $P < 0.05$.

3. Results

The final sample included 42489 individuals aged ≥ 18 years (China $n=14813$; Ghana $n=5110$; India $n=11230$; Mexico $n=2756$; Russia $n=4355$; South Africa $n=4225$). The mean age (SD) of the sample was 43.8 (14.4) years and 50.1% were females (**Table 1**). The percentage of those who were ≥ 50 years was 27.3% (unweighted $N=34,129$), and the age range was 18-114 years. The overall prevalence of

depression was 4.1%. Overall, the prevalence of physical injuries with and without disability were, respectively, 1.4% and 6.5% for any injury, 0.4% and 2.0% for injuries due to road traffic accident, and 1.0% and 4.7% for injuries due to other cause. The most common causes of injury not related with traffic accidents were a fall (59.3%), struck/hit by a person or object (13.1%), and stabbed (7.4%) in the overall sample (**eTable 2** of the supplementary material). There was a wide variation in terms of the major causes of injury not related with traffic accidents between countries. The most common physical disabilities associated with an injury were difficulties using (or unable to use) hand or arm and walking with a limp (**eTable 3** of the supplementary material). The prevalence of male sex, lower levels of wealth and education, rural setting, and alcohol consumption was significantly higher among those who had any injury (**Table 1**). The sample characteristics by country are shown in **Table 2**. The prevalence of any injury was highest in India (10.7%), followed by Ghana (7.2%), and Mexico (6.4%), with the lowest prevalence being observed in South Africa (3.2%). The prevalence of disability due to any injury was highest in Mexico (3.1%) and India (2.4%).

The prevalence of depression was higher among those who had any injury in the overall sample and also country-wise samples (**Figure 1**). The results for the multivariable logistic regression analyses showed that compared to having no injury, any injury without disability (OR=1.72; 95%CI=1.18-2.50) and with disability (OR=3.81; 95%CI=2.16-6.73) were both associated with significantly higher odds for depression (**Table 3**). As for injuries due to road traffic accidents, injury without disability was not associated with depression. The deletion of those who had a self-inflicted injury **had almost no influence in** the results (data not shown). The country-wise association between any injury and depression is shown in **Figure 2**. In all

countries, any injury was associated with higher odds for depression (ORs ranging from 1.45 to 12.00) although the results were not significant in Ghana (OR=1.45; 95%CI=0.64-3.27) and Russia (OR=2.37; 95%CI=0.49-11.41). The overall pooled estimate based on a meta-analysis was 3.28 (95%CI=1.71-6.31) with a moderate level of between-country heterogeneity being observed ($I^2=63.1\%$).

4. Discussion

4.1. Main findings

This study including more than 42000 adults from six LMICs showed that the prevalence of physical injury ranged from 3.2% in South Africa to 10.7% in India. The overall prevalence of depression in our study (4.1%) was comparable to previously reported rates of depression from 36 LMICs (6.6%) and 10 high-income countries (5.5%) (Bromet et al., 2011; Stubbs et al., 2016). Overall, physical injury without disability was associated with a 1.7 times higher odds for depression, whereas injury with disability was associated with a 3.8-fold increased odds. There was a moderate level of between-country heterogeneity, with particularly strong associations being observed in Mexico and South Africa.

4.2. Interpretation of the findings

To the best of our knowledge, this is the first nationally representative study investigating the association between physical injury (including associated disability) and depression in LMICs, although numerous studies of small sample size including patients with some form of physical trauma/injury from a single hospital have been published in recent years (Rasmussen et al., 2007; Ukpong et al., 2007; Hawamdeh

et al., 2008; de Moraes et al., 2010; Husain et al., 2010; Eldin et al., 2012; Dogu et al., 2014; Gandjalikhan-Nassab et al., 2016; Asuquo et al., 2017; Braimah et al., 2017; Chen et al., 2017; Wu et al., 2017). For example, one study from India on survivors of tortures found that chronic injuries were associated with major depression (Rasmussen et al., 2007). Another prospective study from Nigeria showed that 41.2% and 11.8% of patients with facial injury had anxiety and depression, respectively, in the weeks following the trauma (Ukpong et al., 2007). Next, one study found that 40% of injured workers in Egypt were affected by depression or anxiety, and these psychiatric disorders had a negative impact on the perceived health-related quality of life (Eldin et al., 2012). Finally, an Iranian case-control study found that the prevalence of depression was 14% in patients with facial trauma whereas this figure was only 4% in controls (Gandjalikhan-Nassab et al., 2016). Taking together, these results underline the possibility that physical injury may be an important risk factor for depression in LMICs.

Although the exact mechanism linking physical injury and depression is largely unknown, several hypotheses may be proposed. First, pain due to the physical injury may increase the risk for depression. For example, almost 63% of patients with major physical trauma reported injury-related pain 12 months after the injury (Rivara et al., 2008), while pain may induce depression via fatigue (Iversen and Wessely, 2003; Salvetti et al., 2013), sleep problems (Yokoyama et al., 2010; Tang et al., 2015) or functional limitations (Smith et al., 2016; He et al., 2019). In LMICs, pain can potentially be an important cause of depression as it is often poorly managed and access to analgesics is limited in some resource-poor settings (Morris and Roques, 2018). Second, physical injury can also lead to impairments and disabilities

in activities of daily living (Anke et al., 1997), and physical disability in turn increases the risk for depression (Turner and Noh, 1988), possibly via social categorization and stereotyping, abuse, and loss of roles (e.g., at work or home) (Noh et al., 2016). In line with this hypothesis, we found in this study that the physical injury-depression relationship was much stronger in participants with a disability (OR=3.8) than in those without a disability (OR=1.7). It is possible that physical disabilities may have a particularly profound effect on mental wellbeing in LMICs as rehabilitation centers are often underdeveloped or limited in most LMICs (Xiao et al., 2017). Third, stress is a frequent reaction after experiencing physical trauma (Wiseman et al., 2015), and can trigger a depressive episode (Yang et al., 2015). The impact of physical injury on stress has been proposed to be mediated by re-experiencing of the trauma, nightmares and emotional distress when exposed to specific triggers (Wiseman et al., 2015). Another factor that can increase stress after a physical injury is out-of-pocket expenditure or catastrophic expenditure for health care (Leive and Xu, 2008; Vancampfort et al., 2017), which is common in many LMICs where the availability of health insurance is limited. Next, physical injury may also result in job loss due to disability (Doctor et al., 2005), and this can increase risk for subsequent major depression via factors such as poverty (Heflin and Iceland, 2009; Andreeva et al., 2015). Given that many jobs in LMICs are manual work (e.g., agricultural occupation) (Atkinson et al., 2016), people in LMICs may be at a particularly high risk of losing jobs and income as a result of disabilities. The lack of well-established social security schemes in many LMICs (e.g., disability pensions) may further complicate the situation. Finally, it is also possible that depression increases the risk for physical injuries via decreased vigilance, lack of attention, irritability, or insomnia (Peele and Tollerud, 2005; Kim et al., 2008; Patten et al., 2010).

Another interesting result of our study is that there was a moderate level of between-country heterogeneity in the association between physical injury and depression in the six LMICs. The physical injury-depression relationship was strongest in South Africa (OR=12.0) and weakest in Ghana (OR=1.5). This finding is of particular importance because it shows that some factors potentially buffering the impact of physical injury on depression are likely to be differentially distributed between these countries. Indeed, our data showed that the main types of injury vary substantially between countries, and it is possible that they have different effects on mental health. Apart from this, differences in levels of rehabilitation (Furlan et al., 2018), community reintegration (Sekaran et al., 2010) or psychological management (Yatham et al., 2017) between countries may also explain the between-country difference. Furthermore, country differences may also be related to how symptoms of depression are understood, expressed and reported. For example, depression is often expressed as a somatic symptom in Asian cultures (Ryder et al., 2008). Finally, the weak association between physical injury and depression in Ghana, the only low-income country included in this study, suggests that non-economic factors (e.g., resilience, hope, familial environment) may play a role in this relationship.

In our analysis, alcohol consumption was not significantly associated with depression, and thus, this factor is unlikely to have influenced the association between injury and depression. However, this may have been due to the different timeframe used for the assessment of alcohol consumption in comparison with depression and injury. Given that previous studies have found positive associations between alcohol consumption and depression or injury (Paljärvi et al., 2009; Clausen

et al., 2016), future studies should further assess the role of alcohol consumption in the association between injury and depression.

4.3. Clinical and policy implications and areas for future research

The results of our study indicate that victims of physical injury should be screened for depression in the months and years following the trauma. Clinicians should be aware that those who become disabled as the result of an injury are at especially high risk for depression. Close collaboration between health professionals involved in the management of physical injuries such as orthopedic surgeons, specialists of physical medicine and rehabilitation, or physiotherapists and mental health care workers (e.g., psychiatrists, psychologists, social workers) may be essential to improve clinical outcomes when depression is detected in patients with an injury. However, many resource-limited settings lack specialists in these fields and adequate treatment for injuries and mental health problems is often not available, while rehabilitation facilities for disabilities are also limited. This may pose a significant challenge in addressing the injury-depression comorbidity in some LMICs. Previous studies from LMICs have shown that the use of lay community health workers may be key to improve mental wellbeing in people living in resource-limited settings (Ali et al., 2003; Baker-Henningham et al., 2005; Cooper et al., 2009; Tripathy et al., 2010; Niemi et al., 2016). Thus, it is possible that community health workers may be able to address depression or reduce risk for depression of physically injured people by interpersonal psychotherapy, peer support or health information (Yatham et al., 2017).

Next, it is also theoretically possible that pain treatment, adequate rehabilitation for disability, and stress management may reduce risk for future onset of depression in people who experienced an injury. Given that little is known about the effect of these potential interventions on depression in injured individuals, investigating these issues may provide insight on how to prevent depression among trauma patients. Finally, in terms of policy implications, some LMICs lack the basic infrastructure or law enforcement to prevent injuries and this may be particularly important in areas where the consequences of injuries may be particularly severe in terms of disabilities and economic consequences due to factors such as inadequate treatment, loss of jobs, and lack of social security schemes. For example, enforcing laws on speeding, alcohol consumption, and motorcycle helmets are essential to prevent traffic injuries, and in the case of fall-related injuries, tall buildings should be equipped with window guards, and standards should be clearly defined for playground equipments (World Health Organization, 2014).

4.4. Strengths and limitations

The use of nationally representative data and the large sample size are the two main strengths of this study. However, our findings should be interpreted in light of several limitations. First, the survey data were based on self-report, and may have been subject to some level of bias (e.g., recall bias, social desirability bias). **Second, questions on injury lacked details on the exact circumstance, severity, and requirement for hospital admission which could have provided a better understanding of the association between injury and depression. Third,** although some authors have suggested that depression and other psychiatric conditions may be better viewed as a continuous construct (Markon et al., 2011; Haslam et al., 2012;

Chen et al., 2017), we were unable to analyze depression as a continuous measure. **Fourth**, institutionalized and homeless individuals were not included in this study. Thus, the results are not generalizable to these populations which may have a high prevalence of injury and/or depression. **Fifth**, the results of our study were derived only from six LMICs and therefore, our results are not representative of all LMICs. **Next, the response rate for the SAGE survey was comparable to or higher than that of other national surveys on ageing in most of the countries included in our study. Nevertheless, there was some between-country variability and the response rate for Mexico was low. The low response rate of Mexico was partly attributable to the short time for field work where multiple visits were not possible if the respondent was not available at the initial visit. Thus, it is possible that some level of bias may exist due to non-response. However, the use of sampling weights to adjust for the population structure is likely to have partially mitigated the potential bias. Furthermore, other epidemiological studies have found that poor response rates have a minimal effect on the risk estimates and in the identification of risk factors of psychiatric disorders (Batty and Gale, 2009; Bergman et al., 2010).** Finally, it was not possible to investigate causality or temporality of the physical injury-depression relationship because of the cross-sectional design.

4.5. Conclusion

This nationally representative study found that physical injury is associated with an increased odds for depression and that this association is particularly pronounced in those who became disabled as the result of an injury in six LMICs. Clinicians should be aware of this comorbidity, and future studies should investigate the factors that underlie this comorbidity to improve clinical outcomes, and to prevent depression

onset among those who experienced an injury. Furthermore, enforcement of policies at the country-level to reduce injuries may also improve mental wellbeing among people living in LMICs.

5. References

- Agtarap, S., Boals, A., Holtz, P., Roden-Foreman, K., Rainey, E.E., Ruggero, C., Warren, A.M., 2017. The effect of depressive symptoms on social support one year following traumatic injury. *J. Affect. Disord.* 207, 398–405. <https://doi.org/10.1016/j.jad.2016.08.067>
- Ahl, R., Lindgren, R., Cao, Y., Riddez, L., Mohseni, S., 2017. Risk factors for depression following traumatic injury: An epidemiological study from a scandinavian trauma center. *Injury* 48, 1082–1087. <https://doi.org/10.1016/j.injury.2017.03.019>
- Ali, B.S., Rahbar, M.H., Naeem, S., Gul, A., Mubeen, S., Iqbal, A., 2003. The effectiveness of counseling on anxiety and depression by minimally trained counselors: a randomized controlled trial. *Am. J. Psychother.* 57, 324–336. <https://doi.org/10.1176/appi.psychotherapy.2003.57.3.324>
- Andreeva, E., Magnusson Hanson, L.L., Westerlund, H., Theorell, T., Brenner, M.H., 2015. Depressive symptoms as a cause and effect of job loss in men and women: evidence in the context of organisational downsizing from the Swedish Longitudinal Occupational Survey of Health. *BMC Public Health* 15, 1045. <https://doi.org/10.1186/s12889-015-2377-y>
- Anke, A.G., Stanghelle, J.K., Finset, A., Roaldsen, K.S., Pillgram-Larsen, J., Fugl-Meyer, A.R., 1997. Long-term prevalence of impairments and disabilities after multiple trauma. *J. Trauma* 42, 54–61.
- Arokiasamy, J.T., Krishnan, R., 1994. Some epidemiological aspects and economic costs of injuries in Malaysia. *Asia. Pac. J. Public Health* 7, 16–20. <https://doi.org/10.1177/101053959400700103>
- Asuquo, J.E., Edet, B.E., Abang, I.E., Essien, E.A., Osakwe, O.G., Aigbomain, E.J.,

- Chigbundu, K.C., 2017. Depression and posttraumatic stress disorder among road traffic accident victims managed in a Tertiary hospital in Southern Nigeria. *Niger. J. Clin. Pract.* 20, 170–175. <https://doi.org/10.4103/1119-3077.196114>
- Atkinson, K., Lowe, S., Moore, S., 2016. Human development, occupational structure and physical inactivity among 47 low and middle income countries. *Prev. Med. Rep.* 3, 40–45. <https://doi.org/10.1016/j.pmedr.2015.11.009>
- Azizi, M., Khamseh, F., Rahimi, A., Barati, M., 2013. The relationship between self-esteem and depression in nursing students of a selected medical university in Tehran. *Iran. J. Psychiatr. Nurs.* 1, 28–34.
- Baker-Henningham, H., Powell, C., Walker, S., Grantham-McGregor, S., 2005. The effect of early stimulation on maternal depression: a cluster randomised controlled trial. *Arch. Dis. Child.* 90, 1230–1234. <https://doi.org/10.1136/adc.2005.073015>
- Barry, L.C., Soulos, P.R., Murphy, T.E., Kasl, S.V., Gill, T.M., 2013. Association Between Indicators of Disability Burden and Subsequent Depression Among Older Persons. *J. Gerontol. Ser. A* 68, 286–292. <https://doi.org/10.1093/gerona/gls179>
- Bashah, D.T., Dachew, B.A., Tiruneh, B.T., 2015. Prevalence of injury and associated factors among patients visiting the Emergency Departments of Amhara Regional State Referral Hospitals, Ethiopia: a cross-sectional study. *BMC Emerg. Med.* 15, 20. <https://doi.org/10.1186/s12873-015-0044-3>
- Batty, G.D., Gale, C.R., 2009. Impact of resurvey non-response on the associations between baseline risk factors and cardiovascular disease mortality: prospective cohort study. *J. Epidemiol. Community Health* 63, 952–955. <https://doi.org/10.1136/jech.2008.086892>
- Bergman, P., Ahlberg, G., Forsell, Y., Lundberg, I., 2010. Non-Participation in the Second Wave of the Part Study on Mental Disorder and Its Effects on Risk

Estimates. Int. J. Soc. Psychiatry 56, 119–132.

<https://doi.org/10.1177/0020764008098838>

Braimah, R.O., Ukpong, D.I., Ndukwe, K.C., Akinyoola, A.L., 2017. Comparative study of anxiety and depression following maxillofacial and orthopedic injuries. Study from a Nigerian University Teaching Hospital. Clin. Exp. Dent. Res. 3, 215–219. <https://doi.org/10.1002/cre2.90>

Bromet, E., Andrade, L.H., Hwang, I., Sampson, N.A., Alonso, J., de Girolamo, G., de Graaf, R., Demyttenaere, K., Hu, C., Iwata, N., Karam, A.N., Kaur, J., Kostyuchenko, S., Lépine, J.-P., Levinson, D., Matschinger, H., Mora, M.E.M., Browne, M.O., Posada-Villa, J., Viana, M.C., Williams, D.R., Kessler, R.C., 2011. Cross-national epidemiology of DSM-IV major depressive episode. BMC Med. 9, 90. <https://doi.org/10.1186/1741-7015-9-90>

Chen, Y., Bennett, D., Clarke, R., Guo, Y., Yu, C., Bian, Z., Ma, L., Huang, Y., Sun, Q., Zhang, N., Zheng, X., Chen, J., Peto, R., Kendler, K.S., Li, L., Chen, Z., 2017. Patterns and correlates of major depression in Chinese adults: a cross-sectional study of 0.5 million men and women. Psychol. Med. 47, 958–970. <https://doi.org/10.1017/S0033291716002889>

Cheng, Y., Li, X., Lou, C., Sonenstein, F.L., Kalamar, A., Jejeebhoy, S., Delany-Moretlwe, S., Brahmabhatt, H., Olumide, A.O., Ojengbede, O., 2014. The Association Between Social Support and Mental Health Among Vulnerable Adolescents in Five Cities: Findings From the Study of the Well-Being of Adolescents in Vulnerable Environments. J. Adolesc. Health, The Well-Being of Adolescents in Vulnerable Environments Study 55, S31–S38. <https://doi.org/10.1016/j.jadohealth.2014.08.020>

Chin, W.-Y., Wan, E.Y.F., Choi, E.P.H., Chan, K.T.Y., Lam, C.L.K., 2016. The 12-Month Incidence and Predictors of PHQ-9–Screened Depressive Symptoms in

Chinese Primary Care Patients. *Ann. Fam. Med.* 14, 47–53.
<https://doi.org/10.1370/afm.1854>

Clausen, T., Martinez, P., Towers, A., Greenfield, T., Kowal, P., 2016. Alcohol Consumption at Any Level Increases Risk of Injury Caused by Others: Data from the Study on Global AGEing and Adult Health. *Subst. Abuse Res. Treat.* 9, 125–132.
<https://doi.org/10.4137/SART.S23549>

Clausen, T., Martinez, P., Towers, A., Greenfield, T., Kowal, P., 2015. Alcohol Consumption at Any Level Increases Risk of Injury Caused by Others: Data from the Study on Global AGEing and Adult Health. *Subst. Abuse Res. Treat.* 9, 125–132.
<https://doi.org/10.4137/SART.S23549>

Cooper, P.J., Tomlinson, M., Swartz, L., Landman, M., Molteno, C., Stein, A., McPherson, K., Murray, L., 2009. Improving quality of mother-infant relationship and infant attachment in socioeconomically deprived community in South Africa: randomised controlled trial. *BMJ* 338, b974.

Cuijpers, P., Karyotaki, E., Reijnders, M., Purgato, M., Barbui, C., 2018. Psychotherapies for depression in low- and middle-income countries: a meta-analysis. *World Psychiatry Off. J. World Psychiatr. Assoc. WPA* 17, 90–101.
<https://doi.org/10.1002/wps.20493>

Dalal, K., Lin, Z., Gifford, M., Svanström, L., 2013. Economics of Global Burden of Road Traffic Injuries and Their Relationship with Health System Variables. *Int. J. Prev. Med.* 4, 1442–1450.

Das, R., Singh, O., Thakurta, R.G., Khandakar, M.R., Ali, S.N., Mallick, A.K., Roy, P., Bhattacharrya, A.K., 2013. Prevalence of Depression in Patients with Type II Diabetes Mellitus and its Impact on Quality of Life. *Indian J. Psychol. Med.* 35, 284–289. <https://doi.org/10.4103/0253-7176.119502>

De Boni, R., Bozzetti, M.C., Hilgert, J., Sousa, T., Von Diemen, L., Benzano, D., Menegon, G., Holmer, B., Duarte, P. do C.A.V., Pechansky, F., 2011. Factors associated with alcohol and drug use among traffic crash victims in southern Brazil. *Accid. Anal. Prev.* 43, 1408–1413. <https://doi.org/10.1016/j.aap.2011.02.016>

de Moraes, V.Y., Jorge, M.R., Faloppa, F., Belloti, J.C., 2010. Anxiety and depression in Brazilian orthopaedics inpatients: a cross sectional study with a clinical sample comparison. *J. Clin. Psychol. Med. Settings* 17, 31–37. <https://doi.org/10.1007/s10880-009-9184-5>

DeVylder, J.E., Kelleher, I., Lalane, M., Oh, H., Link, B.G., Koyanagi, A., 2018. Association of Urbanicity With Psychosis in Low- and Middle-Income Countries. *JAMA Psychiatry* 75, 678–686. <https://doi.org/10.1001/jamapsychiatry.2018.0577>

Doctor, J.N., Castro, J., Temkin, N.R., Fraser, R.T., Machamer, J.E., Dikmen, S.S., 2005. Workers' risk of unemployment after traumatic brain injury: A normed comparison. *J. Int. Neuropsychol. Soc.* 11, 747–752. <https://doi.org/10.1017/S1355617705050836>

Dogu, B., Kuran, B., Sirzai, H., Sag, S., Akkaya, N., Sahin, F., 2014. The relationship between hand function, depression, and the psychological impact of trauma in patients with traumatic hand injury. *Int. J. Rehabil. Res. Int. Z. Rehabil. Rev. Int. Rech. Readaptation* 37, 105–109. <https://doi.org/10.1097/MRR.0000000000000040>

Eldin, W.S., Hirshon, J.M., Smith, G.S., Kamal, A.-A.M., Abou-El-Fetouh, A., El-Setouhy, M., 2012. Health-related quality of life after serious occupational injury in Egyptian workers: a cross-sectional study. *BMJ Open* 2, e000413. <https://doi.org/10.1136/bmjopen-2011-000413>

Fernández-Niño, J.A., Manrique-Espinoza, B.S., Bojorquez-Chapela, I., Salinas-Rodríguez, A., 2014. Income inequality, socioeconomic deprivation and depressive

symptoms among older adults in Mexico. *PloS One* 9, e108127.
<https://doi.org/10.1371/journal.pone.0108127>

Ferrari, A.J., Charlson, F.J., Norman, R.E., Patten, S.B., Freedman, G., Murray, C.J.L., Vos, T., Whiteford, H.A., 2013. Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010. *PLoS Med.* 10, e1001547. <https://doi.org/10.1371/journal.pmed.1001547>

Furlan, A.D., Irvin, E., Munhall, C., Giraldo-Prieto, M., Fullerton, L., McMaster, R., Danak, S., Costante, A., Pitzul, K., Bhide, R.P., Marchenko, S., Mahood, Q., David, J.A., Flannery, J.F., Bayley, M., 2018. Rehabilitation service models for people with physical and/or mental disability living in low- and middle-income countries: A systematic review. *J. Rehabil. Med.* 50, 487–498. <https://doi.org/10.2340/16501977-2325>

Gandjalikhan-Nassab, S. -a.-H., Samieirad, S., Vakil-Zadeh, M., Habib-Aghahi, R., Alsadat-Hashemipour, M., 2016. Depression and anxiety disorders in a sample of facial trauma: A study from Iran. *Med. Oral Patol. Oral Cirugia Bucal* 21, e477-482.

Garin, N., Koyanagi, A., Chatterji, S., Tyrovolas, S., Olaya, B., Leonardi, M., Lara, E., Koskinen, S., Tobiasz-Adamczyk, B., Ayuso-Mateos, J.L., Haro, J.M., 2016. Global Multimorbidity Patterns: A Cross-Sectional, Population-Based, Multi-Country Study. *J. Gerontol. A. Biol. Sci. Med. Sci.* 71, 205–214.
<https://doi.org/10.1093/gerona/glv128>

Geerlings, S.W., Twisk, J.W.R., Beekman, A.T.F., Deeg, D.J.H., van Tilburg, W., 2002. Longitudinal relationship between pain and depression in older adults: sex, age and physical disability. *Soc. Psychiatry Psychiatr. Epidemiol.* 37, 23–30.
<https://doi.org/10.1007/s127-002-8210-2>

Geldsetzer, P., Vaikath, M., Wagner, R., Rohr, J.K., Montana, L., Gómez-Olivé, F.X.,

Rosenberg, M.S., Manne-Goehler, J., Mateen, F.J., Payne, C.F., Kahn, K., Tollman, S.M., Salomon, J.A., Gaziano, T.A., Bärnighausen, T., Berkman, L.F., 2018. Depressive Symptoms and their Relation to Age and Chronic diseases among middle-aged and Older Adults in rural South Africa. *J. Gerontol. A. Biol. Sci. Med. Sci.* <https://doi.org/10.1093/gerona/gly145>

Gopalakrishnan, S., 2012. A Public Health Perspective of Road Traffic Accidents. *J. Fam. Med. Prim. Care* 1, 144–150. <https://doi.org/10.4103/2249-4863.104987>

Haslam, N., Holland, E., Kuppens, P., 2012. Categories versus dimensions in personality and psychopathology: a quantitative review of taxometric research. *Psychol. Med.* 42, 903–920. <https://doi.org/10.1017/S0033291711001966>

Hawamdeh, Z.M., Othman, Y.S., Ibrahim, A.I., 2008. Assessment of anxiety and depression after lower limb amputation in Jordanian patients. *Neuropsychiatr. Dis. Treat.* 4, 627–633.

He, M., Ma, J., Ren, Z., Zhou, G., Gong, P., Liu, M., Yang, X., Xiong, W., Wang, Q., Liu, H., Zhang, X., 2019. Association between activities of daily living disability and depression symptoms of middle-aged and older Chinese adults and their spouses: A community based study. *J. Affect. Disord.* 242, 135–142. <https://doi.org/10.1016/j.jad.2018.08.060>

Heflin, C.M., Iceland, J., 2009. Poverty, Material Hardship and Depression. *Soc. Sci. Q.* 90, 1051–1071. <https://doi.org/10.1111/j.1540-6237.2009.00645.x>

Higgins, J.P.T., Thompson, S.G., Deeks, J.J., Altman, D.G., 2003. Measuring inconsistency in meta-analyses. *BMJ* 327, 557–560. <https://doi.org/10.1136/bmj.327.7414.557>

Husain, N., Humail, S.M., Chaudhry, I.B., Rahman, R., Robinson, H., Creed, F., 2010. Psychological distress among patients of an orthopaedic outpatient clinic: a

study from a low-income country. *Ann. Gen. Psychiatry* 9, 9.
<https://doi.org/10.1186/1744-859X-9-9>

Hyder, A.A., 2013. Injuries in low- and middle-income countries: a neglected disease in global public health. *Injury* 44, 579–580.
<https://doi.org/10.1016/j.injury.2013.01.028>

Iversen, A., Wessely, S., 2003. Chronic fatigue and depression. *Curr. Opin. Psychiatry* 16, 17.

Jorge, R.E., Starkstein, S.E., Arndt, S., Moser, D., Crespo-Facorro, B., Robinson, R.G., 2005. Alcohol Misuse and Mood Disorders Following Traumatic Brain Injury. *Arch. Gen. Psychiatry* 62, 742–749. <https://doi.org/10.1001/archpsyc.62.7.742>

Kalaiselvana, G., Dongre, A.R., Mahalakshmy, T., 2011. Epidemiology of injury in rural Pondicherry, India. *J. Inj. Violence Res.* 3, 62–67.
<https://doi.org/10.5249/jivr.v3i2.74>

Kessler, R.C., Üstün, T.B., 2004. The World Mental Health (WMH) Survey Initiative Version of the World Health Organization (WHO) Composite International Diagnostic Interview (CIDI). *Int. J. Methods Psychiatr. Res.* 13, 93–121.

Kilzieh, N., Rastam, S., Maziak, W., Ward, K.D., 2008. Comorbidity of depression with chronic diseases: a population-based study in Aleppo, Syria. *Int. J. Psychiatry Med.* 38, 169–184. <https://doi.org/10.2190/PM.38.2.d>

Kim, H.-C., Park, S.-G., Min, K.-B., Yoon, K.-J., 2008. Depressive symptoms and self-reported occupational injury in small and medium-sized companies. *Int. Arch. Occup. Environ. Health* 82, 715. <https://doi.org/10.1007/s00420-008-0365-6>

Kowal, P., Chatterji, S., Naidoo, N., Biritwum, R., Fan, W., Lopez Ridaura, R., Maximova, T., Arokiasamy, P., Phaswana-Mafuya, N., Williams, S., Snodgrass, J.J., Minicuci, N., D'Este, C., Peltzer, K., Boerma, J.T., SAGE Collaborators, 2012. Data

resource profile: the World Health Organization Study on global AGEing and adult health (SAGE). *Int. J. Epidemiol.* 41, 1639–1649. <https://doi.org/10.1093/ije/dys210>

Koyanagi, A., Garin, N., Olaya, B., Ayuso-Mateos, J.L., Chatterji, S., Leonardi, M., Koskinen, S., Tobiasz-Adamczyk, B., Haro, J.M., 2014. Chronic conditions and sleep problems among adults aged 50 years or over in nine countries: a multi-country study. *PloS One* 9, e114742. <https://doi.org/10.1371/journal.pone.0114742>

Koyanagi, A., Stickley, A., Garin, N., Miret, M., Ayuso-Mateos, J.L., Leonardi, M., Koskinen, S., Galas, A., Haro, J.M., 2015. The association between obesity and back pain in nine countries: a cross-sectional study. *BMC Public Health* 15, 123. <https://doi.org/10.1186/s12889-015-1362-9>

Koyanagi, A., Stickley, A., Haro, J.M., 2016. Psychotic symptoms and smoking in 44 countries. *Acta Psychiatr. Scand.* 133, 497–505. <https://doi.org/10.1111/acps.12566>

Kozela, M., Bobak, M., Besala, A., Micek, A., Kubinova, R., Malyutina, S., Denisova, D., Richards, M., Pikhart, H., Peasey, A., Marmot, M., Pająk, A., 2016. The association of depressive symptoms with cardiovascular and all-cause mortality in Central and Eastern Europe: Prospective results of the HAPIEE study. *Eur. J. Prev. Cardiol.* 23, 1839–1847. <https://doi.org/10.1177/2047487316649493>

Kudebong, M., Wurapa, F., Nonvignon, J., Norman, I., Awoonor-Williams, J.K., Aikins, M., 2011. Economic burden of motorcycle accidents in Northern Ghana. *Ghana Med. J.* 45, 135–142.

Leive, A., Xu, K., 2008. Coping with out-of-pocket health payments: empirical evidence from 15 African countries. *Bull. World Health Organ.* 86, 849–856. <https://doi.org/10.2471/BLT.07.049403>

Markon, K.E., Chmielewski, M., Miller, C.J., 2011. The reliability and validity of discrete and continuous measures of psychopathology: a quantitative review.

Psychol. Bull. 137, 856–879. <https://doi.org/10.1037/a0023678>

Martin, L., Byrnes, M., Bulsara, M.K., McGarry, S., Rea, S., Wood, F., 2017. Quality of life and posttraumatic growth after adult burn: A prospective, longitudinal study. *Burns* 43, 1400–1410. <https://doi.org/10.1016/j.burns.2017.06.004>

Mitchell, M.C., Burns, N.R., Dorstyn, D.S., 2008. Screening for depression and anxiety in spinal cord injury with DASS-21. *Spinal Cord* 46, 547–551. <https://doi.org/10.1038/sj.sc.3102154>

Morriss, W.W., Roques, C.J., 2018. Pain management in low- and middle-income countries. *BJA Educ.* 18, 265–270. <https://doi.org/10.1016/j.bjae.2018.05.006>

Niemi, M., Kiel, S., Allebeck, P., Hoan, L.T., 2016. Community-based intervention for depression management at the primary care level in Ha Nam Province, Vietnam: a cluster-randomised controlled trial. *Trop. Med. Int. Health* 21, 654–661. <https://doi.org/10.1111/tmi.12674>

Noh, J.-W., Kwon, Y.D., Park, J., Oh, I.-H., Kim, J., 2016. Relationship between Physical Disability and Depression by Gender: A Panel Regression Model. *PloS One* 11, e0166238. <https://doi.org/10.1371/journal.pone.0166238>

O'Donnell, M.L., Creamer, M., Pattison, P., Atkin, C., 2004. Psychiatric morbidity following injury. *Am. J. Psychiatry* 161, 507–514. <https://doi.org/10.1176/appi.ajp.161.3.507>

Paljärvi, T., Koskenvuo, M., Poikolainen, K., Kauhanen, J., Sillanmäki, L., Mäkelä, P., 2009. Binge drinking and depressive symptoms: a 5-year population-based cohort study. *Addiction* 104, 1168–1178. <https://doi.org/10.1111/j.1360-0443.2009.02577.x>

Patel, V., 2007. Mental health in low- and middle-income countries. *Br. Med. Bull.* 81–82, 81–96. <https://doi.org/10.1093/bmb/ldm010>

- Patten, S.B., Williams, J.V., Lavorato, D.H., Eliasziw, M., 2010. Major depression and injury risk. *Can. J. Psychiatry Rev. Can. Psychiatr.* 55, 313–318. <https://doi.org/10.1177/070674371005500507>
- Peele, P.B., Tollerud, D.J., 2005. Depression and occupational injury: results of a pilot investigation. *J. Occup. Environ. Med.* 47, 424–427.
- Prince, M., Acosta, D., Ferri, C.P., Guerra, M., Huang, Y., Rodriguez, J.J.L., Salas, A., Sosa, A.L., Williams, J.D., Dewey, M.E., Acosta, I., Jotheeswaran, A.T., Liu, Z., 2012. Dementia incidence and mortality in middle-income countries, and associations with indicators of cognitive reserve: a 10/66 Dementia Research Group population-based cohort study. *Lancet* 380, 50–58. [https://doi.org/10.1016/S0140-6736\(12\)60399-7](https://doi.org/10.1016/S0140-6736(12)60399-7)
- Rasmussen, A., Rosenfeld, B., Reeves, K., Keller, A.S., 2007. The effects of torture-related injuries on long-term psychological distress in a Punjabi Sikh sample. *J. Abnorm. Psychol.* 116, 734–740. <https://doi.org/10.1037/0021-843X.116.4.734>
- Rayner, L., Simpson, A., Matcham, F., Shetty, S., Lahoti, O., Groom, G., Hotopf, M., 2016. Mental disorder in limb reconstruction: Prevalence, associations and impact on work disability. *J. Psychosom. Res.* 89, 53–60. <https://doi.org/10.1016/j.jpsychores.2016.08.009>
- Razvodovsky, Y.E., 2012. Contribution of alcohol in accident related mortality in Belarus: a time series approach. *J. Inj. Violence Res.* 4, 58–64. <https://doi.org/10.5249/jivr.v4i2.100>
- Rivara, F.P., Mackenzie, E.J., Jurkovich, G.J., Nathens, A.B., Wang, J., Scharfstein, D.O., 2008. Prevalence of pain in patients 1 year after major trauma. *Arch. Surg. Chic.* 143, 282–287; discussion 288. <https://doi.org/10.1001/archsurg.2007.61>

- Routley, V., Ozanne-Smith, J., Li, D., Hu, X., Wang, P., Qin, Y., 2007. Pattern of seat belt wearing in Nanjing, China. *Inj. Prev.* 13, 388–393. <https://doi.org/10.1136/ip.2007.015701>
- Ryder, A.G., Yang, J., Zhu, X., Yao, S., Yi, J., Heine, S.J., Bagby, R.M., 2008. The cultural shaping of depression: somatic symptoms in China, psychological symptoms in North America? *J. Abnorm. Psychol.* 117, 300–313. <https://doi.org/10.1037/0021-843X.117.2.300>
- Salvetti, M. de G., Pimenta, C.A. de M., Braga, P.E., McGillion, M., 2013. Prevalence of fatigue and associated factors in chronic low back pain patients. *Rev. Lat. Am. Enfermagem* 21, 12–19. <https://doi.org/10.1590/S0104-11692013000700003>
- Sekaran, P., Vijayakumari, F., Hariharan, R., Zachariah, K., Joseph, S.E., Kumar, R.K.S., 2010. Community reintegration of spinal cord-injured patients in rural south India. *Spinal Cord* 48, 628–632. <https://doi.org/10.1038/sc.2010.6>
- Smith, P.D., Becker, K., Roberts, L., Walker, J., Szanton, S.L., 2016. Associations among pain, depression, and functional limitation in low-income, home-dwelling older adults: An analysis of baseline data from CAPABLE. *Geriatr. Nurs. N. Y.* N 37, 348–352. <https://doi.org/10.1016/j.gerinurse.2016.04.016>
- Solanki, H.K., Ahamed, F., Gupta, S.K., Nongkynrih, B., 2016. Road Transport in Urban India: Its Implications on Health. *Indian J. Community Med. Off. Publ. Indian Assoc. Prev. Soc. Med.* 41, 16–22. <https://doi.org/10.4103/0970-0218.170959>
- Stubbs, B., Koyanagi, A., Schuch, F.B., Firth, J., Rosenbaum, S., Veronese, N., Solmi, M., Mugisha, J., Vancampfort, D., 2016. Physical activity and depression: a large cross-sectional, population-based study across 36 low- and middle-income countries. *Acta Psychiatr. Scand.* 134, 546–556. <https://doi.org/10.1111/acps.12654>
- Stubbs, B., Vancampfort, D., Veronese, N., Kahl, K.G., Mitchell, A.J., Lin, P.-Y.,

- Tseng, P.-T., Mugisha, J., Solmi, M., Carvalho, A.F., Koyanagi, A., 2017a. Depression and physical health multimorbidity: primary data and country-wide meta-analysis of population data from 190 593 people across 43 low- and middle-income countries. *Psychol. Med.* 47, 2107–2117. <https://doi.org/10.1017/S0033291717000551>
- Stubbs, B., Vancampfort, D., Veronese, N., Thompson, T., Fornaro, M., Schofield, P., Solmi, M., Mugisha, J., Carvalho, A.F., Koyanagi, A., 2017b. Depression and pain: primary data and meta-analysis among 237 952 people across 47 low- and middle-income countries. *Psychol. Med.* 47, 2906–2917. <https://doi.org/10.1017/S0033291717001477>
- Tang, N.K.Y., McBeth, J., Jordan, K.P., Blagojevic-Bucknall, M., Croft, P., Wilkie, R., 2015. Impact of musculoskeletal pain on insomnia onset: a prospective cohort study. *Rheumatol. Oxf. Engl.* 54, 248–256. <https://doi.org/10.1093/rheumatology/keu283>
- Tao, M., Li, Yihan, Xie, D., Wang, Zhiyang, Qiu, J., Wu, W., Sun, J., Wang, Zhoubing, Tao, D., Zhao, H., Tian, T., Zhang, Jingxuan, Gao, C., Niu, Q., Li, Q., Liu, S., Liu, J., Zhang, Y., He, Q., Rong, H., Gan, Z., Li, J., Chen, X., Pan, J., Li, Yi, Cui, Y., Han, W., Ma, H., Xie, S., Jin, G., Li, L., Zhang, R., Tan, Q., Zhang, Jun, Guan, J., Shi, S., Chen, Y., Kendler, K.S., Flint, J., Gao, J., 2011. Examining the relationship between lifetime stressful life events and the onset of major depression in Chinese women. *J. Affect. Disord.* 135, 95–99. <https://doi.org/10.1016/j.jad.2011.06.054>
- Tomlinson, M., Grimsrud, A.T., Stein, D.J., Williams, D.R., Myer, L., 2009. The epidemiology of major depression in South Africa: Results from the South African Stress and Health study. *South Afr. Med. J. Suid-Afr. Tydskr. Vir Geneeskde.* 99, 367–373.
- Tripathy, P., Nair, N., Barnett, S., Mahapatra, R., Borghi, J., Rath, Shibnand, Rath,

- Suchitra, Gope, R., Mahto, D., Sinha, R., Lakshminarayana, R., Patel, V., Pagel, C., Prost, A., Costello, A., 2010. Effect of a participatory intervention with women's groups on birth outcomes and maternal depression in Jharkhand and Orissa, India: a cluster-randomised controlled trial. *Lancet Lond. Engl.* 375, 1182–1192. [https://doi.org/10.1016/S0140-6736\(09\)62042-0](https://doi.org/10.1016/S0140-6736(09)62042-0)
- Turner, R.J., Noh, S., 1988. Physical Disability and Depression: A Longitudinal Analysis. *J. Health Soc. Behav.* 29, 23–37. <https://doi.org/10.2307/2137178>
- Ukpong, D.I., Ugboko, V.I., Ndukwe, K.C., Gbolahan, O., 2007. Psychological complications of maxillofacial trauma: preliminary findings from a Nigerian university teaching hospital. *J. Oral Maxillofac. Surg. Off. J. Am. Assoc. Oral Maxillofac. Surg.* 65, 891–894. <https://doi.org/10.1016/j.joms.2006.06.292>
- Vancampfort, D., Koyanagi, A., Ward, P.B., Veronese, N., Carvalho, A.F., Solmi, M., Mugisha, J., Rosenbaum, S., De Hert, M., Stubbs, B., 2017. Perceived Stress and Its Relationship With Chronic Medical Conditions and Multimorbidity Among 229,293 Community-Dwelling Adults in 44 Low- and Middle-Income Countries. *Am. J. Epidemiol.* 186, 979–989. <https://doi.org/10.1093/aje/kwx159>
- Wadhvaniya, S., Gupta, S., Mitra, S., Tetali, S., Josyula, L.K., Gururaj, G., Hyder, A.A., 2017. A comparison of observed and self-reported helmet use and associated factors among motorcyclists in Hyderabad city, India. *Public Health, Supplement: Global Road Safety: Monitoring Risks and Evaluating Programs* 144, S62–S69. <https://doi.org/10.1016/j.puhe.2016.11.025>
- Watt, K., Purdie, D.M., Roche, A.M., McClure, R.J., 2004. Risk of injury from acute alcohol consumption and the influence of confounders. *Addict. Abingdon Engl.* 99, 1262–1273. <https://doi.org/10.1111/j.1360-0443.2004.00823.x>
- Wesson, H., Boikhutso, N., Bachani, A., Hofman, K., Hyder, A., 2014. The cost of

injury and trauma care in low- and middle-income countries: a review of economic evidence. *Health Policy Plan.* 29, 795–808. <https://doi.org/10.1093/heapol/czt064>

Wiseman, T.A., Curtis, K., Lam, M., Foster, K., 2015. Incidence of depression, anxiety and stress following traumatic injury: a longitudinal study. *Scand. J. Trauma Resusc. Emerg. Med.* 23, 29. <https://doi.org/10.1186/s13049-015-0109-z>

Wong, K.O., Hunter Rowe, B., Douwes, J., Senthilselvan, A., 2013. Asthma and wheezing are associated with depression and anxiety in adults: an analysis from 54 countries. *Pulm. Med.* 2013, 929028. <https://doi.org/10.1155/2013/929028>

World Health Organization, 2017. Depression and Other Common Mental Disorders: Global Health Estimates. World Health Organization, Geneva.

World Health Organization, 2014. Injuries and violence: the facts 2014.

Wu, H., Zhang, F., Cheng, W., Lin, Y., Wang, Q., 2017. Factors Related to Acute Anxiety and Depression in Inpatients with Accidental Orthopedic Injuries. *Shanghai Arch. Psychiatry* 29, 77–84. <https://doi.org/10.11919/j.issn.1002-0829.216070>

Xiao, Y., Zhao, K., Ma, Z.-X., Li, X., Qiu, Y.-P., 2017. Integrated medical rehabilitation delivery in China. *Chronic Dis. Transl. Med.* 3, 75–81. <https://doi.org/10.1016/j.cdtm.2017.02.003>

Yang, L., Zhao, Y., Wang, Y., Liu, L., Zhang, X., Li, B., Cui, R., 2015. The Effects of Psychological Stress on Depression. *Curr. Neuropharmacol.* 13, 494–504. <https://doi.org/10.2174/1570159X1304150831150507>

Yatham, S., Sivathasan, S., Yoon, R., da Silva, T.L., Ravindran, A.V., 2017. Depression, anxiety, and post-traumatic stress disorder among youth in low and middle income countries: A review of prevalence and treatment interventions. *Asian J. Psychiatry*. <https://doi.org/10.1016/j.ajp.2017.10.029>

Yokoyama, E., Kaneita, Y., Saito, Y., Uchiyama, M., Matsuzaki, Y., Tamaki, T.,

Munezawa, T., Ohida, T., 2010. Association between Depression and Insomnia Subtypes: A Longitudinal Study on the Elderly in Japan. *Sleep* 33, 1693–1702. <https://doi.org/10.1093/sleep/33.12.1693>

Zeng, Q., Xu, Y., Chun Wang, W., 2013. Quality of life in outpatients with depression in China. *J. Affect. Disord.* 150, 513–521. <https://doi.org/10.1016/j.jad.2013.04.052>




Figure 1 Prevalence of depression among those with and without any injury in the past 12 months

Data are based on weighted prevalence.

Any injury included both road traffic accidents and bodily injury for other causes.




Figure 2 Country-wise association between any injury (exposure) and depression (outcome) in the past 12 months estimated with multivariable logistic regression

Abbreviation: OR Odds ratio; CI Confidence interval.

Models are adjusted for sex, age, wealth, education, setting, and alcohol consumption.

Any injury included both road traffic accidents and bodily injury for other causes.

The overall estimate was calculated by random-effects meta-analysis.

Table 1 Sample characteristics (overall and by presence of any injury)

| Characteristic | Category | Overall | Any injury ^a | | P-value ^b |
|---------------------|-----------|-------------|-------------------------|-------------|----------------------|
| | | | No | Yes | |
| Sex | Female | 50.1 | 50.8 | 42.5 | <0.001 |
| | Male | 49.9 | 49.2 | 57.5 | |
| Age | Mean (SD) | 43.8 (14.4) | 43.8 (14.4) | 43.5 (13.8) | 0.512 |
| Wealth | Poorest | 14.9 | 14.6 | 19.5 | <0.001 |
| | Poorer | 17.8 | 17.7 | 19.7 | |
| | Middle | 18.8 | 18.5 | 22.5 | |
| | Richer | 21.1 | 21.3 | 19.7 | |
| | Richest | 27.3 | 28.0 | 18.8 | |
| Education | ≤Primary | 43.0 | 42.2 | 52.4 | <0.001 |
| | Secondary | 46.6 | 47.1 | 39.9 | |
| | ≥Tertiary | 10.4 | 10.6 | 7.7 | |
| Setting | Rural | 55.8 | 54.9 | 66.3 | <0.001 |
| | Urban | 44.2 | 45.1 | 33.7 | |
| Alcohol consumption | Never | 68.3 | 68.7 | 63.4 | 0.007 |
| | Non-heavy | 26.4 | 26.2 | 27.9 | |
| | Heavy | 5.3 | 5.0 | 8.7 | |

Abbreviation: SD Standard deviation

Data are % unless otherwise stated.

Data are based on weighted estimates.

^a Any injury included road traffic accidents and bodily injury for other causes occurring in the past 12 months.

^b P-value was obtained by Chi-squared test and Student's *t*-test for categorical and continuous variables, respectively.

Table 2 Sample characteristics by country

| Characteristic | Category | China | Ghana | India | Mexico | Russia | South Africa |
|------------------------|------------------|-------------|-------------|-------------|-------------|-------------|--------------|
| Sex | Female | 49.13 | 49.99 | 49.12 | 52.00 | 55.01 | 52.77 |
| | Male | 50.87 | 50.01 | 50.88 | 48.00 | 44.99 | 47.23 |
| Age | Mean (SD) | 45.6 (12.8) | 44.4 (14.3) | 41.1 (14.7) | 42.7 (14.7) | 47.3 (16.9) | 41.8 (14.8) |
| Education | ≤Primary | 37.53 | 63.44 | 61.35 | 51.69 | 3.22 | 37.55 |
| | Secondary | 53.15 | 32.04 | 30.28 | 35.85 | 75.56 | 54.45 |
| | ≥Tertiary | 9.32 | 4.52 | 8.38 | 12.46 | 21.21 | 8.00 |
| Setting | Rural | 51.52 | 54.23 | 74.52 | 22.24 | 18.46 | 30.66 |
| | Urban | 48.48 | 45.77 | 25.48 | 77.76 | 81.54 | 69.34 |
| Alcohol consumption | Never | 66.51 | 44.76 | 84.03 | 45.38 | 19.17 | 76.52 |
| | Non-heavy | 24.89 | 52.53 | 15.31 | 45.56 | 70.82 | 15.36 |
| | Heavy | 8.61 | 2.72 | 0.67 | 9.06 | 10.01 | 8.12 |
| Depression | Presence | 0.83 | 4.19 | 7.89 | 5.66 | 2.54 | 3.77 |
| Injury | Any injury | 6.06 | 7.22 | 10.73 | 6.38 | 6.14 | 3.18 |
| | Traffic accident | 1.97 | 1.30 | 2.93 | 3.74 | 2.42 | 2.42 |
| Injury with disability | Any injury | 0.83 | 1.09 | 2.42 | 3.14 | 0.06 | 0.84 |
| | Traffic accident | 0.26 | 0.10 | 0.58 | 1.66 | 0.03 | 0.71 |

Abbreviation: SD Standard deviation

Data are % unless otherwise stated.

Data are based on weighted estimates.

Table 3 Association of any injury, injury due to traffic accident, and injury due to other causes with depression estimated by multivariable logistic regression

| Characteristic | Category | Any injury | | Injury due to traffic accident | | Injury due to other causes | |
|---------------------|---------------------------|------------|-------------|--------------------------------|--------------|----------------------------|-------------|
| | | OR | 95%CI | OR | 95%CI | OR | 95%CI |
| Injury | No injury | 1.00 | | 1.00 | | 1.00 | |
| | Injury without disability | 1.72** | [1.18,2.50] | 0.65 | [0.35,1.22] | 2.05** | [1.33,3.14] |
| | Injury with disability | 3.81*** | [2.16,6.73] | 6.12* | [1.42,26.40] | 3.20*** | [2.02,5.05] |
| Sex | Female | 1.00 | | 1.00 | | 1.00 | |
| | Male | 0.67** | [0.51,0.87] | 0.67** | [0.52,0.86] | 0.67** | [0.52,0.87] |
| Age (years) | Per one year increase | 1.03*** | [1.02,1.03] | 1.03*** | [1.02,1.03] | 1.03*** | [1.02,1.03] |
| Wealth | Poorest | 1.00 | | 1.00 | | 1.00 | |
| | Poorer | 0.91 | [0.68,1.22] | 0.91 | [0.68,1.22] | 0.91 | [0.68,1.22] |
| | Middle | 0.75 | [0.54,1.06] | 0.75 | [0.54,1.05] | 0.76 | [0.54,1.06] |
| | Richer | 0.54*** | [0.39,0.75] | 0.53*** | [0.38,0.74] | 0.54*** | [0.39,0.75] |
| | Richest | 0.53** | [0.35,0.78] | 0.52** | [0.35,0.77] | 0.52** | [0.35,0.78] |
| Education | ≤Primary | 1.00 | | 1.00 | | 1.00 | |
| | Secondary | 1.10 | [0.80,1.51] | 1.09 | [0.79,1.49] | 1.10 | [0.80,1.51] |
| | ≥Tertiary | 0.80 | [0.41,1.58] | 0.79 | [0.40,1.56] | 0.80 | [0.40,1.57] |
| Setting | Rural | 1.00 | | 1.00 | | 1.00 | |
| | Urban | 1.22 | [0.90,1.64] | 1.20 | [0.89,1.62] | 1.21 | [0.89,1.63] |
| Alcohol consumption | Never | 1.00 | [1.00,1.00] | 1.00 | [1.00,1.00] | 1.00 | [1.00,1.00] |
| | Non-heavy | 0.87 | [0.60,1.26] | 0.88 | [0.61,1.28] | 0.87 | [0.60,1.26] |
| | Heavy | 0.72 | [0.27,1.89] | 0.75 | [0.28,1.96] | 0.70 | [0.26,1.84] |

Abbreviation: OR Odds ratio; CI Confidence interval.

Models are adjusted for all variables in the Table and country.

* p<0.05, ** p<0.01, *** p<0.001.