



HAL
open science

Changes in marine phytoplankton diversity: Assessment under the Marine Strategy Framework Directive

I. Rombouts, N. Simon, A. Aubert, T. Cariou, E. Feunteun, L. Guérin, M. Hoebeke, A. Mcquatters-Gollop, F. Rigaut-Jalabert, Luis Felipe Artigas

► To cite this version:

I. Rombouts, N. Simon, A. Aubert, T. Cariou, E. Feunteun, et al.. Changes in marine phytoplankton diversity: Assessment under the Marine Strategy Framework Directive. *Ecological Indicators*, 2019, 102, pp.265-277. 10.1016/j.ecolind.2019.02.009 . hal-03485708

HAL Id: hal-03485708

<https://hal.science/hal-03485708>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Changes in marine phytoplankton diversity: assessment under the Marine Strategy**

2 **Framework Directive**

3 Rombouts ^{1,2,3,4*}, I., Simon¹, N., Aubert², A., Cariou³, T., Feunteun², E., Guérin⁵, L., Hoebeke³,
4 M., McQuatters-Gollop⁶, A., Rigaut-Jalabert³, F., Artigas⁴, L.F.

5

6 ¹CNRS, Sorbonne Université, Station Biologique de Roscoff (SBR), UMR 7144,

7 Ecology of Marine Plankton, place Georges-Teissier, 29688 Roscoff cedex, France.

8

9 ² MNHN, UMR BOREA (MNHN, Sorbonne Universités, CNRS, UniCaen, IRD, Univ

10 Antilles) Station Marine de Dinard, Centre de Recherche et d'Enseignement Sur les Systèmes

11 Côtiers (CRESCO), 38 Rue du Port Blanc, 35800 Dinard, France.

12

13 ³CNRS, Sorbonne Université, Station Biologique de Roscoff (SBR), FR2424,

14 place Georges-Teissier, 29688 Roscoff cedex, France

15

16 ⁴Laboratoire d'Océanologie et de Géosciences (LOG), UMR 8187 CNRS-Université du

17 Littoral Côte d'Opale-Université de Lille, 62930 Wimereux, France.

18

19 ⁵UMS PatriNat (AFB, MNHN, CNRS) Station Marine de Dinard, Centre de Recherche et

20 d'Enseignement Sur les Systèmes Côtiers (CRESCO), 38 Rue du Port Blanc, 35800 Dinard,

21 France.

22

23 ⁶Marine and Conservation Policy Research Group, University of Plymouth, Drake Circus,
24 Plymouth, UK, PL4 8AA

25

26 ***Corresponding author:**

27 Dr Isabelle Rombouts

28 Laboratoire d'Océanologie et de Géosciences (LOG - CNRS - ULCO - U Lille)

29 Maison de la Recherche en Environnement Naturel (MREN)

30 32, Avenue du Maréchal Foch

31 62930 Wimereux

32 France

33 E-mail: isabelle.rombouts@univ-lille1.fr

34

35 **Number of words:**

36 Abstract: 365

37 Main text: 7235

38 Manuscript total: 11065

39 **Number of references: 97**

40 **Number of figures: 5 and tables: 3**

41 **Supplementary material: Number of figures: 5 and tables: 2**

42

43 **Keywords**

44 Community composition – Good Environmental Status – Indicators – Marine Policy – MSFD

45 - OSPAR – Pelagic Habitat – Plankton

46 **Abstract**

47 The Marine Strategy Framework Directive requires EU Member States to assess the Good
48 Environmental Status (GES) of their marine waters in a coherent and strategic manner. For
49 the regional assessment of biodiversity, the OSPAR Intersessional Coordination Group of
50 Biodiversity Assessment and Monitoring (ICG-COBAM) provides substantial advice.
51 Through expert working groups, phytoplankton indicators are currently being developed to
52 measure the state and the change in pelagic diversity, to quantify food web dynamics and to
53 measure the extent of eutrophication impacts. We developed a multi-metric indicator that is
54 compliant with the common OSPAR indicator “Changes in plankton diversity” (PH3). The
55 aim was to describe the structure of the phytoplankton community (alpha diversity) and to
56 detect significant temporal changes (beta diversity) to evaluate the health of pelagic habitats.
57 In this pilot study, we used three coastal time-series in the Western Channel and the north of
58 the Bay of Biscay (North Atlantic, France) to test the efficiency and the performance of
59 several existing diversity indices. We validated two alpha diversity indices, namely the
60 Menhinick Index (D) and the Hulburt Index (δ), based on their complementary ecological
61 information, their strong relationship with habitat characteristics, and their relative ease of
62 interpretation for stakeholders. Temporal shifts or rate of change in community structure
63 were detected by the Local Contributions to Beta Diversity index (LCBD; a beta diversity
64 measure). For the years where significantly high LCBD values were found, the Importance
65 Value Index (IVI) was calculated to potentially identify the taxa (genus) responsible for the
66 “unusual” community structure. For example, at the Ouest Loscolo site in 2008, an elevated
67 LCBD (0.45) coincided with a high dominance value (Hulburt’s Index) caused by the
68 occurrence of a monospecific bloom of *Leptocylindrus* spp. (IVI = 73%) in July (2.2×10^6 cells L⁻¹)

69 and October (8×10^6 cells L^{-1}). In this way, PH3 informs on different aspects of phytoplankton
70 diversity from a community to a genus level. At the current stage of development, however,
71 PH3 acts as a “surveillance” rather than an operational indicator since the relationship to
72 GES is not directly tracked. In the future, by additional testing of PH3 and extending the
73 geographical scope, the robustness of the assessment could be further determined across the
74 OSPAR Maritime Area.

75

76 Introduction

77 The Marine Strategy Framework Directive (MSFD) requires that European Member States
78 that share a marine region or sub-region cooperate when developing their marine strategies
79 (CEC 2008). In this respect, Regional Sea Conventions, like OSPAR (Convention for the
80 Protection of the Marine Environment of the North-East Atlantic), take a key role as a
81 platform for EU Member States to coordinate their approaches in implementing the MSFD at
82 a regional scale. For the ‘biodiversity’ descriptors of the Directive (i.e. D1 Biodiversity, D2
83 Non-indigenous species, D4 Food webs and D6 Seafloor integrity), the OSPAR Intersessional
84 Coordination Group of Biodiversity Assessment and Monitoring (ICG-COBAM) provides
85 substantial regional advices for the North East Atlantic, on the basis of its intersessional
86 work and its seven dedicated working groups each covering an ecosystem component
87 (marine mammals, seabirds, fish and cephalopods, benthic habitats, pelagic habitats, non-
88 indigenous species and food webs). The main tasks of the working groups are to identify a
89 set of common indicators and to coordinate the development of these indicators for their use
90 in regional assessments. To date, common indicators based on plankton communities have

91 been adopted by OSPAR to assess Good Environmental Status (GES) of pelagic habitats at
92 the regional scale of the North East Atlantic ([https://oap.ospar.org/en/ospar-](https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/)
93 [assessments/intermediate-assessment-2017/biodiversity-status/habitats/](https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/)).

94

95 Coastal ecosystems face increasing human disturbances such as pollution and/or
96 eutrophication (i.e. excessive nutrients or organic enrichments) that can drive marked
97 changes in the plankton community dynamics and thus in its structural attributes, such as
98 diversity, dominance or size structure. Phytoplankton, for example, show rapid responses to
99 altered nutrient levels through changes in biomass and composition (Reynolds, 2006).

100 Whereas the use of phytoplankton biomass for water quality assessment has a long history
101 (Paształeniec, 2016), the evaluation of community composition has gained a more recent
102 interest through the implementation of the Water Framework Directive (WFD) (Devlin et al.,
103 2009; Uusitalo et al., 2013). In the WFD, three metrics, namely ‘phytoplankton abundance’,
104 ‘phytoplankton biomass’ and ‘phytoplankton taxonomic composition’, are part of the
105 biological quality elements (BQEs), i.e. organism groups which integrate the effects of
106 various stressors such as nutrient enrichment, acidification, and, to some extent, hypoxia or
107 habitat degradation (Lyche-Solheim et al., 2013). In contrast to measurements for chlorophyll
108 *a* as a proxy for biomass, the assessment of the taxonomic composition of the phytoplankton
109 assemblage could provide information about the whole community, including the
110 importance of the different size-groups such as the pico- and nano-phytoplankton
111 (Domingues et al., 2008).

112

113 Diversity indices summarise the abundance data for multiple species in an assemblage into a
114 single number to describe the state of the community (Kwak and Peterson, 2007). A plethora

115 of indices exist in the scientific literature that focus on different aspects of biodiversity
116 (richness, dominance, evenness) and are usually weighted in different ways, for example, the
117 Simpson's index is more weighted on dominant species compared to the Shannon index
118 (Magurran, 1988). The choice of the most appropriate indices depends on the type of
119 assemblage considered, the objectives of the study and the data availability (e.g. Chiarucci et
120 al., 2011; Morris et al., 2014). In terms of community structure, many natural biotic
121 communities, such as phytoplankton, are characterized by the presence of a few common
122 species with high abundances and many rare species (Wilhm and Dorris, 1968). Over time,
123 abundances of phytoplankton can vary by several orders of magnitude at the seasonal,
124 interannual and interdecadal time scales as a result of variations in natural environmental
125 conditions and/or from anthropogenic pressures (e.g. Zingone et al., 2010; Muñiz et al., 2018).
126 On a seasonal basis, phytoplankton exhibit a distinct succession in species composition, i.e.
127 an ordered sequence of substitutions of species (Margalef, 1978; Reynolds, 2006), and these
128 variations are sometimes even more significant than inter-annual trends in phytoplankton
129 community structure. The causes of succession are complex and have not been totally
130 elucidated (Sommer et al., 2012). Succession can depend on species-interactions and, more
131 importantly, the reactivity to favourable environmental conditions throughout the year, such
132 as seasonal changes in temperature, water column mixing/stratification, nutrient loadings
133 and light availability (Chalar, 2009). Other processes act on time periods of days to weeks,
134 like meteorological (wind, rain and cloudiness) and hydrological events
135 (upwelling/downwelling events). Finally, marked changes in the relative abundances of
136 species can also be a result of environmental perturbations such as pollution or
137 eutrophication (Bužančić et al., 2016; Domingues et al., 2017). In these cases, an increase in

138 dominance occurs because only a subset of species can actively benefit from the new
139 conditions (Ben Othman et al., 2018; Coclet et al., 2018).

140

141 Biodiversity measures can be useful for conservation practice and management purposes
142 (Chiarucci et al., 2011; Scheiner et al., 2017). In this respect, "species richness" was identified
143 as an Essential Biodiversity Variable (EBV), a measurement required for studying, reporting
144 and managing biodiversity change (Pereira et al., 2013; Kissling et al., 2018). Whilst
145 taxonomic richness is a useful biodiversity metric, its applicability to assess the state of
146 pelagic habitats in water quality assessment is debatable and to date no consensus has been
147 achieved about which indices are more appropriate and informative for assessing the state
148 and change in phytoplankton communities. One of the main problems is that the response of
149 phytoplankton communities to anthropogenic pressures is often non-linear, making clear
150 state-pressure relationships difficult to identify (Garmendia et al., 2013; Ninčević-Gladan et
151 al., 2015). As an example, Shannon and Simpson indices are widely used in descriptive
152 studies to quantify community diversity but were found inappropriate as tools for water
153 quality assessment due to their erratic behaviour along a eutrophication gradient (Spatharis
154 et al., 2011). To increase the robustness of assessment using diversity indices, several studies
155 have proposed to modify already existing diversity metrics, for example the Shannon95
156 (Uusitalo et al., 2013), and/or the use of composite indices (Spatharis and Tsirtsis, 2010;
157 Vadrucci et al., 2013; Laplace-Treytoure and Feret, 2016), to date mainly developed for
158 freshwater systems and transitional waters. Whilst these studies agree on the use of
159 phytoplankton community structure as an essential component for water quality assessment
160 (Devlin et al., 2009; Facca et al., 2014), further work is needed in this respect (Caroppo et al.,
161 2013; Garmendia et al., 2013; Varkitzi et al., 2018).

162

163 Within the OSPAR Regional Sea Convention, marine phytoplankton and zooplankton
164 community indicators are currently under development to assess the Environmental Status
165 of Pelagic Habitats (OSPAR 2017a). Pelagic Habitat indicator 1 (PH1) "Changes in
166 phytoplankton and zooplankton communities" uses the relative changes in abundances of
167 lifeform pairs based on functional traits to indicate ecological change (Tett et al., 2008;
168 McQuatters-Gollop et al., 2015; OSPAR, 2017b). For example, in the pairing of diatoms and
169 dinoflagellates, the dominance of the latter could indicate eutrophication resulting in less
170 desirable food webs. Pelagic Habitat indicator 2 (PH2) "Changes in Phytoplankton Biomass
171 and Zooplankton Abundance" provides an indication of deviations in total biomass or
172 abundance of plankton from the assumed natural variability in time-series (OSPAR, 2017c).
173 Finally, Pelagic Habitat indicator 3 (PH3) identifies changes in the community structure
174 using taxonomic diversity indices (OSPAR, 2017d). These three common indicators consider
175 plankton communities at different organizational levels: PH2 at the broadest organizational
176 level since it considers total phytoplankton biomass and total copepod abundance, PH1 at an
177 intermediate level since it considers lifeform pairs, and PH3 at the finest level of
178 organization, if possible down to the species level.

179 This paper summarises the development of the OSPAR common indicator "Changes in
180 plankton diversity" (PH3) for phytoplankton communities. The aim of PH3 is to characterise
181 the phytoplankton community structure and to detect potential temporal shifts, preferably in
182 relation to the environment. Frequently used diversity indices, mainly developed in the
183 context of the Water Framework Directive, were preselected. Microphytoplankton counts
184 obtained from three coastal time-series in the Western Channel and the north of the Bay of
185 Biscay (fig. 1) were used here to test the efficiency and the performance of several diversity

186 indices for assessing GES of pelagic habitats under the MFSD. More specifically, we tested
187 these diversity indices for their ecological relevance, mathematical consistency and link to
188 marine hydrological factors.

189

190 Materials and methods

191 1. Phytoplankton and environmental datasets

192 Microscopic counts of phytoplankton data from the Western Channel and the north of the
193 Bay of Biscay, France, were collated from two sources, namely RESOMAR-Pelagos (Pelagic
194 database of the Réseau National des Stations et Observatoires Marins;
195 <http://resomar.cnrs.fr/Base-de-donnee-Pelagos>) and REPHY (Réseau d'Observation et de
196 Surveillance du Phytoplancton et des
197 Phycotoxines;http://envlit.ifremer.fr/surveillance/phytoplancton_phycotoxines/presentation).

198 The REPHY is implemented and managed by the French Research Institute for the
199 Exploitation of the Sea (IFREMER). The database of RESOMAR-Pelagos hosts plankton data
200 collected from most of the French coastal marine stations and observatories. From the
201 RESOMAR-Pelagos database, we filtered for stations where samples were collected and
202 analysed using consistent methodology, were sampled at a minimum monthly frequency,
203 which contained minimal gaps in the sampling, and which simultaneously sampled
204 nutrients and hydrological factors. This selection resulted in the station of SOMLIT-Astan
205 (2007-2013, fig. 1), a coastal long-term monitoring station situated 4.6 km from the coast that
206 is characterized by permanently mixed waters with limited continental influence. Twice a
207 month, seawater samples are collected at 1 m depth using a 5 liters Niskin bottle for

208 phytoplankton analysis. Samples are fixed with acid Lugol's iodine solution and then stored
209 according to the methods described in Sournia (1978). Cell counts are made under an
210 inverted light microscope at 200-400x magnification. Further details on phytoplankton
211 quantification and identification protocols for SOMLIT-Astan can be found in Guilloux et al.
212 (2013). Environmental data from the site are collected by the Station Biologique de Roscoff
213 and hosted by the SOMLIT (Service d'Observation en Milieu LITtoral, INSU-CNRS)
214 database; they were retrieved from their online platform (<http://somlit.epoc.u->
215 [bordeaux1.fr/fr/](http://somalit.fr/fr/)). Data on salinity (psu), temperature (°C), inorganic nutrients (ammonia,
216 nitrate, nitrites, silicate, phosphates; in $\mu\text{mol L}^{-1}$) and oxygen (ml L^{-1}) were used in the
217 analysis.

218 In the Bay of Biscay, data from two REPHY sites, Ouest Loscolo and Le Croisic, were made
219 available for analyses (Catherine Belin, pers. comm.). These sites are shallow, meso- to
220 macrotidal, with a moderate wave exposure at 2.9 km from the coast for the Ouest Loscolo
221 station and 0.2 km from the coast for Le Croisic station. They are both under the influence of
222 riverine output, namely from the Loscolo and the Loire River. Water samples are collected
223 on a bi-monthly basis at the surface in order to determine phytoplankton cell abundance and
224 taxonomic composition. Phytoplankton samples are fixed with Lugol's solution (neutral or
225 acidic) and counted according to the Utermöhl method (Utermöhl, 1958). Further details
226 about sampling and processing of phytoplankton and physico-chemical parameters are
227 available in the literature (Neaud-Masson, 2015). The level of taxonomic identification
228 depends on the analytical method used and the experience of the phytoplankton analyst.
229 Changes in the taxonomic analyst may lead to heterogeneous data regarding taxonomic
230 classification and hence to a misinterpretation of phytoplankton time-series (Hernández-
231 Fariñas et al., 2013); this is true of many multidecadal datasets. Consequently, although

232 phytoplankton data in SOMLIT-Astan has been collected from the year 2000 onwards, only
233 the period 2007-2014 was considered for analysis since the same two operators worked
234 closely for the analyses of the samples during this time-period. Across datasets, most taxa
235 were identified to the species level but for consistency and again to reduce bias from
236 misidentification, abundance data (expressed as number of cells per liter) of the taxonomic
237 units were grouped to the genus level and pooled monthly. If the identification was at a
238 lower taxonomic level (Class, Phylum, as is the case for the smaller species), then these were
239 also taken into account but cells that were classified as “non-identified” were not used in
240 the analysis.

241

242 2. Data analysis

243 To select the most appropriate indices for the assessment of GES for pelagic habitats,
244 diversity indices were tested on the three sites in a range of simple and multivariate
245 analyses. After pre-selecting diversity indices from the literature, we have adopted some
246 criteria that biodiversity measures should satisfy for their use in quality assessment (van
247 Strien et al., 2012; Buckland et al., 2005). The final indicator should (1) provide ecological
248 information on the state condition of phytoplankton communities using several aspects of
249 biodiversity: richness, dominance, and evenness; and detect significant temporal changes in
250 the structure of the phytoplankton community (2) be mathematically consistent, (3) have a
251 link with environmental conditions.

252

253 2.1. Selection of diversity indices for the quantification of alpha diversity

254 2.1.1. Ecological relevance

255 In terms of ecological information, three aspects of diversity indices, i.e. the number of taxa,
256 their overall abundance and their evenness in the community, are of primary interest to
257 describe community structure and change, and have received an increased interest in
258 environmental management, especially in combination with each other (Buckland et al.,
259 2011). The aim was to select an index from each group so as to describe different aspects of a
260 phytoplankton community. Monthly and annual means in diversity indices were then
261 calculated for the three time-series so as to identify seasonal and annual trends in
262 community structure in terms of abundance of taxa.

263

264 2.1.1.1. Indices based on richness (number of taxa)

265 In phytoplankton studies, the most commonly used indices to describe the number of taxa in
266 the community includes species richness (S), the Margalef (d) Index and the Menhinick (D)
267 Index (Varkitzi et al., 2018). The latter index, in particular, has been found suitable as an
268 indicator of eutrophication in transitional (Facca et al., 2014) and coastal waters (Spatharis
269 and Tsirtsis, 2010; Buzançıç et al., 2016). The Menhinick index (D ; Whittaker, 1977) is a
270 measure of taxonomic richness where S represents the number of taxa, and N , the number of
271 individuals.

$$272 \quad D = \frac{S}{\sqrt{N}} \quad (1)$$

273

274 Whilst species richness (S) is the simplest and most straightforward index to calculate, this
275 estimate is strongly influenced by the sampling process (Peet, 1974; Rodriguez-Samos et al.,
276 2014). To investigate the effect of sampling effort on our estimates of richness, the cumulative
277 number of species as a function of the consecutive number of samples in time, were drawn.

278

279 2.1.1.2. Indices based on dominance and evenness (relative abundance)

280 As mentioned previously, phytoplankton communities are characterized by complex
281 dynamics with a strong seasonal cycle. Hence, indices that provide information on the
282 temporary dominance of species are of particular interest for the development of the
283 indicator, PH3, described here. For this purpose, diversity measures that include a richness
284 and an evenness component were used to express a relative concentration of dominance. In
285 this respect, the Shannon-Wiener and the Simpson's index are frequently used for describing
286 diversity in ecological assessment (Heip, 1998; Kabuta and Duijts, 2000). Additionally,
287 another dominance measure, the Hulburt index (δ ; Hulburt, 1963) has been developed to
288 describe phytoplankton communities in particular and was recently proposed as a suitable
289 indicator of eutrophication in the context of the WFD (Facca et al., 2014). Since this index is
290 expressed as a percentage, it is relatively easy to interpret.

291
$$\delta = 100 (n_1 + n_2) / N \quad (2)$$

292

293 where n_1 is the abundance of the dominant genus; n_2 is the abundance of the second most
294 abundant genus; and N is the total abundance.

295 Classical measures such as Shannon and Simpson's are based on species proportions and fail
296 to measure changes in abundance if all species in a community are declining at the same rate
297 (Buckland et al., 2011). To overcome this issue, the geometric mean index G_j , for example,
298 quantifies the average trend in relative abundance across species in the community
299 (Buckland et al., 2011). Finally, evenness indices express the equitability of species

300 abundance in the sample or the community (Washington, 1984). Here, we applied the
301 Pielou's index (J' ; Pielou, 1975).

302

303 2.1.2. Mathematical consistency

304 Within each index group, however, indices can be mathematically related since they are
305 either using common metrics and/or are derived from similar equations. With these
306 potentially competing indices, it is important to examine their mathematical convergence so
307 as to reduce redundancy in the information and to select only an optimal subset of indices
308 (Lyashevskaya and Farnsworth, 2012; van Strien et al., 2012; Bandeira et al., 2013). To do so,
309 simple statistical correlations (Bravais-Pearson) between all selected diversity indices (based
310 on monthly abundances) were calculated for each sampling site separately to investigate the
311 mathematical redundancies within each group.

312

313 2.1.3. Link with environmental conditions

314 Biodiversity metrics that respond differently to environmental factors can be considered
315 complementary (Gascon et al., 2009; Gallardo et al., 2011). Hence, we investigated to what
316 extent the selected biodiversity measures reflected changes in the environmental conditions
317 and if certain indices are interrelated.

318 A standardized Principal Components Analysis (PCA; Jolliffe, 1986) was applied to the
319 potential environmental correlates of phytoplankton diversity to determine: (1) the
320 environmental variables that explained the largest variation in the data set, (2) the
321 relationships among these potential environmental predictors, and (3)
322 how the scores of the principal components were related to the phytoplankton diversity
323 metrics. The procedure was applied to each single time-series separately. For each

324 environmental variable, the annual mean and the coefficient of variation (COV), used here as
325 an index of seasonal variation, were calculated. The environmental data were first
326 normalized using the omnibus procedure (Legendre and Legendre, 1998). The correlation
327 matrix of all standardized variables was used to calculate the eigenvectors and the Principal
328 Components (PCs). The PCs were then ranked in order of significance and the contribution
329 of each variable to each PC was calculated. To check for nonlinearity among environmental
330 descriptors, the multinormality of the PCs was tested. The outcome of the PCA was used to
331 investigate the relationships of phytoplankton diversity with a combination of
332 environmental factors instead of computing a suite of correlation coefficients of diversity
333 with single factors. Linear Bravais–Pearson’s correlations were calculated to assess the
334 relationship between each PC and the phytoplankton diversity indices.

335

336 2.2. Measuring beta diversity

337 Since considerable community changes can occur without being reflected in alpha diversity,
338 we also used measures of directional turnover to investigate the rate of change in community
339 structure. Here, we applied a beta diversity measure to assess the change in community
340 structure from one sampling unit to another along a temporal gradient (from year to year)
341 (see Andersen et al., 2011 for definitions on beta diversity). According to Legendre and De
342 Cáceres (2013), total beta diversity can be partitioned into Species Contributions (SCBD:
343 degree of variation of individual species across the study area) and Local Contributions
344 (LCBD: comparative indicators of the ecological uniqueness of the sites) to Beta Diversity.
345 For the objective of the study, we were interested in the LCBD indices that indicate how
346 much each observation contributes to the total community variance in time. Where a year

347 with an average species composition would have an LCBD value of 0, large LCBD values
348 may indicate degraded and species-poor sites that are in need of restoration (Legendre and
349 De Cáceres, 2013). High values may also correspond to special ecological conditions, or may
350 result from the disturbance effect of invasive species on communities. Here, temporal beta
351 diversity was computed as the method described in detail by Legendre and De Cáceres
352 (2013). Firstly, the raw abundance data were transformed using the Chord method (Legendre
353 and Galagher, 2001). Secondly, the total variance of the transformed community composition
354 was calculated by taking the squared deviations from the column means. The relative
355 contribution of the sampling unit j to beta or LCBD is the sum of squares for each sampling
356 unit divided by the total sum of squares. The statistical significance of the LCBD values was
357 also calculated. For the years where significant LCBD values were found, the Importance
358 Value Index (IVI; Curtis, 1959) was calculated. In addition to diversity indices, the IVI can be
359 used to indicate the overall importance of a species in a community (Jose, 2012) and here, to
360 potentially identify the taxa (genus) responsible for the “unusual” community structure. For
361 the genera where only one species was identified, the species instead of the genus name was
362 retained. The IVI (Eq. 3) was calculated as the sum of the relative density (RD; Eq. 4) and the
363 relative frequency (RF; Eq. 5) of the taxonomic units in the community.

$$364 \quad \text{IVI} = \text{RD}_i + \text{RF}_i \quad (3)$$

365 Here, the RD reflected the numerical strength of a genus in relation to the total number of
366 individuals of all the genera and can be calculated as:

$$367 \quad \text{RD}_i = (n_i / N) * 100 \quad (4)$$

368 where n_i is the number of individuals of the genus i and N is the total number of individuals
369 of all the genera. The RF is the degree of dispersion of individual genera over time in relation
370 to the number of all the genera which occurred in the time-series.

371
$$RF_i = (f_i/F)*100 \quad (5)$$

372 where f_i is the number of occurrence of the genus i and F is the total number of occurrence of
373 all the genera.

374 For these analyses, only monthly abundance time-series data (at the genus level) from the
375 Ouest Loscolo and Le Croisic site (Bay of Biscay) were considered, as these long time-series
376 (>25 years) provided the most robust analyses compared to the shorter available data set of
377 SOMLIT-Astan. In the graphical representations, only the top 5 genera with the highest IVI
378 values are shown.

379 All analyses were carried out using the software package MATLAB R2015a.

380

381 2. Results

382 Species accumulation curves showed that our observed richness values likely
383 underestimated the total richness of the phytoplankton communities (Figure S1). For the
384 three datasets, there is an increasing trend in the number of species along the time-series and
385 the curves did not reach saturation level indicating that the total community has not been
386 sampled yet.

387

388 Using all nine indices, correlation analyses investigated the likely redundancy between
389 indices from a mathematical perspective. Similar results were obtained for all sampling sites
390 but only the results for SOMLIT-Astan are presented here (Table 1). As expected, strong
391 correlations between diversity measures were found. This is not surprising as they represent
392 aspects of the same phenomenon (Morris et al., 2014). For the richness group, the Margalef's
393 index (d) and the number of genera (S) were highly and positively correlated ($r^2=0.87$). The

394 Menhinick's index (D) was not related to the other indices within the group suggesting that
395 its information is complementary to the two others. For the dominance indices, the Hulburt's
396 index (δ), the Simpson's index (λ), the Shannon index (H') and the Berger Parker's index (BP)
397 were all strongly related ($r^2 > 0.90$). Between categories, D was strongly and negatively related
398 ($r^2 \geq -0.90$) to the Brillouin's index (H_B) and this could suggest that these metrics carry similar
399 information despite not being related mathematically. The Pielou's index (J') was not
400 significantly related to any of the other indices. The behaviour of geometric means (G_j) could
401 not be investigated since it requires that each species is recorded in every year.
402 Unfortunately, relative abundance estimates of many phytoplankton species were equal to
403 zero and thus G_j could not be calculated.

404

405 The Principal Components Analysis (PCA) investigated the relationships among the mean
406 and seasonal variations in physico-chemical factors (Fig. 2), and the relationships of the PC
407 with phytoplankton diversity indices (Table 2). Similar correlations were found for the
408 different test sites, suggesting that the analyses explain the general behaviour of the index
409 and that the responses are not only a function of the prevailing local environmental
410 conditions. In SOMLIT-Astan, for example, the first Principal Component (PC1) explained
411 43% of the variation in the data where temperature, nitrate, phosphate and silicate
412 contributed mostly (Fig. 2a). The PC2 was explained by salinity, oxygen and nitrite and
413 accounted for 26% in the variation. For the seasonal variations in the environmental factors
414 (Fig. 2b), the PC1 explained 28% and the PC2 explained 26%. However, in terms of the
415 correlations with the PC and diversity indices, the seasonal variations in environmental
416 factors are more strongly related to diversity than annual mean conditions (Table 2). For the

417 richness group, D was the metric best explained by the seasonal variations in environmental
418 factors for SOMLIT-Astan ($r^2 = 0.76$; $p < 0.001$).

419 For the dominance metrics, H_B best reflected the seasonal variations in the environment ($r^2 =$
420 0.74 ; $p < 0.001$). This common sensitivity of D and the H_B in relation to changes in the
421 environment might explain the strong interrelationships previously detected (Table 1).

422

423 A summary table describes the performance for each α diversity index in relation to the
424 previously described criteria: ecological relevance, mathematical consistency and link with
425 hydrological conditions (Table 3). The final selection for the indices included D to describe
426 genus richness and δ to describe genus dominance since they have the best scores for the
427 three criteria. Whilst J' described a different aspect of diversity, this measure was not
428 retained for the PH3 indicator since it contained little complementary information for the
429 assessment.

430

431 To investigate the seasonal and annual variations in the three aspects of diversity
432 simultaneously, contour plots of genus richness (expressed here as D), dominance (expressed
433 here as δ) and evenness (J') per sampling site are shown (Fig. 3). Since similar trends in
434 biodiversity change were found for those indices that are strongly interrelated, only the
435 contour plots of the three previously selected indices (indicated in bold in Table 3) are
436 presented here. Here, both richness and dominance were highly variable between years and
437 variations were site-specific. In contrast, the evenness was comparatively less variable and
438 showed trends that were more similar than the ones encountered for dominance. For the
439 longer time-series of Le Croisic and Ouest Loscolo, there was an increase in the number and
440 duration of high dominance events along the period. For Le Croisic, for example, there

441 seemed to be a trend where the start of the dominance period occurred earlier in the year
442 from 2001 onwards. For Ouest Loscolo, the dominance period was nearly extended across all
443 seasons with longer peak periods (from 2007) compared to earlier years in the time-series
444 where the dominance periods were confined to spring and autumn times. This seasonal
445 expansion of high dominance correlated with increased periods of low richness and
446 evenness.

447 For SOMLIT-Astan, a short but high dominance event was recorded in May 2008 with an
448 unusually low dominance in September of the same year (Fig. 3; Fig. S2a). The next year, the
449 dominance period was more spread out from mid-April to October with two peaks in May
450 and September.

451 Whilst the contour plots for α diversity indices informed on the state of the community, the β
452 index was able to detect significant temporal changes at the community (LCBD) and the
453 genus level (IVI) on an annual basis. For Le Croisic, a year of relatively low richness and high
454 dominance (2007) was followed by a year of high richness, with peaks in June-July and
455 September (2008) (Fig. 3, Fig. S2b). The events in 2007 were marked by a relatively elevated
456 value of the LCBD (0.26) indicating a significant shift in the phytoplankton community
457 structure (Fig.4). Upon visual inspection of the IVI for the same year (Fig. 5a), the peak in
458 dominance was due to the blooming of the species *Lepidodinium chlorophorum* (47%) with an
459 abundance of 3.9×10^6 cells L^{-1} in July and to a lesser extent to the genera *Skeletonema* spp. (1.5
460 $\times 10^6$ cells L^{-1}) in April and *Leptocylindrus* spp. in Mai (5.4×10^5 cells L^{-1}) and September (6.13
461 $\times 10^5$ cells L^{-1}). The previous year at the same site was characterised by a community
462 dominated by *Chaetoceros* spp. (32%) and *Gymnodinium* spp. (18%) with lower abundances
463 ($<8 \times 10^5$ cells L^{-1}). In 2014, a value of the LCBD (0.25) similar to that of 2007 was found, that
464 also coincided with a bloom of *Lepidodinium chlorophorum* (77%), with an abundance of

465 1.15×10^7 cells L^{-1} (Fig. 5b). Before and after the bloom, *Leptocylindrus* spp. (13%) was also
466 abundant ($>8 \times 10^5$ cells L^{-1}). Similarly, in the Ouest Loscolo site, high LCBD (0.45) and
467 dominance values were recorded in 2008 (Fig. 3). In this case, a monospecific bloom of
468 *Leptocylindrus* spp. (73%) that peaked in July (2.2×10^6 cells L^{-1}) and October (8×10^6 cells L^{-1})
469 was responsible (Fig. 5c). Earlier in the year, smaller blooms were recorded in April for the
470 genus *Skeletonema* spp. (1.17×10^6 cells L^{-1}) and in June for the Chaetocerotaceae (1.8×10^6 cells
471 L^{-1}). In 2011, an unusually high richness and relatively low dominance was recorded at Ouest
472 Loscolo but this marked change in community structure was not reflected in the LCBD's.
473 This shows the importance to consider both α and β diversity indices together to detect and
474 interpret potential changes in the phytoplankton community structure.

475

476 Discussion

477 Ecological indicators based on key functional groups, such as phytoplankton, can provide
478 sensitive and quantifiable indications of ecological changes and environmental perturbations
479 in marine surface waters (Paerl et al., 2003; Rombouts et al, 2013). The common OSPAR
480 Pelagic Habitat indicator "Changes in plankton diversity" was developed as a surveillance
481 indicator to describe the phytoplankton community structure and to identify temporal
482 changes or "events" within the assessment period. Since biodiversity is multi-dimensional,
483 no single measure can meet all needs for assessing change (Buckland et al., 2017). It is,
484 therefore, important to use PH3 as a composite indicator where the alpha diversity, i.e. the
485 diversity within a site or sample, and the beta diversity that focuses on the rate of change, or
486 turnover, in species composition are being considered. For this purpose, four indices were
487 identified that focus on different aspects of plankton biodiversity from a community to

488 genus level namely the taxon (genus) richness (Menhinick's index, D), dominance (Hulburt
489 index, δ), temporal variation (Local Contributions to Biodiversity, LCBD) and taxa
490 identification (Important Value Index, IVI). Whilst the richness and dominance indices are
491 evaluated on a monthly basis, the temporal variation and taxa identification are assessed on
492 an annual level.

493

494 The final selection of one richness and one dominance index was based on a comparative
495 analysis of the metrics' performances. The performances were mainly evaluated from an
496 ecological perspective and from the sensitivity of the metrics but ultimately, the selected
497 indices were retained on their ability to synthesise relevant information in an understandable
498 and unambiguous manner to stakeholders. The Menhinick's diversity index (D) was selected
499 as the most appropriate metric to describe the number of taxa in the community. In this
500 study, it was found to be the most sensitive to changes in environmental conditions that
501 could be either from a natural or an anthropogenic source. Similar studies agree that D is one
502 of the most efficient tools for the assessment of water quality (e.g. Facca et al., 2014; Spatharis
503 and Tsirtsis, 2010; Buzançıç et al., 2016; Varkitzi et al., 2018). However, caution must be taken
504 when interpreting any index based on estimates of the number of species in the community
505 since these are biased (Heip et al., 1998). An observed increase in the counts of
506 phytoplankton taxa and thus an increase in the biodiversity index can have numerous
507 causes: sampling methods (Rodriguez-Ramos et al., 2014) and effort (Cozzoli et al., 2017),
508 advection of new taxa (Lévy et al., 2014; Sun and Xue, 2016), increased knowledge of the
509 taxonomic analyst (Dromph et al., 2013), etc. Whilst these factors likely underestimate the
510 true taxonomic diversity in the phytoplankton community, here, we are more interested in
511 the overall state and the relative changes in the community composition on a seasonal and

512 annual basis. In any case, considering the highly intra-annual variability of taxa and
513 abundances, consistent monthly monitoring is essential when quantifying phytoplankton
514 community diversity. Also, any taxonomic richness index should be interpreted in
515 conjunction with a dominance index to better understand the overall structure of the
516 phytoplankton community. Here, visual inspection suggests a seasonal expansion of the low
517 diversity in conjunction with high dominance periods over years, especially notable for the
518 longer time-series, Ouest Loscolo and Le Croisic.

519

520 Dominance phenomena and significant changes in phytoplankton community structure can
521 occur in impacted areas (e.g. Buzançıç et al., 2016). Here, as a dominance measure, the
522 Hulburt index (δ) was mainly selected for its ease of interpretation (as a percentage, where a
523 high value indicates high dominance) but also for its recent applications in water quality
524 assessments (Facca et al., 2014). Using the Principal Component Analysis, the Brillouin index
525 (H_B) was found to be the only dominance measure that explained the variations in the
526 environment but since this metric was interrelated with D and thus likely to be redundant,
527 the former was not retained. Periods of relatively high dominance were also identified by the
528 LCBDs as a general period of significant change or turnover. For the stations Ouest Loscolo
529 and Le Croisic in the Bay of Biscay, 2007 and 2008, respectively, were identified as years with
530 a temporary shift to relatively high community variation. The analysis of the Important
531 Value Index (IVI) showed that these observed temporal shifts in community structure were
532 marked by a monospecific bloom from *Leptocylindrus* spp. (a diatom - at Ouest Loscolo, > 8
533 million cells L⁻¹) and *Lepidodinium* spp. (a dinoflagellate - at Le Croisic, > 4 million cells L⁻¹). A
534 high increase of biomass, so called bloom events if the number of cells > 1 million cells L⁻¹,
535 can be a result of nutrient inputs such as nitrate and phosphate (Alves-de-Souza et al., 2006),

536 but also of changing environmental conditions, for example temperature and salinity
537 (Pizarra et al., 1997). *Lepidodinium chlorophorum*, for example, is known to form regular
538 “green” blooms over the French Atlantic Shelf (Sourisseau et al., 2016), but in the year 2007 a
539 unusual high number of events was observed (Chauvin, 2012). In terms of ecological
540 impacts, their blooms can cause anoxia and bright-green coloured waters. For the genus
541 *Leptocylindrus* spp, the unusual high temperatures recorded in 2007 could explain the
542 observed bloom since the genus has an ecological niche of relatively warm temperatures and
543 high light conditions (Hernández-Fariñas et al., 2013). Whilst *Leptocylindrus* spp. has been
544 identified as an indicator of eutrophication (Ninčević-Gladan et al., 2015), there are no
545 records of a similar application in our study area. In this specific case, taxa identification
546 using the IVI index helped to understand the ecological behaviour of the taxa (for example,
547 as a response to environmental conditions). Also, in case a genus would develop into a
548 Harmful Algal Bloom (HAB), the potential effects of blooming taxa on the ecosystem could
549 be investigated. Further analyses of the effects of natural and anthropogenic pressures on
550 phytoplankton communities will help to identify the most effective mechanisms and the
551 actions needed to maintain or to restore GES conditions (Crise et al., 2015).

552

553 Volume indices, such as the geometric mean of relative abundance (G), are increasingly
554 being used to examine trends in biological diversity and to assess whether biodiversity
555 targets are being met (Buckland et al., 2011). In contrast to the Shannon’s and Simpson’s
556 indices, G will decline if all species are declining at the same rate even if there is no trend in
557 evenness. Whilst the concept of this volume index is interesting, the geometric mean has also
558 a number of drawbacks that unfortunately make the index unsuitable for assessing
559 phytoplankton communities. Most importantly, the index is based on within-taxon trends

560 and requires a robust calculation where each taxon is recorded in every year. Since
561 phytoplankton datasets are generally characterized by a small number of abundant species
562 and many rare species, the index is likely to exhibit high variance and unstable behaviour
563 when species are not consistently present in the community. A potential solution would be
564 to calculate the index on only those taxa that are present in every sample but then the index
565 would reflect trends of the subset of taxa and not the whole community, and as such, the
566 index has limited use as a community diversity measure to assess GES of pelagic habitats.

567

568 Compared to phytoplankton biomass indicators, the development of community
569 composition indicators for water quality assessment is in its early stages. Firstly, the
570 responses of phytoplankton community composition to a combination of nutrients is
571 relatively unpredictable and so, establishing significant pressure-state relationships can
572 become difficult (Garmendia et al., 2013; Ochocka and Pasztaleniec, 2016), especially in
573 marine open water systems. Studies of phytoplankton communities in relation to pressure
574 gradients confirmed the intermediate disturbance level hypothesis, which predicts high
575 richness in areas subjected to intermediate levels of disturbance (Sommer et al., 1993;
576 Ninčević-Gladan et al., 2015). So in line with this view, high diversity does not necessarily
577 correlate with “good” environmental conditions. Conversely, the presence of blooms could
578 be perceived as “negative” by societies but can be often driven by natural conditions. As
579 long as the pressure–state relationships are inadequately understood, ecologically
580 meaningful boundaries and thus targets to assess GES cannot be defined for PH3.
581 Unfortunately, we were unable to examine the behaviour of the indicator under different
582 stressor scenarios. Whilst PH3 will need further development to support formal state
583 assessment, the indicator can still be very informative on the health of the environment and

584 act as a “surveillance” indicator rather than an operational one. Although, “surveillance”
585 indicators do not directly track state in relation to GES, they do provide complementary
586 information (highlighting a « specific cause for concern ») that presents a broader and more
587 holistic picture of state, and inform and support science, policy, and management (Shephard
588 et al., 2015; Varkitzi et al., 2018; Bedford et al., 2018). In this respect, PH3, in its current state
589 of development, will act as a warning signal by highlighting unprecedented or directional
590 state shifts in the plankton communities of the marine pelagic habitat.

591

592 Detecting trends in the structure of phytoplankton communities is achievable but requires
593 the collection of suitable data (Ajani et al., 2014). Long-term monitoring networks of
594 sufficient spatial and temporal resolution are needed to distinguish the anthropogenic and
595 natural processes that affect the phytoplankton abundance and composition, and to be able
596 to detect significant changes in the community structure in a robust manner. Several
597 transnational projects and conventions have already highlighted the need for appropriate
598 monitoring programs to feed biodiversity indicators and associated parameters. The
599 PERSEUS project, for example, pointed out the lack of quantitative data on pressures and a
600 lack of spatial coverage, in particular offshore data on nutrients, phytoplankton and
601 dissolved oxygen (Crise et al., 2015). For more complete regional assessments, in particular,
602 better acquisition of region-wide plankton data and coherent monitoring programmes will
603 still be required (Caroppo et al., 2013; OSPAR, 2017d; Varkitzi et al., 2018). In terms of
604 sampling frequency, a minimum of bimonthly sampling is advised for estimating
605 phytoplankton biodiversity (Uusitalo et al., 2013; OSPAR, 2017d). With regards to the
606 analysis of the phytoplankton community data, light microscopy is the most commonly used
607 laboratory technique for the determination of the abundance and species identification

608 (OSPAR, 2016). Whilst this method is time-consuming and requires a high degree of
609 expertise (Havskum et al., 2004), detailed taxonomic data, containing information on the
610 presence/absence and abundance of individual plankton species, are required to underpin
611 the development of sensitive species and community-level indicators (Beaugrand et al., 2005;
612 McQuatters-Gollop et al., 2017). In this respect, well-educated microscopists are necessary for
613 obtaining reliable phytoplankton monitoring results (Lehtinen et al., 2012). Unfortunately,
614 adequate funding to support plankton taxonomy in line with its value to science and
615 decision making remains a key challenge to ensuring the availability of plankton data for
616 marine policy and conservation (McQuatters-Gollop et al., 2017). Innovative analysis
617 techniques exist (OSPAR, 2016; Karlson et al., 2016; Chust et al., 2017; Aubert et al., 2017) but
618 it is difficult to find a “one size fits all” method for counting and characterizing the
619 composition of the phytoplankton communities in marine systems, due to their intrinsically
620 high spatial and temporal variability (Garmendia et al., 2013), and diversity of sizes (Sieburth
621 et al., 1978). In any case, microscopic data will still be required to support and validate new
622 analytical methods and to test indicators derived from these new types of monitoring
623 (McQuatters-Gollop et al., 2017).

624

625 Whilst some authors remain sceptical of the community composition approach (e.g.
626 Ninčević- Gladan et al., 2015), others have demonstrated successful applications of
627 composition based metrics for water quality assessment, mainly developed for use in the
628 WFD (e.g. Tett et al., 2008; Devlin et al., 2009; Facca et al., 2014). In most cases, these
629 assessments were carried out using multimetric indicators because the inclusion of
630 additional metrics can render an index more sensitive and robust (e.g. Hering et al., 2006;
631 Rombouts et al., 2013). When selecting indicators, the aggregation (combined use of several

632 indicators for an ecosystem-based approach) should consider different elements of
633 community response to environmental change, e.g. taxonomic and functional diversity,
634 biomass, species composition and the presence of opportunistic or non-indigenous species
635 (Lehtinen et al., 2012; Zettler et al., 2017). In case of the common OSPAR indicators, this type
636 of aggregation could be achieved by combining each Pelagic Habitat (PH) indicator where
637 the plankton community is considered at different resolutions, PH1 at the life-form level of
638 the community, PH2 for the total biomass/abundance of the community and PH3 at the
639 species level. Hence, by combining the information from these three indicators, a more
640 holistic assessment of plankton dynamics can be obtained than from each indicator
641 individually.

642

643 With the current OSPAR common indicators, the determination of the ecological quality of
644 the pelagic habitat is based on the biological quality elements only, the plankton. According
645 to Article 3 of the MSFD, however, “Good Environmental Status” (GES) for pelagic habitats
646 is defined by “the structure, functions, and processes of the constituent marine ecosystems,
647 together with the associated physiographic, geographic, geological and climatic factors,
648 allow those ecosystems to function fully and to maintain their resilience to human-induced
649 environmental change.” Even with a clear definition of GES, the variability in prevailing
650 conditions of the marine environment makes recognising if we have reached GES
651 challenging, especially for pelagic habitats. Therefore, a more integrated approach that also
652 accounts for the non-biological components of the sea water will need to be developed
653 (Ferreira et al., 2011; Rombouts et al., 2013). Recently, Dickey-Collas and colleagues (2017)
654 discussed the challenges related to the concept of “good” environmental status of pelagic
655 habitats and propose directions for reflection and research to effectively monitor progress

656 towards, or movement from, GES. In summary, the authors propose three conditions that
657 should be met for pelagic habitats to be in GES: (i) all species present under current
658 environmental conditions have access to the pelagic habitats essential to close their life
659 cycles; (ii) biogeochemical regulation is maintained at normal levels; (iii) critical physical
660 dynamics and movements of biota and water masses at multiple scales are not obstructed.

661

662 For now, the current determination of GES for pelagic habitats takes a pragmatic approach
663 and largely relies on existing information, data and methodologies. Especially for pelagic
664 habitats, monitoring all species groups in all pelagic habitat types in all localities is simply
665 not feasible. At best, it is possible to monitor a selection of species groups, preferably species
666 sensitive to environmental change over relatively short time-scales and where data can be
667 collected to ensure regular updates (Van Strien et al., 2012 and references therein). Any
668 outstanding issues can be addressed during subsequent MSFD cycles through, for example,
669 the development of new methodologies (Danovaro et al., 2016), the gathering of additional
670 data through monitoring programmes and further development of indicators (EC, 2011;
671 Padegimas et al., 2017). In line with the ongoing work within OSPAR and other Regional
672 Seas conventions, the further implementation of the MSFD will continue to be agreed with
673 the stakeholders at transnational level and to be based on solid scientific knowledge (Varkitzi
674 et al., 2018). The pilot study for the development of PH3 presented here is based on the
675 outcome of the Intermediate Assessment 2017 and this type of preliminary assessment is the
676 starting point of a long-term iterative process.

677

678

679

680 Acknowledgements

681 I.R. and A.A. received funding from the French Ministry for the Ecological and Solidary
682 Transition (MTES), the French National Centre for Scientific Research (CNRS-INEE, CNRS-
683 INSU), the French National Museum of Natural History (MNHN) and the EU DG
684 ENV/MSFD/Action Plan Project (11.0661/2015/712630/SUB/ENVC.2 OSPAR) Applying an
685 ecosystem approach to (sub) regional habitat assessments (EcApRHA). A.M-G would like to
686 thank the UK National Environmental Research Council for support through the NERC
687 Knowledge Exchange fellowship scheme. We want to thank all members of the REPHY
688 program and the SRN network of the Institute for the Exploitation for the Sea (IFREMER) for
689 the use of the Ouest Loscolo and Le Croisic data and the RESOMAR-Pelagos and SOMLIT
690 team for the use of the SOMLIT-Astan data. In particular, we thank Stéphanie Ristori, Manon
691 Viprey and Loïc Guilloux.

692 We thank the anonymous reviewers for their careful reading of our manuscript and their
693 many insightful comments and suggestions.

694

695 References

696

697 Ajani, P.A., Allen, A.P., Ingleton, T. and Armand, L. (2014) A decadal decline in relative
698 abundance and a shift in microphytoplankton composition at a long-term coastal station
699 off southeast Australia. *Limnology and Oceanography* 59: 519-531.

700

701 Alves-de-Souza, C., Menezes, M. and Huszar, V. (2006) Phytoplankton composition and
702 functional groups in a tropical humic coastal lagoon, Brazil. *Acta Botanica Brasilica* 20:
703 701-708.

704

705 Aubert, A., Rombouts, I., Artigas, L.F., Budria, A., Ostle, C., Padegimas, B. and
706 McQuatters-Gollop, A. (2017) Combining methods and data for a more holistic
707 assessment of the plankton community. EcApRHA deliverable WP 1.2., OSPAR, London.

708

709 Bandeira, B., Jamet, J.-L., Jamet, D. and Ginoux, J.-M. (2013) Mathematical convergences
710 of biodiversity indices. *Ecological Indicators* 29, 522–528.

711 <https://doi.org/10.1016/j.ecolind.2013.01.028>

712

713 Bedford, J., Johns, D., Greenstreet, S. and McQuatters-Gollop, A. (2018) Plankton as
714 prevailing conditions: a surveillance role for plankton indicators within the Marine
715 Strategy Framework Directive. *Marine Policy* 89: 109-115.

716

717 Ben Othman, H., Lanouguère, É., Got, P., Sakka Hlaili, A. and Leboulanger, C. (2018)
718 Structural and functional responses of coastal marine phytoplankton communities to
719 PAH mixtures. *Chemosphere* 209: 908-919.

720

721 Beaugrand, G. (2005) Monitoring pelagic ecosystems using plankton indicators. *ICES*
722 *Journal of Marine Science* 62: 333-338.

723

724 Buckland, S.T., Baillie, S.R., Dick, J.M., Elston, D., Magurran, A., Scott, E.M., Smith, R.,
725 Somerfield, P., Studeny, A. and Watt, A. (2012) How should regional biodiversity be

726 monitored? *Environmental and Ecological Statistics* 19: 601.

727 <https://doi.org/10.1007/s10651-012-0202-7>.

728

729 Buckland, S.T., Magurran, A.E., Green, R.E. and Fewster, R.M. (2005) Monitoring change

730 in biodiversity through composite indices. *Philosophical Transactions of the Royal*

731 *Society B: Biological Sciences* 360: 243-254.

732

733 Buckland, S.T., Studeny, A.C., Magurran, A.E., Illian, J.B. and Newson, S.E. (2011) The

734 geometric mean of relative abundance indices: a biodiversity measure with a difference.

735 *Ecosphere* 2(9):100. <https://doi.org/10.1890/ES11-00186.1>

736

737 Buckland, S.T., Yuan, Y. and Marcon, E. (2017) Measuring temporal trends in biodiversity

738 *A St A Advances in Statistical Analyses* (2017) 101: 461.

739 <https://doi.org/10.1007/s10182-017-0308-1>

740

741 Bužančić, M., Ninčević Gladan, Ž., Marasović, I., Kušpilić, G. and Grbec, B. (2016)

742 Eutrophication influence on phytoplankton community composition in three bays on the

743 eastern Adriatic coast. *Oceanologia* 58: 302-316.

744 <http://dx.doi.org/10.1016/j.oceano.2016.05.003>

745

746 Caroppo, C., Buttino, I., Camatti, E., Caruso, G., De Angelis, R., Facca, C., Giovanardi, F.,

747 Lazzara, L., Mangoni, O. and Magaletti, E. (2013) State of the art and perspectives on the

748 use of planktonic communities as indicators of environmental status in relation to the EU
749 Marine Strategy Framework Directive. 44° Congresso della Società Italiana di Biologia
750 Marina Roma, 14-16 maggio 2013

751

752 CEC (2008) Directive 2008/56/EC of the European Parliament and of the Council of 17
753 June 2008 establishing a framework for community action in the field of marine
754 environmental policy (Marine Strategy Framework Directive). Official Journal of the
755 European Union, L 164, 25/06/2008, 19–40. [http://eur-lex.europa.eu/legal-](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32008L0056&from=EN)
756 [content/EN/TXT/PDF/?uri=CELEX:32008L0056&from=EN](http://eur-lex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32008L0056&from=EN) (accessed 18 April 2017).

757

758 Chalar, G. (2009) The use of phytoplankton patterns of diversity for algal bloom
759 management. *Limnologica - Ecology and Management of Inland Waters* 39: 200-208.

760

761 Chiarucci, A., Bacaro, G. and Scheiner, S.M. (2011) Old and new challenges in using
762 species diversity for assessing biodiversity. *Philosophical Transactions of the Royal*
763 *Society B: Biological Sciences* 366: 2426-2437.

764

765 Chust G., Vogt M., Benedetti F., Nakov T., Villéger S., Aubert A., Vallina S., Righetti D.,
766 Not F., Biard T., Bittner L., Benoiston A.S., Guidi L., Villarino E., Gaborit C., Cornils A.,
767 Buttay L., Irisson J.O., Chiarello M., Lima A., Blanco-B.L., Basconi L. and Ayata S.D.
768 (2017) Mare incognitum: A glimpse into future plankton diversity and ecology research.
769 *Frontiers in Marine Science* 4: 68. <https://doi.org/10.3389/fmars.2017.00068>

770

771 Coclet, C., Garnier, C., Delpy, F., Jamet, D., Durrieu, G., Le Poupon, C., Mayer, M. and
772 Misson, B. (2018) Trace metal contamination as a toxic and structuring factor impacting
773 ultraphytoplankton communities in a multicontaminated Mediterranean coastal area.
774 Progress in Oceanography 163: 196-213.

775

776 Cozzoli, F., Stanca, E., Selmeçy, G.B., Francé, J., Varkitzi, I. and Basset, A. (2017)
777 Sensitivity of phytoplankton metrics to sample-size: A case study on a large transitional
778 water dataset (WISER). Ecological Indicators 82: 558-573.

779

780 Crise, A., Kaberi, H., Ruiz, J., Zatsepin, A., Arashkevich, E., Giani, M., Karageorgis, A.P.,
781 Prieto, L., Pantazi, M. and Gonzalez-Fernandez, D. (2015) A MSFD complementary
782 approach for the assessment of pressures, knowledge and data gaps in Southern
783 European Seas: The PERSEUS experience. Marine Pollution Bulletin 95(1): 28-39.

784

785 Danovaro, R., Carugati, L., Berzano, M., Cahill, A.E., Carvalho, S., Chenuil, A.,
786 Corinaldesi, C., Cristina, S., David, R., Dell'Anno, A., Dzhembekova, N., Garcés, E.,
787 Gasol, J.M., Goela, P., Féral, J.-P., Ferrera, I., Forster, R.M., Kurekin, A.A., Rastelli, E.,
788 Marinova, V., Miller, P.I., Moncheva, S., Newton, A., Pearman, J.K., Pitois, S.G., Reñé, A.,
789 Rodríguez-Ezpeleta, N., Saggiomo, V., Simis, S.G.H., Stefanova, K., Wilson, C., Lo
790 Martire, M., Greco, S., Cochrane, S.K.J., Mangoni, O. and Borja, A. (2016) Implementing
791 and Innovating Marine Monitoring Approaches for Assessing Marine Environmental
792 Status. Frontiers in Marine Science 3.

793

794 Devlin, M., Barry, J., Painting, S. and Best, M. (2009) Extending the phytoplankton tool kit
795 for the UK Water Framework Directive: indicators of phytoplankton community
796 structure. *Hydrobiologia* 633: 151-168.

797

798 Dickey-Collas, M., McQuatters-Gollop, A., Bresnan, E., Kraberg, A. C., Manderson, J. P.,
799 Nash, R. D. M., Otto, S. A., Sell, A. F., Tweddle, J. F. and Trenkel, V. M. (2017) Pelagic
800 habitat: exploring the concept of good environmental status. *ICES Journal of Marine*
801 *Science* 74: 2333–2341.

802

803 Domingues, R.B., Barbosa, A., and Galvão, H. (2008) Constraints on the use of
804 phytoplankton as a biological quality element within the Water Framework Directive in
805 Portuguese waters. *Marine Pollution Bulletin* 56: 1389–1395.

806

807 Domingues, R.B., Guerra, C.C., Galvão, H.M., Brotas, V. and Barbosa, A.B. (2017) Short-
808 term interactive effects of ultraviolet radiation, carbon dioxide and nutrient enrichment
809 on phytoplankton in a shallow coastal lagoon, *Aquatic Ecology* 51: 91.

810 <https://doi.org/10.1007/s10452-016-9601-4>

811

812 Dromph, K.M., Agusti, S., Basset, A., Franco, J., Henriksen, P., Icely, J., Lehtinen, S.,
813 Moncheva, S., Revilla, M., Roselli, L. and Sørensen, K. (2013) Sources of uncertainty in
814 assessment of marine phytoplankton communities. *Hydrobiologia* 704: 253-264.

815 <https://doi.org/10.1007/s10750-012-1353-0>

816

817 EC (2011) Working Group on Good Environmental Status. Common Understanding of
818 (Initial) Assessment, Determination of Good Environmental Status (GES) and
819 Establishment of Environmental Targets (Art. 8, 9 and 10 MSFD).

820

821 Facca, C., Bernardi Aubry, F., Socal, G., Ponis, E., Acri, F., Bianchi, F., Giovanardi, F. and
822 Sfriso, A. (2014) Description of a Multimetric Phytoplankton Index (MPI) for the
823 assessment of transitional waters. *Marine Pollution Bulletin* 79: 145–154.
824 <https://doi.org/10.1016/j.marpolbul.2013.12.025>

825

826 Gallardo, B., Gascón, S., Quintana, X. and Comín, F.A. (2011) How to choose a
827 biodiversity indicator – Redundancy and complementarity of biodiversity metrics in a
828 freshwater ecosystem. *Ecological Indicators* 11: 1177-1184.

829

830 Garmendia, M., Borja, A. Franco, J. and Revilla, M. (2013) Phytoplankton composition
831 indicators for the assessment of eutrophication in marine waters: Present state and
832 challenges within the European directives, *Marine Pollution Bulletin* 66 (1–2): 7-16.

833

834 Gascon, S., Boix, D., and Sala, J. (2009) Are different biodiversity metrics related to the
835 same factors? A case study from Mediterranean wetlands. *Biological Conservation* 142:
836 2602–2612.

837

838 Guilloux L., Rigaut-Jalabert F., Jouenne F., Ristori S., Viprey M., Not F., Vaultot, D and
839 Simon, N. (2013) An annotated checklist of marine phytoplankton taxa at the SOMLIT-
840 Astan time-series off Roscoff (Western English Channel, France) : data collected from
841 2000 to 2010. Cahiers de Biologie Marine 54: 247-256.

842

843 Havskum, H., Schlüter, L., Scharek, R., Berdalet, E. and Jacquet, S. (2004) Routine
844 quantification of phytoplankton groups: microscopy or pigment analyses? Marine
845 Ecology Progress Series 273: 31-42.

846

847 Hernández-Fariñas, T., Soudant, D., Barillé, L., Belin, C., Lefebvre, A., and Bacher, C.
848 (2013). Temporal changes in the phytoplankton community along the French coast of the
849 eastern English Channel and the southern Bight of the North Sea. ICES Journal of Marine
850 Science 71 (4): 821–833. <https://doi.org/10.1093/icesjms/fst192>

851

852 Heip, C.H.R., Herman, P.M.J. and Soetaert, K. (1998) Indices of diversity and evenness.
853 Oceanis 24 (4): 61-87.

854

855 Hering, D., Feld, C.K., Moog, O. and Ofenböck, T. (2006) Cook book for the development
856 of a Multimetric Index for biological condition of aquatic ecosystems: experiences from the
857 European AQEM and STAR projects and related initiatives. Hydrobiologia 566: 311-324.

858

859 Hulburt, E. (1963) The diversity of phytoplanktonic populations in oceanic, coastal, and
860 estuarine regions. Journal of Marine Research 21: 81–93.

861

862 Jolliffe, I.T. (1986) *Principal Components Analysis*. Springer-Verlag, New York.

863

864 Jose, S.K. (2012). *Phytodiversity assessment*. In: *Geospatial characterization and*
865 *conservation potential for Agasthyamala Biosphere Reserve (ABR), Western ghats, India.*

866

867 Kabuta, S. and H. Duijts (2000) *Indicators for the North Sea*. Report Rijksinstituut
868 voor Kust en Zee/RIKZ No 2000.022.

869

870 Karlson, B., Artigas, F., Créach, V., Louchart, A., Wacquet, G., Seppälä, J. (2016). *Novel*
871 *methods for automated in situ observations of phytoplankton diversity*, Joint European
872 *Research Infrastructure network for Coastal Observatory – Novel European eXpertise for*
873 *coastal observatories - JERICO-NEXT, WP3 D3.1, Table 2.1, page 7/69 - www.jerico-ri.eu*

874

875 Kissling, W. D., Ahumada, J. A., Bowser, A., Fernandez, M., Fernández, N., García, E. A.,
876 Guralnick, R. P., Isaac, N. J., Kelling, S., Los, W. , McRae, L., Mihoub, J., Obst, M.,
877 Santamaria, M., Skidmore, A. K., Williams, K. J., Agosti, D., Amariles, D., Arvanitidis, C.,
878 Bastin, L. , De Leo, F., Egloff, W., Elith, J., Hobern, D., Martin, D., Pereira, H. M., Pesole,
879 G., Peterseil, J., Saarenmaa, H., Schigel, D., Schmeller, D. S., Segata, N., Turak, E., Uhlir,
880 P. F., Wee, B. and Hardisty, A. R. (2018) *Building essential biodiversity variables (EBVs)*
881 *of species distribution and abundance at a global scale*. *Biological Review* 93: 600-625.
882 doi:10.1111/brv.12359

883

884 Kwak, T. J., and Peterson, J. T. (2007) Community indices, parameters, and comparisons.
885 In C. S. Guy and M. L. Brown (Eds.) Analysis and interpretation of freshwater fisheries
886 data. (pp. 677-763). Bethesda, MD: American Fisheries Society.

887

888 Laplace-Treyture, C. and Feret, T. (2016) Performance of the Phytoplankton Index for
889 Lakes (IPLAC): A multimetric phytoplankton index to assess the ecological status of
890 water bodies in France. *Ecological Indicators* 69: 686-698.

891

892 Legendre, P. and De Cáceres, M. (2013) Beta diversity as the variance of community data:
893 dissimilarity coefficients and partitioning. *Ecology Letters* 16: 951-963.

894

895 Legendre, P. and Gallagher, E.D. (2001) Ecologically meaningful transformations for
896 ordination of species data. *Oecologia* 129: 271. <https://doi.org/10.1007/s004420100716>.

897

898 Legendre, P. and Legendre, L. (1998) *Numerical Ecology*. Elsevier Science, Amsterdam.

899

900 Lehtinen S., Kauppila P., Kaitala S., Basset A., Lugoli F., Moncheva S., Icely J., Henriksen,
901 P. and Heiskana A.-S. (2012) Deliverable D4.1-4: Manuscript on the review of multi-
902 species indicators synthesised with WP results. SYKE (Finnish Environmental Institute),
903 <http://www.wiser.eu/download/D4.1-4.pdf>

904

905 Lévy, M., Jahn, O., Dutkiewicz, S. and Follows, M. J. (2014) Phytoplankton diversity and
906 community structure affected by oceanic dispersal and mesoscale turbulence. *Limnology
907 and Oceanography: Fluids and Environments* 4. [https://doi.org/10.1215/21573689-
908 2768549](https://doi.org/10.1215/21573689-2768549).

909

910 Lyashevskaya, O. and Farnsworth, K.D. (2012) How many dimensions of biodiversity do
911 we need? *Ecological Indicators* 18: 485-492.

912

913 Lyche-Solheim, A., Feld, C.K., Birk, S., Phillips, G., Carvalho, L., Morabito, G., Mischke,
914 U., Willby, N., Søndergaard, M., Hellsten, S., Kolada, A., Mjelde, M., Böhmer, J., Miler,
915 O., Pusch, M.T., Argillier, C., Jeppesen, E., Lauridsen, T.L. and Poikane, S. (2013)
916 Ecological status assessment of European lakes: a comparison of metrics for
917 phytoplankton, macrophytes, benthic invertebrates and fish. *Hydrobiologia* 704: 57-74.

918

919 Magurran, A.E. (1988) *Ecological diversity and its measurement*. Croom Helm.

920

921 Margalef, R. (1978) Life-forms of phytoplankton as survival alternatives in an unstable
922 environment. *Oceanologica Acta* 1: 493-509.

923

924 McQuatters-Gollop, A., Edwards, M., Helaouët, P., Johns, D.G., Owens, N.J.P., Raitos,
925 D.E., Schroeder, D., Skinner, J. and Stern, R.F. (2015) The Continuous Plankton Recorder
926 survey: How can long-term phytoplankton datasets contribute to the assessment of Good
927 Environmental Status? *Estuarine, Coastal and Shelf Science* 162: 88-97.

928

929 McQuatters-Gollop, A., Johns, D.G., Bresnan, E., Skinner, J., Rombouts, I., Stern, R.,
930 Aubert, A., Johansen, M., Bedford, J. and Knights, A. (2017) From microscope to
931 management: The critical value of plankton taxonomy to marine policy and biodiversity
932 conservation. *Marine Policy* 83: 1-10.

933

934 Menhinick, E.F. (1964) A comparison of some species-individuals diversity indices
935 applied to samples of field insects. *Ecology* 45: 859–861. <https://doi.org/10.2307/1934933>

936

937 Morris, K.E., Caruso, T., Buscot, F., Fischer, M., Hancock, C., Maier, T.S., Meiners, T.,
938 Müller, C., Obermaier, E., Prati, D., Socher, S.A., Sonnemann, I., Wäschke, N., Wubet, T.,
939 Wurst, S. and Rillig, M.C. (2014) Choosing and using diversity indices: insights for
940 ecological applications from the German Biodiversity Exploratories. *Ecology and*
941 *Evolution* 4: 3514-3524.

942

943 Muñoz, O., Rodríguez, J.G., Revilla, M., Laza-Martínez, A., Seoane, S. and Franco, J. (2018)
944 Seasonal variations of phytoplankton community in relation to environmental factors in
945 an oligotrophic area of the European Atlantic coast (southeastern Bay of Biscay). *Regional*
946 *Studies in Marine Science* 17: 59-72.

947

948 Neaud Masson, N. (2015) Observation et dénombrement du phytoplancton marin par
949 microscopie optique photonique : Spécifications techniques et méthodologiques
950 appliquées au REPHY, Ifremer, Nantes, 53 pp.

951

952 Ninčević-Gladan, Ž., Bužančić, M., Kušpilić, G., Grbec, B., Matijević, S., Skejić, S.,
953 Marasović, I. and Morović, M. (2015) The response of phytoplankton community to
954 anthropogenic pressure gradient in the coastal waters of the eastern Adriatic Sea.
955 Ecological Indicators 56: 106-115.

956

957 Ochocka, A. and Pasztaleniec, A. (2016) Sensitivity of plankton indices to lake trophic
958 conditions. Environmental Monitoring and Assessment 188: 622.

959

960 OSPAR (2016) CEMP eutrophication monitoring guidelines: phytoplankton species
961 composition. OSPAR Agreement 2016-06.

962 [http://mcc.jrc.ec.europa.eu/documents/OSPAR/CEMP_GuidelinesPhytoplanktonmonitori](http://mcc.jrc.ec.europa.eu/documents/OSPAR/CEMP_GuidelinesPhytoplanktonmonitoring.pdf)
963 [ng.pdf](http://mcc.jrc.ec.europa.eu/documents/OSPAR/CEMP_GuidelinesPhytoplanktonmonitoring.pdf)

964

965 OSPAR (2017a) Intermediate assessment 2017: Biodiversity Status :
966 Habitats. [https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-](https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/)
967 [2017/biodiversity-status/habitats/](https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/)

968 OSPAR (2017b) Intermediate assessment 2017: Biodiversity Status: Changes in
969 Phytoplankton and Zooplankton Communities. [https://oap.ospar.org/en/ospar-](https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/plankton-biomass/)
970 [assessments/intermediate-assessment-2017/biodiversity-status/habitats/plankton-](https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/plankton-biomass/)
971 [biomass/](https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-status/habitats/plankton-biomass/)

972

973 OSPAR (2017c) Intermediate assessment 2017: Biodiversity Status: Changes in
974 Phytoplankton Biomass and Zooplankton Abundance.
975 [https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-
977 status/habitats/changes-phytoplankton-and-zooplankton-communities/](https://oap.ospar.org/en/ospar-assessments/intermediate-assessment-2017/biodiversity-
976 status/habitats/changes-phytoplankton-and-zooplankton-communities/)

978 OSPAR (2017d) Intermediate assessment 2017: Biodiversity Status: Habitats: Pilot
979 assessment of Changes in Plankton Diversity. [https://oap.ospar.org/en/ospar-
982 assessments/intermediate-assessment-2017/biodiversity-status/habitats/pilot-assessment-
983 changes-plankton/](https://oap.ospar.org/en/ospar-
980 assessments/intermediate-assessment-2017/biodiversity-status/habitats/pilot-assessment-
981 changes-plankton/)

983 Padegimas, B., Artigas, L.F., Arroyo, N.L., Aubert, A., Budria, A., Capuzzo, E. Corcoran,
984 E., S. A. M. Elliott, J. M. González-Irusta, L. Guérin, A. Judd, J. Kromkamp, A.
985 McQuatters-Gollop, B. Meakins, N. Niquil10, N., Ostle, C., Pesch, R., Preciado, I., Safi, G.,
986 Schmitt, P., Serrano, A., Thorpe, R., Torriente, A. and Vina-Herbon, C. (2017) Action Plan
987 for the further implementation of habitat and food web indicators and progressing
988 integrated assessments in OSPAR (sub) regions. EcApRHA Deliverable WP5.6. 20pp.
989 ISBN: 978-1-911458-30-2

990

991 Pasztaleniec, A. (2016) Phytoplankton in the ecological status assessment of European
992 lakes – advantages and constraints. Environmental Protection and Natural Resources;
993 The Journal of Institute of Environmental Protection-National Research Institute 27(1):
994 26-36. <https://doi.org/10.1515/oszn-2016-0004>
995

996 Peet, R.K. (1974) The measurement of species diversity. *Annual Review of Ecology and*
997 *Systematics* 5: 285-307. <https://doi.org/10.1146/annurev.es.05.110174.001441>
998

999 Pereira, H. M., Ferrier, S., Walters, M., Geller, G. N., Jongman, R. H. G., Scholes, R. J. and
1000 Bruford, M. W. (2013) Essential Biodiversity Variables. *Science* 339 (6117): 277–278.
1001 <https://doi.org/10.1126/science.1229931>.
1002

1003 Pielou, E. (1975) *Ecological Diversity*, Wiley & Sons. ed. New York.
1004

1005 Pielou, E.C. (1969) *An introduction to mathematical ecology*. New York, USA, Wiley-
1006 *Inter science*.
1007

1008 Reynolds, C. (2006) *The ecology of phytoplankton*. Cambridge University Press, UK, 535
1009 pp.
1010

1011 Rodríguez-Ramos, T., Dornelas, M., Marañón, E. and Cermeño, P. (2014) Conventional
1012 sampling methods severely underestimate phytoplankton species richness. *Journal of*
1013 *Plankton Research*: 36, 334-343.
1014

1015 Rombouts, I., Beaugrand, G., Artigas, L.F., Dauvin, J.-C., Gevaert, F., Goberville, E.,
1016 Kopp, D., Lefebvre, S., Luczak, C., Spilmont, N., Travers-Trolet, M., Villanueva, M.C.,
1017 Kirby, R.R. (2013) Evaluating marine ecosystem health: Case studies of indicators using

1018 direct observations and modelling methods. *Ecological Indicators* 24: 353–365.
1019 <https://doi.org/10.1016/j.ecolind.2012.07.001>
1020
1021 Scheiner, S.M., Kosman, E., Presley, S.J. and Willig, M.R. (2017) The components of
1022 biodiversity, with a particular focus on phylogenetic information. *Ecology and Evolution*
1023 7: 6444-6454.
1024
1025 Shannon, C.E. and Weaver, W. (1949) *The mathematical theory of communication*.
1026 University of Illinois Press, Urbana.
1027
1028 Shephard, S., Greenstreet, S.P.R., Piet, G.J., Rindorf, A. and Dickey-Collas, M. (2015)
1029 Surveillance indicators and their use in implementation of the Marine Strategy
1030 Framework Directive. *ICES Journal of Marine Science* 72: 2269-2277.
1031
1032 Sieburth, J. M., Smetacek, V., and Lenz, J. (1978). Pelagic ecosystem structure:
1033 Heterotrophic compartments of the plankton and their relationship to plankton size
1034 fractions. *Limnology and Oceanography*, 23, 1256–1263.
1035
1036 Simpson, E.H. (1949) Measurement of diversity. *Nature* 163: 688.
1037
1038 Sommer, U., Padisak, J., Reynolds, C.S. and Juhász-Nagy, P. (1993) Hutchinson's
1039 heritage: diversity-disturbance relationship in phytoplankton. *Hydrobiologia* 249: 1-7.
1040

1041 Sommer, U., Adrian, R., De Senerpont Domis, L., Elser, J.J., Gaedke, U., Ibelings, B.,
1042 Jeppesen, E., Lürling, M., Molinero, J.C., Mooij, W.M., van Donk, E. and Winder, M.
1043 (2012) Beyond the Plankton Ecology Group (PEG) Model: Mechanisms Driving Plankton
1044 Succession. *Annual Review of Ecology, Evolution, and Systematics* 43: 429-448.
1045
1046 Sourisseau, M., Jegou, K., Lunven, M., Quere, J., Gohin, F. and Bryere, P. (2016)
1047 Distribution and dynamics of two species of Dinophyceae producing high biomass
1048 blooms over the French Atlantic Shelf. *Harmful algae* 53: 53-63.
1049
1050 Sournia, A. (1978) *Phytoplankton manual*. UNESCO, Paris.
1051 Spatharis, S. and Tsirtsis, G. (2010) Ecological quality scales based on phytoplankton for
1052 the implementation of Water Framework Directive in the Eastern Mediterranean.
1053 *Ecological Indicators* 10: 840–847.
1054
1055 Spatharis, S., Roelke, D.L., Dimitrakopoulos, P.G. and Kokkoris, G.D. (2011) Analyzing
1056 the (mis)behavior of Shannon index in eutrophication studies using field and simulated
1057 phytoplankton assemblages. *Ecological Indicators* 11: 697-703.
1058
1059 Sun, J. and Xue, B. (2016) Marine phytoplankton diversity and the impact of global
1060 climate change. *Biodiversity Science* 24(7): 739-747.
1061
1062 Tett, P., Carreira, C., Mills, D.K., van Leeuwen, S., Foden, J., Bresnan, E. and Gowen, R.J.
1063 (2008) Use of a Phytoplankton Community Index to assess the health of coastal waters.

1064 ICES Journal of Marine Science: Journal du Conseil 65: 1475–1482.
1065 <https://doi.org/10.1093/icesjms/fsn161>
1066
1067 Utermöhl, H. (1958) Zur Ver vollkommung der quantitativen phytoplankton-methodik.
1068 Mitteilung Internationale Vereinigung Fuer Theoretische unde Amgewandte Limnologie,
1069 9, 39 pp.
1070
1071 Uusitalo, L., Fleming-Lehtinen, V., Hällfors, H., Jaanus A., Hällfors, S., and London, L.
1072 (2013) A novel approach for estimating phytoplankton biodiversity – ICES Journal of
1073 Marine Science 70: 408–417.
1074
1075 Vadrucchi, M.R., Stanca, E., Mazziotti, C., Umani, S.F., Georgia, A., Moncheva, S.,
1076 Romano, A., Bucci, R., Ungaro, N. and Basset, A. (2013) Ability of phytoplankton trait
1077 sensitivity to highlight anthropogenic pressures in Mediterranean lagoons: A size spectra
1078 sensitivity index (ISS-phyto). Ecological Indicators 34: 113-125.
1079
1080 van Strien, A. J., Soldaat, L. and Gregory, R. D. (2012) Desirable mathematical properties
1081 of indicators for biodiversity change. Ecological Indicators 14: 202–208.
1082
1083 Varkitzi, I., Francé, J., Basset, A., Cozzoli, F., Stanca, E., Zervoudaki, S., Giannakourou,
1084 A., Assimakopoulou, G., Venetsanopoulou, A., Mozetič, P., Tinta, T., Skejic, S., Vidjak, O.,
1085 Cadiou, J.F. and Pagou, K. (2018) Pelagic habitats in the Mediterranean Sea: A review of
1086 Good Environmental Status (GES) determination for plankton components and

1087 identification of gaps and priority needs to improve coherence for the MSFD
1088 implementation. *Ecological Indicators* 95: 203-218.

1089

1090 Whittaker, R. H. (1977) Evolution of species diversity in land communities. *Evolutionary*
1091 *Biology* 10: 1-67.

1092

1093 Wilhm, J.L. and Dorris, T.C. (1968) Biological Parameters for Water Quality Criteria.
1094 *BioScience* 18: 477-481.

1095

1096 Zingone, A., Philips, E.J. and Harrison, P.J. (2010) Multiscale Variability of Twenty-Two
1097 Coastal Phytoplankton Time Series: a Global Scale Comparison. *Estuaries and Coasts* 33:
1098 224. <https://doi.org/10.1007/s12237-009-9261-x>

1099

1100 Zettler, M.L., Darr, A., Labrenz, M., Sagert, S., Selig, U., Siebert, U. and Stybel, N. (2017)
1101 Chapter 14: « Biological Indicators » in *Biological Oceanography of the Baltic Sea*; eds.
1102 Snoeijis-Leijonmalm, P., Schubert, H., Radziejewska, T., Springer, Amsterdam.

1103

Tables and figures

Figure 1. Map of the study area and the three sampling sites used for the analyses.

Table 1. Correlation coefficients (Bravais-Pearson, r^2) between diversity indices. Values larger than 0.8 and with a significance level of $p < 0.05$ are in bold. The grey shading indicates that the indices are from the same group: richness (light grey), dominance (white) or evenness (dark grey). The correlation coefficients presented here are from SOMLIT-Astan sampling site only.

Table 2. Correlation coefficients (Bravais-Pearson, r^2) between the Principal Components (PC) that were calculated from monthly mean data and seasonal variability (calculated by the coefficient of variation (COV)) and diversity indices for the SOMLIT-Astan sampling site. Statistically significant values ($p < 0.001$) are indicated with an asterisk (*).

Table 3. List of diversity indices and results' summary of the performance analyses based on three criteria: (1) ecological relevance, (2) mathematical redundancy and (3) link with environmental factors. For the criterion (2), the indices with the same number of C 's are significantly interrelated. The indices in bold are retained for the calculation of the PH3 indicator.

Figure 2a. Contributions of environmental descriptors (monthly means) of the SOMLIT-Astan dataset to the space of the first two principal components. The environmental descriptors included temperature ($^{\circ}\text{C}$), salinity (psu), ammonia (NH_4 , $\mu\text{mol L}^{-1}$), nitrate (NO_3 , $\mu\text{mol L}^{-1}$), nitrites (NO_2 , $\mu\text{mol L}^{-1}$), silicate (SiOH_4 , $\mu\text{mol L}^{-1}$), phosphates (PO_4 , $\mu\text{mol L}^{-1}$), and

oxygen (mL L⁻¹). The circle of equilibrium descriptor contribution was drawn at $\sqrt{2/p} = 0.50$ where $p = 8$ descriptors.

Figure 2b. Contributions of environmental descriptors (seasonal variability) of the SOMLIT-Astan dataset to the space of the first two principal components. The seasonal variability was calculated by the coefficient of variation (COV). The environmental descriptors included temperature (°C), salinity (psu), nitrate (NO₃, μmol L⁻¹), nitrites (NO₂, μmol L⁻¹), silicate (SiOH₄, μmol L⁻¹), phosphates (PO₄, μmol L⁻¹), and oxygen (mL L⁻¹). The circle of equilibrium descriptor contribution was drawn at $\sqrt{2/p} = 0.53$ where $p = 7$ descriptors.

Figure 3. Contourplots of genus richness (Menhinick (D); top), dominance (Hulbert (δ); middle) and evenness (Pielou (J'); bottom) indices in time (monthly values) for the station SOMLIT-Astan (left), Le Croisic (middle) and Ouest Loscolo (right).

Figure 4. Local Contribution to Beta Diversity (LCBD) in time (annual values) for the stations Le Croisic and Ouest Loscolo. Statistically significant values ($p < 0.05$) were found in 2007 and 2014 for the station Le Croisic and in 2008 for the station Ouest Loscolo in 2008.

Figure 5. Temporal distribution of abundances (cells L⁻¹; monthly values) for the five most abundant genera (or species if only one is species identified in the genus) for (a) the station Le Croisic in 2007, (b) the station Le Croisic in 2014, and (c) the station Ouest Loscolo in 2008. The values of the Important Value Index (IVI; expressed as a percentage) for each species/genus are between brackets.

Supplementary material

Figure S1. Species accumulation curves showing the cumulative sum of the number of species along the sampling years for (a) SOMLIT-Astan, (b) Ouest Loscolo and (c) Le Croisic.

Table S1. Correlation coefficients (Bravais-Pearson, r^2) between diversity indices for (a) Ouest Loscolo and (b) Le Croisic. Values larger than 0.8 and with a significance level of $p < 0.05$ are in bold. The grey shading indicates that the indices are from the same group: richness (light grey), dominance (white) or evenness (dark grey).

Figure S2. Boxplots of annual values of genus richness (Menhinick's Index (D ; left)) and dominance (Hulburt's Index (δ ; right)) for (a) SOMLIT-Astan, (b) Le Croisic and (c) Ouest Loscolo.




Figure 1





Figure 2




Figure 3

year




Figure 4




Figure 5a




Figure 5b




Figure 5c

	S	d	D	λ	H'	BP	H_B	δ	J'
Richness (S)	-								
Margalef (d)	0,87	-							
Mehninick (D)	-0,29	0,18	-						
Simpson (λ)	-0,13	0,25	0,70	-					
Shannon (H')	-0,09	0,34	0,80	0,96	-				
Berger-Parker (BP)	-0,14	0,24	0,71	0,97	0,95	-			
Brillouin (H_B)	0,57	0,12	-0,90	-0,69	-0,75	-0,68	-		
Hulburt (δ)	-0,17	0,25	0,80	0,92	0,97	0,94	-0,74	-	
Pielou's evenness (J')	-0,12	0,22	0,53	0,69	0,73	0,69	-0,52	0,73	-

Table 1

Diversity index	Reference	Ecological relevance	Mathematical redundancy	Link with hydrology
Richness (S)	Whittaker 1972	number of taxa	C	-0.63*
Margalef's (d)	Margalef 1951	number of taxa	C	-0.50
Menhinick (D)	Menhinick 1964	number of taxa	CC	0.72**
Simpson (λ)	Simpson 1949	dominance	CCC	-0.42
Shannon (H')	Shannon and Weaver 1949	dominance	CCC	-0.47
Berger-Parker (BP)	Berger and Parker 1970	dominance	CCC	-0.37
Brillouin (HB)	Brillouin 1956	dominance	CC	-0.74**
Hulburt (δ)	Hulburt 1963	dominance	CCC	-0.55
Pielou's evenness (J')	Pielou 1969	heterogeneity	CCCC	0.72
Geometric means (G_j)	Buckland et al. 2011	volume	N/A	N/A

Table 2

Diversity index	Monthly means		Seasonal variability	
	PC1	PC2	PC1	PC2
Richness (S)	-0,27	0,13	-0,63	-0,35
Margalef's (d)	-0,1	0,04	-0,5	-0,33
Menhinick (D)	0,31	-0,23	0,72*	0,36
Simpson (λ)	0,14	-0,06	-0,42	-0,07
Shannon (H')	0,13	0,06	-0,48	-0,26
Berger-Parker (BP)	0,12	-0,07	-0,37	0,09
Brillouin (HB)	-0,42	0,22	-0,74*	-0,42
Hulburt (δ)	0,13	-0,08	-0,55	-0,45
Pielou's evenness (J')	0,2	-0,08	0,72	0,20

Table 3