



HAL
open science

HLA-G expression during hookworm infection in pregnant women

Euripide Avokpaho, Tania C. D'almeida, Ibrahim Sadissou, Léonidas Tokplonou, Rafiou Adamou, Paulin Sonon, Jacqueline Milet, Gilles Cottrell, Amandine Mondière, Achille Massougbodji, et al.

► **To cite this version:**

Euripide Avokpaho, Tania C. D'almeida, Ibrahim Sadissou, Léonidas Tokplonou, Rafiou Adamou, et al.. HLA-G expression during hookworm infection in pregnant women. *Acta Tropica*, 2019, 196, pp.52 - 59. 10.1016/j.actatropica.2019.04.026 . hal-03485711

HAL Id: hal-03485711

<https://hal.science/hal-03485711v1>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **HLA-G expression during hookworm infection in pregnant women**

2 Euripide Avokpaho¹, Tania C. d'Almeida¹, Ibrahim Sadissou^{1,2,3}, Léonidas Tokplonou^{1,2},
3 Rafiou Adamou^{1,2}, Paulin Sonon^{1,2,3}, Jacqueline Milet¹, Gilles Cottrell¹, Amandine Mondière¹,
4 Achille Massougbodji², Kabirou Moutairou², Eduardo A. Donadi³, Celso Teixeira Mendes
5 Junior⁴, Benoit Favier^{5,6}, Edgardo Carosella^{5,6}, Philippe Moreau^{5,6*}, Nathalie Rouas-Freiss^{5,6*},
6 André Garcia^{1*} and David Courtin^{1*}

7 ¹ MERIT IRD, Université Paris 5, Sorbonne Paris Cité, Paris

8 ² Université d'Abomey-Calavi, Cotonou, Bénin

9 ³ Division of Clinical Immunology, School of Medicine of Ribeirão Preto, University of São
10 Paulo, Brazil

11 ⁴ Department of Chemistry, Faculty of Philosophy, Sciences and Letters of Ribeirão Preto,
12 University of São Paulo, Brazil

13 ⁵ CEA, Institut des Maladies Emergentes et des Thérapies Innovantes (IMETI), Service de
14 Recherche en Hémato-Immunologie (SRHI), Hôpital Saint-Louis, IUH, Paris, France ;

15 ⁶ Université Paris Diderot, Sorbonne Paris Cite, IUH, Hôpital Saint-Louis, UMRE5, IUH,
16 Paris, France

17 * Joint senior authors

18

19 **Introduction:** HLA-G plays a key role on immune tolerance. Pathogens can induce soluble
20 HLA-G (sHLA-G) production to down-regulate the host immune response, creating a
21 tolerogenic environment favorable for their dissemination. To our knowledge, no study has
22 yet been conducted to assess the relationship between sHLA-G and geohelminth infections.

23 **Methods:** The study was conducted in Allada, Southeastern Benin, from 2011-2014. The
24 study population encompassed 400 pregnant women, included before the end of the 28th
25 week of gestation and followed-up until delivery. At two antenatal care visits and at delivery,
26 stool and blood samples were collected. Helminths were diagnosed by means of the Kato-
27 Katz concentration technique. We used quantile regression to analyze the association
28 between helminth infections and sHLA-G levels during pregnancy.

29 **Results:** sHLA-G levels gradually increased during pregnancy and reached maximal levels
30 at delivery. Prevalence of helminth infections was low, with a majority of hookworm
31 infections. We found significantly more hookworm-infected women above the 80th quantile
32 (Q80) of the distribution of the mean sHLA-G level ($p < 0.03$, multivariate quantile regression).
33 Considering only women above the Q80 percentile, the mean sHLA-G level was significantly
34 higher in hookworm-infected compared to uninfected women ($p = 0.04$).

35 **Conclusion:** High levels of sHLA-G were associated with hookworm infection in pregnant
36 women. This result is consistent with the potential involvement of sHLA-G in immune
37 tolerance induced by helminths during pregnancy.

38

39 **1. Introduction**

40 Helminths, and particularly geohelminths, infect 1.45 billion people worldwide (Pullan et al.,
41 2014). The prevalence of geohelminth infection during pregnancy in endemic African areas
42 could reach more than 70% (van Eijk et al., 2009). Children born to infected mothers are
43 reported to be more susceptible to helminth infection and to display weaker responses to
44 vaccines (Cooper et al., 1998; Elias et al., 2005; Malhotra et al., 1999; Malhotra et al., 2003;
45 Mehta et al., 2012; Sabin et al., 1996). These last observations could be explained by the
46 phenomenon of immune tolerance. The latter was also suspected in children exposed to
47 malaria *in utero* (Malhotra et al., 2009). Lymphocyte responses to parasite antigens and
48 detection of specific immunoglobulins in cord blood, consistent with *in utero* sensitization,
49 have been described for several helminth species including ascariasis and hookworm
50 (Guadalupe et al., 2009; Larocque et al., 2006; Pit et al., 2000). Considering that the human
51 leukocyte antigen-G (HLA-G) is a well-recognized immune checkpoint molecule, the study of
52 the potential involvement of this molecule in geohelminth infections is of interest.

53 Human Leukocyte Antigen-G (HLA-G) is a non-classical HLA class I antigen that differs from
54 classical class I molecules by its restricted tissue distribution, diversity of protein isoforms
55 and limited gene polymorphism (Donadi et al., 2011). Seven isoforms of the protein are
56 generated by alternative splicing, four are membrane bound (HLA-G1-4) and three are
57 soluble (HLA-G5-7 of which the shed HLA-G1 and the soluble HLA-G5 are usually measured
58 in plasma samples). HLA-G further differs from classical class I molecules through its
59 immune regulatory functions, which is of paramount importance in maternal-fetal immune
60 tolerance by inhibiting the cytotoxic effect of maternal NK cells (Moreau et al., 1998; Rouas-
61 Freiss et al., 1997). A lack or a failure of HLA-G expression may lead to a breach of fetal-
62 maternal tolerance and consequent fetus rejection (Cecati et al., 2011; Papamitsou et al.,
63 2014). Modulation of HLA-G expression has also been described in tumor cell development,
64 allogenic organ engraftment (Carosella et al., 2015) and viral infections (Amiot et al., 2014b).
65 It has been suggested that high HLA-G levels induce tolerance and contribute to immune

66 evasion by tumor cells (Morandi et al., 2016; Pistoia et al., 2007) or viruses (Amiot et al.,
67 2014a; Amiot et al., 2014b).

68 In tropical areas, parasitic infections during pregnancy modify the immune development of
69 the fetus (Dauby et al., 2012). Such infections affect the susceptibility of young children to
70 infectious pathogens, their response to vaccines, and the development of allergic disorders
71 such as asthma and rhinitis (Thornton et al., 2010). Placental *Plasmodium falciparum*
72 infection is associated with susceptibility to malaria during infancy (Bouaziz et al., 2017; Le
73 Hesran et al., 1997; Le Port et al., 2011; Mutabingwa et al., 2005; Schwarz et al., 2008) and it
74 has been hypothesized that immune tolerance may be responsible for the shortening of time
75 to the first occurrence of malaria infection, but a complete explanation of the physiopathology
76 involved has yet to be established (Moya-Alvarez et al., 2014). Furthermore, placental
77 malaria is also associated with non-malarial infections, suggesting that immune tolerance
78 may go beyond malaria-specific immunity (Rachas et al., 2012). The association of HLA-G
79 with the susceptibility to malaria during infancy and with low birth weight could reflect the
80 potential involvement of sHLA-G in immune tolerance mediated by placental malaria
81 (d'Almeida et al., 2016; d'Almeida et al., 2017; Sadissou et al., 2014).

82 Considering that the expression of HLA-G may modulate viral and parasitic infections
83 (Sabbagh et al., 2018), and considering that HLA-G plays an important role on pregnancy, in
84 the present study we investigated the potential association between soluble HLA-G levels
85 and hookworm infections during pregnancy.

86 **2. Material and methods**

87 **2.1. Study area, study design and follow-up**

88 Our study took place in Allada, a town located 56 km North Cotonou, the economic capital of
89 Benin. The study participants were residents of 21 villages surrounding two primary health
90 facilities: Sékou and Attogon. This study is part of a larger one encompassing 1,183 human
91 immunodeficiency virus (HIV)-negative pregnant women participating in the randomized

92 Malaria in Pregnancy Preventive Alternative Drugs ([http://clinicaltrials.gov/ct2/show/](http://clinicaltrials.gov/ct2/show/NCT00811421)
93 NCT00811421) trial, designed to compare intermittent preventive malarial treatment (IPTp)
94 with either sulfadoxine-pyrimethamine or mefloquine. The study protocol has been described
95 in detail elsewhere (Accrombessi et al., 2015; Ouedraogo et al., 2012). The first 400 children
96 of these women were followed-up to investigate the impact of sociodemographic, economic,
97 environmental and biologic factors such as the parasitological status (geohelminth and *P.*
98 *falciparum* carriers) during pregnancy on infant health during the two years of life in southern
99 Benin (Accrombessi et al., 2015; d'Almeida et al., 2017). The 400 mothers included in the
100 present study are the mothers of these 400 children. Women were included before a
101 gestational age (GA) of 28 weeks, and two doses of IPTp were administered at antenatal
102 visits (ANV). The second dose was given at least 1 month after the first one. During the first
103 ANV, a total dose of 600 mg of mebendazole was given to the women to be taken at home
104 (100 mg two times per day for 3 days) according to the guidelines of the Beninese Ministry of
105 Health. Between antenatal visits, women were encouraged to attend the health facility for
106 any health complaint. At inclusion, a questionnaire was applied by the midwife to obtain data
107 regarding socio-demographic characteristics, gynecological and obstetrical history. During
108 ANV1 and ANV2 (for IPTp1 and IPTp2, respectively) and at delivery, blood and stool
109 samples were collected before the intake of IPTp, to allow plasma sHLA-G measurements as
110 well as malaria and geohelminth diagnoses. At delivery, a placental blood smear was
111 collected to detect placental malaria. Malarial infection was defined as the presence of
112 asexual *P. falciparum* parasites in a blood smear. All the medications prescribed during the
113 follow-up were free of charge.

114 **Soluble HLA-G quantification**

115 Soluble HLA-G (sHLA-G) was quantified using MEM-G/9 antibody, which recognizes
116 the most abundant soluble isoforms (shed sHLA-G1 and intron 4 containing secreted HLA-
117 G5), and an anti-human β 2-microglobulin antibody, respectively, as capture and detection
118 antibodies (Menier et al., 2003; Rebmann et al., 2005). Microtitration plates were coated with

119 10 µg/mL MEM-G/9 mouse-anti-human HLA-G mAb (Exbio, Praha, Czech Republic) and
120 incubated overnight at 4°C. Plates were saturated with 300 µL ready to use diluent buffer
121 (DAKO, Carpinteria, CA, USA) for 2 hours. Plasma samples were diluted ($\frac{1}{2}$) in diluent
122 buffer, tested in duplicate and incubated for 2 hours. After incubation and washing,
123 monoclonal anti-human beta2-microglobulin antibody (DAKO, Glostrup, Denmark), which
124 recognizes the immobilized antibody sHLA-G complex, was added to the wells and incubated
125 for 1 hour. The plates were then incubated for 1 hour with 100 µL (1:200) envision buffer +
126 system HRP (DAKO, Carpinteria, CA, USA) to obtain anti-β2-microglobulin-horseradish
127 peroxidase complex to improve the efficiency of the reaction. All incubation steps were
128 performed at room temperature and followed by four washes using washing buffer (H₂O,
129 PBS 1X, 0.1% Tween 20). The plates were incubated for 30 min with substrate
130 (Tetramethylbenzidine super sensitive-TMB, Sigma Aldrich, Saint Louis, MO, USA) and
131 absorbance was measured at 450 nm after adding HCL (1 N). Total sHLA-G levels were
132 determined from a five-point standard calibration curve using dilutions (12.5- 200 ng/mL) of
133 HLA-G5 purified from M8-HLA-G5 cell line culture supernatants. Results are presented in
134 ng/mL.

135 **Helminth identification**

136 Women were invited to provide a sample of their fresh morning stool put in a 120 mL plastic
137 container that was distributed beforehand. Stool samples were transferred to the health
138 centre laboratory and were analysed at the same day, using standard 41.7 mg plastic
139 templates, and thick smears were examined under a light microscope by experienced
140 laboratory technicians. Microscopical identification of *Ascaris lumbricoides* and *Trichuris*
141 *trichiura* eggs is easy recognizable using the Kato-Katz method; however, *Ancylostoma*
142 *duodenale* and *Necator americanus* eggs cannot be distinguished. For this reason, we
143 included both *Ancylostoma duodenale* and *Necator americanus* species in a same hookworm
144 group.

145 **Ethics**

146 The protocol received approval from the Comité Consultatif de Déontologie et d’Ethique of
147 the Institut de Recherche pour le Développement (IRD, France) and from the Ethics
148 Committee of the Faculté des Sciences de la Santé of Cotonou
149 (N°43/11/2010/CE/FSS/UAC). Before inclusion, the protocol of the study was explained in the
150 relevant local languages and written consent was obtained. Women who could not read and
151 write provided thumbprints to confirm their agreement to participate in the study after a nurse
152 had explained the study.

153 **Definition and characteristics of the study variables**

154 **Soluble HLA-G levels**

155 The outcome variable is the level of sHLA-G, used as a quantitative variable after a
156 logarithmic transformation to take into account the skewness of the distribution.

157 **Variable of interest: Helminth infections**

158 Both hookworms (*Ancylostoma duodenale*, *Necator americanus*) and round worms (*Ascaris*
159 *lumbricoides* and *Trichuris trichiura*) were detected in faecal samples. However, due to the
160 very low prevalence of both *Ascaris lumbricoides* and *Trichuris trichiura*, our analyses
161 focused on hookworm infection during pregnancy, by creating a variable coded 1 if at least
162 one hookworm egg was detected during follow-up.

163 **Covariates**

164 Information on the following variables was collected: gravidity, IPTp regimen, placental
165 malaria, inclusion health center, ethnicity, marital status, occupation, education level, latrine
166 use at home and “wealth score”. The wealth score was computed using four socio-economic
167 items which were theoretically associated with social welfare: motorbike, television,
168 electricity, and fridge. Each item could take the value 0 or 1 (yes = 1, no = 0), and the score
169 was obtained by summing the values of each item.

170 **Statistical analysis**

171 Categorical variables were described using proportions and 95% confidence interval and the
172 continuous variables were described by the median and interquartile range. To compare
173 proportions, we used the Chi-square, Fisher T and Cuzick trend tests. Continuous variables
174 were compared by means of Student T test, Wilcoxon-Mann-Witney rank-sum test and
175 analysis of variance (ANOVA). We checked for normality of distribution using
176 Skewness/Kurtosis normality test and compared the variances with Breusch-Pagan/Cook-
177 Weisberg heteroscedasticity test. A p-value less than 5% was considered as significant.

178 Classical methods for repeated measurements (mixed models) were not used to analyse
179 sHLA-G level at antenatal visits because each antenatal visit had its particular characteristics
180 and cannot be considered as a repeated measurement of the same variable under the same
181 conditions. Measurements at the first antenatal visit were done before any intervention.
182 Medication was given at ANV1 and ANV2 (IPTp for malaria during the two first antenatal
183 visits and mebendazole during the first one). Furthermore, the third measurement occurred at
184 delivery. Consequently, to create a unique variable accounting for the level of sHLA-G during
185 pregnancy, a mean sHLA-G (MsHLA-G) level was computed. To take into account the fact
186 that not all women were present at each measurement time, and that sHLA-G levels
187 increased significantly at delivery, the MsHLA-G was computed for each woman as the mean
188 of her visit-adjusted sHLA-G level. The visit-adjusted sHLA-G levels were obtained by
189 subtracting the mean sHLA-G of the population at each visit. The association between
190 MsHLA-G and hookworm infection was analyzed by means of quantile regression that takes
191 into account the heteroscedasticity of the distribution (Koenker, 2005; Sims, 1996). Univariate
192 and multivariate analyses were done and variables with p-value less than 0.2 in univariate
193 analysis were considered in the multivariate model. The variable of interest was “at least one
194 hookworm infection during the follow-up”. Hookworm and “inclusion health facility” were
195 included in the multivariate analysis. The other covariates considered were: gravidity,
196 regimen of intermittent preventive treatment, placental malaria, age, ethnicity, marital status,
197 occupation, education, latrines use at home and wealth score. Starting from the initial model

198 we proceeded using a step-by-step backward regression. Only the variables exhibiting p-
199 value<0.05 were maintained in the final model. All analyses were performed using Stata12
200 software (StatCorp LP, College Station, TX, USA).

201 **Results**

202 **Study population**

203 Four hundred pregnant women were included in the study. Their median age was 25 years
204 (range: 18-42; Interquartile range [IQR]: 21-30). The median gestational age (GA) at inclusion
205 was 24 weeks (range: 13-28; IQR: 20-26). The median wealth score based on 4 variables
206 (ownership of motorbike, electricity, television, and refrigerator) was 1, (range: 0-4, IQR: 1-2).
207 The socio-demographic and economic characteristics of the participants are summarized in
208 Table 1.

209 **Helminth infections**

210 The results of parasitic coprology are shown in Figure 1. Stool samples were collected from
211 395 (98.75%) participants at ANV1, 387 (96.75%) at ANV2 and 355 (88.75%) at delivery,
212 giving a total of 1137 stool samples collected. Among the 1137 stool samples, helminth eggs
213 were detected in 88 stool samples (7.7%). The sample positivity rate to helminth infections
214 was 9% at ANV1, 8.8% at ANV2 and 5% at delivery. Hookworm eggs (*Ancylostoma*
215 *duodenale* and *Necator americanus*) were found in 74 samples (27 infections at ANV1, 31 at
216 ANV2 and 16 at delivery), 10 samples contained eggs of *Ascaris lumbricoides* (1 at ANV1, 6
217 at ANV2 and 3 at delivery) and eggs of *Trichuris trichiura* were found in 3 stool samples (1
218 sample positive at ANV1, 1 at ANV2 and 1 at delivery, all from different participants).

219 The prevalence of helminth infection (at least one geohelminth in the stools during the whole
220 follow-up) in the study population was 18.5% (74/400). The majority of the helminth infections
221 detected in the stools were hookworm infections (86.5% (64/74)), while *Ascaris lumbricoides*
222 and *Trichuris trichiura* eggs were respectively detected in 13.5% (10/74) and 4% (3/74) of the

223 positive stool samples. Two participants were co-infected with both hookworm and *Ascaris*
224 *lumbricoides*, one diagnosed at ANV2 and the other diagnosed at delivery.

225 The distribution of hookworm infection according to the different characteristics of the study
226 population is presented in Table 2. The prevalence of hookworm infection was significantly
227 higher in Sekou health facility and in the Aizo ethnic group ($p=0.02$ in both cases, Chi^2). The
228 possession of latrines was associated with a significantly lower prevalence ($p=0.001$, Chi^2) as
229 was a high wealth score ($p=0.024$, Chi^2).

230 **Soluble HLA-G level**

231 Soluble HLA-G measurement was available on three occasions from 85% of mothers and
232 98% were sampled at least twice. sHLA-G levels were stable over the two first visits (mean
233 sHLA-G level at ANV1 = 10.09 ng/ml and mean sHLA-G level at ANV2 = 10.57 ng/ μl) but
234 increased significantly at delivery (mean sHLA-G level = 17.26 ng/ μl ($p<10^{-3}$, ANOVA)).

235 **Relation between hookworm infection and adjusted mean of sHLA-G.**

236 The distribution of adjusted means of sHLA-G (MsHLA-G) is presented in Figure 2. Despite
237 the log-transformation, both the hypotheses of normality ($p<0.0001$, Shapiro Wilk normality
238 test) and of homoscedasticity ($p=0.02$; Breusch-Pagan/Cook-Weisberg heteroscedasticity
239 test) were significantly rejected. To deal with both the skewed distribution and
240 heteroscedasticity, a quantile (5th, 10th, up to 95th quantile) regression (Chamberlain, 1994)
241 was used to analyze the effect of hookworm infection and of other covariates (inclusion
242 health facility, latrines use at home, wealth score, gravidity, regimen of intermittent preventive
243 treatment, placental malaria, age, ethnicity, marital status, occupation, education) on the
244 different quantiles of mean sHLA-G levels.

245 In univariate analysis, inclusion health facility had a significant effect on the distribution of
246 sHLA-G with significantly fewer women above the Q50 of the distribution of the mean level of
247 sHLA-G included at Sekou health center (Table 3). No significant effect of the others
248 covariates was detected on the distribution of sHLA-G. As hookworm infection is the variable

249 of interest, the multivariate model of regression included the inclusion health center and
250 hookworm infection. The effect of inclusion health facility on the distribution of sHLA-G
251 remained significant in multivariate analysis and a significant effect of hookworm infection
252 was observed ($p < 0.03$, multiple quantile regression). Significantly more hookworm-infected
253 women were found above the Q80 percentile of sHLA-G distribution (Table 3 and [Figure 3](#)).
254 Considering only women above the Q80 percentile, the geometric mean level of sHLA-G was
255 significantly higher in hookworm-infected than in uninfected women (1.57 vs 1.41
256 respectively; $p = 0.04$, Student T test).

257 **Discussion**

258 The main results of the present work show a relatively low frequency of helminth infection
259 during pregnancy of Beninese women; however, we observed an association between the
260 plasma levels of the tolerogenic molecule HLA-G and hookworm infection.

261 Prevalence of hookworm infections in our study decreased from 9% at ANV1 to 5% at
262 delivery. The detection of hookworms was performed with the inexpensive and field-
263 applicable Kato-Katz method however the low sensibility of the Kato-Katz method probably
264 underestimates the intestinal helminth infection prevalence in our study population.

265 This finding could be explained by the intake of one dose of mebendazole (600 mg) during
266 antenatal visits after 16th week of gestation, as recommended by the Beninese National
267 Health Program. Anthelmintic treatment during pregnancy aims to fight against maternal
268 anemia (Ouedraogo et al., 2012), which has an important impact on neonatal and infant
269 health (Koura et al., 2012). Helminth infections during pregnancy are also associated with
270 increased susceptibility to geohelminth infections in the children (Mehta et al., 2012) and to
271 weaker anti-vaccine responses (Labeaud et al., 2009). The mechanisms by which maternal
272 helminth infection can influence immune development of the fetus are partially understood
273 (Mpairwe et al., 2014). However, epidemiological evidence suggests that *in utero*
274 sensitization results in down-regulation of immune responses in offspring (Dauby et al.,

275 2012), leading to a phenomenon of immune tolerance, in which HLA-G could play a crucial
276 role.

277 In the multivariate model of regression, sHLA-G levels were associated with health center
278 and hookworm infection. Women included at Sekou health center had lower level of sHLA-G
279 than those from Attogon health center. Significant differences in term of ethnicity group
280 distribution and wealth score were observed between both health centers ($p < 0.001$ and
281 $p < 0.001$ respectively, Chi-Square test). Women from Fon ethnicity group were more
282 represented at Attogon than at Sekou (57 % versus 9 %) while women from Aizo ethnicity
283 group were more frequent at Sekou than Attogon (82 % versus 31 %). Differences in
284 behaviour and/or in genetic mechanism influencing regulation of HLA-G expression between
285 both ethnicity groups could partly explain the modulation of sHLA-G observed between
286 health centers. It is important to emphasize that Sekou is a semi-rural village while Attogon is
287 more rural therefore difference of HLA-G expression could also be related to environmental
288 factors, such as the prevalence of other pathogens that could also influence the expression
289 of sHLA-G. The elevated risk of exposure to mosquito bites was previously associated with
290 an increased expression of sHLA-G (Sadissou et al., 2014). In the present work, no
291 association was observed between peripheral or placental malaria infections during
292 pregnancy and a significant modulation of plasmatic sHLA-G in mothers (d'Almeida et al.,
293 2017). Difference of sHLA-G level observed between the health centers seems not directly
294 related to malaria infection.

295 Our results show that there are more hookworm infected women above the Q80 and that
296 these women have significantly higher levels of sHLA-G than not infected women from the
297 same quantiles. The use of Kato-Katz is a potential limitation in our study. Some intestinal
298 helminth infections could be failed in case of light-intensity hookworm infections and we have
299 potentially included false negatives in the group of "hookworm uninfected women". Such
300 misclassification should not significantly affect our result because a higher level of sHLA-G is
301 expected in presence of false positives in the group of women with no hookworm egg. Such
302 situation of misclassification could typically lead to a lack of statistical power, which could

303 reduce the chance to find an association between hookworm infection and sHLA-G levels.
304 The use of more sensitive tools such as the PCR or coproculture methods could reinforce the
305 result of association between hookworm infection and sHLA-G levels obtained in our study
306 and give a better insight of the hookworm prevalence in the study areas.

307 It is known that hookworm infection results firstly in a T helper 2 cell type polarization,
308 characterized by the proliferation of T cells that produce IL-4, IL-5 and IL-13, activation of
309 basophils, eosinophils, mast cells and secretion of high levels of IgE. Then, a progressive
310 and generalized immunosuppression, characterized by induction of regulatory responses
311 (regulatory T, B and dendritic cells) with increased expression of the regulatory cytokines IL-
312 10 and TGF- β , often occurs. The progressive emergence of an immunosuppressive state
313 could be a mechanism by which hookworms escape the effective anti-helminthic Th2
314 response (Borkow et al., 2000; Maizels et al., 2004). HLA-G can participate in this
315 immunoregulatory state (Carosella et al., 2015) that can lead to a tolerogenic environment.

316 HLA-G is a very complex molecule with well-recognized biological features, and its immune
317 functions have recently been reviewed (Carosella et al., 2015). HLA-G is known to inhibit the
318 function of T, NK and B cells through direct interaction with ILT2. Recently, Naji et al., (2014)
319 showed that HLA-G inhibits B cell proliferation, differentiation, and immunoglobulin secretion
320 in both T cell-dependent and T cell-independent models of B cell activation (Naji et al.,
321 2014). It has been recently shown that children born to mother with high sHLA-G levels are
322 predisposed to have high sHLA-G levels during their two first years of life (d'Almeida et al.,
323 2016). We can then hypothesise that the high levels of sHLA-G found in hookworm infected
324 women could be associated with high levels in their newborns that can be involved in the
325 immune tolerance phenomenon.

326 In tropical areas, immune tolerance is not restricted to helminth infections (Raberg, 2014).
327 Placental malaria is classically described as a risk factor for malaria in early infancy (Le
328 Hesran et al., 1997; Le Port et al., 2011; Malhotra et al., 2009; Mutabingwa et al., 2005) and
329 although not totally understood, induction of a tolerant state has also been proposed to be an
330 explanation to this susceptibility. However, we have recently demonstrated that children born

331 to mothers with placental malaria are not only more susceptible to malaria but also to non-
332 malarial fever (Rachas et al., 2012). These findings underline the fact that immune tolerance
333 could impact on immunity in a more general manner rather than simply malaria-specific
334 immunity. The involvement of HLA-G in tropical diseases has been recently shown for both
335 malaria and human African trypanosomiasis at both genetic and protein levels (Courtin et al.,
336 2013; d'Almeida et al., 2016; Garcia et al., 2013; Gineau et al., 2016; Sadissou et al., 2014).
337 Some associations between antibody responses to malarial antigens and some
338 polymorphisms of *HLA-G* strengthen the concept of a potential role of HLA-G in the control of
339 host immunity in tropical diseases (Sabbagh et al., 2013).

340 Our results also showed a significant increase of sHLA-G levels at delivery compared to the
341 sHLA-G levels quantified during antenatal visits. Similar result was previously observed in a
342 Swedish population (Rizzo et al., 2009). Rizzo and coll. found significantly higher plasmatic
343 sHLA-G level at delivery than during the 3rd trimester of pregnancy in allergic and non-
344 allergic pregnant women. Moreover the progression of labour was associated with a
345 continuous and significant increase in the sHLA-G level (Knafel et al., 2009). In Africa, the
346 accessibility of health services is often difficult in rural area and most often pregnant women
347 reach the maternity late (Thaddeus and Maine, 1994). In this context, our population study
348 could contain a greater proportion of women with advanced progression of labor. The
349 elevation of plasmatic sHLA-G could also be influence by others factors such as stress and
350 hormonal changes induced during delivery.

351 In conclusion, our results argue for the first time for the potential involvement of HLA-G in the
352 complex phenomenon of helminth-mediated immune tolerance during pregnancy.
353 Investigation of this immune tolerance might help in furthering our understanding of the
354 development of infant immune response.

355 **Acknowledgements**

356 The authors thank Dr. Adrian Luty for helpful corrections to the manuscript. This study was
357 supported by the Agence Nationale de la Recherche (ANR – TOLIMMUNPAL project) and by

358 Brazil-France Research Cooperation Program USP/COFECUB (Grant numbers Uc Me 169-
359 17). Tania C d'Almeida was partly supported by grants from « Fondation pour la Recherche
360 Médicale » and Institut de Recherche pour le Développement (IRD). PhD scholarships were
361 awarded by IRD and Ambassade de France, Cotonou to Ibrahim Sadissou, Rafiou Adamou
362 and Léonidas Tokplonou.

363

364 **References**

- 365 Accrombessi, M., Ouedraogo, S., Agbota, G.C., Gonzalez, R., Massougbdji, A., Menendez, C., Cot, M.,
366 2015. Malaria in Pregnancy Is a Predictor of Infant Haemoglobin Concentrations during the First Year
367 of Life in Benin, West Africa. *PloS one* 10, e0129510.
- 368 Amiot, L., Vu, N., Rauch, M., L'Helgoualc'h, A., Chalmel, F., Gascan, H., Turlin, B., Guyader, D.,
369 Samson, M., 2014a. Expression of HLA-G by mast cells is associated with hepatitis C virus-induced
370 liver fibrosis. *Journal of hepatology* 60, 245-252.
- 371 Amiot, L., Vu, N., Samson, M., 2014b. Immunomodulatory properties of HLA-G in infectious diseases.
372 *Journal of immunology research* 2014, 298569.
- 373 Borkow, G., Leng, Q., Weisman, Z., Stein, M., Galai, N., Kalinkovich, A., Bentwich, Z., 2000. Chronic
374 immune activation associated with intestinal helminth infections results in impaired signal
375 transduction and energy. *The Journal of clinical investigation* 106, 1053-1060.
- 376 Bouaziz, O., Courtin, D., Cottrell, G., Milet, J., Nuel, G., Garcia, A., 2018. Is placental malaria a long
377 term risk factor for mild malaria attack in infancy? Revisiting a paradigm. *Clinical infectious diseases*
378 66, 930-935.
- 379 Carosella, E.D., Rouas-Freiss, N., Tronik-Le Roux, D., Moreau, P., LeMaout, J., 2015. HLA-G: An
380 Immune Checkpoint Molecule. *Advances in immunology* 127, 33-144.
- 381 Cecati, M., Giannubilo, S.R., Emanuelli, M., Tranquilli, A.L., Saccucci, F., 2011. HLA-G and pregnancy
382 adverse outcomes. *Medical hypotheses* 76, 782-784.
- 383 Chamberlain, C.A., 1994. Quantile regression, censoring, and the structure of wages. Cambridge
384 University Press.
- 385 Cooper, P.J., Espinel, I., Paredes, W., Guderian, R.H., Nutman, T.B., 1998. Impaired tetanus-specific
386 cellular and humoral responses following tetanus vaccination in human onchocerciasis: a possible
387 role for interleukin-10. *The Journal of infectious diseases* 178, 1133-1138.
- 388 Courtin, D., Milet, J., Sabbagh, A., Massaro, J.D., Castelli, E.C., Jamonneau, V., Bucheton, B., Sese, C.,
389 Favier, B., Rouas-Freiss, N., Moreau, P., Donadi, E.A., Garcia, A., 2013. HLA-G 3' UTR-2 haplotype is
390 associated with Human African trypanosomiasis susceptibility. *Infection, genetics and evolution* 17,
391 1-7.
- 392 d'Almeida, T.C., Sadissou, I., Cottrell, G., Tahar, R., Moreau, P., Favier, B., Moutairou, K., Donadi, E.A.,
393 Massougbdji, A., Rouas-Freiss, N., Courtin, D., Garcia, A., 2016. Evolution of the levels of human
394 leukocyte antigen G (HLA-G) in Beninese infant during the first year of life in a malaria endemic area:
395 using latent class analysis. *Malaria journal* 15, 78.
- 396 d'Almeida, T.C., Sadissou, I., Milet, J., Cottrell, G., Mondiere, A., Avokpaho, E., Gineau, L., Sabbagh, A.,
397 Massougbdji, A., Moutairou, K., Donadi, E.A., Favier, B., Carosella, E., Moreau, P., Rouas-Freiss, N.,
398 Courtin, D., Garcia, A., 2017. Soluble human leukocyte antigen -G during pregnancy and infancy in
399 Benin: Mother/child resemblance and association with the risk of malaria infection and low birth
400 weight. *PloS one* 12, e0171117.
- 401 Dauby, N., Goetghebuer, T., Kollmann, T.R., Levy, J., Marchant, A., 2012. Uninfected but not
402 unaffected: chronic maternal infections during pregnancy, fetal immunity, and susceptibility to
403 postnatal infections. *The Lancet. Infectious diseases* 12, 330-340.
- 404 Donadi, E.A., Castelli, E.C., Arnaiz-Villena, A., Roger, M., Rey, D., Moreau, P., 2011. Implications of the
405 polymorphism of HLA-G on its function, regulation, evolution and disease association. *Cellular and*
406 *molecular life sciences : CMLS* 68, 369-395.
- 407 Elias, D., Akuffo, H., Pawlowski, A., Haile, M., Schon, T., Britton, S., 2005. *Schistosoma mansoni*
408 infection reduces the protective efficacy of BCG vaccination against virulent *Mycobacterium*
409 *tuberculosis*. *Vaccine* 23, 1326-1334.
- 410 Garcia, A., Milet, J., Courtin, D., Sabbagh, A., Massaro, J.D., Castelli, E.C., Migot-Nabias, F., Favier, B.,
411 Rouas-Freiss, N., Donadi, E.A., Moreau, P., 2013. Association of HLA-G 3'UTR polymorphisms with
412 response to malaria infection: a first insight. *Infection, genetics and evolution* 16, 263-269.

413 Gineau, L., Courtin, D., Camara, M., Ilboudo, H., Jamonneau, V., Dias, F.C., Tokplonou, L., Milet, J.,
414 Mendonca, P.B., Castelli, E.C., Camara, O., Favier, B., Rouas-Freiss, N., Moreau, P., Donadi, E.A.,
415 Bucheton, B., Sabbagh, A., Garcia, A., 2016. Human Leukocyte Antigen-G: A Promising Prognostic
416 Marker of Disease Progression to Improve the Control of Human African Trypanosomiasis. *Clinical*
417 *infectious diseases* 63, 1189-1197.

418 Guadalupe, I., Mitre, E., Benitez, S., Chico, M.E., Nutman, T.B., Cooper, P.J., 2009. Evidence for in
419 utero sensitization to *Ascaris lumbricoides* in newborns of mothers with ascariasis. *The Journal of*
420 *infectious diseases* 199, 1846-1850.

421 Knafel, A., Basta, P., Pitynski, K., Mach, P., Bednarek, W., Klimek, M., Zietek, J., Zajac, K., Danciewicz,
422 L., Iwaniec, M., Basta, A., Wicherek, L., 2009. Soluble HLA-G changes in maternal blood serum during
423 the progression of labor. *Neuroendocrinol Lett* 30, 67-73.

424 Koenker, R., 2005. *Quantile Regression*. Cambridge University Press,
425 <https://doi.org/10.1017/CBO9780511754098>.

426 Koura, K.G., Ouedraogo, S., Cottrell, G., Le Port, A., Massougbdji, A., Garcia, A., 2012. Maternal
427 anaemia at delivery and haemoglobin evolution in children during their first 18 months of life using
428 latent class analysis. *PLoS one* 7, e50136.

429 Labeaud, A.D., Malhotra, I., King, M.J., King, C.L., King, C.H., 2009. Do antenatal parasite infections
430 devalue childhood vaccination? *PLoS neglected tropical diseases* 3, e442.

431 Larocque, R., Casapia, M., Gotuzzo, E., MacLean, J.D., Soto, J.C., Rahme, E., Gyorkos, T.W., 2006. A
432 double-blind randomized controlled trial of antenatal mebendazole to reduce low birthweight in a
433 hookworm-endemic area of Peru. *Tropical medicine & international health : TM & IH* 11, 1485-1495.

434 Le Hesran, J.Y., Cot, M., Personne, P., Fievet, N., Dubois, B., Beyeme, M., Boudin, C., Deloron, P.,
435 1997. Maternal placental infection with *Plasmodium falciparum* and malaria morbidity during the
436 first 2 years of life. *American journal of epidemiology* 146, 826-831.

437 Le Port, A., Watier, L., Cottrell, G., Ouedraogo, S., Dechavanne, C., Pierrat, C., Rachas, A., Bouscaillou,
438 J., Bouraima, A., Massougbdji, A., Fayomi, B., Thiebaut, A., Chandre, F., Migot-Nabias, F., Martin-
439 Prevel, Y., Garcia, A., Cot, M., 2011. Infections in infants during the first 12 months of life: role of
440 placental malaria and environmental factors. *PLoS one* 6, e27516.

441 Maizels, R.M., Balic, A., Gomez-Escobar, N., Nair, M., Taylor, M.D., Allen, J.E., 2004. Helminth
442 parasites--masters of regulation. *Immunological reviews* 201, 89-116.

443 Malhotra, I., Dent, A., Mungai, P., Wamachi, A., Ouma, J.H., Narum, D.L., Muchiri, E., Tisch, D.J., King,
444 C.L., 2009. Can prenatal malaria exposure produce an immune tolerant phenotype? A prospective
445 birth cohort study in Kenya. *PLoS Med* 6, e1000116.

446 Malhotra, I., Mungai, P., Wamachi, A., Kioko, J., Ouma, J.H., Kazura, J.W., King, C.L., 1999. Helminth-
447 and *Bacillus Calmette-Guerin*-induced immunity in children sensitized in utero to filariasis and
448 schistosomiasis. *J Immunol* 162, 6843-6848.

449 Malhotra, I., Ouma, J.H., Wamachi, A., Kioko, J., Mungai, P., Njzovu, M., Kazura, J.W., King, C.L., 2003.
450 Influence of maternal filariasis on childhood infection and immunity to *Wuchereria bancrofti* in
451 Kenya. *Infection and immunity* 71, 5231-5237.

452 Mehta, R.S., Rodriguez, A., Chico, M., Guadalupe, I., Broncano, N., Sandoval, C., Tupiza, F., Mitre, E.,
453 Cooper, P.J., 2012. Maternal geohelminth infections are associated with an increased susceptibility to
454 geohelminth infection in children: a case-control study. *PLoS neglected tropical diseases* 6, e1753.

455 Menier, C., Saez, B., Horejsi, V., Martinozzi, S., Krawice-Radanne, I., Bruel, S., Le Danff, C., Reboul, M.,
456 Hilgert, I., Rabreau, M., Larrad, M.L., Pla, M., Carosella, E.D., Rouas-Freiss, N., 2003. Characterization
457 of monoclonal antibodies recognizing HLA-G or HLA-E: new tools to analyze the expression of
458 nonclassical HLA class I molecules. *Human immunology* 64, 315-326.

459 Morandi, F., Rizzo, R., Fainardi, E., Rouas-Freiss, N., Pistoia, V., 2016. Recent Advances in Our
460 Understanding of HLA-G Biology: Lessons from a Wide Spectrum of Human Diseases. *Journal of*
461 *immunology research* 2016, 4326495.

462 Moreau, P., Paul, P., Rouas-Freiss, N., Kirszenbaum, M., Dausset, J., Carosella, E.D., 1998. Molecular
463 and immunologic aspects of the nonclassical HLA class I antigen HLA-G: evidence for an important
464 role in the maternal tolerance of the fetal allograft. *Am J Reprod Immunol* 40, 136-144.

465 Moya-Alvarez, V., Abellana, R., Cot, M., 2014. Pregnancy-associated malaria and malaria in infants: an
466 old problem with present consequences. *Malaria journal* 13, 271.

467 Mpairwe, H., Tweyongyere, R., Elliott, A., 2014. Pregnancy and helminth infections. *Parasite*
468 *immunology* 36, 328-337.

469 Mutabingwa, T.K., Bolla, M.C., Li, J.L., Domingo, G.J., Li, X., Fried, M., Duffy, P.E., 2005. Maternal
470 malaria and gravidity interact to modify infant susceptibility to malaria. *PLoS medicine* 2, e407.

471 Naji, A., Menier, C., Morandi, F., Agaogue, S., Maki, G., Ferretti, E., Bruel, S., Pistoia, V., Carosella,
472 E.D., Rouas-Freiss, N., 2014. Binding of HLA-G to ITIM-bearing Ig-like transcript 2 receptor suppresses
473 B cell responses. *J Immunol* 192, 1536-1546.

474 Ouedraogo, S., Koura, G.K., Accrombessi, M.M., Bodeau-Livinec, F., Massougbdji, A., Cot, M., 2012.
475 Maternal anemia at first antenatal visit: prevalence and risk factors in a malaria-endemic area in
476 Benin. *The American journal of tropical medicine and hygiene* 87, 418-424.

477 Papamitsou, T., Toskas, A., Papadopoulou, K., Sioga, A., Lakis, S., Chatzistamatiou, M., Economou, Z.,
478 Adriopoulou, L., 2014. Immunohistochemical study of immunological markers: HLAG, CD16, CD25,
479 CD56 and CD68 in placenta tissues in recurrent pregnancy loss. *Histology and histopathology* 29,
480 1047-1055.

481 Pistoia, V., Morandi, F., Wang, X., Ferrone, S., 2007. Soluble HLA-G: Are they clinically relevant?
482 *Seminars in cancer biology* 17, 469-479.

483 Pit, D.S., Polderman, A.M., Schulz-Key, H., Soboslay, P.T., 2000. Prenatal immune priming with
484 helminth infections: parasite-specific cellular reactivity and Th1 and Th2 cytokine responses in
485 neonates. *Allergy* 55, 732-739.

486 Pullan, R.L., Smith, J.L., Jasrasaria, R., Brooker, S.J., 2014. Global numbers of infection and disease
487 burden of soil transmitted helminth infections in 2010. *Parasites & vectors* 7, 37.

488 Raberg, L., 2014. How to live with the enemy: understanding tolerance to parasites. *PLoS biology* 12,
489 e1001989.

490 Rachas, A., Le Port, A., Cottrell, G., Guerra, J., Choudat, I., Bouscaillou, J., Massougbdji, A., Garcia, A.,
491 2012. Placental malaria is associated with increased risk of nonmalaria infection during the first 18
492 months of life in a Beninese population. *Clinical infectious diseases : an official publication of the*
493 *Infectious Diseases Society of America* 55, 672-678.

494 Rebmann, V., Lemaoult, J., Rouas-Freiss, N., Carosella, E.D., Grosse-Wilde, H., 2005. Report of the
495 Wet Workshop for Quantification of Soluble HLA-G in Essen, 2004. *Human immunology* 66, 853-863.

496 Rizzo, R., Stignani, M., Amoudruz, P., Nilsson, C., Melchiorri, L., Baricordi, O., Sverremark-Ekstrom, E.,
497 2009. Allergic women have reduced sHLA-G plasma levels at delivery. *Am J Reprod Immunol* 61, 368-
498 376.

499 Rouas-Freiss, N., Goncalves, R.M., Menier, C., Dausset, J., Carosella, E.D., 1997. Direct evidence to
500 support the role of HLA-G in protecting the fetus from maternal uterine natural killer cytotoxicity.
501 *Proceedings of the National Academy of Sciences of the United States of America* 94, 11520-11525.

502 Sabbagh, A., Courtin, D., Milet, J., Massaro, J.D., Castelli, E.C., Migot-Nabias, F., Favier, B., Rouas-
503 Freiss, N., Moreau, P., Garcia, A., Donadi, E.A., 2013. Association of HLA-G 3' untranslated region
504 polymorphisms with antibody response against *Plasmodium falciparum* antigens: preliminary results.
505 *Tissue antigens* 82, 53-58.

506 Sabbagh, A., Sonon, P., Sadissou, I., Mendes-Junior, C.T., Garcia, A., Donadi, E.A., Courtin, D., 2018.
507 The role of HLA-G in parasitic diseases. *HLA* 91, 255-270.

508 Sabin, E.A., Araujo, M.I., Carvalho, E.M., Pearce, E.J., 1996. Impairment of tetanus toxoid-specific
509 Th1-like immune responses in humans infected with *Schistosoma mansoni*. *The Journal of infectious*
510 *diseases* 173, 269-272.


511 Sadissou, I., d'Almeida, T., Cottrell, G., Luty, A., Krawice-Radanne, I., Massougbdji, A., Moreau, P.,
512 Moutairou, K., Garcia, A., Favier, B., Rouas-Freiss, N., Courtin, D., 2014. High plasma levels of HLA-G
513 are associated with low birth weight and with an increased risk of malaria in infancy. *Malaria journal*
514 13, 312.

515 Schwarz, N.G., Adegnika, A.A., Breitling, L.P., Gabor, J., Agnandji, S.T., Newman, R.D., Lell, B., Issifou,
516 S., Yazdanbakhsh, M., Luty, A.J., Kremsner, P.G., Grobusch, M.P., 2008. Placental malaria increases

517 malaria risk in the first 30 months of life. *Clinical infectious diseases : an official publication of the*
518 *Infectious Diseases Society of America* 47, 1017-1025.
519 Sims, C.A., 1996. *Advances in Econometrics*. Cambridge University Press
520 <https://doi.org/10.1017/CCOL0521444594>.
521 Thaddeus, S., Maine, D., 1994. Too Far to Walk - Maternal Mortality in Context. *Soc Sci Med* 38,
522 1091-1110.
523 Thornton, C.A., Macfarlane, T.V., Holt, P.G., 2010. The hygiene hypothesis revisited: role of materno-
524 fetal interactions. *Current allergy and asthma reports* 10, 444-452.
525 van Eijk, A.M., Lindblade, K.A., Odhiambo, F., Peterson, E., Rosen, D.H., Karanja, D., Ayisi, J.G., Shi,
526 Y.P., Adazu, K., Slutsker, L., 2009. Geohelminth Infections among pregnant women in rural western
527 Kenya; a cross-sectional study. *PLoS neglected tropical diseases* 3, e370 doi:
528 10.1371/journal.pntd.0000370.

529

Figure 1: Percentage of stool samples with helminth eggs during the pregnancy follow-up



A positive sample was defined as follow: stool sample with detection of helminth eggs by Kato-Katz method. Respectively 395, 387 and 355 stool samples were collected and examined at antenatal visit (AV)1, AV2 and at delivery.

Figure 2: Distribution of the adjusted mean of sHLA-G according to hookworm infection status of the women in the study





Figure 3: Effect of hookworm infection on the adjusted mean of sHLAG controlled for inclusion center (Adjusted coefficients of quantile regression).



*Quantiles with significant difference between coefficients of infected or uninfected women.

The curves show the regression coefficients on 20 quantiles of sHLA-G distribution during the follow-up, in women infected or non-infected with hookworms, controlled for the inclusion health facility. The graphic represents the effect of helminth infection on the quantiles of log HLA-G distribution. Significantly more hookworm-infected women were found above the Q80 percentile of sHLA-G distribution.

Table 1: Socio-demographic and economic characteristics of the participants

Variables		N (%)
Inclusion health facility	Attogon	96 (24)
	Sékou	304 (76)
Marital status	Single	9 (2.3)
	Married	391 (97.7)
Occupation	Not working	223 (55.8)
	Working	177 (29.2)
Education	Not educated	283 (70.8)
	Educated	117 (29.2)
Ethnicity	Fon	83 (20.8)
	Aizo	278 (69.5)
	Others	39 (9.7)
Gravidity	Multigravidae	337 (84.3)
	Primigravidae	63 (15.7)
Placental malaria	No	353 (88.3)
	Yes	43 (10.7)
	Unknown	4 (1)
IPTp regimen	SP	138 (34.5)
	MQ single dose	142 (35.5)
	MQ standard dose	120 (30)
Household Latrines	No	231 (57.8)
	Yes	169 (42.2)
Electricity	No	262 (65.5)
	Yes	138 (34.5)
Refrigerator	No	384 (96)
	Yes	16 (4)
Television	No	267 (66.7)
	Yes	133 (33.2)
Motorbike	No	90 (22.5)
	Yes	310 (77.5)

Table 2: Distribution of hookworm infections according to socio-demographic and economic characteristics of the study population

Variable		Hookworm infection*	
		N(%)	p-value
Inclusion			
Health facility	Attogon	8(8.3)	0.02
	Sékou	56(18.4)	
Marital status	Single	2(22.2)	0.61
	Married	62(15.9)	
Occupation	Not working	41(18.4)	0.14
	Working	23(13)	
Education	Not educated	48(17)	0.41
	Educated	16(13.7)	
Ethnicity	Fon	10(12.1)	0.02
	Aizo	53(19.1)	
	Others	1	
Gravidity	Multigravidae	54(16)	0.98
	Primigravidae	10(15.9)	
Placental malaria	No	54(15.3)	0.57
	Yes	8(18.6)	
IPTp regimen	SP	17(12.3)	0.30
	MQ single dose	24(16.9)	
	MQ standard dose	23(19.2)	
Household Latrines	No	49(21.2)	0.001
	Yes	15(8.9)	
Wealth score	<1	48(19.2)	0.024
	≥1	16(10.7)	

* At least one hookworm infection during the follow-up

Table 3: Effect of hookworm infection on 20 quantiles of sHLAG distribution during the follow-up. Crude coefficients and adjusted coefficients (controlled for inclusion center) of simultaneous quantile regression.

Quantiles	Variable	Crude β^1	95% CI	P-value	adjusted β^2	95% CI	P-value
Q10	Hookworm infection	0	-0.01; 0.01	1	0	-0.07; 0.07	1
	Sekou health facility#	0	-0.19; 0.19	1	0	-0.21; 0.21	1
Q15	Hookworm infection	-0.02	-0.31; 0.26	0.87	0	-0.18; 0.18	1
	Sekou health facility	-0.21	-0.54; 0.12	0.21	-0.21	-0.53; 0.11	0.19
Q20	Hookworm infection	-0.36	-0.76; 0.05	0.08	-0.34	-0.73; 0.05	0.09
	Sekou health facility	-0.27	-0.85; 0.31	0.36	-0.21	-0.71; 0.29	0.42
Q25	Hookworm infection	-0.55	-1.22; 0.11	0.1	-0.55	-1.15; 0.05	0.07
	Sekou health facility	-0.3	-0.91; 0.31	0.34	-0.29	-0.79; 0.22	0.27
Q30	Hookworm infection	-0.3	-1.02; 0.42	0.41	-0.2	-0.99; 0.60	0.63
	Sekou health facility	-0.4	-1.14; 0.33	0.28	-0.39	-1.07; 0.29	0.26
Q35	Hookworm infection	-0.18	-1.02; 0.66	0.67	-0.13	-0.97; 0.71	0.76
	Sekou health facility	-0.66	-1.43; 0.11	0.09	-0.69	-1.42; 0.03	0.06
Q40	Hookworm infection	-0.04	-0.93; 0.85	0.93	-0.14	-1.10; 0.81	0.77
	Sekou health facility	-0.72	-1.54; 0.11	0.09	-0.77	-1.54; 0.01	0.05
Q45	Hookworm infection	0.004	-0.87; 0.88	0.99	-0.06	-1.14; 1.01	0.91
	Sekou health facility	-0.62	-1.31; 0.08	0.08	-0.57	-1.30; 0.17	0.13
Q50	Hookworm infection	-0.1	-0.82; 0.62	0.78	-0.06	-1.03; 0.90	0.9
	Sekou health facility	-0.65	-1.28; -0.03	0.04	-0.63	-1.22; -0.04	0.04
Q55	Hookworm infection	-0.09	-0.75; 0.57	0.78	0.06	-0.81; 0.93	0.89
	Sekou health facility	-0.42	-0.83; -0.01	0.05	-0.42	-0.87; 0.02	0.06
Q60	Hookworm infection	-0.003	-0.66; 0.47	0.74	0.09	-0.56; 0.74	0.79
	Sekou health facility	-0.37	-0.68; -0.06	0.02	-0.39	-0.72; -0.06	0.02
Q65	Hookworm infection	0.2	-0.49; 0.47	0.99	0.12	-0.38; 0.62	0.64
	Sekou health facility	-0.3	-0.57; -0.03	0.03	-0.32	-0.62; -0.03	0.03
Q70	Hookworm infection	0.2	-0.22; 0.61	0.35	0.27	-0.13; 0.66	0.19
	Sekou health facility	-0.25	-0.47; -0.03	0.02	-0.26	-0.49; -0.03	0.03

Q75	Hookworm infection	0.19	-0.18; 0.56	0.31	0.26	-0.10; 0.62	0.15
	Sekou health facility	-0.17	-0.42; 0.07	0.17	-0.21	-0.47; 0.04	0.10
Q80	Hookworm infection	0.22	-0.10; 0.54	0.18	0.27	-0.03; 0.58	0.08
	Sekou health facility	-0.23	-0.45; -0.01	0.04	-0.27	-0.51; -0.03	0.03
Q85	Hookworm infection	0.24	-0.02; 0.50	0.07	0.35	0.06; 0.64	0.02
	Sekou health facility	-0.22	-0.43; -0.01	0.04	-0.3	-0.52; -0.08	0.01
Q90	Hookworm infection	0.15	-0.08; 0.38	0.21	0.27	0.01; 0.53	0.04
	Sekou health facility	-0.23	-0.41; -0.06	0.01	-0.31	-0.51; -0.11	0.01
Q95	Hookworm infection	0.17	-0.11; 0.44	0.24	0.31	0.04; 0.58	0.02
	Sekou health facility	-0.13	-0.33; 0.07	0.21	-0.23	-0.41; -0.06	0.01

¹: From univariate quantile regression

²: From multivariate quantile regression

#: Reference: Attagon health facility