



HAL
open science

Introductory Chapter: Feed Your Mind - How Does Nutrition Modulate Brain Function throughout Life?

Clémentine Bosch-Bouju

► **To cite this version:**

Clémentine Bosch-Bouju. Introductory Chapter: Feed Your Mind - How Does Nutrition Modulate Brain Function throughout Life?. Feed Your Mind - How Does Nutrition Modulate Brain Function throughout Life?, IntechOpen, 2019, 10.5772/intechopen.89349 . hal-03485932

HAL Id: hal-03485932

<https://hal.science/hal-03485932>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,300

Open access books available

117,000

International authors and editors

130M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Introductory Chapter: Feed Your Mind - How Does Nutrition Modulate Brain Function throughout Life?

Clémentine Bosch-Bouju

1. Introduction

Nutrition is a transdisciplinary science that aims at studying the role of food and physical exercise on health and diseases. It includes notably biochemistry, metabolism, behavioural and social studies. Nutrition science is highly diverse and constantly evolving for several reasons:

- The control of energy balance is a primary need for humans and for all animals, which implies it is of high importance.
- A high diversity of dietary models throughout the world exists, which is closely related to the diversity of culture, even in one single country.
- In human societies, alimentation participates to social behaviour.
- Modern societies have been hugely transformed since food industrialization.

The last point has critically changed the way of life for people in industrialised countries. Food industrialisation was originally designed to ease life and feed the growing population following the First and Second World Wars. Unexpectedly, industrialisation has engendered a profound dietary imbalance.

For the first time in history, people with industrialised diet are facing food with high quantities of sugar and saturated fats. This diet can also lead to deficiency in essential nutrients, such as vitamins, minerals and polyunsaturated fatty acids. This change in diet has been paralleled by an increase in sedentariness, which contributes to energy imbalance. This change in the last 70 years leads to global epidemics as obesity, cardiovascular disorders and depressive disorders.

In this context, the brain appears as a structure highly sensitive to dietary imbalance [1]. In addition, the brain can participate to dietary imbalance by controlling feeding behaviours and through its interaction with peripheral organs and hormones involved in the control of energy balance. Thus, the impact of nutrition on brain function is a specific field of research that has gained interest in the recent years, and the role of the brain appears central in nutrition. Nutrition will impact the brain from early-developmental stages to adolescence, when the brain tissue is built and modelled. Nutrition also appears critical at adult age, to maintain optimal brain function. Finally, nutrition appears as an important environmental factor that

should be controlled through ageing, to prevent neurodegenerative processes. In this introductory chapter, we will take the example of vitamin A, which is a nutrient that has been demonstrated as critical for brain function throughout life [2–5].

2. Vitamin A and its active metabolite: retinoic acid

Vitamin A is a lipophilic vitamin that cannot be synthesised *de novo* by organism. Food sources are animal products for the preformed vitamin A and vegetables for provitamin A. Preformed vitamin A and provitamin A are found in sufficient quantities in most diets, but strong deficiency may occur in undernourished populations, mainly in sub-Saharan Africa. Vitamin A deficiency in children can cause blindness and growth retardation and can ultimately lead to death. Besides, vitamin A deficiency may occur with ageing, since its bioavailability can be reduced. In both cases, vitamin A supplementation can reverse this deficiency, which emphasises the need to study the impact of vitamin A supplementation for pregnant women and children, as well as for elderly.

The active metabolite of vitamin A in the brain is the all-*trans* retinoic acid (*atRA*), synthesised from vitamin A (retinol) by aldehyde dehydrogenase enzymes [6, 7]. Retinoic acid and its derivatives bind to nuclear receptors, namely, RAR and RXR. These receptors act as homo- or heterodimers to control gene transcription [3, 6]. Therefore, retinoic acid is a powerful morphogen factor, highly involved in embryonic development and growth [8].

3. The role of vitamin A during brain development

During brain development, retinoic acid synthesised from dietary vitamin A binds to nuclear receptors to drive transcription of hundreds of genes. Retinoic acid is a morphogen that acts through a concentration gradient [8]. To carry out the precision of retinoic acid gradient during embryonic development, metabolism and signalling pathways for retinoic acid are multiple and highly robust [8]. Remarkably, retinoic acid is the only morphogen that comes from dietary sources, which highlights the importance of vitamin A intake during pregnancy. Through its action on nuclear receptors, retinoic acid is implicated in the patterning of neural axis along the dorsoventral and anteroposterior axes [2].

In addition to development and growth, retinoic acid in the brain is critical for neuronal differentiation and neurite outgrowth. This is particularly true in the basal ganglia network, which is a group of brain structures involved in the control of action. Indeed, retinoic acid is key in the differentiation of specific subpopulations of neurons, such as the medium size spiny neurons in the striatum, or the dopaminergic neurons from the substantia nigra pars compacta [9–13].

4. The role of vitamin A in the adult brain

At adult age, vitamin A-derived retinoic acid has two remarkable roles; it is involved in homeostatic plasticity and in neurogenesis [3, 5, 14, 15]. First, retinoic acid induces molecular and morphological changes in neuronal network, to maintain overall synaptic activity constant, despite synaptic plasticity, a process named homeostatic synaptic plasticity [16]. For this, retinoic acid binds to nuclear RAR and RXR receptors, but evidence suggests that non-genomic mechanisms are also

involved [17–19]. Changes induced by retinoic acid to produce homeostatic plasticity include transcription of synaptic proteins, such as AMPA receptors, but also modulation of neuronal morphology, through outgrowth or retraction of synapse boutons and dendrites.

Second, retinoic acid controls neurogenesis. In rodent deficient for vitamin A, neurogenesis in the hippocampus is strongly dampened and can be restored by replenishment with vitamin A or derivatives [20–23]. However, in physiological conditions, retinoic acid applied exogenously can inhibit neurogenesis in the hippocampus and facilitate cell differentiation [3]. The apparent discrepancy between physiological and deficient states may be explained by the finely tuned concentration and timing of retinoic acid that is needed for its control of neurogenesis. In the hypothalamus, neurogenesis is highly dynamic to orchestrate circadian rhythms that govern several hypothalamic functions. In this structure, a similar role has been demonstrated for retinoic acid, inhibiting proliferation of cells and thus neurogenesis [5].

5. The role of vitamin A in the ageing brain

Ageing process is a physiological and unavoidable phenomenon involving the whole body. Related to brain function, ageing is characterised by a cognitive decline, which comprises all molecular and cellular alterations that lead ultimately to decreased performance in cognitive and executive functions [24]. Evidence suggests that bioavailability of retinoic acid in the brain reduces with ageing, likely due to peripheral alterations [4]. Yet several studies have highlighted the parallel between ageing, decrease in brain retinoid signalling and cognitive decline [4, 25–27]. Mechanisms linking retinoid metabolism and cognitive decline are not yet completely unravelled, but evidence suggests the involvement of glucocorticoids [20] and morphological changes of neuronal trees (unpublished data).

Beyond normal ageing, decreased retinoid signalling in the brain with ageing may be involved in neurodegenerative processes, such as in Alzheimer's and Parkinson's diseases. For Alzheimer's disease, accumulating evidence suggests that the lack of retinoic acid in the hippocampus is a risk factor that precipitates the deposition of β -amyloid plaques, a histological landmark of the disease [28–31]. Therefore, vitamin A and retinoids appear as potential therapeutics [30, 32]. However, mechanisms are not yet sufficiently understood to elaborate such therapeutics. Additionally, retinoids are lipophilic molecules that oxidise rapidly, which render their pharmacological use challenging.

In the context of Parkinson's disease, a similar process has been proposed. Reduced signalling of retinoids in the brain, and particularly in the substantia nigra compacta, may be a risk factor that accelerates the degeneration of dopaminergic neurons [33, 34]. This hypothesis is supported by the fact that rodents deficient for vitamin A or for retinoid receptors display motor impairments close to Parkinson's models [35, 36]. Some studies revealed encouraging results with retinoid supplementation to prevent neurodegeneration in cell culture or in vivo [37, 38], but research is still preliminary to envisage soon vitamin A-based treatments for this disease.

6. Conclusion

Through the solely example of vitamin A, we can appreciate the importance of nutrition for brain physiology, from development to adult age. Remarkably, a same

nutrient can have different roles depending on the timing, the brain structure and even the cell type. The field of research on nutrition applied to neurosciences is growing, to better encompass the impact of industrialised diet on brain function and also to consider nutrients and their derivatives as potential therapeutic strategies. In this way, it is to expect that nutritional approaches will give rise in the future to new therapeutics to prevent, and in some case cure, neurological disorders.

IntechOpen

IntechOpen

Author details

Clémentine Bosch-Bouju

INRA, Bordeaux INP, NutriNeuro, University of Bordeaux, Bordeaux, France

*Address all correspondence to: clementine.bosch-bouju@inra.fr

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Gómez-Pinilla F. Brain foods: The effects of nutrients on brain function. *Nature Reviews. Neuroscience*. 2008;**9**(7):568-578
- [2] Maden M. Retinoic acid in the development, regeneration and maintenance of the nervous system. *Nature Reviews. Neuroscience*. 2007;**8**(10):755-765
- [3] McCaffery P, Zhang J, Crandall JE. Retinoic acid signaling and function in the adult hippocampus. *Journal of Neurobiology*. 2006;**66**(7):780-791
- [4] Sauvant P, Feart C, Sauvant J, Pallet V. Vitamin A and cognitive impairment. In: *Diet and Nutrition in Dementia and Cognitive Decline. Advances in Clinical Neuroscience and Rehabilitation*; 2015. pp. 623-635. Available from: <https://linkinghub.elsevier.com/retrieve/pii/B9780124078246000586>
- [5] Shearer KD, Stoney PN, Morgan PJ, McCaffery PJ. A vitamin for the brain. *Trends in Neurosciences*. 2012;**35**(12):733-741
- [6] Blomhoff R, Blomhoff HK. Overview of retinoid metabolism and function. *Journal of Neurobiology*. 2006;**66**(7):606-630
- [7] Koppaka V, Thompson DC, Chen Y, Ellermann M, Nicolaou KC, Juvonen RO, et al. Aldehyde dehydrogenase inhibitors: A comprehensive review of the pharmacology, mechanism of action, substrate specificity, and clinical application. *The American Society for Pharmacology and Experimental Therapeutics. Pharmacological Reviews*. 2012;**64**(3):520-539
- [8] Schilling TF, Nie Q, Lander AD. Dynamics and precision in retinoic acid morphogen gradients. *Current Opinion in Genetics and Development*. 2012;**22**(6):562-569
- [9] Chatzi C, Brade T, Duyster G. Retinoic acid functions as a key GABAergic differentiation signal in the basal ganglia. *PLoS Biology*. 2011;**9**(4):e1000609
- [10] Occhi G, Regazzo D, Albiger NM, Ceccato F, Ferasin S, Scanarini M, et al. Activation of the dopamine receptor Type-2 (DRD2) promoter by 9-cis retinoic acid in a cellular model of Cushing's disease mediates the inhibition of cell proliferation and ACTH secretion without a complete corticotroph-to-melanotroph transdifferentiation. *Endocrinology*. 2014;**155**(9):3538-3549
- [11] Pedrini S, Bogush A, Ehrlich ME. Phosphatidylinositide 3-kinase and protein kinase C zeta mediate retinoic acid induction of DARPP-32 in medium size spiny neurons in vitro. *Journal of Neurochemistry*. 2008;**106**(2):917-924
- [12] Podleśny-Drabiniok A, Sobska J, de Lera AR, Gołombiowska K, Kamińska K, Dollé P, et al. Distinct retinoic acid receptor (RAR) isotypes control differentiation of embryonal carcinoma cells to dopaminergic or striatopallidal medium spiny neurons. *Scientific Reports*. 2017;**7**(1):13671
- [13] Rataj-Baniowska M, Niewiadomska-Cimicka A, Paschaki M, Szyszka-Niagolov M, Carramolino L, Torres M, et al. Retinoic acid receptor β controls development of Striatonigral projection neurons through FGF-dependent and Meis1-dependent mechanisms. *Journal of Neuroscience: The Official Journal of the Society for Neuroscience*. 2015;**35**(43):14467-14475
- [14] Lane MA, Bailey SJ. Role of retinoid signalling in the adult brain. *Progress in Neurobiology*. 2005;**75**(4):275-293

- [15] Olson CR, Mello CV. Significance of vitamin A to brain function, behavior and learning. *Molecular Nutrition and Food Research*. 2010;**54**(4):489-495
- [16] Turrigiano GG, Nelson SB. Homeostatic plasticity in the developing nervous system. *Nature Reviews. Neuroscience*. 2004;**5**(2):97-107
- [17] Aoto J, Nam CI, Poon MM, Ting P, Chen L. Synaptic signaling by all-trans retinoic acid in homeostatic synaptic plasticity. *Neuron*. 2008;**60**(2):308-320
- [18] Arendt KL, Zhang Y, Jurado S, Malenka RC, Südhof TC, Chen L. Retinoic acid and LTP recruit postsynaptic AMPA receptors using distinct SNARE-dependent mechanisms. *Neuron*. 2015;**86**(2):442-456
- [19] Chen L, Lau AG, Sarti F. Synaptic retinoic acid signaling and homeostatic synaptic plasticity. *Neuropharmacology*. 2014;**78**:3-12
- [20] Bonhomme D, Minni AM, Alfos S, Roux P, Richard E, Higuieret P, et al. Vitamin A status regulates glucocorticoid availability in Wistar rats: Consequences on cognitive functions and hippocampal neurogenesis? *Frontiers in Behavioral Neuroscience*. 2014;**20**:1-13
- [21] Bonnet E, Touyarot K, Alfos S, Pallet V, Higuieret P, Abrous DN. Retinoic acid restores adult hippocampal neurogenesis and reverses spatial memory deficit in vitamin A deprived rats. In: McCabe BD, editor. *PLoS One*. 2008;**3**(10):e3487
- [22] Jacobs S, Lie DC, DeCicco KL, Shi Y, DeLuca LM, Gage FH, et al. Retinoic acid is required early during adult neurogenesis in the dentate gyrus. *Proceedings of the National Academy of Sciences of the United States of America*. 2006;**103**(10):3902-3907
- [23] Touyarot K, Bonhomme D, Roux P, Alfos S, Lafenêtre P, Richard E, et al. A mid-life vitamin A supplementation prevents age-related spatial memory deficits and hippocampal neurogenesis alterations through CRABP-I. *PLoS One*. 2013;**8**(8):e72101
- [24] Hedden T, Gabrieli JDE. Insights into the ageing mind: A view from cognitive neuroscience. *Nature Reviews. Neuroscience*. 2004;**5**(2):87-96
- [25] Bonhomme D, Pallet V, Dominguez G, Servant L, Henkous N, Lafenêtre P, et al. Retinoic acid modulates intrahippocampal levels of corticosterone in middle-aged mice: Consequences on hippocampal plasticity and contextual memory. *Frontiers in Aging Neuroscience*. 2014;**6**:6
- [26] Etchamendy N, Enderlin V, Marighetto A, Pallet V, Higuieret P, Jaffard R. Vitamin A deficiency and relational memory deficit in adult mice: Relationships with changes in brain retinoid signalling. *Behavioural Brain Research*. 2003;**145**(1-2):37-49
- [27] Mingaud F, Mormede C, Etchamendy N, Mons N, Niedergang B, Wietrzyk M, et al. Retinoid hyposignaling contributes to aging-related decline in hippocampal function in short-term/working memory organization and long-term declarative memory encoding in mice. *Journal of Neuroscience: The Official Journal of the Society for Neuroscience*. 2008;**28**(1):279-291
- [28] Corcoran JPT, So PL, Maden M. Disruption of the retinoid signalling pathway causes a deposition of amyloid beta in the adult rat brain. *The European Journal of Neuroscience*. 2004;**20**(4):896-902
- [29] Goncalves MB, Clarke E, Hobbs C, Malmqvist T, Deacon R, Jack J, et al. Amyloid β inhibits retinoic acid

synthesis exacerbating Alzheimer disease pathology which can be attenuated by an retinoic acid receptor α agonist. *The European Journal of Neuroscience*. 2013;**37**(7):1182-1192

[30] Goodman AB. Retinoid receptors, transporters, and metabolizers as therapeutic targets in late onset Alzheimer disease. *Journal of Cellular Physiology*. 2006;**209**(3):598-603

[31] Goodman AB, Pardee AB. Evidence for defective retinoid transport and function in late onset Alzheimer's disease. *Proceedings of the National Academy of Sciences of the United States of America*. 2003;**100**(5):2901-2905

[32] Sodhi RK, Singh N. Retinoids as potential targets for Alzheimer's disease. *Pharmacology, Biochemistry, and Behavior*. 2014;**120**:117-123

[33] Cai H, Liu G, Sun L, Ding J. Aldehyde dehydrogenase 1 making molecular inroads into the differential vulnerability of nigrostriatal dopaminergic neuron subtypes in Parkinson's disease. *Translational Neurodegeneration*. 2014;**3**:27

[34] Liu G, Yu J, Ding J, Xie C, Sun L, Rudenko I, et al. Aldehyde dehydrogenase 1 defines and protects a nigrostriatal dopaminergic neuron subpopulation. *The Journal of Clinical Investigation*. 2014;**124**(7):3032-3046

[35] Carta M, Stancampiano R, Tronci E, Collu M, Usiello A, Morelli M, et al. Vitamin A deficiency induces motor impairments and striatal cholinergic dysfunction in rats. *Neuroscience*. 2006;**139**(4):1163-1172

[36] Krezel W, Ghyselinck N, Samad TA, Dupé V, Kastner P, Borrelli E, et al. Impaired locomotion and dopamine signaling in retinoid receptor mutant mice. *Science*. 1998;**279**(5352):863-867

[37] Esteves M, Cristóvão AC, Saraiva T, Rocha SM, Baltazar G, Ferreira L, et al. Retinoic acid-loaded polymeric nanoparticles induce neuroprotection in a mouse model for Parkinson's disease. *Frontiers in Aging Neuroscience*. 2015;**7**:20

[38] Reiner DJ, Yu S-J, Shen H, He Y, Bae E, Wang Y. 9-cis retinoic acid protects against methamphetamine-induced neurotoxicity in nigrostriatal dopamine neurons. *Neurotoxicity Research*. 2014;**25**(3):248-261