



HAL
open science

Simply connected open 3-manifolds with slow decay of positive scalar curvature

Jian Wang

► **To cite this version:**

Jian Wang. Simply connected open 3-manifolds with slow decay of positive scalar curvature. Comptes Rendus. Mathématique, 2019, 357, pp.284 - 290. 10.1016/j.crma.2019.02.001 . hal-03485967

HAL Id: hal-03485967

<https://hal.science/hal-03485967>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

SIMPLY-CONNECTED OPEN 3-MANIFOLDS WITH SLOW DECAY OF POSITIVE SCALAR CURVATURE

JIAN WANG

ABSTRACT. The goal of this paper is to investigate the topological structure of open simply-connected 3-manifolds whose scalar curvature has a slow decay at infinity. In particular, we show that the Whitehead manifold does not admit a complete metric, whose scalar curvature decays slowly, and in fact that any contractible complete 3-manifolds with such a metric is diffeomorphic to \mathbb{R}^3 . Furthermore, using this result, we prove that any open simply-connected 3-manifold M with $\pi_2(M) = \mathbb{Z}$ and a complete metric as above, is diffeomorphic to $\mathbb{S}^2 \times \mathbb{R}$.

1. INTRODUCTION

Thanks to G.Perelman's proof of W.Thurston's Geometrisation conjecture in [12, 13, 14], the topological structure of compact 3-manifolds is now well understood. However, it is known from the early work [21] of J.H.C Whitehead that the topological structure of non-compact 3-manifolds is much more complicated. For example, there exists a contractible open 3-manifold, called Whitehead manifold, which is not homeomorphic to \mathbb{R}^3 . An interesting question in differential geometry is whether the Whitehead manifold admits a complete metric with positive scalar curvature.

The study of manifolds of positive scalar curvature has a long history. Among many results, we mention the topological classification of compact manifolds of positive scalar curvature and the Positive Mass Theorem. There are two methods which have achieved many breakthroughs: minimal hypersurfaces and K-theory.

The K-theory method is pioneered by A.Lichnerowicz and is systemically developed by M.Gromov and H.Lawson in [6], based on the Atiyah-Singer Index theorem in [1]. Furthermore, combined with some results about the Novikov conjecture, S.Chang, S.Weinberger and G.L.Yu [3] investigated the topological structure of open 3-manifolds with uniformly positive scalar curvatures and finitely generated fundamental groups. Precisely, they proved that any contractible 3-manifold whose scalar curvature is bounded away from zero is \mathbb{R}^3 , which implies that the Whitehead manifold does not admit a metric with uniformly positive scalar curvature.

The origin of the minimal hypersurfaces method is the article of R.Schoen and S.T.Yau [18]. For open 3-manifolds, there are many applications, such

as Positive Mass Theorem[19, 20] and 3-dimensional Milnor Conjecture[7]. In [6], Gromov and Lawson applied this method to open 3-manifolds.

In this paper, we extend Gromov-Lawson's Theorem[6] to open 3-manifolds whose scalar curvature has a decay at infinity:

Theorem 1.1. *Assume that (M^3, g) is a contractible complete 3-manifold. Let $0 \in M$ and $r(x)$ be the distance function from x to 0 . If there exists a number $\alpha \in [0, 2)$ such that*

$$\liminf_{r(x) \rightarrow \infty} r^\alpha(x) \text{Scal}(x) > 0,$$

then M^3 is diffeomorphic to \mathbb{R}^3 .

Our main tool comes from the solution to the Plateau Problem. Let us give a brief review of the existence of the solution to the Plateau problem and their regularity. In the case of \mathbb{R}^3 , the existence is due to Douglas [4] and Radó [15] : For any smooth curve γ in \mathbb{R}^3 , there exists a surface with minimal area, spanning γ , which is parametrized by a disc D^2 . Furthermore, Osserman [10] and Gulliver [5] proved that this solution has no interior branch point. In 1948, Morrey [8, 9] devised a new method to solve the Plateau problem for a map from a disc to a "homogeneously regular" Riemannian manifold. In addition, Osserman and Gulliver's arguments show that Morrey's solution also has no interior branch point.

To sum up, for any null-homotopic smooth curve γ in a complete Riemannian manifold (M^3, g) , there exists a continuous map $f : D^2 \rightarrow M$ such that f induces a homomorphism between ∂D and γ and the interior of f is a minimal immersion.

Let us explain the scheme of Gromov-Lawson's proof in [6]. From J.Stalling's results in [17], it is sufficient to prove that M is simply-connected at infinity. We argue by contradiction and suppose that M is not simply-connected at infinity. Namely, there exists a compact set K in M , satisfying that for any geodesic ball $B_R(0)$ containing K , there is a smooth closed curve γ in $M \setminus B_R(0)$, which is not null-homotopic in $M \setminus K$. Choosing R large enough and considering the solution S to the Plateau problem for γ , S can not be contained in a "small" neighborhood of some circle(s), which contradicts to a result of Rosenberg in [16]. Before stating Rosenberg's result, we firstly define the notion of stable H -surface.

Definition 1.2. Let (M, g) be a 3-dimensional Riemannian manifold and Σ a surface immersed in (M, g) . Σ is said to be a H -surface if the mean curvature of Σ is constant and equals to H , where $H \in \mathbb{R}$. Furthermore, we consider the operator L of Σ defined as

$$L = \Delta_\Sigma + |A|^2 + Ric_M(\mathbf{n})$$

where $|A|^2$ is the square length of the second fundamental form of Σ in M , \mathbf{n} is a unit normal vector field along Σ . We say that Σ is a *stable H -surface*

if

$$-\int_{\Sigma} uL(u) \geq 0$$

for any smooth function u with compact support over Σ .

In [16], applying the stable condition, Rosenberg obtained the following:

Theorem 1.3. [16] *Let (N, g) be a complete Riemannian 3-manifold satisfying*

$$3H^2 + S(x) \geq c, \text{ for any } x \in N,$$

where $S(x)$ is the scalar curvature, $H \in \mathbb{R}$ and $c > 0$. Suppose that Σ is a stable H -surface immersed in N . Then one has for any $x \in \Sigma$

$$d_{\Sigma}(x, \partial\Sigma) \leq 2\pi/(3c)^{1/2},$$

where d_{Σ} is the intrinsic distance in Σ .

Remark 1.4. From the proof of Theorem 1.3 in [16], we do not need the condition that the scalar curvature $S(x)$ is uniformly bounded.

When trying to generalize Gromov-Lawson's arguments to contractible 3-manifolds whose scalar curvature decay at infinity, one encounters an obstacle—the lack of uniform lower-bound of the scalar curvature on the surface S spanning the curve γ . However, this can be overcome by choosing a new curve σ near the boundary of $B_{R/2}(0)$, which is homotopic to γ in $M \setminus K$. More precisely, using the regularity property of S , after a small deformation, the intersection of S and the boundary of $B_{R/2}(0)$ is some circle(s). Therefore, the intersection can be chosen as a curve σ , which satisfies the required property. This leads to Theorem 1.1, by applying Gromov-Lawson's argument[6] to this curve σ .

2. THE PROOF OF THEOREM 1.1

Before the proof of Theorem 1.1, we introduce an important notion and recall a classical result about open 3-manifolds.

Definition 2.1. A topological space X is said to be *simply-connected at infinity*, if for any compact set C , there exists a compact set V containing C , such that the induced map $i_* : \pi_1(X \setminus V) \rightarrow \pi_1(X \setminus C)$ is trivial, where $i : X \setminus V \rightarrow X \setminus C$ is an inclusion map.

It is well-known from [21] that the Whitehead manifold is not simply-connected at infinity. In fact, there is a unique non-compact 3-manifold which is both simply-connected at infinity and contractible.

Theorem 2.2. [Stallings] [17] *Let X be a contractible 3-manifold, then X is homeomorphic to \mathbb{R}^3 if and only if X is simply-connected at infinity.*

Here is the idea of proof of Theorem 1.1. In the following, we assume that (M, g) is a contractible 3-manifold as in Theorem 1.1. According to Theorem 2.2, it is sufficient to show that M is simply-connected at infinity.

From the assumption about the scalar curvature in Theorem 1.1, there are two positive constants C and R_0 , such that for $r(x) \geq R_0$, one has,

$$\text{Scal}(x) \geq \frac{C}{r(x)^\alpha}.$$

Compared with the argument of Gromov-Lawson in [6], our main difficulty is the lack of the uniform lower-bound of the scalar curvature. However, using Theorem 1.3, we obtain the following:

Proposition 2.3. *If $R > 2 \max \left\{ R_0, \left(\frac{4^{1+\alpha/2}\pi}{(3C)^{1/2}} \right)^{\frac{2}{2-\alpha}} \right\}$, then the induced map $\pi_1(M \setminus B_{4R}(0)) \rightarrow \pi_1(M \setminus B_R(0))$ is trivial.*

Then, due to Proposition 2.3, M is simply-connected at infinity, which implies that M is homeomorphic to \mathbb{R}^3 . It implies Theorem 1.1.

We now prove Proposition 2.3. Let us consider a smooth closed curve γ in $M \setminus B_{4R}(0)$. Because γ may be far away from the compact set $B_R(0)$, there does not exist a “good” estimate of the lower-bound of the scalar curvature along γ . In order to overcome it, we establish the following lemma:

Lemma 2.4. *For any smooth circle γ in $M \setminus B_{4R}(0)$, exactly one of the following holds:*

- γ is contractible in $M \setminus B_R(0)$,
- for any $\epsilon > 0$ and any $R < Q < 4R$, there exists a curve $\hat{\sigma}$ in $B_{Q+\epsilon}(0) \setminus B_Q(0)$, which is not contractible in $M \setminus B_R(0)$.

Proof. Let $\hat{f} : D^2(1) \rightarrow M$ be the solution to the Plateau problem for γ , where $D^2(1)$ is the unit disc. Then $\hat{f}|_{\partial D^2(1)}$ is a homeomorphism between $\partial D^2(1)$ and γ . Furthermore, the interior of \hat{f} is an immersion.

Suppose that γ is non-contractible in $M \setminus B_R(0)$. Therefore, $\hat{f}(D^2(1)) \cap \partial B_Q(0)$ is nonempty. Furthermore, $\hat{f}^{-1}(\hat{f}(D^2(1)) \cap B_Q(0))$ belongs to the interior of $D^2(1)$. Then, after a small variation of \hat{f} in the interior of $D^2(1)$, there exists a map $f : D^2(1) \rightarrow M$, such that

- the interior of f is still an immersion,
- $f|_{\partial D^2(1)}$ is a homeomorphic map from $\partial D^2(1)$ to γ ,
- f intersects transversally with $\partial B_Q(0)$.

Therefore, the pre-image of $f(D^2(1)) \cap \partial B_Q(0)$ is a 1-dimensional compact submanifold in $D^2(1)$. That is to say: $f^{-1}(f(D^2(1)) \cap \partial B_Q(0))$ is a disjoint union of some circle(s) in $D^2(1)$.

Let $f^{-1}(B_Q(0))$ be the disjoint union of $\{\gamma_i\}_{i \in I}$, where each γ_i is diffeomorphic to a circle in $D^2(1)$ and I is a finite set. Let D_i be the unique disc bounded by γ_i in $D^2(1)$. Let us consider the set $\{D_i\}_{i \in I}$ and define the partially ordered relationship on it, induced by inclusion. For any maximal element D_j in $(\{D_i\}_{i \in I}, \subset)$, $\gamma_j = \partial D_j$ is defined as an outmost circle.

Let $\{\gamma_j\}_{j \in I_0}$ be outmost circles in $\{\gamma_i\}_{i \in I}$, where $I_0 \subset I$. For each outmost circle γ_j , we assume that σ_j is the boundary of a tubular neighborhood

of γ_i in $D^2(1) \setminus \cup_{i \in I} D_i$, contained in $f^{-1}(B_{Q+\epsilon}(0) \setminus B_Q(0))$ and \hat{D}_j is the unique disc bounded by σ_j in $D^2(1)$.

Claim: some element in $\{f(\sigma_j)\}_{j \in I_0}$ is not contractible in $M \setminus B_R(0)$.

We argue it by contradiction. Suppose that each $f(\sigma_j)$ is contractible in $M \setminus B_R(0)$. In other words, for each $j \in I_0$ there exists a continuous map $g_j : D^2(1) \rightarrow M$, such that (1) $g_j(D^2(1)) \cap B_R(0) = \emptyset$; (2) $g_j|_{\partial D^2(1)}$ is a homomorphism from $\partial D^2(1)$ to $f(\sigma_j)$.

After changing the coordinate over $D^2(1)$, we may suppose that for each $j \in I_0$, \hat{D}_j is a disc with center at x_j and radius r_j . We will construct a new map $g : D^2(1) \rightarrow M$, such that (1) $g|_{\partial D^2(1)}$ is a homeomorphic map from $\partial D^2(1)$ to γ ; $g(D^2(1)) \cap B_R(0) = \emptyset$. It implies that γ is contractible in $M \setminus B_R(0)$, which contradicts our assumption that γ is not contractible in $M \setminus B_R(0)$.

Let us describe the construction of g as follows:

$$g(x) = \begin{cases} f(x), & x \in D^2(1) \setminus \cup_{j \in I_0} \hat{D}_j \\ g_j\left(\frac{x-x_j}{r_j}\right), & x \in \hat{D}_j. \end{cases} \quad (2.1)$$

g is a required map described as above. We finish the proof of the claim.

We may suppose that $f(\sigma_{j_0})$ is non-contractible in $M \setminus B_R(0)$ and choose $\hat{\sigma} = f(\sigma_{j_0})$. It is the required candidate in the assertion. This completes the proof. \square

Remark 2.5. In the proof of Lemma 2.4, \hat{f} is only required to be an immersion. Because we just need the property that after perturbing \hat{f} , we obtain a new immersion f satisfying that f intersects transversally with the surface Σ . Therefore, the pre-image of f is a submanifold. Hence, there are many other choices for \hat{f} .

We now give the proof of Proposition 2.3.

Proof of Proposition 2.3. We prove it by contradiction. First, we assume that for some $R > 2 \max\{R_0, (\frac{4^{1+\alpha/2}\pi}{(3C)^{1/2}})^{\frac{2}{2-\alpha}}\}$, there is a curve $\gamma \subset M \setminus B_{4R}(0)$ such that γ is nontrivial in $\pi_1(M \setminus B_R(0))$.

We take $Q = 2R$ and $\epsilon = 1$ and apply Lemma 2.4 to γ . There is a non-contractible curve $\hat{\sigma} \subset B_{2R+1}(0) \setminus B_{2R}(0)$ in $M \setminus B_R(0)$.

Let $f : D^2(1) \rightarrow M$ be the solution to the Plateau problem for the circle $\hat{\sigma}$ in M . Then $f(D^2(1)) \cap \partial B_R(0)$ is non-empty, since $\hat{\sigma}$ is non-contractible in $M \setminus B_R(0)$. Let us consider the set $\Sigma := f(D^2(1)) \cap (B_{3R}(0) \setminus B_R(0))$. By the assumption of the scalar curvature, i.e. $\liminf_{r(x) \rightarrow \infty} r^\alpha(x) \text{Scal}(x) > 0$, one has,

$$\text{Scal}(x) \geq \frac{C}{(4R)^\alpha}, \text{ over } \Sigma.$$

By Theorem 1.3 and Remark 1.4, we deduce that

Σ is contained in the $\frac{(4R)^{\alpha/2}\pi}{(3C)^{1/2}}$ -neighborhood of $\partial\Sigma$.

However, $\partial\Sigma = \hat{\sigma} \Pi (\partial B_R(0) \cap \Sigma)$ is contained in a union of $M \setminus B_{2R}(0)$ and the closure of $B_R(0)$. The above fact gives

$$R \leq 2 \frac{(4R)^{\alpha/2}\pi}{(3C)^{1/2}}$$

That is to say, $R \leq 2 \left(\frac{4^{\alpha/2}\pi}{(3C)^{1/2}} \right)^{\frac{2}{2-\alpha}} < 2 \left(\frac{4^{1+\alpha/2}\pi}{(3C)^{1/2}} \right)^{\frac{2}{2-\alpha}}$. This contradicts the choice of R . \square

As a corollary, we have the following

Corollary 2.6. *The Whitehead manifold does not admit a complete metric with $\liminf_{r(x) \rightarrow \infty} r^\alpha(x) \text{Scal}(x) > 0$, where r is a distance function from a given point and $\alpha \in [0, 2)$.*

3. APPLICATION

In this section, we use Theorem 1.1 to prove that any complete open 3-manifold M such that (1) $\pi_1(M) = \{0\}$ and $\pi_2(M) = \mathbb{Z}$; (2) its scalar curvature decays slowly at infinity, is homeomorphic to $\mathbb{R} \times \mathbb{S}^2$. Let us review the sphere theorem and some classical applications of the sphere theorem.

Theorem 3.1 (Sphere Theorem [11, 2]). *Any orientable 3-manifold M with non-trivial $\pi_2(M)$ has an embedded sphere.*

The non-trivial embedded sphere plays a crucial role in the prime decomposition of 3-manifolds. In a simply-connected 3-manifold, the non-trivial embedded sphere always separates the 3-manifold into two connected components.

Lemma 3.2. *Let M^3 be a simply-connected open manifold and $S \subset M$ a non-trivial sphere, then $M \setminus S$ is a disjoint union of two non relatively compact components.*

Proof. First, we claim that S is a separating sphere in M . If $M \setminus S$ is connected, then there exists a circle $\alpha \subset M$, such that α intersects S transversally at one point. Hence, the intersection number (S, α) is 1 or -1 . However, the intersection number between S and α is zero, since α is contractible in M and the intersection number is a homotopic invariant. It leads to a contradiction. Therefore, S separates M into two components.

Second, we will show that each component of $M \setminus S$ is non relatively compact. If one component, denoted by M' , is relatively compact, we use the Van Kampen Theorem and $\pi_1(M) = \{1\}$ to obtain that M' is also simply connected. We define a 3-manifold M'' by gluing a 3-ball along the boundary sphere $\partial M'$. Hence, by the Van Kampen theorem, the compact 3-manifold

M'' is also simply connected. By the Poincaré conjecture [12, 13, 14], M'' is a 3-sphere. Therefore, M' is homeomorphic to a 3-ball, which contradicts our assumption that S is non-trivial in $\pi_2(M)$. \square

The second homotopy group is an important homotopic invariant for the classification of CW complexes. We will use the following lemma frequently.

Lemma 3.3. *An open 3-manifold M with $\pi_1(M) = \pi_2(M) = 0$ is contractible.*

Proof. Because M is an open manifold, the 0-dimensional cohomology group of M with compact support, denoted by $H_c^0(M, \mathbb{Z})$, is trivial. The Poincaré duality gives $H_c^0(M, \mathbb{Z}) \cong H_3(M, \mathbb{Z}) = \{0\}$. Since $\pi_1(M) = \pi_2(M) = \{0\}$, we use the Hurewicz Theorem to see that $\pi_3(M) \cong H_3(M) = \{0\}$. Then, using the Hurewicz Theorem inductively, we have that $\pi_n(M) \cong H_n(M) = 0$, for any $n \geq 3$. By Whitehead Theorem, M is contractible. \square

We now consider a simply-connected open 3-manifold M with nontrivial second homotopy group. From Theorem 3.1 and Lemma 3.2, there exists a non-trivial sphere S separating M into two non relatively compact parts: M_1 and M_2 . We define M_i^0 as a union of B^3 and M_i along S , where B^3 is a 3-ball. Then M can be viewed as the connected sum of M_0^0 and M_1^0 .

The pair (M, \bar{M}_i) is a CW pair, where \bar{M}_i is the closure of M_i in M , for $i \in I$, where $I = (\mathbb{Z}/2\mathbb{Z}, +)$. For each CW pair (M, \bar{M}_i) , one has a continuous map $f_i : M \rightarrow M/\bar{M}_i$. Because ∂M_i is a sphere S , M/\bar{M}_i is homeomorphic to M_{i+1}^0 , for $i \in I = (\mathbb{Z}/2\mathbb{Z}, +)$. Therefore, f_i can be viewed as a continuous map from M to M_{i+1}^0 . Furthermore, the induced map $(f_i)_* : \pi_2(M) \rightarrow \pi_2(M_{i+1}^0)$ verifies the following:

Proposition 3.4. *For each $i \in I$, the induced map $(f_i)_* : \pi_2(M) \rightarrow \pi_2(M_{i+1}^0)$ is a surjective map with nontrivial kernel.*

Proof. In the above statement, S is a non-trivial sphere in M . M_{i+1}^0 is obtained by gluing a 3-ball B^3 along S to M_{i+1} , i.e. $M_{i+1}^0 = M_{i+1} \cup_S B^3$, for $i \in I$.

The image of S is bounded by a 3-ball, which implies that $[f_i(S)]$ is trivial in $\pi_2(M_{i+1}^0)$. Therefore, the kernel of $(f_i)_*$ is non-trivial.

For each $i \in I$, any continuous map $g : \mathbb{S}^2 \rightarrow M_{i+1}^0$ is homotopic to another continuous map $g' : \mathbb{S}^2 \rightarrow M_{i+1}$ in $M_{i+1}^0 = M_{i+1} \cup_S B^3$. Let $i : M_{i+1} \rightarrow M$ be the inclusion map, $i \circ g'$ is a map from \mathbb{S}^2 to M and $(f_i)_*([g']) = [g]$ in $\pi_2(M_{i+1}^0)$. Therefore, $(f_i)_*$ is surjective. \square

In the following, we will use the sphere theorem to analyze the topological structure of a simply-connected open 3-manifold M with $\pi_2(M) = \mathbb{Z}$. Together with the surgery as described above, we prove :

Theorem 3.5. *Assume that (M^3, g) is a simply-connected open 3-manifold with $\pi_2(M) = \mathbb{Z}$. Let $0 \in M$ be a point and $r(x)$ a distance function from x*

to 0. If there exists a real number $\alpha \in [0, 2)$, such that,

$$\liminf_{r(x) \rightarrow \infty} r^\alpha(x) \text{Scal}(x) > 0,$$

then M^3 is diffeomorphic to $\mathbb{R} \times \mathbb{S}^2$.

Proof. By the Sphere theorem, there exists a non-trivial embedded sphere S cutting M into two non relatively compact components M_0 and M_1 . M can be viewed as the connected sum of two simply-connected 3-manifolds M_0^0 and M_1^0 , where $M_0^0 = M_0 \cup_S B^3$ and $M_1^0 = M_1 \cup_S B^3$.

Furthermore, we will construct a metric over M_i^0 satisfying the curvature condition as assumed in Theorem 1.1.

First, let $N_r(S)$ be the tubular neighborhood of S in \overline{M}_i , with radius r . For any constant $\epsilon > 0$, there exists a smooth function $\phi(x)$ supported in $N_{2\epsilon}(S)$, such that, $\phi(x) = 1$ on $N_\epsilon(S)$. Meanwhile, we may find a smooth function $\tau(x)$ on M_i^0 , satisfying the following:

- (1) $\tau(x)$ has support in the compact set $M_i^0 \setminus (\overline{M}_i \setminus N_{4\epsilon}(S))$,
- (2) $\tau(x) = 1$ on $M_i^0 \setminus (\overline{M}_i \setminus N_{2\epsilon}(S))$

Second, choose any smooth metric g' over $M_i^0 \setminus (\overline{M}_i \setminus N_{4\epsilon}(S))$. Define a new metric g_i over M_i^0 :

$$g_i = \begin{cases} g, & \overline{M}_i \setminus N_{4\epsilon}(S); \\ g', & M_i^0 \setminus M_i; \\ (1 - \phi)g + \tau g', & \text{otherwise.} \end{cases} \quad (3.1)$$

where g is a metric in the assumption of Theorem 3.5.

If $[S]$ is a generator of $\pi_2(M)$, from the proof of Proposition 3.4, $[S]$ is contained in the kernel of the induced map $(f_i)_*$. Then $\pi_2(M_i^0) = \{0\}$ for each $i \in I$. Thanks to Lemma 3.3, M_i^0 is contractible. Therefore, (M_i^0, g_i) satisfies the curvature condition as assumed in Theorem 3.5. Due to Theorem 1.1, each M_i^0 is homeomorphic to \mathbb{R}^3 . M is the connected sum of two \mathbb{R}^3 s. Hence, M is diffeomorphic to $\mathbb{S}^2 \times \mathbb{R}$.

If $[S]$ is not a generator, by Proposition 3.4, then the induced map maps from $\pi_2(M)$ onto $\pi_2(M_i^0)$ with non-trivial kernel, which indicates that $\pi_2(M_i^0)$ is finite. In this case, there is a unique topological structure of M_i^0 as follows. By Proposition 3.6 (in the following), each M_i^0 is homeomorphic to \mathbb{R}^3 which implies that M is homeomorphic to $\mathbb{S}^2 \times \mathbb{R}$. \square

Proposition 3.6. *Let (M^3, g) be a simply-connected open 3-manifold satisfying that $\pi_2(M)$ is a finite group. For $0 \in M$, assume that r is the distance function from 0. If there exists a number $\alpha \in [0, 2)$, such that,*

$$\liminf_{r(x) \rightarrow \infty} r^\alpha(x) \text{Scal}(x) > 0,$$

then M^3 is homeomorphic to \mathbb{R}^3 .

Proof. Suppose that M is not homeomorphic to \mathbb{R}^3 . In addition, we observe that $\pi_2(M)$ is non-trivial. (Otherwise, by Lemma 3.3, M is contractible.

Combined with Theorem 1.1, M is \mathbb{R}^3 , which contradicts our above assumption.)

By the sphere theorem, there exists a non-trivial embedded sphere S cutting M into two non relatively compact components. Two simply-connected non-compact manifolds M_0^0 and M_1^0 are obtained by the process described in the proof of Theorem 3.5. Therefore, M_0^0 and M_1^0 satisfy the following:

- M is the connected sum of M_0^0 and M_1^0
- $|\pi_2(M_i^0)| < |\pi_2(M)|$, since the map $\pi_2(M) \rightarrow \pi_2(M_i^0)$ is surjective with non-trivial kernel (Proposition 3.4).

We observe that one of $\{\pi_2(M_i^0)\}_{i \in I}$ is nontrivial. (Otherwise, from Lemma 3.3, M_i^0 is contractible. From Theorem 1.1, M_i^0 is \mathbb{R}^3 , which implies that M is $\mathbb{S}^2 \times \mathbb{R}$. This contradicts the assumption that $\pi_2(M)$ is finite.)

Without loss of generality, we may assume that $\pi_2(M_0^0)$ is nontrivial. Set $M_1 = M_0^0$. M_0^1 and M_1^1 can be constructed as the above process in the proof of Theorem 3.5. We may suppose that one of $\pi_2(M_i^1)$ is non-trivial. (Otherwise, a similar argument for $\pi_2(M_i^0)$ as above will work.) Then we can repeat this process, until the second homotopy groups of two new manifolds are both trivial.

After repeating the above process several times, three families of manifolds M_k , M_0^k and M_1^k are constructed as in the above process. These manifolds satisfy the following:

- M_k is the connected sum of M_0^k and M_1^k ,
- $|\pi_2(M_k)| < |\pi_2(M_{k-1})|$.

Because $\pi_2(M)$ is finite and $|\pi_2(M_k)| < |\pi_2(M_{k-1})|$, this process will stop after finite steps. There exists an integer k_0 such that $\pi_2(M_{k_0})$ is nontrivial and $\pi_2(M_i^{k_0})$ is trivial for each i . From Lemma 3.3, this implies that $M_i^{k_0}$ is contractible. Due to Theorem 1.1, $M_i^{k_0}$ is homeomorphic to \mathbb{R}^3 . However, M_{k_0} is the connected sum of $M_0^{k_0}$ and $M_1^{k_0}$. Hence, M_{k_0} is $\mathbb{S}^2 \times \mathbb{R}$, which contradicts the fact that $\pi_2(M_{k_0})$ is finite. □

ACKNOWLEDGEMENT

The author would like to express his deep gratitude to his advisor, Professor Gérard Besson for his useful suggestions, also for his constant encouragement and support. The work is supported by ERC GETOM project and ANR-CCEM project.

REFERENCES

- [1] Atiyah, M.; Singer, I. *The index of elliptic operators on compact manifolds*, Bull. Amer. Math. Soc. **69** (1962), 422-433.
- [2] Batude, J. L. *Singularité générique des applications différentiables de la 2-sphère dans une 3-variété différentiable*. Annales de l'Institut Fourier. **21** (1971), no. 3, 151-172.
- [3] Chang, S.; Weinberger, S.; Yu, G. *Taming 3-manifolds using scalar curvature*. Geom. Dedicata. **148** (2010), 3-14.

- [4] Douglas, J. *Solution of the problem of Plateau*, Trans. Am. Math. Soc. **32** (1932), 731-756.
- [5] Gulliver, R. D. *Regularity of minimizing surfaces of prescribed mean curvature*, Ann. Math. **97** (1973), 275-305.
- [6] Gromov, M.; Lawson, H. *The positive scalar curvature and the Dirac operator on the complete Riemannian manifolds*, Inst. Hautes Études Sci. Publ. Math. **58** (1984), 83-196.
- [7] Liu, G. *3-Manifolds with nonnegative Ricci curvature*. Invent. Math. **193** (2013), no. 2, 367-375.
- [8] Morrey, C. B. *The problem of Plateau in a Riemannian manifold*, Ann. of Math. **49** (1948), 807-851.
- [9] Morrey, C. B. *Multiple Integrals in the Calculus of Variations*, Springer-Verlag, Berlin (1966).
- [10] Osserman, R. *A proof of the regularity everywhere of the classical solution to Plateau's problem*, Ann of Math. **91** (1970), 550-569.
- [11] Papakyriakopoulos, C. *On Dehn's lemma and asphericity of knots*. Ann. of Math. **66** (1957), no. 1, 1-26.
- [12] Perelman, G. *The entropy formula for Ricci flow and its geometric applications*, arXiv: math/0211159.
- [13] Perelman, G. *Ricci flow with surgery on the three manifolds*, arXiv: math/0303109.
- [14] Perelman, G. *Finite extinction time for the solution to Ricci flow on certain three-manifolds*. arXiv: math/0307245.
- [15] Rado, T. *On Plateau's problem*, Ann. of Math. **31** (1930), 457-469.
- [16] Rosenberg, H. *Constant mean curvature surface in Homogenously regular 3-manifold*. Bull. Austral. Math. Soc. **74** (2006), 227-238.
- [17] Stallings, J. *Group theory and Three dimensional manifolds*. Yale University Press, New Haven (1971).
- [18] Schoen, R.; Yau, S.-T. *Complete three-dimensional manifold with positive Ricci curvature and scalar curvature*, Seminar on Differential Geometry, Ann. of Math Stud., Vol.102,209-228. Princeton University Press, Princeton(1982)
- [19] Schoen, R.; Yau, S.-T. *On the proof of the positive mass conjecture in general relativity*, Commun. Math. Phys. **65** (1979), 45-76.
- [20] Schoen, R.; Yau, S.-T. *Proof of the positive mass theorem. II*, Commun. Math. Phys. **79** (1981), 231-260.
- [21] Whitehead, J.H.C. *A certain open manifold whose group is unity*. Quart.J.Math. Oxford, **6** (1935), 268-279.

UNIVERSITÉ GRENOBLE ALPES, INSTITUT FOURIER, 100 RUE DES MATHS, 38610
GIÈRES, FRANCE

E-mail address: jian.wang1@univ-grenoble-alpes.fr