



HAL
open science

Impact of turning, pod storage and fermentation time on microbial ecology and volatile composition of cocoa beans

Yasmine Hamdouche, Jean Christophe Meile, Marc Lebrun, Tagro Guehi, Renaud Boulanger, Corinne Teyssier, Didier Montet

► To cite this version:

Yasmine Hamdouche, Jean Christophe Meile, Marc Lebrun, Tagro Guehi, Renaud Boulanger, et al.. Impact of turning, pod storage and fermentation time on microbial ecology and volatile composition of cocoa beans. Food Research International, 2019, 119, pp.477 - 491. 10.1016/j.foodres.2019.01.001 . hal-03486184

HAL Id: hal-03486184

<https://hal.science/hal-03486184v1>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Impact of turning, pod storage and fermentation time on microbial ecology and volatile**
2 **composition of cocoa beans.**

3 **Yasmine Hamdouche^{1,2,3}, Jean Christophe Meile⁴, Marc Lebrun², Tagro Guehi⁵, Renaud**
4 **Boulangier², Corinne Teyssier², Didier Montet².**

5 ¹ University of M'Hamed Bougara of Boumerdes, Avenue de l'indépendance, 35000,
6 Boumerdes, Algeria.

7 ² UMR 95 QualiSud / Cirad/ University of Montpellier, TA B-95 /16, 73 rue J-F Breton,
8 34398 Montpellier cedex 5, France.

9 ³ LBSM laboratory (Laboratory of Biology and microbial systems), ENS (Normal Superior
10 School), Kouba, 16 000, Algiers, Algeria

11 ⁴ UMR 95 QualiSud / Cirad Station de Ligne-Paradis 7 chemin de l'IRAT, 97410 Saint-Pierre
12 Reunion, France.

13 ⁵ Laboratory of Microbiology and Molecular Biology, Department of Food Science and
14 Technology, University of Nangui Abrogoua, Abidjan, Ivory Coast.

15

16 Corresponding author: yasminehamdouche@gmail.com

17

18 **Key words:** cocoa beans fermentation, microbial communities, volatile compounds, SPME-
19 GC-MS, PCR-DGGE.

20

21

22

23 **Abstract**

24 Cocoa quality depends on several parameters, such as cocoa variety, environmental growth
25 conditions, cultivation technique, and post-harvest treatments applied to cocoa beans. In this
26 work, we studied the impact of cocoa post-harvest processing on both microbial communities
27 structure and volatile composition. Cocoa beans samples were fermented in wooden boxes in
28 Ivory Coast at different time intervals with turning and without turning, and derived from
29 pods stored for two different duration times. Cocoa beans were analyzed using a molecular
30 fingerprinting method (PCR-DGGE) in order to detect variations in microbial communities'
31 structure; this global analysis was coupled to SPME-GC-MS for assessing cocoa **volatile**
32 **profiles**. The results showed that the main parameter that influenced microbial communities
33 structure was fermentation time, followed by turning, whereas, pods storage duration had a
34 minor impact. Similar results were obtained for aromatic profile, except for pods storage
35 duration that significantly affected volatile compound production. Global statistical analysis
36 using Canonical Correspondence Analysis (CCA), showed the relationship between microbial
37 communities and volatile composition. Furthermore, this study allowed the identification of
38 discriminating microbial and chemical markers of cocoa post-harvest processing.

39

40

41

42

43

44

45

46 **1. Introduction**

47 Cocoa is ranked among the most traded agricultural products in the world. Cocoa beans
48 production is estimated to 4,552,000 tons in 2016/2017; 73% of which is produced by Africa,
49 mainly by Ivory Coast, which is the cocoa first producer, followed by Ghana, Cameroon and
50 Nigeria (ICCO, 2017).

51 Cocoa has an intrinsic quality potential that depends on several parameters, such as cocoa
52 variety, age of the cocoa tree, soil chemical composition, post-harvest treatments, and
53 industrial process (Kongor et al., 2016). In particular, postharvest processing applied on beans
54 can enhance or degrade the sanitary and/or organoleptic qualities of cocoa (Saltini et al.,
55 2013). The transformation of cocoa goes through three main stages (pods opening,
56 fermentation, and drying). The crucial step is fermentation, which is a spontaneous process
57 involving microbial flora succession responsible for aroma precursor's production inside the
58 beans through biochemical reactions. Fermentation lasts usually between 5 to 7 days. Yeasts
59 are the first type of microorganism involved followed by Lactic Acid Bacteria (LAB), and
60 then Acetic Acid Bacteria (AAB) (Schwan and Wheals, 2004). Several recent studies tried to
61 understand the role of these microorganism groups in cocoa fermentation and to control this
62 phenomenon by inoculating selected strains previously isolated from cocoa (Lefeber et al.,
63 2012, Crafacck et al., 2013). The major results of these works showed that yeast strains have a
64 greater impact on the improvement of cocoa aromatic quality when compared to bacterial
65 strains (Crafack et al., 2013; 2014; Ho et al., 2014; Koné et al., 2016; Hamdouche et al.,
66 2017).

67 Several studies on the impact of postharvest treatments on cocoa beans composition showed
68 that fermentation types (Papalexandratou et al., 2011; Visintin et al., 2016) and times
69 (Hamdouche et al., 2015) significantly affected the microbiota associated with cocoa beans,

70 while pods storage (pulp preconditioning) and turning affected chemical composition and
71 physical appearance of cocoa (Nazaruddin et al., 2006; Guehi et al., 2010).

72 Aroma profile significantly contributes to cocoa quality and thus is crucial for the
73 acceptability of chocolate. This profile varied mainly with fermentation and drying time
74 (Rodriguez-Campos et al., 2011; 2012). Aroma precursors previously formed during
75 fermentation and drying steps were highly developed during roasting. Roasting reduces
76 acidity by volatile acids evaporation and removes bitter, acidic and astringent flavors (Beckett
77 2000, Ramli et al., 2006). Maillard reactions generate a valuable source of aromatic
78 compounds as pyrazines, pyrroles, furans and thiazoles (Belitz et al., 2009). The 2-
79 methylpyrazine and 2, 3, 5, 6 tetramethylpyrazine are the main volatile compounds present in
80 roasted cocoa. Both are responsible for the sweet and strong cocoa flavor respectively
81 (Lefeber et al., 2012).

82 The most used techniques for volatile compounds extraction are distillation, direct solvent
83 extraction, solid phase extraction (SPE), headspace methods and solid phase micro-extraction
84 (SPME). After concentration, gas chromatography is the most suitable analytical method.
85 This technique allows the separation of a very large number of compounds on a fused silica
86 capillary column. Different detection methods coupled with gas chromatography allow
87 volatile compounds detection and identification: flame ionization, flame photometry, atomic
88 emission, and mass spectrometry. SPME-GC-MS for the headspace analysis of volatile
89 compounds allowed sensitive and representative analysis of cocoa products with high
90 reproducibility (Ducki et al., 2008).

91 A recent work revealed that cocoa aroma compounds composition varied according to cocoa
92 variety (Menezes et al., 2016), but no study showed the impact of pod storage and turning on
93 microbial ecology and the associated impact on aroma profiles. The main aims of this original
94 study were: (i) to measure the impact of the variation of three parameters (pod storage,

95 fermentation time and turning) during cocoa post-harvest processing on microbial and volatile
96 profiles; (ii) to identify discriminating microbial and/or aroma markers of each process; (iii) to
97 establish relationship between microbial communities associated with cocoa beans and
98 volatile compounds produced.

99 **2. Material and Methods**

100 **2. 1. Cocoa beans sampling**

101 Four hundred cocoa pods were harvested in November 2014 in Akoupé (Ivory Coast). The
102 number of harvested pods was divided in two batches (200 pods each), the first batch was
103 stored during 2 days and the other one during 8 days. After storage, all pods were opened
104 using a machete. For each pods opening delay, approximately 30 kg of pulp-bean mass were
105 extracted manually, recovered and distributed in two wooden boxes for fermentation during 7
106 days. Two types of processing were applied on these cocoa beans: fermentation with and
107 without turning. The turning was done at 2 and 4 fermentation days, this choice was made
108 following the artisanal practices applied. One kilogram of cocoa beans was collected at 2, 4,
109 5, 6 and 7 days of fermentation and sun dried during 7 days on wooden drying racks. During
110 drying step, no turning was done. The sampling was performed in sterile conditions by using
111 sterile gloves and plastic bags. Samples were sent by plane to Cirad laboratory, Montpellier,
112 France and stored at room temperature. The microbial and chemical analyses were carried out
113 immediately.

114 **2. 2. Microbial ecology analysis**

115 **2. 2. 1. Microbial DNA extraction**

116 Microbial DNA was extracted from 10 g of fermented and dried cocoa beans. Two technical
117 repetitions were assessed for each sample. The extraction was performed according to the
118 method described previously by Hamdouche et al. (2015). The quantity and purity of

119 extracted DNA were measured by a UV spectrophotometer (Biospec-Nano), and by
120 electrophoresis through a 0.8% agarose gel in 1X TAE buffer (containing 40 mM Tris-HCl,
121 pH 7.4, 20 mM sodium acetate, 1.0 mM Na₂-EDTA; Eppendorf) with a 1 Kb molecular
122 weight ladder (Promega). The agarose gels were run at 100 V for 30 min and stained for 15
123 min in Gel Red solution (50 mg.mL⁻¹; Biotium), then observed and photographed on a UV
124 transilluminator, using a black and white CCD camera (Scion Company) and Gel Smart 7.3
125 system software (Clara Vision).

126 2. 2. 2. PCR amplification of extracted DNA

127 For bacterial communities analyses: A 160 bp fragment of the V3 variable region of 16S
128 rDNA was amplified using gc-338f (5'-GCG CCG CCG CGC GCG GCG GGC GGG GCG
129 GGG GCA CGG GGG GCA TCC TAC GGG AGG CAG CAG-3', Sigma) and 518r (5'-ATT
130 ACC GCG GCT GCT GG-3', Sigma) DNA primers (Leesing, 2005; Hamdouche et al., 2015;
131 2016). A 40-bp GC-clamp (Sigma) was added to the forward primer in order to ensure that the
132 DNA fragment will remain partially double stranded and that the region screened is in the
133 lowest melting domain (Sheffield et al., 1989). Each mixture (final volume 50 µL) contained
134 about 100 ng of DNA template, DNA primers at 0.2 mM, deoxyribonucleotide triphosphate
135 (dNTPs) at 200 µM, 1.5 mM MgCl₂, 5 µL of 10X of Taq reaction buffer MgCl₂ free and 1.25
136 unit of a-Taq polymerase (Promega). In order to increase the specificity of amplification and
137 to reduce the formation of spurious by-products, DNA was amplified by a “touchdown” PCR
138 previously described by Hamdouche et al. (2015).

139 For fungal communities analyses: a 250 bp fragment of D1/D2 region of the 26S rDNA was
140 amplified using universal DNA primers gc-NL1 (5'-CGC CCG CCG CGC GCG GCG GGC
141 GGG GCG GGG GCC ATA TCA ATA AGC GGA GGA AAA G-3') and LS2 (5'-ATT CCC
142 AAA CAA CTC GAC TC-3', Sigma) (Cocolin et al., 2000). A 30-bp GC-clamp was added to

143 the forward primer. PCR mixture performed as previously described for bacterial DNA
144 analyses. The amplification program was described previously by Hamdouche et al. (2015).

145 Electrophoresis in 2% (w/v) agarose gel with TAE 1X buffer stained with Gel Red (as
146 described above) was used to verify and quantify PCR products using a standard (DNA mass
147 ladder 100 bp, Promega). The migration lasted about 30 min at 100 V.

148 **2. 2. 3. Denaturing gradient gel electrophoresis (DGGE) analysis**

149 PCR products were separated by DGGE, using a Bio-Rad Dcode universal mutation detection
150 system (Bio-Rad Laboratories, USA) according to the procedure described by Muyzer et al.
151 (1993) and improved by Leesing (2005). Forty microliters (μL) of each PCR product were
152 loaded onto 8% (w/v) polyacrylamide gels (acrylamide/N,N'-methylene bisacrylamide,
153 37.5/1, Promega) in 1X TAE.

154 Electrophoresis experiments were performed at 60°C using a denaturing gradient ranging
155 from 30 to 60% (100% corresponded to 7 M urea and 40% (v/v) formamide, Promega). The
156 migration was carried out at 20 V for 10 min and then at 80 V for 12 h. Then the gel was
157 stained for 30 min with Gel Red solution and then photographed as described above in 2.2.1
158 section.

159 **2. 2. 4. Purification and identification of DGGE bands**

160 Detected DNA bands were cut from the DGGE gel with a sterile scalpel. The DNA of each
161 band was then eluted in 100 μL TE buffer (10 mM Tris-HCl; pH 8.0, Promega) at 4°C
162 overnight. The DNA purification was carried out according to the protocol described by
163 Hamdouche et al. (2015; 2016). Bacterial and fungal purified DNA bands were amplified
164 using 338f and NL-1 (without GC clamp) and 518r and LS-2 primers respectively. PCR
165 amplicons were sent for sequencing to GATC Biotech (Germany).

166 The raw sequences data were edited using Bioedit Software and compared to the GenBank
167 database using the BLAST program (Altschul et al., 1997) (<http://www.ncbi.nlm.nih.gov/BLAST/>). Sequences having a percentage of identity of 97% or greater were considered
168 to belong to the same species (Palys et al., 1997).
169

170 **2. 2. 5. Image and statistical analysis**

171 In each lane, banding patterns were standardized with two bacterial reference DNA patterns
172 (*Lactobacillus plantarum* and *Escherichia coli* DNAs) for bacteria and two yeast DNA
173 reference patterns of yeasts (*Lodderomyces elongisporus* and *Candida apicola* DNAs). Each
174 band relative positions to reference DNA were measured and recorded.

175 The images of DGGE gels were analyzed using ImageQuant TL software V. 2003
176 (Amersham Biosciences, USA). In DGGE analysis, the revealed banding pattern is considered
177 as an image of all the major microbial populations. An individual band refers to a unique
178 ‘sequence type’ or phytotype (Kowalchuk et al., 1997). DGGE fingerprinting was marked by
179 the presence and absence of co-migrating bands and their intensity using ImageQuant TL
180 software V. 2003.

181 Pair wise community similarities were quantified using Dice similarity coefficient
182 (Heyndrickx et al., 1996), then generated data were exploited by Principal Component
183 Analysis (PCA) and Canonical analysis of correspondence (CAC) using XLSTAT (V. 2014).
184 The bands intensities (or volume as expressed in “grey level x pixel” unit) measured by
185 ImageQuant were statistically compared using ANOVA.

186 CAC is a method that encompasses several statistical analyses; Ter Braak (1986) developed it
187 for the first time. In our case, it was used to describe the linear relationships that exist between
188 two groups of variables measured on the same individuals. The first group represents the
189 variables to be explained (pod storage duration, fermentation time and turning) and the second

190 group shows the explanatory variables that are the microbial communities (DNA bands
191 identified by PCR-DGGE).

192

193

194 **2. 3. Volatile compound analysis**

195 **2. 3. 1. Sample preparation and SPME-HS conditions**

196 The shell and the bean were both analysed, because the objective of the work was to evaluate
197 the volatile composition of exported merchant cocoa (fermented and dried beans). In food
198 industry, before the shelling, fermented and dried beans are roasted in order to develop the
199 chocolate aromas, these aromas depend essentially on the precursors born during
200 fermentation.

201 Two hundred grams of unshelled dried cocoa beans from Ivory Coast were put in liquid
202 nitrogen and ground using a waring blender (SEB, France), to obtain cocoa flour, which was
203 stored at - 80°C until analysis. Three grams of cocoa powder were introduced into vial (10
204 mL), 30 µL of internal standard of 1-butanol (202.5 mg/L) (Sigma) were added, and the vial
205 was sealed with a septum cap. The extraction of three technical repetitions was performed
206 using a 50/30 µm divinylbenzene/carboxene/polydimethylsiloxane fiber (DVB/CAR/PDMS,
207 Supelco), and using the technique of solid phase micro-extraction in the headspace (SPME-
208 HS) previously described by Rodriguez-Campos et al. (2012) and modified in our lab by
209 Koné et al. (2016). These authors had conditioned the fiber at 250°C for 3 min then exposed it
210 to the sample headspace at 50°C for 45 min.

211 **2. 3. 2. Gas Chromatography Analytical Conditions**

212 The analysis of the volatile compounds extracted from cocoa powder was performed using an
213 Agilent 6890N gas chromatography–mass spectrometer (GC-MS) in automatic injection
214 mode, on a polar capillary column DBWAX, 30 m length \times 0.25 mm internal diameter \times 0.25
215 μm film thickness (Agilent, Palo Alto, CA, USA). The GC-MS conditions have been
216 established in Cirad laboratory, the injection temperature was 250°C. The GC oven
217 temperature was initially set at 40°C for 5 min, increased to 140°C at a rate of 2°C/min and
218 then increased at a rate of 10°C/min to 250°C. The carrier gas was high purity Helium at 1
219 $\text{mL}\cdot\text{min}^{-1}$. Splitless injection mode was used at 250°C for 2 min. The selective mass detector
220 was a quadrupole (Hewlett Packard, Model 5973), with an electronic impact ionization
221 system at 70 eV and at the temperature source at 230°C.

222 **2. 3. 3. Volatile compounds identification**

223 For the identification of compounds, pure standards were not used, **this is why the**
224 **identification was just tentative.** The identification was done: (i) by comparing the mass
225 spectra with commercial database Wiley275.L, HP product no. G1035A); (ii) and by
226 comparing of the linear retention index calculated (RI) with those found in the literature data.
227 Retention index (RI) was calculated in the same chromatographic conditions using series of n-
228 alkane C8-C20 (Supelco, Bellefonte, USA). Peak areas were used for relative quantification
229 of compounds using the MSD Chemstation software (version E.02.02.1431, Agilent
230 Technologies).

231 **2. 3 .4. Statistical analysis**

232 The chromatographic data obtained by SPME-GC-MS were statistically analyzed using
233 ANOVA and Principal Component Analysis (PCA). As described previously, Canonical
234 analysis of correspondence (CAC) was applied to GC-MS and DGGE data, in order to

235 describe relationship between microbial communities (explanatory variables) and volatile
236 compounds produced (variables to be explained).

237 **3. Non-volatile organic acids analysis**

238 **3.1. Organic acids extraction**

239 The extraction was carried out according to the protocol of Holm et al. (1993). Five grams of
240 cocoa powder were mixed with 50 mL of warm distilled water (75°C) in a 100 mL flask. The
241 resulting suspension was well mixed and then cooled to room temperature. The volume was
242 supplemented with distilled water up to 100 mL, then filtered through a pleated filter paper
243 (Whatman) and a 0.45 µm filter (Sigma-Aldrich), then transferred to an HPLC vial (1 mL)
244 (Thomson).

245 **3.2. Organic acids quantification by HPLC**

246 Concentrations of acetic acid, citric acid, lactic acid, malic acid, oxalic acid and tartaric acid
247 were determined by a High Performance Liquid Chromatography (HPLC) with conductivity
248 using AS11-HC column (Dionex). The mobile phase, at a flow rate of 0.6 mL.min⁻¹ consisted
249 of ultrapure water (eluent A), NaOH 0.1 N (eluent B) and NaOH 0.002 N (eluent C). A
250 volume of 0.02 mL for each sample was injected and run together with external standards. All
251 samples were analyzed in triplicate and the data were controlled and processed with
252 ClarityChrom software and analyzed using ANOVA.

253 **3. Results and discussion**

254 **3. 1. Microbial community structures**

255 **3. 1. 1. Variation in bacterial community structures**

256 Microbial DNA profiles were generated by DGGE from all cocoa fermentations carried out,
257 with and without turning, with a delay of pod breaking of 2 and 8 days, respectively. Bacterial

258 DNA profile bands diversity at different time intervals of fermentation (Fig. 1.a and b). After
259 two fermentation days, a few DNA bands (Y, B V, W) were observed and identified by
260 sequencing as belonging to *Enterobacteriaceae*, *Lactobacillus fermentum*, and *Acetobacter*
261 *pasteurianus* taxa respectively (Fig. 1.a,b ; Table 1). At days 4 and 5, additional DNA bands
262 (A, E, F, J, L, M, N) were detected and identified as belonging to *Lactobacillus curvatus*,
263 *Bacillus megaterium*, *Bacillus sp*, *Acetobacter sp* and *Providencia sp species*, respectively.
264 All eight bacterial taxa remained detected until the last day of fermentation. This is
265 concordant with previously reported data (Lefeber et al., 2011; Hamdouche et al., 2015),
266 which showed that bacterial communities of cocoa undergo drastic changes after 4 days of
267 fermentation and remain pretty similar throughout the process length.

268 Moreover, at the shortest fermentation time, only *L. fermentum*, *A. pasteurianus* and species
269 belonging to *Enterobacteriaceae* taxa were detected and confirmed the presence of LAB and
270 AAB in the initial phase of fermentation (Schwan and Wheals, 2004; Papalexandratou et al.,
271 2013; Hamdouche et al., 2015).

272 Comparison of bacterial DNA profiles of fermented cocoa beans with and without turning
273 showed that the majority of revealed bands were common in the two types of fermentation
274 processing whatever the pod storage time. However, some bands (M, E, F) identified as
275 *Bacillus* genus were mainly revealed in DGGE profiles obtained from cocoa beans fermented
276 with turning (Fig. 1.a,b ; Table 1). Furthermore, some bands (G, H) were detected only in
277 fermentation with turning and were identified as belonging to *Acetobacter sp*. These
278 observations were explained by the better development of aerobic microorganisms due to
279 aeration obtained by turning. The abundance of *Bacillus* genera populations seems to be the
280 consequence of temperature elevation after 2 days and at the turning time, which promoted
281 the development of thermophilic and aerobic species (Lima et al., 2011). Previous works,
282 carried out on fermentation of different cocoa beans masses, showed that the fermentation of

283 the large cocoa bean-pulp mass decreased the aeration, which slows down microbial activities
284 (Camu et al., 2008; Guehi et al., 2010). It is important to note that *A. pasteurianus* species (V,
285 W) was not present during fermentation with turning, at 5, 6 and 7 days (Fig. 1.b-Table 1).
286 This result was unexpected because *A. pasteurianus* was the most prevalent AAB species
287 involved in spontaneous cocoa from Ivory coast, Brazil and Malaysia (Hamdouche et al.,
288 2015, Papalexandratou et al., 2011; 2013) and was tested as starter culture in cocoa
289 fermentation (Moens et al., 2014; Lefeber et al., 2010 ; 2012). First of all, we thought that this
290 unexpected result came from technical error, but the most surprising thing about this
291 observation, was the correlation of this result with the one obtained from volatile compounds
292 analysis. The disappearance of *A. pasteurianus* species was correlated with acetic acid content
293 decreases during fermentations with turning. If the technical error is sparing, the result could
294 be explained by the fact that the temperature increased during fermentation with turning (by
295 increasing activity AAB). In our fermentation assays, the temperatures could have risen such
296 as it became lethal for this thermosensible species. However, since, temperatures were not
297 measured in our studies, we could not conclude about this hypothesis.

298 When comparing DGGE bacterial profiles obtained from stored pods during 2 and 8 days,
299 DNA bands (R, K, X) specific to cocoa pods stored 8 days were detected and identified as
300 belonging to *Klebsiella pneumoniae*, *Bacillus sp*, and *Acinetobacter sp* (Fig. 1.b compared to
301 Fig. 1.a). These bacterial taxa could be present on pods surface and developed during storage
302 time, which rendered them detectable by PCR-DGGE. To our knowledge, no work has
303 investigated the pod storage effect on cocoa microbial ecology except the study of our
304 colleagues Kedjebo et al. (2016) who studied the effect of post-harvest treatment on the
305 occurrence of ochratoxin A (OTA) in raw cocoa beans from Ivory Coast. They showed that
306 OTA content in raw cocoa beans increased with pods storage duration (0, 4 and 7 days) which
307 reflected the increase of ochratoxinogenic fungi activity (*Aspergillus* strains) during cocoa

308 pods storage time. Therefore, their conclusion confirmed our results that showed a highest
309 microbial contamination with long pod storage after fermentation.

310 Principal Component Analysis (PCA) was applied on bacterial DGGE profiles to represent
311 the dataset variance. We observed that fermented cocoa samples tended to group according to
312 fermentation time along the first component axis (F1 representing 53.44% of the total
313 variance). For 2 day storage samples, two main groups containing cocoa beans samples turned
314 fermented during 2 and 4 days on the left side of the axis and those fermented during 5, 6 and
315 7 days on the right side of the axis could be observed (Fig. 2). The picture is similar for 8 days
316 storage of non-turned samples. Eight-day storage of turned samples appeared as two main
317 groups: 2 day fermented on the left side of the axis and the rest of the samples on the right
318 side. 2 days storage unturned samples tend to group according to the fermentation duration.

319 The secondary axis F2 (12.58% variance) separated groups of cocoa beans samples fermented
320 by turning (bottom part) from those fermented without turning (upper part, Fig.2). This result
321 showed that bacterial communities varied mainly according to fermentation time followed by
322 processing (turning). PCA did not show an obvious effect of storage time on bacterial ecology
323 of cocoa comparing with the impact of the two other parameters (fermentation time and
324 turning). The strong dynamics and activity of the microbial populations observed during the
325 fermentation does not seem to be linked to any other environmental parameter.

326 **3. 1. 2. Variation in fungal community structures**

327 During cocoa fermentations, dominant species (1, 10, and 11) were detected on DGGE
328 profiles and identified as belonging to *Hanseniaspora opuntiae*, *Pichia kudriavzevii* and
329 *Pichia manshurica* species (Fig. 3 - Table 2). These species had previously been detected in
330 fermented cocoa beans from the same location in Ivory Coast (Hamdouche et al., 2015; Koné
331 et al., 2016) and in other countries and continents (Crafack et al., 2013; de Melo Periera et al.,

332 2013; Papalexandratou et al., 2013). Minor variations were observed comparing DGGE
333 profiles obtained from cocoa fermented with the two processes due to intrinsic variations in
334 fungal diversity during fermentation (Fig. 3.a.b). Generally, we did not observe specific
335 fungal communities associated to a process probably due to the fact that turning or aeration do
336 not drastically favor fungal populations growth.

337 In DGGE profiles obtained from fermented cocoa beans previously stored longer (8 days),
338 four specific bands were detected (4, 5, 7, 9) and identified as *Hanseniaspora uvarum* and
339 *Pichia sp* (Fig. 3.b). The results of PCR amplifications coupled to DGGE profiles of both
340 bacterial and yeast communities showed more DNA bands in cocoa pods stored longer (8
341 days), which probably means that microbial species abundance increased with storage time
342 (as discussed above with bacterial data).

343 PCA on fungal communities associated with cocoa, represented by F1 and F2 axis (58. 26%)
344 did not allow to discriminate cocoa beans samples, this result was expected, as mentioned
345 before (Fig. 1 in supplementary material).

346 Canonical analysis of correspondence (CAC) was applied in order to describe the relation
347 between variations in microbial community structures and changes in cocoa post-harvest
348 parameters. CAC results showed that microbial communities associated with turning
349 parameter consisted mainly of bacterial communities: *Bacillus*, LAB and *Acetobacter* species
350 (Fig. 4). This result was expected for AAB because they are aerobic microorganisms, which
351 are usually affected by the aeration. For *Bacillus*, it could be explained by the increasing of
352 the temperature after turning due to the thermophilic aptitude of this genus. No microbial
353 species was found linked to the parameter of pods storage time (Fig. 4) as pods storage
354 duration was not related to the composition of microbiota.

355 **3. 2. Volatile compounds analysis**

356 The results obtained on the volatile compounds analysis might not be accurate because of the
357 limit of the identification and quantification method used.

358 Twenty-six volatile compounds were detected in fermented and dried cocoa beans by SPME-
359 GC-MS (Table 3). These compounds were separated in 7 families: aldehydes, ketones, esters,
360 alcohols, acids, pyrazines and phenols.

361

362

363 **3. 2. 1. Variation of aldehyde compounds**

364 Four aldehydes compounds were identified on all fermented cocoa beans analyzed, which
365 were isobutanal, 2 and 3-methylbutanal, benzaldehyde. Benzaldehyde was the dominant
366 aldehyde, followed by 2 and 3-methylbutanal in all cocoa fermentations (Fig. 5.a.g.m and Fig.
367 6.a.g). These aldehydes were previously identified on fermented cocoa beans (Serra-Bonvehi,
368 2005; Rodriguez-Campos et al., 2012; Ho et al., 2014). In all fermentations, aldehydes content
369 increased with fermentation time. These results were correlated with those obtained by
370 Rodriguez-Campos et al. (2012). Statistical test (ANOVA) showed significant differences in
371 aldehydes level ($p < 0.05$) between the first stage of fermentation (2 and 4 days) and the last
372 stage (5, 6 and 7 days) (Fig. 5.a.g.m and Fig. 6.a.g).

373 ANOVA test showed significant differences for aldehydes content between cocoa beans
374 fermented with and without turning (Fig. 7.a.b), with the highest level obtained in the case of
375 fermentation with turning at 7 days. These observations were probably related to the presence
376 of LAB that were detected throughout the fermentation stages mostly in the case of turning
377 (Fig. 1.a and b). Aldehydes compounds could be produced by enzymatic reactions from
378 amino acids mainly by LAB (Jinap et al., 1994). The 2 and 3-methylbutanal are usually
379 produced from isoleucine and leucine, while benzaldehyde is produced from phenylalanine by

380 LAB (Jinap et al., 1994; Smit et al., 2005; Bonnarme et al., 2004). During beans mixing,
381 temperature increased and reached its maximum (Camu et al., 2008), which increased the
382 enzymatic activity. As regard to pod storage time effect, the aldehydes level had not
383 significant difference ($p>0.05$) between the two pod storage delays (Fig. 7.a.b).

384 **3. 2. 2. Variation of ketone compounds**

385 For this family, the detected compounds were 2-heptanone, 2-pentanone, 2-nonanone,
386 acetophenone, and acetoin. This later had been found as the dominant ketone in all cocoa
387 samples (Fig. 5.b.h.n and Fig.6.b.h) as reported by Rodriguez-Campos et al. (2011). In
388 general, acetoin quantity decreased with fermentation time. This compound is mainly
389 produced from pyruvate by alcoholic fermentation and from 2,3-butanediol by yeast
390 (Pretorius, 2000). Acetoin abundance in the first stage of fermentation could be correlated to
391 yeasts intervention in this stage. The quantity of others ketones varied slightly during
392 fermentation. These compounds were naturally present in cocoa beans (Ho et al., 2014) and
393 their production was not affected by fermentation, except for 2-pentanone, which was
394 probably used as substrate by microorganisms during fermentation. Ketones were
395 significantly important ($p<0.05$) in cocoa beans fermented without turning (Fig. 7.a,b). Some
396 ketones that were probably present naturally (Afoakwa *et al.*, 2008, Rodriguez et al., 2012),
397 were degraded and catabolized in the case of the aerated fermentation process, where more
398 diversity and abundance of bacteria were observed by PCR-DGGE (Fig. 1.a). There was no
399 difference in ketones level for the two pod storage durations, except in the case of
400 fermentation with turning, whither the dominance of ketones was observed in pods stored at
401 the shorter time (2 days) (Fig. 7.a.b).

402 **3. 2. 3. Variation of alcohol compounds**

403 The alcohols compounds found in cocoa beans samples were: ethanol, 2-pentanol,
404 isopentanol, 2-heptanol, 2,3-butanediol, and 2-phenethylethanol. Among them, 2-
405 phenethylethanol and 2,3 -butanediol were dominant during the four types of fermentations
406 (Fig. 5.c.i.o and Fig. 6.c.i); as already showed by Rodriguez-Campos et al. (2011). Ethanol
407 was surely produced by the majority of yeasts (Ho et al., 2014, Koné et al., 2016). It was
408 probably the same in this study where it was generated by all detected yeasts by PCR-DGGE
409 (Fig. 3). Some alcohols could be elaborated by both bacteria and yeasts such as 2,3-
410 butanediol. Ethanol content presented significant differences ($p < 0.05$) between fermentation
411 times (2 and 4 days) and (5, 6, and 7 days) (Fig. 5.c.i.o and Fig. 6.c.i). Its concentration was
412 higher during the first four days of fermentation and then decreased rapidly, because this
413 compound is produced at the first (anaerobic) fermentation stage, which lasted 48 h,
414 subsequently it was oxidized in acetic acid (Schwan and Wheals, 2004) mainly by AAB
415 during the second (aerobic) stage.

416 The abundance of these alcohols was significantly different ($p < 0.05$) during the different
417 fermentation process, with the highest level obtained in cocoa fermented beans without
418 turning mainly for ethanol (Fig 7.a,b, Fig. 5.c.i.o and Fig. 6.c.i). In aerated conditions, the
419 contribution of oxygen allows alcohols oxidation. For example, most alcoholic compounds
420 are oxidized to ester compounds, as phenethylethanol that is a precursor of aldehyde and is
421 oxidized to phenylethyl acetate with the presence of oxygen by LAB (Smit et al., 2005).
422 Furthermore, alcohol compounds could be evaporated with the aeration brought by turning
423 process. This result agreed with those obtained by Camu et al. (2008) that showed the
424 evaporation of ethanol during the aeration of cocoa beans mass, as well as favoring the AAB
425 growth that oxidized ethanol into acetic acid.

426 Total alcohols level was significantly higher in cocoa obtained from pods stored longer (Fig
427 7.a.b) as 2,3-butanediol (Fig 5.c.i.o and Fig 6.c.i). This may be explained by the studies of

428 Afoakwa et al. (2013a; 2013b) that showed the conversion of total sugars with pod storage in
429 alcohol. Furthermore, Meyer et al. (1989) observed that increasing pod storage time before
430 fermentation improved subsequent microbial aeration, prevented LAB development/growth
431 and increased yeasts abundance and activity. This result could be explained by the fact that
432 alcoholic fermentation may begin during the long storage of the pods injured during the
433 harvest, which are naturally or subsequently contaminated by yeasts.

434 **3. 2. 4. Variation of ester compounds**

435 The six detected esters were ethyl acetate, isobutyl acetate, isoamyl acetate, phenylethyl
436 acetate, methyl isopentanoate, and methyl isovalerate (Fig .5.d.j.p and Fig. 6.d.j). Ethyl
437 acetate was the dominant ester in all fermentations. According to Rodriguez-Campos et al.
438 (2012), esters concentration decreased with fermentation time, but in our case, no significant
439 difference was observed ($p>0.05$) depending on fermentation time except for ethyl acetate,
440 which dominated in the first stage of fermentation (2 days) (Fig .5.d.j.p and Fig. 6.d.j). This
441 ester seems to be produced only at the beginning of cocoa fermentation as reported previously
442 (Rodriguez-Campos et al., 2011), and also in wine fermentation (Rojas et al., 2003) mainly in
443 the first day of fermentation. This may be due to the abundance of glucose and ethanol during
444 this stage, because they are the main substrates for ethyl acetate synthesis (Yong et al., 1981).
445 Moreover, Ethyl acetate is a product of esterification from acetic acid and ethanol (Pretorius,
446 2000); this concept explains the correlation between the amount of this ester, acetic acid and
447 ethanol quantities mainly in the case of fermentation with aeration, in which a logical
448 evolution of the quantities was observed for the most of volatile compounds detected in this
449 study. A large quantity of these compounds was observed at the beginning of fermentation
450 and decreased sharply after 2 days (Fig. 5.c.i.o.e.k.q and 6.c.i.e.k). The concentration of this
451 ester was significantly higher in cocoa fermented without turning (Fig 7.a,b). Esters were
452 synthesized by yeasts from alcohols during anaerobic phase (Rojas et al., 2002), which

453 explained their dominance in fermentation without aeration. This observation could explain
454 the high level of esters in this process. Concerning pods storage time, esters were higher as
455 mainly methyl isovalerate in cocoa obtained from pods, which were stored less time (Fig.
456 7.a.b). That could be explained by the decrease of sugar and protein during cocoa pod storage
457 (Guehi et al., 2010, Afoakwa et al., 2013) that could be substrates for esters synthesis.
458 Substrates reduction during long storage (8 days) allowed probably the decrease of the esters
459 content.

460 3. 2. 5. Variation of acid compounds

461 Three acids were detected in fermented cocoa beans, of which acetic acid was largely
462 dominant comparing with isobutyric and isovaleric acids. Acetic acid quantity decreased
463 significantly during fermentation realized with turning (Fig. 5.e.k.q and Fig. 6.e.k). In the case
464 of fermentation without turning, the evolution of acetic acid was irregular because its quantity
465 was not stable, decreased and increased during fermentation (Fig. 7.a,b). The illogical
466 evolution of acetic acid amount in this case, could be explained by the fact that there was no
467 beans turning, therefore no homogenization in the metabolites produced and beans
468 composition. Whatever the pods storage delay, a normal evolution of acetic acid amount
469 during fermentation with turning was observed. According to the literature, acetic acid
470 increased and reached its maximum around 72h of cocoa fermentation then decreased after 88
471 h (Shwan and Wheal, 2004; Lefeber and al., 2012). This data confirmed our result, which
472 showed that turning of cocoa beans allowed to realize fermentation with a normal evolution
473 over time. The concentration of this acid was significantly higher in cocoa beans fermented
474 without turning. To explain this, one possibility is that residual or minimal acetic fermentation
475 (with AAB growth and/or activity) occurs in unmixed cocoa, and the rise in temperature plus
476 the aeration favors the volatilization nutrients depletion (Camu et al., 2008) during
477 fermentation with turning.

478 During fermentation, assimilation of citric acid by LAB generates a slight rise in pH to 4.5
479 and temperature to 37°C of cocoa pulp (Camu et al., 2007) which create a favorable
480 environment for the growth of AAB involved in the second phase of fermentation (acetic
481 fermentation). During this phase, AAB are responsible of the oxidation of ethanol to acetic
482 acid (Schwan and Wheals, 2004). Our results on acetic acid bacterial communities structure
483 was related to acetic acid production, because low acid acetic level was observed in parallel
484 with the disappearance of the main AAB, which was *A. pasteurianus* (band V and W) (Fig. 1)
485 in the absence of aeration.

486 We also observed that volatile acids content decreased among time in cocoa beans and were
487 very low after a pod storage of 8 days (Fig. 7.a.b). This observation is in agreement with the
488 works of Tomlins et al. (1993) and Jinap et al. (1994), who observed a lower acetic acid
489 content in stored cocoa compared with un-stored cocoa pods. As mentioned previously from
490 the literature, acetic acid is produced by oxidation of the ethanol during the second
491 fermentation phase (aerobic). These results were consistent with those obtained on ethanol
492 (section 3.2.3) which was low after pod storage of 8 days. Therefore, acetic acid content,
493 which is volatile, could vary according to the ethanol content and time.

494 **3. 2. 6. Variation of others volatile compounds**

495 *Pyrazines:* Tetramethylpyrazine was the unique compound belonging to pyrazine family that
496 was detected (Fig. 5.f.l.r and Fig. 6.f.l). In most cases, tetramethylpyrazine compound has
497 been reported to produce characteristics notes of roasted cocoa that is important for flavor
498 quality (Afoakwa et al., 2008; Serra-Bonvehi, 2005). Furthermore, various studies reported on
499 the microbial origin of pyrazines in fermented food as cocoa and soybeans (Selamat et al.,
500 1994; Besson et al., 1997). In our case, tetramethylpyrazine was probably produced by
501 *Bacillus* as *Bacillus megaterium* species that was detected by PCR-DGGE (Fig. 1). They play
502 an important role in producing alkylypyrasines in association with *B. subtilis* during cocoa

503 fermentation. Our results showed that tetramethylpyrazine level had no significant difference
504 during fermentation times, pods storage times and between fermentation processing. Pyrazine
505 compounds were not affected by any of the three processes.

506 *Methoxyphenol*: This volatile phenolic compound was only detected in fermented cocoa beans
507 obtained from pods stored for 8 days and it was significantly ($p < 0.05$) higher in the case of
508 fermentation with turning (Fig. 5.f.l.r, Fig. 6.f.l). Some phenolic compounds were previously
509 detected at the end of cocoa fermentation (6 and 8 days) (Rodriguez-Campos et al., 2012).

510 The presence of this compound only in fermented cocoa beans obtained from long pods
511 storage when more microbial diversity was detected, could be explained by the increase of
512 nutrient degradation, including phenolic acids, thus 2-methoxyphenol elaboration. We also
513 connected the production of this compound to the natural heating during sun drying. In
514 general, phenolic acids are degraded thermically or decomposed by microorganisms into
515 phenols, as 2-methoxyphenol that results from a heat degradation of ferulic acid during coffee
516 and cocoa roasting or kiln drying (Berlitz, 2009).

517 3. 3. Organic acids composition

518 Five non-volatile organic acids were detected and quantified: citric, lactic, malic, oxalic and
519 tartaric acids. The total of non-volatile acids was significantly ($p < 0.05$) higher on cocoa beans
520 fermented without turning (when compared to turned cocoa), especially at the end of
521 fermentation (5, 6 and 7 days) (Fig. 8). This confirmed the results obtained on volatile acids,
522 which are less abundant during fermentation with aeration. The amount of acids is
523 significantly greater ($p < 0.05$) on cocoa beans with short pod storage durations (2 days) (Fig.
524 8). This is consistent with the results obtained on volatile acids by Afoakwa et al. (2013a;
525 2013b) who showed consistent decreases in non-volatile acidity during pod storage.

526 The concentration of volatile and non-volatile acids identified was affected during
527 fermentation probably under microbial activities. Those naturally present in cocoa beans were
528 degraded, mainly citric acid, while others were produced during fermentation, such as acetic
529 and lactic acids, which could be considered as cocoa quality markers.

530 **3. 4. Correlations between microbial ecology and volatile compounds produced in cocoa** 531 **fermentation**

532 In order to measure the impacts of turning during fermentation (fermentation time and pods
533 storage time), PCA was applied on all volatiles compounds detected in fermented cocoa beans
534 (Fig. 9). Statistical analysis showed a global (on the volatile groups) and specific (on volatile
535 compounds) effects. We showed that ketones, alcohols, esters and acids contents were higher
536 in cocoa fermented without turning, while aldehydes were significantly abundant in the
537 process with turning. Pod storage duration had an effect on all chemical families except for
538 aldehydes and pyrazines. ACP separated 4 samples groups according to fermentation time,
539 fermentation process and pod storage time. Principal axis F1 (38.86%) separated cocoa beans
540 samples fermented (5, 6 and 7 days) without turning from the beans fermented with turning
541 (Fig. 9). ACP analysis allowed identifying discriminant compounds that were affected by
542 studied parameters. Discriminant and significant compounds in fermentation without turning
543 were acetic acid, isobutyric acid, 2,3-butanediol, acetoin, phenethyl acetate, isoamyl acetate.
544 In the case of turning, all aldehydes compounds detected and some esters were identified as
545 discriminant compounds. The same axis F1 regrouped samples according to fermentation
546 time, the higher level of acetic acid, ethyl acetate, isobutyl acetate and ethanol at the first
547 stage of fermentation (2 and 4 days) discriminated beans fermented at short term from the
548 other samples. F2 axis (18.45%), separated the two pods storage times starting from 5 days of
549 fermentation, by 2-methoxyphenol and acetophenone that were significant (abundant) in long
550 storage (8 days).

551 Variations in volatile composition were compared with literature and with microbial ecology
552 analyses carried out in our study. Some hypotheses were given about the relationship that
553 could exist between microbial community structures and volatile composition of cocoa beans.
554 CAC was applied on all data obtained in order to determine these relationships (Fig. 10).
555 Microbial taxa detection by DGGE and their correlation with volatile compounds had a
556 similar evolution whatever the fermentation time and the processing applied. It was not
557 possible or rather difficult to relate each volatile group and/or compound to a given microbial
558 taxa, which is explained by the fact that several yeasts or bacteria could produce the same
559 compounds (Ho et al., 2014; Koné et al., 2016). Moreover, the CAC described the relation
560 between *A. pasteurianus* (V, W) acetic acid, and acetoin and between *H. opuntiae* to ethyl
561 acetate and ethanol (Fig. 10). This suggested that acetic acid level was related to *A.*
562 *pasteurianus* abundance. This hypothesis could be verified by the identification of **volatile**
563 **profile** for each microbial species detected during fermentation.

564 **4. Conclusion**

565 This study measured the effect of three cocoa process parameters on both microbial
566 community structures and volatile composition of cocoa beans, and showed that fermentation
567 time has a major impact on the aroma, followed by turning. In our study, pod storage duration
568 (2 days vs 8 days) had a minor impact on cocoa microbial communities. Contrariwise, the
569 effect was pronounced on volatile composition when it was associated to turning. Fermented
570 cocoa beans with desired flavors were obtained with short pod storage times, which promoted
571 a significant presence of esters. Consequently, as expected a short fermentation without
572 aeration could have a negative impact on the cocoa volatile composition, because of the high
573 levels of undesirable compounds such as acetic acid and acetoin.

574 The molecular approach coupled to volatile and statistical analysis provided insight into the
575 relationship between the microbial communities' structure and aroma compound production

576 in fermenting cocoa beans. To better describe the relationship, it would be interesting to carry
577 out the study in a specific way by identifying the aromatic profile of each microbial strains
578 involved in the processes. Consequently, this global analysis allowed the identification of
579 molecular and volatile markers, which could be good indicators of cocoa post-harvest
580 processing. Acetic acid, acetoin, benzaldehyde, 2 and 3-methylbutanal content are related to
581 cocoa fermentation time and aeration type, whereas, the evaluation of esters concentration
582 could be used to distinguish pods storage duration of fermented beans.

583 At this stage, the impact of post-harvest processing was measured on cocoa volatile
584 composition but not specifically on aroma quality. That aspect yet remains to be explored in
585 the future.

586 **Acknowledgments**

587 This work was supported by DESI grant from CIRAD France (support of PhD students) and
588 Y. Hamdouche was supported by a PhD fellowship from HUBBARD company.

589

590 **References**

591 Afoakwa, E. O., Kongor, J. E., Takrama, J. F., & Budu, A. S. (2013a). Changes in
592 acidification, sugars and mineral composition of cocoa pulp during fermentation of pulp pre-
593 conditioned cocoa (*Theobroma cacao*) beans. *International Food Research Journal*, 20(3),
594 1215-1222

595 Afoakwa, E. O., Paterson, A., Fowler, M., & Ryan, A. (2008). Flavor formation and character
596 in cocoa and chocolate: a critical review. *Critical reviews in food science and nutrition*, 48(9),
597 840-857.

598 Afoakwa, E. O., Quao, J., Takrama, J., Budu, A. S., & Saalia, F. K. (2013b). Chemical
599 composition and physical quality characteristics of Ghanaian cocoa beans as affected by pulp
600 pre-conditioning and fermentation. *Journal of Food Science and Technology*, *50*(6), 1097-
601 1105.

602 Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., & Lipman,
603 D.J. (1997). Gapped BLAST and PSI-BLAST: a new generation of protein databases search
604 programs. *Nucleic Acids Research*, *25*, 3389-3402.

605 Beckett, S. T. (2000). *The Science of Chocolate*. Royal Society of Chemistry Paperbacks.

606 Belitz, H. D., Grosch, W., & Schieberle, P. (2009). Coffee, tea, cocoa. *Food chemistry*, 938-
607 970.

608 Besson, I., Creuly, C., Gros, J. B., & Larroche, C. (1997). Pyrazine production by *Bacillus*
609 *subtilis* in solid-state fermentation on soybeans. *Applied microbiology and biotechnology*,
610 *47*(5), 489-495.

611 Bonnarne, P., Amarita, F., Chambellon, E., Semon, E., Spinnler, H. E., & Yvon, M. (2004).
612 Methylthioacetaldehyde, a possible intermediate metabolite for the production of volatile
613 sulphur compounds from L-methionine by *Lactococcus lactis*. *FEMS microbiology letters*,
614 *236*(1), 85-90.

615 Camu, N., Gonzalez, A., De Winter, T., Van Schoor, A., De Bruyne, K., Vandamme, P.
616 (2008). Influence of turning and environmental contamination on the dynamics of populations
617 of lactic acid and acetic acid bacteria involved in spontaneous cocoa bean heap fermentation
618 in Ghana. *Applied and Environmental Microbiology*, *74*, 86-98.

619 Cocolin, L., Bisson, L. F., & Mills, D. A. (2000). Direct profiling of the yeast dynamics in
620 wine fermentations. *FEMS Microbiology Letters*, *189*, 81-87.

621 Crafac, M., Keul, H., Eskildsen, C. E., Petersen, M. A., Saerens, S., Blennow, A., & Nielsen,
622 D. S. (2014). Impact of starter cultures and fermentation techniques on the volatile aroma and
623 sensory profile of chocolate. *Food research international*, 63, 306-316.

624 Crafac, M., Mikkelsen, M. B., Saerens, S., Knudsen, M., Blennow, A., Lowor, S., ... &
625 Nielsen, D. S. (2013). Influencing cocoa flavour using *Pichia kluyveri* and *Kluyveromyces*
626 *marxianus* in a defined mixed starter culture for cocoa fermentation. *International journal of*
627 *food microbiology*, 167(1), 103-116.

628 de Melo Pereira, G. V., Magalhães, K. T., de Almeida, E. G., da Silva Coelho, I., & Schwan,
629 R. F. (2013). Spontaneous cocoa bean fermentation carried out in a novel-design stainless
630 steel tank: Influence on the dynamics of microbial populations and physical–chemical
631 properties. *International journal of food microbiology*, 161(2), 121-133.

632 Ducki, S., Miralles-Garcia, J., Zumbé, A., Tornero, A., & Storey, D. M. (2008). Evaluation of
633 solid-phase micro-extraction coupled to gas chromatography–mass spectrometry for the
634 headspace analysis of volatile compounds in cocoa products. *Talanta*, 74(5), 1166-1174.

635 Guehi, S. T., Dabonne, S., Ban-Koffi, L., Kedjebo, D. K., & Zahouli, G. I. B. (2010). Effect
636 of turning beans and fermentation method on the acidity and physical quality of raw cocoa
637 beans. *Advance Journal of Food Science and Technology*, 2(3), 163-171.

638 Hamdouche, Y., Guehi, T., Durand, N., Kedjebo, K. B. D., Montet, D., & Meile, J. C. (2015).
639 Dynamics of microbial ecology during cocoa fermentation and drying: Towards the
640 identification of molecular markers. *Food Control*, 48, 117-122.

641 Hamdouche, Y., Nganou D. N., Meile, J.C., Durand, N., Teyssier, C., Montet, D. (2016).
642 Discrimination of post-harvest coffee processing methods by microbial ecology analyses.
643 *Food Control*, 65, 112-120.

644 Hamdouche, Y., Teyssier, C., & Didier, M. (2017). Impact of Acetic Bacteria on Cocoa
645 Fermentation. *In Acetic Acid Bacteria: Fundamentals and Food Applications*. Editor: Ilkin
646 Yucel Sengun. CRC Press.

647 Heyndrickx, M., Vauterin, L., Vandamme, P., Kersters, K., & De Vos, P. (1996).
648 Applicability of combined amplified ribosomal DNA restriction analysis (ARDRA) patterns
649 in bacterial phylogeny and taxonomy. *Journal of Microbiological Methods*, 26, 247-259.

650 Ho, V. T., Zhao, J., & Fleet, G. (2014). Yeasts are essential for cocoa bean fermentation.
651 *International journal of food microbiology*, 174, 72-87.

652 Holm, C. S., Aston, J. W., & Douglas, K. (1993). The effects of the organic acids in cocoa on
653 the flavour of chocolate. *Journal of the Science of Food and Agriculture*, 61(1), 65-71.

654 ICCO 2017 : International Cocoa Organization.

655 Jinap, S., Thien, J., & Yap, T. N. (1994). Effect of drying on acidity and volatile fatty acids
656 content of cocoa beans. *Journal of the Science of Food and Agriculture*, 65(1), 67-75.

657 Kedjebo, K. B. D., Guehi, T. S., Kouakou, B., Durand, N., Aguilar, P., Fontana, A., &
658 Montet, D. (2016). Effect of post-harvest treatments on the occurrence of ochratoxin A in raw
659 cocoa beans. *Food Additives & Contaminants: Part A*, 33(1), 157-166.

660 Koné, M. K., Guéhi, S. T., Durand, N., Ban-Koffi, L., Berthiot, L., Tachon, A. F., & Montet,
661 D. (2016). Contribution of predominant yeasts to the occurrence of aroma compounds during
662 cocoa bean fermentation. *Food Research International*, 89, 910-917.

663 Kongor, J. E., Hinneh, M., Van de Walle, D., Afoakwa, E. O., Boeckx, P., & Dewettinck, K.
664 (2016). Factors influencing quality variation in cocoa (*Theobroma cacao*) bean flavour
665 profile—A review. *Food Research International*, 82, 44-52.

666 Kowalchuk, G.A., Stephen, J.R., de Boer, W., Prosser, J.I., Embley, T.M., & Woldendorp,
667 J.W. (1997). Analysis of ammonia-oxidizing bacteria of the beta subdivision of the class
668 Proteobacteria in coastal sand dunes by denaturing gradient gel electrophoresis and
669 sequencing of PCR amplified 16S ribosomal DNA fragments. *Applied and Environmental*
670 *Microbiology*, 63, 1489-1497.

671 Leensing, R. (2005). Identification and validation of specific markers for traceability of
672 aquaculture fish for import/export (PhD thesis). France: University of Montpellier 2.

673 Lefeber, T., Gobert, W., Vrancken, G., Camu, N., & De Vuyst, L. (2011). Dynamics and
674 species diversity of communities of lactic acid bacteria and acetic acid bacteria during
675 spontaneous cocoa bean fermentation in vessels. *Food Microbiology*, 28(3), 457-464.

676 Lefeber, T., Janssens, M., Camu, N., & De Vuyst, L. (2010). Kinetic analysis of strains of
677 lactic acid bacteria and acetic acid bacteria in cocoa pulp simulation media toward
678 development of a starter culture for cocoa bean fermentation. *Applied and environmental*
679 *microbiology*, 76(23), 7708-7716.

680 Lefeber, T., Papalexandratou, Z., Gobert, W., Camu, N., & De Vuyst, L. (2012). On-farm
681 implementation of a starter culture for improved cocoa bean fermentation and its influence on
682 the flavour of chocolates produced thereof. *Food microbiology*, 30(2), 379-392.

683 Lefeber, T., Papalexandratou, Z., Gobert, W., Camu, N., & De Vuyst, L. (2012). On-farm
684 implementation of a starter culture for improved cocoa bean fermentation and its influence on
685 the flavour of chocolates produced thereof. *Food microbiology*, 30(2), 379-392.

686 Lima, L. J., Kamphuis, H. J., Nout, M. R., & Zwietering, M. H. (2011). Microbiota of cocoa
687 powder with particular reference to aerobic thermoresistant spore-formers. *Food*
688 *microbiology*, 28(3), 573-582.

689 Menezes, A. G. T., Batista, N. N., Ramos, C. L., Silva, A. R. D. A., Efraim, P., Pinheiro, A.
690 C. M., & Schwan, R. F. (2016). Investigation of chocolate produced from four different
691 Brazilian varieties of cocoa (*Theobroma cacao* L.) inoculated with *Saccharomyces cerevisiae*.
692 *Food Research International*, *81*, 83-90.

693 Meyer, B., Biehl, B., Said, M. B., & Samarakoddy, R. J. (1989). Post-harvest pod storage: A
694 method for pulp preconditioning to impair strong nib acidification during cocoa fermentation
695 in Malaysia. *Journal of the Science of Food and Agriculture*, *48*(3), 285-304.

696 Moens, F., Lefeber, T., & De Vuyst, L. (2014). Oxidation of metabolites highlights the
697 microbial interactions and role of *Acetobacter pasteurianus* during cocoa bean fermentation.
698 *Applied and environmental microbiology*, *80*(6), 1848-1857.

699 Muyzer, G., De Waal, E. C., & Uitterlinden, A. G. (1993). Profiling of complex microbial
700 populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-
701 amplified genes coding for 16S rRNA. *Applied Environmental Microbiology*, *59*, 695-700.

702 Nazaruddin, R., Seng, L. K., Hassan, O., & Said, M. (2006). Effect of pulp preconditioning on
703 the content of polyphenols in cocoa beans (*Theobroma cacao*) during fermentation. *Industrial*
704 *Crops and Products*, *24*(1), 87-94.

705 Palys, T., Nakamura, L. K., & Cocha, F. M. (1997). Discovery and classification of ecological
706 diversity in the bacterial world: the role of DNA sequence data. *International Journal of*
707 *Systematic Bacteriology*, *47*, 1145-1156.

708 Papalexandratou, Z., Camu, N., Falony, G., & De Vuyst, L. (2011). Comparison of the
709 bacterial species diversity of spontaneous cocoa bean fermentations carried out at selected
710 farms in Ivory Coast and Brazil. *Food microbiology*, *28*(5), 964-973.

711 Papalexandratou, Z., Lefeber, T., Bahrim, B., Lee, O. S., Daniel, H. M., & De Vuyst, L.
712 (2013). *Hanseniaspora opuntiae*, *Saccharomyces cerevisiae*, *Lactobacillus fermentum*, and
713 *Acetobacter pasteurianus* predominate during well-performed Malaysian cocoa bean box
714 fermentations, underlining the importance of these microbial species for a successful cocoa
715 bean fermentation process. *Food microbiology*, 35(2), 73-85.

716 Papalexandratou, Z., Vrancken, G., De Bruyne, K., Vandamme, P., & De Vuyst, L. (2011).
717 Spontaneous organic cocoa bean box fermentations in Brazil are characterized by a restricted
718 species diversity of lactic acid bacteria and acetic acid bacteria. *Food microbiology*, 28(7),
719 1326-1338.

720 Pretorius, I. S. (2000). Tailoring wine yeast for the new millennium: novel approaches to the
721 ancient art of winemaking. *Yeast*, 16(8), 675-729.

722 Ramli, N., Hassan, O., Said, M., Samsudin, W., & Idris, N. A. (2006). Influence of roasting
723 conditions on volatile flavor of roasted Malaysian cocoa beans. *Journal of Food Processing*
724 *and Preservation*, 30(3), 280-298.

725 Rodriguez-Campos, J., Escalona-Buendía, H. B., Contreras-Ramos, S. M., Orozco-Avila, I.,
726 Jaramillo-Flores, E., & Lugo-Cervantes, E. (2012). Effect of fermentation time and drying
727 temperature on volatile compounds in cocoa. *Food Chemistry*, 132(1), 277-288.

728 Rodriguez-Campos, J., Escalona-Buendía, H. B., Orozco-Avila, I., Lugo-Cervantes, E., &
729 Jaramillo-Flores, M. E. (2011). Dynamics of volatile and non-volatile compounds in cocoa
730 (*Theobroma cacao* L.) during fermentation and drying processes using principal components
731 analysis. *Food Research International*, 44(1), 250-258.

732 Rojas, V., Gil, J. V., Manzanares, P., Gavara, R., Piñaga, F., & Flors, A. (2002). Measurement
733 of alcohol acetyltransferase and ester hydrolase activities in yeast extracts. *Enzyme and*
734 *Microbial Technology*, 30(2), 224-230.

735 Rojas, V., Gil, J. V., Piñaga, F., & Manzanares, P. (2003). Acetate ester formation in wine by
736 mixed cultures in laboratory fermentations. *International Journal of Food Microbiology*,
737 *86(1)*, 181-188.

738 Saltini, R., Akkerman, R., & Frosch, S. (2013). Optimizing chocolate production through
739 traceability: A review of the influence of farming practices on cocoa bean quality. *Food*
740 *control*, *29(1)*, 167-187.

741 Schwan, R. F., & Wheals, A. E. (2004). The microbiology of cocoa fermentation and its role
742 in chocolate quality. *Critical reviews in food science and nutrition*, *44(4)*, 205-221.

743 Selamat, J. I. N. A. P., & Mordingah Harun, S. (1994). Formation of methyl pyrazine during
744 cocoa bean fermentation. *Pertanika*, *17*, 27-27.


745 Serra-Bonvehí, J. (2005). Investigation of aromatic compounds in roasted cocoa powder.
746 *European Food Research and Technology*, *221(1-2)*, 19-29.

747 Sheffield, V. C., Beck, J. S., Stone, E. M., & Myers, R. M. (1989). Attachment of a 40 bp
748 G+C rich sequence (GC-clamp) to genomic DNA fragments by polymerase chain reaction
749 results in improved detection of single-base changes. *Proceeding of the National Academy of*
750 *Sciences of the United States of American*, *86*, 232-236.


751 Smit, G., Smit, B. A., & Engels, W. J. (2005). Flavour formation by lactic acid bacteria and
752 biochemical flavour profiling of cheese products. *FEMS microbiology reviews*, *29(3)*, 591-
753 610.

754 Smit, G., Smit, B. A., & Engels, W. J. (2005). Flavour formation by lactic acid bacteria and
755 biochemical flavour profiling of cheese products. *FEMS Microbiology Reviews*, *29(3)*, 591-
756 610.

- 757 Ter Braak, C. J. (1986). Canonical correspondence analysis: a new eigenvector technique for
758 multivariate direct gradient analysis. *Ecology*, 67(5), 1167-1179.
- 759 Tomlins, K. I., Baker, D. M., Daplyn, P., & Adomako, D. (1993). Effect of fermentation and
760 drying practices on the chemical and physical profiles of Ghana cocoa. *Food Chemistry*,
761 46(3), 257-263.
- 762 Visintin, S., Alessandria, V., Valente, A., Dolci, P., & Cocolin, L. (2016). Molecular
763 identification and physiological characterization of yeasts, lactic acid bacteria and acetic acid
764 bacteria isolated from heap and box cocoa bean fermentations in West Africa. *International
765 journal of food microbiology*, 216, 69-78.
- 766 Yong, F. M., Lee, K. H., & Wong, H. A. (1981). The production of ethyl acetate by soy yeast
767 *Saccharomyces rouxii* NRRL Y-1096. *International Journal of Food Science & Technology*,
768 16(2), 177-184.



(a)



(b)

Figure 1: Bacterial DNA DGGE profiles obtained from fermented cocoa beans (with and without turning) during 2, 4, 5, 6 and 7 days fermentation, previously stored during 2 days (a) and 8 days (b). Identification of DGGE bands for bacterial species are given in Table 1.




Figure 2: Principal component analysis (PCA) of bacterial 16S rDNA DGGE band profiles obtained from cocoa beans stored during 2 days, fermented with turning (2S2F T, 2S2F T, 2S4F T, 2S4F T, 2S5F T, 2S5F T, 2S6F T, 2S6F T, 2S7F T and 2S7F T) and without turning (2S2F, 2S2F, 2S4F, 2S4F, 2S5F, 2S5F, 2S6F, 2S6F, 2S7F and 2S7F) during 2, 4, 5, 6 and 7 days fermentation and from cocoa beans stored during 8 days, fermented with turning (8S2F T, 8S2F T, 8S4F T, 8S4F T, 8S5F T, 8S5F T, 8S6F T, 8S6F T, 8S7F T and 8S7F T) and without turning (8S2F, 8S2F, 8S4F, 8S4F, 8S5F, 8S5F, 8S6F, 8S6F, 8S7F and 8S7F) during 2, 4, 5, 6 and 7 days fermentation.




Figure 3: Yeasts DNA DGGE profiles obtained from fermented cocoa beans (with and without turning) during 2, 4, 5, 6 and 7 days fermentation, previously stored during 2 days (a) and 8 days (b) . Identification of DGGE bands for yeast species are given in Table 2.







Figure 4: Canonical analysis of correspondence (CAC) of bacterial and yeast DGGE band profiles obtained from cocoa beans stored during 2 days, fermented with turning (2S2F T, 2S2F T, 2S4F T, 2S4F T, 2S5F T, 2S5F T, 2S6F T, 2S6F T, 2S7F T and 2S7F T) and without turning (2S2F, 2S2F, 2S4F, 2S4F, 2S5F, 2S5F, 2S6F, 2S6F, 2S7F and 2S7F) during 2, 4, 5, 6 and 7 days fermentation and from cocoa beans stored during 8 days, fermented with turning (8S2F T, 8S2F T, 8S4F T, 8S4F T, 8S5F T, 8S5F T, 8S6F T, 8S6F T, 8S7F T and 8S7F T) and without turning (8S2F, 8S2F, 8S4F, 8S4F, 8S5F, 8S5F, 8S6F, 8S6F, 8S7F and 8S7F) during 2, 4, 5, 6 and 7 days fermentation.




a




b




c




g




h




i




m




n




o




d




j




p




e




k




q



f



l




r

■ 2F (Storage 2 days) ■ 2F+T (Storage 2 days)
 ■ 2F (Storage 8 days) ■ 2F+T (Storage 8 days)


■ 4F (Storage 2 days) ■ 4F+T (Storage 2 days)
 ■ 4F (Storage 8 days) ■ 4F+T (Storage 8 days)

■ 5F (Storage 2 days) ■ 5F+T (Storage 2 days)
 ■ 5F (Storage 8 days) ■ 5F+T (Storage 8 days)


Figure 5: Concentration of aldehydes (a,g,m), ketones (b,h,n), alcohols (c,i,o), esters (d,j,p), acids (e,k,q) and other volatile compounds (f,l,r) produced in cocoa beans previously stored during 2 and 8 days, fermented 2 days (2F), 4 days (4F) , 5 days (5F) with turning (+T) and without turning.▲ indicates significant differences between fermentation with and without turning.● indicates significant differences between the two pods storage duration. ◆ indicates significant differences between fermentation times.




a




g




b




h




c




i




d




j




e



k



f



l

■ 6F (Storage 2 days) ■ 6F+T (Storage 2 days)
 ■ 6F (Storage 8 days) ■ 6F+T (Storage 8 days)

■ 7F (Storage 2 days) ■ 7F+T (Storage 2 days)
 ■ 7F (Storage 8 days) ■ 7F+T (Storage 8 days)

Figure 6: Concentration of aldehydes (a,g), ketones (b,h), alcohols (c,i), esters (d,j), acids (e,k) and other volatile compounds (f,l) produced in cocoa beans previously stored during 2 and 8 days, fermented 6 days (6F) and 7 days (7F) with turning (+T) and without turning. ▲ indicates significant differences between fermentation with and without turning. ● indicates significant differences between the two pods storage duration. ◆ indicates significant differences between fermentation times.








Figure 7: Total Concentration of volatile compounds produced in cocoa beans previously stored during 2 days (a) and 8 days (b), fermented without turning at 2, 4, 5, 6, 7 days (2F, 4F, 5F, 6F, 7F) and with turning (2F+T, 4F+T, 5F+T, 6F+T, 7F+T). ▲ indicates significant differences between fermentation with and without turning. ● indicates significant differences between the two pods storage duration. ◆ indicates significant differences between fermentation times.



(a)



(b)



(c)

Figure 8: Concentration of organic acids present in cocoa beans previously stored during 2 days (a) and 8 days (b), fermented with turning and without turning. In (c) the two storage durations (2 and 8 days) were compared. ▲ indicates significant differences between fermentation with and without turning. ● indicates significant differences between the two pods storage duration.




Figure 9 : Principal component analysis (PCA) of peak areas of volatiles compounds obtained from cocoa beans stored during 2 days, fermented with turning (2S2F T, 2S2F T, 2S4F T, 2S4F T, 2S5F T, 2S5F T, 2S6F T, 2S6F T, 2S7F T and 2S7F T) and without turning (2S2F , 2S2F , 2S4F , 2S4F , 2S5F , 2S5F , 2S6F , 2S6F , 2S7F and 2S7F) during 2, 4, 5, 6 and 7 days fermentation and from cocoa beans stored during 8 days, fermented with turning (8S2F T, 8S2F T, 8S4F T, 8S4F T, 8S5F T, 8S5F T, 8S6F T, 8S6F T, 8S7F T and 8S7F T) and without turning (8S2F , 8S2F , 8S4F , 8S4F , 8S5F , 8S5F , 8S6F , 8S6F , 8S7F and 8S7F) during 2, 4, 5, 6 and 7 days fermentation.




Figure 10: Canonical analysis of correspondence (CAC) applied on microbial DGGE band profiles and peak areas obtained from cocoa beans stored during 2 and 8 days and fermented with and without turning during 2, 4, 5, 6 and 7 days.

Table 1: Identification of bacterial DNA bands isolated from DGGE experiment

Band	Family/ Genus/species	Percent identity	Query cover	E-value	References
A	<i>Lactobacillus curvatus</i>	93%	99%	5e-74	FJ609221
Y	<i>Enterobacteriaceae</i>	100%	100%	2e-97	KM021355
B	<i>Lactobacillus fermentum</i>	99%	100%	1e-95	LC042465
C	<i>Enterococcus casseliflavus</i>	99%	100%	1e-78	KJ571214
E	<i>Bacillus megaterium</i>	94%	100%	2e-77	KJ004421
F	<i>Bacillus sp</i>	99%	100%	8e-97	HG794259
G	<i>Acetobacter sp</i>	98%	100%	1e-89	AB853266
H		98%	100%	6e-88	
J		99%	100%	2e-92	
T		90%	99%	1e-53	
X	<i>Acinetobacter sp</i>	92%	100%	2e-51	KJ814994
K	<i>Bacillus sp</i>	100%	100%	2e-97	KM983003
L	<i>Providencia sp</i>	90%	97%	5e-64	AB920789
M	<i>Bacillus sp</i>	90%	100%	5e-43	FJ235680
N	<i>Providencia sp</i>	97%	100%	2e-67	KM059194
O	<i>Enterobacter aerogenes</i>	99%	100%	1e-95	LN623623
P	<i>Bacillus sp</i>	91%	97%	1e-53	KC236480
Q	<i>Enterobacter sp</i>	99%	100%	8e-97	GU944492
S	<i>Klebsiella sp</i>	92%	99%	7e-57	GQ416571
V	<i>Acetobacter pasteurianus</i>	100%	100%	1e-83	KM983001
W				5e-83	
R	<i>Klebsiella pneumoniae</i>	100%	100%	2e-97	KP761422
D, I	NI				

NI: not identified band

Table 2: Identification of DGGE bands for yeast species

Band	Family/ Genus/ species	Percent identity	Query cover	E-value	References
1	<i>Hanseniaspora opuntiae</i>	98%	100%	6e-93	KC111446
2	<i>Candida ethanolica/ Pichia deserticola</i>	93%	100%	8e-56	KM234475
3	<i>Pichia sp</i>	99%	100%	1e-79	JX408867
4	<i>Pichia galeiformis</i>	93%	100%	3e-76	HM212622
5	<i>Hanseniaspora uvarum</i>	93%	100%	4e-65	KM816746
9	<i>Pichia sp</i>	95%	100%	4e-65	EU884437
10	<i>Pichia kudriavzevii</i>	99%	100%	2e-47	KC494718
11	<i>Pichia manshurica</i>	99%	100%	7e-11	JQ419868
12	<i>Candida ethanolica/ Pichia deserticola</i>	93%	99%	2e-64	KM005182
6, 7, 13	NI				

NI: not identified band

Table 3: Volatiles compounds identified in cocoa beans fermented with and without turning

Group	RT (min)	RI	RI*	Compound
Aldehydes	1.50	811	821	Isobutanal
	2.15	910	912	2-Methylbutanal
	2.19	913	910	3-Methylbutanal
	28.06	1497	1495	Benzaldehyde
Ketones	2.94	971	984	2-Pentanone
	9.51	1153	1170	2-Heptanone
	21.06	1367	1388	2-Nonanone
	35.21	1629	1645	Acetophenone
	14.83	1252	1272	Acetoin
Esters	1.94	886	885	Ethyl acetate
	2.28	920	916	Methyl isovalerate
	3.60	1011	1005	Isobutyl acetate
	3.72	1016	1015	Methyl isopentanoate
	6.92	1105	1117	Isoamyl acetate
	44.55	1802	1803	Phenylethyl acetate
Alcohols	2.46	934	929	Ethanol
	7.42	1115	1118	2-Pentanol
	11.44	1189	1206	Isopentanol
	17.96	1310	1332	2-Heptanol
	33.03	1588	1582	2,3-Butanediol
	49.58	1895	1925	2-Phenylethanol
Acids	24.26	1426	1449	Acetic acid
	32.01	1570	1588	Isobutyric acid
	37.58	1680	1691	Isovaleric acid
Others	26.06	1460	1468	Tetramethylpyrazine
	46.73	1842	1848	2-Methoxy-phenol

*: Odour description and Retention Index (RI) obtained from literature: (Serra-Bonvehí, 2005; Rodriguez-Campos et al., 2011; 2012).

RI: Retention Index calculated

RT: Retention time

