



HAL
open science

Spike discharge characteristic of the caudal mesencephalic reticular formation and pedunculo pontine nucleus in MPTP-induced primate model of Parkinson disease

Laurent Goetz, Brigitte Piallat, Manik Bhattacharjee, Hervé Mathieu, Olivier David, Stéphan Chabardès

► To cite this version:

Laurent Goetz, Brigitte Piallat, Manik Bhattacharjee, Hervé Mathieu, Olivier David, et al.. Spike discharge characteristic of the caudal mesencephalic reticular formation and pedunculo pontine nucleus in MPTP-induced primate model of Parkinson disease. *Neurobiology of Disease*, 2019, 128, pp.40 - 48. 10.1016/j.nbd.2018.08.002 . hal-03486198

HAL Id: hal-03486198

<https://hal.science/hal-03486198>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Spike discharge characteristic of the caudal mesencephalic reticular formation and**
2 **pedunculopontine nucleus in MPTP-induced primate model of Parkinson disease**

3
4 Laurent Goetz^{1,2}, Brigitte Piallat^{1,2}, Manik Bhattacharjee^{1,2}, Hervé Mathieu^{1,2,3,4},
5 Olivier David^{1,2}, Stéphan Chabardès^{1,2,5,6}

6
7
8 ¹ *Grenoble Institute of Neurosciences-INSERM U1216 CEA-UJF-CHUGA, Grenoble, France*

9 ² *University Grenoble Alpes, Grenoble, France*

10 ³ *Unité Mixte de Service IRMaGe, Grenoble Alpes Hospital, F-38000 Grenoble, France*

11 ⁴ *Unité Mixte de Service 3552, CNRS, F-38000 Grenoble, France*

12 ⁵ *Department of Neurosurgery-Centre Hospitalier Universitaire Grenoble Alpes, Grenoble, France*

13 ⁶ *CEA Clnatec-Minatec, Grenoble, France*

14
15 **Abstract**

16
17 The pedunculopontine nucleus (PPN) included in the caudal mesencephalic reticular
18 formation (cMRF) plays a key role in the control of locomotion and wake state. Regarding its
19 involvement in the neurodegenerative process observed in Parkinson disease (PD), deep brain
20 stimulation of the PPN was proposed to treat levodopa-resistant gait disorders. However, the
21 precise role of the cMRF in the pathophysiology of PD, particularly in freezing of gait and
22 other non-motor symptoms is still not clear.

23 Here, using micro electrode recording (MER) in 2 primates, we show that dopamine depletion
24 did not alter the mean firing rate of the overall cMRF neurons, particularly the putative non-
25 cholinergic ones, but only a decreased activity of the regular neurons sub-group (though to be
26 the cholinergic PPN neurons). Interestingly, a significant increase in the relative proportion
27 of cMRF neurons with a burst pattern discharge was observed after MPTP intoxication. The
28 present results question the hypothesis of an over-inhibition of the CMRF by the basal ganglia
29 output structures in PD. The decreased activity observed in the regular neurons could explain
30 some non-motor symptoms in PD regarding the strong involvement of the cholinergic neurons
31 on the modulation of the thalamo-cortical system. The increased burst activity under
32 dopamine depletion confirms that this specific spike discharge pattern activity also observed
33 in other basal ganglia nuclei and in different pathologies could play a major role in the
34 pathophysiology of the disease and could explain several symptoms of PD including the

35 freezing of gait. The present data will have to be replicated in a larger number of animals and
36 will have to investigate more in details how the modification of the spike discharge of the
37 cMRF neurons in the parkinsonian state could alter functions such as locomotion and
38 attentional state. This will ultimately allow a better comprehension of the pathophysiology of
39 freezing of gait.

40

41 **Keywords:**

42

43 Pedunculopontine nucleus

44 Parkinson disease

45 Mirco electrode recording

46 Mesencephalic reticular formation

47 Non-human primate

48 MPTP

49

50 **Abbreviations:**

51

52 cMRF: caudal mesencephalic reticular formation ;

53 DBS: deep brain stimulation ;

54 MER: micro electrode recording ;

55 PD: Parkinson disease ;

56 PPN: pedunculopontine nucleus ;

57 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine: MPTP

58

59

60 **Funding:**

61

62 This work was supported by grants from Medtronic and by France Parkinson Association.

63 Sponsors had no role in the study design, data collection, data analysis, or data interpretation.

64 The neurophysiology facility and the Grenoble MRI facility IRMaGe were partly funded by

65 the French program Investissement d’Avenir run by the Agence Nationale pour la Recherche

66 (Grant Infrastructure d’avenir en Biologie Santé, ANR-11-INBS-0006

67

68

69

70 **Introduction**

71 Improving therapeutic strategies to treat gait disorders in neurodegenerative diseases requires
72 a better understanding of the pathophysiology at the level of brainstem structures. The caudal
73 mesencephalic reticular formation (cMRF) contains the pedunculopontine nucleus (PPN)
74 (*nucleus tegmentalis pedunculopontinus*) and the cuneiform nucleus thought to be involved in
75 several functions such as the supra-spinal control of locomotion, postural tone, waking state
76 and sleep (Gut and Winn, 2016; Mena-Segovia and Bolam, 2017; Garcia-Rill, 2015).

77 In addition to the neurodegenerative process affecting dopaminergic neurons of the *substantia*
78 *nigra pars compacta* occurring in Parkinson disease (PD), the loss of cholinergic PPN
79 neurons is thought to be involved in the PD pathophysiology (Braak et al., 2003; Coelho and
80 Ferreira, 2012; Galvan and Wichmann, 2008; Hirsch et al., 1987; Jellinger, 1988; Karachi et
81 al., 2010; Pienaar et al., 2013; Rinne et al., 2008; Zweig et al., 1989). Based on experimental
82 data in Parkinsonian animal models, it was suggested that over-activity of the basal ganglia
83 (BG) output structures (internal segment of the *globus pallidus* and *substantia nigra pars*
84 *reticulata*) could lead to hypoactivity of the PPN and finally could explain akinetic symptoms
85 and rigidity (Aziz and Stein, 2008; Gomez-Gallego et al., 2006; Hamani et al., 2007; Pahapill
86 and Lozano, 2000; Takakusaki et al., 2008). On the contrary, an increased
87 electrophysiological activity of PPN neurons was observed in the 6-hydroxydopamine PD
88 model in rat (Breit et al., 2001), in agreement with a previous metabolic study in rat (Orioux
89 et al., 2000).

90 Altogether, these results highlighted a potential role of the PPN neurons in the
91 pathophysiology of the PD. In 2005, deep brain stimulation (DBS) of the PPN was proposed
92 as a new therapeutic strategy to treat levodopa-resistant gait disorders in PD such as freezing
93 of gait (Mazzone et al., 2005; Plaha and Gill, 2005). Despite a growing interest in the
94 neurological community, the precise role of the PPN in the pathophysiology of PD and related
95 gait troubles is still lacking (Mena-Segovia and Bolam, 2017; Pienaar et al., 2017). Regarding
96 the overall considerations on the difficulty to study PPN activity in human beings, we
97 initiated experiments in the behaving primate to study PPN activity during locomotion (Goetz
98 et al., 2016a) and to investigate the role of PPN in the pathophysiology of PD. Regarding the
99 lack of precise anatomical delimitation of the PPN in primate and human, we decided to refer
100 to the cMRF to describe the PPN area, assuming that it encompasses the PPN (cholinergic
101 and non-cholinergic neurons) and the cuneiform nucleus (Goetz et al., 2016a).

102 For several decades, intoxication with the neurotoxin 1-methyl-4-phenyl-1,2,3,6-
103 tetrahydropyridine (MPTP) has proved to be a very robust non-human primate model of PD
104 (Fox and Brotchie, 2010; Morissette and Paolo, 2017; Przedborski et al., 2001; Wichmann et
105 al., 2017). This is the model we considered here, with the goal to evaluate the spike discharge
106 characteristics of the cMRF neurons in the PD context. Using micro electrode
107 recording (MER), we investigated how dopaminergic depletion induced by MPTP
108 intoxication could modify the activity of cMRF neurons by evaluating changes in firing rate
109 and pattern. In particular, we tested the hypothesis of a hypoactivity of the MRF neurons in
110 PD because of basal ganglia dysfunction.

111

112 **Materials and methods**

113 The methods used in the present study were described in detail in our previous study (Goetz
114 et al., 2016a) and followed methods commonly used in primate MER studies of the basal
115 ganglia or PPN area in our lab and by other groups using ventricular landmarks to calculate
116 electrode trajectories (Devergnas et al., 2012; Matsumura et al., 1997; Wichmann et al.,
117 1994).

118

119 *Animals*

120 The same two cynomolgus monkeys (*Macaca fascicularis* - CRP Port Louis, Mauritius) that
121 were used in the previous report (Goetz et al., 2016a) were also used in this present study
122 (Primate K: male, 9 years old, weight: 9 kg ; Primate T : male, 3 years old, weight: 5.5 kg).
123 All experiments were carried out in accordance with the recommendations of the European
124 Community Council Directives of 1986 (86/609/EEC), the National Institutes of Health
125 Guide for the Care and Use of Laboratory Animals. The experimental protocol was approved
126 by the Ethics Committee of Région Rhône-Alpes. Animals were kept with other congeners
127 allowing social behavior, in an air-conditioned room under standard conditions of temperature
128 ($23 \pm 1^\circ\text{C}$), humidity ($65 \pm 4\%$) and light (12 h light/dark cycle). They had access *ad libitum*
129 to food and water and were given fresh fruits and vegetables every day. No food or water
130 restriction was used in the present experiment.

131

132 *Surgical procedures and assessment of MER trajectories*

133 Briefly, under ventriculographical X-Ray control, MRI-compatible titanium footed head
134 holder and recording chamber (Crist Instrument Company, Hagerstown, USA) were

135 stereotactically fixed to the skull with titanium screws. The precise assessment of micro
136 electrode trajectories and recording site were performed with methods commonly used in
137 primate electrophysiological studies combined with MRI (pre-, post-operative and *post*
138 *mortem*) and histological data obtained for each primate (electrolytic lesion and
139 immunohistological staining processed for either a Nissl stain or immunocytochemistry for
140 acetylcholine transferase to localize the cholinergic cMRF neurons).

141

142 *MER procedures and data acquisition*

143 MER were performed using a microdrive set-up (FlexMT-tm and Multi Drive-Tm, Alpha
144 Omega Engineering, Nazareth, Israel). Bevelled mini-tubes were lowered through the dura
145 matter to avoid any damage of the micro electrode when piercing the dura-mater. Two to
146 three tungsten micro electrodes (Impedance: 2-3 MOhm) (FHC Inc, Bowdoin, USA) were
147 lowered independently. The extracellular signal was amplified, analog band-pass filtered
148 (300-6000 Hz) and sampled at 50 kHz. The first cortical neuronal activity defined the null
149 depth of the trajectory. In both primates, the same trajectories were performed in normal and
150 after MPTP intoxication.

151

152 *MPTP treatment*

153 Primates were intoxicated by an intramuscular (IM) injection of MPTP (0.6 mg.kg^{-1} in
154 primate K and 0.5 mg.kg^{-1} in primate T). Primates developed severe Parkinsonian symptoms
155 that necessitated a complete nursing. In primate T, supplementary injection was necessary to
156 induced severe symptoms (0.4 mg.kg^{-1}). L-Dopa therapy (Modopar 50 mg) was provided in
157 the two primates to alleviate the symptoms. Evaluation of the Parkinsonian symptoms was
158 performed every two days, off medication using a modified version of the Imbert scale
159 (Imbert et al., 2000) (See Supplementary Material). Four weeks after the intoxication,
160 primates recovered and symptoms were stable enough to start the electrophysiological
161 protocol. To evaluate dopaminergic degeneration, tyrosine hydroxylase immunostaining was
162 done using a standard immunoperoxidase method, on adjacent coronal serial sections of $50 \mu\text{m}$
163 as previously described (Song and Haber, 2000) (See Supplementary Material).
164 Electrophysiological recordings were always performed off medication.

165

166 *Signal processing*

167 Spike sorting was performed using a template-matching algorithm (Spike 2, CED Software,
168 Cambridge, United Kingdom). Semi-automatic spike isolation was systematically verified and

169 corrected on the basis of the quality of the spike detection (waveforms overdraw) and by a
170 principal component analysis to discriminate single-unit from other multi-unit activity.
171 Calculation of the neuronal refractory period on interspike interval (ISI) was systematically
172 performed and used as an exclusion criterion when it was found to be less than 2 ms (Fee et
173 al., 1996). Firing rate and firing pattern analysis were performed using homemade software
174 developed under Matlab R2009b (The MathWorks Inc., Natick, Massachusetts, USA). A
175 minimum of 500 events in the resulting spike train representing the activity of a single-unit
176 was required to include a neuron in the study.

177 For each neuron, we used the distribution of the inverse of interspike intervals (ISIs) to
178 calculate the mean, standard deviation (SD) and median of the FR.

179

180 *Firing pattern analysis*

181 We determined the firing pattern characteristics by a combined approach using data based on
182 the interval ISI and autocorrelogram analysis further completed with burst detection methods
183 (Abeles, 1982; Wichmann et al., 1994; Baron et al., 2011; Piallat et al., 2011).
184 Autocorrelation histograms of spike trains were calculated for 1s (1 ms bin width) using
185 MATLAB software. Peaks were considered significant if they were found to be above the
186 confidence line set at mean \pm 2 SDs. We classified single-unit activities as regular, irregular,
187 or burst as follows:

- 188 • Regular activity was characterized by a tonic discharge with a narrow peaked interval
189 interspike distribution and a significant peak on the autocorrelation function usually
190 equal to the mean firing period (i.e., 1/FR).
- 191 • Irregular activity was characterized by a wide interval interspike distribution and a flat
192 autocorrelogram.
- 193 • Burst activity showed a wide or bimodal interval interspike distribution and a
194 significant single peak on the autocorrelation function. In addition, two different burst
195 detection methods were used to complete the detection: the “Poisson surprise” method
196 of Legendy and Salcman (1985). Applied to our data, using the Poisson surprise
197 method, bursts were defined as segments including at least 3 spikes with an *S* value of
198 at least 3. Due to the fact that some neurons displayed some long periods of pause that
199 affected the mean FR, thus affecting the Poisson surprise *S* value, we complemented
200 the burst detection with the interval method that calculates the percentage of spikes in
201 burst derived from the study of Baron et al. (2011). For each neuron, we selected the

202 following final parameters: maximum initial interval signifying burst onset: the
203 median ISI ; longest ISI allowed within a burst: median ISI/2 ; Minimum number of
204 events in a burst: 3. This allowed us to use a burst detection method based on the
205 baseline median rather than the mean FR. Based on these two methods and the
206 complementary ISI and autocorrelogram analysis, the definition of the burst pattern
207 group was statistically significant in the two primates using the two methods (See
208 Figure 2 C-D).

209 For each burst neuron, auto-correlogram analysis was used to assess periodic oscillations
210 defined by the presence of two or more consecutive significant peaks in the autocorrelogram.
211 The main frequency of bursts was defined using the spectral analysis of the transformation of
212 the spike train from discrete series to a continuous function. A power spectrum was computed
213 from the new continuous function (resolution of 0.4 Hz) using a fast Fourier transform within
214 the 0.5–15Hz range (Hanning window, resolution 0.75 Hz). A significant oscillatory activity
215 was considered if the spectrum had one or more peaks above the confidence line
216 (mean + 2 SD).

217

218 *Statistics*

219 Statistical analyses were performed using SPSS 20 (IBM, Armonk, New York) and Prism
220 (GraphPad Software, Inc, La Jolla, USA). We decided to analyse the data obtained in the two
221 primates in normal state and after MPTP intoxication separately as we could not guarantee
222 that the level of neuronal degeneration was identical between the primates. Thus, we decided
223 to replicate the analysis in the two animals and to evaluate the similarities between the
224 findings. Identification of significant differences of mean firing rate between groups were
225 performed using non-parametric Mann-Whitney U-test (significance was assumed for
226 $P \leq 0.05$).

227 Firing rate Chi-2 analysis without correction was performed to test the independence between
228 spiking pattern variables (burst, irregular, regular) and primate status (normal versus MPTP)
229 for each primate. More specifically, to test the association between burst pattern and primate
230 status, we performed the Cochran-Mantel-Haenszel test with 2 stratum (Primates K and T).
231 The null hypothesis was rejected if the P-value of the Mantel-Haenszel test was less than
232 0.05.

233

234

Figure 1 about here

235

236 **Results**

237 We recorded 235 single-unit activities in the cMRF in normal state (91 neurons in Primate K
238 and 144 neurons in primate T). Using the same trajectories, we recorded 152 neuronal
239 activities (single-unit) under MPTP condition (67 neurons in Primate K and 85 neurons in
240 primate T). The electrophysiological mapping of the cMRF extended antero-posteriorly from
241 2 to 6 mm from the anterior border of PC, between 1 and 7 mm laterally from the midline and
242 rostro-caudally from the caudal level of the IC to the pontine nuclei below the superior
243 cerebellar peduncle.

244

245 *Firing rate*

246 Figures 1 A-B show the distributions of the baseline mean firing rate in the two primates in
247 normal and MPTP states. In both primates and in both states, the distributions are wide, with a
248 mean firing rate varying from very low frequencies up to 100 Hz.

249 When comparing the firing rate between the 2 states, Normal *versus* MPTP, in the two
250 primates, the overall mean firing rate were not significantly different between normal and
251 MPTP state (Figure 2A). In primate K, the mean firing rate between normal and MPTP state
252 changed from 16.0 Hz (SD: 15.7) to 13.8 Hz (SD: 10)(Mann-Whitney U; $p= 0.63$). In primate
253 T, the mean firing rate between normal and MPTP state changed from 20.5 (SD: 17.7) Hz to
254 21.8 (SD: 17.8)(Mann-Whitney U; $p= 0.43$).

255 *By comparing the mean firing rate of the pattern sub-groups between normal and MPTP*
256 *(Figure 2B), we did not find significant difference in the mean firing rate of the burst neurons*
257 *between the 2 states (15.7 Hz (SD: 10.1) and 12.3 Hz (SD: 8.6) in primate K (Mann-Whitney*
258 *U; $p= 0.11$) and (18.1 Hz (SD: 13.5) and 22.4 Hz (SD: 15.1) in primate T (Mann-Whitney U;*
259 *$p= 0.17$). Similarly, no significant difference was observed in the irregular pattern sub-group*
260 *in the 2 primates between the 2 states: 9.7 Hz (SD: 10.5) and 9.3 Hz (SD: 11.9) in primate K*
261 *(Mann-Whitney U; $p= 0.84$) and 11.37 Hz (SD: 11.4) and 10.6 Hz (SD: 7.9) in primate T*
262 *(Mann-Whitney U; $p= 0.87$). On contrary, when considering the regular pattern sub-group, a*
263 *significant difference was observed in Primate K (Mann-Whitney U; $p= 0.045$) with a*
264 *decrease of the mean firing rate from 33.5 Hz (SD: 22.7) in normal state to 21.4 Hz (SD:*
265 *16.0) after MPTP intoxication. In Primate T, no significant difference (Mann-Whitney U; $p=$*
266 *0.9) was observed in this pattern sub-group 32.6 Hz (SD: 19.4) in normal state and 33.8 Hz*
267 *(SD: 22.0) under MPTP condition.*

268

269 Figure 2 about here

270 Figure 3 about here

271

272 *Firing pattern*

273 Significant differences in firing pattern were observed in the two primates between normal
274 and MPTP states (χ^2 -test: $p < 0.05$). More precisely, in primate K, between normal and
275 MPTP states, the percentage of burst neurons increased from 34.1% to 47.8% while the
276 percentage of the irregular ones decrease from 48.4 % to 26.9% of the overall number of
277 neurons. Similarly, in primate T, the evolution of the percentages of burst and irregular
278 neurons were from 23.6% to 42.4%, and from 41.7% to 30.6%, respectively. In both primates,
279 the change of percentage of regular neurons was less than 10% (Figure 3A). In addition, by
280 comparing the relative proportions of the burst neurons compared with non-burst neurons
281 (irregular + regular) simultaneously in the two primates (treated in the Cochran-Mantel-
282 Haenszel test as two stratum), we also found significant difference in the percentage of burst
283 versus non-burst neurons between normal and MPTP (Cochran-Mantel-Haenszel χ^2 -test: p
284 = 0.004) (Figure 3B). Interestingly, the relative proportion of neurons with a burst oscillatory
285 activity did not change significantly between normal and PD states (Figure 4A). However,
286 under MPTP condition, those burst neurons displayed an oscillatory frequency around 20 Hz
287 in both primates, which was significantly higher than normal condition in primate K only
288 (Figures 4B).

289 Figure 4 about here

290 **Discussion**

291 We could perform the same mapping of the cMRF, using the same micro electrode
292 trajectories, between normal and Parkinsonian states in two non-human primates to
293 investigate how neuronal activities in the cMRF were affected by dopamine depletion.
294 Recording sessions under MPTP conditions provided two main results. i) No significant
295 difference in mean firing rate between normal and MPTP conditions of the overall cMRF
296 recorded neurons but a decrease of the mean firing rate observed in the regular neurons
297 pattern sub-group in 1 primate. ii) A significant increase in the relative proportion of cMRF
298 neurons with a busrt discharge pattern after MPTP intoxication.

299

300 *Firing rate in MPTP condition*

301 We did not observe a significant overall change in the mean firing rate of cMRF neurons in
302 the 2 primates, as was first hypothesized. Indeed, convincing datasets obtained in primate by

303 lesion or chemical manipulation suggested that hyperactive GABAergic afferences from
304 GPi/SNr could lead to hypoactivity of the PPN and consequently to the development of
305 akinetic symptoms (Gomez-Gallego et al., 2006; Kojima et al., 1997; Nandi et al., 2002b;
306 Palombo et al., 1990). This led to the well-known hypothesis supported by the “rate model”:
307 in the context of PD, hyperactivity of the basal ganglia output structures leads to
308 inactivation/hypoactivity of the PPN (Hamani et al., 2007; Matsumura, 2005; Mena-Segovia
309 et al., 2004; Pahapill and Lozano, 2000; Stein and Aziz, 2012; Takakusaki et al., 2008). Of
310 interest, DBS of the human PPN was proposed in 2005 as a new therapeutic strategy to treat
311 gait disorders in PD based on those results and hypothesis (Mazzone et al., 2005; Plaha and
312 Gill, 2005). However, some electrophysiological data obtained in a rodent model of PD
313 provided opposite results, in favor of an hyperactivity of the PPN following dopaminergic
314 depletion (Breit et al., 2001; Orieux et al., 2000). Similarly, this latter result was not observed
315 in the present data in primate treated with MPTP.

316 When considering the pattern sub-groups, our previous observations on the evolution of the
317 mean firing rate between the 2 conditions were confirmed. Indeed, in primate, projections
318 from the BG output structures (GPi and SNr) preferentially target non-cholinergic neurons in
319 the cMRF (M. Parent et al., 1999; Rolland et al., 2011; Shink et al., 1997).

320 In the present study, it was not possible to determine the neurochemical nature of the
321 recording neurons but assuming that neurons with a regular firing pattern could be the
322 putative cholinergic type II neurons (Takakusaki et al., 1996; Leonard and Llinás, 1994;
323 Takakusaki:1996jw; Kang and Kitai, 1990; Gut and Winn, 2016; Mena-Segovia and Bolam,
324 2017) we neither observed a modification of the non-regular neurons (putative non-
325 cholinergic) known to receive projections from the BG output structures projections.

326 However, we observed a significant decrease in firing rate of the regular neurons in one
327 primate while in the other primate, a similar decrease was observed but not significantly. One
328 possible explanation is that only the firing rate of the PPN cholinergic neurons were affected
329 by dopamine depletion. This is surprising because it was observed in primate that DAT-
330 positive fibers tended to avoid PPN cholinergic cell bodies (Rolland et al., 2009). However, a
331 dopaminergic modulation of the cholinergic neurons projections via passive transmission can
332 still be hypothesized as it was demonstrated in several species from vertebrates to mammals
333 including human, that TH+/DAT+ fibers and varicosities can be observed in the vicinity of
334 PPN cholinergic neurons (Ryczko et al., 2016).

335

336 *Firing pattern in MPTP condition*

337 On the other hand, in the two primates, we found a significant increase in the relative number
338 of neurons that discharged in bursts after MPTP intoxication. Furthermore, the proportion of
339 burst neurons with an oscillatory activity remained stable after dopamine depletion but a trend
340 to shift from α to β band frequency was evident. In normal primate, the only studies that
341 investigated the spike discharge of PPN neurons did not perform a detailed analysis of the
342 firing pattern (Kobayashi et al., 2002; Matsumura et al., 1997). But when considering the
343 raster plot provided in the aforementioned studies, it is reasonable to consider that some of the
344 recorded neurons discharged in burst mode. In human, MER during DBS electrode
345 implantation in the PPN area were performed by several groups (Piallat et al., 2009;
346 Shimamoto et al., 2010; Tattersall et al., 2014; Weinberger et al., 2008). They all described a
347 bursting activity in the PPN neurons but the paradigms associated with MER in PD patients
348 makes difficult the comparisons with the present data.

349 The precise mechanisms responsible for the increase in burst pattern and the role of dopamine
350 depletion in this phenomenon remain to be determined. Could an excessive hyperpolarisation
351 of the cMRF neurons by BG output structures lead to an increase bursting rebound is an
352 opened question (Beurrier et al., 1999; Galvan and Wichmann, 2008; Plenz and Kitai, 1999;
353 Wichmann and Soares, 2006). This could explain some effects observed in primate after
354 neurochemical modulation of PPN neurons by GABAergic agent (agonist and antagonist)
355 (Nandi et al., 2002a). This would be in accordance with the rebound spikes observed in PPN
356 neurons (type I & II) after SNr stimulation in rat (Kang and Kitai, 1990). Also, the close
357 interconnection between PPN and STN should be considered in this perspective as an increase
358 bursting activity in the STN has been observed in primate after MPTP intoxication (Bergman
359 et al., 1994).

360 *Considerations associated with spike discharge analysis*

361 As mentioned by Israel and colleagues (Israel and Bergman, 2008), caution should be
362 observed regarding the large variability of firing rate analysis in the BG studies that led to
363 consider that BG output structures are hyperactive in PD context. Indeed, two studies in
364 primate failed to demonstrate a significant increase in firing rate of the GPi and SNr after
365 MPTP intoxication (Raz et al., 2000; Wichmann et al., 1999). [Similarly, in our results, we
366 found a large variability of the mean firing rate of the cMRF neurons in both primates, thus
367 caution should also be observed in the present study when interpreting the evolution of the
368 firing rate after MPTP intoxication.](#) Moreover, as pointed out by several studies, PPN area is
369 composed by a heterogeneous neuronal population (Martinez-Gonzalez et al., 2012; Pienaar
370 et al., 2017; Rye et al., 1987; Wang and Morales, 2009). Thus, the evaluation of the effect of

371 GABAergic projections on the heterogeneous cMRF neuronal population in primate should
372 be considered with caution. Finally, given the constraints related to experimentation on
373 primate, it was not possible to define the neurochemical nature of the recorded neurons. Thus,
374 interpretation of the discharge characteristics was based on studies leaded on other animal
375 models and therefore requires caution.

376

377 *Functional implications*

378 The decrease activity of the regular neurons (putative PPN cholinergic type II cells) must be
379 confirmed on larger number of cells and animals. This could explain some non-motor troubles
380 including sleep disorders and attentional symptoms observed in the Parkinsonian state
381 (Schapira et al., 2017). Indeed, the PPN cholinergic neurons considered as the cholinergic arm
382 of the reticular ascending system (Garcia-Rill, 2015; Jones, 2005; Shute and Lewis, 1967;
383 Steriade, 1996) are known to project on intralaminar and reticular thalamic nuclei (Lavoie and
384 A. Parent, 1994a), especially on parafascicular thalamic neurons (inhibitory effect) and
385 thalamocortical neurons (excitatory effect) (Ye et al., 2010). Decreased thalamic input by the
386 PPN neurons in the wake state could induce changes in the firing pattern of thalamic neurons,
387 switching spontaneously from a tonic to a bursting discharge mode leading to low-frequency
388 rhythmicity of thalamic neurons (Llinás et al., 2005). Ultimately, a decrease in PPN input
389 observed after MPTP intoxication could cause dysrhythmic state in the thalamo-cortical
390 system, *i.e.* the thalamocortical dysrhythmia (Llinás and Steriade, 2006). In line with the
391 previous arguments, in a previous study in normal behaving primate, we found some neurons
392 in the cMRF mostly with a regular discharge pattern, which modulated their activity during
393 transition from wakefulness to sleep, suggesting an active role in brain state as previously
394 found in other species (Datta and Siwek, 2002; Mena-Segovia et al., 2008; Steriade and
395 McCarley, 2013). Regarding the present data, change in the spiking discharge after MPTP
396 intoxication could lead to disturbance in brain state transitions.

397 Cholinergic and non-cholinergic efferents from the PPN on dopaminergic neurons of the
398 *substantia nigra* has been described in several species including primate and may play a
399 pivotal modulation of the dopaminergic transmission on BG (Dautan et al., 2016). Regarding
400 the location of the synaptic contact between cholinergic axons and dopaminergic SNc neurons
401 (perisomatic and on proximal dendrites) (Lavoie and A. Parent, 1994b), the spike discharge
402 modifications of the cMRF observed in the present study, *i.e.* decrease firing rate of regular
403 neurons and increase bursting activity following MPTP intoxication, could lead to dramatic

404 modification of the dopaminergic innervation of the BG and thus could be a key feature of the
405 pathophysiology of PD.

406 Finally, regarding the potential effect of spike discharge disturbance of the cMRF neurons
407 after MPTP intoxication on both the thalamocortical system and the BG circuitry, the
408 involvement of the cMRF neurons in the pathophysiology of freezing of gait is likely. Indeed,
409 in a previous study, we described the involvement of the same cMRF neurons in the control
410 of locomotion via 2 types of neuronal response, *i.e.* a phasic and/or a tonic discharge mode
411 (Goetz et al., 2016a). We also found that some cMRF neurons were involved in both
412 locomotion and wake state transition (Goetz et al., 2016b), suggesting that the cMRF could be
413 considered as an integrative structure (Lau et al., 2015) for both motor and attentional
414 functions in order to adapt locomotion in the environment via its ascending and descending
415 projections.

416 Using our model of bipedal locomotion (Goetz et al., 2012), we could perform recordings in
417 the cMRF after MPTP intoxication while the bipedal locomotion primate was blocked
418 (impossibility for the primate to initiate the swing phase). Although this cannot be considered
419 as an episode of freezing of gait in primate, in few neurons, we observed abnormal neuronal
420 activities in the cMRF during blocking of the swing phase associated with an abnormal EMG
421 activity of the quadriceps muscle group (see Supplementary Data). These preliminary data
422 will have to be confirmed in further experiments in order to precisely characterize a possible
423 role of spike discharge abnormality of cMRF neurons in the occurrence of freezing of gait.

424

425 *Implication for DBS*

426 While the mechanisms of action of DBS are still not clearly understood at the level of the BG,
427 the investigation of such mechanisms in the reticular formation to treat gait disorders should
428 be regarded very cautiously. If the present data were confirmed on a larger neuronal
429 population and animals, then it is reasonable to consider the hypothesis that DBS could have
430 two effects of the neuronal activity in the cMRF: first stimulation at low frequency (from 10
431 to 30 Hz) could drive the remaining PPN cholinergic neurons to fire with a higher frequency,
432 then restoring a normal PPN-thalamic transmission resulting in the suppression or reduction
433 of thalamocortical dysrhythmia. Second, it has been suggested that the excessive burst activity
434 observed in several nuclei of the BG could play a critical role in the pathophysiology of
435 related symptoms in different pathologies (Galvan and Wichmann, 2008; Lobb, 2014; Piallat
436 et al., 2011; Welter et al., 2011). Thus, we can suggest that a possible mechanism of DBS in
437 the BG and in the cMRF is associated with the reduction of pathological burst activity by

438 acting on the membrane hyperpolarization and the restoration of a physiological discharge
439 activity.

440

441 **Conclusion**

442 Our results did not confirm the hypothesis of an over-inhibition of the PPN by the SNr/GPi
443 complex because we did not observe any significant decrease in the mean firing rate of the
444 overall cMRF neurons, nor on putative non-cholinergic neurons, after MPTP intoxication.
445 However, the decreased activity of the regular neurons (putative PPN cholinergic neurons)
446 could have dramatic consequences on the thalamocortical system and finally could explain
447 some of the non-motor symptoms in PD. In parallel, the increase in bursting activity observed
448 in the 2 primates provides new insights on the key role of the cMRF neurons in the
449 pathophysiology of PD and gait disorders. It is now of special interest in further studies to
450 consider the cMRF as an integrative structure associated with motor function and the reticular
451 activating system regarding its dense ascending and descending projections. This will allow
452 investigating complex symptoms such as freezing of gait or sleep/arousal disturbance.

453

454 **Acknowledgements**

455 We are grateful to Vincente Dicalogero for animal care, Yann Thibaudier for assistance
456 during experiments; Patrick Mouchet in Grenoble for insightful discussions and
457 Prof. Suzanne N. Haber and her team in the Department of Pharmacology and Physiology,
458 University of Rochester Medical Center for the valuable tissue processing and
459 immunohistochemistry.

460

461 **References**

- 462 Abeles, M., 1982. Quantification, smoothing, and confidence limits for single-units'
463 histograms. *J Neurosci Methods* 5, 317–325.
- 464 Aziz, T.Z., Stein, J.F., 2008. Editorial commentary: Oscillatory activity and deep brain
465 stimulation in the pedunculopontine nucleus 212, 247–250.
466 doi:10.1016/j.expneurol.2008.04.029
- 467 Baron, M.S., Chaniary, K.D., Rice, A.C., Shapiro, S.M., 2011. Multi-neuronal recordings in
468 the Basal Ganglia in normal and dystonic rats. *Front Syst Neurosci* 5, 67.
469 doi:10.3389/fnsys.2011.00067
- 470 Bergman, H., Wichmann, T., Karmon, B., DeLong, M.R., 1994. The primate subthalamic
471 nucleus. II. Neuronal activity in the MPTP model of parkinsonism. *J Neurophysiol* 72,
472 507–520.
- 473 Beurrier, C., Congar, P., Bioulac, B., Hammond, C., 1999. Subthalamic nucleus neurons
474 switch from single-spike activity to burst-firing mode. *J Neurosci* 19, 599–609.

475 Braak, H., Del Tredici, K., Rüb, U., de Vos, R.A.I., Jansen Steur, E.N.H., Braak, E., 2003.
476 Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol. Aging* 24,
477 197–211.

478 Breit, S., Bouali-Benazzouz, R., Benabid, A.-L., Benazzouz, A., 2001. Unilateral lesion of the
479 nigrostriatal pathway induces an increase of neuronal activity of the pedunculopontine
480 nucleus, which is reversed by the lesion of the subthalamic nucleus in the rat. *Eur J*
481 *Neurosci* 14, 1833–1842.

482 Coelho, M., Ferreira, J.J., 2012. Late-stage Parkinson disease. *Nat Rev Neurol*.
483 doi:10.1038/nrneurol.2012.126

484 Datta, S., Siwek, D.F., 2002. Single cell activity patterns of pedunculopontine tegmentum
485 neurons across the sleep-wake cycle in the freely moving rats. *J Neurosci Res* 70, 611–
486 621. doi:10.1002/jnr.10405

487 Dautan, D., Souza, A.S., Huerta-Ocampo, I., Valencia, M., Assous, M., Witten, I.B.,
488 Deisseroth, K., Tepper, J.M., Bolam, J.P., Gerdjikov, T.V., Mena-Segovia, J., 2016.
489 Segregated cholinergic transmission modulates dopamine neurons integrated in distinct
490 functional circuits. *Nat Neurosci* 19, 1025–1033. doi:10.1038/nn.4335

491 Devergnas, A., Piallat, B., Prabhu, S., Torres, N., Louis Benabid, A., David, O., Chabardes,
492 S., 2012. The subcortical hidden side of focal motor seizures: evidence from micro-
493 recordings and local field potentials 135, 2263–2276. doi:10.1093/brain/aws134

494 Fee, M.S., Mitra, P.P., Kleinfeld, D., 1996. Variability of extracellular spike waveforms of
495 cortical neurons. *J Neurophysiol* 76, 3823–3833.

496 Fox, S.H., Brotchie, J.M., 2010. The MPTP-lesioned non-human primate models of
497 Parkinson's disease. Past, present, and future. *Prog Brain Res* 184, 133–157.
498 doi:10.1016/S0079-6123(10)84007-5

499 Galvan, A., Wichmann, T., 2008. Pathophysiology of parkinsonism. *Clin Neurophysiol* 119,
500 1459–1474. doi:10.1016/j.clinph.2008.03.017

501 Garcia-Rill, E., 2015. Waking and the Reticular Activating System in Health and Disease.
502 Academic Press. doi:10.1016/B978-0-12-801385-4.09989-4

503 Goetz, L., Piallat, B., Bhattacharjee, M., Mathieu, H., David, O., Chabardes, S., 2016a. On the
504 Role of the Pedunculopontine Nucleus and Mesencephalic Reticular Formation in
505 Locomotion in Nonhuman Primates. *J Neurosci* 36, 4917–4929.
506 doi:10.1523/JNEUROSCI.2514-15.2016

507 Goetz, L., Piallat, B., Bhattacharjee, M., Mathieu, H., David, O., Chabardès, S., 2016b. The
508 primate pedunculopontine nucleus region: towards a dual role in locomotion and waking
509 state. *Journal of Neural Transmission* 123, 667–678. doi:10.1007/s00702-016-1577-7

510 Goetz, L., Piallat, B., Thibaudier, Y., Montigon, O., David, O., Chabardes, S., 2012. A non-
511 human primate model of bipedal locomotion under restrained condition allowing gait
512 studies and single unit brain recordings. *J Neurosci Methods* 204, 306–317.
513 doi:10.1016/j.jneumeth.2011.11.025

514 Gomez-Gallego, M., Fernandez-Villalba, E., Fernandez-Barreiro, A., Herrero, M.T., 2006.
515 Changes in the neuronal activity in the pedunculopontine nucleus in chronic MPTP-
516 treated primates: an in situ hybridization study of cytochrome oxidase subunit I, choline
517 acetyl transferase and substance P mRNA expression. *J Neural Transm* 114, 319–326.
518 doi:10.1007/s00702-006-0547-x

519 Gut, N.K., Winn, P., 2016. The pedunculopontine tegmental nucleus-A functional hypothesis
520 from the comparative literature. *Mov Disord n/a–n/a*. doi:10.1002/mds.26556

521 Hamani, C., Stone, S., Laxton, A., Lozano, A.M., 2007. The pedunculopontine nucleus and
522 movement disorders: anatomy and the role for deep brain stimulation. *Parkinsonism Relat*
523 *Disord* 13 Suppl 3, S276–80. doi:10.1016/S1353-8020(08)70016-6

524 Hirsch, E.C., Graybiel, A.M., Duyckaerts, C., Javoy-Agid, F., 1987. Neuronal loss in the

525 pedunclopontine tegmental nucleus in Parkinson disease and in progressive supranuclear
526 palsy. *Proc Natl Acad Sci USA* 84, 5976–5980.

527 Imbert, C., Bezard, E., Guitraud, S., Boraud, T., Gross, C.E., 2000. Comparison of eight
528 clinical rating scales used for the assessment of MPTP-induced parkinsonism in the
529 Macaque monkey. *J Neurosci Methods* 96, 71–76.

530 Israel, Z., Bergman, H., 2008. Pathophysiology of the basal ganglia and movement disorders:
531 from animal models to human clinical applications. *Neurosci Biobehav Rev* 32, 367–377.
532 doi:10.1016/j.neubiorev.2007.08.005

533 Jellinger, K., 1988. The pedunclopontine nucleus in Parkinson“s disease, progressive
534 supranuclear palsy and Alzheimer”s disease. *J Neurol Neurosurg Psychiatr* 51, 540–543.
535 doi:10.1136/jnnp.51.4.540

536 Jones, B.E., 2005. From waking to sleeping: neuronal and chemical substrates. *Trends*
537 *Pharmacol. Sci.* 26, 578–586. doi:10.1016/j.tips.2005.09.009

538 Kang, Y., Kitai, S.T., 1990. Electrophysiological properties of pedunclopontine neurons and
539 their postsynaptic responses following stimulation of substantia nigra reticulata 535, 79–
540 95. doi:10.1016/0006-8993(90)91826-3

541 Karachi, C., Grabli, D., Bernard, F.A., Tandé, D., Wattiez, N., Belaid, H., Bardinnet, E.,
542 Prigent, A., Nothacker, H.-P., Hunot, S., Hartmann, A., Lehéricy, S., Hirsch, E.C.,
543 François, C., 2010. Cholinergic mesencephalic neurons are involved in gait and postural
544 disorders in Parkinson disease. *The Journal of clinical investigation* 120, 2745–2754.
545 doi:10.1172/JCI42642

546 Kobayashi, Y., Inoue, Y., Yamamoto, M., Isa, T., Aizawa, H., 2002. Contribution of
547 pedunclopontine tegmental nucleus neurons to performance of visually guided saccade
548 tasks in monkeys. *J Neurophysiol* 88, 715–731.

549 Kojima, J., Yamaji, Y., Matsumura, M., Nambu, A., Inase, M., Tokuno, H., Takada, M., Imai,
550 H., 1997. Excitotoxic lesions of the pedunclopontine tegmental nucleus produce
551 contralateral hemiparkinsonism in the monkey. *Neurosci Lett* 226, 111–114.

552 Lau, B., Welter, M.-L., Belaid, H., Fernandez Vidal, S., Bardinnet, E., Grabli, D., Karachi, C.,
553 2015. The integrative role of the pedunclopontine nucleus in human gait. *Brain* 138,
554 1284–1296. doi:10.1093/brain/awv047

555 Lavoie, B., Parent, A., 1994a. Pedunclopontine nucleus in the squirrel monkey: projections
556 to the basal ganglia as revealed by anterograde tract-tracing methods. *J Comp Neurol* 344,
557 210–231. doi:10.1002/cne.903440204

558 Lavoie, B., Parent, A., 1994b. Pedunclopontine nucleus in the squirrel monkey: cholinergic
559 and glutamatergic projections to the substantia nigra. *J Comp Neurol* 344, 232–241.
560 doi:10.1002/cne.903440205

561 Legéndy, C.R., Salcman, M., 1985. Bursts and recurrences of bursts in the spike trains of
562 spontaneously active striate cortex neurons. *J Neurophysiol* 53, 926–939.

563 Leonard, C.S., Llinás, R., 1994. Serotonergic and cholinergic inhibition of mesopontine
564 cholinergic neurons controlling REM sleep: an in vitro electrophysiological study.
565 *Neuroscience* 59, 309–330.

566 Llinás, R., Urbano, F.J., Leznik, E., Ramírez, R.R., van Marle, H.J.F., 2005. Rhythmic and
567 dysrhythmic thalamocortical dynamics: GABA systems and the edge effect. *Trends in*
568 *Neurosciences* 28, 325–333. doi:10.1016/j.tins.2005.04.006

569 Llinás, R.R., Steriade, M., 2006. Bursting of Thalamic Neurons and States of Vigilance. *J*
570 *Neurophysiol* 95, 3297–3308. doi:10.1152/jn.00166.2006

571 Lobb, C., 2014. Abnormal Bursting as a Pathophysiological Mechanism in Parkinson's
572 Disease. *Basal Ganglia* 3, 187–195. doi:10.1016/j.baga.2013.11.002

573 Martinez-Gonzalez, C., Wang, H.-L., Micklem, B.R., Bolam, J.P., Mena-Segovia, J., 2012.
574 Subpopulations of cholinergic, GABAergic and glutamatergic neurons in the

575 pedunclopontine nucleus contain calcium-binding proteins and are heterogeneously
576 distributed. *Eur J Neurosci* 35, 723–734. doi:10.1111/j.1460-9568.2012.08002.x

577 Matsumura, M., 2005. The pedunclopontine tegmental nucleus and experimental
578 parkinsonism. *A review* 252 Suppl 4, IV5–IV12. doi:10.1007/s00415-005-4003-x

579 Matsumura, M., Watanabe, K., Ohye, C., 1997. Single-unit activity in the primate nucleus
580 tegmenti pedunclopontinus related to voluntary arm movement. *Neurosci Res* 28, 155–
581 165.

582 Mazzone, P., Lozano, A., Stanzione, P., Galati, S., Scarnati, E., Peppe, A., Stefani, A., 2005.
583 Implantation of human pedunclopontine nucleus: a safe and clinically relevant target in
584 Parkinson's disease. *NeuroReport* 16, 1877–1881.

585 Mena-Segovia, J., Bolam, J.P., 2017. Rethinking the Pedunclopontine Nucleus: From
586 Cellular Organization to Function. *Neuron* 94, 7–18. doi:10.1016/j.neuron.2017.02.027

587 Mena-Segovia, J., Bolam, J.P., Magill, P.J., 2004. Pedunclopontine nucleus and basal
588 ganglia: distant relatives or part of the same family? *J Neurosci* 27, 585–588.
589 doi:10.1016/j.tins.2004.07.009

590 Mena-Segovia, J., Sims, H.M., Magill, P.J., Bolam, J.P., 2008. Cholinergic brainstem neurons
591 modulate cortical gamma activity during slow oscillations. *The Journal of Physiology*
592 586, 2947–2960. doi:10.1113/jphysiol.2008.153874

593 Morissette, M., Paolo, T., 2017. Non-human primate models of PD to test novel therapies.
594 *Journal of Neural Transmission* 1–34. doi:10.1007/s00702-017-1722-y

595 Nandi, D., Aziz, T.Z., Giladi, N., Winter, J., Stein, J.F., 2002a. Reversal of akinesia in
596 experimental parkinsonism by GABA antagonist microinjections in the pedunclopontine
597 nucleus *J Neurosci* 22, 2418–2430.

598 Nandi, D., Aziz, T.Z., Liu, X., Stein, J.F., 2002b. Brainstem motor loops in the control of
599 movement. *Mov Disord* 17 Suppl 3, S22–7.

600 Orieux, G., François, C., Féger, J., Yelnik, J., Vila, M., Ruberg, M., Agid, Y., Hirsch, E.C.,
601 2000. Metabolic activity of excitatory parafascicular and pedunclopontine inputs to the
602 subthalamic nucleus in a rat model of Parkinson's disease. *J Neurosci* 20, 79–88.

603 Pahapill, P.A., Lozano, A.M., 2000. The pedunclopontine nucleus and Parkinson's disease.
604 *J Neurosci* 20 (Pt 9), 1767–1783.

605 Palombo, E., Porrino, L.J., Bankiewicz, K.S., Crane, A.M., Sokoloff, L., Kopin, I.J., 1990.
606 Local cerebral glucose utilization in monkeys with hemiparkinsonism induced by
607 intracarotid infusion of the neurotoxin MPTP. *J Neurosci* 10, 860–869.

608 Parent, M., Lévesque, M., Parent, A., 1999. The pallidofugal projection system in primates:
609 evidence for neurons branching ipsilaterally and contralaterally to the thalamus and
610 brainstem. *J Chem Neuroanat* 16, 153–165.

611 Piallat, B., Chabardes, S., Torres, N., Fraix, V., Goetz, L., Seigneuret, E., Bardin, E.,
612 Ferraye, M., Debu, B., Krack, P., Yelnik, J., Pollak, P., Benabid, A.-L., 2009. Gait is
613 associated with an increase in tonic firing of the sub-cuneiform nucleus neurons.
614 *Neuroscience* 158, 1201–1205. doi:10.1016/j.neuroscience.2008.10.046

615 Piallat, B., Polosan, M., Fraix, V., Goetz, L., David, O., Fenoy, A., Torres, N., Quesada, J.-L.,
616 Seigneuret, E., Pollak, P., Krack, P., Bougerol, T., Benabid, A.L., Chabardès, S., 2011.
617 Subthalamic neuronal firing in obsessive-compulsive disorder and Parkinson disease.
618 *Ann Neurol* 69, 793–802. doi:10.1002/ana.22222

619 Pienaar, I.S., Elson, J.L., Racca, C., Nelson, G., Turnbull, D.M., Morris, C.M., 2013.
620 Mitochondrial Abnormality Associates with Type-Specific Neuronal Loss and Cell
621 Morphology Changes in the Pedunclopontine Nucleus in Parkinson Disease. *The*
622 *American Journal of Pathology* 183, 1826–1840. doi:10.1016/j.ajpath.2013.09.002

623 Pienaar, I.S., Vernon, A., Winn, P., 2017. The Cellular Diversity of the Pedunclopontine
624 Nucleus. *The Neuroscientist* 107385841668247–17. doi:10.1177/1073858416682471

625 Plaha, P., Gill, S.S., 2005. Bilateral deep brain stimulation of the pedunculopontine nucleus
626 for Parkinson's disease. *NeuroReport* 16, 1883–1887.

627 Plenz, D., Kital, S.T., 1999. A basal ganglia pacemaker formed by the subthalamic nucleus
628 and external globus pallidus. *Nature* 400, 677–682. doi:10.1038/23281

629 Przedborski, S., Jackson-Lewis, V., Naini, A.B., Jakowec, M., Petzinger, G., Miller, R.,
630 Akram, M., 2001. The parkinsonian toxin 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine
631 (MPTP): a technical review of its utility and safety. *J Neurochem* 76, 1265–1274.

632 Raz, A., Vaadia, E., Bergman, H., 2000. Firing patterns and correlations of spontaneous
633 discharge of pallidal neurons in the normal and the tremulous 1-methyl-4-phenyl-1,2,3,6-
634 tetrahydropyridine vervet model of parkinsonism. *Journal of Neuroscience* 20, 8559–
635 8571.

636 Rinne, J.O., Ma, S.Y., Lee, M.S., Collan, Y., Røyttä, M., 2008. Loss of cholinergic neurons in
637 the pedunculopontine nucleus in Parkinson's disease is related to disability of the patients.
638 *Parkinsonism Relat Disord* 14, 553–557. doi:10.1016/j.parkreldis.2008.01.006

639 Rolland, A.-S., Karachi, C., Muriel, M.-P., Hirsch, E.C., François, C., 2011. Internal pallidum
640 and substantia nigra control different parts of the mesopontine reticular formation in
641 primate. *Mov Disord* 26, 1648–1656. doi:10.1002/mds.23705

642 Rolland, A.-S., Tandé, D., Herrero, M.-T., Luquin, M.-R., Vazquez-Claverie, M., Karachi, C.,
643 Hirsch, E.C., François, C., 2009. Evidence for a dopaminergic innervation of the
644 pedunculopontine nucleus in monkeys, and its drastic reduction after MPTP intoxication.
645 *J Neurochem* 110, 1321–1329. doi:10.1111/j.1471-4159.2009.06220.x

646 Ryczko, D., Cone, J.J., Alpert, M.H., Goetz, L., Auclair, F., Dubé, C., Parent, M., Roitman,
647 M.F., Alford, S., Dubuc, R., 2016. A descending dopamine pathway conserved from
648 basal vertebrates to mammals. *Proc Natl Acad Sci USA* 113, E2440–9.
649 doi:10.1073/pnas.1600684113

650 Rye, D.B., Saper, C.B., Lee, H.J., Wainer, B.H., 1987. Pedunculopontine tegmental nucleus
651 of the rat: cytoarchitecture, cytochemistry, and some extrapyramidal connections of the
652 mesopontine tegmentum. *J Comp Neurol* 259, 483–528. doi:10.1002/cne.902590403

653 Schapira, A.H.V., Chaudhuri, K.R., Jenner, P., 2017. Non-motor features of Parkinson
654 disease. *Nat Rev Neurosci* 18, 435–450. doi:10.1038/nrn.2017.62

655 Shimamoto, S.A., Larson, P.S., Ostrem, J.L., Glass, G.A., Turner, R.S., Starr, P.A., 2010.
656 Physiological identification of the human pedunculopontine nucleus. *J Neurol Neurosurg*
657 *Psychiatr* 81, 80–86. doi:10.1136/jnnp.2009.179069

658 Shink, E., Sidibé, M., Smith, Y., 1997. Efferent connections of the internal globus pallidus in
659 the squirrel monkey: II. Topography and synaptic organization of pallidal efferents to the
660 pedunculopontine nucleus. *J Comp Neurol* 382, 348–363.

661 Shute, C.C., Lewis, P.R., 1967. The ascending cholinergic reticular system: neocortical,
662 olfactory and subcortical projections. 90, 497–520.

663 Song, D.D., Haber, S.N., 2000. Striatal responses to partial dopaminergic lesion: evidence for
664 compensatory sprouting. *Journal of Neuroscience* 20, 5102–5114.

665 Stein, J.F., Aziz, T.Z., 2012. Basal ganglia output to the PPN, a commentary 233, 745–746.
666 doi:10.1016/j.expneurol.2011.11.023

667 Steriade, M., 1996. Arousal: revisiting the reticular activating system. *Science* 272, 225–226.

668 Steriade, M.M., McCarley, R.W., 2013. *Brainstem Control of Wakefulness and Sleep*.
669 Springer Science & Business Media.

670 Takakusaki, K., Shiroyama, T., Yamamoto, T., Kitai, S.T., 1996. Cholinergic and
671 noncholinergic tegmental pedunculopontine projection neurons in rats revealed by
672 intracellular labeling. *J Comp Neurol* 371, 345–361. doi:10.1002/(SICI)1096-
673 9861(19960729)371:3<345::AID-CNE1>3.0.CO;2-2

674 Takakusaki, K., Tomita, N., Yano, M., 2008. Substrates for normal gait and pathophysiology

675 of gait disturbances with respect to the basal ganglia dysfunction. 255 Suppl 4, 19–29.
676 doi:10.1007/s00415-008-4004-7

677 Tattersall, T.L., Stratton, P.G., Coyne, T.J., Cook, R., Silberstein, P., Silburn, P.A., Windels,
678 F.C.O., Sah, P., 2014. Imagined gait modulates neuronal network dynamics in the human
679 pedunculopontine nucleus. *Nat Neurosci* 1–8. doi:10.1038/nn.3642

680 Wang, H.-L., Morales, M., 2009. Pedunculopontine and laterodorsal tegmental nuclei contain
681 distinct populations of cholinergic, glutamatergic and GABAergic neurons in the rat. *Eur*
682 *J Neurosci* 29, 340–358. doi:10.1111/j.1460-9568.2008.06576.x

683 Weinberger, M., Hamani, C., Hutchison, W.D., Moro, E., Lozano, A.M., Dostrovsky, J.O.,
684 2008. Pedunculopontine nucleus microelectrode recordings in movement disorder
685 patients. *Exp Brain Res* 188, 165–174. doi:10.1007/s00221-008-1349-1

686 Welter, M.L., Burbaud, P., Fernandez-Vidal, S., Bardinnet, E., Coste, J., Piallat, B., Borg, M.,
687 Besnard, S., Sauleau, P., Devaux, B., Pidoux, B., Chaynes, P., Tézenas du Montcel, S.,
688 Bastian, A., Langbour, N., Teillant, A., Haynes, W., Yelnik, J., Karachi, C., Mallet, L.,
689 2011. Basal ganglia dysfunction in OCD: subthalamic neuronal activity correlates with
690 symptoms severity and predicts high-frequency stimulation efficacy. *Transl Psychiatry* 1,
691 e5. doi:10.1038/tp.2011.5

692 Wichmann, T., Bergman, H., DeLong, M.R., 1994. The primate subthalamic nucleus. I.
693 Functional properties in intact animals. *J Neurophysiol* 72, 494–506.

694 Wichmann, T., Bergman, H., DeLong, M.R., 2017. Basal ganglia, movement disorders and
695 deep brain stimulation: advances made through non-human primate research. *Journal of*
696 *Neural Transmission* 1–12. doi:10.1007/s00702-017-1736-5

697 Wichmann, T., Bergman, H., Starr, P.A., Subramanian, T., Watts, R.L., DeLong, M.R., 1999.
698 Comparison of MPTP-induced changes in spontaneous neuronal discharge in the internal
699 pallidal segment and in the substantia nigra pars reticulata in primates. *Exp Brain Res*
700 125, 397–409.

701 Wichmann, T., Soares, J., 2006. Neuronal firing before and after burst discharges in the
702 monkey basal ganglia is predictably patterned in the normal state and altered in
703 parkinsonism. *J Neurophysiol* 95, 2120–2133. doi:10.1152/jn.01013.2005

704 Ye, M., Hayar, A., Strotman, B., Garcia-Rill, E., 2010. Cholinergic modulation of fast
705 inhibitory and excitatory transmission to pedunculopontine thalamic projecting neurons. *J*
706 *Neurophysiol* 103, 2417–2432. doi:10.1152/jn.01143.2009

707 Zweig, R.M., Jankel, W.R., Hedreen, J.C., Mayeux, R., Price, D.L., 1989. The
708 pedunculopontine nucleus in Parkinson's disease. *Ann Neurol* 26, 41–46.
709 doi:10.1002/ana.410260106

710
711

712 **FIGURE LEGENDS**

713

714 **Figure 1:** Base line mean firing rate distribution in primate K and primate T in normal
715 and MPTP states. (a) : Normal state. (b) MPTP state. Y axis represents the number of
716 neurons and X axis represents the mean firing rate lustered in bins. The center of the
717 first bin is at 5 Hz and bin width is 10 Hz.

718

719 **Figure 2:** A: Evolution of the mean firing rate of cMRF neurons after MPTP intoxication.
720 B: Evolution of the mean firing rate of cMRF pattern sub-groups after MPTP

721 intoxication. The box plots summarize the distribution of the firing rate, with the limits
722 of the box representing the 25th and 75th quartiles and the central line showing the
723 median sample value. The whiskers extend to the min and max values. (* Significant $p <$
724 0.05 Mann-Whitney U-test). C: Burst discharge characteristics: a) Mean Poisson surprise
725 value and b) mean percentage of spikes in burst per pattern group, per primate, in
726 normal and MPTP state according to the methods developed by Legendy and Salcman
727 (Legendy and Salcman, 1985) and (Baron et al., 2011) respectively (** Significant $p <$
728 0.01 - * Significant $p < 0.05$ - Mann-Whitney U-test).


729


730 **Figure 3:** Spike discharge pattern in normal and MPTP state in the two primates:
731 relative proportion of neurons in the different pattern sub-groups (in percentage). A:
732 Comparison between 3 pattern sub-groups (* Significant $p < 0.05$ - χ^2 -test). B:
733 Comparison between burst *versus* non-burst neurons. Primates are treated as stratum in
734 statistical test (* Significant $p < 0.05$ - Cochran-Mantel-Haenszel χ^2 -test).


735


736 **Figure 4:** Relative proportion of oscillatory *versus* non-oscillatory burst neurons in A
737 and mean burst oscillatory frequency in normal and MPTP state in B. The box plots in B
738 summarize the distribution of the neurons coordinates, with the limits of the box
739 representing the 25th and 75th quartiles and the central line showing the median
740 sample value. The whiskers extend to the min and max values. (** Significant $p < 0.01$ -
741 Mann-Whitney U-test).


742




















■ Oscillatory burst
□ Non-oscillatory burst

