



HAL
open science

Contribution of a better maxillofacial phenotype in Silver–Russell syndrome to define a better orthodontics and surgical management

S. Vo Quang, E. Galliani, S. Eche, C. Tomat, B. Fauroux, A. Picard, N. Kadlub

► To cite this version:

S. Vo Quang, E. Galliani, S. Eche, C. Tomat, B. Fauroux, et al.. Contribution of a better maxillofacial phenotype in Silver–Russell syndrome to define a better orthodontics and surgical management. *Journal of Stomatology, Oral and Maxillofacial Surgery*, 2019, 120, pp.110 - 115. 10.1016/j.jormas.2018.10.011 . hal-03486629

HAL Id: hal-03486629

<https://hal.science/hal-03486629>

Submitted on 20 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Contribution of a better maxillofacial phenotype in Silver Russell Syndrome to define a better orthodontics and surgical management

Auteurs

Solène Vo Quang¹, Eva Galliani^{1, 2}, Sophie Eche^{1, 2}, Catherine Tomat^{1, 2}, Brigitte Fauroux^{3, 4}, Arnaud Picard^{1, 2, 4}, Natacha Kadlub^{1, 2, 4}

1. APHP, Necker Enfants Malades, service de Chirurgie Maxillo-faciale et Chirurgie Plastique, 75015 Paris, France
2. APHP, Necker Enfants Malades, Centre de Reference des fentes et malformations faciales, 75015 Paris, France
3. APHP, Necker Enfants Malades, Unité de ventilation non invasive et du sommeil de l'enfant, 75015 Paris, France
4. Université Paris Descartes, 75006 Paris, France

Corresponding author

Natacha Kadlub

Service de Chirurgie Maxillofaciale et Chirurgie Plastique

APHP, Necker Enfants Malades

149 rue de Sevres

75015 Paris, France

Phone: +33 1 7139 67 54

@: natacha.kadlub@aphp.fr

Contribution of a better maxillofacial phenotype in Silver Russell Syndrome (SRS) to define a better orthodontics and surgical management

Abstract

Introduction: Maxillofacial phenotype for SRS is incompletely described in literature. The aim of this study was to describe a maxillofacial phenotype for SRS, to determine a better treatment.

Materials and methods: A retrospective study was conducted including 37 patients with SRS. 24-control patients had been included and appereled. The subjective clinical examination included analyzes of SRS defined criteria. Frontal and lateral photographs had been reviewed, according to Farkas analysis; dental photographs had been examined for the deep-bite and the crowding severity. Radiologic cephalometric analysis had been reviewed.

Results: Maxillofacial examination showed protruding forehead (55%), anteverted ears (55%) and low set ears (16%), small triangular face (48%); retrognathia (29%) and micrognathia (13%). SSR patients presented a lower forehead transverse growth, forehead height, and higher sagittal and transverse mandibular growth than control patients. Deep-bite was present in 21 patients of patient, and crowding in 17 patients. Cephalometric analysis showed 18 patients with the skeletal class II. We did not note a correlation between sleep apnea and retrognathia, neither between genetic anomalies and craniofacial phenotype.

Conclusion: In this study we showed new SRS characteristics: small forehead, small mandible, skeletal class II and a dental phenotype, leading to a specific maxillofacial and orthopedic management.

Key words: Silver Russell syndrome; maxilla-facial phenotype; retrognathia

Introduction

In 1953 Silver et al.¹ described a syndrome associating congenital hemihypertrophia, shortness of stature and elevated urinary gonadotropins. One year later Russell² published five cases associating craniofacial abnormalities and severe intra-uterine growth retardation. Subsequently, Black³ and Rimoin⁴ defined Silver Russell Syndrome (SRS), as the combination of intrauterine and/or postnatal growth retardation, a facial dimorphism and asymmetry of members. SSR is an extremely rare disease with an incidence estimated at 1-30/100 000 per birth⁵. SRS is a heterogeneous condition based on the identification of clinical features. Recently, molecular genetic anomalies associated to SRS had been identified, from 35 to 50% of individuals with SRS presents a hypomethylation of the paternal imprinting center 1 (IC1) of chromosome 11p15.5 (region regulation IGF2/H19 locus) and 10% have maternal disomy for chromosome 7 (mUPD7)⁶⁻⁹. SRS is considered as an imprinting disease. A new clinical scoring system (sensitivity = 98%) for the detection of patients with molecular anomalies associated SRS had been recently described¹⁰. Diagnosis of SRS imposed prenatal growth retardation associated to three of this five following criteria: post natal growth retardation (>2DS), relative macrocephaly, prominent forehead, and body asymmetry, feeding difficulties during early childhood. This score described 2 craniomaxillofacial features whereas SRS initial definition included a facial dimorphism. A new Consensus Statement summarizes recommendations for clinical diagnosis, investigation and management of patients with Silver-Russell syndrome¹¹. Although the definition of SSR includes facial dimorphism, craniomaxillofacial characteristics of SRS are not precisely described in literature^{9,10}.

Otherwise, micro-mandible associated syndromes are often responsible of obstructive sleep apnea^{12,13}, moreover, in other short stature imprinting disease, as Willy-Prader syndrome, central and obstructive sleep apnea had been identified^{14,15}. Sleep apnea syndrome had never been explored in SRS condition.

The aim of this study was to describe and define a maxillofacial phenotype for Silver Russell syndrome, to study its association with sleep apnea troubles, and finally to propose a guideline for orthodontic and maxillofacial management.

Materials and methods

Patients

We conducted a retrospective study from 2009 to 2015, including 37 patients with SRS syndrome before growth hormone therapy registered in the database of the French reference centre for rare maxillofacial disease.

All patients had a genetic examination to seek for a molecular diagnosis.

Patients were evaluated at the age of their first consultation in the reference centre.

Control patients

We included 24 control patients, apperelled to SSR patient according to their civil age, dental age and gender. Control patient were included from non-rare disease patient database of the same centre. Patients with facial malformation, genetic diseases, facial trauma, protruding ear, and central or obstructive apnea were excluded.

Clinical Evaluation

All patients had a clinical evaluation carried out by the one independent senior maxillofacial surgeon.

The subjective and independant clinical examination included the analyse of the following defined criteria: facial dimorphism, a bulging forehead, a small triangular face, retrognathia, micrognathia, bluish sclera, anteverted ears, low-set ears, a thin upper lip, a shrill voice, according to the phenotypic characteristics described in literature (Figure 1.A-D)¹⁰. Maxillofacial clinical evaluation included occlusal description with the determination of dental Angle class, the vertical, sagittal and transversal anomalies. Furthermore the volume of the tonsils was noted to exclude local cause of obstructive apnea.

Frontal and lateral facial Photographs analysis

At each consultation, one professional photographer had performed frontal and lateral photographs. Frontal and lateral photographs had been analysed, according to Farkas analysis (Figure 1C-D)¹⁶⁻¹⁸ by two different and independent investigators, at the age of the first medical consultation in the reference centre (the values of the measures were weighted to the age of the patient). The transverse forehead growth ((ft-ft/eu-eu) was evaluated by the ratio of the outer canthi distance (from the left exocathion to the right exocanthion: ft-ft) distance and

the width of the head (distance from the left eurion to the right eurion: eu-eu). The height of the forehead (tr-g/eu-eu) is estimated by the ratio of the vertical distance of the forehead (distance from the hair planting line to the glabella: tr-g) and the width of the head. Transverse mandibular growth (go-go/n-gn) was assessed by the ratio of the lower facial width (distance from the left gonion to the right gonion: go-go) and total facial height (distance from the nasion to the gnathion: n-gn). Sagittal mandibular growth (sn-gn/t-gn) was assessed by the ratio of the lower facial height (distance from the subnasal point to the gnathion: sn-gn) and facial depth (distance from the tragion to the gnathion: t-gn) on left profile photography.

For photographic analysis, each patient had been apperelled to two control patients.

Dental Photographs analysis

From dental photographs, we analysed the deep-bite (figure 2.A) (the maxillary incisor overlapped the lower incisive), and the crowding severity (figure 2.B-E): no crowding, light crowding (few crowding was present or/and Bogue's diastema were absent in lacteal denture), moderate crowding (one incisive was lingual positioned or lacteal canine was lost earlier), important crowding (there were no space for one incisive), severe crowding (teeth were arranged in two rows).

Cephalometric Analysis

Frontal and lateral cephalometric X-Ray had been analysed according to Delaire with Orqualceph software (Informer®, France). The following data were analysed: skeletal class, dental class, respective position of the maxillar and the mandibular bone, the position of the hyoid bone, the existence of a tonsil hypertrophy.

Sleep evaluation

An air recording polysomnography evaluation was performed for each patient after the first consultation and before the Growth hormone treatment initiation. Apnea hypopnea per hour index (AHI), desaturation index, snoring index were evaluated. The respiratory rate, the duration of the breathing analysis and the results of cutaneous blood gas were noted for each patient. Obstructive sleep apnea syndrome was defined by an AHI superior to 5 per hour and sub classified into mild obstructive (AHI from 5 to 15), moderate (from 16 to 30), and severe (>30).

Statistical Analysis

Statistical comparisons were made to compare SSR patient to control using Fisher test, with $p < 0.05$

being considered significant.

Results

Patients

Among the 37-SRS patients included, 6 had been excluded (absence of photographs n=5, absence of apparelled control n=1). Age of the first maxillofacial consultation in the reference centre varied from 2 to 14 Years-old (average, 6.13 YO). Among the 31 patients analysed, 16 were boys. All patients presented the clinical criteria of SRS⁸. Genetic molecular diagnostics has been tested for all children: 17 had hypomethylation of 11p15 chromosome (ICR1); 6 presented a disomy of chromosome 7 (mUPD7); 2 had 11p15 duplication of maternal material; and no mutation had been identified in 6 cases.

Clinical evaluation (Table 2)

Maxillofacial examination showed protruding forehead in 17 (55%) patients; anteverted ears in 17 patients (55%) and low set ears in 5 patients (16%); small triangular face in 15 patients (48%); retrognathia in 9 patients (29%) and micrognathia in 4 (13%); thin upper lip in 5 patients (16%); and shrilly voice in 2 cases (6%). None presented bluish sclera.

Occlusal evaluation showed Angle class 2 in 28 cases (90%), and class 1 in the 3 other patients (10%). Four (13%) patients had hypertrophic tonsils (Table 2).

Patients with hypomethylation of 11p15 chromosome and patient with disomy of chromosome 7 had similar facial phenotype except for the small triangular face that seemed to be more frequent in patients with mUPD7 (Table2). However patients without mutations or with other seemed to present a less pronounced facial phenotype, with principally anteverted ears.

Frontal and lateral facial Photographs analysis (Table 1)

SSR patients presented a significant lower forehead transverse length, forehead height, and sagittal mandibular height than control patients; and a non-significant higher transverse mandibular length than control (figure 3, Table 1).

Dental Photographs analysis

Among the 31 patients analysed, 26 intra oral photographs were available.

Deep-bite was present in 21 patients (80%), and crowding in 17 patients (67%). Crowding was considered as severe in 3 cases (17%), important in 3 cases (17%), moderate in 7 cases (41%), and light in 4 patients (25%).

Cephalometric

A cephalometric X-Ray was performed in 22/31 patients. Cephalometric analysis showed 18 patients with the skeletal class II (81%), 3 with skeletal class I (14%), and 1 with skeletal class III (5%). All the patients with a skeletal class II (n=18) had a mandibular retrognathism with a short ramus and short horizontal branch. Among the patients from the skeletal class I (n=3), 2 presented a hypomaxilly and mandibular retrognathism and 1 patient presented a maxillary and mandibular prognathism. The sole patient with skeletal class III showed a mandibular retromicrognathism associated with hypomaxilly. Almost all patients (n=21/22) displayed a molar dental class II. The hyoid bone was in a normal position in 15 cases, accessioned (from 2 to 4 mm up) in 4 cases, and lowered (from 1 to 9 mm down) in 3 cases.

Sleep apnea

A sleep apnea study was performed on 25 patients, 18 patients presented a mild SAOS (72%), 5 patients presented mild sleep disorders without SAOS, and 2 patients presented a normal polysomnography. Sleep apnea studies were compared to cephalometry analysis (table 3).

Among the 18 patients with mild SAOS, 18 presented a mandibular retromicrognathism, 7 presented an abnormal position of the hyoid bone, 4 showed a tonsil hypertrophy and 2 a long uvula.

Discussion

This study demonstrated the difficulty to describe a typical maxillofacial phenotype in SRS. In this large series, we demonstrated that the facial dimorphism spectrum is often incomplete in SRS patients. However, we showed new SRS characteristics: small forehead, small mandible, skeletal class II and a dental phenotype, leading to a specific maxillofacial and orthopaedic management.

This study represents the largest series of maxillofacial phenotyping published, in our knowledge. Despite the rarity of the disease (incidence: 1-30/100 000)⁷, 31 patients had been

included. As previously described by Netchine^{9, 10}, we confirmed the absence of gender predisposition.

In our study, most patients presented a genetic molecular anomaly, principally a hypomethylation of 11p15 chromosome (ICR1) followed by a disomy of chromosome 7 (mUPD7), these results were consistent with the genetic molecular analysis described in the international literature^{7, 11, 19}. In our series, we showed few phenotypic differences between ICR1 and mUPD7 patients; ICR1 patients presented a more typical dimorphism with the small triangular face. However, patient without either ICR1 or mUPD7 mutation displayed a milder phenotype. In a larger series, Eggerman also noticed differences of phenotype according to genotype, but showed the ICR1 patients presented a higher rate of prominent forehead, and a lower rate of microretrognathia than mUPD7 patients⁸. Recently Azzi et al. also showed this milder facial phenotype in patients without the two usual mutations⁹.

In our series, diagnosis of SRS had been made late, around 6 year-old. SRS syndrome is a rare disease without pathognomonic sign^{1, 2, 11}; indeed its clinical diagnosis is difficult, and often delayed. Since the absence of pathognomonic sign, many clinical scoring had been described¹⁰. Recently, Netchine et al published the first sensitive scoring system including the following criteria: small for gestational age, birth length and/or weight $\leq -2S$ DS; postnatal growth retardation (height $\leq -2SDS$), relative macrocephaly at birth, body asymmetry, feeding difficulties and/or body mass index (BMI) $\leq -2SDS$ in toddlers; protruding forehead at the age of 1–3 years¹¹. The presence of 4 symptoms predicted the diagnosis of SRS with a 98% of sensitivity. From this study, no typical craniofacial phenotype could be described. All patients have facial dimorphism, principally protruding forehead (55%) and anteverted ears (55%). Most of the SSR patients had a skeletal class 2 (81%). SSR children tend to have smaller forehead (transverse diameter and height) and smaller mandible (transverse and sagittal) compared to control patients.

In our series, we analysed the craniofacial phenotype with objective evaluations; farkas analysis in photography and cephalometric analysis. From Farkas analysis¹⁶⁻¹⁸, we showed that SSR patients tended to present a lower forehead transverse length, forehead height, and sagittal mandibular height than control patients. In our study, whereas subjective clinical evaluation showed a low rate of micrognathism and retrognathism; cephalometric analysis showed that all patients presented a short mandible ramus and horizontal branch. Finally, cephalometric analysis demonstrated that microretrognathism was underestimated. Likewise, in literature, the incidence of microretrognathism is probably unevaluated, since most authors did

not evaluate the cephalometric Xray¹². Kotilainen *et al*²⁰, also described from a cephalometric analysis a smaller mandible in SRS patient compared to their control.

Our results showed that 18 patients presented skeletal class II with short ramus and short body; only one patient presented maxillary and mandibular prognathism; and one had a class skeletal III but with a hypomaxilla and a microretrognathism. The lower arch may present, in some cases (3 for this study), a severe crowding with a lower lateral incisor being displaced in a lingual position. The dental development is symmetrical with a delayed dental eruption. Class II malocclusion was present in 90% of the patients and supraclusion in 87% cases, which was higher rate than in standard population²¹.

In 2003, Bergman and coll. studied the craniofacial characteristics, and the malocclusion characteristics of 16 SRS children. They showed that SRS children had smaller linear facial dimensions and deviations in the facial proportions, maxillary and mandibular repositioning, a high rate of malocclusion and a teeth eruption delay²⁵.

Therefore, retromicrognathism may be responsible of SAOS^{12, 13}. Indeed, in our series, 60% of the patients presented a mild SAOS, probably caused by microretrognathism and majored in some cases by tonsil hypertrophy. However, despite this microretrognathism none presented a severe SAOS.

Conclusion

SRS is extremely rare disease characterized by a polymorphic phenotype. Craniomaxillofacial analysis is often incomplete and subjective in literature. We proposed a systematic and normative evaluation with Farkas and cephalometric analysis in SRS patients in a larger series of SRS patients.

We showed new SRS characteristics: small forehead, small mandible, skeletal class II and a dental phenotype (crowding on the lower arch/ supraclusion), leading to a specific maxillofacial and orthopaedic management.

The orthopaedics, orthodontics and orthognathic surgery treatment can be protocolled as described:

1. For upper crowding: we can propose palatal expansion or rapid palatal expansion, which permit a gradually widens the palate and the upper arch, once the first definitive molars are erupted.

2. For light to moderate lower crowding: we can propose stripping (enamel reduction) or mandibular orthodontic widening
3. For important to severe crowding: dental extraction should be avoided (worsening of the skeletal class2 malocclusion), surgery (symphyseal distraction) should be considered.

References :

1. Silver HK, Kiyasu W, George J, Deamer WC. Syndrome of congenital hemihypertrophy, shortness of stature, and elevated urinary gonadotropins. *Pediatrics* 1953; 12:368-76.
2. Russell A. A syndrome of intra-uterine dwarfism recognizable at birth with cranio-facial dysostosis, disproportionately short arms, and other anomalies (5 examples). *Proc R Soc Med* 1954;47:1040-4.
3. Black J. Low birth weight dwarfism. *Arch Dis Child* 1961;6: 633-44,.
4. Rimoin DL. The Silver syndrome in twins. In Bergsma D (ed): "Malformation Syndromes" New York: National Foundation-March of Dimes, BD:OAS V (2) 1969: 183-7.
5. Christoforidis A, Maniadaki I, Stanhope R. Managing children with Russell-Silver syndrome: more than just growth hormone treatment? *J Pediatr Endocrinol Metab.* 2005;18:651-2.
6. Gicquel C, Rossignol S, Cabrol S, Houang M, Steunou V, Barbu V, et al. Epimutation of the telomeric imprinting center region on chromosome 11p15 in Silver-Russell syndrome. *Nat Genet* 2005;37:1003-7.
7. Bartholdi D, Krajewska WM, Ounap K, Gaspar H, Chrzanowska KH, Ilyana H, et al. Epigenetic mutations of the imprinted IGF2-H19 domain in Silver-Russell syndrome (SRS): results from a large cohort of patients with SRS and SRS-like phenotypes. *J Med Genet* 2009;46:192-7.

8. Eggermann T, de Nanclares GP, Maher ER, Temple IK, Tümer Z, Monk D, Mackay DJ, Grønskov K, Riccio A, Linglart A, Netchine I. Imprinting disorders: a group of congenital disorders with overlapping patterns of molecular changes affecting imprinted loci. *Clin Epigenetics* 2015;7:123.
9. Azzi S, Salem J, Thibaud N, Chantot-Bastaraud S, Lieber E, Netchine I, *et al.* A prospective study validating a clinical scoring system and demonstrating phenotypical-genotypical correlations in Silver-Russell syndrome. *J Med Genet* 2015;52:446-53.
10. Netchine I, Rossignol S, Dufourg MN, Azzi S, Rousseau A, Perin L, *et al.* 11p15 imprinting center region 1 loss of methylation is a common and specific cause of typical Russell-Silver syndrome: clinical scoring system and epigenetic-phenotypic correlations. *J Clin Endocrinol Metab* 2007;92:3148-54,.
11. Wakeling EL, Brioude F, Lokulo-Sodipe O, O'Connell SM, Salem J, Bliet J, Canton AP, *et al.* Diagnosis and management of Silver-Russell syndrome: first international consensus statement. *Nat Rev Endocrinol* 2017;13:105-24.
12. Lowe AA, Santamaria JD, Fleetham JA, Price C. Facial morphology and obstructive sleep apnea. *AM J Orthod Dentofacial Orthop* 1986;90:484-91.
13. Lyberg T, Krogstad O, Djupesland G. Cephalometric analysis in patients with obstructive sleep apnoea syndrome. I. Skeletal Morphology. *J Laryngol Otol*, 1989;103:287-292.
14. Festen DA, Wevers M, De Weerd AW, Van Den Bossche RA, Duivenvoorden HJ, Otten BJ, Wit JM, Hokken-Koelega AC. Psychomotor development in infants with Prader-Willi syndrome and associations with sleep-related breathing disorders. *Pedi Res* 2007; 2:221-4.
15. Sedky K, Bennett DS, Pumariega A. Prader Willi syndrome and obstructive sleep apnea: co-occurrence in the pediatric population. *Journal of clinical sleep medicine: J Clin Sleep Med* 2014; 10:403-9.
16. Farkas LG, Hajniš K, Posnick JC. Anthropometric and anthroposcopic findings of the nasal and facial region in cleft patients before and after primary lip and palate repair. *Cleft Palate Craniofac J* 1993;30:1-12.

17. Farkas LG, Katic MJ, Forrest CR. International anthropometric study of facial morphology in various ethnic groups/races. *J Craniofac Surg* 2005;16:615-46.
18. Farkas LG, Katic MJ, Forrest CR. Comparison of craniofacial measurements of young adult African-American and North American white males and females. *Ann Plast Surg* 2007;59:692-8.
19. Bonaldi A, Mazzeu JF, Costa SS, Honjo RS, Bertola DR, Albano LM, Furquim IM, Kim CA, Vianna-Morgante AM. Microduplication of the ICR2 domain at chromosome 11p15 and familial Silver–Russell syndrome. *Am J Med Genet A* 2011;155:2479-83.
20. Kotilainen J, Hölttä P, Mikkonen T, Arte S, Sipilä I, Pirinen S. Craniofacial and dental characteristics of Silver-Russell syndrome. *Am J Med Genet A* 1995;56:229-236.
21. Bergman A, Kjellberg H, Dahigren J. Craniofacial morphology and dental age in children with Silver-Russell Syndrome. *Orthod Craniofac Res* 2003;6:54-62.

Tables

	SRS	Control	p
Forehead transverse growth (ft-ft/eu-eu)	0.72 [0.58-0.84]	0.81 [0.71-0.93]	6.7 x10⁻⁷
Forehead height (tr-g/eu-eu)	0.47 [0.31-0.61]	0.51 [0.34-0.65]	0.048
Transverse mandibular growth (go-go/n-gn)	1.13 [0.84-1.4]	1.09 [0.85-1.29]	0.1
Sagittale mandibular growth (sn-gn/t-gn)	0.53 [0.43-0.68]	0.59 [0.47-0.73]	0.0022

Table 1 : Farkas analysis. Comparison between SSR patients and their controls

	11p15 ICR1 hypomethylation (n=17)	Disomy Chromosome 7 (n=6)	Other mutations or no mutation(n=8)
protruding forehead	n=11 (65%)	n=4 (67%)	n=1 (12.5%)
small triangular face	n=11 (65%)	n=2 (33.3%)	n=2 (25%)
Clinical retrognathia	n=5 (29.5%)	n=2 (33.3%)	n=2 (25%)
Clinical micrognathia	n=5 (29.5%)	n=1 (16.7%)	n=1 (12.5%)
bluish sclera	n=3 (0%)	n=3 (0%)	n=3 (0%)
anteverted ears	n=9 (52.9%)	n=3 (50%)	n=5 (62.5%)
low set ears	n=3 (17.6%)	n=2 (33.3%)	n=3 (0%)
thin upper lip	n=3 (17.6%)	n=1 (16.7%)	n=1 (12.5%)
talking shrilly	n=3 (0%)	n=2 (33.3%)	n=3 (0%)
skeletal class 2	n=17 (100%)	n=6 (100%)	n=5 (62.5%)
hyoid bone ascension	n=4 (23.5%)	n=3 (50%)	n=2 (25%)
Radiological retromicrognathism	n=17 (100%)	n=6 (100%)	n=8 (100%)
tonsils hypertrophy	n= 2 (11.7%)	n=1 (16.7%)	n=1 (12.5%)

Table 2 : Clinical maxillo-facial phenotype and genetic anomalies

skeletal class	I	II	III
Normal sleep evaluation	0	2	0
Sleeping troubles without SAOS	1	3	1
Mild sleep apnea syndroma	2	15	1

Table 3: Sleep apnea studies and cephalometric analysis

Legends:

Figure 1: Phenotypic characteristics and Farkas analysis

- A: bulging forehead, retrognathia, anteverted ears
- B: small triangular face, low-set ears
- C: sagittal mandibular growth (sn-gn/t-gn)
- D: transverse forehead growth (ft-ft/eu-eu), height of the forehead (tr-g/eu-eu), transverse mandibular growth

Figure 2: Dental Photographs analysis

- A: the deep bite (the maxillary incisor overlapped the lower incisive)
- B: the crowding severity -no crowding,
- C: light crowding
- D: moderate crowding
- E: important and severe crowding

Figure 3: Phenotypic characteristics such as four Farkas analysis of our population of SSR patients





