



## **Correction of thoracic adolescent idiopathic scoliosis via a direct convex rod manoeuvre**

Philippe Violas, Cyril Bryand, Constantin Gomes, Paul Sauleau, Grégory Lucas

### **► To cite this version:**

Philippe Violas, Cyril Bryand, Constantin Gomes, Paul Sauleau, Grégory Lucas. Correction of thoracic adolescent idiopathic scoliosis via a direct convex rod manoeuvre. *Orthopaedics & Traumatology: Surgery & Research*, 2019, 105, pp.1171 - 1174. <10.1016/j.otsr.2019.05.007>. <hal-03487515>

**HAL Id: hal-03487515**

**<https://hal.science/hal-03487515v1>**

Submitted on 20 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

## **Correction of Thoracic Adolescent Idiopathic Scoliosis via a Direct Convex Rod**

**Manoeuvre**<sup>1</sup>

Philippe **Violas**<sup>1\*</sup>, Cyril **Bryand**<sup>1</sup>, Constantin **Gomes**<sup>2</sup>, Paul **Sauleau**<sup>2</sup>, Grégory **Lucas**<sup>1</sup> and the **Orthopedics and Traumatology Society of Western France (SOO)**<sup>3</sup>

<sup>1</sup>Service de Chirurgie Pédiatrique, Hôpital Sud, CHU de Rennes, 16 Boulevard de Bulgarie, 35200 Rennes, France

<sup>2</sup>Département de neurophysiologie, Hôpital Pontchaillou, Avenue Henri Le Guilloux, 35200 Rennes, France

<sup>3</sup> 18, Rue de Bellinière, 49800 Trélazé, France

**\*Corresponding author:** Philippe Violas

Tel.: +33 299 284 321

E-mail: [philippe.violas@chu-rennes.fr](mailto:philippe.violas@chu-rennes.fr)

---

<sup>1</sup> Article issued from the SOO (the Orthopedics and Traumatology Society of Western France)

## **ABSTRACT**

The various techniques available for scoliosis surgery via the posterior approach involve positioning implants on either side of the curve and reducing the deformity by manoeuvres on the concave rod or simultaneously on both rods. Correction solely via a direct convex rod manoeuvre would eliminate the need for implants on the concave side. This technique was used to treat thoracic adolescent idiopathic scoliosis in 23 patients with a mean age of 14 years and 9 months. Low-dose biplanar EOS radiographs were obtained before surgery, on post-operative day 7, and at last follow-up (at least 2 years after surgery) to allow comparisons of Cobb's angle (72°, 33°, and 35°, respectively), thoracic kyphosis (21°, 29°, and 26°), lumbar lordosis (58°, 50°, and 55°), and apical vertebra rotation (-26°, -12°, and -11°). Although scoliosis requires corrections in all three dimensions, this technique seems to produce satisfactory outcomes while obviating the need for implants on the concave side, thereby decreasing the risk of iatrogenic adverse events.

**Level of evidence:** IV

**Key words:** Thoracic adolescent idiopathic scoliosis. Posterior instrumentation. Convex rod corrective manoeuvre. In situ contouring.

## **Introduction**

Adolescent idiopathic scoliosis (AIS) usually affects the thoracic spine and is often characterized in the sagittal plane by decreased kyphosis. When performing Cotrel-Dubousset instrumentation, the derotation rod is positioned on the concave side of the curve then rotated to produce sufficient kyphosis [1]. Reports of insufficient vertebral rotation correction with Cotrel-Dubousset instrumentation led to the development of an in situ contouring technique using a pre-contoured concave rod [2]. The correction occurs on the concave side, where implants must therefore be placed. The more recently described technique involving simultaneous translation on two rods combined with derotation on the concave side requires a high density of concave implants or connectors [3]. Another recently reported method uses translation of sublaminar bands and reduction on the concave rod [4-5]. These techniques aim to reduce the coronal deformity and improve kyphosis correction while also derotating the apical vertebra. The same applies to the simultaneous double-rod rotation technique, which also requires implants on the concave side [6].

On the concave side of the curve, the vertebral and pedicular deformities increase the risk of iatrogenic injury due to implants. To decrease this risk, we suggest dispensing with implants on the concave side (at the apical vertebra and at least the two supra- and infra-jacent vertebrae) and evaluating the possibility of deformity correction using only a direct manoeuvre on the convex rod.

## **Technique**

Spinal cord monitoring was performed by an electrophysiology physician throughout the procedure. As shown in the video, the operative technique involved arthrectomy of the

exposed vertebral levels and instrumentation using mono-axial pedicle screws at the levels nearest the apex of the convexity. On the concave side, the apical vertebra and at least the two supra- and infra-jacent vertebrae were left free of instrumentation. The first rod, made of malleable titanium (U1-R6xxHT, diameter 6) was inserted into the in situ contouring implants on the convex side then locked into the most distal implant. Contouring irons were used to correct the deformity from one level to the next, in the coronal plane then in the sagittal plane. The rod was then locked into the most proximal implant. Direct vertebral derotation was applied to the screws at the apex of the convexity, from one level to the next. A second rod, made of rigid titanium (L2-R6xxHT) or cobalt-chrome (U1-R6xxHC) (diameter 6), was inserted on the concave side to stabilise the correction. After bone freshening, transverse connectors were implanted and bone grafting performed using the bone removed during neural arch freshening [7].

## **Results**

This technique was used in 23 patients, 19 females and 4 males, between April 2014 and December 2015. Mean age at surgery was 14 years and 9 months (range, 12-18 years) and mean Risser stage was 4. Curve distribution according to Lenke was as follows: 1, n=5; 2, n=15; and 3, n=3. Low-dose biplanar EOS radiographs were obtained before surgery, on post-operative day 7, and at last follow-up to allow comparisons of Cobb angle, thoracic kyphosis (T4-T12), and lumbar lordosis (L1-L5); in addition, 3D reconstruction using sterEOS software (EOS Imaging, Paris, France) was performed to enable comparisons of apical vertebra rotation (AVR) (Figure 1). A follow-up of at least 2 years was required; mean follow-up was 37 months (range, 27-36 months). Table 1 reports the radiographic parameter values at the three time points. Mean instrumentation length was 12 levels. Mean operative

time was 230 minutes and mean blood loss was 200 mL. No patient required blood transfusion.

## **Discussion**

With this technique, the initial correction from one implant to the next was performed first in the coronal plane then in the sagittal plane in order to create sufficient kyphosis. Then, once the rod was locked at both ends, a direct derotation manoeuvre applied to the screws at the apex from one implant to the next induced no loss of the previously established kyphosis. Compared to other studies in patients with large pre-operative angles ( $72^\circ$ ), we obtained similar quantitative correction in the coronal plane, as well as in the horizontal plane (from  $-21^\circ$  to  $-11^\circ$  at last follow-up) [8-9]. Despite the improved kyphosis restoration ( $21^\circ$  before surgery and  $26^\circ$  at last follow-up), our technique did not allow as much correction in the sagittal plane compared to simultaneous translation on two rods with pre-contouring [10].

Correction techniques involving manoeuvres on the convex side have been described previously but consistently involved instrumentation of the vertebrae at the apex of the concavity [11-13]. Pedicle screw implantation on the convex side is easier and carries less risk of malposition within the vertebral pedicles [14]. Constraints during contouring are less marked, allowing better preservation of the mechanical properties of the rod. Finally, decreasing implant density was recently shown to decrease costs [15].

The main limitations of this study are the retrospective design and small sample size. In the future, follow-up data from the patients will add to the information in this technical note. More specifically, data are needed to determine whether the correction could be significantly improved depending on the evaluation of curve rigidity. Nevertheless, this technique is

simple, and during the procedure the surgeon can decide to instrument the concave side if the use of a more conventional method seems desirable.

### **Disclosure of interest**

None of the authors has any conflicts of interest to declare.

Funding sources: none

Authors' contribution: All authors contributed to the manuscript

### **FIGURE LEGENDS**

Figure 1. Female patient aged 14 years at surgery

Antero-posterior radiographs before surgery (a), 7 days after surgery (b), and 26 months after surgery (c)

Lateral radiographs before surgery (d), 7 days after surgery (e), and 26 months after surgery (f)

Figure 2. Photographs of the same patient as in Figure 1: view from the back before (a) and after (b) surgery; view from the side before (c) and after (d) surgery; view of the gibbus before (e) and after (f) surgery

## References

- [1] Cotrel Y, Dubousset J. A new technic for segmental spinal osteosynthesis using the posterior approach. *Rev Chir Orthop* 1984;70:489-94.
- [2] Steib JP, Dumas R, Mitton D, et al. Surgical correction of scoliosis by in situ contouring: a detorsion analysis. *Spine* 2004;29:193-9.
- [3] Clément JL, Chau E, Vallade MJ, et al. Simultaneous translation on two rods is an effective method for correction of hypokyphosis in AIS: radiographic results of 24 hypokyphotic thoracic scoliosis with 2 years follow-up. *Eur Spine J* 2001;20:1149-56.
- [4] Ilharreborde B, Even J, Lefevre Y, et al. Hybrid constructs for tridimensional correction of the thoracic spine in adolescent idiopathic scoliosis : a comparative analysis of universal clamps versus hooks. *Spine* 2010;35:306-14.
- [5] Sales de Gauzy J, Jouve JL, Ilharreborde B, et al. Use of the universal clamp in adolescent idiopathic scoliosis s. *Eur Spine J* 2014;23(Suppl4):S446-51.
- [6] Sudo H, Ito M, Abe Y et al: Surgical treatment of Lenke 1 thoracic adolescent idiopathic scoliosis with maintenance of kyphosis using the simultaneous double-rod rotation technique. *Spine* 2014, 39:1163-9.
- [7] Violas P, Chapuis M, Bracq H. Local autograft bone in the surgical management of adolescent idiopathic scoliosis. *Spine* 2004;29:189-92.
- [8] Clements DH, Betz RR, Newton PO, et al. Correlation of scoliosis curve correction with the number and type of fixation anchors. *Spine* 2009;34:2147–50.
- [9] Asher M, Lai SM, Burton D, et al. Safety and efficacy of Isola instrumentation and arthrodesis for adolescent idiopathic scoliosis: two- to 12-year follow-up. *2004 Spine*; 29:2013–23.



- [11] Terai H, Toyoda H, Suzuki A, et al. A new corrective technique for adolescent idiopathic scoliosis: convex manipulation using 6.35 mm diameter pure titanium rod followed by concave fixation using 6.35 mm diameter titanium alloy. *Scoliosis* 2015 11;(Suppl 2):S14.
- [10] Clement JL, Chau E, Geoffray A, Suisse G. Restoration of thoracic kyphosis by simultaneous translation on two rods for adolescent idiopathic scoliosis. *Eur Spine J*. 2014, 23 Suppl 4:S438-45.
- [12] Chang KW. Cantilever bending technique for treatment of large and rigid scoliosis. *Spine* 2003;28:2452-8.
- [13] Yang J, Huang Z, Grevitt MP et al. Double-curve synchronous derotation with convex correction: a new corrective technique for adolescent idiopathic scoliosis with double curves. *J Spinal Disord Tech* 2014, 27:E32-6.
- [14] Uehara M, Takahashi J, Ikegami S, et al. Pedicle screw loosening after posterior spinal fusion for adolescent idiopathic scoliosis in upper and lower instrumented vertebrae having major perforation. *Spine* 2017, 42:1895-900.
- [15] Tannous OO, Banagan KE, Belin EJ, et al. Low-density pedicle screw constructs for adolescent idiopathic scoliosis: Evaluation of effectiveness and cost. *Global Spine J*. 2018, 82:114-20.

Figure 1.




Figure 2




Table 1: Angles measured on radiographs before and after surgery

Mean (range)	Before surgery	Post-operative D7	Last follow-up
Cobb angle	72° (56°;95°)	33° (24°;52°)	35° (24°;52°)
Thoracic kyphosis	21° (-13°;65°)	29° (-4°;46°)	26° (-6°;49°)
Lumbar lordosis	58° (37°;77°)	50° (36°-65°)	55° (40°;68°)
Apical vertebra rotation	-26° (-4°;48°)	-12° (5°;-28°)	-11° (-23°;6°)