



HAL
open science

Can the amniotic membrane be used to treat peripheral nerve defects? A review of literature

M. Bourgeois, F. Loisel, L. Obert, I. Pluvy, F. Gindraux

► To cite this version:

M. Bourgeois, F. Loisel, L. Obert, I. Pluvy, F. Gindraux. Can the amniotic membrane be used to treat peripheral nerve defects? A review of literature. *Hand Surgery and Rehabilitation*, 2019, 38, pp.223 - 232. <10.1016/j.hansur.2019.05.006>. <hal-03487696>

HAL Id: hal-03487696

<https://hal.science/hal-03487696v1>

Submitted on 20 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

Can the amniotic membrane be used to treat peripheral nerve defects? A review of literature

La membrane amniotique peut-elle être utilisée pour traiter les anomalies nerveuses périphériques? Une revue de littérature

Abstract

Human amniotic membrane is currently being used in ophthalmology and dermatology applications. The objective of this review was to establish proof-of-concept for using amniotic membrane to treat peripheral nerve defects. We performed a search using: 1) PubMed with the keywords/MeSH terms: "amnion", "amniotic membrane", "angiogenesis", "anti-microbial", "characteristic", "chorion", "epithelialization", "fibrosis", "gap", "growth factors", "use", "nerve; 2) the American clinical trials registry with "amniotic membrane"; 3) Lim Jeremy's book "A Primer on Amniotic Membrane Regenerative Healing"; 4) the search engine Google. Our findings pointed to the amniotic membrane being a biodegradable and bioactive scaffold that contains many growth factors important for efficient nerve regeneration. Multiple animal studies and the single human clinical trial performed up to now have highlighted its role in preventing recurrence of perineural adhesions, reducing fibrosis, accelerating nerve repair and improving nerve function. Thus, the amniotic membrane has ideal properties for treating peripheral nerve injuries. It could very likely prevent neuroma formation. The best format would be a freeze-dried one containing the amnion and chorion layers in order to preserve all its growth factors, and facilitate its handling and storage in the operating room.

Résumé

L'objectif consistait à établir une preuve de concept sur l'utilisation de la membrane amniotique humaine, déjà largement exploitée en ophtalmologie et en dermatologie, pour traiter une lésion nerveuse périphérique avec perte de substance. Notre méthode de

recherche a utilisé : 1) Le site "PubMed" avec comme mots clefs/MeSH : "amnion", "amniotic membrane", "angiogenesis", "anti-microbial", "characteristic", "chorion", "epithelialization", "fibrosis", "gap", "growth factors", "use", "nerve" ; 2) le site "Clinical trials", avec : "amniotic membrane" ; 3) l'ouvrage de Lim Jeremy "A Primer on membrane regenerative healing" ; 4) le moteur de recherche Google. Les résultats ont montré que la membrane amniotique est un support biodégradable et bioactif possédant de nombreux facteurs de croissance propices à une régénération nerveuse efficace. Les modèles animaux répertoriés et le seul essai clinique ont révélé son action dans la prévention de la récurrence des adhérences périmébrales, la réduction de la fibrose, l'accélération de la régénération de la structure nerveuse et l'amélioration de la récupération de la fonction nerveuse. Ainsi la membrane amniotique possède des propriétés optimales pour l'indication ciblée et empêcherait très certainement la formation d'un névrome. Son format de prédilection serait une forme lyophilisée, avec ses deux feuillets originels afin de conserver le maximum de facteurs de croissance et faciliter sa manipulation et son stockage au bloc opératoire.

Keywords: Antifibrotic; Scaffold; Growth factors; Proof of concept; Amniotic membrane

Mots clefs : Anti-fibrosant ; Support ; Facteurs de croissances ; Preuve de concept

1. Introduction

Traumatic peripheral nerve lesions are a genuine public health problem as there are more than 300,000 cases per year in Europe [1]. The incidence of finger lacerations with nerve injury is up to 6.2/100,000 inhabitants, resulting in a significant cost to society [2].

Several techniques, such as autologous nerve graft, autograft vein conduits, nerve transfer, collagen or silicone tubes, and neurotubes, have been described for repairing peripheral nerve defects [3]. However, these techniques have drawbacks, [4] such as donor site morbidity for autologous grafts, development of fibrosis and limitations on defect size for neurotubes [5] and vein conduits [6].

One of the goals of peripheral nerve repair is to optimize the healing process in order to prevent neuroma development. Posttraumatic neuromas are a benign, disorganized tissue stemming from an unsuccessful attempt at repairing a peripheral nerve, which results in a “pseudotumor”. It appears 1 to 12 months after the injury event. Yütsel et al. believe neuromas develop because of injury to the perineurium [6]. Axons that are not contained in their envelope can escape into adjacent tissues. This phenomenon is accompanied by significant cell proliferation (fibroblasts, Schwann cells) and by blood vessel development. The result is a disorganized network of axons, which are deviated and curled up [7].

In recent years, the amniotic membrane has had tremendous success in repairing soft tissues, particularly in the ophthalmology and dermatology context [8], making it a relevant option in nerve repair. Riccio et al. reported encouraging results in humans for repairing posttraumatic nerve defects 3 to 5 cm in length [9]. This tissue is mainly composed of types I, III, IV and V collagen, laminin, proteoglycans, elastin and fibronectin [10]. It is made up of two layers: the amnion (which is in contact with the amniotic fluid) and the chorion. The amnion is 0.02–0.5 mm thick while the chorion is 1 mm thick. The amnion mainly consists of an epithelium, which is rich in epithelial cells, a basal membrane and a fibroblast layer, which is rich in amniotic mesenchymal stromal cells. The amnion itself is sometimes incorrectly called the “amniotic membrane”. The chorion consists of a reticular layer, which is rich in chorionic mesenchymal stem cells, a basal membrane, and a

trophoblast, which is rich in trophoblast stem cells.

The amniotic membrane contributes to re-epithelialization and has antifibrotic, antiinflammatory, antiangiogenic, antimicrobial and antiviral properties [11,12]. It can be used as an allograft because the absence of immunogenicity. Furthermore, it contributes to the healing of wounds and has analgesic properties. Its mechanical properties—viscoelasticity due to its high levels of collagen and elastin—and its high levels of several growth factors [13] make it a biologically active scaffold well-suited to the challenge of nerve repair [4].

We performed a review of literature to establish proof of concept on the use of this highly innovative and promising tissue for treating peripheral nerve defects.

2. Material and methods

2.1. Search strategy

The review of literature encompassed several different sources:

- PubMed using the keywords/MeSH entry terms: "amnion", "amniotic membrane", "angiogenesis", "anti-microbial", "characteristic", "chorion", "epithelialization", "fibrosis", "gap", "growth factors", "use", "nerve". The Boolean operator "and" was used to joint "Amniotic membrane" with "Angiogenesis"; "Antimicrobial"; "Characteristic"; "Epithelialization"; "Fibrosis"; "Growth factors"; "Use"; "Nerve". Additional articles were added after manually reviewing the reference list of all the included publications.
- The United States clinical trials registry (<https://clinicaltrials.gov/ct2/search>), using the keywords/MeSH entry terms "amniotic membrane" and capturing completed or on-going clinical trials.
- The book by Jeremy Lim: A Primer on Amniotic Membrane Regenerative Healing. MiMedx Group, 2015.
- The Google search engine, mainly to search for marketed products.

One of our research group's prior publications describing the clinical use, patents and marketed products was also consulted [14].

2.2. Selection criteria

For preclinical proof of concept, studies were included if they analyzed amniotic membrane use in animal models of nerve defect or nerve transection.

For clinical trials, studies were included:

- from PubMed if they reported on amniotic membrane use in humans for treating nerve defects.
- from the clinical trials registry when they referred to amniotic membrane use in a rheumatology, orthopedic or trauma context.

In both cases (animal models and clinical trials), only articles in which the amniotic membrane was used as a scaffold were retained. In vitro studies or studies using cells isolated from the tissue were excluded.

For evidence in support of nerve repair/regeneration, the most relevant articles, often the most recent ones and those cited as a reference, were used in our review.

Generally, only articles published in English were retained.

2.3. Screening of studies and data extraction

The article selection and data collection was performed independently by two investigators (MB and FG). The title and abstract were selected based on the following question: Is amniotic membrane use in an animal model or clinical study beneficial for treating nerve defects? An article was evaluated in its entirety if the title and abstract pertained to this screening question. Discrepancies between the two investigators were discussed to select the final articles. Tables were generated and used to collect the relevant information.

3. Results

The PubMed search yielded the following (before 01-OCT-2018):

- Amniotic membrane “and” angiogenesis: 85 articles

- Amniotic membrane “and” antimicrobial: 894 articles
- Amniotic membrane “and” characteristic: 178 articles
- Amniotic membrane “and” epithelialization: 237 articles
- Amniotic membrane “and” fibrosis: 348 articles
- Amniotic membrane “and” growth factors: 1566 articles
- Amniotic membrane “and” use: 456 articles
- Amniotic membrane “and” nerve: 204 articles

The clinical trials registry search (before 01-OCT-2018) yielded 186 studies.

3.1 Amniotic membrane format

The human amniotic membrane, and more specifically the amnion (separated from the chorion), is collected from placenta following Cesarean birth. Tissue from vaginal delivery is not used since it is contaminated by the normal vaginal flora. Several methods exist to store this product: cryopreservation, lyophilization and preservation in dried form [15]. Fresh human amniotic membrane, which is sometimes used in animal models, corresponds to tissue that has not been treated after being collected.

In cryopreserved form, the amniotic membrane is packaged in a solution including a cryoprotectant (glycerol or dimethyl sulfoxide – DMSO) and Dulbecco’s modified Eagle’s medium. The tissue is then stored at –80°C and can be used for up to 1 to 2 years. With this type of preservation, 20% to 50% of the cells survive [16]. This is the benchmark format in current practice and clinical trials [8].

In its lyophilized form, the amniotic membrane is quickly frozen at –50 to –80°C, then freeze-dried in a lyophilizer under vacuum. In some cases, the tissue will undergo pretreatment to destroy any residual viable cells; sterilization by gamma irradiation very often follows the lyophilization step. These two additional processing steps offer new hope for ensuring the safety of amniotic membrane from vaginal delivery, thereby increasing the number of placentas collected. This also allows the allograft to be stored in its original format (two layers). In other cases, the chorion (which is in contact with the maternal tissue) is

generally separated from the amnion to avoid risk of contamination by the mother's cells. The resulting "amnion–chorion" product is thicker and stronger than amnion alone. Lastly, lyophilization allows the tissue to be stored at room temperature (e.g. in an operating room).

In its dehydrated form, the amniotic membrane is stored at room temperature in a biosafety cabinet and exposed to air for varying lengths of time (12 to 24 hours). This is usually followed by gamma irradiation for sterilization purposes.

Whether the tissue is used in its fresh, cryopreserved, lyophilized or dehydrated form, it may be subjected to de-epithelialization immediately after being harvested in order to remove the epithelial cell layer.

Table 1 lists the features of the amniotic membrane based on its format and suitability for the target clinical indication.

3.2. Preclinical proof of concept

From the PubMed search and after the inclusion and exclusion criteria were applied, 15 articles on animal models were analyzed and relevant data were extracted [17-31] (Table 2). Among these 15 studies, 10 were published after 2009, evidence of the recent interest in using amniotic membrane for nerve repair. The studies appear to be highly relevant given the number of animals used, the control groups, comparison with other treatments and functional outcomes.

In these animal models, the human amniotic membrane was used in its fresh or cryopreserved form and was very often de-epithelialized. The dehydrated form was used in only two studies [19,31]. In most studies, the model used is one where a localized 1–1.5 cm nerve defect or lesion is made in the sciatic nerve of rats.

The amniotic membrane typically measures 1–2 cm and is secured at each end by an epineural suture [17,19,24-26,28,30,31], photochemical tissue bonding [22,23,27] or is left unsutured [21,29]. Only a few studies specified the orientation of the applied amnion: mesenchymal side against the nerve [21] or, more recently, epithelial side [29,31], after simulation of nerve wrapping *in vitro* [29].

The literature search showed that its use as a perineural wrap appears conducive to repairing a nerve lesion or defect (Table 2). First, this increases the number of axons, nerve fibers and myelinated fibers [18,19,22,25]. For example, for a 1-cm nerve defect, the number of large-diameter axons is higher when an amniotic membrane is used instead of a silicone tube or nerve autograft [18]. It also appears that the diameter of the configured amnion tube greatly affects the repair; a 1–2 mm diameter tube appears ideal [19]. The G-ratio, which is defined as the ratio of the axon diameter to the nerve fiber diameter, is improved [22,25] and the functional recovery, estimated by the Sciatic Functional Index (SFI), is significant [18,25]. From an electrophysiology perspective, Meng et al. showed that amnion combined with nerve autograft helps to increase the amplitude and reduce conduction latency at 8 weeks [25]. Second, the amniotic membrane has a noteworthy effect on fibrosis and perineural adhesions [21,24,25]. Lemke et al. specified that the tissue prevents the recurrence of severe perineural adhesions, reduces perineural and intraneural fibrosis, accelerates nerve structure regeneration and improves recovery of nerve function [29].

Some authors report the benefits of the amniotic membrane in a vascularized form [17], or combined with skeletal muscle [31], hyaluronic acid [21], granulocyte colony stimulating factor (G-CSF) [26] or betamethasone [28] to potentiate its effect. Lastly, photochemical tissue bonding of the amnion compared to fixation using epineural sutures positively affects axon size, axonal migration, nerve fiber diameter, G-ratio index, muscle mass innervated by the transected then repaired nerve, and myelin sheath thickness [22,23,27].

3.3. Clinical trials

A search on the clinical trials registry reveals that more than 100 clinical studies with human amniotic membrane or its derivatives have been performed or are ongoing. However, we did not find any clinical trials that specifically addressed our research question. Thus we decided to compile the 40 clinical trials related to the musculoskeletal system (surgical and medical pathology) (Table 3): 25 describe treatments for osteoarthritis, rotator cuff disease,

osteochondral lesions, tendinitis or low back pain; 15 relate to tendon lesions in the hand or ankle, anterior cruciate ligament injuries, reducing adhesions during total knee arthroplasty and spinal fusion surgery.

We found a single clinical trial on PubMed describing use of the amniotic membrane as a nerve conduit for treating nerve defects. Riccio et al. reported very promising results in a prospective single-center study performed between March 2012 and September 2013 in the Department of Reconstructive Surgery and Hand Surgery, AOU "Ospedali Riuniti" di Ancona, Italy [9]. The trial involved 5 patients between 22 and 42 years of age (mean of 33.6 years) who had a median nerve injury at the wrist with a 3–5 cm defect (4 cm on average). The time to surgical nerve repair was 0 to 5 months. Fresh or dehydrated amnion was filled with skeletal muscle harvested from an adjacent area. This sheet was then shaped into a tube and secured with 9-0 epineural suture consisting of two stitches at the back and three stitches in front. They found that sensory and motor functions were restored based on the Lister muscular test, Jamar test and Sakellarides classification: one patient had a sensory result of S3, two were S3+ and two were S4; one patient had a motor result of M3 and four had a motor result of M4. This clinical proof of concept was substantiated by a recently published experimental model [31].

The PubMed search also identified a clinical trial that we chose to include in our analysis: treatment of recurrent ulnar nerve syndrome at the elbow in which the patients were treated with an amniotic membrane nerve wrap sutured on itself with 8-0 suture [32]. This retrospective study included 8 patients treated between December 2012 and March 2014. The average pain level on a visual analog scale was lower by 3.5 points after the procedure than before it; the QuickDASH score improved by 30 points, the grip strength by 25 lbs., or 38% compared with the opposite side. Three patients returned to work full time, one patient was able to go from light duty to a full workload, one unemployed patient began a new job and three remained unemployed. No complications were reported.

Nevertheless, this lack of clinical data means that clinical proof of concept studies on amniotic membrane use for treating nerve defects still need to be done.

3.4. Evidence in favor of nerve repair/regeneration

3.4.1. Presence of growth factors

One of the main reasons amniotic membrane is used in tissue engineering is its bioactive nature. In fact, no matter which format is used, this tissue contains a multitude of growth factors [13] including neurotrophic factors such as nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), neurotrophin (NT-3), glial-derived neurotrophic factor (GDNF) and ciliary neurotrophic factor (CNTF) [4].

However, certain conditions impact the type and amount of growth factors in the tissue:

- Maternal and gestational age: membranes from a full-term pregnancy and from donors under 35 years of age have higher growth factor levels [33].
- Harvest site: epidermal growth factor (EGF) concentration can vary two-fold depending on where the amniotic membrane is harvested in the placenta [34]. This variation in concentration may be related to different epithelial cell morphology and different mitochondrial activity depending on the harvest site [35]. A recent study has shown the region opposite the placenta (reflected region) is most suitable for preventing recurrent adhesions and fibrosis [29].
- Preservation of both layers (amnion and chorion): the chorion layer contains 82% of the growth factors, except for EGF, of which 20% is contained in the amnion and 80% in the chorion [13,36].
- Preservation of epithelium in amnion: Fresh amniotic membranes with epithelium are very high in basic fibroblast growth factor (bFGF), EGF, hepatic growth factor receptor (HGFR), keratinocyte growth factor (KGF), keratinocyte growth factor receptor (KGFR), transforming growth factor (TGF) - β 1, 2, 3 and tumor necrosis factor (TNF) - α [37].
- Method of tissue preservation: the fresh form has the most growth factors, followed by the cryopreserved form and then the lyophilized form [38-40].

However, this aspect is controversial in the literature.

3.4.2. Antifibrotic, antiinflammatory and pro- or antiangiogenic properties

These growth factors have specific functions. The antifibrotic effect is one of the primary properties that ensures good nerve healing. Fibroblasts are naturally involved in healing and are activated during the process by TGF- β . The antifibrotic effect of the amniotic membrane is achieved by suppression of TGF- β signaling and reduction in the expression of TGF- β 1, 2 and 3 isoforms and TGF- β receptor II [15]. These antifibrotic properties have been demonstrated in various animal models during peripheral nerve repair (Table 2).

The antiinflammatory effect of the amniotic membrane is guided by inhibition of the expression of proinflammatory cytokines such as bFGF, interferon (IFN) - γ , interleukin (IL) -1 α , 2, 8, 10, platelet derived growth factor (PDGF) and TNF- β on the damaged ocular surface [15].

Lastly, the tissue's mesenchyme has fibroblasts that synthesize proangiogenic growth factors: angiopoietin (Ang) -2, bFGF, EGF, heparin binding epidermal growth factor (HB-EGF), HGF, PDGF-BB, placental growth factor (PLGF) and vascular epidermal growth factor (VEGF) [41]. These trigger proliferation of endothelial cells and guide the formation of new blood vessels. Paradoxically, the amniotic membrane can also have an antiangiogenic effect [42] through the production of other factors such as tissue inhibitor of metalloproteinase (TIMP) -1, 2 or pigment epithelium-derived factor (PEDF) [42,43]. The microenvironment in which the allograft is implanted will drive the pathway toward pro- or antiangiogenic activity.

3.4.3. Semipermeable and absorbable scaffold properties

Whether the amniotic membrane is used in its cryopreserved or lyophilized form, it is an excellent biodegradable scaffold with biochemical and mechanical properties that make it well-suited to nerve regeneration. It has unique nerve conduit capacities because of its structural properties and specific protein elements such as collagen, laminin and proteoglycans. This tissue has been exploited in patents and marketed products [8,16]

(Table 4). It has other specific characteristics that are similar to another human tissue type—the induced membrane—and can function as a bioreactor [44].

As for its degradation, the amniotic membrane (depending on the format used) is still visible when animals are euthanized at 4 weeks [29], 6 weeks [45] and 12 weeks [21,23] after its implantation, but is no longer visible at 4 months (16 weeks) [18]. Lemke et al. point out that the membrane can no longer be differentiated from neighboring tissues at 3 months [29]. Thus, beyond this point in time, we cannot expect the tissue to have a positive impact, although it no longer matters, given the speed of the adhesion process in their study.

4. Discussion

There is little data on using the amniotic membrane to repair peripheral nerve defects. This literature review describes the exceptional ability of this tissue to regenerate nerves and/or reduce adhesions and fibrosis and serves as proof of concept for the targeted clinical indication. It supplements the similar evaluation done for oral surgery [46].

As shown in our review's findings, in addition to its fundamental antifibrotic nature, this bioactive tissue has several growth factors and angiogenic factors that can effectively guide nerve regeneration. In fact, certain growth factors such as NGF, BDNF, NT-3 and TGF- β are involved in the formation of fibrosis [47]. For example, TGF- β is produced by cells recruited to the site of the local damage. It converts fibroblasts into myofibroblasts to repair the extracellular matrix components. It also regulates the production of enzymes such as matrix metalloproteinases (MMPs) that catalyze the degradation of extracellular matrix. In the case of excessive scarring, myofibroblasts synthesize an abnormally high amount of extracellular matrix components and MMPs [48]. The amniotic membrane contributes to the synthesis of MMP inhibitors, TIMP-1,-2,-3 and -4 [49], and has an anti-inflammatory action through modulation of cytokines such as IL-6, IL-10 IL-1 β , IL-9, IL-17a, IL-22, TNF α , IFN γ and prostaglandins [50,51]. In animal models, xenografting of this tissue helps to reduce fibrosis in various contexts: nerve lesions [25], burns and chronic wounds [52], peritoneal [53] or uterine horn adhesions [54] and tendon repair surgery [55]. These properties allow the

amniotic membrane to act as a scaffold for the development of organized perineural connective tissue, which is a fundamental requirement for good quality axon regeneration.

Our review also highlighted the scaffold properties of this human allograft, particularly strength and resorption, which are compatible with the target clinical indication. While the amnion is no longer clearly distinguishable from other tissues 3–4 months after its implantation in animal models, its resorption rate does not seem to interfere with nerve repair. For clinical use, we theoretically will need to master the physical stability of the membrane so it can prevent adhesion from recurring over time. Along the same lines, we cannot expect consistent production of growth factors by amniotic cells but rather early release of these factors immediately after implantation, from the cells or matrix being degraded [29]. We propose using a multilayer product (superimposition of several membranes) and/or the original format (both layers: amnion and chorion) to increase its thickness and thus slow its resorption rate to meet this physical stability requirement.

The last element is that this tissue has very little immunogenicity based on the HLA (human leukocyte antigen) system [56]. On one hand, it has been stated that an allograft of amniotic membrane in animals provides a better environment for nerve regeneration than a xenograft [30]. On the other hand, the foreign body reaction is minimal when fresh human amniotic membrane xenograft is used in animal models [29,30]. Thus it is not much of a leap to think that using this tissue will induce little to no foreign body reaction as an allograft in humans. This will also reduce the likelihood of making the fibrosis worse, which sometimes happens when slightly immunogenic neurotubes are used.

5. Conclusion

The use of amniotic membrane to repair peripheral nerve defects appears very promising based on in vivo animal studies and very early clinical data. This tissue has inherent properties such as bioactivity, immunotolerance and antifibrotic effect that we believe, based on extrapolation, to be very likely to prevent neuroma formation. With the appropriate regulatory approvals, this scaffold could be used in humans, preferably in the

lyophilized form consisting of the two original layers (amnion, chorion). The goals would be to 1) preserve as many growth factors as possible, 2) make it easy to handle and store at room temperature in the operating room, and 3) slow down its resorption rate. Its lyophilized form, often supplemented with a virus inactivation process, is theoretically a safer product. Its optimal format appears to be as a tubular membrane, possibly in multiple layers, secured by epineural sutures at both two ends of the nerve defect.

Conflict of interest:

The authors have no conflict of interest to disclose relative to this article.

Acknowledgements:

This work contributes to the COST Action CA17116 International Network for Translating Research on Perinatal Derivatives into Therapeutic Approaches (SPRINT), supported by COST (European Cooperation in Science and Technology). The authors wish to thank Joanne Archambault, PhD and Joan Wallace for English-language assistance.

References

1. Ichihara S, Inada Y, Nakamura T. Artificial nerve tubes and their application for repair of peripheral nerve injury: an update of current concepts. *Injury* 2008;39 Suppl 4:29-39.
2. Thorsén F, Rosberg HE, Steen Carlsson K, Dahlin LB. Digital nerve injuries: epidemiology, results, costs, and impact on daily life. *J Plast Surg Hand Surg* 2012;46:184-90.
3. Jardin E, Huard S, Chastel R, Uhring J, Obert L. Large diameter nerve conduits use in the upper limb: report of four cases and literature review. *Chir Main* 2011;30:393-9.
4. Riccio M, Marchesini A, Pugliese P, De Francesco F. Nerve repair and regeneration: Biological tubulization limits and future perspectives. *J Cell Physiol* 2019;234:3362-75.
5. Den Dunnen WF, Van der Lei B, Schakenraad JM, Blaauw EH, Stokroos I, Pennings AJ, et al. Long-term evaluation of nerve regeneration in a biodegradable nerve guide. *Microsurgery* 1993;14:508-15.
6. Boyd KU, Nimigan AS, Mackinnon SE. Nerve reconstruction in the hand and upper extremity. *Clin Plast Surg* 2011;38:643-60.
7. Yüksel F, Kışlaoğlu E, Durak N, Uçar C, Karacaoğlu E. Prevention of painful neuromas by epineural ligatures, flaps and grafts. *Br J Plast Surg* 1997;50:182-5.
8. Silini AR, Cargnoni A, Magatti M, Pianta S, Parolini O. The long path of human placenta, and its derivatives, in regenerative medicine. *Front Bioeng Biotechnol* 2015;3:162.
9. Riccio M, Pangrazi PP, Parodi PC, Vaienti L, Marchesini A, Neuendorf AD, et al. The amnion muscle combined graft (AMCG) conduits: a new alternative in the repair of wide substance loss of peripheral nerves. *Microsurgery* 2014;34:616-22.
10. Niknejad H, Peirovi H, Jorjani M, Ahmadiani A, Ghanavi J, Seifalian AM. Properties of the amniotic membrane for potential use in tissue engineering. *Eur Cell Mat* 2008;15:88-99.
11. Dua HS, Gomes JA, King AJ, Maharajan VS. The amniotic membrane in ophthalmology. *Surv Ophthalmol* 2004;49:51-77.

12. Baradaran-Rafii A, Aghayan HR, Arjmand B, Javadi MA. Amniotic membrane transplantation. *Iran J Ophthalmic Res* 2007;2:58-75.
13. Koob TJ, Lim JJ, Zabek N, Masee M. Cytokines in single layer amnion allografts compared to multilayer amnion/chorion allografts for wound healing. *J Biomed Mater Res B Appl Biomater* 2015;103:1133-40.
14. Gindraux F, Romain L, Nicod L, de Billy B, Meyer C, Zwetyenga N, et al. Human amniotic membrane: clinical uses, patents and marketed products. *Recent Patents on Regenerative Medicine* 2013;3:193-214.
15. Jirsova K, Jones GLA. Amniotic membrane in ophthalmology: properties, preparation, storage and indications for grafting-a review. *Cell Tissue Bank* 2017;18:193-204.
16. Laurent R, Nallet A, Obert L, Nicod L, Gindraux F. Storage and qualification of viable intact human amniotic graft and technology transfer to a tissue bank. *Cell Tissue Bank* 2014;15:267-75.
17. Ozcan G, Shenaq S, Spira M. Vascularized nerve tube: an experimental alternative for vascularized nerve grafts over short gaps. *J Reconstr Microsurg* 1993;9:405-13.
18. Mohammad J, Shenaq J, Rabinovsky E, Shenaq S. Modulation of peripheral nerve regeneration: a tissue-engineering approach. The role of amnion tube nerve conduit across a 1-centimeter nerve gap. *Plast Reconstr Surg* 2000;105:660-6.
19. Mligiliche N, Endo K, Okamoto K, Fujimoto E, Ide C. Extracellular matrix of human amnion manufactured into tubes as conduits for peripheral nerve regeneration. *J Biomed Mater Res* 2002;63:591-600.
20. Duan XL, Xu YZ, Zeng ZC. [Bridging rat sciatic nerve defects with the composite nerve-muscle autografts wrapped with human amnion matrix membrane]. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 2004;29:279-83.
21. Ozgenel GY, Fílíz G. Combined application of human amniotic membrane wrapping and hyaluronic acid injection in epineurectomized rat sciatic nerve. *J Reconstr Microsurg* 2004;20:153-7.

22. Henry FP, Goyal NA, David WS, Wes D, Bujold KE, Randolph MA, et al. Improving electrophysiologic and histologic outcomes by photochemically sealing amnion to the peripheral nerve repair site. *Surgery* 2009;145:313-21.
23. O'Neill AC, Randolph MA, Bujold KE, Kochevar IE, Redmond RW, Winograd JM. Preparation and integration of human amnion nerve conduits using a light-activated technique. *Plast Reconstr Surg* 2009;124:428-37.
24. Kim SS, Sohn SK, Lee KY, Lee MJ, Roh MS, Kim CH. Use of human amniotic membrane wrap in reducing perineural adhesions in a rabbit model of ulnar nerve neuroorrhaphy. *J Hand Surg Eur* 2010;35:214-9.
25. Meng H, Li M, You F, Du J, Luo Z. Assessment of processed human amniotic membrane as a protective barrier in rat model of sciatic nerve injury. *Neurosci Lett* 2011;496:48-53.
26. Fesli A, Sari A, Yilmaz N, Comelekoglu U, Tasdelen B. Enhancement of nerve healing with the combined use of amniotic membrane and granulocyte-colony-stimulating factor. *J Plast Reconstr Aesthet Surg JPRAS* 2014;67:837-43.
27. Fairbairn NG, Ng-Glazier J, Meppelink AM, Randolph MA, Valerio IL, Fleming ME, et al. Light-activated sealing of nerve graft coaptation sites improves outcome following large gap peripheral nerve injury. *Plast Reconstr Surg* 2015;136(4):739-50.
28. Sadraie SH, Parivar K, Arabi F, Moattari M, Kaka G, Mansouri K. Study of transected sciatic nerve repair by amniotic membrane with betamethasone in adult albino wistar rats. *Arch Iran Med* 2016;19(9):612-7.
29. Lemke A, Ferguson J, Gross K, Penzenstadler C, Bradl M, Mayer RL, et al. Transplantation of human amnion prevents recurring adhesions and ameliorates fibrosis in a rat model of sciatic nerve scarring. *Acta Biomater* 2018;66:335-49.
30. Hasturk AE, Yilmaz ER, Hayirli N, Kayalar AE, Akyildiz S, Gökce EC, et al. Stereologic and ultrastructural comparison of human and rat amniotic membrane wrapping for rat sciatic nerve repair. *J Clin Neurosci* 2018;57:157-61.

31. Marchesini A, Raimondo S, Zingaretti N, Riccio V, Battiston B, Provinciali M, et al. The amnion muscle combined graft (AMCG) conduits in nerves repair: an anatomical and experimental study on a rat model. *J Mater Sci Mater Med* 2018;2:120.
32. Gaspar MP, Abdelfattah HM, Welch IW, Vosbikian MM, Kane PM, Rekant MS. Recurrent cubital tunnel syndrome treated with revision neurolysis and amniotic membrane nerve wrapping. *J Shoulder Elbow Surg* 2016;25:2057-65.
33. Lopez-Valladares MJ, Teresa Rodriguez-Ares M, Tourino R, Gude F, Teresa Silva M, Couceiro J. Donor age and gestational age influence on growth factor levels in human amniotic membrane. *Acta Ophthalmol* 2010;88:e211-6.
34. Gicquel JJ, Dua HS, Brodie A, Mohammed I, Suleman H, Lazutina E, et al. Epidermal growth factor variations in amniotic membrane used for ex vivo tissue constructs. *Tissue Eng Part A* 2009;15:1919-27.
35. Banerjee A, Weidinger A, Hofer M, Steinborn R, Lindenmair A, Hennerbichler-Lugscheider S, et al. Different metabolic activity in placental and reflected regions of the human amniotic membrane. *Placenta* 2015;36:1329-32.
36. McQuilling JP, Vines JB, Kimmerling KA, Mowry KC. Proteomic comparison of amnion and chorion and evaluation of the effects of processing on placental membranes. *Wounds* 2017;29:E38-E42.
37. Koizumi NJ, Inatomi TJ, Sotozono CJ, Fullwood NJ, Quantock AJ, Kinoshita S. Growth factor mRNA and protein in preserved human amniotic membrane. *Curr Eye Res* 2000;20:173-7.
38. Wolbank S, Hildner F, Redl H, van Griensven M, Gabriel C, Hennerbichler S. Impact of human amniotic membrane preparation on release of angiogenic factors. *J Tissue Eng Regen Med* 2009;3:651-4.
39. Russo A, Bonci P, Bonci P. The effects of different preservation processes on the total protein and growth factor content in a new biological product developed from human amniotic membrane. *Cell Tissue Bank* 2012;13:353-61.

40. Paolin A, Trojan D, Leonardi A, Mellone S, Volpe A, Orlandi A, et al. Cytokine expression and ultrastructural alterations in fresh-frozen, freeze-dried and gamma-irradiated human amniotic membranes. *Cell Tissue Bank* 2016;17:399-406.
41. Koob TJ, Lim JJ, Masee M, Zabek N, Rennert R, Gurtner G, et al. Angiogenic properties of dehydrated human amnion/chorion allografts: therapeutic potential for soft tissue repair and regeneration. *Vasc Cell* 2014;6:10.
42. Hao Y, Ma DH, Hwang DG, Kim WS, Zhang F. Identification of antiangiogenic and antiinflammatory proteins in human amniotic membrane. *Cornea* 2000;19:348-52.
43. Shao C, Sima J, Zhang SX, Jin J, Reinach P, Wang Z, et al. Suppression of corneal neovascularization by PEDF release from human amniotic membranes. *Invest Ophthalmol Vis Sci* 2004;45:1758-62.
44. Gindraux F, Rondot T, de Billy B, Zwetyenga N, Fricain JC, Pagnon A, et al. Similarities between induced membrane and amniotic membrane: Novelty for bone repair. *Placenta* 2017;59:116-23.
45. Tao H, Fan H. Implantation of amniotic membrane to reduce postlaminectomy epidural adhesions. *Eur Spine J* 2009;18:1202-12.
46. Fénelon M, Catros S, Fricain JC. What is the benefit of using amniotic membrane in oral surgery? A comprehensive review of clinical studies. *Clin Oral Investig* 2018;22:1881-91.
47. Tseng SC, Espana EM, Kawakita T, Di Pascuale MA, Li W, He H, et al. How does amniotic membrane work? *Ocul Surf* 2004;2:177-87.
48. Silini A, Parolini O, Huppertz B, Lang I. Soluble factors of amnion-derived cells in treatment of inflammatory and fibrotic pathologies. *Curr Stem Cell Res Ther* 2013;8:6-14.
49. Kim JS, Kim JC, Na BK, Jeong JM, Song CY. Amniotic membrane patching promotes healing and inhibits proteinase activity on wound healing following acute corneal alkali burn. *Exp Eye Res* 2000;70:329-37.
50. Rossi D, Pianta S, Magatti M, Sedlmayr P, Parolini O. Characterization of the conditioned medium from amniotic membrane cells: prostaglandins as key effectors of its immunomodulatory activity. *PLoS One* 2012;7:e46956.

51. Pianta S, Bonassi Signoroni P, Muradore I, Rodrigues MF, Rossi D, Silini A, et al. Amniotic membrane mesenchymal cells-derived factors skew T cell polarization toward treg and downregulate Th1 and Th17 cells subsets. *Stem Cell Rev* 2014;11:394-407.
52. Tseng SC, Li DQ, Ma X. Suppression of transforming growth factor-beta isoforms, TGF-beta receptor type II, and myofibroblast differentiation in cultured human corneal and limbal fibroblasts by amniotic membrane matrix. *J Cell Physiol* 1999;179:325-35.
53. Yetkin G, Uludag M, Citgez B, Karakoc S, Polat N, Kabukcuoglu F. Prevention of peritoneal adhesions by intraperitoneal administration of vitamin E and human amniotic membrane. *Int J Surg* 2009;7:561-5.
54. Arora M, Jaroudi KA, Hamilton CJ, Dayel F. Controlled comparison of interceed and amniotic membrane graft in the prevention of postoperative adhesions in the rabbit uterine horn model. *Eur J Obstet Gynecol Reprod Biol* 1994;55:179-82.
55. Luo J, Yang Z, Li X. [Effect of human acellular amnion membrane on tendon adhesion in rat]. *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi* 2004;18:431-4.
56. Adinolfi M, Akle CA, McColl I, Fensom AH, Tansley L, Connolly P, et al. Expression of HLA antigens, beta 2-microglobulin and enzymes by human amniotic epithelial cells. *Nature* 1982;295:325-7.

Table titles

Table 1. Summary of the properties of the amniotic membrane in its various formats [8,10-15,35,37-39]

Table 2. Use of human amniotic membrane in animal models for repairing nerve defects or nerve transection.

Table 3. Clinical trials (completed or on-going) reporting use of the human amniotic membrane in rheumatology, orthopedic and trauma conditions. *Source:*

<https://clinicaltrials.gov/ct2/search>

Table 4. Summary of the marketed amniotic membrane products, including their composition and features.

Table 1. Summary of the properties of the amniotic membrane in its various configurations
[8,10-15,35,37-39]

Features	Fresh amniotic membrane	Cryopreserved amniotic membrane	Lyophilized or dehydrated amniotic membrane (+ gamma irradiation)
Need for cold chain after collection and during handling	Yes	Yes	No
Cell viability	+++ (controversial)	++ (controversial)	None
Growth factors	Amnion: ++ Chorion: +++	Amnion: + Chorion: ++	Amnion: + Chorion: ++
Possibility of room temperature storage (e.g. in operating room)	No	No	Yes
Ease of handling and cutting tissue	No	No	Yes
Use in two-layer configuration (amnion + chorion)	No	No	Yes
Additional gamma irradiation	No	Not usually	Yes

+: moderate; ++: high; +++: very high

Table 2. Use of human amniotic membrane in animal models for repairing nerve defects or nerve transection.

Reference	Animal species (strain) / number	Configuration of human amniotic membrane	Nerve insult / type of nerve	Main findings
Ozcan et al. [17]	Rats (<i>Lewis</i>) 25	Manufactured multilayer amnion tube	1-cm defect Femoral nerve	Vascularized amnion tube achieved nerve regeneration similar to vascularized nerve graft and better than non-vascularized amniotic membrane tube or non-vascularized nerve graft
Mohammad et al. [18]	Rats (<i>Sprague Dawley</i>) 66	De-epithelialized, dehydrated and gamma-sterilized amnion tube	1-cm defect Sciatic nerve	Advantage of amnion tube: - Better histological results than with nerve autograft or silicone tube - Better function in the early stages
Mligiliche et al. [19]	Rats (<i>Wistar</i>) 24	De-epithelialized and dehydrated amnion tubes of different diameters	0.5-cm defect Sciatic nerve	Advantage of 1–2 mm diameter amnion tube: - Same number of myelinated axons as healthy nerve, but smaller in diameter - Gastrocnemius muscle is reinnervated after 9 months
Duan et al. [20]	Rats (<i>Wistar</i>) 54	Information not available	1-cm defect Sciatic nerve	Advantage of composite nerve-muscle autografts wrapped with human amnion matrix membrane: - More neurofilaments in nerve root; neurofilaments are denser and more regular - More axons; axons are larger in diameter and thicker
Ozgenel et al. [21]	Rats (<i>Sprague-Dawley</i>) 72	Fresh de-epithelialized amnion	3-cm defect Sciatic nerve	Nerve wrapping with amnion reduced fibrosis and perineural adhesions This action was bolstered by hyaluronic acid injection
Henry et al. [22]	Rabbits (<i>New Zealand</i>) 24	Cryopreserved amnion	Transection 5 cm from distal end Peroneal nerve	Amnion fixation on nerve using photochemical tissue bonding better than simple suture
O'Neill et al. [23]	Rats (<i>Sprague-Dawley</i>) 24	Cryopreserved amnion	1-cm defect Sciatic nerve	Advantage of photochemical tissue bonding of the amnion over epineural fixation: - Better functional outcomes (SFI) at 8 and 12 weeks - Greater muscle mass in re-innervated gastrocnemius

				- Larger number of fibers, larger diameter and thicker myelin layer
Kim et al. [24]	Rabbits (<i>New Zealand</i>) 10	Cryopreserved amnion	Transection Ulnar nerve	Nerve wrapping with amnion significantly reduced fibrosis and perineural adhesions compared to contralateral control without wrapping
Meng et al. [25]	Rats (<i>Sprague Dawley</i>) 36	Cryopreserved amnion	1 cm defect Sciatic nerve	Advantage of nerve wrapping with amnion autograft: - Improved functional recovery in the early phase after nerve lesion - Fewer adhesions and less fibrosis
Fesli et al. [26]	Rats (<i>Wistar</i>) 70	Fresh amnion	Transection Sciatic nerve	Nerve wrapping with amnion reduced perineural fibrosis, improved clinical response and improved electrophysiology data Intraperitoneal GCSF injection potentiated these results
Fairbairn et al. [27]	Rats (<i>Lewis</i>) 100	Cryopreserved de-epithelialized amnion	1.5 cm defect Sciatic nerve	Advantage of nerve wrapping with amnion secured by photochemical tissue bonding versus simple suture or biological glue: - Less perineural fibrosis - Better SFI results - Increased muscle mass in reinnervated gastrocnemius - Larger fiber and axon diameter - Thicker myelin layer
Sadraie et al. [28]	Rats (<i>Wistar</i>) 42	Fresh de-epithelialized amnion	Transection Sciatic nerve	Advantage of nerve wrapping with amnion combined with betamethasone injection (0.2 mL betamethasone (4 mg/mL)): - Better SFI at 4 weeks - Improved paw withdrawal reflex (seconds) - Reduced conduction latency (ms) - Stronger conduction amplitude (mV) - More large-diameter (> 6 μ m) fibers

Lemke et al. [29]	Rats (<i>Sprague Dawley</i>), 54	Fresh amnion with epithelial side placed against nerve	Chemical induction of fibrosis Sciatic nerve	Significant advantage of amnion nerve conduit: - Better nerve healing - Reduced perineural fibrosis and adhesions - Accelerated nerve regeneration: less demyelination, intraneural structure regenerated more quickly, better nerve function (SFI) - Regulation of immune response
Hasturk et al. [30]	Rats (<i>Wistar</i>) 25	Cryopreserved human and rat amnion	Transection Sciatic nerve	Advantage of nerve wrapping with amnion allograft over xenograft
Marchesini et al. [31]	Rats (<i>Wistar</i>) 14	Dehydrated amnion filled with skeletal muscle harvested from neighboring tissue	1.5-cm defect Median nerve of forelimb	Experimental support for clinical proof-of-concept [9]





SFI: Sciatic Functional Index; GCSF: granulocyte colony stimulating factor

Table 3. Clinical trials (completed or on-going) reporting use of the human amniotic membrane in rheumatology, orthopedic and trauma conditions. Source:

<https://clinicaltrials.gov/ct2/search>


Scope of application	Medical condition	Study number (NCT)	Nb. of studies
	Shoulder osteoarthritis	NCT03770546	1
	Knee osteoarthritis	NCT02318511, NCT02768155, NCT02767492, NCT03074526, NCT03337243, NCT03408145, NCT03441607, NCT03485157	8
	Hip osteoarthritis	NCT03063099	1
Musculoskeletal system:	Rotator cuff tear	NCT03379324	1
Medical treatment	Osteochondral lesion	NCT02837484, NCT03036878	2
	Low back pain	NCT02932020, NCT03644251	2
	Tendinitis: Achilles, plantar fascia	NCT01357187, NCT01659827, NCT01996111, NCT02427191, NCT02982226, NCT03414255, NCT03414268	7
	Tendinitis of epicondylar muscles, musculoskeletal problems, trigger finger	NCT01921569, NCT03390920, NCT03583151	3
	Hand tendon surgery	NCT02361814, NCT03013582	2
	Ankle tendon surgery: Achilles tendon, fibular tendons	NCT01708187, NCT02719288	2
Musculoskeletal system:	Knee surgery: arthroplasty	NCT02088567	1
Surgical treatment	Knee surgery: anterior cruciate ligament reconstruction	NCT03294759, NCT03294759	2
	Foot surgery: hallux rigidus	NCT01825356	1
	Spine surgery	NCT02023372, NCT02070484, NCT02300909, NCT02381067, NCT02808234, NCT03113786	6
	Nonunion	NCT03031509	1
		Total studies	40

Table 4. Summary of the commercialized amniotic membrane products, including their composition and features.

Company	Product name	Size	Format / Process	Indications	Similar products
Alliqua Biomedical	Biovance® 	1 x 2 to 6 x 6 cm	Decellularized, dehydrated amnion	Acute and chronic wounds (ulcers, burns...) including wounds with exposed tendon, muscle, bone or other vital structures	
AediCell®	Plurivest® and Demavest® 	1.5 x 1 ; 2 x 3 cm	Freeze-dried and gamma-irradiated placental tissue matrix (placenta disc, amnion/chorion and umbilical cord)	Wounds, ulcers, Burns, surgical needs	
AFCell - Liventa Bioscience	AmnioClear™ 	2 x 2 to 4 x 6 cm	Amnion/Chorion processed aseptically and sterilized	Soft tissue covering, Diabetic foot ulcers	
Alon Source Group	ASGBarrier™ 	2 x 3 to 4 x 8 cm	Wet or dry amnion	Wound healing	ASGFluid™ and AmnioELITE™: amnion and amniotic fluid-derived cells

Alamo Tissue Service

AmnioBioGraft (+)



1 - 75 sq cm
0.8 to 1.4 cm
2 x 3 to 4 x 8 cm


Amniotic membrane cleaned and sterilized by PASCO2® technology
AmnioBioGraft: amnion
AmnioBioGraft +: amnion and chorion

Wound healing,
Ophthalmology

AmnioBioGraft Cord: derived from the umbilical cord

Allosource

Allowrap®




2 x 2 to 4 x 8 cm

Amnion/Chorion cleaned by AlloTrue™ technology
Moist (AlloWrap DS) or dry (AlloWrap Dry) configurations

Biological barrier following surgical repair

Alphatec Spine

AmnioShield




No information

Dehydrated amnion disinfected by PurionSM process

Chronic wounds
Scars

Amnio Medical

NEOX® 100



NEOX® CORD 1K/RT



2 x 2 to 7 x 7 cm
2 x 1 to 8 x 3 cm

NEOX® 100: Cryopreserved amnion
NEOX® CORD 1K: Cryopreserved umbilical cord + amnion
NEOX® CORD RT: room temperature storage
NEOX® FLO: Lyophilized umbilical cord-Based particulate
CRYOTEK® preservation process

Wound healing,
Diabetic foot and chronic ulcers

CLARIX®: Reconstruction, tendon repair, nerve decompression, trauma, and arthroplasty

RESPINA®: Lumbar interbody fusion

PalenGen® membrane & hydromembrane

Amnio
Technology
LLC



1 x 1 to 8 x 8 cm

Air dried amnion using Pinnacle
technologies

Dry or Wet forms

Wound covering and
support for native tissues

XPlus Membrane & XPlus
Hydromembrane: wet form providing
greater tensile strength, shape
manipulation, and slower resorption in
vivo

PalinGen® Flow: amniotic tissue
allografts containing cellular
components cytokines, growth factors
and extracellular matrix proteins

Xwrap™

Applied
Biologics



2 x 2 to 4 x 8 cm


Dehydrated amnion
Dry, ECM, Hydro PLUS forms

Wound healing

FloGraft™:
Cryopreserved injectable amnion

BioDfence® G3

BioDlogics
(Integra
LifeSciences)



2 x 3 to 10 x 10 cm

Dehydrated, sterilized amnion/chorion
stored at room temperature

Wound healing
Ophthalmology
Nerve, tendon surgery

BioDDryFlex®, BioDOptix®:
Dehydrated amnion using patent-
pending DryFlex® processing
technology

BioDFence® Sentry™: Dehydrated
amnion, spongy (intermediate) layer
and the chorion

BioDFactor®: Flowable tissue allograft
derived from morselized amniotic
tissue and components of the amniotic
fluid

PROKERA®

Bio-tissue



No information


Cryopreserved amnion using CryoTek®
processing method, held under tension
inside a thermoplastic circle

Ophthalmology

PROKERA SLIM, PROKERA, and
PROKERA PLUS

AmnioGraft®
AmnioGuard®

Amnio-care™



Biocover
Laboratories

No information

Dry amnion

Ophthalmology

Acelagraft™



Celgene
Cellular
Therapeutics

No information

Dehydrated and decellularized amniotic
membrane, stored at room temperature

Wound healing
Chronic ulcers

AmnioExcel



Derma
Sciences
(Integra)

Disks: 1.2 to 2.4 cm
1.5 x 1.5 to 10 x 10
cm

Dehydrated using the proprietary
DryFlex™ process

Chronic wounds

AMNIOMATRIX®:
Cryopreserved suspension allograft
derived from the amniotic membrane
and components of the amniotic fluid
using the patented CryoPrime™

FlowerPatch™



Flower
Orthopedics™

2 x 2 to 4 x 8 cm

Dehydrated dual layer amnion

Wound healing

FLOWERFLO™: Flowable amnion
tissue allograft stored at ambient
temperatures

AmbioDisk™



IOP
Ophthalmics

Disk: 15 mm
1.5 x 2 to 4 x 4 cm

Cleaned, dehydrated and sterilized
amnion
Visual identification of amniotic
membrane orientation

Ocular surface
reconstruction

AmbioDry2™: Dry state

Keera

AMX: Amniotic Membrane eXtract

Soluble powder

Lyophilized extract of the fresh amniotic
membrane

Ophthalmology

AmnioFix®



MiMedx Group


Disk: 16 mm
Sheets: 2 x 3 to 4 x
6 cm
Wraps: 2 x 2 to 4 x
6 cm

Dehydrated amnion/chorion
membrane using PurionSM process,
terminally sterilized

Wound healing
Skin Substitute product,
Diabetic Foot and venous
leg ulcers

EpiFix®, Epiburn®
AmnioFill®: minimally manipulated,
non-viable cellular tissue matrix
allograft
AmnioCord®, Epicord®: minimally
manipulated, dehydrated, non-viable
cellular umbilical cord

AmnioBand®



AmnioBand®
Particulate



Musculoskeletal
Transplant
Foundation
(MTF)
biologics

Disks: 10 to 18 mm
2 x 2 to 7 x 7cm
Particulate: 40 to 80
mg

Dehydrated sterilized amnion/chorion

Acute or chronic wound
covering

AmnioBand® Viable Membrane:
cryopreserved allograft

Amniotic membrane small or large

NHS Blood and Transplant




2 x 2 or 3 x 3 cm

Cryopreserved amnion at a minimum of -40°C in 50% glycerol / 50% Hanks solution

Ophthalmic surgery, primarily ophthalmic surface dressing

Affinity®

Organogenesis



No information

The only fresh hypothermically stored amnion with viable cells (including stromal cells, fibroblasts, and epithelial cells), growth factors and cytokines; processed by AlloFresh® method


Wound healing
Soft tissue repairs

Nushield®: Dehydrated sterilized amnion/chorion processed by BioLoc™ method (tendons and nerves)

NuCel®: cryopreserved amniotic membrane with cells from amniotic fluid; Powder in suspension (orthopedic procedures)

Versashield®

Orthofix (MTF)



2 x 2 to 4 x 6 cm
2 x 3 ; 3 x 8 cm

Dehydrated sterilized amnion/chorion

General Surgery, Corneal Plastic Surgery, Burn and Wound Care, Sports Foot and Ankle Procedures, Spine and Dura repair

Stavux®

Osiris Therapeutics







2 x 4 to 3 x 6 cm

Cryopreserved human placental tissue composed of umbilical amnion and Wharton's jelly


Tendon Repair
Achilles Tendon Rupture
Bunionectomy
Hallux Rigidus Correction
Foot Amputations
Fibromatosis
Arthrodesis

Graftix®: Cryopreserved amnion/chorion (wound healing, Bone and tendon)

GrafixPLPime®: Lyophilized amnion/chorion

Seedbiotech	<p>Aril®</p> 	<p>Disks: 5 to 15 mm Ellipse: 1 x 2 to 8 x 10 cm</p>	<p>Decellularized and gamma irradiated amnion</p>	<p>Orthopedic, surgical, ophthalmology, spinal and wound covering applications</p>	
SurgiLogix	<p>SX Barrier™</p> 	<p>2 x 3 to 4 x 8 cm</p>	<p>Wet or dry amnion</p>	<p>Wound healing Tissue repair</p>	<p>AmnioELITE: amniotic fluid allograft comprised of amnion SXFluid™: amniotic tissue allograft comprised of ground amnion and amniotic fluid-derived components</p>
Snoasis Medical	<p>BioXclude®</p> 	<p>0.8 x 0.8 to 2 x 3 cm</p>	<p>Washed, dehydrated and sterilized amnion/chorion using PurionSM process</p>	<p>Dental and maxillofacial surgery Ophthalmology</p>	
SKY orthobiologics	<p>ActiveBarrier® 45, 200, 2000</p> 	<p>Various thicknesses: ActiveBarrier® 45: 45 µm ActiveBarrier® 200: 200 µm ActiveBarrier® 2000: 2000 µm 2 x 2 to 4 x 8 cm</p>	<p>ActiveBarrier® 45: Amnion ActiveBarrier® 200: Chorion ActiveBarrier® 2000: entire placental membrane ActiveBarrier® 45 & 200: Dehydrated, HydraTek® process ActiveBarrier® 2000: Fresh frozen</p>	<p>Chronic wounds, nerve conduit, nerve defect, tendon sleeve</p>	<p>OculoMatrix® & VisiDisc® Thin / WoundEx®45: Dehydrated amnion OculoMatrix® & VisiDisc® Thick / WoundEx®200: Dehydrated chorion ReVive™ Membrane: Dehydrated placental membrane</p>

Vivex
Biomedical



Cygnus®

Various sizes
1 x 1 to 12 x 12 cm

Dehydrated allograft

Cygnus® Solo (amnion)
Cygnus® Dual (dual layer amnion)
Cygnus® Matrix (amnion/chorion)
Cygnus® Max (umbilical cord)

Spine & Neurosurgery

Foot & Ankle
Wound Care
Burn Care
Dermatology
Ophthalmology
Oral Surgery

AlloGen® (frozen or liquid): all-natural
liquid matrix allograft derived from
amniotic fluid