



HAL
open science

Variability of shoulder girdle proprioception in 44 healthy volunteers

Camille Echali er, Julien Uhring, Julie Ritter, Pierre-Bastien Rey, Emmanuelle Jardin, S everin Rochet, Laurent Obert, Fran ois Loisel

► To cite this version:

Camille Echali er, Julien Uhring, Julie Ritter, Pierre-Bastien Rey, Emmanuelle Jardin, et al.. Variability of shoulder girdle proprioception in 44 healthy volunteers. *Orthopaedics & Traumatology: Surgery & Research*, 2019, 105, pp.825 - 829. <10.1016/j.otsr.2019.01.019>. <hal-03488080>

HAL Id: hal-03488080

<https://hal.science/hal-03488080v1>

Submitted on 20 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

Original article

Variability of Shoulder Girdle Proprioception in 44 Healthy Volunteers

Camille **Echalier**^a, Julien **Uhring**^b, Julie **Ritter**^a, Pierre-Bastien **Rey**^c, Emmanuelle **Jardin**^d, Séverin **Rochet**^a, Laurent **Obert**^a, François **Loisel**^a

- a. Service d'orthopédie, de traumatologie, de chirurgie plastique, reconstructrice et assistance main, pôle innovation et technique chirurgicale, EA 4268 innovation, Nano médecine, université de Franche Comté, IFR 133 Inserm, CHU Jean Minjoz, boulevard Fleming, 25000 Besançon, France
- b. Hôpital-Clinique Claude Bernard, 97, rue Claude-Bernard, BP 45050, 57072 Metz, France
- c. Hôpital Privé de la Miotte, 15 avenue de la Miotte, CS 30109, 90002 Belfort, France
- d. Clinique du Diaconat-Roosevelt, 14 boulevard Roosevelt, 68067 Mulhouse, France

Corresponding author:

Camille ECHALIER, CHRU Minjoz, 3 bd Fleming, 25000 Besançon, FRANCE

Tel.: +33 688 558 810

E-mail: camille_echalier@live.fr

ABSTRACT

Background: Improved knowledge of normal shoulder girdle proprioception should benefit the treatment and rehabilitation of shoulder disorders. Whereas many of the available methods for assessing joint position sense (JPS) are costly and complex, Balke et al. have described a simple test. The primary objective of this study was to use this test to identify factors that influence shoulder-girdle JPS evaluation in healthy individuals. The secondary objective was to determine reference values based on the values obtained and on the factors associated with their variability.

Hypothesis: Age and dominant limb influence the results of shoulder girdle JPS evaluation, creating a need for reference values that take these factors in account.

Patients and methods: A single-centre prospective study of healthy volunteers was performed between September 2012 and January 2013. In each volunteer, shoulder repositioning accuracy was assessed bilaterally as described by Balke et al. A line was drawn on the floor parallel to and 1 metre away from a wall. A target on the wall showed marks corresponding to three angles of arm elevation: 55°, 90°, and 125°. The volunteer was asked to raise the arm to the three positions, in abduction and in forward flexion, while memorising the joint positions, then to replicate the same positions with the eyes closed. The absolute differences between the replicated positions and the reference positions, designated angle deviations, were measured in degrees. Age, sex, and dominant upper limb were recorded.

Results: The 88 shoulders of 44 healthy volunteers were studied. No significant difference was found between males and females ($p>0.05$). The only significant difference between sides was better replication by the dominant arm of flexion at 55° ($p=0.03$). By univariate analysis, age was the only factor significantly associated with repositioning errors ($p=0.003$); neither dominant limb nor sex were significant ($p=0.29$ and $p=0.59$, respectively). In flexion, the angle deviation increased significantly with movement amplitude. The measurements

were used to create a chart of expected angle deviations in healthy individuals according to age and to plane and amplitude of movement.

Discussion: The test described by Balke et al. is simple and feasible in everyday practice. Age, dominance, and plane of movement should be considered when evaluating abnormal shoulders. Further studies in larger numbers of individuals are needed to better define normal angle deviations related to these factors.

Level of evidence: IV

Key words: Shoulder. Proprioception. Proprioception test. Joint position sense. Instability.

Introduction

The shoulder girdle is a mobile biomechanical complex whose anatomical configuration results in considerable instability. Dynamic stability is provided by various passive and active restraints. Several studies have highlighted the key role of proprioceptive information supplied by specific mechanoreceptors (Golgi, Pacini, and Ruffini) that reside within the joints, ligaments, tendons, and skin [1-5]. These mechanoreceptors may be damaged in patients with conditions such as rotator cuff tears or gleno-humeral instability [6, 7].

Shoulder proprioception has been a focus of growing attention in sports medicine in recent years. Impaired proprioception has been reported to result in musculo-skeletal lesions, whose recurrence is more likely if the impairments persist [8]. To determine whether surgical procedures and/or rehabilitation programmes are effective in improving proprioception, accurate methods for measuring proprioception are needed.

Proprioception can be divided into three components, namely, joint position sense (JPS), kinaesthesia, and sensitivity to resistance. The tests currently being developed focus on JPS and often require specific measurement tools, many of which are both costly and challenging to use [9, 10]. JPS testing during active movements provides a more accurate picture, as the muscle contraction generates afferent input, contributes to stability, and replicates events that occur during the activities of daily living [11].

Balke et al. [12] have developed a simple and inexpensive test for measuring JPS at the shoulder. This test requires no specific equipment and can be performed in everyday clinical practice. The primary objective of this study was to use this test to identify factors that influence shoulder girdle JPS in healthy individuals. The secondary objective was to determine reference values based on the measurements obtained and factors associated with

variability. The working hypothesis was that age and dominant limb influence the results of shoulder girdle JPS evaluation, creating a need for reference values that take these factors in account.

Patients and Methods

A single-centre prospective study was performed over the 4-month period from September 2012 to January 2013 at the university hospital in Besançon, France. The study participants were healthy volunteers with no constitutional or acquired shoulder girdle abnormalities, normal active and passive range of motion at both shoulders, and no shoulder pain. All volunteers received oral and written information about the study. In each, the ability to replicate shoulder positions was assessed bilaterally.

All measurements were performed by the same examiner as described by Balke et al. [12] (Figure 1). The volunteer stood on a line drawn parallel to and 1 metre away from the wall. A target hung on the wall showed the points the volunteer had to aim for to achieve 55°, 90°, and 125° of arm elevation in forward flexion and in abduction. The point corresponding to 90° was at the centre of the target. The height of the target on the wall was adjustable so that the centre could be aligned with the gleno-humeral joint of the volunteer. To avoid stimulation of the skin, the volunteers did not wear clothes that covered the shoulders. The wrist was immobilised in a removable brace to avoid errors due to wrist movements. A laser pointer was affixed to the wrist on the side being tested. The volunteer was then asked to memorise the joint positions while raising the arm to each of the three levels (55°, 90°, and 125°), in forwards flexion and abduction. With the eyes covered, the volunteer then attempted to replicate each position, on three different occasions, and the degree of forward elevation or abduction was measured at each attempt. Thus, 18 measurements were obtained

for each shoulder and 36 measurements for each volunteer. The difference between each value obtained with the eyes covered and the theoretical value (55°, 90°, or 125°) was computed and designated by the term 'angle deviation'. For each volunteer, age, sex, and dominant limb were recorded.

Statistical analysis

Quantitative variables were described as mean±SD and qualitative variables as n (%). Quantitative variables were compared between groups by applying the Wilcoxon-Mann-Whitney test. Univariate linear regression was used to assess associations between repositioning errors and age, sex, and dominance. The statistical analyses were performed using R version 3.5.0 and RStudio version 1.1.453 (R Foundation for Statistical Computing, Vienna, Austria). Values of *p* lower than 0.05 were taken to indicate significant differences.

Results

Population

The 88 shoulders of 44 healthy volunteers were studied (Table 1). There were 23 (52.3%) males and 21 (47.7%) females) with a mean age of 31.5 years (range, 16-54 years). Of the 44 volunteers, 39 (88.6%) were right-handed and 5 (11.4%) were left-handed.

Statistical associations

No differences were found between males and females for any of the levels of forwards elevation or abduction (Figure 2). The only significant difference according to handedness occurred for forwards flexion at 55°, for which repositioning was significantly more accurate on the dominant side (mean angle deviation, 8° vs. 9° on the non-dominant

side, $p=0.03$) (Figure 3). By univariate analysis, only age was associated with repositioning errors ($p=0.003$); no significant association was found for dominant side or sex ($p=0.29$ and $p=0.59$, respectively).

In abduction (Figure 4), mean angle deviations differed significantly between 125° and 90° and between 125° and 55° . No significant differences in repositioning errors were found between 55° and 90° of abduction. In forwards flexion, mean angle deviations showed significant differences across the three levels.

Reference values

The values obtained were used to determine the expected angle deviations in healthy individuals according to age, plane of motion, and range of motion (Table 2).

Discussion

In this study, age significantly influenced the results of shoulder JPS testing using the method described by Balke et al. Expected repositioning errors in healthy individuals were determined.

To date, no consensus exists about the best method for evaluating shoulder proprioception. Many different methods have been described [10]. The link established in several studies between instability and SLAP tears or Bankart's tear indicate an association between impaired proprioception and damage to the joint capsule [13-15]. Gleno-humeral proprioception is impaired in patients with chronic shoulder instability but improves gradually after stabilising surgery [16, 17]. However, in one study, surgery after a single episode of anterior gleno-humeral dislocation failed to improve proprioception [18]. Studies

of tests to assess JPS were done in small numbers of patients, all of whom were young, and failed to consider inter- and intra-individual variations.

The testing method devised by Balke et al. [12] is simple and inexpensive, requires no specific equipment, and exhibits satisfactory intra-observer repeatability [10]. We used it to conduct a larger study of gleno-humeral JPS in healthy volunteers and to establish reference values for purposes of comparison when assessing abnormal shoulders (glenoid impingement, instability, post-operative recovery) and the effects of treatment.

Age-related impairments in proprioception may be partly ascribable to muscle fatigability [19]. An increase in muscle fatigue with advancing age seems plausible. Studies of individuals with similar physical capabilities divided into different age groups would be needed. JPS testing during passive movements would eliminate any bias due to muscle fatigue, as the muscles would then serve merely as proprioception effectors and not as movement effectors. However, this approach does not seem satisfactory, as proprioception is less effective during passive than active movements, as muscles supply proprioceptive information when they are contracted [11, 20].

Sex was not significantly associated with JPS, as previously reported [1, 12]. Consequently, data from males and females can be pooled.

Proprioception seemed more effective on the dominant side, although the difference was statistically significant only for forwards flexion at 55°. Aydin et al. [16] contended that the dominant limb performed better due to its more frequent use providing greater opportunities for learning. Furthermore, the repeated use of forwards flexion at 55° to carry out activities of daily living may improve performance as it can be likened to intensive training. Nevertheless, in our study, the only significant difference between the dominant and non-dominant sides was a 1° difference in forwards flexion at 55°. Such a small difference is not clinically significant. A study in a larger number of individuals would

improve our knowledge of the performance of the dominant limb in this sector of the motion range.

The ability to achieve accurate repositioning in forwards flexion decreased as the amplitude of the movement increased. In abduction, a non-significant trend towards diminished accuracy was seen between 55° and 90°. In contrast, in earlier studies performance was best in the plane of the gleno-humeral joint (90°) [1, 12]. This finding was ascribed to the greater tension on muscle and tendons as amplitude increased, translating into greater recruitment of tendon mechanoreceptors, which are the main proprioceptive organs [5]. In our study, repositioning errors increased (i.e., accuracy decreased) with increasing amplitude. Lönn et al. [21] reported that repositioning accuracy improved when the starting point was an intermediate position, that is, when the movement arc was shorter. In addition, muscle fatigability increases with movement amplitude and impairs JPS, thereby decreasing repositioning accuracy [22, 23].

Dvir et al. [24] underlined the importance of assessing the clinical significance of statistically significant differences between measured values. A statistically significant difference is not necessarily associated with a change that is clinically meaningful to the patient. Based on the values obtained in our study, we suggest that angle deviations may be clinically significant if greater than 15°.

Conclusion

The evaluation of shoulder proprioception raises both practical and analytical challenges. The test devised by Balke et al. is simple and inexpensive. JPS at the shoulder is associated with age but not with sex or dominance. Repositioning errors increase with the amplitude of forwards flexion. Further studies in larger numbers of individuals are required.

Given its simplicity, the Balke test should prove useful for monitoring proprioception after the treatment of shoulder conditions. Additional work is needed to improve our understanding of normal angle deviations in healthy individuals. Finally, it would be of interest to assess potential correlations linking functional shoulder scores to JPS in order to shed light on the functional impact of impaired proprioception.

Disclosure of interest

Laurent Obert has no conflicts of interest to declare in relation to this work but has financial ties to FX solutions, Zimmer, Medartis, Evolutis, Biotech Wright, Lilly, the Besançon university hospital, the Bourgogne Franche Comté university, Elsevier, and Springer.

None of the other authors has any conflicts of interest to declare.

Funding

None

Contributions of each author:

Echalier C: corrected the data and drafted the manuscript

Ritter J: corrected the data

Uhring J, Rey PB, Jardin E, and Rochet S: evaluated the results

Obert L and Loisel F: conceived the study and revised the manuscript

FIGURE LEGENDS

Figure 1: The individual stands 1 metre away from the wall, in front of a target, and raises the arm in abduction and forwards elevation, aiming at the three angles marked on the target (55°, 90° et 125°).

Figure 2: Mean angle deviations in males and females for the three positions tested, in abduction and forwards flexion. No significant difference was found.

Figure 3: Mean angle deviations for the dominant and non-dominant limbs in the three positions tested, in abduction and forwards flexion. Accuracy was significantly better for the dominant limb at 55° of forwards flexion.

Figure 4: Mean angle deviations for each level of forwards flexion and abduction: in forwards flexion, angle deviation increased with amplitude

Table 1. Main features of the 44 healthy volunteers

Age, years, mean\pmSD	31.5 \pm 10.4
Males, n	23
Right-handed, n	39

Table 2. Expected angle deviation according to age, plane of motion, and movement amplitude in healthy individuals

	Age <25 years	Age 25 to 35 years	Age >35 years
Abduction			
55°	10°	11°	11°
90°	11°	12°	12°
125°	16°	11°	16°
Flexion			
55°	6°	9°	10°
90°	9°	9°	13°
125°	11°	9°	17°

References

- [1] Jerosch J, Thorwesten L, Steinbeck J, Reer R. Proprioceptive function of the shoulder girdle in healthy volunteers. *Knee Surg Sports Traumatol Arthrosc* 1996;3(4):219-25.
- [2] Riemann BL, Lephart SM. The sensorimotor system, part I: the physiologic basis of functional joint stability. *J Athl Train* 2002;37(1):71-9.
- [3] Myers JB, Lephart SM. Sensorimotor deficits contributing to glenohumeral instability. *Clin Orthop Relat Res* 2002(400):98-104.
- [4] Myers JB, Wassinger CA, Lephart SM. Sensorimotor contribution to shoulder stability: effect of injury and rehabilitation. *Man Ther* 2006;11(3):197-201.
- [5] Myers JB, Lephart SM. The role of the sensorimotor system in the athletic shoulder. *J Athl Train* 2000;35(3):351-63.
- [6] Gumina S, Camerota F, Celletti C, Venditto T, Candela V. The effects of rotator cuff tear on shoulder proprioception. 2018.
- [7] Contemori S, Biscarini A, Botti FM, Busti D, Panichi R, Pettorossi VE. Sensorimotor Control of the Shoulder in Professional Volleyball Players With Isolated Infraspinatus Muscle Atrophy. *J Sport Rehabil* 2018;27(4):371-9.
- [8] Lephart SM, Warner JJ, Borsa PA, Fu FH. Proprioception of the shoulder joint in healthy, unstable, and surgically repaired shoulders. *J Shoulder Elbow Surg* 1994;3(6):371-80.
- [9] Contu S, Hussain A, Kager S, Budhota A, Deshmukh VA, Kuah CWK, et al. Proprioceptive assessment in clinical settings: Evaluation of joint position sense in upper limb post-stroke using a robotic manipulator. *PLoS One* 2017;12(11):e0183257.
- [10] Ager AL, Roy JS, Roos M, Belley AF, Cools A, Hebert LJ. Shoulder proprioception: How is it measured and is it reliable? A systematic review. *J Hand Ther* 2017;30(2):221-31.
- [11] Lin YL, Karduna A. Errors in Shoulder Joint Position Sense Mainly Come from the Glenohumeral Joint. *J Appl Biomech* 2017;33(1):32-8.
- [12] Balke M, Liem D, Dedy N, Thorwesten L, Balke M, Poetzl W, et al. The laser-pointer assisted angle reproduction test for evaluation of proprioceptive shoulder function in patients with instability. *Arch Orthop Trauma Surg* 2011;131(8):1077-84.
- [13] Hantes M, Raoulis V. Arthroscopic Findings in Anterior Shoulder Instability. *Open Orthop J* 2017;11:119-32.
- [14] Durban CM, Kim JK, Kim SH, Oh JH. Anterior Shoulder Instability with Concomitant Superior Labrum from Anterior to Posterior (SLAP) Lesion Compared to Anterior Instability without SLAP Lesion. *Clin Orthop Surg* 2016;8(2):168-74.
- [15] Clavert P. Glenoid labrum pathology. *Orthop Traumatol Surg Res* 2015;101(1 Suppl):S19-24.
- [16] Aydin T, Yildiz Y, Yanmis I, Yildiz C, Kalyon TA. Shoulder proprioception: a comparison between the shoulder joint in healthy and surgically repaired shoulders. *Arch Orthop Trauma Surg* 2001;121(7):422-5.
- [17] Zuckerman JD, Gallagher MA, Cuomo F, Rokito A. The effect of instability and subsequent anterior shoulder repair on proprioceptive ability. *J Shoulder Elbow Surg* 2003;12(2):105-9.

- [18] Edmonds G, Kirkley A, Birmingham TB, Fowler PJ. The effect of early arthroscopic stabilization compared to nonsurgical treatment on proprioception after primary traumatic anterior dislocation of the shoulder. *Knee Surg Sports Traumatol Arthrosc* 2003;11(2):116-21.
- [19] Iida N, Kaneko F, Aoki N, Shibata E. The effect of fatigued internal rotator and external rotator muscles of the shoulder on the shoulder position sense. *J Electromyogr Kinesiol* 2014;24(1):72-7.
- [20] Lonn J, Djupsjobacka M, Johansson H. Replication and discrimination of limb movement velocity. *Somatosens Mot Res* 2001;18(1):76-82.
- [21] Lonn J, Crenshaw AG, Djupsjobacka M, Pedersen J, Johansson H. Position sense testing: influence of starting position and type of displacement. *Arch Phys Med Rehabil* 2000;81(5):592-7.
- [22] Skinner HB, Wyatt MP, Hodgdon JA, Conard DW, Barrack RL. Effect of fatigue on joint position sense of the knee. *J Orthop Res* 1986;4(1):112-8.
- [23] Myers JB, Guskiewicz KM, Schneider RA, Prentice WE. Proprioception and neuromuscular control of the shoulder after muscle fatigue. *J Athl Train* 1999;34(4):362-7.
- [24] Dvir Z. Difference, significant difference and clinically meaningful difference: The meaning of change in rehabilitation. *J Exerc Rehabil* 2015;11(2):67-73.







