

# Superficial temporal artery-middle cerebral artery anastomosis patency correlates with cerebrovascular reserve in adult moyamoya syndrome patients

R. Aboukais, B. Verbraeken, X. Leclerc, C. Gautier, H. Henon, M. Vermandel,

T. Menovsky, J.-P. Lejeune

# ▶ To cite this version:

R. Aboukais, B. Verbraeken, X. Leclerc, C. Gautier, H. Henon, et al.. Superficial temporal arterymiddle cerebral artery anastomosis patency correlates with cerebrovascular reserve in adult moyamoya syndrome patients. Neurochirurgie, 2019, 65, pp.146 - 151. 10.1016/j.neuchi.2019.05.001 . hal-03488163

# HAL Id: hal-03488163 https://hal.science/hal-03488163

Submitted on 20 Dec 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Superficial temporal artery-middle cerebral artery anastomosis patency correlates with cerebrovascular reserve in adult moyamoya syndrome patients

Rabih Aboukais, MD<sup>1,2\*</sup>, Barbara Verbraeken, MD<sup>4</sup>, Xavier Leclerc, MD, PhD<sup>3</sup>, Corinne Gautier, MD<sup>3</sup>, Hilde Henon, MD, PhD<sup>5</sup>, Maximilien Vermandel, PhD<sup>2</sup>, Tomas Menovsky, MD, PhD<sup>4</sup>, Jean-Paul Lejeune MD, PhD<sup>1,2</sup>,

1 Department of Neurosurgery, Lille University Hospital, Hôpital Nord, Lille, France

2 Univ. Lille, INSERM, CHU Lille, U1189-ONCO-THAI-Image Assisted Laser Therapy for Oncology, F-59000, Lille, France.

3 Department of Neuroradiology, Lille University Hospital, Hôpital Nord, Lille, France

4 Department of Neurosurgery, Antwerp University Hospital, University of Antwerp, Belgium

5 Univ. Lille, INSERM U 1171 (Degenerative & Vascular Cognitive Disorders). CHU-Lille, Department of Neurology, Lille, France

Counts: Abstract: 225 words Total word count: 3,334 words References: 24 Tables + figures: 7

\* Corresponding author: rabihdoc@hotmail.com
Rabih Aboukais
Department of Neurosurgery, Lille University Hospital, Rue E. Laine, 59037 Lille cedex,
France
email: rabihdoc@hotmail.com
Telephone: +33.3.20.44.54.55
Fax: +33.3.20.44.55.11

Conflicts of interest: none

#### Abstract

**Objectives:** To evaluate the effectiveness of superficial temporal artery-middle cerebral artery (STA-MCA) bypass in improving cerebrovascular reserve (CVR) in Moyamoya syndrome.

**Patients and Methods:** This prospective study included 10 consecutive patients treated for Moyamoya syndrome by STA-MCA bypass in our institution between June 2016 and January 2018. Perfusion MRI, transcranial Doppler and 99m Tc-HMPAO SPECT with acetazolamide challenge were performed before and after treatment to evaluate perfusion and cerebrovascular reserve. STA-MCA bypass was indicated for patients with history of ischemic or hemorrhagic stroke and when CVR was diminished on both transcranial Doppler and 99m Tc-HMPAO SPECT with acetazolamide challenge or brain perfusion was deteriorated on MRI.

**Results:** Bypass anastomosis was patent in all patients at end of surgery. One patient presented partial postoperative sensorimotor deficit related to an ischemic lesion in the frontal cortical area. One patient presented regressive chronic subdural hematoma without neurological deficit. Three months after treatment, CVR was significantly improved in 8 patients and unchanged in 2, probably related to low flow. Further follow-up found CVR deterioration in 1 patient, with anastomosis occlusion at 1 year.

**Conclusion:** Our data suggest that improvement in cerebral perfusion and CVR depends on flow in the STA-MCA anastomosis in patients with Moyamoya syndrome. Systematic long-term follow-up of anastomosis flow, brain perfusion and CVR improves quantification of the benefit of STA-MCA anastomosis in terms of disease progression.

Keywords: anastomosis, cerebrovascular reserve, STA-MCA bypass, Moyamoya, brain perfusion

**Mots clés:** anastomose, réserve cérébrovasculaire, pontage temporo-sylvien, Moyamoya, perfusion cérébrale

## Introduction

Cerebrovascular reserve (CVR) is defined as the ability of brain parenchyma to adjust cerebral blood flow in response to altered metabolic demand or vasoactive stimulation. It may be used to identify new or recurrent stroke risk in patients with cerebrovascular disease[1]. Deterioration of CVR plays an important role in the occurrence of stroke in patients with Moyamoya syndrome[2]. As CVR is diminished in these patients, the risk of ischemic or hemorrhagic stroke increases. Some authors[3, 4] have reported an annual risk of ischemic stroke close to 13% and an annual risk of hemorrhagic stroke close to 8% in patients with Moyamoya syndrome when CVR is impaired. CVR and cerebral perfusion can be evaluated by transcranial Doppler[5], 99m Tc-HMPAO SPECT[6] with acetazolamide challenge, MRI[7] and CT scan[8] perfusion. Monitoring CVR is essential for treatment decision-making and treatment evaluation in Moyamoya syndrome[2]. The purpose of the treatment is to improve CVR in order to reduce risk of stroke[9]. Superficial temporal artery-middle cerebral artery (STA-MCA) bypass is the main treatment in adult patients with Moyamoya syndrome[10].

The goal of this study was to evaluate the effectiveness of STA-MCA bypass in improving CVR in patients with Moyamoya syndrome.

#### **Materials and Methods**

This prospective study included 10 consecutive patients treated for Moyamoya syndrome by STA-MCA bypass in our institution between June 2016 and January 2018. Approval was obtained from the local institutional review board (ref. DEC19-002). The series comprised 5 men and five women, with mean age at diagnosis of 44.8 years (range: 19-62 years).

#### Preoperative evaluation

All patients were clinically evaluated at our institution prior to treatment. Age at diagnosis, medical history, neurological status and American Society of Anesthesiology (ASA) score[11] were recorded. Moyamoya syndrome was diagnosed in all patients using conventional cerebral angiography and cranial magnetic resonance imaging (MRI). The STA main trunk and frontal and parietal branches were studied on conventional external carotid angiogram. Perfusion MRI, transcranial Doppler with acetazolamide challenge and 99m Tc-

HMPAO SPECT with acetazolamide challenge were performed before treatment to evaluate baseline perfusion and CVR.

CVR was assessed on HMPAO SPECT. After keeping the patient in a dark and quiet area for 30 minutes, SPECT and planar images were acquired with intravenous injection of 732 mBq 99mTc HMPAO. Scans were repeated within 3 to 7 days with acetazolamide challenge. Acetazolamide was injected intravenously at 20 mg/kg body weight and, 20 minutes later, the patient received 732 mBq 99mTc HMPAO. SPECT and planar images were acquired 20 minutes after administration of 99mTc HMPAO. Images were compared with and without acetazolamide challenge, to obtain the differences between gamma-ray counts in 2 cerebral hemispheres to assess CVR, a difference of more than 6% being defined as impaired CVR. Two of the 10 patients (20%) did not have baseline perfusion MRI, and CVR was evaluated by SPECT HMPAO and transcranial Doppler.

The treatment strategy was systematically discussed and decided upon by a multidisciplinary team including neurologists, neurosurgeons, neuroradiologists and intensivists. The STA-MCA bypass procedure was indicated for patients with history of ischemic or hemorrhagic stroke and when CVR was diminished on both transcranial Doppler and 99m Tc-HMPAO SPECT with acetazolamide challenge and on perfusion MRI. However, 1 patient without history of stroke nevertheless underwent the STA-MCA procedure because of severely deteriorated CVR. In accordance with the COSS study[12], the bypass procedure was never performed within 3 months of an acute episode of stroke[13]. Antiplatelet medication was not discontinued perioperatively.

#### Surgery

The course of the STA is marked on the skin by palpation and portable continuous-wave Doppler ultrasonography. A frontotemporal incision is carefully made, conserving STA integrity. A frontotemporal craniotomy is performed over the Sylvian fissure and the dura is opened to identify suitable recipient M3-M4 cortical arteries. Once a recipient artery has been identified, the STA is dissected under an operating microscope.

Depending on STA length and diameter, the frontal, parietal or both branches are carefully dissected, distally cut and proximally clipped with a temporary aneurysm clip. The recipient cortical artery or arteries are dissected from the overlying arachnoid and isolated with a rubber-leaf/foil layer. The cortical artery is temporarily clipped with a small aneurysm clip and arteriotomy is performed. Subsequently, end-to-side anastomosis of the STA to the cortical artery is performed using microneurosurgical techniques with 10-0 interrupted nylon

sutures. During bypass anastomosis clamping time, blood pressure is raised to normotensive values. STA-MCA anastomosis patency is confirmed by intraoperative indocyanine green video-angiography.

#### Postoperative evaluation

Early postoperative clinical examination was performed within the first week, with transcranial Doppler, MRI with diffusion and perfusion sequences, and CT angiography for each patient 1) to confirm bypass patency, 2) to measure flow in each anastomosis, 3) to detect any ischemic lesions and 4) to assess brain perfusion.

All patients were re-evaluated 3 months after bypass surgery. Clinical examination findings and modified Rankin Scale score (mRS) were recorded. MRI screened for ischemic lesions and assessed brain perfusion. Transcranial Doppler assessed flow in the anastomoses. At 3, 6 and 12 months after treatment, transcranial Doppler with acetazolamide challenge, 99m Tc-HMPAO SPECT with acetazolamide challenge and MRI with perfusion sequence were performed to evaluate CVR and brain perfusion. In case of discordance between test results, conventional CT angiography was performed to assess the internal and external carotid arteries.

#### Results

Mean age at diagnosis was 44.8 years (range, 19-62 years) and mean follow-up was 24 months (range, 6-30 months). ASA score was 1 in 2 patients, 2 in 7 and 3 in 1. Functional outcomes are presented in Table 1.

#### Intraoperative findings

STA-MCA bypass anastomosis was systematically feasible. In 2 patients, double STA-MCA anastomosis was performed because both frontal and parietal branches of STA were suitable. Mean occlusion time per anastomosis was 55 minutes (range, 33-85 minutes). Intraoperative indocyanine green video-angiography demonstrated anastomosis patency in all

patients at end of procedure. Mean microsurgical procedure time was 5.4 hours (range, 3.1-8.7 hours).

#### Microsurgical morbidity

One of the 10 patients (Patient 4) showed postoperative partial sensorimotor deficit related to an ischemic lesion in a frontal cortical area. Another patient (Patient 8) presented chronic subdural hematoma without neurological deficit.

#### Long-term clinical outcome

One patient (Patient 1) showed a frontoparietal cortical ischemic lesion with hemiparesis and aphasia 1 year after left-sided STA-MCA anastomosis. This stroke occurred during femoral vascular bypass surgery and was related to low intraoperative blood pressure (70 mmHg). There was no change in mRS score in the other patients at 1 year (Table 1).

#### Radiological outcome

Three months after treatment, brain perfusion and CVR improved significantly in 8 patients and remained unchanged in 2 (Table 2). STA-MCA anastomosis occlusion occurred in 1 patient (Patient 2) at 1 year, without neurological deterioration; CVR had improved after the bypass procedure and decreased after anastomosis occlusion (Figure 1). On perfusion MRI, transit time was 0 seconds when the STA-MCA anastomosis was patent, but 4 seconds when occluded.

In 1 patient (Patient 8), chronic subdural hematoma occurred after treatment, stretching the STA-MCA bypass; however, the bypass was not definitely occluded, as patency and improved CVR were recorded after regression of the hematoma (Figure 2).

One patient (Patient 9) showed no significant improvement in CVR after treatment; transcranial Doppler revealed only weak flow (19 mL/min) in the anastomosis.

#### Discussion

STA-MCA anastomosis patency correlated strongly with CVR improvement. Therefore, systematic long-term monitoring of flow and CVR on transcranial Doppler, SPECT HMPAO and perfusion MRI seems mandatory in patients with Moyamoya syndrome.

#### STA-MCA anastomosis morbidity

STA-MCA bypass is a valuable microsurgical procedure that demonstrated efficacy in many intracranial vascular diseases such as Moyamoya disease (MMD), aneurysm and skull base tumor[14]. STA-MCA bypass is usually described as a "low-flow" bypass, but conversion to a "high-flow" state in the late postoperative period has been reported by many authors [2, 15]. In our study, flow was >50 mL/min in 6 patients by 6 months post-operatively.

STA-MCA bypass shows low morbidity[16] and patency is approximately 95% at end of procedure[17-19]. In his systematic review, Schaller[20] reported that long-term bypass patency was excellent, with 1% failure per year following the first year after surgery. In his series of 264 MMD patients, including 450 revascularization procedures, Guzman[16] reported 3.5% surgical morbidity and 0.7% mortality per treated hemisphere. Complications were mainly related to MMD pathophysiology, implicating fragile vessels and hemodynamic instability[21]. Thus, risk of bypass thrombosis is slightly higher in case of MMD. Moreover, MMD can involve extracranial arteries, including the STA[22], which perhaps explains late STA-MCA anastomosis occlusion in our series.

#### Prognostic value of cerebrovascular reserve evaluation

Perfusion and CVR can be evaluated by various techniques, including perfusion MRI, transcranial Doppler with acetazolamide challenge and 99m Tc-HMPAO SPECT with acetazolamide challenge. In our study, the results of these techniques correlated[23, 24]: when transcranial Doppler showed anastomosis occlusion in Patients 2 and 8, SPECT HMPAO and perfusion MRI showed immediate deterioration of CVR.

In his review, Garrett[2] demonstrated that patients with severe hemodynamic failure were at higher risk of cerebral infarction than those with mild disease, and that patients with severe failure responded better to surgery. In a series of 23 patients who underwent STA-MCA bypass, O[25] reported that (99m)Tc HMPAO SPECT images of the territory supplied by the bypass graft showed increase in both cerebrovascular flow and reserve at baseline, and that the increase was significantly higher following administration of acetazolamide. Low[10] reported that transcranial Doppler and (99m)Tc HMPAO SPECT acetazolamide-challenge images repeated at  $4 \pm 1$  months showed significant improvement in CVR in the 46 patients who underwent STA-MCA bypass. Moreover, during a mean follow-up of 34 months (range, 18-39 months), only 6 of the 46 patients (13%) developed cerebral ischemic events. In our series, CVR improved significantly in 8 patients 3 months after STA-MCA bypass.

Patient 9 showed no significant improvement in CVR, probably due to weak bypass-flow as measured on transcranial Doppler. In Patient 8, CVR remained diminished because of postoperative chronic SDH, probably stretching the STA and reducing vessel lumen, with no flow seen on Doppler. A sudden deterioration of CVR was recorded in 1 patient 9 months after surgery, due to late bypass occlusion, without clinical signs of stroke; this late occlusion could be related to MMD involvement of the external carotid artery network, which includes the STA[22]. A cerebral ischemic event was noted in 1 patient with patent STA-MCA bypass 1 year after treatment, during femoral vascular bypass surgery with low intraoperative blood pressure: i.e., iatrogenic; CVR had been satisfactory after STA-MCA bypass, but presumably insufficient to protect the patient against an acute episode during a sudden drop in blood pressure (SBP 70 mmHg).

### Conclusion

STA-MCA bypass is a valid method to improve the brain perfusion and CVR in Moyamoya syndrome. Improvement in CVR depends on the flow and the patency of the STA-MCA anastomosis. Likewise, the success of the anastomosis may be related to the need for bypass to supplement the low blood flow demonstrated on CVR. Our data suggest that improvement in cerebral perfusion and CVR depends on the flow in the STA-MCA anastomosis in patients with Moyamoya syndrome. Systematic long-term follow-up evaluation of anastomosis flow, brain perfusion and CVR should better quantify the benefit of STA-MCA anastomosis in terms of disease progression.

## Legends

 Table 1. Functional outcomes

Table 2. Cerebrovascular reserve evaluation on different examinations

**Figure 1.** Patient 2, 19 years old, with a deterioration of CVR in left middle cerebral artery territory (red arrow) related to unilateral left-side Moyamoya disease. STA-MCA anastomosis was performed in June 2016. CVR increased after treatment, as demonstrated on 99m Tc-HMPAO SPECT with acetazolamide challenge in December 2016. STA-MCA anastomosis occlusion occurred one 1 year after treatment and CVR decreased as demonstrated on 99m Tc-HMPAO SPECT with acetazolamide challenge in July 2017.

Figure 2. Patient 8, 55 years old, with deterioration of CVR in left middle cerebral artery territory related to unilateral left-side Moyamoya disease. STA-MCA anastomosis was

performed but persistent CVR deterioration was noted on 99m Tc-HMPAO SPECT with acetazolamide challenge 3 months after treatment (Figure 2A). MRI demonstrated a chronic subdural hematoma, with the STA stretched and thinned (Figure 2B) and no flow was detected in the STA-MCA anastomosis by cranial Doppler. After administration of corticosteroids, MRI demonstrated regression of the chronic subdural hematoma (Figure 2C) and transcranial Doppler revealed recanalization of the STA-MCA anastomosis. Improvement in cerebrovascular reserve was noted on 99m Tc-HMPAO SPECT with acetazolamide challenge (Figure 2D).

Funding-No funding was received for this research.

Conflict of Interest: All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest; and expert testimony or patent-licensing arrangements), or non-financial interest (such as personal or professional relationships, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript.

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee (Lille University Hospital/ Ethics committee) and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards

# References

- [1] Juttukonda MR, Donahue MJ. Neuroimaging of vascular reserve in patients with cerebrovascular diseases. NeuroImage 2017.
- [2] Garrett MC, Komotar RJ, Starke RM, Merkow MB, Otten ML, Sciacca RR, et al. The efficacy of direct extracranial-intracranial bypass in the treatment of symptomatic hemodynamic failure secondary to athero-occlusive disease: a systematic review. Clinical Neurology and Neurosurgery 2009;111(4):319-26.
- [3] Kraemer M, Heienbrok W, Berlit P. Moyamoya disease in Europeans. Stroke 2008;39(12):3193-200.
- [4] Duan L, Bao XY, Yang WZ, Shi WC, Li DS, Zhang ZS, et al. Moyamoya disease in China: its clinical features and outcomes. Stroke 2012;43(1):56-60.
- [5] Douvas I, Moris D, Karaolanis G, Bakoyiannis C, Georgopoulos S. Evaluation of cerebrovascular reserve capacity in symptomatic and asymptomatic internal carotid stenosis with transcranial Doppler. Physiological research 2016;65(6):917-25.
- [6] Acker G, Lange C, Schatka I, Pfeifer A, Czabanka MA, Vajkoczy P, et al. Brain Perfusion Imaging Under Acetazolamide Challenge for Detection of Impaired Cerebrovascular Reserve Capacity: Positive Findings with (15)O-Water PET in Patients with Negative (99m)Tc-HMPAO SPECT Findings. Journal of Nuclear Medicine 2018;59(2):294-8.
- [7] Kawano T, Ohmori Y, Kaku Y, Muta D, Uekawa K, Nakagawa T, et al. Prolonged Mean Transit Time Detected by Dynamic Susceptibility Contrast Magnetic Resonance Imaging Predicts Cerebrovascular Reserve Impairment in Patients with Moyamoya Disease. Cerebrovasc Dis 2016;42(1-2):131-8.
- [8] Takahashi S, Tanizaki Y, Akaji K, Kimura H, Katano T, Suzuki K, et al. Identification of hemodynamically compromised regions by means of cerebral blood volume mapping utilizing computed tomography perfusion imaging. Journal of Clinical Neuroscience 2017;38:74-8.
- [9] Qian C, Yu X, Li J, Chen J, Wang L, Chen G. The Efficacy of Surgical Treatment for the Secondary Prevention of Stroke in Symptomatic Moyamoya Disease: A Meta-Analysis. Medicine 2015;94(49):e2218.
- [10] Low SW, Teo K, Lwin S, Yeo LL, Paliwal PR, Ahmad A, et al. Improvement in cerebral hemodynamic parameters and outcomes after superficial temporal arterymiddle cerebral artery bypass in patients with severe stenoocclusive disease of the intracranial internal carotid or middle cerebral arteries. Journal of Neurosurgery 2015;123(3):662-9.
- [11] Aronson WL, McAuliffe MS, Miller K. Variability in the American Society of Anesthesiologists Physical Status Classification Scale. AANA Journal 2003;71(4):265-74.
- [12] Powers WJ, Clarke WR, Grubb RL, Jr., Videen TO, Adams HP, Jr., Derdeyn CP. Extracranial-intracranial bypass surgery for stroke prevention in hemodynamic cerebral ischemia: the Carotid Occlusion Surgery Study randomized trial. JAMA 2011;306(18):1983-92.
- [13] Grubb RL, Jr., Powers WJ, Clarke WR, Videen TO, Adams HP, Jr., Derdeyn CP. Surgical results of the Carotid Occlusion Surgery Study. Journal of Neurosurgery 2013;118(1):25-33.
- [14] Sekhar LN, Natarajan SK, Ellenbogen RG, Ghodke B. Cerebral revascularization for ischemia, aneurysms, and cranial base tumors. Neurosurgery 2008;62(6 Suppl 3):1373-408; discussion 408-10.

- [15] Cherian J, Srinivasan V, Kan P, Duckworth EAM. Double-Barrel Superficial Temporal Artery-Middle Cerebral Artery Bypass: Can It Be Considered "High-Flow?". Oper Neurosurg (Hagerstown) 2018;14(3):288-94.
- [16] Guzman R, Lee M, Achrol A, Bell-Stephens T, Kelly M, Do HM, et al. Clinical outcome after 450 revascularization procedures for moyamoya disease. Clinical article. Journal of Neurosurgery 2009;111(5):927-35.
- [17] Charbel FT, Meglio G, Amin-Hanjani S. Superficial temporal artery-to-middle cerebral artery bypass. Neurosurgery 2005;56(1 Suppl):186-90; discussion -90.
- [18] Newell DW, Vilela MD. Superficial temporal artery to middle cerebral artery bypass. Neurosurgery 2004;54(6):1441-8; discussion 8-9.
- [19] Gross BA, Du R. STA-MCA bypass. Acta Neurochirurgica 2012;154(8):1463-7.
- [20] Schaller B. Extracranial-intracranial bypass surgery to reduce the risk of haemodynamic stroke in cerebroocclusive atherosclerotic disease of the anterior cerebral circulation a systematic review. Neurologia i Neurochirurgia Polska 2007;41(5):457-71.
- [21] Brandicourt P, Bonnet L, Bejot Y, Drouet C, Moulin T, Thines L. Moya-Moya syndrome after cranial radiation for optic glioma with NF1. Case report and literature review of syndromic cases. Neuro-Chirurgie 2018;64(1):63-7.
- [22] Sun SJ, Zhang JJ, Li ZW, Xiong ZW, Wu XL, Wang S, et al. Histopathological features of middle cerebral artery and superficial temporal artery from patients with moyamoya disease and enlightenments on clinical treatment. Journal of Huazhong University of Science and Technology Medical sciences = Hua zhong ke ji da xue xue bao Yi xue Ying De wen ban = Huazhong ke ji daxue xuebao Yixue Yingdewen ban 2016;36(6):871-5.
- [23] Kim JH, Lee SJ, Shin T, Kang KH, Choi PY, Gong JC, et al. Correlative assessment of hemodynamic parameters obtained with T2\*-weighted perfusion MR imaging and SPECT in symptomatic carotid artery occlusion. AJNR 2000;21(8):1450-6.
- [24] Kikuchi K, Murase K, Miki H, Kikuchi T, Sugawara Y, Mochizuki T, et al. Measurement of cerebral hemodynamics with perfusion-weighted MR imaging: comparison with pre- and post-acetazolamide 133Xe-SPECT in occlusive carotid disease. AJNR 2001;22(2):248-54.
- [25] O JH, Jang KS, Yoo Ie R, Kim SH, Chung SK, Sohn HS, et al. Assessment of cerebrovascular reserve before and after STA-MCA bypass surgery by SPECT and SPM analysis. Korean Journal of Radiology 2007;8(6):458-65.





















































%













































Patient	Age at	Disease	Preoperative	Preop	mRS score 3 months after
	diagnosis		symptoms	erativ	bypass
	C			e	• •
				mRS	
				score	
1		Atheromato	3 ischemic		
		us	strokes with		
	50	Moyamoya	motor aphasia	2	2
2		Unilateral	Headache and		
		Moyamoya	paresthesia, no		
	19		stroke	2	2
3		Bilateral	2 left ischemic		
		Moyamoya	strokes with		
			transient motor		
	49		aphasia	2	2
4		Unilateral	1 right ischemic		
		Moyamoya	stroke without		
	49		symptoms	2	2
5		Unilateral	2 left ischemic		
		Moyamoya	strokes with		
	43		hemiparesis	2	2
6		Cavernous	1 left ischemic		
		sinus	stroke with		
		meningioma	transient motor		
		with ICA	aphasia		
	62	occlusion		0	0
7		Bilateral	1 right ischemic		
		Moyamoya	stroke and 1 left		
			brain hemorrhage		
	30		with hemiparesis	2	2
8		Unilateral	1 left ischemic		
		Moyamoya	stroke with motor		
	55		aphasia	1	2
9		Bilateral	Headache without		2
	47	Moyamoya	stroke	1	
10	45	Bilateral	Right ischemic	3	3
		Moyamoya	stroke with		
			hemiparesis		

Table 1. Functional outcomes

Patien	Preoperati	Flow in	Early	Perfusion	99m Tc-	Transcranial
t	ve	anastomosi	postoperati	MRI 3	HMPAO	Doppler with
	perfusion	s on early	ve	months	SPECT with	acetazolamid
	MRI	transcrania	perfusion	after	acetazolamid	e challenge 3
	(mean	l Doppler	MRI (mean	anastomos	e challenge 3	months after
	transit	(mL/minut	transit time)	is (mean	months after	anastomosis
	time)	e)		transit	anastomosis	(improvement
				time)	(improvement	in
					in	cerebrovascul
					cerebrovascul	ar reserve)
					ar reserve)	
1	5 seconds	85	5 seconds	5 seconds	+	+
2	2 seconds	82	2 seconds	0 seconds	+	+
3	4 seconds	70	2 seconds	1 second	+	+
4	2 seconds	55	2 seconds	-	+	+
5	-	45	-	3 seconds	+	+
6	7 seconds	35	3 seconds	3 seconds	+	+
7	-	120	-	0 seconds	+	+
8	4 seconds	45	3 seconds	3 seconds	-	-
9	7 seconds	19	5 seconds	4 seconds	-	-
10	4 seconds	48	3 seconds	2 seconds	+	+

Table 2. Cerebrovascular reserve evaluation on different examinations