



HAL
open science

Response of dung beetle assemblages to grazing intensity in two distinct bioclimatic contexts

William Perrin, Marco Moretti, Alan Vergnes, Daniel Borcard, Pierre Jay-Robert

► **To cite this version:**

William Perrin, Marco Moretti, Alan Vergnes, Daniel Borcard, Pierre Jay-Robert. Response of dung beetle assemblages to grazing intensity in two distinct bioclimatic contexts. *Agriculture, Ecosystems & Environment*, 2020, 289, pp.106740 -. 10.1016/j.agee.2019.106740 . hal-03488621

HAL Id: hal-03488621

<https://hal.science/hal-03488621>

Submitted on 21 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 Response of dung beetle assemblages to grazing intensity in
2 two distinct bioclimatic contexts

3 William Perrin ^a, Marco Moretti ^b, Alan Vergnes ^a, Daniel Borcard ^c, Pierre Jay-Robert ^a

4

5 ^a Centre d'Écologie Fonctionnelle et Évolutive (UMR 5175 CEFE), Université Paul Valéry
6 Montpellier 3, Université de Montpellier, École Pratique des Hautes Études (EPHE), Centre
7 National de la Recherche Scientifique (CNRS), Institut de Recherche pour le Développement
8 (IRD), Montpellier, France

9 ^b Biodiversity and Conservation Biology, Swiss Federal Research Institute WSL, Birmensdorf,
10 Switzerland

11 ^c Département de sciences biologiques, Université de Montréal, C.P. 6128, succursale
12 Centre-ville, Montréal (Québec), H3C 3J7, Canada

13

14 William Perrin: william.perrin@univ-montp3.fr (**corresponding author**)

15 Marco Moretti: marco.moretti@wsl.ch

16 Alan Vergnes: alan.vergnes@univ-montp3.fr

17 Daniel Borcard: daniel.borcard@umontreal.ca

18 Pierre Jay-Robert: pierre.jay-robert@univ-montp3.fr

19

20 **Abstract**

21 Given the impact of livestock on ecosystems worldwide, it is necessary to understand the
22 effects of grazing practices on biodiversity in order to improve the sustainability of pasture
23 management practices. In a pasture, spatio-temporal variability in livestock activity results in
24 a heterogeneous distribution of defoliation, trampling and excreta. To date, fine-scale
25 analyses of grazing intensity have been rare, and the geographical extent of the studies often
26 limited. In this study, we addressed this gap by analysing the influence of contrasting intra-
27 pasture grazing intensity on the structure and composition of dung beetle assemblages. To
28 do this, we studied a three-level grazing intensity gradient in two distinct bioclimatic
29 contexts, a Mediterranean steppe and the Alps, which also allowed us to determine if dung
30 beetle responses to grazing intensity are related to bioclimatic conditions. The observed
31 dung beetle responses showed an imprint of the bioclimatic context and the local pasture
32 conditions, and species composition and relative abundance showed strong variations along
33 the grazing intensity gradient in both study areas. Species assemblages from the most and
34 least grazed parts of pastures differed strongly. By altering habitat conditions, changing dung
35 availability and modifying competitive interactions, fine-scale heterogeneity in grazing
36 intensity led to substantial variations in the abundance of dung beetle nesting guilds. In both
37 study areas an increase in grazing intensity was detrimental to the largest species and the
38 soil-digging species (which bury dung in underground nests), whereas dung-dwelling species
39 (which reproduce inside dung pads) were favoured. We discuss the combined use of nesting
40 guilds and body mass as potential features to generalize the application of dung beetles as
41 indicators of grazing practices.

42 Keywords: Pastoralism; IndVal; Nesting guilds; Scarabaeoidea; Alps; Mediterranean

43 **1. Introduction**

44 Livestock grazing has been shown to be one of the main anthropogenic factors that have
45 structured biodiversity for centuries in many regions worldwide (Alkemade et al., 2013). In
46 European grasslands today, large wild ungulates have been largely replaced by domestic
47 mammals (Bradshaw et al., 2003), which are now the most important grazers and one of the
48 largest land-use sectors in rural economies. In the European Union, pastureland covers one-
49 third of the area used for agriculture (European Commission, 2015). The extensive
50 management of livestock supports rich biodiversity (Redecker et al., 2002) and is
51 consequently promoted by European policy (Kerven and Behnke, 2011). However, the
52 socioeconomic context drives polarization in livestock activities, with large areas left
53 abandoned while practices are intensified in concentrated locations (Mazoyer and Roudart,
54 1997; Steinfeld et al., 2010). These trends in grazing intensity and livestock management are
55 likely to cause profound changes in ecosystems (Squires et al., 2018; Stoate et al., 2009) and
56 may have negative effects on biodiversity. The abandonment of the traditional practice of
57 extensive grazing on rangelands reduces landscape heterogeneity, which leads to a decline
58 in species diversity, specifically for specialist species of open habitats (García-Tejero et al.,
59 2013; Hodgson et al., 2005; Sirami et al., 2010; Wenzel et al., 2006). Additionally, increasing
60 livestock stocking rates may reduce species diversity (e.g. in insects, Kruess and Tschardtke,
61 2002; and plants, Pizzio et al., 2016), create unfavourable conditions for grassland breeding
62 species (e.g. in birds, Paine et al., 1996) and disrupt ecosystem functioning, for example, by
63 altering soil properties (Zhou et al., 2017).

64 In terrestrial ecosystems, arthropods are among the most diverse and numerous organisms,
65 with an importance in ecosystem function inversely proportional to their small size (Wilson,

66 1987). A review by van Klink et al. (2015) showed that livestock effects on arthropod
67 diversity are modulated by grazing intensity. For example, the most diverse arthropod
68 assemblages seem to be found in heterogeneous pastures created by intermediate grazing
69 intensity. Most of the reported results concerned predators (e.g. spiders and ground beetles)
70 and phytophagous insects (e.g. butterflies and grasshoppers) directly affected by grazing
71 mammals through the modification of the physical habitat (vegetation structure and cover)
72 and plant food resources (diversity and abundance of consumed plant species) (van Klink et
73 al., 2015). In addition to changing the structure and composition of vegetation, herbivorous
74 mammals provide faeces (dung pads) used by many organisms for food or reproduction
75 (Hanski and Cambefort, 1991). Of these organisms, dung beetles (Coleoptera: Scarabaeinae,
76 Aphodiinae, Geotrupinae) are an important group involved in the key ecosystem processes
77 of dung removal and burial (Milotić et al., 2019; Slade et al., 2007) as well as secondary seed
78 dispersal (Griffiths et al., 2016). Without such recycling, pastures become covered by
79 patches of grass with lower digestibility and nutritive value, rapidly becoming unsuitable for
80 grazing (Gittings et al., 1994). To exploit ephemeral and patchily dispersed dung pads, dung
81 beetle assemblages display a diversity of feeding and breeding strategies, reducing
82 competition between co-occurring species (Hanski and Cambefort, 1991). In this way, dung
83 beetles can be classified into two types of nesting guilds, based on the main strategy used by
84 a species to exploit dung resources (Jay-Robert et al., 2008a): dung-dwellers, which develop
85 entirely inside the dung, or soil-diggers, which relocate dung fragments in the soil for feeding
86 and breeding. These guilds are a useful way to understand the processes by which dung
87 beetles respond to land-use management gradients (Campos and Hernández, 2015;
88 Jankielsohn et al., 2001; Numa et al., 2012; Tonelli et al., 2019).

89 Compared to herbivorous and predator arthropods, few studies have addressed the effects
90 of grazing intensity on dung beetle assemblages in temperate regions (van Klink et al., 2015),
91 although dung beetles are known to be good indicators of changes in environmental
92 conditions and of various anthropogenic pressures (Audino et al., 2014; Bicknell et al., 2014;
93 McGeoch et al., 2002; Spector, 2006). In Western Europe, these insects are closely linked to
94 domestic ungulates in open landscapes (Buse et al., 2015; Jay-Robert et al., 2008b; Lobo et
95 al., 2006; Macagno and Palearini, 2009). The preference of some species for shrubby and
96 woody habitats can allow them to be used as indicators of changes in landscape structure
97 due to grazing abandonment (Tonelli et al., 2019; Verdú et al., 2011). The studies conducted
98 by Tonelli et al. (2019, 2018) in Italian pastures highlighted the negative effects of grazing
99 abandonment on dung beetles, reducing alpha diversity, evenness and dung beetle biomass,
100 as a result of insufficient dung inputs to sustain diverse assemblages. Equally, an increase in
101 intensive farming has also been shown to be detrimental to dung beetle diversity, mainly
102 due to habitat deterioration (soil trampling) from high grazing intensity (Hutton and Giller,
103 2003; Negro et al., 2011). These studies used a research approach that compared distinct
104 pastures with different stocking rates or grazing management methods. However, distinct
105 pastures may differ by more properties than stocking rate (e.g. the possible presence of wild
106 ungulates in abandoned pastures, differences in vegetation structure, use of anthelmintic
107 drugs), which makes the effects of grazing mammals per se difficult to assess. Within a
108 pasture, spatio-temporal variability in livestock activity (e.g. foraging, resting areas), which
109 depends on the type of livestock management employed, results in a spatially irregular
110 distribution of defoliation, trampling and nutrient input (from dung and urine production).
111 One might expect that the heterogeneous availability of dung and varying perturbation
112 intensity within a pasture may affect the structure and composition of dung beetle

113 assemblages. But while this intra-pasture variation in grazing intensity may be a more
114 appropriate scale of analysis, it is rarely considered (however, see Chillo et al., 2017;
115 Jerrentrup et al., 2014). Likewise, the potential use of dung beetles as indicators of local
116 grazing practices at this fine scale of analysis has not been assessed.

117 In this study, we investigated for the first time the effects of contrasting intra-pasture
118 grazing intensity on dung beetle assemblages, addressing the following questions: (i) How
119 does grazing intensity affect the composition and structure (richness, evenness) of dung
120 beetle assemblages at this scale? (ii) How does grazing intensity affect the abundance of the
121 two nesting guilds, i.e. dung-dwelling and soil-digging species? (iii) Can we define indicator
122 species for the impact of local grazing practices?

123 To explore these questions, we used a three-level grazing intensity gradient within
124 extensively managed pastures with grazing practices that were as similar as possible. To our
125 knowledge, very few previous studies have compared the effects of grazing on arthropod
126 communities in distinct bioclimatic regions (Báldi et al., 2013; Batáry et al., 2010, 2007;
127 Kőrösi et al., 2012; and one on dung beetles in a tropical context, Barragán et al., 2014) so
128 we also compared two very distinct bioclimatic areas in France: a Mediterranean semi-arid
129 lowland steppe, and mountain pastures (2000 m a.s.l.) in the Western Alps. These two areas
130 support different dung beetle communities: one typical of Mediterranean dry steppe
131 grasslands (Tatin et al., 2014) and one adapted to the high-altitude climate of the Alps
132 (Lumaret and Stiernet, 1991). By comparing the dung beetle assemblage/grazing intensity
133 relationship between the two areas, we wanted to determine if the effects of grazing
134 depend on local conditions and species identity or if they are similar across ecological

135 contexts. We also expected that investigating contrasting contexts would improve the ability
136 of using dung beetles as indicators for grazing intensity.

137 **2. Material & methods**

138 *2.1. Study areas and sampling design*

139 The study was carried out in pastures in two protected areas in France that differ markedly
140 in terms of bioclimatic conditions: the Coussouls de Crau National Nature Reserve and the
141 Vanoise National Park (**Figure 1**). The Coussouls de Crau Reserve (hereafter, “the steppe”) is
142 a vast area of dry grasslands (11,000 ha) located near the Mediterranean Sea (43°33’N,
143 4°51’E) at an altitude of less than 50 m a.s.l. (Tatin et al., 2013). The climate is typically
144 Mediterranean, with a dry season in summer, two periods of rainfall in spring and autumn,
145 and winds above 50 km•h⁻¹ more than 110 days a year (Devaux et al., 1983). In this plain,
146 evidence of sheep grazing goes back to antiquity (Badan et al., 1995). Today, some 40,000
147 sheep still graze this semi-arid rangeland from the end of winter to early summer (Tatin et
148 al., 2013). The grazing activity is organized in a patchwork of 70 contiguous pastures (grazing
149 areas) through which a shepherd conducts his/her flock. During the night, the sheep are kept
150 in a barn or in an outdoor fenced enclosure, depending on the flock size. The Vanoise
151 National Park (hereafter, “the Alps”) is located in the western Alps (45°14’N, 6°43’E) and has
152 a typical alpine climate, with snowy winters and mild summers. The topography of the Alps
153 has favoured high-altitude pastoralism (above 2000 m in elevation). Every year 61,000 sheep
154 graze the alpine pastures of the National Park, following a transhumant cycle: herds are
155 brought up to the alpine pastures in early summer and moved down to the valleys and plains
156 in autumn. Shepherds conduct their flocks across the pasture during the day, and the sheep
157 are kept in outdoor fenced enclosures at night. In both areas, grazing follows traditional

158 management practices that have been used for centuries: the livestock primarily graze on
159 the native vegetation of natural grasslands, except during the cold season when the animals
160 stay in barns.

161 In each study area we selected two distinct pastures (**Figure 1**) with a comparable long-term
162 grazing regime (**Appendix A**). The small number of pasture replicates is due to the scarcity of
163 sites where we were sure that no antiparasitic treatment had been administered to the
164 sheep for at least three months prior to sampling (Beynon et al., 2012; Sands and Wall, 2018;
165 Verdú et al., 2018). The two pastures were located 6.2 km apart in the steppe, and 3.2 km
166 apart in the Alps; within a study area, both pastures had similar environmental conditions
167 (**Appendix A**). One distinct sheep flock grazed each pasture during a four-month period on
168 average (March–June in the steppe and July–October in the Alps).

169 Based on detailed knowledge of local grazing practices provided by herders and shepherds,
170 in each pasture, we selected three zones characterized by different grazing intensity (GI),
171 which we defined as Low (LGI), Moderate (MGI) and High (HGI) (**Figure 1**). In all the studied
172 pastures, certain zones were little exploited by shepherds. In the Alps, LGI zones
173 corresponded to areas that are usually difficult to access or had limited visibility, principally
174 due to steep terrain. In the steppe, there are no physical boundaries between contiguous
175 pastures, so the LGI zones were located at pasture edges, where shepherds only occasionally
176 take their flock to avoid the risk of grazing on neighbouring grazing areas. In these LGI zones,
177 we assumed that the availability of sheep droppings was low. The MGI zones were located in
178 main diurnal grazing zones, where shepherds regularly herd their flock each day to graze.
179 There, dropping distribution was temporally regular but spatially heterogeneous, depending
180 on the preferential foraging areas, which vary throughout the grazing season (depending on

181 forage availability). The HGI zones were located within or near the overnight site of the flock.
182 These zones are characterized by intensive and repeated trampling and a high load of
183 droppings. As each pasture contained only one overnight site (i.e. one HGI zone available to
184 sample), it was not possible to replicate the different grazing conditions within a pasture.
185 The mean distances between the different GI zones are shown in **Appendix B (Tables B.1,**
186 **B.2).**

187 In all the pastures, both in the steppe and the Alps, the vegetation was herbaceous and
188 homogeneous across the different GI zones; isolated shrubs were found only in the least
189 grazed zones of pastures. To validate our a priori characterization of the different GI zones,
190 we measured the maximum height of herbaceous vegetation and the amount of sheep
191 droppings in the representative plots where dung beetles were sampled (the protocol is
192 detailed in **Appendix C**). We assumed that these descriptors (height of herbaceous
193 vegetation and quantity of droppings) were good surrogates of grazing intensity levels
194 between the different GI zones (Noy-Meir et al., 1989; Rotem, 2016; Xu et al., 2014).
195 Moreover, they were relevant to the purpose of our study: (i) dropping availability is a key
196 factor determining the coexistence of dung beetle species at a fine spatial scale (Finn and
197 Giller, 2000; Horgan, 2005) and (ii) herbaceous height depends on the intensity of herbivory
198 and on the severity of disturbance induced by sheep herds. The results showed that the
199 maximum height of herbaceous vegetation significantly decreases while the amount of
200 sheep droppings significantly increases in line with higher grazing intensity (**Appendix C;**
201 **Figures C.1, C.2; Table C.1**).

202 *2.2. Dung beetle sampling*

203 Within each LGI, MGI and HGI zone in each pasture we selected one sampling plot where
204 dung beetles were collected using five pitfall traps (type CSR, described by Veiga et al., 1989)
205 **(Figure 1)**. In both study areas, HGI zones had a small surface area (less than 5000 m²).
206 Therefore, in order to sample as homogeneous a habitat as possible in terms of grazing
207 pressure, we placed the CSR traps 10 m apart in accordance with the standard design used in
208 temperate contexts (Frank et al., 2017; Jay-Robert et al., 2008a, 2008b; Lobo et al., 2006;
209 Lobo et al., 1998). The pitfall traps consisted of plastic basins (Ø 20 cm, depth 15 cm) buried
210 to the rim in the soil, filled with water and a few drops of neutral soap, and covered with a
211 grid, on top of which we placed 300 g of fresh sheep droppings. The traps were left open for
212 72 hours on sunny days during the time when dung beetle species richness and abundance is
213 highest in the French Mediterranean (Jay-Robert et al., 2008a) and the Alps (Jay-Robert et
214 al., 2008b): the sampling was conducted from 23 to 26 April 2018 in the steppe and from 28
215 July to 4 August 2017 in the Alps. Overall, 60 pitfall traps were set up (5 per level of grazing
216 intensity × 3 GI levels per pasture × 2 pastures × 2 areas). The 15 traps used in each pasture
217 were assumed to allow a close to exhaustive representation of the composition of local
218 species pools (Lobo et al., 1998). All captured individuals were identified to species level
219 using the taxonomic key provided by Paulian and Baraud (1982) and the nomenclature
220 provided by Löbl and Löbl (2016).

221 *2.3. Sampling completeness*

222 Since the number of species captured in a given trap is expected to be sensitive to the
223 number of individuals sampled (Chao et al., 2005), we calculated two estimators of
224 asymptotic species richness based on species-incidence data, in order to test the
225 completeness of our assemblage samples. We calculated Chao2 (Chao, 1987), which is

226 considered a robust estimator of minimum richness, and ICE based on estimated sampling
227 coverage, i.e. the proportion of assemblage richness represented by the species in a set of
228 replicated incidence samples (Lee and Chao, 1994). We used improved versions (iChao2 and
229 ICE-1) of these indices as these increase accuracy when the heterogeneity of species
230 abundance is relatively high (this was the case with our dataset) (Chiu et al., 2014; Lee and
231 Chao, 1994). We used the SpadeR software (Chao et al., 2015) to compute these two species
232 richness estimators.

233 *2.4. Patterns in species assemblage composition*

234 We analysed the assemblage-level responses using a Multivariate ANOVA based on a
235 Redundancy Analysis (RDA) (Legendre and Legendre, 2012) calculated on species abundance
236 matrices from the steppe and the Alps separately, using pitfall traps as sampling units.
237 Redundancy Analysis is a constrained ordination method related to principal component
238 analysis (Legendre and Anderson, 1999; Legendre and Legendre, 2012). An RDA-based
239 MANOVA allows testing the influence of selected factors and their interaction on
240 multivariate response matrices (in this case, species assemblages) and representing the
241 results in triplots as in a standard RDA. The difference in community composition between
242 the steppe and the Alps was very large, requiring no statistical tests (only 4 common species
243 out of 52; **Appendix D**). So in order to avoid a uselessly complicated ANOVA design that
244 would have included problematic permutation testing procedures (Anderson and Braak,
245 2003), we ran the test separately for each study area. In this test, we analysed the influence
246 of two predictors on the structure of dung beetle assemblages: (i) “*pasture identity*” (two
247 levels), and (ii) “*grazing intensity*” (three levels: LGI, MGI or HGI). We recoded these two
248 factors as orthogonal Helmert contrasts, enabling us to test the predictors and their

249 interaction (Borcard et al., 2018) and reflecting our crossed sampling design within each
250 study area (levels of grazing intensity crossed with the two pastures in each study area;
251 **Figure 1**). Before applying the RDA-based MANOVA, we Hellinger-transformed the
252 abundance of species to reduce the effect of extreme values and minimize the effect of
253 double absences in the species matrices (Legendre and Gallagher, 2001). We tested the
254 assumption of homogeneity of group dispersions (variance–covariance matrices) for the two
255 factors with Anderson's (2006) permutation test. The significance of “*pasture identity*” and
256 “*grazing intensity*”, as well as their interaction, was assessed with pseudo-*F* permutation
257 tests. In our case, in the presence of only two replicates for the “*pasture identity*” factor, the
258 application of a crossed design and the consequent definition of an interaction term was a
259 technical choice that allowed us to verify, through the interaction test, whether the effect of
260 grazing intensity was the same in the two pastures. Since the interaction was significant, the
261 effect of grazing intensity had to be assessed separately in each pasture. Consequently,
262 separate one-way RDA-based MANOVAs were performed on each pasture, with “*grazing*
263 *intensity*” as an explanatory factor, and R-squared values were calculated to show the
264 percentage of variance explained. We used the R package “vegan” (Oksanen et al., 2018) (R
265 Core Team, 2019) to test multivariate homogeneity of variance (using the *betadisper*
266 function) and fit the RDA-based MANOVAs (using the *rda* and *anova* functions).

267 2.5. Number of individuals, rarefied species richness, Pielou's evenness and nesting 268 guilds

269 For each pitfall trap, we measured (i) the number of captured individuals, (ii) the rarefied
270 species richness based on the minimum number of dung beetles sampled in a trap (i.e. 15 in
271 the steppe and 94 in the Alps), allowing us to minimize density effects (Gotelli and Colwell,

272 2010), and (iii) Pielou's evenness (1966). For the calculation of rarefied species richness, we
273 removed one trap that captured only five individuals to avoid standardizing species richness
274 from an insufficient number of individuals. However, we retained this trap to estimate
275 variation in the other response variables studied, since removing or keeping it had no effect
276 on models parameters. We also recorded the number of individuals belonging to the two
277 nesting guilds, dung-dwellers and soil-diggers. Dung-dwellers develop entirely inside the
278 dung, which they use as both a food and nesting resource (Hanski and Cambefort, 1991).
279 Several dung-dwelling species may also lay eggs at the dung–soil interface, enabling larvae to
280 move into the top layer of soil depending on the environmental conditions (Gittings and
281 Giller, 1997; Landin, 1961; Lumaret and Kirk, 1991). Soil-diggers relocate dung fragments and
282 pack them into the soil for feeding and breeding purposes. Underground nests can either be
283 located directly beneath a dung pad or at some distance from it, as the dung can be
284 extracted and the dung balls rolled to another location (Hanski and Cambefort, 1991). The
285 larvae develop in a dung brood ball, or brood mass, which is packed in a chamber or at the
286 end of a tunnel, depending on the species. In European dung beetles, small and medium-
287 sized soil-diggers usually nest at a depth between 3 and 15 cm, while the largest species can
288 dig nests deeper than 30 cm (Klemperer, 1979; Lumaret, 1983).

289 We assessed the effect of grazing intensity on the five response variables (number of
290 captured individuals, rarefied species richness, Pielou's evenness, number of dung-dwellers
291 and number of soil-diggers) using regressions, with pitfall traps as sampling units. Our
292 sampling design featured two nested levels: two studied areas (steppe and Alps), each of
293 which had two pastures (**Figure 1**). As mentioned above, the three levels of grazing intensity
294 (LGI, MGI and HGI) were crossed with the pasture identity. Because our aim was to compare
295 the response of dung beetle communities to grazing between the two study areas and the

296 difference in the community composition between these areas was evident, the models
297 were fitted separately in each study area (as was done for the RDA-based MANOVAs). We
298 thus removed the highest level of the hierarchical structure (i.e. the study area). In each
299 model, we tested the effects of “*grazing intensity*” (fixed factor with three levels: LGI, MGI,
300 HGI) on the five response variables described above. One can consider the “*pasture identity*”
301 as a random factor, as the pastures we sampled were chosen from a large number of
302 pastures in each study area. However, the insufficient number of independent pastures in
303 each study area (n = 2) prevented us from applying mixed models with this factor as a
304 random effect, as mixed model estimation requires a reasonable number of random-effects
305 levels. Gelman and Hill (2007) argue that when the number of groups is small (less than five),
306 there is not enough information to accurately estimate group-level variation. As a result,
307 multilevel models in this context typically gain little relative to standard models. Thus, we
308 used standard linear and generalized linear models, considering the “*pasture identity*” as a
309 fixed effect. In the Results section, we focus on the effects of the “*grazing intensity factor*” –
310 the results for the “*pasture identity*” factor are presented in **Appendix E**. Generalized linear
311 models allow the use of non-Gaussian error distribution, which was especially useful for
312 count data in our case (i.e. number of individuals, dung-dwellers and soil-diggers). We
313 systematically corrected for Poisson overdispersion using a negative binomial family
314 (Gardner et al., 1995). We used the R packages “stats” (using the *lm* and *glm* functions) and
315 “MASS” (Venables and Ripley, 2002) (using the *glm.nb* function) to fit the models.

316 2.6. Indicator species analysis

317 To identify indicator species for grazing intensity, we calculated the IndVal index (Dufrêne
318 and Legendre, 1997). The IndVal combines the degree of specificity (highest when the

319 species is present in a given habitat type but not elsewhere) and fidelity (highest when the
320 species is present in all sites of a given habitat type). The closer the IndVal value is to 1, the
321 higher the specificity and/or fidelity of a species to a given level or pair of levels of grazing
322 intensity. We calculated this index on non-transformed species abundance matrices, using
323 pitfall traps as sampling units, to individual levels of grazing intensity and their combinations
324 (i.e. LGI/MGI and MGI/HGI) based on the approach by De Cáceres and Legendre (2009). We
325 ran the analysis separately for each studied area (steppe and Alps), and assessed the
326 statistical significance of the IndVal values by means of a permutation test with a 5%
327 rejection threshold (Dufrêne and Legendre, 1997). We associated each selected indicator
328 species with its species-specific nesting guild and mean body mass. The latter was measured
329 by randomly selecting between 4 and 10 individuals per species (depending on the number
330 available) and weighing them after drying them at 70 °C for 24 h. We used this information
331 to improve the ecological interpretation of the results and make it more accessible for non-
332 specialists of dung beetles. We used the R package “indicspecies” (using the *multipatt*
333 function) to find indicator species for different grazing intensity levels (De Cáceres and
334 Legendre, 2009).

335

336 **3. Results**

337 *3.1. Dung beetle inventories*

338 We collected a total of 11,733 dung beetles belonging to 52 species (see **Table 1** and
339 **Appendix D** for the complete list of the sampled species). While species richness was higher
340 in the steppe, the number of individual beetles captured was seven times higher in the Alps
341 (**Table 1**). The species composition in the two study areas was very different, with only four

342 species in common (*Aphodius cardinalis*, *Calamosternus granarius*, *Coloboferus erraticus*
343 and *Otophorus haemorrhoidalis*; **Appendix D**). The five most abundant species were *C.*
344 *granarius*, *C. erraticus*, *Euorodalus paracoenosus*, *Onthophagus ruficapillus* and *O. vacca* in
345 the steppe; and *Amidorus obscurus*, *Euheptaulacus carinatus*, *Oromus alpinus*, *Onthophagus*
346 *fracticornis* and *Parammoecius corvinus* in the Alps (**Appendix D**). In the Alps, we collected
347 four large species of the Geotrupinae subfamily (*Geotrupes stercorarius*, *Anoplotrupes*
348 *stercorosus*, *Trypocopris vernalis* and *T. alpinus*), which are absent in the steppe ecosystem.
349 The iChao 2 and ICE-1 species richness values show that our sampling was relatively
350 exhaustive, representing 75–98% of the estimated species richness in the study area
351 considered (**Table 1**).

352 3.2. Patterns in species assemblage composition

353 For all the subsequent analyses, we removed 9 singleton species, retaining 43 species. In
354 both study areas, the RDA-based MANOVAs showed a significant effect on dung beetle
355 species assemblages of “pasture identity” (steppe, $F_{1, 28} = 7.67$, $p = 0.001$; Alps, $F_{1, 28} = 60.75$,
356 $p = 0.001$) and of “grazing intensity” (steppe, $F_{2, 27} = 22.10$, $p = 0.001$; Alps, $F_{2, 27} = 42.85$, $p =$
357 0.001). The interaction between these two factors was also significant in the steppe ($F_{2, 27} =$
358 4.46 , $p = 0.001$) and the Alps ($F_{2, 27} = 29.93$, $p = 0.001$), meaning that the effect of grazing
359 intensity on species assemblages differed significantly according to the pasture considered.
360 Thus, we ran RDAs constrained by the grazing intensity factor and plotted the results
361 (triplots) separately for each pasture in each study area. In the steppe, grazing intensity
362 explained 74.8% and 58.7% of the total variation in species assemblages respectively in
363 pastures A and B (**Figure 2**). Most of the variation was explained by the first axis of each
364 RDA, which gathered 78–93% of the variation explained by the constraining factor. In both

365 steppe pastures, the first axis of the triplots clearly differentiated the two endpoints of the
366 grazing intensity gradient, showing the HGI plots on the left, and the MGI and LGI plots on
367 the right of the ordination planes. In steppe pasture A, species assemblages of the HGI plot
368 were more distinct from the other plots than in pasture B. Moreover, the MGI and LGI plots
369 were not particularly contrasted along the second axis, which explained a low percentage of
370 variation (5%). In pasture B, the second axis explained a larger percentage of variation (13%)
371 and showed contrast between the MGI and LGI plots.

372 In the Alps, grazing intensity explained 89.9% and 82.9% of the total variation in species
373 assemblages respectively in pastures C and D (**Figure 3**). As in the steppe, most of the
374 variation was explained by the first RDA axis (70–80%), which clearly differentiated the LGI
375 plots on the left and HGI plots on the right of the ordination planes. The second axis of each
376 RDA triplot differentiated the MGI plots from the other grazing intensity levels.

377 A comparative analysis of the results indicated that (i) the grazing intensity gradient
378 structured the dung beetle species assemblages in all the studied pastures; (ii) in each
379 studied area, the level of differentiation between assemblages along the gradient was
380 dependent on the pasture considered; and (iii) species assemblages showed more contrast
381 due to grazing intensity in the Alps than in the steppe.

382 *3.3. Number of individuals, rarefied species richness and Pielou's evenness according* 383 *to grazing intensity*

384 When the "*pasture identity*" factor was held constant, "*grazing intensity*" had differing
385 effects on the response variables reflecting dung beetle community structure. In the steppe,
386 the number of captured individuals and the rarefied species richness did not show any
387 significant response to grazing intensity (**Figure 4A**). Pielou's evenness was higher in HGI

388 plots compared to LGI plots, but values of MGI plots were no different from the other two
389 levels of grazing intensity. In the Alps, the number of captured individuals was higher in MGI
390 and HGI plots and lower in LGI plots (**Figure 4B**). The rarefied species richness was higher in
391 LGI plots than in MGI and HGI plots. An inverse relationship was observed for Pielou's
392 evenness, which progressively and significantly increased with grazing intensity.

393 When the "*grazing intensity*" factor was held constant, there were no significant differences
394 in the number of captured individuals, rarefied species richness or Pielou's evenness
395 between the two sampled pastures in the steppe (**Appendix E, Figure E.1**). In the Alps, there
396 were significantly more individuals sampled in pasture C, whereas the rarefied species
397 richness was higher in pasture D (**Appendix E, Figure E.1**).

398 *3.4. Nesting guild abundance according to grazing intensity*

399 When the "*pasture identity*" factor was held constant, "*grazing intensity*" had similar effects
400 on the nesting guild abundance in the two study areas. The results of the models showed
401 that, in both study areas, the abundance of dung-dwellers significantly increased between
402 LGI and HGI plots (**Figure 5**). The MGI plots occupied an intermediate position between the
403 two other levels in the steppe, but MGI and HGI plots did not differ from each other in the
404 Alps. The abundance of soil-diggers significantly decreased between LGI and HGI plots in the
405 Alps, whereas in the steppe, no significant variation was observed (**Figure 5**). However,
406 because the p value between MGI and HGI plots was close to 0.05 (0.0573), we cannot
407 completely exclude a possible effect of grazing, and thus a tendency towards a decrease in
408 soil-diggers between moderate and high grazing intensity zones (Amrhein, Greenland, &
409 McShane, 2019).

410 When the “grazing identity” factor was held constant, there were more dung-dwellers
411 caught in pasture B than in pasture A in the steppe, while there was no difference in the
412 number of soil-diggers (**Appendix E, Figure E.2**). In the Alps, there were significantly more
413 dung-dwellers and soil-diggers captured in pasture C than in pasture D (**Appendix E, Figure**
414 **E.2**). This pattern can be explained by the greater overall abundance observed in pasture C
415 (**Appendix E, Figure E.1**).

416 3.5. Indicator species

417 In the steppe, half of the species collected (excluding singletons) were retained as indicators
418 of grazing intensity (**Table 2**). Six species were indicators for individual grazing intensity
419 levels, and six others for combined levels. *Copris hispanus hispanus*, by far the largest beetle
420 of the 12 indicator species identified, was the only indicator species for LGI plots. We found
421 that the body mass of indicator species belonging to the soil-digger guild decreased between
422 LGI/MGI plots and MGI/HGI plots. Including both nesting guilds, indicator species for the
423 LGI/MGI combination covered a relatively large range of body mass values, from 2.8 ± 0.8 g
424 to 67.7 ± 13 g. Only soil-diggers were associated with MGI plots and the MGI/HGI
425 combination, while indicators for HGI plots were mostly dung-dwellers.

426 Of the 22 species collected in the Alps (excluding singletons), only 4 (18%) were selected as
427 indicators of grazing intensity (**Table 2**). In this area, we only found species indicators for LGI
428 plots and the combination MGI/HGI plots. Two large soil-digging species (*Geotrupes*
429 *stercorarius* and *Trypocopris vernalis*) were associated with low grazing intensity, while the
430 only species associated with MGI/HGI plots was a small dung-dweller (*Oromus alpinus*), 30 to
431 50 times lighter. One species, *Otophorus haemorrhoidalis*, collected in both study areas, was

432 found to be an indicator for LGI plots only in the Alps. The position of some of the indicator
433 species found is shown on the ordination planes of the RDAs in **Figures 2 and 3**.

434

435 **Discussion**

436 Overall, our results show that the intra-pasture variation in grazing intensity exerted by
437 domestic ungulates strongly structured dung beetle species assemblages. Beyond the effects
438 of the bioclimatic context and the local pasture conditions, the dung beetle communities
439 showed strong response patterns to variation in grazing intensity in the two study areas.
440 Whereas species assemblages were more contrasted along the grazing intensity gradient in
441 the alpine pastures, the observed differences reflected substantial variation in nesting guild
442 abundance in both areas. This result is all the more significant since less than 10% of the
443 species were common between the steppe and the Alps. Furthermore, taking nesting guild
444 into account offers deeper insights about the effects of grazing on dung beetle assemblages.

445 *4.1. Sampling effectiveness*

446 The iChao2 and ICE-1 estimates showed that by sampling during the peak of dung beetle
447 activity we were able to obtain a quite exhaustive representation of local species
448 assemblages inhabiting the studied pastures. According to Lobo et al. (2006), as the
449 emergence of dung beetles during their maximum activity period depends on the habitat
450 conditions of the previous year at the same period, we assumed that targeting these activity
451 peaks would be relevant for investigating the effects of various grazing conditions. In the
452 steppe, we collected 64% of the total dung beetle species richness known to occur in the
453 area, based on inventories conducted in the Coussouls de Crau National Nature Reserve over

454 five years of successive sampling in eight different pastures during spring and autumn (Tatin
455 et al., 2014 and unpublished results). In the Alps, we recorded 70% of the total species
456 richness previously inventoried in the total area of the Vanoise National Park based on
457 intensive sampling carried out across all seasons (Lumaret and Stiernet, 1991). Previous large
458 surveys conducted in France both in Mediterranean ecosystems (Jay-Robert et al., 2008a)
459 and in the western Alps (Jay-Robert et al., 2008b) have shown that the seasonal activity of
460 Scarabaeinae species is mainly restricted to spring/early summer (depending on elevation)
461 and that only Aphodiinae species were temporally segregated, with an activity period from
462 early spring to late autumn (and even winter at the lowest elevations). Consequently,
463 sampling in spring/summer was effective for studying diverse assemblages composed of a
464 mixture of coexisting dung-dweller and soil-digger species.

465 *4.2. Intra-pasture variation in grazing intensity shapes the structure and composition* 466 *of dung beetle assemblages*

467 In central Italy, Tonelli et al. (2017) previously found significant contrasts in the composition
468 of dung beetle assemblages between pastures subjected to low and moderate grazing
469 intensity. Our results reveal that such differences also exist within pastures. This
470 corroborates studies showing that differences in the composition and structure of terrestrial
471 arthropod assemblages often reflect fine-scale habitat heterogeneity (Gonçalves-Souza et
472 al., 2015; Weiss et al., 2016). In our case, as the vegetation in the different GI zones was
473 similar (grasslands without shrubs, see “Material & methods”), we suggest that fine-scale
474 variation in habitat conditions in terms of droppings availability and disturbance intensity
475 (especially trampling) might be the main cause of the observed changes. The difference in
476 species composition between pastures might explain the significant interaction between

477 pasture identity and grazing intensity. However, the similar patterns showed by the
478 multivariate analyses between the steppe and the Alps suggest that the grazing intensity
479 gradient had a similar effect whatever the bioclimatic context. In each pasture, the
480 multivariate analyses show that species assemblages were significantly different across the
481 three levels of the grazing intensity gradient. In addition, the RDA-based MANOVAs show
482 that the sampled zones that were spatially closest (**Appendix B**) were clearly distinguished
483 from one another. Dung beetles are known to have a high dispersal capacity, necessary to
484 find ephemeral and patchily distributed dung pads (Roslin and Viljanen, 2011). They are able
485 to forage over hundreds of metres (Roslin, 2000), across distances exceeding those
486 separating our sampling plots. However, despite the capacity of the observed species to
487 colonize the spatially closest sampled zones, thus possibly inducing positive spatial
488 autocorrelation between them, the observed patterns show a greater imprint of the local
489 grazing conditions.

490 The increased droppings availability in moderately and highly grazed zones of pastures
491 allowed more individuals to coexist in the Alps. This corroborates the results of Lumaret et
492 al. (1992) showing that the abundance and biomass of dung beetles strongly depend on the
493 quantity of trophic resources. If ungulate grazing is considered a disturbance process (Hobbs,
494 2006), one might expect that an increase in grazing intensity would promote the coexistence
495 of a few well-adapted and numerically dominant species, with the other species remaining in
496 low numbers (Grime, 1973), resulting in decreasing evenness. Surprisingly, we observed that
497 the evenness of species assemblages significantly increased along the grazing intensity
498 gradient in both study areas. This result may be explained by a level of disturbance resulting
499 from grazing that was still not strong enough, even in the most intensively grazed zones
500 sampled. However, resource scarcity favoured the dominance of a lower number of species

501 in the least intensively grazed zones, which is likely to explain the decrease in species
502 evenness. Theoretically, the most competitive and/or efficient species are more suited for
503 exploiting limited resources (Connell, 1978). In our surveys, two dung beetles belonging to
504 the *Onthophagus* genus were particularly abundant in the least grazed habitats: *O. vacca*
505 accounted for almost 70% of the abundance in LGI plots in the steppe, and *O. fracticornis*
506 represented half of the abundance in LGI plots in the Alps (**Appendix D**). In central Italy,
507 Tonelli et al. (2018) obtained similar results, finding that *Onthophagus* were among the most
508 common species in an abandoned pasture. Taken together, these findings suggest that
509 *Onthophagus* species are efficient foragers, allowing them to maintain their population in
510 habitats where dung is relatively scarce, such as in the least grazed zones of pastures.

511 Although both study areas shared many similar results, the species assemblages observed in
512 the Alps were more differentiated along the grazing intensity gradient than in the steppe,
513 reflecting differences in livestock management. While herd sizes were similar in both study
514 areas, sheep density was more than twice as high in the Alps as in the steppe (11.8 vs 4.7
515 sheep/ha; **Appendix A**). As higher livestock density results in higher droppings input and soil
516 trampling, this is likely to increase habitat heterogeneity within the pasture, leading to
517 contrasting habitat conditions between moderate and low grazing zones, and thus increasing
518 the differences in dung beetle assemblages. Hence, the lower livestock density in the steppe
519 is likely to explain that dung beetle assemblages inhabiting low and moderate grazing zones
520 were more similar to each other than they were to assemblages in high grazing zones.

521 Previous studies have reported a decrease in observed species richness with an increase in
522 grazing intensity for different arthropod taxa such as grasshoppers, butterflies,
523 hymenopterans and moths, mainly due to homogenization of vegetation structure and

524 diversity, but also to changes in trophic interactions (Enkhtur et al., 2017; Kruess and
525 Tschardtke, 2002). On this question, we found contrasting results between the two study
526 areas. While in the Alps rarefied species richness decreased with an increase in grazing
527 intensity in moderate and high grazing zones of pastures, in the steppe, the turnover in
528 species composition along the grazing intensity gradient did not lead to significant changes
529 in rarefied species richness. Several processes may explain the differences we observed
530 between the two study areas. Of these, specific historical anthropogenic processes (i.e. the
531 history of local grazing practices) and abiotic constraints (e.g. climate) are very likely to
532 separately drive the establishment of current dung beetle species assemblages and the
533 interspecific interactions that take place in the steppe and the Alps. In the Cévennes
534 National Park, an intermediate location between the steppe and the Alps (44°20'N, 3°50'E;
535 800 m a.s.l.), Jay-Robert et al. (2008c) observed higher species richness and temporal
536 turnover in the dung beetle assemblages of grazed shrubland than in those of grazed
537 grasslands, where dung resources were twice as high. In this region of low mountains, dung
538 beetle diversity was thus more limited by the homogeneous structure of the herbaceous
539 habitat than by resource availability. Considering that short vegetation covers the steppe
540 whatever the grazing intensity (Devaux et al., 1983), one may hypothesize that habitat
541 heterogeneity was too low to make a difference to the number of coexisting species
542 throughout the grazing intensity gradient.

543 *4.3. Nesting guilds reflect the effects of grazing intensity*

544 Differences in dung beetle nesting guild abundance in abandoned and moderately grazed
545 pastures have been reported in southern Europe (Tonelli et al., 2019). Our findings showed
546 that such differences also occur in response to fine-scale variation in grazing intensity within

547 pastures. The shifts observed in nesting guild abundance support the previously discussed
548 differences in the overall structure and composition of dung beetle assemblages along the
549 grazing intensity gradient. The similarity in the trends observed between the steppe and the
550 Alps indicates that the filtering effect exerted by grazing pressure selects species according
551 to particular traits, allowing them to persist at different levels of grazing intensity.

552 Dung-dwellers were significantly more abundant in zones of high and moderate grazing
553 intensity than in those of low grazing intensity, whereas the abundance of soil-diggers
554 tended to decrease when grazing intensity increased. In overgrazed habitats, vegetation is
555 permanently short. Hard trampling increases soil compactness and creates patches of bare
556 soil, leading to a warmer underground microclimate and promoting water evaporation
557 (Greenwood and McKenzie, 2001; Lu et al., 2017; Yan et al., 2018; Zhou et al., 2010).
558 Significant changes in soil moisture and physical conditions may compromise the ability of
559 some soil-digger species to dig efficiently (Dabrowski et al., 2019) or the suitability of the
560 underground habitat for their larvae (Sowig, 1996, 1995). Hutton and Giller (2003) and
561 Negro et al. (2011) also observed a decrease in the abundance of soil-diggers in overgrazed
562 and intensively managed pastures in lowland Ireland and the Italian Alps, respectively.
563 Because of their autecological requirements, dung-dwellers might be less sensitive to soil
564 properties than soil-diggers; however, the quantity of available resources is an important
565 factor driving their abundance (Lumaret et al., 1992). Their dwelling behaviour does not
566 involve food relocation, so the maintenance of local populations depends solely on the
567 presence of dung and the preservation of this resource during larval development (Gittings
568 and Giller, 1999, 1997).

569 Typically, dung pads are an ephemeral and patchily distributed resource. Several studies
570 support the hypothesis that strong competition for local resources occurs both between
571 coexisting individuals and dung beetle species (Finn and Gittings, 2003; Giller and Doube,
572 1994, 1989; Gittings and Giller, 1999). In addition, sheep droppings are a relatively scarce
573 resource for dung beetles, occurring in much smaller quantities and drying out rapidly
574 compared to cattle dung, which can gather hundreds of individuals (Finn and Gittings, 2003;
575 Lumaret, 1995; Lumaret et al., 1992). Soil-diggers are usually known to be faster at resource
576 acquisition than dung-dwellers (Hanski and Cambefort, 1991), which could explain their
577 dominance in the least grazed parts of pastures. Highly abundant at low grazing intensity,
578 the previously mentioned *Onthophagus* species are soil-diggers that build ramified tunnels
579 below droppings or dung pads. The foraging efficiency of soil-diggers allows them to rapidly
580 access and secure the few resources available, conferring an advantage in resource-scarce
581 environments. Because of their high competitiveness in resource acquisition, an increase in
582 soil-digger abundance is likely to lead to more competitive interactions (Connell, 1978) and
583 to prevent the installation of dung-dwellers. In the Alps, the largest soil-diggers we collected
584 – *Geotrupes stercorarius* and *Anoplotrupes stercorosus* (Geotrupinae) – were much more
585 abundant in low grazing zones (**Appendix D**). Geotrupinae species are fast burrowers able to
586 build deep tunnels filled with large faecal masses in only 24 hours (Klemperer, 1979).
587 Gittings and Giller (1999) showed that, by rapidly removing dung pads from the surface of
588 the ground, Geotrupinae abundance can negatively impact the reproductive success of
589 dung-dwellers. This intense exploitative competition may be one of the factors explaining
590 the decrease of dung-dwellers we observed in the least grazed parts of pastures.

591 For arthropods, the heterogeneity in vegetation structure and abiotic conditions created by
592 a moderate level of grazing often promotes more diverse assemblages than at low and high

593 grazing levels, favouring the occurrence of species with different habitat requirements (van
594 Klink et al., 2015). Our results partially support this general finding: although species richness
595 did not peak at intermediate grazing intensity, both dung-dwellers and soil-diggers were
596 found in high abundance in moderate grazing zones of pastures. Moderate sheep grazing
597 results in a patchy distribution of droppings and heterogeneous soil conditions and
598 vegetation cover, creating a variety of habitats suitable for species with diverse nesting
599 strategies. With a moderate level of habitat disturbance, one could hypothesize that a
600 balance exists between competitive exclusion and loss of competitive dominants through
601 disturbance, these conditions favouring the coexistence of competitive species (soil-diggers)
602 and disturbance-tolerant species (dung-dwellers) (Connell, 1978; Mackey and Currie, 2001).

603 *4.4. Can dung beetles be relevant indicators of grazing intensity effects?*

604 As expected, given the strong compositional difference in dung beetle assemblages between
605 the steppe and the Alps, we found different indicator species in each study area.
606 Nevertheless, the body mass and nesting behaviour of indicator species highlighted a similar
607 trend: irrespective of species identity, the largest soil-digger indicator species were
608 associated with relatively low grazing intensity, while the indicator species for higher grazing
609 intensity were mostly either small soil-diggers or small dung-dwellers. The indicator species
610 for moderate grazing levels covered a large range of body mass values and belonged to both
611 nesting guilds. This pattern was especially well defined in the steppe since soil-diggers were
612 almost twice as diverse in the Mediterranean lowland as in the alpine highland (14 vs 8 soil-
613 digger species in the steppe and in the Alps, respectively; Jay-Robert et al., 1997).

614 Generally, the smallest dung beetle species are known to be more fecund than the largest
615 (Halffter and Edmonds, 1982; Hanski and Cambefort, 1991). Compared to soil-diggers, dung-

616 dwellers show higher fecundity and ovipositing rates (Halffter and Edmonds, 1982; Hanski
617 and Cambefort, 1991; Gittings and Giller, 1997). In Scarabaeinae soil-diggers, the largest
618 species are usually less fecund, compensated by a high level of parental care. In the steppe,
619 the largest soil-digger indicator species, *Copris hispanus hispanus* and *Gymnopleurus*
620 *flagellatus flagellatus*, were indicators of low and low/moderate grazing intensity,
621 respectively. They have a lower reproductive rate compared to the smaller soil-diggers
622 (*Onthophagus* species; Halffter and Edmonds, 1982) that we found as indicators of higher
623 grazing intensities. It is possible that these large soil-diggers are less able to produce a
624 replacement clutch if repeated trampling destroys the previous one (as Simons et al. (2016)
625 concluded about the response of various arthropods to flooding intensity). Additionally, the
626 largest species are known to forage across larger areas than the smallest species, thus
627 suggesting greater dispersal capacity for large dung beetles (Peck and Howden, 1984; Roslin,
628 2000, 2001). This may confer these beetles with a greater ability to maintain their
629 populations in the least grazed parts of pastures where resource availability is low.
630 Nevertheless, there are very few quantitative descriptions of dung beetle dispersal and
631 foraging behaviour, preventing detailed hypotheses. As we did not analyse body-size
632 patterns exactly along the grazing intensity gradient, these interpretations should be taken
633 with caution. Further studies are needed to more accurately determine the relationship
634 between the biology of dung beetle species and their ability to persist subject to distinct
635 levels of grazing.

636 Other studies have previously identified several of our indicator species as characteristic of
637 comparable habitat conditions in Europe. In Ireland, Hutton and Giller (2003) found that
638 *Geotrupes stercorarius*, one of the largest soil-diggers occurring in northern temperate dung
639 beetle communities, was significantly more abundant in extensively managed pastures (on

640 organic farms). In the French Alps, we found this species was an indicator for the least
641 grazed zones of pastures. We can hypothesize that *Geotrupes stercorarius* is able to maintain
642 its population in resource-scarce environments thanks to its foraging efficiency, avoiding the
643 most trampled habitats with high soil disturbance. In the Mediterranean region, (J.-P.
644 Lumaret & Iborra, 1996) found high numbers of *Aphodius foetidus* and *Calamosternus*
645 *granarius* inhabiting dropping accumulations where sheep rest. We corroborated this
646 observation in the steppe, where we found that these two species were indicators for the
647 most grazed zones of pastures. *Calamosternus granarius*, which is a cosmopolitan species
648 originating from the Palearctic region, ranks among the smallest of the dung-dwellers in
649 Europe and is considered a generalist feeder (Gittings and Giller, 1997). Brown (1940), Sears
650 (1978) and Ratcliffe (1991) showed that this species feeds on a variety of faeces, carrion and
651 compost material. The generalist diet of *Calamosternus granarius* is likely to allow
652 individuals to consume and to breed on the accumulated manure produced by livestock in
653 overgrazed areas.

654 Given the wide distribution of *Geotrupes stercorarius* (introduced in North America; Brown,
655 1940), *Calamosternus granarius* (introduced in North America, Australia and the Afro-
656 tropical region; de Jong et al., 2014) and *Aphodius foetidus* (widely distributed in Western
657 Europe; Lumaret, 1990), we would recommend extending their use as indicators of grazing
658 practices to other regions. However, species found to be indicators in a given area may not
659 be transposable elsewhere because of changes in species pool composition across distinct
660 bioclimatic contexts. Practically speaking, the use of a set of indicator species may thus be
661 restricted to a particular locality or region (Zettler et al., 2013), so our IndVal results are
662 likely difficult to generalize if only species identity is considered. Integrating species traits (as
663 we did with nesting guild and body mass) improves the ecological interpretation of IndVal

664 results. This suggests a way to bypass dung beetle taxonomy, which requires specialist
665 knowledge. Dung beetles collected in the field can be easily classified as dung-dwellers or
666 soil-diggers in different weight classes (or size, which is easier to measure and correlates
667 with body mass), avoiding the need for difficult species identification training. The use of
668 body size and nesting guilds as indicator measures would be an easier method for
669 conservation practitioners, improving the use of dung beetles as indicators for impacts of
670 local grazing practices. To increase the scope of our results and the usability of this method,
671 we recommend comparing findings from studies carried out in different bioclimatic contexts
672 and across a higher number of pastures.

673

674 **Conclusion**

675 Working at an intra-pasture scale and selecting sheep-grazed pastures with similar grazing
676 histories, we were able to analyse the processes by which the pressure of domestic
677 ungulates can substantially modify the composition and structure of dung beetle
678 assemblages. Our results suggest that, by altering habitat conditions, changing droppings
679 availability and modifying competitive interactions, fine-scale variation in grazing intensity
680 leads to substantial variations in nesting guild abundance. High grazing intensity is
681 detrimental to soil-diggers and the largest dung beetle species. Whereas some of these
682 species (e.g. *Gymnopleurus flagellatus flagellatus*) have historically declined in Western
683 Europe due to a progressive decrease in extensive grazing practices (Caillol, 1908; Carpaneto
684 et al., 2007; Lobo, 2001), local intensification of livestock farming may not compensate for
685 grazing abandonment on surrounding lands. Conversely, small dung-dwellers are favoured in
686 highly grazed habitats, where sheep droppings are predictable and highly available. These

687 results show that grazing intensity acts as an environmental filter on dung beetle
688 assemblages, selecting species according to particular traits. In order to better understand
689 the mechanisms by which grazing practices influence the structure of these assemblages,
690 further studies could integrate life-history trait approaches (e.g. Chillo et al., 2017).

691

692 **Acknowledgements**

693 William Perrin received a PhD grant from University Paul-Valéry Montpellier 3. The project
694 also received financial support from the Vanoise National Park. The authors are very grateful
695 to the staff at the Provence-Alpes-Côtes-d'Azur Conservatory of Natural Areas and the
696 Vanoise National Park, especially Laurent Tatin, Fanny Sauguet, Guy-Noël Grosset and Franck
697 Parchoux, for their cooperation and assistance in this study. We would also like to thank the
698 shepherds and herders for their cooperation; this study could not have been carried out
699 without them. The comments of the two anonymous reviewers were also very valuable and
700 contributed to improving the manuscript.

701

702 **Appendices**

703 **Appendix A.** General information for each pasture in each study area regarding: grazing
704 practices (historical and present) and environmental conditions (vegetation, soil type,
705 climate).

706 **Appendix B.** Distances between the different GI zones in each pasture and in each study
707 area.

708 **Appendix C.** Characterization of the grazing intensity gradient.

709 **Appendix D.** Abundance of dung beetle species collected in the two study areas, the steppe
710 and the Alps.

711 **Appendix E.** Effect of “*pasture identity*” on dung beetle assemblages.

712

713 **References**

714 Alkemade, R., Reid, R.S., Berg, M. van den, Leeuw, J. de, Jeuken, M., 2013. Assessing the
715 impacts of livestock production on biodiversity in rangeland ecosystems. PNAS 110,
716 20900–20905. <https://doi.org/10.1073/pnas.1011013108>

717 Amrhein, V., Greenland, S., McShane, B., 2019. Scientists rise up against statistical
718 significance. Nature 567, 305–307. <https://doi.org/10.1038/d41586-019-00857-9>

719 Anderson, M., Braak, C.T., 2003. Permutation tests for multi-factorial analysis of variance. J.
720 Stat. Comput. Simul. 73, 85–113. <https://doi.org/10.1080/00949650215733>

721 Anderson, M.J., 2006. Distance-based tests for homogeneity of multivariate dispersions.
722 Biometrics 62, 245–253. <https://doi.org/10.1111/j.1541-0420.2005.00440.x>

723 Audino, L.D., Louzada, J., Comita, L., 2014. Dung beetles as indicators of tropical forest
724 restoration success: Is it possible to recover species and functional diversity? Biol.
725 Conserv. 169, 248–257. <https://doi.org/10.1016/j.biocon.2013.11.023>

726 Badan, O., Congès, G., Brun, J.-P., 1995. Les bergeries romaines de la Crau d’Arles. Les
727 origines de la transhumance en Provence. Gallia 52, 263–310.
728 <https://doi.org/10.3406/galia.1995.3152>

729 Báldi, A., Batáry, P., Kleijn, D., 2013. Effects of grazing and biogeographic regions on
730 grassland biodiversity in Hungary – analysing assemblages of 1200 species. *Agric.*
731 *Ecosyst. Environ.* 28–34. <https://doi.org/10.1016/j.agee.2012.03.005>

732 Barragán, F., Moreno, C.E., Escobar, F., Bueno-Villegas, J., Halffter, G., 2014. The impact of
733 grazing on dung beetle diversity depends on both biogeographical and ecological
734 context. *J. Biogeogr.* 41, 1991–2002. <https://doi.org/10.1111/jbi.12351>

735 Batáry, P., Báldi, A., Sárospataki, M., Kohler, F., Verhulst, J., Knop, E., Herzog, F., Kleijn, D.,
736 2010. Effect of conservation management on bees and insect-pollinated grassland plant
737 communities in three European countries. *Agric. Ecosyst. Environ.* 136, 35–39.
738 <https://doi.org/10.1016/j.agee.2009.11.004>

739 Batáry, P., Báldi, A., Szél, G., Podlussány, A., Rozner, I., Erdős, S., 2007. Responses of
740 grassland specialist and generalist beetles to management and landscape complexity.
741 *Divers. Distrib.* 13, 196–202. <https://doi.org/10.1111/j.1472-4642.2006.00309.x>

742 Beynon, S.A., Peck, M., Mann, D.J., Lewis, O.T., 2012. Consequences of alternative and
743 conventional endoparasite control in cattle for dung-associated invertebrates and
744 ecosystem functioning. *Agric. Ecosyst. Environ.* 162, 36–44.
745 <https://doi.org/10.1016/j.agee.2012.08.010>

746 Bicknell, J.E., Phelps, S.P., Davies, R.G., Mann, D.J., Struebig, M.J., Davies, Z.G., 2014. Dung
747 beetles as indicators for rapid impact assessments: Evaluating best practice forestry in
748 the neotropics. *Ecol. Indic.* 43, 154–161. <https://doi.org/10.1016/j.ecolind.2014.02.030>

749 Borcard, D., Gillet, F., Legendre, P., 2018. *Numerical Ecology with R*, 2nd ed, Use R! Springer
750 International Publishing.

751 Bradshaw, R.H.W., Hannon, G.E., Lister, A.M., 2003. A long-term perspective on ungulate–
752 vegetation interactions. *For. Ecol. Manag.* 181, 267–280.
753 [https://doi.org/10.1016/S0378-1127\(03\)00138-5](https://doi.org/10.1016/S0378-1127(03)00138-5)

754 Brown, W.J., 1940. NOTES ON THE AMERICAN DISTRIBUTION OF SOME SPECIES OF
755 COLEOPTERA COMMON TO THE EUROPEAN AND NORTH AMERICAN CONTINENTS*.
756 *Can. Entomol.* 72, 65–78. <https://doi.org/10.4039/Ent7265-4>

757 Buse, J., Šlachta, M., Sladeczek, F.X.J., Pung, M., Wagner, T., Entling, M.H., 2015. Relative
758 importance of pasture size and grazing continuity for the long-term conservation of
759 European dung beetles. *Biol. Conserva.* 187, 112–119.
760 <https://doi.org/10.1016/j.biocon.2015.04.011>

761 Caillol, H., 1908. Catalogue des Coléoptères de Provence, 2e Partie. Société des Sciences
762 Naturelles de Provence, Marseille, 359-456.

763 Campos, R.C., Hernández, M.I.M., 2015. Changes in the dynamics of functional groups in
764 communities of dung beetles in Atlantic forest fragments adjacent to transgenic maize
765 crops. *Ecol. Indic.* 49, 216–227. <https://doi.org/10.1016/j.ecolind.2014.09.043>

766 Carpaneto, G.M., Mazziotta, A., Valerio, L., 2007. Inferring species decline from collection
767 records: roller dung beetles in Italy (Coleoptera, Scarabaeidae). *Divers. Distrib.* 13, 903–
768 919. <https://doi.org/10.1111/j.1472-4642.2007.00397.x>

769 Chao, A., 1987. Estimating the Population Size for Capture-Recapture Data with Unequal
770 Catchability. *Biometrics* 43, 783–791. <https://doi.org/10.2307/2531532>

771 Chao, A., Ma, K.H., Hsieh, T. C., 2015. SpadeR: Species Prediction and Diversity
772 Estimation with R. R package version 0.1.0. URL
773 <http://chao.stat.nthu.edu.tw/blog/software-download/>

774 Chao, A., Chazdon, R.L., Colwell, R.K., Shen, T.-J., 2005. A new statistical approach for
775 assessing similarity of species composition with incidence and abundance data. *Ecol.*
776 *Lett.* 8, 148–159. <https://doi.org/10.1111/j.1461-0248.2004.00707.x>

777 Chillo, V., Ojeda, R.A., Capmourteres, V., Anand, M., 2017. Functional diversity loss with
778 increasing livestock grazing intensity in drylands: the mechanisms and their
779 consequences depend on the taxa. *J. Appl. Ecol.* 54, 986–996.
780 <https://doi.org/10.1111/1365-2664.12775>

781 Chiu, C.-H., Wang, Y.-T., Walther, B.A., Chao, A., 2014. An improved nonparametric lower
782 bound of species richness via a modified good–turing frequency formula. *Biometrics* 70,
783 671–682. <https://doi.org/10.1111/biom.12200>

784 Connell, J.H., 1978. Diversity in Tropical Rain Forests and Coral Reefs. *Science* 199, 1302–
785 1310. <https://doi.org/10.1126/science.199.4335.1302>

786 Dabrowski, J., Venter, G., Truter, W.F., Scholtz, C.H., 2019. Dung beetles can tunnel into
787 highly compacted soils from reclaimed mined sites in eMalahleni, South Africa. *Appl. Soil*
788 *Ecol.* 134, 116–119. <https://doi.org/10.1016/j.apsoil.2018.10.015>

789 De Cáceres, M.D., Legendre, P., 2009. Associations between species and groups of sites:
790 indices and statistical inference. *Ecology* 90, 3566–3574. [https://doi.org/10.1890/08-](https://doi.org/10.1890/08-1823.1)
791 1823.1

792 Devaux, J.P., Archiloque, A., Borel, L., Bourrelly, M., Palluel, J.L., 1983. Notice de la carte
793 phytosociologique de la Crau. *Biologie-Ecologie méditerranéenne* 10, 5-54.

794 Dufrêne, M., Legendre, P., 1997. Species Assemblages and Indicator Species: the Need for a
795 Flexible Asymmetrical Approach. *Ecol. Monogr.* 67, 345–366.
796 [https://doi.org/10.1890/0012-9615\(1997\)067\[0345:SAAI\]2.0.CO;2](https://doi.org/10.1890/0012-9615(1997)067[0345:SAAI]2.0.CO;2)

797 Enkhtur, K., Pfeiffer, M., Lkhagva, A., Boldgiv, B., 2017. Response of moths (Lepidoptera:
798 Heterocera) to livestock grazing in Mongolian rangelands. *Ecol. Indic.* 72, 667–674.
799 <https://doi.org/10.1016/j.ecolind.2016.08.053>

800 European Commission, 2015. Statistiques sur la structure des exploitations agricoles.
801 Available from: [https://ec.europa.eu/eurostat/statistics-](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Farm_structure_statistics/fr#Informations_suppl.C3.A9mentaires_Eurostat)
802 [explained/index.php?title=Farm_structure_statistics/fr#Informations_suppl.C3.A9ment](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Farm_structure_statistics/fr#Informations_suppl.C3.A9mentaires_Eurostat)
803 [aires_Eurostat](https://ec.europa.eu/eurostat/statistics-explained/index.php?title=Farm_structure_statistics/fr#Informations_suppl.C3.A9mentaires_Eurostat). (accessed 01 January 2019).

804 Finn, J.A., Giller, P.S., 2000. Patch Size and Colonisation Patterns: An Experimental Analysis
805 Using North Temperate Coprophagous Dung Beetles. *Ecography* 23, 315–327.

806 Finn, J.A., Gittings, T., 2003. A review of competition in north temperate dung beetle
807 communities. *Ecol. Entomol.* 28, 1–13. [https://doi.org/10.1046/j.1365-](https://doi.org/10.1046/j.1365-2311.2002.00487.x)
808 [2311.2002.00487.x](https://doi.org/10.1046/j.1365-2311.2002.00487.x)

809 Frank, K., Hülsmann, M., Assmann, T., Schmitt, T., Blüthgen, N., 2017. Land use affects dung
810 beetle communities and their ecosystem service in forests and grasslands. *Agric.*
811 *Ecosyst. Environ.* 243, 114–122. <https://doi.org/10.1016/j.agee.2017.04.010>

812 García-Tejero, S., Taboada, Á., Tárrega, R., Salgado, J.M., 2013. Land use changes and ground
813 dwelling beetle conservation in extensive grazing dehesa systems of north-west Spain.
814 Biol. Conserv. 161, 58–66. <https://doi.org/10.1016/j.biocon.2013.02.017>

815 Gardner, W., Mulvey, E.P., Shaw, E.C., 1995. Regression analyses of counts and rates:
816 Poisson, overdispersed Poisson, and negative binomial models. Psychol. Bull. 118, 392–
817 404.

818 Gelman, A., Hill, J., 2006. Data Analysis Using Regression and Multilevel/Hierarchical Models.
819 Cambridge, England: Cambridge University Press.

820 Giller, P.S., Doube, B.M., 1994. Spatial and Temporal Co-Occurrence of Competitors in
821 Southern African Dung Beetle Communities. J. Animal Ecol. 63, 629–643.
822 <https://doi.org/10.2307/5229>

823 Giller, P.S., Doube, B.M., 1989. Experimental Analysis of Inter- and Intraspecific Competition
824 in Dung Beetle Communities. J. Animal Ecol. 58, 129–142. <https://doi.org/10.2307/4990>

825 Gittings, T., Giller, P.S., 1999. Larval dynamics in an assemblage of *Aphodius* dung beetles.
826 Pedobiologia 43, 439–452.

827 Gittings, T., Giller, P.S., 1997. Life History Traits and Resource Utilisation in an Assemblage of
828 North Temperate *Aphodius* Dung Beetles (Coleoptera: Scarabaeidae). Ecography 20, 55–
829 66.

830 Gittings, T., Giller, P.S., Stakelum, G., 1994. Dung decomposition in contrasting temperate
831 pastures in relation to dung beetle and earthworm activity. Pedobiologia 38, 455-474.

832 Gonçalves-Souza, T., Araújo, M.S., Barbosa, E.P., Lopes, S.M., Kaminski, L.A., Shimizu, G.H.,
833 Santos, A.J., Romero, G.Q., 2015. Fine-scale Beta-diversity Patterns Across Multiple

834 Arthropod Taxa Over a Neotropical Latitudinal Gradient. *Biotropica* 47, 588–594.
835 <https://doi.org/10.1111/btp.12242>

836 Gotelli, N.J., Colwell, R.K., 2010. Estimating Species Richness, in: Magurran, A.E., McGill, B.J.
837 (Eds.), *Biological Diversity: Frontiers in Measurement and Assessment*. Oxford University
838 Press, United Kingdom, pp. 39-54.

839 Greenwood, K.L., McKenzie, B.M., 2001. Grazing effects on soil physical properties and the
840 consequences for pastures: a review. *Aust. J. Exp. Agric.* 41, 1231–1250.
841 <https://doi.org/10.1071/ea00102>

842 Griffiths, H.M., Bardgett, R.D., Louzada, J., Barlow, J., 2016. The value of trophic interactions
843 for ecosystem function: dung beetle communities influence seed burial and seedling
844 recruitment in tropical forests. *Proc. Biol. Sci.* 283.
845 <https://doi.org/10.1098/rspb.2016.1634>

846 Grime, J.P., 1973. Competitive Exclusion in Herbaceous Vegetation. *Nature* 242, 344.
847 <https://doi.org/10.1038/242344a0>

848 Halffter, G., Edmonds, W.D., 1982. The nesting behavior of dung beetles (Scarabaeinae): An
849 ecological and evolutive approach. Instituto de Ecología Publication 10 México D.F., Man
850 and the Biosphere Program UNESCO.

851 Hanski, I., Cambefort, Y., 1991. *Dung Beetle Ecology*. Princeton University Press.

852 Hobbs, N.T., 2006. Large herbivores as sources of disturbance in ecosystems, in: Danell, K.,
853 Bergstrom, R., Duncan, P., Pastor, J. (Eds.), *Large Herbivore Ecology, Ecosystem*
854 *Dynamics and Conservation*. Cambridge University Press, Cambridge, pp. 261–288.

855 Hodgson, J.G., Montserrat-Martí, G., Tallowin, J., Thompson, K., Díaz, S., Cabido, M., Grime,
856 J.P., Wilson, P.J., Band, S.R., Bogard, A., Cabido, R., Cáceres, D., Castro-Díez, P., Ferrer,
857 C., Maestro-Martínez, M., Pérez-Rontomé, M.C., Charles, M., Cornelissen, J.H.C.,
858 Dabbert, S., Pérez-Harguindeguy, N., Krimly, T., Sijtsma, F.J., Strijker, D., Vendramini, F.,
859 Guerrero-Campo, J., Hynd, A., Jones, G., Romo-Díez, A., de Torres Espuny, L., Villar-
860 Salvador, P., Zak, M.R., 2005. How much will it cost to save grassland diversity? *Biol.*
861 *Conserv.* 122, 263–273. <https://doi.org/10.1016/j.biocon.2004.07.016>

862 Horgan, F.G., 2005. Aggregated distribution of resources creates competition refuges for
863 rainforest dung beetles. *Ecography* 28, 603–618. [https://doi.org/10.1111/j.2005.0906-](https://doi.org/10.1111/j.2005.0906-7590.04177.x)
864 [7590.04177.x](https://doi.org/10.1111/j.2005.0906-7590.04177.x)

865 Hutton, S.A., Giller, P.S., 2003. The effects of the intensification of agriculture on northern
866 temperate dung beetle communities. *J. Appl. Ecol.* 40, 994–1007.
867 <https://doi.org/10.1111/j.1365-2664.2003.00863.x>

868 Jankielsohn, A., Scholtz, C.H., Louw, S.V., 2001. Effect of Habitat Transformation on Dung
869 Beetle AssemblagesA Comparison Between a South African Nature Reserve and
870 Neighboring Farms. *Env. Entomol.* 30, 474–483. [https://doi.org/10.1603/0046-225X-](https://doi.org/10.1603/0046-225X-30.3.474)
871 [30.3.474](https://doi.org/10.1603/0046-225X-30.3.474)

872 Jay-Robert, P., Errouissi, F., Lumaret, J.P., 2008a. Temporal coexistence of dung-dweller and
873 soil-digger dung beetles (Coleoptera, Scarabaeoidea) in contrasting Mediterranean
874 habitats. *Bull. Entomol. Res.* 98, 303–316. <https://doi.org/10.1017/S0007485307005615>

875 Jay-Robert, P., Lobo, J.M., Lumaret, J.-P., 1997. Altitudinal Turnover and Species Richness
876 Variation in European Montane Dung Beetle Assemblages. *Arct. Antarct. Alp. Res.* 29,
877 196–205. <https://doi.org/10.1080/00040851.1997.12003232>

878 Jay-Robert, P., Lumaret, J.-P., Lebreton, J.-D., 2008b. Spatial and Temporal Variation of
879 Mountain Dung Beetle Assemblages and Their Relationships with Environmental Factors
880 (Aphodiinae: Geotrupinae: Scarabaeinae). *Ann. Entomol. Soc. Am.* 101, 58–69.
881 [https://doi.org/10.1603/0013-8746\(2008\)101\[58:SATVOM\]2.0.CO;2](https://doi.org/10.1603/0013-8746(2008)101[58:SATVOM]2.0.CO;2)

882 Jay-Robert, P., Niogret, J., Errouissi, F., Labarussias, M., Paoletti, É., Luis, M.V., Lumaret, J.-P.,
883 2008c. Relative efficiency of extensive grazing vs. wild ungulates management for dung
884 beetle conservation in a heterogeneous landscape from Southern Europe (Scarabaeinae,
885 Aphodiinae, Geotrupinae). *Biol. Conserv.* 141, 2879–2887.
886 <https://doi.org/10.1016/j.biocon.2008.09.001>

887 Jerrentrup, J.S., Wrage-Mönnig, N., Röver, K.-U., Isselstein, J., 2014. Grazing intensity affects
888 insect diversity via sward structure and heterogeneity in a long-term experiment. *J.*
889 *Appl. Ecol.* 51, 968–977. <https://doi.org/10.1111/1365-2664.12244>

890 de Jong, Y., Verbeek, M., Michelsen, V., Bjørn, P. de P., Los, W., Steeman, F., Bailly, N., Basire,
891 C., Chylarecki, P., Stloukal, E., Hagedorn, G., Wetzal, F., Glöckler, F., Kroupa, A., Korb, G.,
892 Hoffmann, A., Häuser, C., Kohlbecker, A., Müller, A., Güntsch, A., Stoev, P., Penev, L.,
893 2014. Fauna Europaea – all European animal species on the web. *Biodivers. Data J.* 2,
894 e4034. <https://doi.org/10.3897/BDJ.2.e4034>

895 Kerven, C., Behnke, R., 2011. Policies and practices of pastoralism in Europe. *Pastoralism* 1,
896 28. <https://doi.org/10.1186/2041-7136-1-28>

897 Klemperer, H.G., 1979. An analysis of the nesting behaviour of *Geotrupes spiniger* Marsham
898 (Coleoptera, Scarabaeidae). *Ecol. Entomol.* 4, 133–150. [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2311.1979.tb00569.x)
899 [2311.1979.tb00569.x](https://doi.org/10.1111/j.1365-2311.1979.tb00569.x)

900 van Klink, R., van der Plas, F., van Noordwijk, C.G.E., Wallis De Vries, M.F., Olf, H., 2015.
901 Effects of large herbivores on grassland arthropod diversity. *Biol. Rev.* 90, 347–366.
902 <https://doi.org/10.1111/brv.12113>

903 Kőrösi, Á., Batáry, P., Orosz, A., Rédei, D., Báldi, A., 2012. Effects of grazing, vegetation
904 structure and landscape complexity on grassland leafhoppers (Hemiptera:
905 Auchenorrhyncha) and true bugs (Hemiptera: Heteroptera) in Hungary. *Insect Conserv.*
906 *Divers.* 5, 57–66. <https://doi.org/10.1111/j.1752-4598.2011.00153.x>

907 Kruess, A., Tschardt, T., 2002. Grazing Intensity and the Diversity of Grasshoppers,
908 Butterflies, and Trap-Nesting Bees and Wasps. *Conserv. Biol.* 16, 1570–1580.
909 <https://doi.org/10.1046/j.1523-1739.2002.01334.x>

910 Landin, B.O., 1961. Ecological studies on dung beetles (Col. Scarabaeidae). *Opuscula*
911 *Entomologica (Supplementum)* 19, 1-277.

912 Lee, S.M., Chao, A., 1994. Estimating population size via sample coverage for closed capture-
913 recapture models. *Biometrics* 50, 88–97.

914 Legendre, P., Anderson, M.J., 1999. Distance-Based Redundancy Analysis: Testing
915 Multispecies Responses in Multifactorial Ecological Experiments. *Ecol. Monogr.* 69, 1–
916 24. [https://doi.org/10.1890/0012-9615\(1999\)069\[0001:DBRATM\]2.0.CO;2](https://doi.org/10.1890/0012-9615(1999)069[0001:DBRATM]2.0.CO;2)

917 Legendre, P., Legendre, L., 2012. *Numerical ecology*, 3rd Edition. Elsevier, Amsterdam.

918 Legendre, P., Gallagher, E.D., 2001. Ecologically meaningful transformations for ordination of
919 species data. *Oecologia* 129, 271–280. <https://doi.org/10.1007/s004420100716>

920 Löbl, I., Löbl, D., 2016. Catalogue of Palaearctic Coleoptera. Scarabaeoidea, Scirtoidea,
921 Dascilloidea, Buprestoidea and Byrrhoidea. Volume 3. Revised and Updated Edition.
922 Brill, Leiden-Boston.

923 Lobo, J.M., 2001. Decline of roller dung beetle (Scarabaeinae) populations in the Iberian
924 peninsula during the 20th century. *Biol. Conserv.* 97, 43–50.
925 [https://doi.org/10.1016/S0006-3207\(00\)00093-8](https://doi.org/10.1016/S0006-3207(00)00093-8)

926 Lobo, J.M., Hortal, J., Cabrero-Sañudo, F.J., 2006. Regional and local influence of grazing
927 activity on the diversity of a semi-arid dung beetle community. *Divers. Distrib.* 12, 111–
928 123. <https://doi.org/10.1111/j.1366-9516.2006.00194.x>

929 Lobo, J.M., Lumaret, J.P., Jay-Robert, P., 1998. Sampling dung beetles in the French
930 Mediterranean area: effects of abiotic factors and farm practices. *Pedobiologia* 42, 252-
931 266.

932 Lu, X., Kelsey, K.C., Yan, Y., Sun, J., Wang, X., Cheng, G., Neff, J.C., 2017. Effects of grazing on
933 ecosystem structure and function of alpine grasslands in Qinghai–Tibetan Plateau: a
934 synthesis. *Ecosphere* 8, e01656. <https://doi.org/10.1002/ecs2.1656>

935 Lumaret, J.P., 1990. Atlas des Coléoptères Scarabeides Laparosticti de France. Muséum
936 National d’Histoire Naturelle. Secrétariat de la Faune et la Flore 1.

937 Lumaret, J.P., 1983. Structure des peuplements de coprophages Scarabaeidae en région
938 méditerranéenne française: relations entre les conditions écologiques et quelques
939 paramètres biologiques des espèces (Col.). *Bull. Soc. entomol. Fr.* 88, 481-495.

940 Lumaret, J.P., Iborra, O., 1996. Separation of trophic niches by dung beetles (Coleoptera,
941 Scarabaeoidea) in overlapping habitats. *Pedobiologia* 40, 392–404.

942 Lumaret, J.P., Kirk, A.A., 1991. South Temperate Dung Beetles, in: Hanski, I., Cambefort, Y.
943 (Eds.), *Dung Beetle Ecology*. Princeton University Press, pp. 97-115.

944 Lumaret, J.P., Stiernet, N., 1991. Montane Dung Beetles, in: Hanski, I., Cambefort, Y. (Eds.),
945 *Dung Beetle Ecology*. Princeton University Press, pp. 242-254.

946 Lumaret, J.P., Kadiri, N., Bertrand, M., 1992. Changes in Resources: Consequences for the
947 Dynamics of Dung Beetle Communities. *J. Appl. Ecol.* 29, 349–356.
948 <https://doi.org/10.2307/2404504>

949 Macagno, A.L.M., Palestrini, C., 2009. The maintenance of extensively exploited pastures
950 within the Alpine mountain belt: implications for dung beetle conservation (Coleoptera:
951 Scarabaeoidea). *Biodivers. Conserv.* 18, 3309. [https://doi.org/10.1007/s10531-009-](https://doi.org/10.1007/s10531-009-9643-1)
952 9643-1

953 Mackey, R.L., Currie, D.J., 2001. The Diversity-Disturbance Relationship: Is It Generally Strong
954 and Peaked? *Ecology* 82, 3479–3492. <https://doi.org/10.2307/2680166>

955 Mazoyer, M., Roudart, L., 2002. *Histoire des agricultures du monde : Du néolithique à la crise*
956 *contemporaine*. Seuil, Paris.

957 McGeoch, M.A., Rensburg, B.J.V., Botes, A., 2002. The verification and application of
958 bioindicators: a case study of dung beetles in a savanna ecosystem. *J. Appl. Ecol.* 39,
959 661–672. <https://doi.org/10.1046/j.1365-2664.2002.00743.x>

960 Milotić, T., Baltzinger, C., Eichberg, C., Eycott, A.E., Heurich, M., Müller, J., Noriega, J.A.,
961 Menendez, R., Stadler, J., Ádám, R., Bargmann, T., Bilger, I., Buse, J., Calatayud, J.,
962 Ciubuc, C., Boros, G., Jay-Robert, P., Kruus, M., Merivee, E., Miessen, G., Must, A., Ardali,
963 E., Preda, E., Rahimi, I., Rohwedder, D., Rose, R., Slade, E.M., Somay, L., Tahmasebi, P.,

964 Ziani, S., Hoffmann, M., 2019. Functionally richer communities improve ecosystem
965 functioning: Dung removal and secondary seed dispersal by dung beetles in the Western
966 Palaeartic. *J. Biogeogr.* 46, 70–82. <https://doi.org/10.1111/jbi.13452>

967 Negro, M., Rolando, A., Palestrini, C., 2011. The impact of overgrazing on dung beetle
968 diversity in the Italian Maritime Alps. *Environ. Entomol.* 40, 1081–1092.
969 <https://doi.org/10.1603/EN11105>

970 Noy-Meir, I., Gutman, M., Kaplan, Y., 1989. Responses of Mediterranean Grassland Plants to
971 Grazing and Protection. *J. Ecol.* 77, 290–310. <https://doi.org/10.2307/2260930>

972 Numa, C., Verdú, J.R., Rueda, C., Galante, E., 2012. Comparing Dung Beetle Species
973 Assemblages Between Protected Areas and Adjacent Pasturelands in a Mediterranean
974 Savanna Landscape. *Rangeland Ecol. Manag.* 65, 137–143.
975 <https://doi.org/10.2111/REM-D-10-00050.1>

976 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P.R.,
977 O’Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Eduard, Szoecs, E., Wagner, H.,
978 2018. *vegan: Community Ecology Package*. R package version 2.5-2. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)
979 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan).

980 Paine, L., Undersander, D.J., Sample, D.W., Bartelt, G.A., Schatteman, T.A., 1996. Cattle
981 trampling of simulated ground nests in rotationally grazed pastures. *Rangeland Ecol.*
982 *Manag.* 49, 294–300.

983 Paulian, R., Baraud, J., 1982. Faune des Coléoptères de France II - Lucanoidea et
984 Scarabaeoidea. *Encyclopédie Entomologique XLIII*, Editions Lechevalier SARL, Paris.

985 Peck, S.B., Howden, H.F., 1984. Response of a dung beetle guild to different sizes of dung
986 bait in a Panamanian rainforest. *Biotropica* 16, 235-238.

987 Pielou, E.C., 1966. The measurement of diversity in different types of biological collections. *J.*
988 *Theor. Biol.* 13, 131–144. [https://doi.org/10.1016/0022-5193\(66\)90013-0](https://doi.org/10.1016/0022-5193(66)90013-0)

989 Pizzio, R., Herrero-Jáuregui, C., Pizzio, M., Oesterheld, M., 2016. Impact of stocking rate on
990 species diversity and composition of a subtropical grassland in Argentina. *Appl. Veg. Sci.*
991 19, 454–461. <https://doi.org/10.1111/avsc.12229>

992 R Development Core Team, 2019. R: A language and environment for statistical computing. R
993 Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

994 Ratcliffe, B.C., 1991. The scarab beetles of Nebraska. *Bulletin of the University of Nebraska*
995 *State Museum* 12.

996 Redecker, B., Härdtle, W., Finck, P., Riecken, U., Schröder, E. (Eds.), 2002. *Pasture*
997 *Landscapes and Nature Conservation*. Springer-Verlag, Berlin Heidelberg.

998 Roslin, T., 2001. Large-scale spatial ecology of dung beetles. *Ecography* 24, 511-524.

999 Roslin, T., 2000. Dung beetle movements at two spatial scales. *Oikos* 91, 323-335.

1000 Roslin, T., Viljanen, H., 2011. Dung Beetle Populations: Structure and Consequences, in:
1001 Simmons, L.W., Ridsdill-Smith, T.J., (Eds.), *Ecology and Evolution of Dung Beetles*. Wiley-
1002 Blackwell, pp. 220-243.

1003 Rotem, G., Gavish, Y., Shacham, B., Giladi, I., Bouskila, A., Ziv, Y. 2016. Combined effects of
1004 climatic gradient and domestic livestock grazing on reptile community structure in a
1005 heterogeneous agroecosystem. *Oecologia* 180(1), 231-242.

1006 Sands, B., Wall, R., 2018. Sustained parasiticide use in cattle farming affects dung beetle
1007 functional assemblages. *Agric. Ecosyst. Environ.* 265, 226–235.
1008 <https://doi.org/10.1016/j.agee.2018.06.012>

1009 Sears, M.K., 1978. Damage to golf course fairways by *Aphodius granarius* (L.) (Coleoptera:
1010 Scarabaeidae). *Proc. Entomol. Soc. Ont.* 109:48.

1011 Simons, N.K., Weisser, W.W., Gossner, M.M., 2016. Multi-taxa approach shows consistent
1012 shifts in arthropod functional traits along grassland land-use intensity gradient. *Ecology*
1013 97, 754–764. <https://doi.org/10.1890/15-0616.1>

1014 Sirami, C., Nespoulous, A., Cheylan, J.-P., Marty, P., Hvenegaard, G.T., Geniez, P., Schatz, B.,
1015 Martin, J.-L., 2010. Long-term anthropogenic and ecological dynamics of a
1016 Mediterranean landscape: Impacts on multiple taxa. *Landsc. Urban Plan.* 96, 214–223.
1017 <https://doi.org/10.1016/j.landurbplan.2010.03.007>

1018 Slade, E.M., Mann, D.J., Villanueva, J.F., Lewis, O.T., 2007. Experimental evidence for the
1019 effects of dung beetle functional group richness and composition on ecosystem function
1020 in a tropical forest. *J. Anim. Ecol.* 76, 1094–1104. [https://doi.org/10.1111/j.1365-
1021 2656.2007.01296.x](https://doi.org/10.1111/j.1365-2656.2007.01296.x)

1022 Sowig, P., 1996. Brood care in the dung beetle *Onthophagus vacca* (Coleoptera:
1023 Scarabaeidae): the effect of soil moisture on time budget, nest structure, and
1024 reproductive success. *Ecography* 19, 254–258. [https://doi.org/10.1111/j.1600-
1025 0587.1996.tb01252.x](https://doi.org/10.1111/j.1600-0587.1996.tb01252.x)

1026 Sowig, P., 1995. Habitat selection and offspring survival rate in three paracoprid dung
1027 beetles: the influence of soil type and soil moisture. *Ecography* 18, 147–154.
1028 <https://doi.org/10.1111/j.1600-0587.1995.tb00335.x>

1029 Spector, S., 2006. Scarabaeine Dung Beetles (coleoptera: Scarabaeidae: Scarabaeinae): An
1030 Invertebrate Focal Taxon for Biodiversity Research and Conservation. *Coleopt. Bull.* 60,
1031 71–83. [https://doi.org/10.1649/0010-065X\(2006\)60\[71:SDBCSS\]2.0.CO;2](https://doi.org/10.1649/0010-065X(2006)60[71:SDBCSS]2.0.CO;2)

1032 Squires, V.R., Dengler, J., Hua, L., Feng, H., 2018. *Grasslands of the World: Diversity,*
1033 *Management and Conservation.* CRC Press.

1034 Steinfeld, H., Mooney, H. A., Schneider, F., Neville, L.E., 2013. *Livestock in a changing*
1035 *landscape, volume 1: drivers, consequences, and responses.* Island Press.

1036 Stoate, C., Báldi, A., Beja, P., Boatman, N.D., Herzon, I., van Doorn, A., de Snoo, G.R., Rakosy,
1037 L., Ramwell, C., 2009. Ecological impacts of early 21st century agricultural change in
1038 Europe – A review. *J. Environ. Manag.* 91, 22–46.
1039 <https://doi.org/10.1016/j.jenvman.2009.07.005>

1040 Tatin, L., Becker, E., Sauguet, F., Jay-Robert, P., 2014. First study focused on dung beetles in
1041 the Crau steppe. *Nature de Provence* 1 – 12. [http://www.cen-](http://www.cen-paca.org/images/5_publications/nature/n2/14CoprophagesTAP01_2014.pdf)
1042 [paca.org/images/5_publications/nature/n2/14CoprophagesTAP01_2014.pdf](http://www.cen-paca.org/images/5_publications/nature/n2/14CoprophagesTAP01_2014.pdf). (accessed
1043 01 January 2019).

1044 Tatin, L., Wolff, A., Boutin, J., Colliot, E., Dutoit, T., coordinateurs., 2013. *Ecologie et*
1045 *conservation d’une steppe méditerranéenne – La plaine de Crau.* Editions QUAE, ISI
1046 Print.

1047 Tocco, C., Probo, M., Lonati, M., Lombardi, G., Negro, M., Nervo, B., Rolando, A., Palestrini,
1048 C., 2013. Pastoral Practices to Reverse Shrub Encroachment of Sub-Alpine Grasslands:
1049 Dung Beetles (Coleoptera, Scarabaeoidea) Respond More Quickly Than Vegetation. PLoS
1050 ONE 8, e83344. <https://doi.org/10.1371/journal.pone.0083344>

1051 Tonelli, M., Verdú, J.R., Zunino, M., 2019. Grazing abandonment and dung beetle
1052 assemblage composition: Reproductive behaviour has something to say. Ecol. Indic. 96,
1053 361–367. <https://doi.org/10.1016/j.ecolind.2018.09.010>

1054 Tonelli, M., Verdú, J.R., Zunino, M., 2018. Effects of the progressive abandonment of grazing
1055 on dung beetle biodiversity: body size matters. Biodivers. Conserv. 27, 189–204.
1056 <https://doi.org/10.1007/s10531-017-1428-3>

1057 Tonelli, M., Verdú, J.R., Zunino, M.E., 2017. Effects of grazing intensity and the use of
1058 veterinary medical products on dung beetle biodiversity in the sub-mountainous
1059 landscape of Central Italy. PeerJ 5, e2780. <https://doi.org/10.7717/peerj.2780>

1060 Veiga, C.M., Lobo, J.M., Martin-Piera, F., 1989. Las trampas pitfall con cebo, sus posibilidades
1061 en el estudio de las comunidades coprofagas de Scarabaeoidea (Col.). II; Analisis de
1062 efectividad. Rev. Ecol. Biol. Sol. 26(1): 91–109.

1063 Venables, W.N., Ripley, B.D., 2002. Modern Applied Statistics with S. Fourth Edition.
1064 Springer, New York.

1065 Verdú, J.R., Lobo, J.M., Sánchez-Piñero, F., Gallego, B., Numa, C., Lumaret, J.-P., Cortez, V.,
1066 Ortiz, A.J., Tonelli, M., García-Teba, J.P., Rey, A., Rodríguez, A., Durán, J., 2018.
1067 Ivermectin residues disrupt dung beetle diversity, soil properties and ecosystem

1068 functioning: An interdisciplinary field study. *Sci. Total Environ.* 618, 219–228.
1069 <https://doi.org/10.1016/j.scitotenv.2017.10.331>

1070 Verdú, J.R., Numa, C., Hernández-Cuba, O., 2011. The influence of landscape structure on
1071 ants and dung beetles diversity in a Mediterranean savanna—Forest ecosystem. *Ecol.*
1072 *Indic.* 11, 831–839. <https://doi.org/10.1016/j.ecolind.2010.10.011>

1073 Weiss, M., Procházka, J., Schlaghamerský, J., Cizek, L., 2016. Fine-Scale Vertical Stratification
1074 and Guild Composition of Saproxyllic Beetles in Lowland and Montane Forests: Similar
1075 Patterns despite Low Faunal Overlap. *PLoS ONE* 11, e0149506.
1076 <https://doi.org/10.1371/journal.pone.0149506>

1077 Wenzel, M., Schmitt, T., Weitzel, M., Seitz, A., 2006. The severe decline of butterflies on
1078 western German calcareous grasslands during the last 30 years: A conservation
1079 problem. *Biol. Conserv.* 128, 542–552. <https://doi.org/10.1016/j.biocon.2005.10.022>

1080 Wilson, E.O., 1987. The Little Things That Run the World (The Importance and Conservation
1081 of Invertebrates). *Conserv. Biol.* 1, 344–346.

1082 Xu, M., Xie, F., Wang, K., 2014. Response of Vegetation and Soil Carbon and Nitrogen Storage
1083 to Grazing Intensity in Semi-Arid Grasslands in the Agro-Pastoral Zone of Northern
1084 China. *PLoS ONE* 9, e96604. <https://doi.org/10.1371/journal.pone.0096604>

1085 Yan, Y., Yan, R., Chen, Jiquan, Xin, X., Eldridge, D.J., Shao, C., Wang, X., Lv, S., Jin, D., Chen,
1086 Jinqiang, Guo, Z., Chen, B., Xu, L., 2018. Grazing modulates soil temperature and
1087 moisture in a Eurasian steppe. *Agric. For. Meteorol.* 262, 157–165.
1088 <https://doi.org/10.1016/j.agrformet.2018.07.011>


1089 Zettler, M.L., Proffitt, C.E., Darr, A., Degraer, S., Devriese, L., Greathead, C., Kotta, J., Magni,
1090 P., Martin, G., Reiss, H., Speybroeck, J., Tagliapietra, D., Hoey, G.V., Ysebaert, T., 2013.
1091 On the Myths of Indicator Species: Issues and Further Consideration in the Use of Static
1092 Concepts for Ecological Applications. PLoS ONE 8, e78219.
1093 <https://doi.org/10.1371/journal.pone.0078219>

1094 Zhou, G., Zhou, X., He, Y., Shao, J., Hu, Z., Liu, R., Zhou, H., Hosseinibai, S., 2017. Grazing
1095 intensity significantly affects belowground carbon and nitrogen cycling in grassland
1096 ecosystems: a meta-analysis. Glob. Change Biol. 23, 1167–1179.
1097 <https://doi.org/10.1111/gcb.13431>


1098 Zhou, Z.C., Gan, Z.T., Shanguan, Z.P., Dong, Z.B., 2010. Effects of grazing on soil physical
1099 properties and soil erodibility in semiarid grassland of the Northern Loess Plateau
1100 (China). Catena 82, 87–91. <https://doi.org/10.1016/j.catena.2010.05.005>

1101

1102




18 **Figure 1.** Location of the two study areas representing two distinct bioclimatic contexts, (1)
 19 the Coussouls de Crau National Nature Reserve (steppe) near the Mediterranean Sea, and
 20 (2) the Vanoise National Park (Alps) in the northern French Alps (map of France:
 21 <https://upload.wikimedia.org/>). Illustration of sheep flocks in these two areas (photographs
 22 ©William Perrin) and schematic representation of the sampling design and the grazing
 23 intensity gradient. In each study area, two distinct pastures were sampled. Within each
 24 pasture, we defined three distinct zones representing three levels of grazing intensity,
 25 ranging from low (LGI) to moderate (MGI) to high (HGI) (although we show the grazing
 26 intensity gradient for just one pasture, the same sampling design was carried out in each of
 27 the four pastures). Within each level of grazing intensity, we selected a sampling plot where
 28 dung beetles were collected with five pitfall traps.




8 **Figure 2.** Triplots of RDA-based MANOVAs for the two pastures (A and B) sampled in the steppe area. Numbers represent the traps, whose “wa”
 9 scores are related to the centroids of the grazing intensity factor (three levels: Low = LGI, Moderate = MGI and High = HGI) by means of straight
 10 lines, in scaling 1. Ellipses represent the three distinct grazing intensity levels in each pasture. Letters represent the species. The percentage of
 11 variation explained by the RDAs is shown for each axis. Underlined letters correspond to indicator species found in the IndVal analysis (**Table 2**):
 12 *Acrossus luridus* (a); *Aphodius foetidus* (c), *Caccobius schreberi* (f), *Calamosternus granarius* (g), *Copris hispanus hispanus* (i), *Gymnopleurus*
 13 *flagellatus flagellatus* (n), *Onthophagus furcatus* (p), *Onthophagus opacicollis* (r), *Onthophagus ruficapillus* (s), *Onthophagus vacca* (u). See
 14 **Appendix D** for letters identifying the other species on the triplots.

1
2
3
4
5
6
7
8
9
10
11
12
13
14



15 **Figure 3.** Triplots of RDA-based MANOVAs for the two pastures (C and D) sampled in the Alps area. Numbers represent the traps, whose “wa” scores are
16 related to the centroids of the grazing intensity factor (three levels: Low = LGI, Moderate = MGI and High = HGI) by means of straight lines, in scaling 1. Ellipses
17 represent the three distinct grazing intensity levels in each pasture. Letters represent the species. The percentage of variation explained by the RDAs is
18 shown for each axis. Underlined letters correspond to indicator species found in the IndVal analysis (Table 2): *Geotrupes stercorarius* (m), *Oromus alpinus* (p).
19 See **Appendix D** for letters identifying the other species on the triplots.



19 **Figure 4.** Variation in the number of individuals, rarefied species richness and Pielou's evenness
 20 between the three levels of grazing intensity in the steppe (A) and the Alps (B). Different letters (a, b,
 21 c) indicate significant differences between grazing intensity levels at $\alpha = 0.05$ based on the models.
 22 Vertical bars represent standard errors produced by the generalized linear models with a negative
 23 binomial family (for the number of individuals) and linear models with a Gaussian family (for rarefied
 24 species richness and Pielou's evenness).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25




Figure 5. Variation in the number of dung-dwellers and soil-diggers between the three levels of grazing intensity in the steppe (A) and the Alps (B). Different letters (a, b, c) indicate significant differences between grazing intensity levels at $\alpha = 0.05$ based on the models. Vertical bars represent standard errors produced by the generalized linear models with a negative binomial family.

1 **Table 1.** Summarized statistics for dung beetles sampled in the steppe and the Alps. Mean values
2 represent the mean number of individuals or species sampled per pitfall trap. *Percentage of species
3 richness (in parentheses) inventoried according to estimates based on the iChao2 and ICE-1 indices.

Study area	Nb. indiv.	Mean indiv. \pm SD	Nb. sp.	Mean sp. \pm SD	iChao2 (%) *	ICE-1 (%) *
Steppe	1,465	48.8 \pm 22.2	30	8 \pm 3.0	40.2 (75)	35.8 (84)
Alps	10,268	342.3 \pm 232.3	26	11.1 \pm 2.3	26.4 (98)	28.2 (92)

4

5

1 **Table 2.** Results of the IndVal analysis with indicator species for each level of grazing intensity (GI) or
 2 combinations of two levels (Low–Moderate = LGI-MGI and Moderate–High = MGI-HGI). Only species
 3 with a significant IndVal value ($p \leq 0.05$) are shown. The values for specificity and fidelity are indicated,
 4 as well as nesting guild and mean dry body mass.

Steppe							
Grazing intensity	Species	Nesting guild	Mean body mass (mg) \pm SD	Specificity	Fidelity	IndVal	p-Value
LGI	<i>Copris hispanus hispanus</i>	Soil-digger	452.2 \pm 108.4	0.80	0.60	0.69	0.0052
LGI - MGI	<i>Onthophagus vacca</i>	Soil-digger	34.1 \pm 5.0	0.94	1.00	0.97	0.0020
	<i>Acrossus luridus</i>	Dung-dweller	12.0 \pm 2.8	0.92	0.70	0.80	0.0356
	<i>Gymnopleurus flagellatus flagellatus</i>	Soil-digger	67.7 \pm 13.0	1.00	0.55	0.74	0.0104
	<i>Euoniticellus fulvus</i>	Soil-digger	13.8 \pm 4.1	1.00	0.50	0.71	0.0262
	<i>Eudolus quadrigutattus</i>	Dung-dweller	2.8 \pm 0.8	1.00	0.45	0.67	0.0400
MGI	<i>Onthophagus furcatus</i>	Soil-digger	4.9 \pm 1.4	0.86	0.90	0.88	0.0002
	<i>Caccobius schreberi</i>	Soil-digger	9.1 \pm 2.2	0.85	0.60	0.71	0.0118
MGI - HGI	<i>Onthophagus ruficapillus</i>	Soil-digger	6.4 \pm 1.6	0.98	0.80	0.89	0.0020
HGI	<i>Calamosternus granarius</i>	Dung-dweller	3.1 \pm 0.9	0.99	1.00	0.99	0.0002
	<i>Aphodius foetidus</i>	Dung-dweller	6.9 \pm 2.4	0.95	0.60	0.76	0.0022
	<i>Onthophagus opacicollis</i>	Soil-digger	15.8 \pm 3.0	1.00	0.50	0.71	0.0048
Alps							
Grazing intensity	Species	Nesting group	Mean body mass (mg) \pm SD	Specificity	Fidelity	IndVal	p-Value
LGI	<i>Geotrupes stercorarius</i>	Soil-digger	163.5 \pm 60.2	0.81	0.80	0.80	0.0006
	<i>Trypocopris vernalis</i>	Soil-digger	92.1 \pm 26.2	0.83	0.70	0.76	0.0016
	<i>Otophorus haemorrhoidalis</i>	Dung-dweller	3.1 \pm 1.0	0.78	0.60	0.68	0.0122
MGI - HGI	<i>Oromus alpinus</i>	Dung-dweller	3.1 \pm 0.7	1.00	0.60	0.68	0.0190