



Removal of metronidazole from aqueous media by *C. vulgaris*

Sufia Hena, Leo Gutierrez, Jean-Philippe Croué

► To cite this version:

Sufia Hena, Leo Gutierrez, Jean-Philippe Croué. Removal of metronidazole from aqueous media by *C. vulgaris*. Journal of Hazardous Materials, 2020, 384, pp.121400. <10.1016/j.jhazmat.2019.121400>. <hal-03488732>

HAL Id: hal-03488732

<https://hal.science/hal-03488732v1>

Submitted on 21 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

Removal of metronidazole from aqueous media by *C. vulgaris*

Sufia Hena^a, Leo Gutierrez^b and Jean-Philippe Croué^{a, c, *}

^a Curtin Water Quality Research Centre, Department of Chemistry, Curtin University, Australia

^b Facultad del Mar y Medio Ambiente, Universidad del Pacifico, Ecuador

^c Institut de Chimie des Milieux et des Matériaux IC2MP UMR 7285 CNRS, Université de Poitiers, France

* Corresponding author

Email : jean-philippe.croue@curtin.edu.au (Jean-Philippe Croué)

25 **Highlights**

26

- 27 1. Metronidazole is efficiently removed during algal treatment.
- 28 2. Metronidazole is removed via sorption onto the biomass.
- 29 3. Metronidazole showed toxicity to *C. vulgaris* at stationary phase of life cycle.
- 30 4. The removal efficiency of *C. vulgaris* decreased with increase of antibiotic.

Abstract

This current study investigated the removal of metronidazole from aqueous media by *C. vulgaris*. Two different initial sizes of inoculum (0.05 and 0.1mg/L) were tested for a wide concentration range of metronidazole (1-50 μ M). The effect of metronidazole concentrations on biomass production was studied for 20 days. The exopolymeric substances (EPS) were quantified and correlated with the removal of antibiotics from aqueous media. Specifically, MDZ stimulated the production of EPS in *C. vulgaris*, which played the major role in the adsorption of this antibiotic. Also, metronidazole significantly influenced the zeta potential of *C. vulgaris* in the test cultures, indicating a change in surface characteristics. This decrease in surface negative charge caused auto-flocculation phenomena at a stationary phase. Chronic and acute toxicity experiments showed that metronidazole was harmful to *C. vulgaris* at stationary phase. Results from this study would advance our knowledge on the treatment of metronidazole-contaminated waters with *C. vulgaris* as a green technology-oriented process.

Keywords: Adsorption; *C. vulgaris*; EC50; metronidazole

1. Introduction

The excessive use of pharmaceuticals and their improper disposal pose risks to the environment and public health [1]. Especially, antibiotics (i.e., widely used for preventing bacterial diseases in humans and livestock) are continuously introduced in natural aquatic environments [2, 3]. Despite their low concentrations, antibiotics potentially affect the quality of drinking water, human health, and ecosystems [4]. The residual antibiotics occurring in aquatic environments would lead to the emergence of resistance among targeted and non-targeted strains of microorganisms [5-7]. Also, surface waters contaminated with antibiotics expand the variety of resistant microorganisms and the transfer of genetic material between multidrug-resistant bacteria [8, 9].

Metronidazole is a first-generation derivative of 5-nitroimidazole drugs developed for therapeutic uses [10, 11]. Metronidazole is an effective prescription medication used for treating several infectious diseases caused by anaerobic bacteria and protozoa in humans and livestock [12, 13], showing minimal side effects and a broad spectrum of applications [14]. In general, antibiotics have a relatively short half-life in open environments (e.g., water bodies or soils) [2]; however, metronidazole is highly persistent in nature due to its high solubility [15] and low biodegradability [16]. Also, the removal of metronidazole from water and wastewater during the treatment train is not efficient [17]. Deblonde et al., [18], reported a mere 38.9% removal of metronidazole from a wastewater treatment plant. Nevertheless, because of its high toxicity and mutagenic effect on targeted and non-targeted microorganisms in the environment [12], it is highly necessary to develop efficient protocols and (bio) technologies to remove this antibiotic from aqueous media.

Several methods have been reported for the removal of metronidazole from aqueous media. Briefly, chemical processes such as reduction [19], Fenton and photo-Fenton process [20], UV-photolysis [21], and physical treatments (e.g., adsorption) [22, 23] have been mainly tested. However, high energy and chemical consumptions, as well as the generation of secondary waste are the main constraints for their successful (economic) implementation [24, 25]. On the other hand, Rivera-Utrilla et al., [22] had reported that microorganisms in wastewater treatment plants cannot efficiently degrade metronidazole. Activated sludge process is the most common biological treatment method to remove most of the drugs from aqueous media; however, this process fails to efficiently remove most of the antibiotics [26]. In general, antibiotics are not readily biodegradable [27]. Antibiotics are mainly removed by sorption onto activated sludges in wastewater treatment plants [4, 28]. However, antibiotics desorb from the activated sludge to the aqueous media during its reuse in aeration tanks to maintain a fixed concentration of activated microorganisms [29]. According to Grandclement et al., [30] wastewater treatment plants using activated sludge are not specifically designed to remove antibiotics.

Simultaneously, the activated sludge in wastewater treatment plants promotes the occurrence of antibiotic-resistant bacteria due to the exposure of bacteria to antibiotics present in wastewater [31]. Therefore, the efficient treatment of antibiotics present in wastewater to ensure biosafety for humans and their environment is a difficult task [32].

The use of microalgae as green technology is a viable option to remove antibiotics from water. Moreover, microalgal cultures mitigate atmospheric CO₂ while the biomass can be used for biofuel feedstocks. Briefly, microalgae are known for efficiently removing xenobiotic compounds from water, including a range of pharmaceuticals. Garcia-Rodríguez et al., [33] studied the role of microalga *Spyrogira* sp. in photolysis degradation and the contribution of light

in the total removal of six antibiotics from the sulfonamides and tetracyclines families. Du et al., [34] investigated the response of green algae *Chlorella pyrenoidosa* towards the removal of ceftazidime antibiotic. Guo and Chen [2] showed the removal of cephalosporins from wastewater using *Chlorella pyrenoidosa*. Conversely, Magalhães et al., [16] unsuccessfully attempted to remove metronidazole from water using *Microcystis protocystis* (cyanobacteria). From these previous studies, chlorophytes have clearly shown more resistant to stress conditions during water treatment. *Chlorella sp.* is a chlorophyte which has been extensively used for the removal of antibiotics such as 7-amino cephalosporanic acid [32], cephalosporins [2], norfloxacin [35], levofloxacin [36], and enrofloxacin [37]. However, to the best of our knowledge, metronidazole removal using *C. vulgaris* has not yet been reported.

This current study has investigated the metronidazole removal capacity of *C. vulgaris* and the dominant mechanisms involved in the removal. The impact of metronidazole at different concentrations on the growth rate of *C. vulgaris* was examined. Additionally, the tolerance and chronic toxicity (EC50) of *C. vulgaris* to this antibiotic as a function of its stage of life cycle was investigated. Finally, the exo-polymeric substances (EPS) production by *C. vulgaris* during exposure to metronidazole, its influence on the surface properties of *C. vulgaris*, and its role during the removal of this antibiotic were comprehensively studied. Results from this study will advance our knowledge of the treatment of metronidazole-contaminated waters with *C. vulgaris*. The protocols followed in the current investigation can be extended to the treatment of different antibiotics-contaminated aquifers using green technology-oriented processes. Simultaneously, a dual-approach can be used in the system to treat wastewater and to produce valuable biomass as feedstock for biofuel, through anaerobic digestion. This latter process would aim to completely degrade any antibiotic residue [61].

2. Materials and Methods

2.1. Chemical reagents

Metronidazole (HPLC grade; >99% purity) was obtained from Sigma-Aldrich. All the chemicals, solvents, and reagents used were of analytical grade. A stock solution of 1mM metronidazole (MDZ) was prepared with 30% methanol and 70% MilliQ water (v/v) and stored in a dark airtight glass vial at 4°C for no longer than 30 days for further experiments.

2.2. Algae Culturing and batch experiments

Chlorella vulgaris (*C. vulgaris*) was obtained from CSIRO Marine Research Hobart, Tasmania. *C. vulgaris* was re-cultured in 2 L conical flasks with a working volume of 1L in BG11 media (Table 1), at 25±4°C. The cultures were exposed to fluorescent lamps to provide the irradiance of 120 µmol photon /m²/s until the stationary phase was attained.

Two sets of batch experiments were conducted. The freshly prepared culture media were inoculated with two different sizes of inoculum of *C. vulgaris* for obtaining 0.05 g/L and 0.5 g L⁻¹ biomass (i.e., dry biomass vs optical density calibration curve) in 3 L of Erlenmeyer flasks with 2 L of working volume. The cultures were spiked with increasing concentration of MDZ (1-50 µM) except for the control cultures of each batch. All experiments were conducted in triplicate.

2.3. Determination of biomass and chlorophyll-a

The determination of the biomass of the test and control cultures was conducted every day by measuring the optical density at 680 nm as described in [38]. A batch study was conducted to evaluate the specific growth rate using the following formula;

138
$$\mu = (\ln N_2 - \ln N_1) / (t_2 - t_1)$$

139 Where, N_1 and N_2 are biomass concentrations (mg L^{-1}) at day t_1 and t_2 , respectively, during the
140 exponential phase of growth. All experiments were conducted in triplicate.

141 The extraction of chlorophyll-a and its quantification was conducted as reported by Xiong et al.,
142 [39]. A 10 mL of *C. vulgaris* was centrifuged at 4500 rpm for 10 min. The supernatant was
143 discarded and the pellet was collected. The pellet was re-suspended in 10 mL of methanol:water
144 (9:1; v/v) and incubated in 20 mL centrifuge tubes for 10 min. The incubated culture was
145 centrifuged again at 4500 rpm for 10 min. The absorbance of the supernatant was measured at
146 665 nm and 652 nm and the chlorophyll-a concentration of the extract was calculated using the
147 following equation:

148
$$\text{Chlorophyll-a } (\mu\text{g L}^{-1}) = 16.29 \times A_{665} - 8.54 \times A_{652}$$

149 Where, A_{665} and A_{652} are the absorbance of the supernatant at wavelength 665 nm and 652 nm,
150 respectively.

151
152 **2.4. Toxicity of MDZ for *C. vulgaris***

153 The toxicity of MDZ was calculated as the Effective Concentration (EC50) of the stressor
154 (MDZ) at which 50% of the growth rate of *C. vulgaris* was inhibited. The inhibition in growth
155 was calculated as a function of chlorophyll-a content. To determine the chronic toxicity of MDZ,
156 a batch experiment was conducted at different concentrations of MDZ (65-90 μM) with an initial
157 biomass concentration of 0.1 g L^{-1} . The microalgae were exposed to antibiotic for 240 hours (10
158 days). However, for acute toxicity experiments, a 0.1 g L^{-1} culture at stationary phase (20 days
159 old culture) was spiked with 1-50 μM antibiotics for 96 hours. The growth rate was calculated in
160 terms of chlorophyll-a ($\mu\text{g L}^{-1}$).

2.5. Extraction and analysis of Exo-polymeric substances (EPS)

Samples of 50 mL of *C. vulgaris* cultures at late exponential phase (i.e., which started approximately from day 10, Fig 1) were kept in the fridge at 4°C to settle the cells. The supernatants were decanted and used for EPS analysis, while the settled cells were re-suspended in 50 mL of MilliQ water in a beaker. The culture was stirred for 2 hours at 1000 rpm using a magnetic stirrer to extract the EPS from the cell wall of *C. vulgaris*. Then, the cells were separated by centrifugation at 1000 rpm for 10 minutes. The second supernatants were collected for analysis of bound EPS. The pellets of the centrifuged biomass were dried at 60°C. The EPS were extracted at every alternate day starting from day 10 until day 20.

30 mL of all supernatants were used for measuring the dry weight of EPS. Every supernatant of the tests and control cultures of age 10 days and beyond were used for EPS determination. 30 mL of supernatants were dried in an oven at 60°C until a stable weight was achieved to calculate the ratio of the dry weight of EPS to biomass.

2.6. Zeta potential

The Zeta (ζ) potentials of *C. vulgaris* from the test (low and high inoculum spiked with 5 μ M MDZ) and control cultures were analyzed on every alternate day from day 0 till day 20 using a Zetasizer (Malvern). 10mL of cultures were collected from the Erlenmeyer flasks. The biomass was allowed to settle down. The supernatants were discarded and biomass was re-suspended in MilliQ water. The suspended *C. vulgaris* was pipetted in a folded cuvette. The zeta potential was measured 5 times and the average and standard deviation were calculated.

2.7. Analysis of MDZ

Samples of 5 mL from test bioreactors were collected and filtered by a syringe filter (0.45 μ m) and analyzed using an HPLC equipped with UV/VIS detector (Agilent 1100 series, USA) at 320 nm. A 250 mm*4.6 mm C18 5- μ m reverse phase HPLC column (AlltimaTM, GRACE) was used and the mobile phase consisted of 0.1% phosphoric acid solution and methanol (80:20 v/v) with a 1 mL min⁻¹ flow rate. To analyze the removal mechanism and removal efficiency of MDZ via *C. vulgaris*, flocculated biomasses of microalgae were collected from the bottom of a bioreactor (i.e., a random selection was conducted from a 10 μ M MDZ high biomass bioreactor on day 20) in triplicate. The biomasses were washed thrice with 5 mL MilliQ to remove the unbounded molecules of MDZ from biomass using filter paper (GF/C; 1.2 μ m) and the filtrates were collected in HPLC vial for analysis after every wash. The absence of MDZ peak after third wash approved the adequacy of three washes of collected biomasses for further steps of analysis. The washed biomasses were collected on glass Petri dishes separately and vacuum dried at 60°C in temp controlled oven for 24 hours till they achieved constant weight. 0.5 mg of dried biomass was suspended into 20 mL of ethyl acetate followed by sonication for 15 minutes and centrifuging at 12,000 rpm for 10 minutes [16]. The biomass was separated from ethyl acetate using a syringe filter (GF/C; 1.2 μ m). The filtrate was vacuum dried at 40°C, followed by diluting the remaining residue in 20 mL fresh culture media. The final diluted residue (i.e., expecting desorbed MDZ in solution) was filtered with a 1.2 μ m GF/C syringe filter and collected in HPLC vial for analysis. The results were used to elucidate the MDZ removal efficiency of *C. vulgaris* and the mechanism involved during the removal process.

3. Results and discussion:

3.1. Effects of MDZ on the growth of *C. vulgaris*:

207 Two sizes of inoculum (0.05 and 0.5 g L⁻¹) were selected to study the effect of MDZ on *C.*
208 *vulgaris* concentration in solution. These results were compared with those of control culture. As
209 depicted in Fig 1a-b (Refer to Table S1a-b for standard deviations), the growth of the cultures
210 can be divided into three phases. The early exponential phase was calculated for 10 days from
211 the beginning of the culture due to the highest growth rate during this period for all tests and
212 control culture regardless of inoculum size. Interestingly, the specific growth rates of *C. vulgaris*
213 during the early exponential phase were almost similar for cultures with the same inoculum size
214 but different concentrations of MDZ (Table 2). Clearly, there was no impact of MDZ on *C.*
215 *vulgaris* during this early exponential phase. Remarkably, there was no removal of MDZ from
216 any of the test bioreactors during the early exponential phase (Figure 2a-b, Table S2a-b). During
217 the late exponential phase (following the first 10 days of growth), the biomass increased with
218 similar specific growth rate for the same inoculum size. These results indicate that MDZ does not
219 impact the specific growth rate of the culture. However, MDZ hampered the duration of the late
220 exponential growth phase, where MDZ decreased the surface charge of the cells causing them to
221 flocculate and settling down (Fig 3).

222 The duration of the late exponential phases was different in different test bioreactors. The longest
223 duration was observed for control cultures as 8 days; nevertheless, the control culture of high
224 inoculum attained a lower specific growth rate than control culture and low inoculum (Table 2).
225 The test cultures with low inoculum remained in late exponential phase for 5-6 days, while high
226 inoculum test cultures showed the shortest duration of 2-3 days. The longer late exponential
227 phase for low inoculum cultures might be due to: 1) a lower concentration of microalgal cells
228 present in the media than higher inoculum bioreactors for the same amount of nutrients, and 2)
229 late auto-flocculation in low inoculum cultures. The late auto-flocculation in low inoculum test

bioreactors caused biomass to remain in suspension media for a longer time and hence increased the duration of the late exponential phase. The concentrations of MDZ start decreasing during late exponential phase in all cases regardless of inoculum sizes (0.05 and 0.5 g L⁻¹) or initial concentrations (1-50 µM) of MDZ as shown in Fig 2a-b. These results confirmed that MDZ played a major role in auto-flocculation as further explained in next section. In all test bioreactors, the stationary phase did not last for more than 24 hours, which was then followed by a quick decline in biomass content. The reason for this rapid decrease in suspended biomass was due to auto-flocculation; however, the removal of MDZ continued even during the decline in biomass phase.

3.2. *Effect of MDZ on zeta potential of C. vulgaris:*

Fig 3 revealed the electronegative zeta potential of *C. vulgaris* from control culture and test bioreactor spiked with 5 µM MDZ (i.e., which was randomly selected as a representative case). In both cases, the electronegative zeta potential decreased with the age of the culture. Throughout the growth cycle of the microalgae, the concentration of the functional groups present on the cell membrane and cell wall changes and influences the surface properties of the cells [40]. However, during the active growth phase the lower intercellular interaction caused a higher net electronegative zeta potential [41]. At late exponential growth phase, the metabolism of the cells and the mobility decreased. Simultaneously, the changes on the cell surface caused intercellular interactions which resulted in a lower electronegative zeta potential [41]. Xia et al., [42] reported that the decrease in the electronegative zeta potential of the *C. vulgaris* was also due to the increase in the effective (hydrodynamic) size of the cells (i.e., cell with bound EPS) at

a later stage of its life cycle. Decho and Gutierrez [43] considered EPS as an extended part of the cell.

Fig 3 showed that the electronegative zeta potential of the *C. vulgaris* decreased more significantly in the test bioreactors than their respective control cultures. Among the test bioreactors, the cultures with low inoculum size showed a higher electronegative zeta potential than the high inoculum size culture at day 12 and beyond. The high inoculum culture contained more biomass, which attained a stationary phase earlier than low inoculum cultures. Since the EPS forms at late exponential or stationary phase of life cycle [43], high inoculum cultures start forming EPS earlier than the low inoculum culture (Table 3).

As mentioned earlier, the net negative zeta potential of the microalgal cell decreased due to change in functional groups present on the cell wall, which would be directly associated with the metabolic conditions of the cells [40]. However, in the current study, the additional decrease in zeta potential compared to their control cultures could be due to the charge dispersion interaction of MDZ with algal cells (discussed in section 3.4) as well as the increase in the effective size of the cell. The effective size of the cells would be larger in test cultures compared to their respective control cultures due to the formation of bound EPS onto the cells in test cultures (Table 3).

A drop in zeta potential in the test cultures indicated a decrease in the repulsive electrostatic forces between the cells which led to auto-flocculation. The auto-flocculation takes place when van der Waals forces overcome the repulsive electrostatic forces [44]. In this study, the repulsive electrostatic forces were most likely decreased due to the adsorption of EPS around the cells and also because of its interaction with MDZ in its scaffolds.

3.3. *Effect of MDZ on the EPS production of C. vulgaris*

EPS is produced by microalgae at a later stage of life cycle [43]. Microalgae produce EPS for cell protection as a waste of metabolic processes [45]. There are two types of EPS produced by *C. vulgaris*; a) EPS soluble (free) in media, and b) loosely bound (adsorbed) EPS which assists in auto-flocculation of cells [46]. EPS is generally discussed as a three-dimensional architecture scaffold providing space for trapping foreign compounds and slowing the diffusion rate, physical stability of cells, digestion of xenobiotic substances, and communication between cells (i.e., quorum sensing) [43].

Control culture produced more soluble (free) EPS than bound (adsorbed) EPS (Table 3). This result explains why control culture cells did not undergo auto-flocculation. However, in test bioreactors the auto-flocculation phenomenon was common. A slight increase in bound EPS was observed with an increase of MDZ concentration in both test cultures (low and high inoculum). The bioreactor spiked with 1 μM MDZ produced significantly less EPS ($47.7 \pm 1.23 \text{ mg g}^{-1} \text{ DW}$ of *C. vulgaris*) than that with 5 μM ($151.6 \pm 3.45 \text{ mg g}^{-1} \text{ DW}$ of *C. vulgaris*), which indicated that MDZ stimulated the EPS production in *C. vulgaris*. However, concentrations higher than 5 μM MDZ produced almost the same amount of EPS, which varied from 151.6-145.5 mg g^{-1} dry weight (DW) and 139.3-146.1 mg g^{-1} DW for low and high biomass bioreactors, respectively. These results revealed that 5 μM MDZ was the optimum concentration of antibiotics to produce maximum EPS for this specific system.

3.4. *Removal of MDZ*

MDZ was analyzed from day 0 until day 20 in test bioreactors. The removal of MDZ by *C. vulgaris* from aqueous media under different conditions is shown in Figure 2a-b. MDZ was not

298 removed until the early exponential growth phase. This result proved that MDZ did not adsorb
299 onto the cell surface of *C. vulgaris* possibly due to the low log K_{ow} (-2.3) value of MDZ. Also,
300 the pH of the *C. vulgaris* culture varied between 9-10, where MDZ is expected to behave as a
301 neutral molecule [47]. Therefore, electrostatic interactions between MDZ and negatively-charged
302 *C. vulgaris* would not be expected. However, π -electron of the heterocyclic aromatic ring of the
303 MDZ molecule would interact with different sites of EPS (Figure 4). EPS are highly hydrated
304 complex polymers mainly composed of polysaccharides, proteins, and DNA [48]. The CH group
305 of the polysaccharides form CH- π bond with the π -electrons of aromatic compounds [49, 50].
306 Consequently, the adsorption of MDZ molecules in the scaffold of the EPS matrix would involve
307 van der Waals interaction between π -electron of the heterocyclic aromatic ring of the MDZ
308 molecule and the hydrogen atom of CH group of the polysaccharides to form a CH- π bond.
309 Another possible mechanism would involve OH- π bond with OH groups of saccharides.
310 However, EPS has a complex structure where the OH groups of polysaccharides involved in
311 intermolecular interactions, might have no free OH to make bonds with MDZ [49]. Nishio [51]
312 has reported that saccharides are overall polar in nature but include some hydrophobic moieties
313 depending upon the orientation of OH groups, which have an affinity to form hydrogen bonds
314 with hydrophobic compounds [49]. The hydrophobic imidazole ring in MDZ would form OH- π
315 bond with OH groups of hydrophobic moieties. Previous studies have reported that MDZ
316 interacts with its π electron of heterocyclic aromatic ring to form π - π dispersion bonds with other
317 aromatic compounds [22]. Protein containing aromatic amino acids would be expected to favor
318 the formation of π - π dispersion bonds for MDZ to interact with EPS. Sanvordeker et al., [52]
319 confirmed that the alkyl side chain of MDZ participates in the binding to proteins. Besides, EPS

contains a diverse range of functional groups, including amine, carboxylate, phosphate, carbonyl, amide, and hydroxyl which can form H-bonds with MDZ.

The removal of MDZ started after day 10 in all cases regardless of MDZ or biomass concentration. Remarkably, MDZ removal and EPS formation occurred almost simultaneously. These results indicated that the MDZ might be removed by EPS solely by adsorption mechanism. Neither metabolites nor antimicrobial biotransformation products were observed in HPLC chromatogram; suggesting that the EPS of *C. vulgaris* might not contain any metabolizing enzyme for MDZ. Therefore, it is postulated that the decrease in MDZ concentration in the bioreactors would only be associated with adsorption phenomena, which eventually played a role in increasing the concentration of EPS. As a control experiment, bound EPS was extracted from 10 μ M high biomass bioreactor, sonicated for 15 minutes, followed by filtration and analysis, where the presence of MDZ was clearly recorded. A bioreactor has a complex system of suspended and flocculated biomass; thus, achieving an accurate mass balance of MDZ was not possible. However, these results showed that 0.69 mg MDZ adsorbed per 100 mg of dry weight of EPS (i.e., corresponding to 59% of initial DMZ) in a 10 μ M MDZ-high biomass bioreactor after 20 days of incubation. The small molecular size of MDZ would favor its adsorption into the scaffold of EPS, while the concentration gradient would play a major role in the diffusion of MDZ into EPS [53].

A comparison between the MDZ removal of low and high biomass-containing bioreactor revealed that the longer the late exponential growth phase, the lower the removal efficiency. Specifically, a longer late exponential growth phase delayed the EPS production. In the case of 5 μ M MDZ bioreactors, the low biomass reactor was in active phase until day 15, while the active or growth phase was over on day 12 in the high biomass reactor. Taking into account both

biomass and EPS, it was clear that EPS initiated the adsorption; however, the higher the biomass, the better was the removal.

3.5. Effect of initial concentrations of MDZ in culture media

Wastewaters from different geographic locations contain different concentrations of pharmaceuticals. Consequently, it is important to examine the MDZ removal from bioreactor as a function of the initial concentration of antibiotic. Two sets of bioreactors with lower and higher biomass concentrations were spiked with different initial concentrations (1-50 μM) of MDZ as shown in Figure 5. In general, the efficiency of bioreactors decreased with an increase in the initial concentration of MDZ for a fixed period of time. The adsorption phenomena in live biological adsorbent are complex processes, especially when the adsorbate had to stimulate the production of adsorbent. As depicted in Fig.5, only 50.80% of MDZ was removed in a low biomass reactor initially containing 1 μM of MDZ. However, MDZ was completely removed in both low as well as high biomass reactors originally containing 5 μM MDZ in 20 and 18 days of experiments, respectively. Nevertheless, concentrations higher than 5 μM of MDZ showed the same trend as the higher the initial concentration of antibiotic, the lower the removal (%) under constant conditions (same inoculum size and incubation period).

It is evident from Table 3 that the EPS production increases with increasing concentration of MDZ in culture; however, this becomes almost constant above 5 μM MDZ. Probably, this was the reason why 1 μM MDZ in low inoculum size test culture was not able to completely remove MDZ, whereas a 5 μM MDZ in low as well as in high inoculum size test cultures were capable to completely remove MDZ from culture media.

3.6. Mechanism of MDZ removal

The removal of MDZ from the bioreactor was confirmed by analyzing the liquid phase using HPLC, where a decrease in concentration was recorded. The absence of new chromatographic peaks during the course of the experiments and the respective analysis validated the hypothesis that *C. vulgaris* might not participate in any biodegradation mechanism. To confirm the mechanism, ethyl acetate was used to desorb the adsorbed MDZ from the flocculated biomass. The sole presence of an MDZ peak from the desorbed sample confirmed that MDZ was only removed by sorption mechanism. These results showed that 0.93 mg of MDZ was adsorbed in 1 g dry weight of *C. vulgaris*. However, comparing this result with those obtained for MDZ removal per 1 g of dry weight of EPS for the same test bioreactor on the same day (i.e. 0.69 mg MDZ/100 mg of dry weight of EPS in 10 μ M MDZ high biomass bioreactor after 20 days of incubation; as mentioned in section 3.4) the amount recovered from dry weight of *C. vulgaris* showed a 9.13% loss in mass balance of MDZ. This decrease in the amount of MDZ suggested its diffusion into the microalgal cell.

It is a general observation that dead microalgae cells have a higher adsorption capacity than live cells [54]. Peng et al., [55] demonstrated that live cells of freshwater algae: *S. obliquus* and *C. pyrenoidosa* adsorbed less progesterone and norgestrel in aqueous media than their dead biomass. Yu et al., [56] explained that the better adsorption capacity of dead biomass was due to loss of permeability control when the cell membrane structures were destroyed.

In this study, the adsorption capacity of live cells with EPS has been compared with live cells without EPS. The first 10 days of culture (Figure 2 (a) and (b)) were considered as cells without EPS while after day 10 cells started to form EPS (especially, attached or bonded) which were considered as live cells with EPS. It is clear from Figure 2 (a) and (b) that there was no removal of MDZ during the initial 10 days (live cells without EPS). The removal of MDZ started with the

EPS formation (i.e. after day 10 in all test bioreactors), which proved that live cells with EPS played a major role in MDZ removal.

However, to confirm if suspended, attached, or both types of EPS were participating in removing MDZ from water, the removal of MDZ from water was compared against the EPSs (suspended, attached, and total EPS) obtained from 1 μ M MDZ test bioreactors (i.e., selected as a representative case) of low and high inoculum (Figure 6). As depicted in Figure 6, it was observed that the attached EPS showed the best fit (i.e., expressed by the highest R^2 value) compared to the other two EPSs (R^2 of suspended EPS=0.49, and R^2 of total EPS=0.88). This high value of correlation coefficient ($R^2 = 0.95$) of the attached EPS indicated a dominant contribution of MDZ removal and good consistency between the 3D scaffold of the attached EPS and the adsorbed antibiotic.

3.7. EC_{50} of *C. vulgaris* for MDZ

The most common toxicity assay is to calculate the effective concentration (EC_{50}) of a stressor (i.e., metronidazole) at which 50% of the growth rate of an organism (microalgae) is inhibited. Photosynthesis is an essential process for the survival of microalgae, which is related to growth and several other metabolic activities. Thus, any disruption in the growth of microalgae during stressors exposure is related to the changes in chlorophyll (Fig. 7a-b). Antibiotics damage thylakoid and chloroplast lipids as they disrupt membranes and other organelles in cells; thus, destroying the photosystem II (PSII) and consequently chlorophylls and carotenoids [57]. Low chlorophyll content, as a stress response to pharmaceuticals, has been previously observed in microalgae *Tetraselmis chui*, *Isochrysis galbana*, and *C. pyrenoidosa* in the presence of thiamphenicol, florfenicol, and chloramphenicol [58].

Two approaches were followed to study the growth inhibition in *C. vulgaris* by MDZ; chronic and acute toxicity of MDZ. The chronic toxicity assay was conducted to assess its adverse effects on an organism as the result of long term exposure, while the acute toxicity is performed for short term exposure. The chronic toxicity of MDZ on *C. vulgaris* was only observed when the antibiotic is added during its growth phase, while acute toxicity was visible when the antibiotic came in contact with microalgal cells at its stationary state. The results showed that the chronic toxicity EC50 for *C. vulgaris* was 83.64 μM (Figure 8a), while for acute toxicity it was 13.63 μM (Figure 8b). The growth inhibition of *C. vulgaris* exposed to increasing concentrations of MDZ (i.e., as measured by the chlorophyll-a yield) is shown in Figure 7a-b for chronic and acute toxicity, respectively.

According to Commission of the European Community (1996), pollutants with an $\text{EC}_{50} > 100 \text{ mg L}^{-1}$ fall under the category of harmless substances, while those with EC_{50} from 10-100 mg L^{-1} are included under the harmful category. Also, substances with EC_{50} between 1-10 mg L^{-1} , and $\text{EC}_{50} < 1 \text{ mg L}^{-1}$ are considered as very toxic. Results from this study indicate that MDZ is harmful to *C. vulgaris* (Figures 8a-b).

Besides the efficiency of *C. vulgaris* for removing DMZ from wastewater, the biomass generated during this treatment would be a valuable feedstock for producing biofuels. Algal biomass has been commonly used as a soil conditioner, plant fertilizer, and animal feed [30]. However, the harvested biomass of *C. vulgaris* would not be recommended as a fertilizer because of its MDZ content. A scheme for biofuel production using MDZ-mediated water is proposed in Figure 9. Rawat et al., [59] have previously demonstrated the production of biogas using algal biomass through anaerobic digestion. The temperatures of these processes vary from 35-80°C [60].

Remarkably, MDZ completely decomposes at 180°C [61], which is significantly lower than the abovementioned lowest temperature of anaerobic digestion processes.

4. Conclusions

In the current study, *C. vulgaris* was used as a novel treatment to remove metronidazole from aqueous media. These results indicated that MDZ stimulated the production of EPS in *C. vulgaris*, which played a major role in the adsorption of this antibiotic. *C. vulgaris* displayed an excellent removal capacity for the target antibiotic at 5µM concentration within the range of 0.05 to 0.5g L⁻¹ inoculum of *C. vulgaris*. These results showed that MDZ was only removed by adsorption mechanism onto the scaffold of the EPS, which slowly diffused into the cells. The removal efficiency decreased with increasing initial concentration of antibiotics. The tolerance of *C. vulgaris* towards MDZ depended upon the stage of the life cycle, and on the exposure time to the antibiotic. However, MDZ is marked as a harmful pollutant to *C. vulgaris*. Nevertheless, the recovery of biofuel from the harvested biomass through anaerobic digestion shows promising for future biofuel applications.

Competing interest

The authors declare no competing interest.

References:

- [1] D. Kang, Q. Zhao, Y. Wu, C. Wu, W. Xiang, Removal of nutrients and pharmaceuticals and personal care products from wastewater using periphyton photobioreactors, *Bioresour. Technol.* 248 (2018) 113–119.
- [2] R. Guo, J. Chen, Application of alga-activated sludge combined system (AASCS) as a novel treatment to remove cephalosporins, *Chem. Eng. J.* 260 (2015) 550–556
- [3] B. Yang, J. Zuo, P. Li, K. Wang, X. Yu, M. Zhang, Effective ultrasound electrochemical degradation of biological toxicity and refractory cephalosporin pharmaceutical wastewater. *Chem. Eng. J.* 287 (2016) 30–37.
- [4] A. Hom-Diaz, Z.N. Norvill, P. Blaquez, T. Vicent, B. Guieysse, Ciprofloxacin removal during secondary domestic wastewater treatment in high rate algal ponds, *Chemosphere* 180 (2017) 33–41.

- [5] J. Tanwar, S. Das, Z. Fatima, S. Hameed, Multidrug resistance: an emerging crisis. *Interdiscip. Perspect. Infect. Dis.* (2014):541340. doi: 10.1155/2014/541340
- [6] C. Bouki, D. Venieri, E. Diamadopoulos, Detection and fate of antibiotic resistant bacteria in wastewater treatment plants: a review. *Ecotoxicol. Environ. Saf.* 91 (2013) 1–9.
- [7] X.H. Wang, A.Y. Lin, Phototransformation of cephalosporin antibiotics in an aqueous environment results in higher toxicity. *Environ. Sci. Technol.* 46 (2012) 12417–12426.
- [8] V. Singh, “Antimicrobial resistance,” in *Microbial Pathogens and Strategies for Combating Them: Science, Technology and Education*, 1 (2013) 291–296.
- [9] M. Popeda, E. Pluciennik, A.K. Bednarek, “Proteins in cancer resistance,” *Postępy Higieny i Medycyny Doświadczalnej*, 68 (2014) 616–632.
- [10] M.J. Ahmed, S.K. Theydan, Microporous activated carbon from Siris seed pods by microwave-induced KOH activation onidazole adsorption, *J. Anal. Appl. Pyrolysis* 99 (2013) 101–109.
- [11] L.A. Dunn, A.G. Burgess, K.G. Krauer, L. Eckmann, P. Vanelle, M.D. Crozet, F.D. Gillin, P. Upcroft, J.A. Upcroft, A new-generation 5-nitroimidazole can induce highly metronidazole-resistant *Giardia lamblia* *In vitro*, *Int. J. Antimicrob. Agents* 36 (2010) 37–42.
- [12] A. Bendesky, D. Menéndez, P. Ostrosky-Wegman, Is metronidazole carcinogenic? *Mutation Res.* 511 (2002) 133–144.
- [13] M. Sakamoto, K. Takeba, T. Sasamoto, T. Kusano, H. Hayashi, S. Kanai, M. Kanda, T. Nagayama, Determination of dimetridazole, metronidazole and ronidazole in salmon and honey by liquid chromatography coupled with tandem mass spectrometry. *Shokuhin Eiseigaku Zasshi*, 52 (2011) 51–58.
- [14] A. Hernandez Ceruelos, L.C. Romero-Quezada, J.C. Ruvalcaba Ledezma, L. Lopez Contreras, Therapeutic uses of metronidazole and its side effects: an update, *Eur Rev Med Pharmacol Sci.* 23 (2019) 397–401.
- [15] W. Cheng, M. Yang, Y. Xie, B. Liang, Z. Fang, E.P. Tsang, Enhancement of mineralization of metronidazole by the electro-Fenton process with a Ce/SnO₂-Sb coated titanium anode, *Chem. Eng. J.* 220 (2013) 214–220.
- [16] S.M.S. Magalhães, C.M. Brêtas, J.M. Brêtas, G.A. Pianetti, M.W. Franco, F.A.R. Barbosa, Toxic concentrations of metronidazole to *Microcystis protocystis*, *Braz. J. Biol.*, 74 (2014) 120–124.
- [17] Q. Jiang, H.H. Ngo, L.D. Nghiem, F.I. Hai, W.E. Price, J. Zhang, S. Liang, L. Deng, W. Guo, Effect of hydraulic retention time on the performance of a hybrid moving bed biofilm reactor-membrane bioreactor system for micropollutants removal from municipal wastewater, *Bioresour Technol.* (2017) Doi.org/10.1016/j.biortech.2017.09.114
- [18] T. Deblonde, C. Cossu-Leguille, P. Hartemann, Emerging pollutants in waste water: A review of the literature, *Int. J. Hygiene and Environ. Health* 214 (2011) 442–448
- [19] Z. Fang, J. Chen, X. Qiu, X. Qiu, W. Cheng, L. Zhu, Effective removal of antibiotic metronidazole from water by nanoscale zero-valent iron particles, *Desalination* 268 (2011) 60–67.
- [20] H. Shemer, Y.K. Kunukcu, K.G. Linden, Degradation of the pharmaceutical Metronidazole via UV, Fenton and photo-Fenton processes, *Chemosphere* 63 (2006) 269–276.

- [21] R.F. Dantas, O. Rossiter, A.K.R. Teixeira, A.S.M. Simoes, V.L. da Silva, Direct UV photolysis of propranolol and metronidazole in aqueous solution, *Chem. Eng. J.* 158 (2010) 143–147.
- [22] J. Rivera-Utrilla, G. Prados-Joya, M. Sanchez-Polo, M.A. Ferro-García, M.I. Bautista-Toledo, Removal of nitroimidazole antibiotics from aqueous solution by adsorption/bioadsorption on activated carbon, *J. Hazard. Mater.* 170 (2009) 298–305.
- [23] D.H. Carrales-Alvarado, R. Ocampo-Perez, R. Leyva-Ramos, J. Rivera-Utrilla, Removal of antibiotic metronidazole by adsorption on various carbon materials from aqueous phase, *J. Colloid Interface Sci.* 436, (2014) 276–285.
- [24] V. Homem, L. Santos, Degradation and removal methods of antibiotics from aqueous matrices – a review, *J. Environ. Manage.* 92 (2011) 2304–2347
- [25] W. Li, Y. Shi, L. Gao, J. Liu, Y. Cai, Occurrence and removal of antibiotics in a municipal wastewater reclamation plant in Beijing, China, *Chemosphere* 92 (2013) 435–444.
- [26] H.Y. Kim, J. Jeon, S. Yu, M. Lee, T.H. Kim, S.D. Kim, Reduction of toxicity of antimicrobial compounds by degradation processes using activated sludge, gamma radiation, and UV, *Chemosphere* 93 (2013) 2480–2487.
- [27] P. Verlicchi, M. Al Aukidy, E. Zambello, Occurrence of pharmaceutical compounds in urban wastewater: removal, mass load and environmental risk after a secondary treatment —a review. *Sci Total Environ* 429 (2012) 123–55.
- [28] F. Polesel, K. Lehnberg, W. Dott, S. Trapp, K.V. Thomas, B.G. Plosz, Factors influencing sorption of ciprofloxacin onto activated sludge: Experimental assessment and modelling implications. *Chemosphere* 119 (2015) 105–111.
- [29] F. Polesel, H.R. Andersen, S. Trapp, B.G. Plosz, Removal of Antibiotics in Biological Wastewater Treatment Systems—A Critical Assessment Using the Activated Sludge Modeling Framework for Xenobiotics (ASM-X). *Environ. Sci. Technol.* 50 (2016) 10316–10334.
- [30] C. Grandclement, I. Seyssiecq, A. Píram, P.W.W. Chung, G. Vanot, N. Tiliacos, N. Roche, P. Doumenq, From the conventional biological wastewater treatment to hybrid processes, the evaluation of organic micropollutant removal: A review, *Water Res.* 111 (2017) 297–317.
- [31] Y. Che, Y. Xia, L. Liu, A.D. Li, Y. Yang, T. Zhang, Mobile antibiotic resistome in wastewater treatment plants revealed by Nanopore metagenomic sequencing, *Microbiome*, 7 (2019) <https://doi.org/10.1186/s40168-019-0663-0>
- [32] W.Q. Guo, Z. He-Shan, L. Shuo, D. Juan-Shan, F. Xiao-Chi, Y. Ren-Li, W. Qing-Lian, R. Nan-Qi, C. Jo-Shu, Removal of cephalosporin antibiotics 7-ACA from wastewater during the cultivation of lipid-accumulating microalgae. *Bioresour. Technol.* 221 (2016) 284–290.
- [33] A. García-Rodríguez, V. Matamoros, C. Fontàs, V. Salvadó, The influence of light exposure, water quality and vegetation on the removal of sulfonamides and tetracyclines: A laboratory-scale study, *Chemosphere* 90 (2013) 2297–2302.
- [34] Y. Du, Y. Feng, R. Guo, J. Chen, Enhancement by the artificial controlled culture for the algal treatment of antibiotic ceftazidime: a three-step response performance and high-removal efficiency, *RSC Adv.*, 5 (2015) 72755–72763
- [35] J. Zhang, D. Fu, J. Wu, Photodegradation of Norfloxacin in aqueous solution containing algae. *J Environ Sci* 24 (2012) 743–749.

- [36] J.Q. Xiong, M.B. Kurade, B.H. Jeon, Biodegradation of Levofloxacin by an acclimated freshwater microalga, *Chlorella vulgaris*. Chem. Eng. J., 313 (2017) 1251-1257.
- [37] J.Q. Xiong, M.B. Kurade, B.H. Jeon, Ecotoxicological effects of enrofloxacin and its removal by monoculture of microalgal species and their consortium, Environ. Pollut. 226 (2017) 486-493.
- [38] S. Hena, N. Fatihah, S. Tabassum, N. Ismail, Three stage cultivation process of facultative strain of *Chlorella sorokiniana* for treating dairy farm effluent and lipid enhancement, Water Res. 80 (2015) 346-356.
- [39] J.Q. Xiong, M.B. Kurade, R.A.I. Abou-Shanab, M.K. Ji, J. Choi, J.O. Kim, B.H. Jeon, Biodegradation of carbamazepine using freshwater microalgae *Chlamydomonas mexicana* and *Scenedesmus obliquus* and the determination of its metabolic fate, Bioresour. Technol. 205 (2016) 183-190.
- [40] X. Zhang, P. Amendola, J.C. Hewson, M. Sommerfeld, Q. Hu, Influence of growth phase on harvesting of *Chlorella zofingiensis* by dissolved air flotation. Bioresour Technol. 116 (2012) 477-84.
- [41] M.K. Danquah, B. Gladman, N. Moheimani, G.M. Forde, Microalgal growth characteristics and subsequent influence on dewatering efficiency. Chem. Eng. J. 151(2009) 73-78.
- [42] L. Xia, R. Huang, Y. Li, S. Song, The effect of growth phase on the surface properties of three oleaginous microalgae (*Botryococcus* sp. FACGB-762, *Chlorella* sp. XJ-445 and *Desmodesmus bijugatus* XJ-231). Front. Microbiol. 8 (2017) doi: 10.3389/fmicb.2017.00922
- [43] A.W. Decho, T. Gutierrez, Microbial Extracellular Polymeric Substances (EPSs) in Ocean Systems Systems. Front. Microbiol. 8 (2017) doi: 10.3389/fmicb.2017.00922.
- [44] R. Henderson, S.A. Parsons, B. Jefferson, The impact of algal properties and pre-oxidation on solid-liquid separation of algae. Water Res. 42 (2008) 1827-1845.
- [45] R. Xiao, Y. Zheng, Overview of microalgal extracellular polymeric substances (EPS) and their applications, Biotechnol Adv. 15 (2016) 1225-1244.
- [46] S. Salim, N.R. Kosterink, N.D. Tchetskoua Wacka, M.H. Vermuë, R.H. Wijffels, Mechanism behind auto-flocculation of unicellular green microalgae *Ettlia texensis*, J.Biotechnol. 174 (2014) 34-38.
- [47] E. Caliskan, S. Gokturk, Adsorption characteristics of sulfamethoxazole and metronidazole on activated carbon, Separation Sci. Technol. 45 (2010) 244-255.
- [48] H.C. Flemming, J. Wingender, The biofilm matrix. Nat. Rev. Microbiol. 8 (2010) 623-633.
- [49] J.L. Asensio, A. Arda, F.J. Canada, J. Jimenez-Baebero, Carbohydrate-aromatic interactions. Acc. Chem. Res. 46 (2013) 946-954.
- [50] M. Kumari, R.B. Sunoj, P.V. Balaji, Exploration of CH- π mediated stacking interactions in saccharide: aromatic residue complexes through conformational sampling. Carbohydr. Res. 361 (2012) 133-140.
- [51] M. Nishio, The CH/ π hydrogen bond in chemistry. Conformation, supramolecules, optical resolution and interactions involving carbohydrates. Phys. Chem. Chem. Phys. 13 (2011) 13873-13900.
- [52] D.R. Sanvordeker, Y.W. Chien, T.K. Lin, H.J. Lambert, Binding of metronidazole and its derivatives to plasma proteins: An assessment of drug binding phenomenon, J. Pharmaceutical Sci. 64 (1975) Doi.org/10.1002/jps.2600641111.

- [53] A.W. Decho, "Localization of quorum sensing by extracellular polymeric substances (EPS): considerations of in situ signaling," in *The Physical Basis of Bacterial Quorum Communication*, ed. S. J. Hagen (New York, NY: Springer), (2015) doi: 10.1007/978-1-4939-1402-9_6
- [54] Y. Tao, W. Li, B. Xue, J. Zhong, S. Yao, Q. Wu, Different effects of copper (II), cadmium (II) and phosphate on the sorption of phenanthrene on the biomass of cyanobacteria, *J. Hazard. Mater.* 261 (2013) 21–28.
- [55] F.Q. Peng, G.G. Ying, B. Yang, S. Liu, H.J. Lai, Y.S. Liu, Z.F. Chen, G.J. Zhou, Biotransformation of progesterone and norgestrel by two freshwater microalgae (*Scenedesmus obliquus* and *Chlorella pyrenoidosa*): transformation kinetics and products identification. *Chemosphere* 95 (2014) 581–588.
- [56] Y. Yu, Y. Zhou, Z. Wang, O. Lopez Torres, R. Guo., J. Chen, Investigation of the removal mechanism of antibiotic ceftazidime by green algae and subsequent microbic impact assessment. *Scientific Report*, (2017) DOI:10.1038/s41598-017-04128-3
- [57] A. Magdaleno, M.E. Saenz, A.B. Juárez, J. Moreton, Effects of six antibiotics and their binary mixtures on growth of *Pseudokirchneriella subcapitata*, *Ecotoxicol. Environ. Safety* 113 (2015) 72–78.
- [58] H.T. Lai, J.H. Hou, C.I. Su, C.L. Chen, Effects of chloramphenicol, florfenicol, and thiamphenicol on growth of algae *Chlorella pyrenoidosa*, *Isochrysis galbana*, and *Tetraselmis chui*, *Ecotoxicol. Environ. Saf.* 72 (2009) 329-334.
- [59] R. Rawat, K. Ranjith, T. Mutanda, F. Bux, Dual role of microalgae: Phycoremediation of domestic wastewater and biomass production for sustainable biofuels production, *Appl. Energy*, 88 (2011) 3411-3424.
- [60] D. Vamvuka, Bio-oil, solid and gaseous biofuels from biomass pyrolysis processes-an overview, *International J. Energy Res.* 35 (2011) 835-862.
- [61] N.A.B. de Souza, A.C.D. Medeiros, A.F.O. Santos, R.O. Macedo, Thermal stability of metronidazole drug and tablets, *Journal of Thermal and Calorimetry*, 72 (2003) 535-538.




Figure 1a: Growth curve of low inoculum size of *C. vulgaris* for control and test cultures at different concentrations (1-50 μM) of MDZ

- The data are the mean of three replicates.
- Refer Supplementary Table 1a for the data and SD




Figure 1b: Growth curve of high inoculum size of *C. vulgaris* for control and test cultures at different concentrations (1-50 μM) of MDZ

- The data are the mean of three replicates.
- Refer Supplementary Table 1b for the data and SD




Figure 2a: Evolution of MDZ concentration during treatment using low inoculum size of *C. vulgaris* containing different concentration (1-50 µM) of MDZ in aqueous media

- The data are the mean of three replicates.
- Refer Supplementary Table 2a for the data and SD




Figure 2b: Evolution of MDZ concentration during treatment using high inoculum size of *C. vulgaris* containing different concentration (1-50 µM) of MDZ in aqueous media

- The data are the mean of three replicates.
- Refer Supplementary Table 2a for the data and SD







Figure 3: Zeta potential of low and high inoculum size of *C. vulgaris* in control culture and culture spiked with 5 µM MDZ on alternate days



H bond between saccharide and MDZ



π - π dispersion bond between aromatic compound of protein (tryptophan) and MDZ



CH- π bond between saccharide and MDZ

Figure 4: Possible bonding between MDZ and components of EPS




Figure 5: Effect of initial MDZ concentration on MDZ removal




Figure 6: Correlations between the amount of attached, suspended and total EPS with MDZ removal from 1 μM test bioreactors with low and high inoculum from day 10 to day20




Figure 7a: Chlorophyll a in *C. vulgaris* during a 240-hour metronidazole chronic toxicity test. The control culture (culture without MDZ) possessed 160.22 µg L⁻¹ after 240 hours




Figure 7b: Chlorophyll a in *C. vulgaris* during a 96-hour metronidazole acute toxicity test. The control culture (culture without MDZ) possessed 240.1 µg L⁻¹ after 96 hours




Figure 8a: Growth inhibition (%) of *C. vulgaris* by MDZ during chronic toxicity test.




Figure 8b: Growth inhibition (%) of *C. vulgaris* by MDZ during chronic toxicity test




Figure 9: Proposed scheme for biogas production using MDZ-containing wastewater.

Table 1: Compositions of BG-11 media in 1L MilliQ

Chemicals	Weight (mg)
K_2HPO_4	75
KH_2PO_4	175
$MgSO_4 \cdot 7H_2O$	75
$NaNO_3$	250
$CaCl_2 \cdot 2H_2O$	25
$NaCl$	25
$Na_2EDTA \cdot 2H_2O$	50
$FeSO_4 \cdot 7H_2O$ in 0.1mL H_2SO_4	4.9
H_3BO_3	11.42
$ZnSO_4 \cdot 7H_2O$	1.412
$MnCl_2 \cdot 4H_2O$	0.232
$CuSO_4 \cdot 5H_2O$	0.252
$Co(NO_3)_2 \cdot 6H_2O$	0.08
$Na_2MoO_4 \cdot 2H_2O$	0.192

Table 2: Specific growth rate of *C. vulgaris* in control and test culture with different concentration of MDZ


Concentration of MDZ	Low inoculum size		High inoculum size	
	Specific growth rate of early exponential phase (d ⁻¹)	Specific growth rate of late exponential phase (d ⁻¹)	Specific growth rate of early exponential phase (d ⁻¹)	Specific growth rate of late exponential phase (d ⁻¹)
Control (without MDZ)	0.279	0.055	0.095	0.036
1µM	0.271	0.065	0.094	0.034
5µM	0.273	0.060	0.096	0.029
10 µM	0.272	0.058	0.097	0.019
25 µM	0.271	0.046	0.093	0.035
50 µM	0.271	0.053	0.091	0.028

Table 3: EPS production in control culture and test bioreactors

Days	Control culture		Low inoculum bioreactors									
			1 μ M		5 μ M		10 μ M		25 μ M		50 μ M	
	EPS (S)*	EPS (A)*	EPS (S)	EPS (A)	EPS (S)	EPS (A)	EPS (S)	EPS (A)	EPS (S)	EPS (A)	EPS (S)	EPS (A)
10	2.31 \pm 0.11	1.45 \pm 0.09	1.62 \pm 0.33	3.01 \pm 0.08	1.71 \pm 0.08	5.54 \pm 0.16	1.76 \pm 0.63	5.12 \pm 0.53	1.97 \pm 0.16	5.62 \pm 0.11	1.88 \pm 0.09	6.32 \pm 1.66
12	7.40 \pm 0.26	7.92 \pm 0.22	2.89 \pm 0.78	8.84 \pm 0.65	2.19 \pm 0.22	15.33 \pm 1.11	2.11 \pm 0.08	15.54 \pm 0.38	2.21 \pm 0.36	15.78 \pm 0.76	2.08 \pm 0.23	16.34 \pm 1.89
14	20.52 \pm 0.46	14.91 \pm 0.28	6.79 \pm 0.54	18.77 \pm 0.54	4.23 \pm 0.23	37.87 \pm 1.29	4.19 \pm 0.43	37.1 \pm 0.97	4.56 \pm 0.69	38.2 \pm 1.06	4.21 \pm 0.25	38.33 \pm 1.27
16	34.82 \pm 0.25	19.28 \pm 0.78	11.19 \pm 1.09	26.54 \pm 0.43	7.88 \pm 0.18	76.45 \pm 2.66	7.12 \pm 0.65	77.15 \pm 1.86	7.19 \pm 0.46	75.29 \pm 1.64	6.89 \pm 0.61	73.32 \pm 2.76
18	42.95 \pm 1.67	30.65 \pm 0.64	21.32 \pm 1.68	38.41 \pm 0.79	20.65 \pm	113.62 \pm 3.54	19.9 \pm 1.06	111.3 \pm 1.69	14.06 \pm 0.87	121.23 \pm 1.39	18.88 \pm 0.28	110.12 \pm 3.18
20	48.76 \pm 1.86	37.50 \pm 2.09	39.98 \pm 1.87	47.72 \pm 1.23	33.67 \pm 1.26	146.1 \pm 2.34	31.04 \pm 0.38	142.32 \pm 2.21	24.76 \pm 1.11	148.99 \pm 2.43	30.03 \pm 0.53	139.31 \pm 5.16
High inoculum bioreactors												
10	4.27 \pm 0.22	2.11 \pm 0.13	3.12 \pm 0.08	9.32 \pm 0.65	3.25 \pm 0.11	12.25 \pm 1.07	3.29 \pm 0.38	12.11 \pm 1.09	3.41 \pm 0.09	12.43 \pm 1.66	3.52 \pm 0.33	13.11 \pm 0.87
12	9.16 \pm 0.31	8.99 \pm 0.76	4.89 \pm 0.18	20.36 \pm 0.23	4.22 \pm 0.18	26.48 \pm 1.76	4.15 \pm 0.79	25.54 \pm 1.24	4.55 \pm 0.97	26.98 \pm 1.45	4.23 \pm 0.27	27.23 \pm 1.56
14	29.12 \pm 1.55	21.34 \pm 1.56	8.77 \pm 0.54	49.42 \pm 1.43	6.78 \pm 0.16	55.15 \pm 1.18	7.12 \pm 0.68	57.75 \pm 1.76	7.67 \pm 0.76	56.67 \pm 2.12	7.45 \pm 0.87	58.43 \pm 0.98
16	37.45 \pm 1.42	34.77 \pm 1.82	13.23 \pm 0.21	59.23 \pm 1.54	11.32 \pm 0.64	107.92 \pm 2.45	12.34 \pm 0.45	78.92 \pm 2.06	12.65 \pm 1.14	76.78 \pm 2.32	10.54 \pm 1.09	76.67 \pm 1.67
18	48.72 \pm 2.64	46.34 \pm 2.44	23.45 \pm 0.67	86.43 \pm 2.69	14.43 \pm 0.98	121.4 \pm 2.27	13.98 \pm 0.88	119.12 \pm 2.46	21.33 \pm 1.12	110.43 \pm 2.34	13.78 \pm 0.95	118.34 \pm 2.65
20	54.66 \pm 1.98	51.86 \pm 1.87	39.18 \pm 2.11	110.9 \pm 2.45	26.76 \pm 1.17	151.6 \pm 3.45	25.08 \pm 1.11	148.78 \pm 2.65	32.65 \pm 1.45	140.01 \pm 4.11	24.44 \pm 1.22	145.55 \pm 3.56

(S)*—Suspended EPS; (A)*--Attached EPS; EPS were measured in mg g⁻¹ dry weight of *C. vulgaris*

Graphical Abstract:



Schematic diagram of MDZ removal in *Chlorella vulgaris* culture