



HAL
open science

Derivative-based generalized sensitivity indices and Sobol' indices

Matieyendou Lamboni

► **To cite this version:**

Matieyendou Lamboni. Derivative-based generalized sensitivity indices and Sobol' indices. Mathematics and Computers in Simulation, 2020, 170, pp.236 - 256. 10.1016/j.matcom.2019.10.017. hal-03488769

HAL Id: hal-03488769

<https://hal.science/hal-03488769v1>

Submitted on 21 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Derivative-based generalized sensitivity indices and Sobol' indices

Matieyendou Lamboni^{a,b}

^aUniversity of Guyane, Department DFRST, 97346 Cayenne, French Guiana, France

^b228-UMR Espace-Dev, University of Guyane, University of Réunion, IRD, University of Montpellier.

Abstract

In uncertainty quantification, multivariate sensitivity analysis (MSA), including variance-based sensitivity analysis, and derivative global sensitivity measure (DGSM) are widely used for assessing the effects of input factors on the model outputs. While MSA allows for identifying the order and the strength of interactions among inputs, DGSM provides only a global effect of inputs by making use of model derivatives. It is interesting to combine the advantages of both approaches and to come up with generalized sensitivity indices (GSIs) from MSA based on model derivatives. First, we derive the mathematical expressions of the total effect and total-interaction effect functionals based on derivatives. Second, we construct minimum variance unbiased estimators (MVUEs) of the total-effect and total-interaction effect covariance matrices, and third, we provide the estimators of the total and total-interaction GSIs as well as their consistency and asymptotic normality. Finally, we demonstrate the applicability of these new results by means of simulations.

Keywords: Derivatives, Generalized sensitivity indices, Matrix norms, MVU estimators, U-statistics.

1. Introduction

Complex mathematical models (either multivariate or single response) are widely used as experimental tools for supporting decision making in natural or human-induced phenomena. They often include numerous uncertain input factors, and it is interesting to assess the effects of input factors on the whole model outputs prior to scenario-building, model-reducing, or model diagnostic activities. Multivariate sensitivity analysis (MSA) ([1; 2; 3; 4; 5]), including variance-based sensitivity analysis (VbSA) ([6; 7; 8; 9]), is the standard way of assessing the impor-

1 tance of input factors on the model output(s) as well as interactions among input
2 factors by making use of the model runs. The estimations of generalized sensitivity
3 indices (GSIs) from MSA, including Sobol' indices, have been largely investigated
4 ([10; 1; 11; 2; 4; 12; 13; 14; 15; 16; 17; 18; 19; 20; 21; 22]). Among the sample-based
5 methods, the estimators of the first-order and total indices proposed in [1; 10; 12]
6 allow for improving the estimates of sensitivity indices, as they are based on mini-
7 mum variance unbiased estimators.

8

9 For response models with available gradients, derivative global sensitivity mea-
10 sure (DGSM) ([23; 24; 25]) is an appropriate way of assessing a global impact of
11 input factors, as it is computationally more attractive than VbSA or MSA (sample-
12 based methods). Given that the DGSM index and the total index (from VbSA)
13 can provide different ranking of input factors, upper bounds and lower bounds of
14 the total index and the total-interaction index based on derivatives were proposed
15 ([26; 27; 28]). An upper bound of the total index (resp. total-interaction index),
16 which is a (known) constant times the DGSM index (resp. cross-derivatives index),
17 is used for the screening purpose. Indeed, while a small value of the upper bound
18 of the Sobol total index means that the associated input does not really act in the
19 model; a big value of the upper bound does not bring much information, regarding
20 factors classification, and big values of upper bounds can happen especially in the
21 case of the total-interaction index ([16; 27]).

22

23 In this paper, we combine the advantages of DGSM and MSA approaches to come
24 up with a new way of computing the GSIs by making use of model derivatives. First,
25 we derive the mathematical expressions of the total effect and total-interaction effect
26 functionals using the model derivatives, cumulative distribution functions (CDFs),
27 and probability density functions (PDFs). Second, we construct minimum variance
28 unbiased estimators (MVUEs) of the total-effect and total-interaction effect covari-

1 ance matrices, and third, we provide estimators of the total and total-interaction
2 GSIs as well as their consistency and asymptotic normality. Finally, we demonstrate
3 the applicability of these new results by means of numerical tests.

4 The paper is organized as follows: in Section 2, we recall two definitions of GSIs
5 using the sensitivity functionals and the Frobenius norm. The sensitivity functionals
6 are the functionals that will lead to access the first-order, total and total-interaction
7 GSIs. Section 3 provides the derivative-based GSIs by deriving the mathematical
8 expressions of the sensitivity functionals using the model derivatives, CDFs, and
9 PDFs. The derivation of the derivative-based total-interaction GSI motivates the
10 proposition of a new Hoeffding decomposition in Section 2. While in Section 4,
11 we propose and study estimators of the GSIs based on model derivatives, Section
12 5 deals with the computational issues. It provides an algorithm for computing the
13 GSIs values and illustrates our approach on test cases. We conclude this work in
14 Section 6.

15 **Notation**

16 This section defines the symbols that will be used throughout the paper. For
17 integer $d \in \mathbb{N}^*$ and $j = 1, \dots, d$, we use $\mu_j(x_j) = \rho_j(x_j) dx_j$ for an absolutely
18 continuous probability measure on an open interval $\Omega_j \subseteq \mathbb{R}$ w.r.t. the Lebesgue
19 measure, X_j for a random variable or factor from μ_j , x_j for a sample value of X_j ,
20 and $\mathbf{X} = \{X_j, j \in \{1, \dots, d\}\}$ for a random vector. We use ρ_j (resp. F_j) for the
21 positive and continuous probability density function (PDF) (resp. the cumulative
22 distribution function: CDF) with $j = 1, \dots, d$. We use $\mu(\mathbf{x})$, Ω , $\rho(\mathbf{x})$, $F(\mathbf{x})$ for
23 the joint probability measure, the joint support, the joint PDF and the joint CDF
24 of \mathbf{X} , respectively, and \mathbb{E} and \mathbb{V} for the expectation and variance taking w.r.t. \mathbf{X} .
25 We use \xrightarrow{D} and \xrightarrow{P} for the convergence in distribution and in probability, respectively.

26

27 For integer $n \in \mathbb{N}^*$, the function $\mathbf{f} : \mathbb{R}^d \rightarrow \mathbb{R}^n$ denotes a deterministic function

1 that includes d input factors \mathbf{X} . We use $u \subseteq \{1, 2, \dots, d\}$ for a non-empty subset
2 of $\{1, 2, \dots, d\}$, $\bar{u} = \{1, 2, \dots, d\} \setminus u$ for the complement of u w.r.t. $\{1, 2, \dots, d\}$,
3 and $|u|$ for its cardinality (i.e., the number of elements in u). For a given u , we
4 use $\mathbf{X}_u = \{X_j, j \in u\}$ for a subset of input factors and $\mathbf{X}_{\sim u} = \{X_j, j \in \bar{u}\}$ for
5 the vector containing all input factors, except \mathbf{X}_u . We have the following partition:
6 $\mathbf{X} = (\mathbf{X}_u, \mathbf{X}_{\sim u})$.

7

For an $n \times n$ square matrix $\Sigma = (\sigma_{ij}, i, j \in \{1, \dots, n\})$, the trace (Tr), the
Frobenius norm ($\|\Sigma\|_F$), and the vectorization ($\text{Vec}(\Sigma)$) of Σ are defined as follows:

$$\text{Tr}(\Sigma) = \sum_{i=1}^n \sigma_{ii},$$

$$\|\Sigma\|_F^2 = \sum_{i=1}^n \sum_{j=1}^n |\sigma_{ij}|^2 = \text{Tr}(\Sigma \Sigma^T),$$

$$\text{Vec}(\Sigma) = [\sigma_{i1} \forall i \in \{1, \dots, n\}; \sigma_{i2} \forall i \in \{1, \dots, n\}; \dots; \sigma_{in} \forall i \in \{1, \dots, n\}]^T.$$

8 By definition, the vectorization of Σ is a vector containing the first column of Σ ,
9 followed by the second column, and so on.

10

The function $\mathbf{f}(\cdot)$ may be subjected to the transformation of the form:

$$D_{\mathbf{w}} \mathbf{f}(\mathbf{X}),$$

11 where $D_{\mathbf{w}}$ is a weighting matrix. In the case of the model outputs with different
12 units, some transformations may be used to obtain unit-less outputs. A classical
13 way to accomplish this is to divide each output by its standard deviation, that is,
14 $D_{\mathbf{w}}^{-1} = \text{diag}(\mathbf{w})$ is a diagonal matrix with \mathbf{w} representing the vector of the standard
15 deviations of the outputs. In the following text, we use $\mathbf{f}(\cdot)$ as either the original
16 function or a given transformation of the latter.

1

2 In what follows, we consider only independent input factors (assumption A1) and
 3 measurable and differentiable functions $\mathbf{f} : \mathbb{R}^d \rightarrow \mathbb{R}^n$ having finite second moments,
 4 that is, $\mathbb{E} [\|\mathbf{f}(\mathbf{X})\|_{L^2}^2] < +\infty$.

5 **2. Multivariate sensitivity analysis: two types of generalized sensitivity** 6 **indices**

7 This section gives two definitions of generalized sensitivity indices, including
 8 Sobol' indices, using the sensitivity functionals. We propose i) a new Hoeffding
 9 decomposition; ii) a link between the sensitivity functionals; and iii) the two main
 10 definitions of GSIs.

11 *2.1. Hoeffding decomposition*

12 Under the independence assumption A1, the multivariate Hoeffding decomposi-
 13 tion ([29]; [30]) is given by

$$\begin{aligned}
 \mathbf{f}(\mathbf{X}) &= \mathbf{f}_\emptyset + \sum_{j=1}^d \mathbf{f}_j(X_j) + \sum_{j_1 < j_2}^d \mathbf{f}_{j_1 j_2}(X_{j_1}, X_{j_2}) + \dots + \mathbf{f}_{1\dots d}(X_1, \dots, X_d) \\
 &= \mathbf{f}_\emptyset + \sum_{\substack{w \subseteq \{1, 2, \dots, d\} \\ |w| > 0}} \mathbf{f}_w(\mathbf{X}_w), \tag{2.1}
 \end{aligned}$$

14 where $\mathbf{f}_\emptyset = \mathbb{E}[\mathbf{f}(\mathbf{X})]$ is the expectation of the model output, $\mathbf{f}_j(X_j) = \mathbb{E}[\mathbf{f}(\mathbf{X})|X_j] - \mathbf{f}_\emptyset$,
 15 and $\mathbf{f}_w(\mathbf{X}_w) = \mathbb{E}[\mathbf{f}(\mathbf{X})|\mathbf{X}_w] - \sum_{v, v \subset w} \mathbf{f}_v(\mathbf{X}_v)$ for a non-empty subset $w \subseteq \{1, 2, \dots, d\}$.

16

17 The functional $\mathbb{E}[\mathbf{f}(\mathbf{X})|\mathbf{X}_u] - \mathbf{f}_\emptyset$ generalizes $\mathbf{f}_j(X_j) = \mathbb{E}[\mathbf{f}(\mathbf{X})|X_j] - \mathbf{f}_\emptyset$ from (2.1)
 18 to cope with any subset \mathbf{X}_u of input factors. It allows for quantifying the single
 19 contribution of the input \mathbf{X}_u . We refer to the latter as the first-order functional,
 20 that is, $\mathbf{f}_u^{fo}(\mathbf{X}_u) = \mathbb{E}[\mathbf{f}(\mathbf{X})|\mathbf{X}_u] - \mathbf{f}_\emptyset$.

1 2.2. *New Hoeffding decomposition: link between the total effect and total-interaction*
2 *effect functionals*

3 The total-effect and total-interaction effect functionals can be easily derived by
4 re-organizing the Hoeffding decomposition. The new decomposition aims at manag-
5 ing the Hoeffding decomposition with only $2^{|u|}$ components, with $u \subseteq \{1, 2, \dots, d\}$
6 and $|u|$ its cardinality. The idea consists in expanding the function $\mathbf{f}(\cdot)$ as a sum
7 of collections of functions defined in Equation (2.1). For a given subset u , a collection
8 (of functions) relies on one component of a partition of the set $\{w, w \subseteq \{1, 2, \dots, d\}\}$
9 into $2^{|u|}$ subsets.

10 **Definition 1.** *Let u be a non-empty subset of $\{1, 2, \dots, d\}$, and $\bar{u} = \{1, 2, \dots, d\} \setminus u$*
11 *be a set containing all the elements of $\{1, 2, \dots, d\}$, except those of u .*
12 *For a given v with $v \subseteq u$, we define the set \mathcal{A}_v as a set containing elements of the*
13 *form $\{v, w\}$ with $w \subseteq \bar{u}$, that is,*

$$\mathcal{A}_v = \{\{v, w\}, w \subseteq \bar{u}\}. \quad (2.2)$$

14 By definition, it is obvious that $v \in \mathcal{A}_v$ and $\mathcal{A}_v \subset \{w, w \subseteq \{1, 2, \dots, d\}\}$. In
15 particular, $\mathcal{A}_\emptyset = \{w, w \subseteq \bar{u}\}$ and $\mathcal{A}_u = \{\{u, w\}, w \subseteq \bar{u}\}$. While \mathcal{A}_u contains all the
16 super-sets of u (i.e., sets that contain u), \mathcal{A}_v contains some super-sets of v but not
17 all of them. Furthermore, $\forall A \in \mathcal{A}_v$, A satisfies the following properties:

- 18 • $\forall j \in v$, then $j \in A$;
- 19
- 20 • $\forall j \in u \setminus v$, then $j \notin A$.

21 Using the set \mathcal{A}_v , with $v \subseteq u$, Lemma 1 gives a partition of $\{w, w \subseteq \{1, 2, \dots, d\}\}$.

22

23 **Lemma 1.** *Let u be a non-empty subset of $\{1, 2, \dots, d\}$, v_1, v_2 be two subsets of u*
24 *(i.e., $v_1 \subseteq u, v_2 \subseteq u$).*

25

26 (i) *If $v_1 \neq v_2$ then we have*

$$\mathcal{A}_{v_1} \cap \mathcal{A}_{v_2} = \emptyset. \quad (2.3)$$

1 (ii) The partition of the set $\{w, w \subseteq \{1, 2, \dots, d\}\}$ is given by

$$\{w, w \subseteq \{1, 2, \dots, d\}\} = \bigcup_{v, v \subseteq u} \mathcal{A}_v. \quad (2.4)$$

2 (iii) If we use \mathcal{B}_{v_1} for the set containing all the super-sets of v_1 , we have

$$\mathcal{B}_{v_1} = \bigcup_{v, v_1 \subseteq v \subseteq u} \mathcal{A}_v. \quad (2.5)$$

3 **Proof.** See Appendix A.

4 □

5 Lemma 1 gives an interesting tool for managing and controlling 2^d elements with
6 only $2^{|u|}$ elements. For instance, when $u = \{j\}$, the set $\{w, w \subseteq \{1, 2, \dots, d\}\}$ is
7 completely controlled by \mathcal{A}_u and \mathcal{A}_\emptyset , and this result was obtained in [26; 13]. When
8 $u = \{j_1, j_2\}$, $\{w, w \subseteq \{1, 2, \dots, d\}\}$ can be managed with the following four sets
9 $\mathcal{A}_\emptyset, \mathcal{A}_{\{j_1\}}, \mathcal{A}_{\{j_2\}}, \mathcal{A}_u$. The set \mathcal{A}_u with $|u| = 1$ (resp. $|u| = 2$) is particularly inter-
10 esting in SA, as it can lead to assess the total effect (resp. the total-interaction effect
11 of the second order) of input factor(s) ([12; 10; 1; 27; 16; 17]). Indeed, $\mathcal{A}_{\{j\}}$ is a set
12 containing all the super-sets of $\{j\}$, and it is sufficient to assess the total effect of
13 X_j . In the same sense, while \mathcal{A}_u can lead to assess the total-interaction effect of the
14 $|u|^{\text{th}}$ order, \mathcal{B}_{v_1} allows for quantifying the total-interaction effect of the $|v_1|^{\text{th}}$ order,
15 with $v_1 \subseteq u$. If $|u| = 1$, the total-interaction effect of X_u comes down to the total
16 effect of X_u .

17

18 Now, if we define $\mathbf{f}_{\mathcal{A}_v}(\mathbf{X}) = \sum_{w, w \in \mathcal{A}_v} \mathbf{f}_w(\mathbf{X}_w)$ with $\mathbf{f}_w(\mathbf{X}_w)$ defined in Equation
19 (2.1), we can give a new decomposition of \mathbf{f} (see Proposition 1).

20 **Proposition 1.** Let u be a non-empty subset of $\{1, 2, \dots, d\}$, $v_1 \subseteq u$ be a subset
21 of u . If assumption A1 (independence of input factors) holds, then

22

23 (i) a new Hoeffding decomposition of \mathbf{f} is given by

$$\mathbf{f}(\mathbf{X}) = \sum_{v, v \subseteq u} \mathbf{f}_{\mathcal{A}_v}(\mathbf{X}_v, \mathbf{X}_{\sim v}), \quad (2.6)$$

1 where the components $\mathbf{f}_{A_v}(\mathbf{x})$, $v \subseteq u$ are mutually orthogonal.

2

3 (ii) The functional

$$\begin{aligned} \mathbf{f}_{v_1}^{sup}(\mathbf{X}) &= \mathbf{f}_{B_{v_1}}(\mathbf{X}) \\ &= \sum_{v, v_1 \subseteq v \subseteq u} \mathbf{f}_{A_v}(\mathbf{X}), \end{aligned} \quad (2.7)$$

4 allows for quantifying the total-interaction effect of \mathbf{X}_{v_1} .

5

6 (iii) The functional

$$\mathbf{f}_u^{tot}(\mathbf{X}) = \sum_{\substack{v, v \subseteq u \\ |v| > 0}} \mathbf{f}_{A_v}(\mathbf{X}_v, \mathbf{X}_{\sim v}), \quad (2.8)$$

7 allows for quantifying the total effect of \mathbf{X}_u .

8 **Proof.** See Appendix B.

9

□

10 In what follows, we refer to $\mathbf{f}_u^{sup}(\mathbf{X})$ as the total-interaction effect functional
 11 (TIEF) of \mathbf{X}_u and \mathbf{f}_u^{tot} as the total effect functional (TEF) of \mathbf{X}_u . Lemma 2 provides
 12 a link between TIEF and TEF.

13 **Lemma 2.** Let $u \subseteq \{1, 2, \dots, d\}$ be a non-empty subset. If assumption A1 holds,
 14 then we have

$$\mathbf{f}_u^{tot}(\mathbf{X}_u, \mathbf{X}_{\sim u}) = \sum_{v, v \subseteq u} (-1)^{|u|-|v|+1} \mathbf{f}_{u \setminus v}^{sup}(\mathbf{X}), \quad (2.9)$$

15 and

$$\mathbf{f}_u^{sup}(\mathbf{X}) = \sum_{v, v \subseteq u} (-1)^{|u|-|v|+1} \mathbf{f}_{u \setminus v}^{tot}(\mathbf{X}). \quad (2.10)$$

16 **Proof.** See Appendix C.

17

□

18 2.3. Definition of generalized sensitivity indices

19 When we use the variance as a measure of the variability of the model outputs,
 20 a definition of the sensitivity indices for the multivariate-response models should be

1 based on the following covariance matrices.

2

3 The first-order covariance matrix of \mathbf{X}_u is given by

$$\Sigma_u = \mathbb{V} [\mathbf{f}_u^{fo}(\mathbf{X})] . \quad (2.11)$$

4 Further, the total-effect covariance matrix of \mathbf{X}_u is given by

$$\Sigma_u^{tot} = \mathbb{V} [\mathbf{f}_u^{tot}(\mathbf{X})] . \quad (2.12)$$

5 Likewise, the total-interaction covariance matrix of \mathbf{X}_u is given by

$$\Sigma_u^{sup} = \mathbb{V} [\mathbf{f}_u^{sup}(\mathbf{X})] . \quad (2.13)$$

6 For the single-response models ($n = 1$), the prioritization of input factors based
7 on the covariance matrices is straightforward, as the covariance matrices are scalars.

8 In the case of the multivariate-response models with $n > 1$, Lamboni [1] proposed to
9 apply matrix norms on the covariance matrices in order to prioritize input factors.

10 In this paper, we consider two types of generalized sensitivity indices from [1].

11

12 **Definition 2.** Let Σ , Σ_u , Σ_u^{sup} , and Σ_u^{tot} be the covariance matrices of the model
13 outputs, the first-order, the total-interaction effect and the total-effect functionals,
14 respectively.

15

16 The first-type GSIs are defined below ([3; 2; 4]).

17 The first-order GSI of \mathbf{X}_u is defined as follows:

$$GSI_u^F = \frac{\left\| \left\| \Sigma_u^{1/2} \right\|_F^2 \right\|_F^2}{\left\| \left\| \Sigma^{1/2} \right\|_F^2 \right\|_F^2} . \quad (2.14)$$

18 Further, the total GSI of \mathbf{X}_u is given by

$$GSI_{T_u}^F = \frac{\left\| \left\| (\Sigma_u^{tot})^{1/2} \right\|_F^2 \right\|_F^2}{\left\| \left\| \Sigma^{1/2} \right\|_F^2 \right\|_F^2} , \quad (2.15)$$

1 and the total-interaction GSI of \mathbf{X}_u is given by

$$GSI_{sup,u}^F = \frac{\left\| (\Sigma_u^{sup})^{1/2} \right\|_F^2}{\left\| \Sigma^{1/2} \right\|_F^2}. \quad (2.16)$$

2 Likewise, the second-type GSIs are defined as follows ([1]):

$$GSI_u^{l_2} = \frac{\left\| \Sigma_u \right\|_F}{n \left\| \Sigma^{1/2} \right\|_F^2}; \quad (2.17)$$

$$3 \quad GSI_{T_u}^{l_2} = \frac{\left\| \Sigma_u^{tot} \right\|_F}{n \left\| \Sigma^{1/2} \right\|_F^2}; \quad (2.18)$$

4 and

$$GSI_{sup,u}^{l_2} = \frac{\left\| \Sigma_u^{sup} \right\|_F}{n \left\| \Sigma^{1/2} \right\|_F^2}. \quad (2.19)$$

Remark 1. The first-type GSIs such as GSI_u^F , $GSI_{T_u}^F$, $GSI_{sup,u}^F$ are equivalent to the classical definition, that is,

$$GSI_u^F = \frac{Tr(\Sigma_u)}{Tr(\Sigma)}, \quad GSI_{T_u}^F = \frac{Tr(\Sigma_u^{tot})}{Tr(\Sigma)}, \quad GSI_{sup,u}^F = \frac{Tr(\Sigma_u^{sup})}{Tr(\Sigma)}.$$

5 In the case of single response models ($n = 1$), the two types GSIs come down to
 6 Sobol' indices. Thus, both types of GSIs extend Sobol' indices to cope with multivari-
 7 ate response models. The l_2 -based definition of GSIs (second-type GSIs) explicitly
 8 includes the off-diagonal elements of the covariance matrices. Therefore, it accounts
 9 for the correlations among the component of sensitivity functionals.

10 3. Generalized sensitivity and Sobol' indices using model derivatives

11 Section 2 provides the definitions of generalized sensitivity indices using the main
 12 sensitivity functionals. To propose the GSIs and Sobol' indices based on model
 13 derivatives, we are going to express the TIEF and TEF as functions of the model
 14 derivatives.

15

16 In what follows, we assume that the function $\mathbf{f} : \mathbb{R}^d \rightarrow \mathbb{R}^n$ is a measurable and
 17 differentiable function with respect to each input (assumption A3). Namely, we use
 18 \mathbf{X} for d input factors and \mathbf{x} for a value of \mathbf{X} ; u for a non-empty subset of $\{1, \dots, d\}$;
 19 $(\mathbf{x}_u, \mathbf{x}_{\sim u})$, $(\mathbf{y}_u, \mathbf{x}_{\sim u})$ and $(\mathbf{z}_u, \mathbf{x}_{\sim u})$ for three sample values of \mathbf{X} . The usual total
 20 differential of \mathbf{f} ($d\mathbf{f}$) with higher-order terms is given as follows ([31]):

$$\begin{aligned}
d\mathbf{f} &= \sum_{j=1}^d \frac{\partial \mathbf{f}}{\partial x_j}(\mathbf{x}) dx_j + \sum_{1 \leq i < j \leq d} \frac{\partial^2 \mathbf{f}}{\partial \mathbf{x}_{\{i,j\}}}(\mathbf{x}) d\mathbf{x}_{\{i,j\}} + \dots + \frac{\partial^d \mathbf{f}}{\partial \mathbf{x}}(\mathbf{x}) d\mathbf{x} \\
&= \sum_{\substack{v, v \subseteq \{1, \dots, d\} \\ |v| > 0}} \frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v}(\mathbf{x}) d\mathbf{x}_v, \tag{3.20}
\end{aligned}$$

1 where $\frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v}(\mathbf{x})$ stands for the $|v|^{\text{th}}$ cross-partial derivatives of each component of \mathbf{f}
2 with respect to each x_j , with $j \in v$. By integrating Equation (3.20), we obtain the
3 increment of \mathbf{f} ([31]), that is,

$$\mathbf{f}(\mathbf{z}_u, \mathbf{x}_{\sim u}) - \mathbf{f}(\mathbf{y}_u, \mathbf{x}_{\sim u}) = \sum_{j \in u} \int_{y_j}^{z_j} \frac{\partial \mathbf{f}}{\partial x_j}(\mathbf{x}) dx_j + \sum_{\substack{v, v \subseteq u \\ |v| > 1}} \int_{\mathbf{y}_v}^{\mathbf{z}_v} \frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v}(\mathbf{x}) d\mathbf{x}_v \tag{3.21}$$

4 When we multiply Equation (3.21) by the probability density function $\rho_u(\mathbf{y}_u)$ and
5 integrate it over the joint support Ω_u , we obtain the total-effect functional given by
6 (see Equation (B.1))

$$\begin{aligned}
\mathbf{f}_u^{\text{tot}}(\mathbf{z}_u, \mathbf{x}_{\sim u}) &= \mathbf{f}(\mathbf{z}_u, \mathbf{x}_{\sim u}) - \int_{\Omega_u} \mathbf{f}(\mathbf{y}_u, \mathbf{x}_{\sim u}) d\mu(\mathbf{y}_u) \\
&= \sum_{j \in u} \int_{\Omega_j} \int_{y_j}^{z_j} \frac{\partial \mathbf{f}}{\partial x_j}(\mathbf{x}) \rho_j(y_j) dx_j dy_j \\
&\quad + \sum_{\substack{v, v \subseteq u \\ |v| > 1}} \int_{\Omega_v} \int_{\mathbf{y}_v}^{\mathbf{z}_v} \frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v}(\mathbf{x}) \rho_v(\mathbf{y}_v) d\mathbf{x}_v d\mathbf{y}_v. \tag{3.22}
\end{aligned}$$

7 Under assumption A1 (independent input factor), $\rho_v(\mathbf{y}_v) = \prod_{j \in v} \rho_j(y_j)$, and the
8 right-hand terms of Equation (3.22) become an iteration of one-dimensional integral.
9 For $j \in u$, consider the measurable function $t_j : \Omega_j \rightarrow [0, 1]$ given by $t_j(x_j) = \frac{x_j - y_j}{z_j - y_j}$.
10 Using (3.22), a change of variables gives

$$\begin{aligned}
\mathbf{f}_u^{tot}(\mathbf{z}_u, \mathbf{x}_{\sim u}) &= \sum_{j \in u} \int_{\Omega_j} \int_0^1 \frac{\partial \mathbf{f}}{\partial x_j} (t_j z_j + y_j(1 - t_j), \mathbf{x}_{\sim j}) \times (z_j - y_j) \times \rho_j(y_j) dt_j dy_j \\
&+ \sum_{\substack{v, v \subseteq u \\ |v| > 1}} \int_{\Omega_v} \int_{[0, 1]^{|v|}} \frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v} ([t_j z_j + y_j(1 - t_j) \forall j \in v], \mathbf{x}_{\sim v}) \\
&\times \prod_{j \in v} (z_j - y_j) \times \rho_j(y_j) dt_v dy_v.
\end{aligned} \tag{3.23}$$

1 Now, we can express the TIEF and TEF as functions of the model derivatives,
2 the cumulative distribution functions (CDF), and the probability density functions
3 (PDF). Theorem 1 provides these results.

4 **Theorem 1.** *Let \mathbf{X} be d input factors, $\mathbf{X}^{(1)}$ be an i.i.d copy of \mathbf{X} , F_j (resp. ρ_j)
5 be the CDF (resp. PDF) of X_j with $j = 1, \dots, d$, and u be a non-empty subset of
6 $\{1, 2, \dots, d\}$. If assumptions*

7
8 *A1: the input factors \mathbf{X} are independent,*

9
10 *A2: the model outputs $\mathbf{f}(\mathbf{X})$ have finite second moments,*

11
12 *A3: each component of \mathbf{f} is measurable and differentiable with respect to \mathbf{x}_u ,*

13
14 *A4: for all $v \subseteq u$, $\frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v}(\mathbf{X})$ are measurable and have finite second moments,*

15
16 *A5: the PDF ρ_j is continuous on the support Ω_j and $\rho_j(x_j) \in]0, +\infty[\forall x_j \in \Omega_j$*

17
18 *hold, then*

19
20 *(i) we have the following expressions of the TEF and TIEF, respectively.*

$$\mathbf{f}_u^{tot}(\mathbf{X}) = \sum_{\substack{v, v \subseteq u \\ |v| > 0}} \mathbb{E}_{\mathbf{X}_u^{(1)}} \left[\frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v} (\mathbf{X}_v^{(1)}, \mathbf{X}_{\sim v}) \prod_{j \in v} \frac{F_j(X_j^{(1)}) \left(1 - \mathbb{I}_{[X_j^{(1)} = X_j]}\right) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \right], \tag{3.24}$$

21 *and*

$$\mathbf{f}_u^{sup}(\mathbf{X}) = (-1)^{|u|+1} \mathbb{E}_{\mathbf{X}_u^{(1)}} \left[\frac{\partial^{|u|} \mathbf{f}}{\partial \mathbf{x}_u} (\mathbf{X}_u^{(1)}, \mathbf{X}_{\sim u}) \prod_{j \in u} \frac{F_j(X_j^{(1)}) \left(1 - \mathbb{I}_{[X_j^{(1)} = X_j]}\right) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \right], \tag{3.25}$$

1 where $\mathbb{E}_{\mathbf{X}_u^{(1)}}$ means that the expectation is taken with respect to $\mathbf{X}_u^{(1)}$, $\prod_{j \in v}$ is the
2 product of a sequence depending on j with $j \in v$, and $\mathbb{I}_{[X_j^{(1)}=X_j]} = 1$ if $X_j^{(1)} = X_j$
3 and 0 otherwise.

4
5 (ii) We have the following expansion of the function \mathbf{f} .

$$6 \quad \mathbf{f}(\mathbf{X}) = \mathbf{f}_\emptyset + \sum_{\substack{v, v \subseteq \{1, \dots, d\} \\ |v| > 0}} \mathbb{E}_{\mathbf{X}^{(1)}} \left[\frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v} (\mathbf{X}_v^{(1)}, \mathbf{X}_{\sim v}) \prod_{j \in v} \frac{F_j(X_j^{(1)}) (1 - \mathbb{I}_{[X_j^{(1)}=X_j]}) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \right], \quad (3.26)$$

7 with $\mathbf{f}_\emptyset = \mathbb{E}[\mathbf{f}(\mathbf{X})]$.

8
9 (iii) The total-interaction effect covariance matrix becomes

$$10 \quad \Sigma_u^{sup} = \mathbb{V} \left[\mathbb{E} \left(\frac{\partial^{|u|} \mathbf{f}}{\partial \mathbf{x}_u} (\mathbf{X}_u^{(1)}, \mathbf{X}_{\sim u}) \prod_{j \in u} \frac{F_j(X_j^{(1)}) (1 - \mathbb{I}_{[X_j^{(1)}=X_j]}) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \mid \mathbf{X} \right) \right]. \quad (3.27)$$

Likewise, the total effect covariance matrix is given by

$$11 \quad \Sigma_u^{tot} = \mathbb{V} \left[\mathbb{E} \left(\sum_{\substack{v, v \subseteq u \\ |v| > 0}} \frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v} (\mathbf{X}_v^{(1)}, \mathbf{X}_{\sim v}) \prod_{j \in v} \frac{F_j(X_j^{(1)}) (1 - \mathbb{I}_{[X_j^{(1)}=X_j]}) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \mid \mathbf{X} \right) \right]. \quad (3.28)$$

12 **Proof.** See Appendix D.

□

From Theorem 1, it comes out that the total effect and total-interaction effect covariance matrices are the first-order covariance matrices of the functions

$$\sum_{\substack{v, v \subseteq u \\ |v| > 0}} \frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v} (\mathbf{X}_v^{(1)}, \mathbf{X}_{\sim v}) \prod_{j \in v} \frac{F_j(X_j^{(1)}) (1 - \mathbb{I}_{[X_j^{(1)}=X_j]}) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})},$$

and

$$\frac{\partial^{|u|} \mathbf{f}}{\partial \mathbf{x}_u} (\mathbf{X}_u^{(1)}, \mathbf{X}_{\sim u}) \prod_{j \in u} \frac{F_j(X_j^{(1)}) (1 - \mathbb{I}_{[X_j^{(1)}=X_j]}) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})},$$

1 respectively. Thus, the estimator of the first-order covariance matrices from [1] can
 2 be adapted to obtain the estimators of GSIs using model derivatives.

3 **Remark 2.** *The derivative-based expressions of the TEF and TIEF in Equations*
 4 *(3.24-3.25) are still suitable for functions that are continuous on the joint support*
 5 *Ω and differentiable almost everywhere.*

6 *It is obvious that the TEF and TIEF are centered, that is,*

$$\mathbb{E} [\mathbf{f}_u^{tot}(\mathbf{X})] = \mathbb{E} [\mathbf{f}_u^{sup}(\mathbf{X})] = \mathbf{0}.$$

7 4. Estimators of generalized sensitivity indices using model derivatives

8 The theory of U-statistics allows for easily deriving the properties of estimators
 9 ([32; 33; 34; 29; 35; 36]). Lamboni [10; 1; 12] introduced the theory of U-statistics in
 10 sensitivity analysis by deriving minimum variance unbiased estimators of (the non-
 11 normalized) GSIs and Sobol' indices as well as the consistency and the asymptotic
 12 normality of such estimators. The main idea consists in i) constructing a kernel
 13 (i.e., a random function which expectation is exactly our parameter of interest);
 14 ii) proposing the estimator of that parameter using the kernel; iii) deriving the
 15 statistical performance of the proposed estimator. In this section, we follow these
 16 main steps to construct estimators of the total-effect, the total-interaction effect
 17 covariance matrices and GSIs by making use of model derivatives.

18 4.1. Kernel functions for the total-effect and the total-interaction effect covariance 19 matrices

20 Let $u \subseteq \{1, \dots, d\}$ be a non-empty subset, $(\mathbf{X}^{(1)}, \mathbf{X}^{(2)})$ be 2 i.i.d copies of \mathbf{X} ,
 21 $(\mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)})$ be 2 i.i.d copies of \mathbf{X}_u . For all $v \subseteq u$ with $|v| > 0$, we define the
 22 weight-derivative function $(\mathbf{d}(\cdot))$ as follows:

$$\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)}) = \frac{\partial^{|\mathbf{v}|} \mathbf{f}}{\partial \mathbf{x}_v}(\mathbf{X}_v^{(3)}, \mathbf{X}_{\sim v}^{(1)}) \prod_{j \in v} \frac{F_j(X_j^{(3)}) \left(1 - \mathbb{I}_{[X_j^{(3)} = X_j^{(1)}]}\right) - \mathbb{I}_{[X_j^{(3)} > X_j^{(1)}]}}{\rho_j(X_j^{(3)})}. \quad (4.29)$$

1 To estimate the total-effect and the total-interaction covariance matrices using
2 model derivatives, we consider two functions with two types of inputs $(\mathbf{X}^{(1)}, \mathbf{X}^{(2)})$
3 and $(\mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)})$. For the total-interaction covariance matrix, we consider the func-
4 tion $\mathbf{K}^{sup}(\cdot)$ given by

$$\begin{aligned} \mathbf{K}^{sup}(\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)}) = & \\ & \frac{1}{4} \left([\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(3)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(3)})] \times [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(4)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(4)})] \right)^T \\ & + \frac{1}{4} \left([\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(4)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(4)})] \times [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(3)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(3)})] \right)^T. \end{aligned} \quad (4.30)$$

5 Likewise, we consider the function $\mathbf{K}^{tot}(\cdot)$ for the total-effect covariance matrix
6 with

$$\begin{aligned} \mathbf{K}^{tot}(\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)}) = & \\ & \frac{1}{4} \left(\sum_{\substack{v, v \subseteq u \\ |v| > 0}} \sum_{\substack{v_1, v_1 \subseteq u \\ |v_1| > 0}} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_v^{(3)})] \times [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_{v_1}^{(4)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_{v_1}^{(4)})] \right)^T \\ & + \frac{1}{4} \left(\sum_{\substack{v, v \subseteq u \\ |v| > 0}} \sum_{\substack{v_1, v_1 \subseteq u \\ |v_1| > 0}} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_{v_1}^{(4)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_{v_1}^{(4)})] \times [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_v^{(3)})] \right)^T. \end{aligned} \quad (4.31)$$

7 The functions $\mathbf{K}^{sup}(\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)})$ and $\mathbf{K}^{tot}(\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)})$ are
8 symmetric under independent permutations of their first arguments $(\mathbf{X}^{(1)}, \mathbf{X}^{(2)})$ and
9 second arguments $(\mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)})$. Indeed, the values of such functions do not change
10 if we permute the position of $\mathbf{X}^{(1)}$ and $\mathbf{X}^{(2)}$ in one hand, and the position of $\mathbf{X}_u^{(3)}$
11 and $\mathbf{X}_u^{(4)}$ in the other hand. Theorem 2 gives other properties of such functions.

12

13 **Theorem 2.** *If assumptions A1-A5 hold, then we have*

$$\mathbb{E} [\mathbf{K}^{sup} (\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)})] = \Sigma_u^{sup}, \quad (4.32)$$

1

$$\mathbb{E} [\mathbf{K}^{tot} (\mathbf{X}^{(1)}, \mathbf{X}^{(2)}, \mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)})] = \Sigma_u^{tot}. \quad (4.33)$$

2 **Proof.** See Appendix E.

3

□

4 Theorem 2 shows that the functions $\mathbf{K}(\cdot)$ (resp. $\mathbf{K}^{tot}(\cdot)$) are unbiased estimators of
 5 Σ_u^{sup} (resp. Σ_u^{tot}). Both functions are called kernels of degree (2, 2) in the theory of
 6 U-statistics of two samples. Theorems 3-4 propose the estimators of Σ_u^{sup} and Σ_u^{tot} .

7 **Theorem 3.** Let $\mathcal{X} = (\mathbf{X}^{(1)}, \mathbf{X}^{(2)})$, $\mathcal{Y} = (\mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)})$, and $\mathcal{X}_i, \mathcal{Y}_i, i = 1, 2, \dots, m$,
 8 be two independent samples of size m from \mathcal{X} and \mathcal{Y} , respectively. If assumptions
 9 A1-A5 hold, then

10

11 (i) the minimum variance unbiased estimator of Σ_u^{sup} for a given m and degree
 12 (2, 2) is given by

$$\begin{aligned} \widehat{\Sigma}_u^{sup} &= \frac{1}{4m} \sum_{i=1}^m \left(\left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,u}^{(3)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,u}^{(3)}) \right] \times \left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,u}^{(4)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,u}^{(4)}) \right]^T \right. \\ &\quad \left. + \left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,u}^{(4)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,u}^{(4)}) \right] \times \left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,u}^{(3)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,u}^{(3)}) \right]^T \right), \end{aligned} \quad (4.34)$$

13 and we have

14

$$\mathbb{E} \left(\widehat{\Sigma}_u^{sup} \right) = \Sigma_u^{sup}. \quad (4.35)$$

15 (ii) If $m \rightarrow +\infty$, $\widehat{\Sigma}_u^{sup}$ is consistent, that is,

$$\widehat{\Sigma}_u^{sup} \xrightarrow{\mathcal{P}} \Sigma_u^{sup}. \quad (4.36)$$

16 (iii) If $m \rightarrow +\infty$, $\text{Vec} \left[\widehat{\Sigma}_u^{sup} \right]$ follows a normal distribution, that is,

$$\sqrt{m} \left(\text{Vec} \left[\widehat{\Sigma}_u^{sup} \right] - \text{Vec} \left[\Sigma_u^{sup} \right] \right) \xrightarrow{\mathcal{D}} \mathcal{N} \left(\mathbf{0}, \mathbb{V} \left(\text{Vec} [\mathbf{K}^{sup} (\mathcal{X}, \mathcal{Y})] \right) \right). \quad (4.37)$$

17 **Proof.** See Appendix F.

1

□

2 **Theorem 4.** Let $\mathcal{X} = (\mathbf{X}^{(1)}, \mathbf{X}^{(2)})$, $\mathcal{Y} = (\mathbf{X}_u^{(3)}, \mathbf{X}_u^{(4)})$, and $\mathcal{X}_i, \mathcal{Y}_i, i = 1, 2, \dots, m$,
 3 be two independent samples of size m from \mathcal{X} and \mathcal{Y} , respectively. If assumptions
 4 A1-A5 hold, then

5

6 (i) the minimum variance unbiased estimator of Σ_u^{tot} for a given m and degree
 7 (2, 2) is given by

$$\begin{aligned} \widehat{\Sigma}_u^{tot} &= \frac{1}{4m} \sum_{i=1}^m \sum_{\substack{v, v \subseteq u \\ |v| > 0}} \sum_{\substack{v_1, v_1 \subseteq u \\ |v_1| > 0}} \\ &\times \left(\left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,v}^{(3)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,v}^{(3)}) \right] \times \left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,v_1}^{(4)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,v_1}^{(4)}) \right]^T \right. \\ &\left. + \left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,v_1}^{(4)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,v_1}^{(4)}) \right] \times \left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,v}^{(3)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,v}^{(3)}) \right]^T \right), \end{aligned} \quad (4.38)$$

8 and we have

9

$$\mathbb{E} \left(\widehat{\Sigma}_u^{tot} \right) = \Sigma_u^{tot}. \quad (4.39)$$

10 (ii) If $m \rightarrow +\infty$, $\widehat{\Sigma}_u^{tot}$ is consistent, that is,

$$\widehat{\Sigma}_u^{tot} \xrightarrow{\mathcal{P}} \Sigma_u^{tot}. \quad (4.40)$$

11 (iii) If $m \rightarrow +\infty$, $\text{Vec} \left[\widehat{\Sigma}_u^{tot} \right]$ follows a normal distribution, that is,

$$\sqrt{m} \left(\text{Vec} \left[\widehat{\Sigma}_u^{tot} \right] - \text{Vec} \left[\Sigma_u^{tot} \right] \right) \xrightarrow{\mathcal{D}} \mathcal{N} \left(\mathbf{0}, \mathbb{V}(\text{Vec}[\mathbf{K}^{tot}(\mathcal{X}, \mathcal{Y})]) \right). \quad (4.41)$$

12 **Proof.** The proof is similar to the proof of Theorem 3.

13

□

14 **Remark 3.** The property of minimum variance is valid for a class of estimators that
 15 are based on the weight-derivative function and make use of the same information
 16 such as the degree (2, 2) and the same sample size (m).

17 Theorems 3-4 give the formulas for computing the TIEF and TEF covariance
 18 matrices. For computing GSIs using matrix norms, we use the formulas proposed in
 19 Corollaries 1-2.

1 **Corollary 1.** *Under assumptions A1-A5,*

2
3 (i) *the minimum variance unbiased estimator of $Tr(\Sigma_u^{sup})$ is given by*

$$\begin{aligned} \widehat{Tr(\Sigma_u^{sup})} &= \frac{1}{2m} \sum_{i=1}^m Tr \left(\left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,u}^{(3)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,u}^{(3)}) \right] \right. \\ &\quad \left. \times \left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,u}^{(4)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,u}^{(4)}) \right]^T \right), \end{aligned} \quad (4.42)$$

4 *and we have*

$$\mathbb{E} \left[\widehat{Tr(\Sigma_u^{sup})} \right] = Tr(\Sigma_u^{sup}); \quad (4.43)$$

$$\widehat{Tr(\Sigma_u^{sup})} \xrightarrow{\mathcal{P}} Tr(\Sigma_u^{sup}); \quad (4.44)$$

$$\sqrt{m} \left(\widehat{Tr(\Sigma_u^{sup})} - Tr(\Sigma_u^{sup}) \right) \xrightarrow{\mathcal{D}} \mathcal{N}(\mathbf{0}, \mathbb{V}(Tr[\mathbf{K}^{sup}(\mathcal{X}, \mathcal{Y})])). \quad (4.45)$$

8 (ii) *A consistent estimator of $\|\Sigma_u^{sup}\|_F$ is given by*

$$\|\widehat{\Sigma_u^{sup}}\|_F = \left\| \widehat{\Sigma_u^{sup}} \right\|_F, \quad (4.46)$$

9 *and we have*

$$\|\widehat{\Sigma_u^{sup}}\|_F \xrightarrow{\mathcal{P}} \|\Sigma_u^{sup}\|_F. \quad (4.47)$$

10 **Proof.** *The proof is obvious bearing in mind Theorem 3, the Slutsky theorem, and*
11 *the linearity of the trace and expectation.*

12 □

13 Likewise, we use the following formulas for computing the non-normalized total
14 GSIs of \mathbf{X}_u .

16 **Corollary 2.** *Under assumptions A1-A5,*

17
18 (i) *the minimum variance unbiased estimator of $Tr(\Sigma_u^{tot})$ is given by*

$$\begin{aligned} \widehat{Tr(\Sigma_u^{tot})} &= \frac{1}{2m} \sum_{i=1}^m \sum_{\substack{v, v \subseteq u \\ |v| > 0}} \sum_{\substack{v_1, v_1 \subseteq u \\ |v_1| > 0}} Tr \left(\left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,v}^{(3)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,v}^{(3)}) \right] \right. \\ &\quad \left. \times \left[\mathbf{d}(\mathbf{X}_i^{(1)}, \mathbf{X}_{i,v_1}^{(4)}) - \mathbf{d}(\mathbf{X}_i^{(2)}, \mathbf{X}_{i,v_1}^{(4)}) \right]^T \right), \end{aligned} \quad (4.48)$$

1 and we have

$$\mathbb{E} \left[\widehat{Tr}(\widehat{\Sigma}_u^{tot}) \right] = Tr(\Sigma_u^{tot}) ; \quad (4.49)$$

$$\widehat{Tr}(\widehat{\Sigma}_u^{tot}) \xrightarrow{\mathcal{P}} Tr(\Sigma_u^{tot}) ; \quad (4.50)$$

$$\sqrt{m} \left(\widehat{Tr}(\widehat{\Sigma}_u^{tot}) - Tr(\Sigma_u^{tot}) \right) \xrightarrow{\mathcal{D}} \mathcal{N} \left(\mathbf{0}, \mathbb{V}(Tr[\mathbf{K}^{tot}(\mathcal{X}, \mathcal{Y})]) \right) . \quad (4.51)$$

5 (ii) A consistent estimator of $\|\Sigma_u^{tot}\|_F$ is given by

$$\|\widehat{\Sigma}_u^{tot}\|_F = \left\| \widehat{\Sigma}_u^{tot} \right\|_F , \quad (4.52)$$

6 and we have

$$\|\widehat{\Sigma}_u^{tot}\|_F \xrightarrow{\mathcal{P}} \|\Sigma_u^{tot}\|_F . \quad (4.53)$$

7 **Proof.** The proof is obvious bearing in mind Theorem 4, the Slutsky theorem, and
8 the linearity of the trace and expectation.

9 □

10 Now, we can derive the estimators of the normalized GSIs proposed in Defini-
11 tion 2. Theorem 5 gives the estimators of both types of GSIs, and it provides the
12 performance of these estimators.

13

14 **Theorem 5.** If assumptions A1-A5 hold, then

15

16 (i) the first-type and total GSI of \mathbf{X}_u is given by

$$\widehat{GSI}_{T_u}^F = \frac{Tr(\widehat{\Sigma}_u^{tot})}{Tr(\widehat{\Sigma})} , \quad (4.54)$$

17 where $\widehat{\Sigma}$ is the classical estimator of the variance-covariance matrix of the model
18 outputs, that is, $\Sigma = \mathbb{V}[\mathbf{f}(\mathbf{X})]$, and it is estimated using M model runs.

19

20 Moreover, if $m \rightarrow +\infty$ and $m/M \rightarrow 0$ then we have

$$\widehat{GSI}_{T_u}^F \xrightarrow{\mathcal{P}} GSI_{T_u}^F , \quad (4.55)$$

21 and

$$\sqrt{m} \left(\widehat{GSI}_{T_u}^F - GSI_{T_u}^F \right) \xrightarrow{\mathcal{D}} \mathcal{N} \left(\mathbf{0}, \frac{\mathbb{V}(Tr[\mathbf{K}^{tot}(\mathcal{X}, \mathcal{Y})])}{(Tr[\Sigma])^2} \right) . \quad (4.56)$$

1 (ii) The first-type and total-interaction GSI of \mathbf{X}_u is given by

$$\widehat{GSI}_{sup,u}^F = \frac{\widehat{Tr}(\widehat{\Sigma}_u^{sup})}{Tr(\widehat{\Sigma})}, \quad (4.57)$$

2 and if $m \rightarrow +\infty$ and $m/M \rightarrow 0$ we have

$$\widehat{GSI}_{sup,u}^F \xrightarrow{\mathcal{P}} GSI_{sup,u}^F. \quad (4.58)$$

$$\sqrt{m} \left(\widehat{GSI}_{sup,u}^F - GSI_{sup,u}^F \right) \xrightarrow{\mathcal{D}} \mathcal{N} \left(\mathbf{0}, \frac{\mathbb{V}(Tr[\mathbf{K}^{sup}(\mathcal{X}, \mathcal{Y})])}{(Tr[\Sigma])^2} \right). \quad (4.59)$$

4 (iii) The second-type (total and total-interaction) GSIs of \mathbf{X}_u are given by

$$\widehat{GSI}_{T_u}^{l_2} = \frac{\|\widehat{\Sigma}_u^{tot}\|_F}{n Tr(\widehat{\Sigma})}, \quad (4.60)$$

5 with

$$\widehat{GSI}_{T_u}^{l_2} \xrightarrow{\mathcal{P}} GSI_{T_u}^{l_2}; \quad (4.61)$$

6 and

$$\widehat{GSI}_{sup,u}^{l_2} = \frac{\|\widehat{\Sigma}_u^{sup}\|_F}{n Tr(\widehat{\Sigma})}, \quad (4.62)$$

7 with

$$\widehat{GSI}_{sup,u}^{l_2} \xrightarrow{\mathcal{P}} GSI_{sup,u}^{l_2}. \quad (4.63)$$

8 **Proof.** See Appendix G.

9

□

10 5. Computational issues

11 This section illustrates the computations of GSIs and Sobol' indices using deriva-
 12 tives of different functions. First, we provide an algorithm for computing the GSIs
 13 using the proposed estimators, and second, we apply such algorithm on different
 14 functions.

15 5.1. Algorithm, design scheme and computational cost

16 For a given sample size m , the following steps allow for computing the total-effect
 17 and total-interaction effect covariance matrices.

1

2 **Algorithm 1.** *Estimations of the d total-effect covariance matrices.*

3

- 4 • *Sample 4 input values (matrices) of type $m \times d$ ($\mathcal{X}_1, \dots, \mathcal{X}_4$)*
- 5 • *For each input factor X_j , replace the j^{th} column of \mathcal{X}_1 and \mathcal{X}_3 with the j^{th}*
6 *column of \mathcal{X}_2 (i.e., X_{2j}) to obtain 2 new matrices ($\mathcal{X}_{1,2j}$, $\mathcal{X}_{3,2j}$)*
- 7 • *Compute the model derivative with respect to X_j for the two input values*
8 *$\mathcal{X}_{1,2j}$, $\mathcal{X}_{3,2j}$ to obtain $\frac{\partial \mathbf{f}}{\partial \mathbf{x}_j}(\mathcal{X}_{1,2j})$ and $\frac{\partial \mathbf{f}}{\partial \mathbf{x}_j}(\mathcal{X}_{3,2j})$*
- *Run the weight-derivative function from Equation (4.29) for ($\mathcal{X}_1 X_{2j}$) (resp. $\mathcal{X}_3 X_{2j}$) as follows:*

$$\mathbf{d}(\mathcal{X}_1 X_{2j}) = \frac{\partial \mathbf{f}}{\partial \mathbf{x}_j}(\mathcal{X}_{1,2j}) \frac{F_j(X_{2j}) (1 - \mathbb{I}_{[X_{2j}=X_{1j}]} - \mathbb{I}_{[X_{2j}>X_{1j}]})}{\rho_j(X_{2j})},$$

9 *with X_{1j} the j^{th} column of \mathcal{X}_1*

- 10 • *Repeat the loop for one more time by replacing \mathcal{X}_2 with \mathcal{X}_4*
- 11 • *Use the estimators in (4.38), (4.54), and (4.60) to obtain the estimates of the*
12 *total-effect covariance matrices and GSIs.*

13 Algorithm 1 is still suitable for computing the total-interaction effect covariance
14 matrices (4.34) and GSIs (4.57, 4.62) of the subset of input factors \mathbf{X}_v , with $|v| > 1$.
15 We should modify the steps ii)-iv) of Algorithm 1 as follows: replace the columns of
16 \mathcal{X}_1 and \mathcal{X}_3 that belong to v with the same columns of \mathcal{X}_2 to obtain 2 new matrices,
17 and compute the weight-derivative function as defined in (4.29).

18

19 The total number of the evaluations of the model derivatives to obtain the values
20 of the d total indices (i.e., computational cost of this algorithm) is $4 \times m \times d$. For
21 the second-order total-interaction indices, up to $4 \times m \times d(d-1)$ evaluations of the
22 model derivatives are necessary for obtaining the indices estimates, as a null partial
23 derivative yields to null cross-partial derivatives. These computational costs depend
24 on the crude estimators used in this paper.

1 *5.2. Test cases*

2 In this section, we performed some numerical tests to assess the effectiveness of
3 our estimations. To illustrate our approach, we considered two types of functions
4 as follows: functions with a small number of inputs ($d = 3$), and functions with a
5 medium number of inputs ($d = 6, d = 10$). We computed the model derivatives
6 using the finite difference method, and we computed the GSIs for two values of the
7 sample size (m), that is, $m = 1000, 5000$ using Sobol's sequence or Quasi-Monte
8 Carlo ([37]). For each sample size (m), we replicated the process of computing the
9 indices $R = 50$ times by changing the seed randomly when sampling input values.

10 *5.2.1. Multivariate Ishigami's function ($d = 3, n = 3$)*

11 The multivariate Ishigami function includes three independent input factors fol-
12 lowing a uniform distribution on $[-\pi, \pi]$, and it provides three outputs ([1]). It is
13 defined as follows:

$$\mathbf{f}(\mathbf{x}) = \begin{bmatrix} \sin(x_1) + 7 \sin^2(x_2) + 0.1 x_3^4 \sin(x_1) \\ \sin(x_1) + 5.896 \sin^2(x_2) + 0.1 x_3^4 \sin(x_1) \\ \sin(x_1) + 6.494 \sin^2(x_2) + 0.125 x_3^4 \sin(x_1) \end{bmatrix}. \quad (5.64)$$

14 The true and estimated values of the GSIs for this function are listed in Table 1.

15 *5.2.2. Block-additive function ($d = 6, n = 1$)*

16 The block-additive function includes six independent inputs following a uniform
17 distribution on $[-1, 1]$ ([27]). It is defined as follows:

$$\mathbf{f}(\mathbf{x}) = \cos(-0.8 - 1.1x_1 + 1.1x_5 + x_3) + \sin(0.5 + 0.9x_4 + x_2 - 1.1x_6). \quad (5.65)$$

18 The true values of Sobol' indices and the estimates are listed in Table 2.

	First-type GSIs			Second-type GSIs		
	$GSI_{T,j}^F$	$\widehat{GSI}_{T,j}^F$		$GSI_{T,j}^{l2}$	$\widehat{GSI}_{T,j}^{l2}$	
		$m = 1000$	5000		$m = 1000$	5000
X_1	0.628	0.639 (0.14)	0.630 (0.06)	0.209	0.213 (0.04)	0.210 (0.02)
X_2	0.372	0.340 (0.35)	0.375 (0.11)	0.125	0.135 (0.09)	0.125 (0.04)
X_3	0.284	0.272 (0.13)	0.277 (0.04)	0.093	0.091 (0.04)	0.092 (0.01)
	$GSI_{sup,u}^F$	$\widehat{GSI}_{sup,u}^F$		$GSI_{sup,u}^{l2}$	$\widehat{GSI}_{sup,u}^{l2}$	
$X_1 : X_2$	0	e-23 (e-22)	3e-24 (e-23)	0	3e-23 (e-22)	9e-24 (e-23)
$X_1 : X_3$	0.284	0.336 (0.74)	0.273 (0.21)	0.093	0.155 (0.22)	0.101 (0.05)
$X_2 : X_3$	0	6e-25 (6e-24)	-6e-25 (3e-24)	0	e-24 (2e-24)	8e-25 (e-24)

Table 1: True and estimated values of two types of generalized sensitivity indices for the multivariate Ishigami function. The estimated GSIs (average over 50 replications) are followed by their standard deviations (in bracket). While the top part of this table focuses on the total GSI, the bottom part deals with the total-interaction GSIs. For concise reporting of small indices, we use e-a for 10^{-a} .

1 *5.2.3. Multivariate Sobol's function ($d = 10, n = 4$)*

- 2 The multivariate Sobol function includes 10 independent input factors following
3 a uniform distribution on $[0, 1]$ ([1]). It is defined as follows:

$$\mathbf{f}(\mathbf{x}) = \begin{bmatrix} \prod_{j=1}^{d=10} \frac{|4x_j - 2| + \mathcal{A}[1,j]}{1 + \mathcal{A}[1,j]} \\ \prod_{j=1}^{d=10} \frac{|4x_j - 2| + \mathcal{A}[2,j]}{1 + \mathcal{A}[2,j]} \\ \prod_{j=1}^{d=10} \frac{|4x_j - 2| + \mathcal{A}[3,j]}{1 + \mathcal{A}[3,j]} \\ \prod_{j=1}^{d=10} \frac{|4x_j - 2| + \mathcal{A}[4,j]}{1 + \mathcal{A}[4,j]} \end{bmatrix}. \quad (5.66)$$

According to the values of \mathcal{A} (matrix of type $4 \times d$), this function has different properties. If

$$\mathcal{A} = \begin{bmatrix} 0 & 0 & 6.52 & 6.52 & 6.52 & 6.52 & 6.52 & 6.52 & 6.52 & 6.52 \\ 0 & 1 & 4.5 & 9 & 99 & 99 & 99 & 99 & 99 & 99 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\ 50 & 50 & 50 & 50 & 50 & 50 & 50 & 50 & 50 & 50 \end{bmatrix},$$

- 4 the values of the GSIs are listed in Table 3.

	First-type GSIs			Second-type GSIs		
	$GSI_{T,j}^F$	$\widehat{GSI}_{T,j}^F$		$GSI_{T,j}^{l2}$	$\widehat{GSI}_{T,j}^{l2}$	
		$m = 1000$	5000		$m = 1000$	5000
X_1	0.231	0.229 (0.02)	0.231 (0.01)	0.231	0.229 (0.02)	0.231 (0.01)
X_2	0.214	0.216 (0.02)	0.213 (0.01)	0.214	0.216 (0.02)	0.213 (0.01)
X_3	0.196	0.196 (0.01)	0.196 (0.004)	0.196	0.196 (0.01)	0.196 (0.004)
X_4	0.176	0.176 (0.01)	0.175 (0.005)	0.176	0.176 (0.01)	0.175 (0.005)
X_5	0.231	0.230 (0.02)	0.231 (0.007)	0.231	0.230 (0.02)	0.231 (0.007)
X_6	0.256	0.254 (0.01)	0.257 (0.006)	0.256	0.254 (0.01)	0.257 (0.006)
	$GSI_{sup,u}^F$	$\widehat{GSI}_{sup,u}^F$		$GSI_{sup,u}^{l2}$	$\widehat{GSI}_{sup,u}^{l2}$	
$X_1 : X_2$	0	2e-20 (e-19)	-9e-22 (6e-21)	0	2e-20 (e-19)	1e-21 (6e-21)
$X_1 : X_3$	0.067	0.069 (0.01)	0.067 (0.006)	0.067	0.069 (0.01)	0.067 (0.006)
$X_1 : X_4$	0	3e-22 (e-21)	-9e-22 (5e-21)	0	8e-22 (e-21)	1e-22 (5e-21)
$X_1 : X_5$	0.078	0.080 (0.01)	0.079 (0.006)	0.078	0.080 (0.01)	0.079 (0.006)
$X_1 : X_6$	0	-4e-21 (3e-20)	-3e-22 (e-21)	0	5e-21 (3e-20)	6e-22 (e-21)
$X_2 : X_3$	0	e-22 (2e-21)	7e-22 (3e-21)	0	9e-22 (2e-21)	9e-22 (3e-21)
$X_2 : X_4$	0.040	0.041 (0.007)	0.040 (0.004)	0.040	0.041 (0.007)	0.040 (0.004)
$X_2 : X_5$	0	6e-23 (e-21)	e-22 (3e-21)	0	5e-22 (e-21)	e-21 (3e-21)
$X_2 : X_6$	0.053	0.054 (0.014)	0.053 (0.005)	0.053	0.054 (0.014)	0.053 (0.005)
$X_3 : X_4$	0	2e-22 (e-21)	-4e-20 (3e-19)	0	4e-22 (e-21)	5e-20 (3e-19)
$X_3 : X_5$	0.067	0.066 (0.010)	0.068 (0.006)	0.067	0.066 (0.010)	0.068 (0.006)
$X_3 : X_6$	0	2e-20 (e-19)	7e-22 (5e-21)	0	2e-20 (e-19)	1e-21 (4e-21)
$X_4 : X_5$	0	-4e-22 (2e-21)	2e-22 (2e-21)	0	9e-22 (2e-21)	1e-21 (2e-21)
$X_4 : X_6$	0.046	0.045 (0.01)	0.045 (0.004)	0.046	0.045 (0.01)	0.045 (0.004)
$X_5 : X_6$	0	-5e-23 (e-21)	e-22 (2e-21)	0	4e-22 (e-21)	8e-22 (2e-21)

Table 2: True and estimated values of Sobol' indices for the block-additive function. The estimated indices (average over 50 replications) are followed by their standard deviations (in bracket). While the top part of this table focuses on the total indices, the bottom part deals with the total-interaction indices. For concise reporting of small indices, we use e-a for 10^{-a} .

1 5.3. Numerical results and discussion

2 Tables 1-3 report the estimated values of sensitivity indices for the three func-
3 tions. From Tables 1-3, it comes out that our estimators give accurate estimates of
4 both GSIs and Sobol' indices in average, and these estimates allow for identifying the
5 most influential input factors. Regarding the precision of our estimates (standard
6 deviations), we obtain better estimates of indices in the case of the block-additive
7 function and the multivariate Sobol function when the sample size is 1000. For the
8 multivariate Ishigami function, we have the worst precision for input X_2 , showing
9 a big impact of the sample size on the estimates and/or a numerical instability of

	First-type GSIs			Second-type GSIs		
	$GSI_{T,j}^F$	$\widehat{GSI}_{T,j}^F$		$GSI_{T,j}^{I2}$	$\widehat{GSI}_{T,j}^{I2}$	
		$m = 1000$	5000		$m = 1000$	5000
X_1	0.605	0.619 (0.19)	0.617 (0.13)	0.147	0.152 (0.05)	0.150 (0.03)
X_2	0.406	0.395 (0.14)	0.409 (0.04)	0.103	0.100 (0.03)	0.101 (0.01)
X_3	0.034	0.034 (0.008)	0.034 (0.003)	0.008	0.008 (0.002)	0.008 (0.001)
X_4	0.021	0.022 (0.006)	0.020 (0.002)	0.005	0.005 (0.002)	0.004 (0.001)
X_5	0.014	0.014 (0.003)	0.015 (0.001)	0.003	0.003 (0.001)	0.004 (0.001)
X_6	0.013	0.012 (0.003)	0.013 (0.002)	0.003	0.003 (0.001)	0.003 (0.0004)
X_7	0.011	0.011 (0.003)	0.011 (0.001)	0.003	0.003 (0.001)	0.003 (0.0003)
X_8	0.011	0.011 (0.003)	0.011 (0.001)	0.002	0.003 (0.001)	0.003 (0.0003)
X_9	0.010	0.010 (0.004)	0.010 (0.002)	0.002	0.002 (0.001)	0.002 (0.0004)
X_{10}	0.009	0.010 (0.002)	0.009 (0.001)	0.002	0.002 (0.001)	0.002 (0.0002)

Table 3: True and estimated values of two types of the total GSIs for the multivariate Sobol function of type A. The estimated GSIs (average over 50 replications) are followed by their standard deviations (in bracket).

1 our estimators. This numerical instability decreases when increasing the sample size
2 ($m = 5000$), and we obtain reasonable precision for all input factors.

3

4 It happens to obtain negative estimates of some null total-interaction indices
5 (Tables 1-2), although the negative estimates are too small (around -10^{-20}). This
6 result seems to suggest the difficulty of our estimator of the total-interaction index
7 to better estimate null values of indices according to the precision required. The
8 second-type GSIs, which are also equivalent to Sobol' indices in the case of single
9 response models, avoid obtaining negatives estimates of indices even for small or
10 null values of indices by definition. Moreover, for the three functions considered in
11 this paper, the second-type GSIs give the same ranking of input factors compared
12 to the first-type GSIs.

13 6. Conclusion

14 In this paper, we propose and study a novel way of computing two types of
15 generalized sensitivity indices, including Sobol' indices, using model derivatives, the
16 cumulative distribution function, and the probability density function. First, we

1 derive the mathematical expressions of the total-effect and total-interaction effect
2 functionals and covariance matrices based on the model derivatives. Second, we con-
3 struct minimum variance unbiased estimators of the total-effect and total-interaction
4 effect covariance matrices, and third, we provide estimators of the total and total-
5 interaction GSIs as well as their consistency and asymptotic normality. Finally, we
6 provide an algorithm for computing these covariance matrices and the GSIs, includ-
7 ing the Sobol indices.

8
9 The numerical tests confirmed the accuracy of our estimates in general, except
10 for some null indices. While a medium value of the sample size ($m = 1000$) gives
11 interesting results for the Sobol and block additive functions, our estimators exhibit
12 a numerical instability in the case of Ishigami's function. Our estimators of the
13 first-type GSIs and Sobol' indices show some difficulties for estimating null indices
14 by providing negative estimates (around -10^{-20}). The second-type GSIs avoid ob-
15 taining negative estimates by definition, and they provide the same ranking of input
16 factors compared to the first-type GSIs for the models considered in this paper. In
17 the next future, it is interesting to i) improve the robustness of our estimators, ii)
18 investigate a dimension-free algorithm for computing our indices using derivatives,
19 iii) compare our estimates with those obtained by using derivatives-free algorithms
20 (i.e., that do not use derivatives), iv) extend our approach to cope with dependent
21 input factors.

22 **Acknowledgments**

23 We would like to thank the two referees for their comments and suggestions that
24 have helped improving this paper. We also thank the Editor in Chief for considering
25 our manuscript.

1 **References**

- 2 [1] M. Lamboni, Multivariate sensitivity analysis: minimum vari-
3 ance unbiased estimators of the first-order and total-effect covari-
4 ance matrices, *Reliability Engineering & System Safety* (2018) –
5 doi:<https://doi.org/10.1016/j.ress.2018.06.004>.
- 6 [2] M. Lamboni, H. Monod, D. Makowski, Multivariate sensitivity analysis to mea-
7 sure global contribution of input factors in dynamic models, *Reliability Engi-
8 neering and System Safety* 96 (2011) 450–459.
- 9 [3] M. Lamboni, D. Makowski, S. Lehuger, B. Gabrielle, H. Monod, Multivariate
10 global sensitivity analysis for dynamic crop models, *Fields Crop Reasearch* 113
11 (2009) 312–320.
- 12 [4] F. Gamboa, A. Janon, T. Klein, A. Lagnoux, Sensitivity indices for multivariate
13 outputs, *Comptes Rendus de l'Académie des Sciences* (2014) In press.
- 14 [5] S. Xiao, Z. Lu, L. Xu, Multivariate sensitivity analysis based on the direction
15 of eigen space through principal component analysis, *Reliability Engineering &
16 System Safety* 165 (2017) 1 – 10.
- 17 [6] I. M. Sobol, Sensitivity analysis for non-linear mathematical models, *Mathe-
18 matical Modelling and Computational Experiments* 1 (1993) 407–414.
- 19 [7] A. Saltelli, K. Chan, E. Scott, *Variance-Based Methods, Probability and Statis-
20 tics*, John Wiley and Sons, 2000.
- 21 [8] R. Ghanem, D. Higdon, H. Owhadi, *Handbook of Uncertainty Quantification*,
22 Springer International Publishing, 2017.
- 23 [9] A. B. Owen, Variance components and generalized Sobol' indices, *SIAM/ASA
24 Journal on Uncertainty Quantification* 1 (1) (2013) 19–41.

- 1 [10] M. Lamboni, Uncertainty quantification: a minimum variance unbiased
2 (joint) estimator of the non-normalized sobol indices, *Stat Papers* (2018) –
3 doi:<https://doi.org/10.1007/s00362-018-1010-4>.
- 4 [11] O. Garcia-Cabrejo, A. Valocchi, Global sensitivity analysis for multivariate out-
5 put using polynomial chaos expansion, *Reliability Engineering & System Safety*
6 126 (2014) 25 – 36.
- 7 [12] M. Lamboni, Global sensitivity analysis: a generalized, unbiased and
8 optimal estimator of total-effect variance, *Statistical Papers* (2016) 1–
9 26doi:10.1007/s00362-016-0768-5.
10 URL <http://dx.doi.org/10.1007/s00362-016-0768-5>
- 11 [13] M. Lamboni, Global sensitivity analysis: an efficient numerical method for
12 approximating the total sensitivity index, *International Journal for Uncertainty*
13 *Quantification* 6 (1) (2016) 1–17.
- 14 [14] E. Plischke, E. Borgonovo, C. L. Smith, Global sensitivity measures from given
15 data, *European Journal of Operational Research* 226 (3) (2013) 536 – 550.
- 16 [15] E. Borgonovo, S. Tarantola, E. Plischke, M. D. Morris, Transformations and
17 invariance in the sensitivity analysis of computer experiments, *Journal of the*
18 *Royal Statistical Society Series B* 76 (5) (2014) 925–947.
- 19 [16] J. Fruth, O. Roustant, S. Kuhnt, Total interaction index: A variance-based
20 sensitivity index for second-order interaction screening, *Journal of Statistical*
21 *Planning and Inference* 147 (2014) 212 – 223.
- 22 [17] R. Liu, A. B. Owen, Estimating mean dimensionality of analysis of variance de-
23 compositions, *Journal of the American Statistical Association* 101 (474) (2006)
24 712–721.

- 1 [18] A. Saltelli, P. Annoni, I. Azzini, F. Campolongo, M. Ratto, S. Tarantola, Vari-
2 ance based sensitivity analysis of model output. Design and estimator for the
3 total sensitivity index, *Computer Physics Communications* 181 (2) (2010) 259–
4 270.
- 5 [19] A. Saltelli, Making best use of model evaluations to compute sensitivity indices,
6 *Computer Physics Communications* 145 (2002) 280–297.
- 7 [20] I. M. Sobol, Global sensitivity indices for nonlinear mathematical models and
8 their Monte Carlo estimates, *Mathematics and Computers in Simulation* 55
9 (2001) 271 – 280.
- 10 [21] A. B. Owen, Better estimation of small Sobol’ sensitivity indices, *ACM Trans.*
11 *Model. Comput. Simul.* 23 (2013) 111–1117.
- 12 [22] B. Sudret, Global sensitivity analysis using polynomial chaos expansions, *Reli-*
13 *ability Engineering & System Safety* 93 (7) (2008) 964 – 979.
- 14 [23] M. Morris, Factorial sampling plans for preliminary computational experiments,
15 *Technometrics* 33 (1991) 161–174.
- 16 [24] I. M. Sobol, S. Kucherenko, Derivative based global sensitivity measures and the
17 link with global sensitivity indices, *Mathematics and Computers in Simulation*
18 79 (2009) 3009–3017.
- 19 [25] S. Kucherenko, M. Rodriguez-Fernandez, C. Pantelides, N. Shah, Monte Carlo
20 evaluation of derivative-based global sensitivity measures, *Reliability Engineer-*
21 *ing and System Safety* 94 (2009) 1135–1148.
- 22 [26] M. Lamboni, B. Iooss, A.-L. Popelin, F. Gamboa, Derivative-based global sen-
23 sitivity measures: General links with Sobol’ indices and numerical tests, *Math-*
24 *ematics and Computers in Simulation* 87 (0) (2013) 45 – 54.

- 1 [27] O. Roustant, J. Fruth, B. Iooss, S. Kuhnt, Crossed-derivative based sensitivity
2 measures for interaction screening, *Mathematics and Computers in Simulation*
3 105 (2014) 105 – 118.
- 4 [28] S. Kucherenko, S. Song, Derivative-based global sensitivity measures and their
5 link with sobol' sensitivity indices. in: Cools R., Nuyens D. (eds) *Monte*
6 *Carlo and Quasi-Monte Carlo methods*, Springer Proceedings in Mathematics
7 & Statistics 163. doi:https://doi.org/10.1007/978-3-319-33507-0_23.
- 8 [29] W. Hoeffding, A class of statistics with asymptotically normal distribution,
9 *Annals of Mathematical Statistics* 19 (1948) 293–325.
- 10 [30] B. Efron, C. Stein, The jackknife estimate of variance, *The Annals of Statistics*
11 9 (1981) 586–596.
- 12 [31] R. Courant, *Differential and integral calculus*, Vol. II, Blackie and Son Limited,
13 Great Britain, 1936.
- 14 [32] E. L. Lehmann, Consistency and unbiasedness of certain nonparametric tests,
15 *Annals of Mathematical Statistics* 22 (1951) 165–179.
- 16 [33] T. S. Ferguson, *A Course in Large Sample Theory*, Chapman-Hall, New York,
17 1996.
- 18 [34] E. L. Lehmann, *Elements of Large Sample Theory*, Springer, 1999.
- 19 [35] W. Hoeffding, A non-parametric test for independence, *Annals of Mathematical*
20 *Statistics* 19 (1948) 546–557.
- 21 [36] N. Sugiura, Multisample and multivariate nonparametric tests based on U-
22 statistics and their asymptotic efficiencies, *Osaka J. Math.* 2 (2) (1965) 385–426.
23 URL <http://projecteuclid.org/euclid.ojm/1200691466>
- 24 [37] C. Dutang, P. Savicky, *randtoolbox: Generating and Testing Random Numbers*,
25 R package version 1.13 (2013).

1 **AppendixA. Proof of Lemma 1**

2 For point (i), as $v_1 \neq v_2$, there exists at least one element $j \in v_2$ and $j \notin v_1$.

3 Bearing in mind the definition and properties of \mathcal{A}_v (see Definition 1), we have:

4 $\forall A_2 \in \mathcal{A}_{v_2}, j \in A_2$. As $j \in (u \setminus v_1)$, we can write $j \notin A_1, \forall A_1 \in \mathcal{A}_{v_1}$. Therefore,

5 $\forall A_1 \in \mathcal{A}_{v_1}, A_1 \notin \mathcal{A}_{v_2}$.

6 For point (ii), it is obvious that $\cup_{v \subseteq u} \mathcal{A}_v \subseteq \{w, w \subseteq \{1, 2, \dots, d\}\}$ by definition. To

7 show that $\{w, w \subseteq \{1, 2, \dots, d\}\} \subseteq \cup_{v \subseteq u} \mathcal{A}_v$, let us consider a set $w_0 \in \{w, w \subseteq \{1, 2, \dots, d\}\}$.

8 If $w_0 \subseteq u$ then there exists $v \subseteq u$ such that $v = w_0$, and therefore $w_0 \in \mathcal{A}_v$. If

9 $w_0 \cap u = \emptyset$ then $w_0 \in \mathcal{A}_\emptyset$ ($v = \emptyset$). If $w_0 = w_1 \cup w_2$ with $w_1 \subseteq u$ and $w_2 \cap u = \emptyset$ then

10 $w_0 \in \mathcal{A}_{w_1} \cup \mathcal{A}_\emptyset$.

11

12 Point (iii) is obvious using point (i).

13 **AppendixB. Proof of Proposition 1**

14 Point (i) is a consequence of Lemma 1 and the fact that the terms in Equation

15 (2.1) are mutually orthogonal and centered.

16

17 Point (ii) is obvious using the set \mathcal{B}_{v_1} from Lemma 1.

18

19 For Point (iii), the functional $\mathbf{f}_u^{tot}(\mathbf{X}) = \sum_{\substack{v \subseteq u \\ |v| > 0}} \mathbf{f}_{\mathcal{A}_v}(\mathbf{X}_v, \mathbf{X}_{\sim v})$ contains all in-

20 formation brought by \mathbf{X}_u to the model, and it is sufficient to assess the overall

21 contribution of input \mathbf{X}_u . We also have ([1])

$$\mathbf{f}_u^{tot}(\mathbf{X}) = \mathbf{f}(\mathbf{X}) - \mathbb{E}[\mathbf{f}(\mathbf{X}) | \mathbf{X}_{\sim u}]. \quad (\text{B.1})$$

1 **AppendixC. Proof of Lemma 2**

2 Bearing in mind (2.7), we have

$$\begin{aligned}
\sum_{v, v \subseteq u} (-1)^{|u|-|v|+1} \mathbf{f}_{u \setminus v}^{sup}(\mathbf{X}) &= \sum_{v, v \subseteq u} (-1)^{|u|-|v|+1} \sum_{w, (u \setminus v) \subseteq w \subseteq u} \mathbf{f}_{\mathcal{A}_w}(\mathbf{X}) \\
&= \sum_{v, v \subseteq u} \sum_{w, (u \setminus v) \subseteq w \subseteq u} (-1)^{|u|-|v|+1} \mathbf{f}_{\mathcal{A}_w}(\mathbf{X}) \quad (\text{C.1})
\end{aligned}$$

3 For $v_1 \subseteq u$ and $|v_1| > 0$, the sum $\sum_{w, (u \setminus v) \subseteq w \subseteq u}$ in Equation (C.1) contains v_1 for
4 $\{v, (u \setminus v) \subseteq v_1\} \equiv \{v = u \setminus t, t \subseteq v_1 \text{ and } t \neq \emptyset\}$. Thus, Equation (C.1) becomes

$$\begin{aligned}
\sum_{v, v \subseteq u} (-1)^{|u|-|v|+1} \mathbf{f}_{u \setminus v}^{sup}(\mathbf{X}) &= \sum_{\substack{v_1, v_1 \subseteq u \\ |v_1| > 0}} \sum_{\substack{t \subseteq v_1 \\ |t| > 0}} (-1)^{|t|+1} \mathbf{f}_{\mathcal{A}_{v_1}}(\mathbf{X}) \\
&= \sum_{\substack{v_1, v_1 \subseteq u \\ |v_1| > 0}} \mathbf{f}_{\mathcal{A}_{v_1}}(\mathbf{X}),
\end{aligned}$$

5 as $\sum_{|t|=1}^{|v_1|} (-1)^{|t|+1} \binom{|v_1|}{|t|} = -\sum_{|t|=1}^{|v_1|} (-1)^{|t|} \binom{|v_1|}{|t|} = 1$. The first result holds using Equa-
6 tion (2.8).

7

8 The last result in (2.10) is obtained using Equation (2.9). Indeed,

$$\begin{aligned}
\sum_{v, v \subseteq u} (-1)^{|u|-|v|+1} \mathbf{f}_{u \setminus v}^{tot}(\mathbf{X}) &= \sum_{v, v \subseteq u} (-1)^{|u|-|v|+1} \sum_{v_1, v_1 \subseteq (u \setminus v)} (-1)^{|u|-|v|-|v_1|+1} \mathbf{f}_{(u \setminus v) \setminus v_1}^{sup}(\mathbf{X}) \\
&= \sum_{v, v \subseteq u} \sum_{v_1, v_1 \subseteq (u \setminus v)} (-1)^{-|v_1|} \mathbf{f}_{u \setminus (v \cup v_1)}^{sup}(\mathbf{X}).
\end{aligned}$$

9 For $w = \{v_1 \cup v\}$, we have $|w| = |v_1| + |v|$ as $v_1 \cap v = \emptyset$. We can see that

1 $\{(v, v_1), v \subset u, v_1 \subset (u \setminus v)\} \equiv \{(w, v_1), w \subset u, v_1 \subseteq w\}$. Thus, we have

$$\begin{aligned} \sum_{v, v \subset u} (-1)^{|v|+|u|-1} \mathbf{f}_{u \setminus v}^{tot}(\mathbf{X}) &= \sum_{w, w \subset u} \sum_{v_1, v_1 \subseteq w} (-1)^{-|v_1|} \mathbf{f}_{u \setminus w}^{sup}(\mathbf{X}) \\ &= \sum_{w, w \subset u} \sum_{|v_1|=0}^{|w|} \binom{|w|}{|v_1|} (-1)^{-|v_1|} \mathbf{f}_{u \setminus w}^{sup}(\mathbf{X}) \\ &= \mathbf{f}_u^{sup}(\mathbf{X}), \end{aligned}$$

2 as $\sum_{|v_1|=0}^{|w|} \binom{|w|}{|v_1|} (-1)^{-|v_1|} = 0$ if $|w| > 0$ and 1 otherwise.

3 AppendixD. Proof of Theorem 1

4 Let start with Point (i). Replacing \mathbf{z}_u with \mathbf{x}_u in Equation (3.23) yields to

$$\begin{aligned} \mathbf{f}_u^{tot}(\mathbf{x}) &= \sum_{j \in u} \int_{\Omega_j} \int_0^1 \frac{\partial \mathbf{f}}{\partial x_j} (t_j x_j + y_j(1-t_j), \mathbf{x}_{\sim j}) \times (x_j - y_j) \times \rho_j(y_j) dt_j dy_j \\ &\quad + \sum_{\substack{v, v \subseteq u \\ |v| > 1}} \int_{\Omega_v} \int_{[0,1]^{|v|}} \frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v} ([t_j x_j + y_j(1-t_j) \forall j \in v], \mathbf{x}_{\sim v}) \\ &\quad \times \prod_{j \in v} (x_j - y_j) \times \rho_j(y_j) d\mathbf{t}_v d\mathbf{y}_v. \end{aligned} \tag{D.1}$$

5 For $j \in u$, consider the measurable function $r_j(y_j) = t_j x_j + y_j(1-t_j)$. We can see
6 that $y_j = \frac{r_j - t_j x_j}{1-t_j}$, $dy_j = \frac{dr_j}{1-t_j}$, and $x_j - y_j = \frac{x_j - r_j}{1-t_j}$. By making a change of variables
7 and bearing in mind Fubini-Lebesgue's theorem, we have

$$\begin{aligned} \mathbf{f}_u^{tot}(\mathbf{x}) &= \sum_{j \in u} \int_{\Omega_j} \frac{\partial \mathbf{f}}{\partial x_j} (r_j, \mathbf{x}_{\sim j}) \left[\int_0^1 \frac{x_j - r_j}{(1-t_j)^2} \rho_j \left(\frac{r_j - t_j x_j}{1-t_j} \right) dt_j \right] dr_j \\ &\quad + \sum_{\substack{v, v \subseteq u \\ |v| > 1}} \int_{\Omega_v} \frac{\partial^{|v|} \mathbf{f}}{\partial \mathbf{x}_v} ([r_j \forall j \in v], \mathbf{x}_{\sim v}) \\ &\quad \times \left[\int_{[0,1]^{|v|}} \prod_{j \in v} \frac{x_j - r_j}{(1-t_j)^2} \rho_j \left(\frac{r_j - t_j x_j}{1-t_j} \right) d\mathbf{t}_v \right] d\mathbf{r}_v. \end{aligned}$$

1 As the integral

$$\begin{aligned}
\int_0^1 \frac{x_j - r_j}{(1 - t_j)^2} \rho_j \left(\frac{r_j - t_j x_j}{1 - t_j} \right) dt_j &= \left[-F_j \left(\frac{r_j - t_j x_j}{1 - t_j} \right) \right]_0^1 \\
&= F_j(r_j) - \lim_{t_j \rightarrow 1} F_j \left(\frac{r_j - t_j x_j}{1 - t_j} \right) \\
&= F_j(r_j) - \mathbb{1}_{[r_j > x_j]} - F_j(r_j) \mathbb{1}_{[r_j = x_j]},
\end{aligned}$$

2 the first result holds, that is,

$$\begin{aligned}
\mathbf{f}_u^{tot}(\mathbf{x}) &= \sum_{j \in u} \int_{\Omega_j} \frac{\partial \mathbf{f}}{\partial x_j}(r_j, \mathbf{x}_{\sim j}) [F_j(r_j)(1 - \mathbb{1}_{[r_j = x_j]}) - \mathbb{1}_{[r_j > x_j]}] dr_j \\
&\quad + \sum_{\substack{v, v \subseteq u \\ |v| > 1}} \int_{\Omega_v} \frac{\partial^{|\mathbf{f}|} \mathbf{f}}{\partial \mathbf{x}_v}([r_j \forall j \in v], \mathbf{x}_{\sim v}) \prod_{j \in v} [F_j(r_j)(1 - \mathbb{1}_{[r_j = x_j]}) - \mathbb{1}_{[r_j > x_j]}] d\mathbf{x}_v,
\end{aligned}$$

3 or equivalently

$$\begin{aligned}
\mathbf{f}_u^{tot}(\mathbf{X}) &= \sum_{j \in u} \mathbb{E}_{X_j^{(1)}} \left[\frac{\partial \mathbf{f}}{\partial X_j}(X_j^{(1)}, \mathbf{X}_{\sim j}) \frac{F_j(X_j^{(1)})(1 - \mathbb{1}_{[X_j^{(1)} = X_j]}) - \mathbb{1}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \right] \\
&\quad + \sum_{\substack{v, v \subseteq u \\ |v| > 1}} \mathbb{E}_{\mathbf{X}_v^{(1)}} \left[\frac{\partial^{|\mathbf{f}|} \mathbf{f}}{\partial \mathbf{x}_v}(\mathbf{X}_v^{(1)}, \mathbf{X}_{\sim v}) \prod_{j \in v} \frac{F_j(X_j^{(1)})(1 - \mathbb{1}_{[X_j^{(1)} = X_j]}) - \mathbb{1}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \right].
\end{aligned}$$

4 To derive the expression of TIEF based on model derivatives, we combine Equa-

1 tion (2.10) and Equation (3.24). Indeed,

$$\begin{aligned}
\mathbf{f}_u^{sup}(\mathbf{X}) &= \sum_{v, v \subset u} (-1)^{|u|-|v|+1} \sum_{\substack{v_1, v_1 \subseteq (u \setminus v) \\ |v_1| > 0}} \mathbb{E}_{\mathbf{X}_{v_1}^{(1)}} \left[\frac{\partial^{|v_1|} \mathbf{f}}{\partial \mathbf{x}_{v_1}} (\mathbf{X}_{v_1}^{(1)}, \mathbf{X}_{\sim v_1}) \right. \\
&\quad \times \left. \prod_{j \in v_1} \frac{F_j(X_j^{(1)})(1 - \mathbb{I}_{[X_j^{(1)}=X_j]}) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \right] \\
&= \sum_{v, v \subset u} \sum_{\substack{v_1, v_1 \subseteq (u \setminus v) \\ |v_1| > 0}} (-1)^{|u|-|v|+1} \mathbb{E}_{\mathbf{X}_{v_1}^{(1)}} \left[\frac{\partial^{|v_1|} \mathbf{f}}{\partial \mathbf{x}_{v_1}} (\mathbf{X}_{v_1}^{(1)}, \mathbf{X}_{\sim v_1}) \right. \\
&\quad \times \left. \prod_{j \in v_1} \frac{F_j(X_j^{(1)})(1 - \mathbb{I}_{[X_j^{(1)}=X_j]}) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \right].
\end{aligned}$$

2 For a given $v_1 \subseteq u$ with $v_1 \neq \emptyset$, we can see that $\{v, v_1 \subseteq (u \setminus v)\} \equiv \{v = u \setminus t, v_1 \subseteq t \subseteq u\}$.

3 Thus, we have

$$\begin{aligned}
\mathbf{f}_u^{sup}(\mathbf{X}) &= \sum_{\substack{v_1, v_1 \subseteq u \\ |v_1| > 0}} \sum_{t, v_1 \subseteq t \subseteq u} (-1)^{|t|+1} \mathbb{E}_{\mathbf{X}_{v_1}^{(1)}} \left[\frac{\partial^{|v_1|} \mathbf{f}}{\partial \mathbf{x}_{v_1}} (\mathbf{X}_{v_1}^{(1)}, \mathbf{X}_{\sim v_1}) \right. \\
&\quad \times \left. \prod_{j \in v_1} \frac{F_j(X_j^{(1)})(1 - \mathbb{I}_{[X_j^{(1)}=X_j]}) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \right] \\
&= (-1)^{|u|+1} \mathbb{E}_{\mathbf{X}_u^{(1)}} \left[\frac{\partial^{|v_1|} \mathbf{f}}{\partial \mathbf{x}_u} (\mathbf{X}_u^{(1)}, \mathbf{X}_{\sim u}) \prod_{j \in u} \frac{F_j(X_j^{(1)})(1 - \mathbb{I}_{[X_j^{(1)}=X_j]}) - \mathbb{I}_{[X_j^{(1)} > X_j]}}{\rho_j(X_j^{(1)})} \right],
\end{aligned}$$

4 due to the fact that

$$\begin{aligned}
\sum_{t, v_1 \subseteq t \subseteq u} (-1)^{|t|+1} &= \sum_{|t|=|v_1|}^{|u|} \binom{|u| - |v_1|}{|t| - |v_1|} (-1)^{|t|+1} = \sum_{i=0}^{|u|-|v_1|} \binom{|u| - |v_1|}{i} (-1)^{i+|v_1|+1} \\
&= (-1)^{|u|+1} \mathbb{I}_{[v_1=u]}.
\end{aligned}$$

5 Point (ii) is a particular case of Equation (3.24). Indeed, if we choose $u =$
6 $\{1, \dots, d\}$, we have $\mathbf{f}_u^{tot}(\mathbf{X}) = \mathbf{f}(\mathbf{X}) - \mathbb{E}[\mathbf{f}(\mathbf{X})]$ (see Equation (B.1)).

7

8 Point (iii) is obvious.

1 **AppendixE. Proof of Theorem 2**

2 First, we establish some useful equalities for the proofs, and second, we derive
3 the two results of Theorem 2. We know that $\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)})$ is centered, that is,
4 $\mathbb{E} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)})] = \mathbf{0}$ (see Remark 2).
5 Bearing in mind that $\mathbf{X}^{(j_1)}$ and $\mathbf{X}_u^{(j_2)}$ are independent for $j_1 = 1, 2$ and $j_2 = 3, 4$,
6 $\mathbf{X}^{(1)}$ and $\mathbf{X}^{(2)}$ (resp. $\mathbf{X}_u^{(3)}$ and $\mathbf{X}_u^{(4)}$) have the same distribution, we have the following
7 equalities.

$$\mathbb{E} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_v^{(3)}) | \mathbf{X}^{(j_1)}] = \mathbb{E}_{\mathbf{X}_v^{(3)}} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_v^{(3)})] = \mathbb{E}_{\mathbf{X}_v^{(4)}} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_v^{(4)})] , \quad (\text{E.1})$$

8 where $\mathbb{E}_{\mathbf{X}_v^{(j_2)}} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_v^{(j_2)})]$ means that the expectation is taken with respect to
9 $\mathbf{X}_u^{(j_2)}$ with $j_1 \in \{1, 2\}$ and $j_2 \in \{3, 4\}$.

10

11 According to Equation (3.25) and using Equation (E.1), the total-interaction
12 covariance matrix is given by

$$\Sigma_u^{sup} = \mathbb{E} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(3)}) \mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(4)})^T] , \quad (\text{E.2})$$

13 with $j_1 \in \{1, 2\}$. Indeed,

$$\begin{aligned} \Sigma_u^{sup} &= \mathbb{E} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(3)}) \mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(4)})^T] \\ &= \mathbb{E} (\mathbb{E} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(3)}) \mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(4)})^T | \mathbf{X}^{(j_1)}]) \\ &= \mathbb{E} (\mathbb{E} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(3)}) | \mathbf{X}^{(j_1)}] \mathbb{E} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(4)})^T | \mathbf{X}^{(j_1)}]) \\ &= \mathbb{E} (\mathbb{E}_{\mathbf{X}_u^{(3)}} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(3)})] \mathbb{E}_{\mathbf{X}_u^{(4)}} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(4)})^T]) \\ &= \mathbb{E} (\mathbb{E}_{\mathbf{X}_u^{(3)}} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(3)})] \mathbb{E}_{\mathbf{X}_u^{(3)}} [\mathbf{d}(\mathbf{X}^{(j_1)}, \mathbf{X}_u^{(3)})^T]) \\ &= \mathbb{V} [\mathbf{f}_u^{sup}(\mathbf{X}^{(j_1)})] \end{aligned}$$

1 To prove Equation (4.32), it is sufficient to show that

$$\begin{aligned} A &= \frac{1}{4} \mathbb{E} \left[[\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(3)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(3)})] \times [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(4)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(4)})]^T \right] \\ &= \frac{1}{2} \Sigma_u^{sup}, \end{aligned}$$

2 bearing in mind the symmetry of Equation (4.30). Indeed,

$$\begin{aligned} A &= \frac{1}{4} \mathbb{E} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(3)}) \mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(4)})^T] - \frac{1}{4} \mathbb{E} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(3)}) \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(4)})^T] \\ &\quad - \frac{1}{4} \mathbb{E} [\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(3)}) \mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(4)})^T] + \frac{1}{4} \mathbb{E} [\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(3)}) \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(4)})^T] \\ &= \frac{1}{2} \Sigma_u^{sup}, \end{aligned}$$

3 as $\mathbb{E} [\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(3)}) \mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(4)})^T] = \mathbb{E} [\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(3)})] \mathbb{E} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(4)})^T] = \mathbf{0}$

4 and $\mathbb{E} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_u^{(3)}) \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_u^{(4)})^T] = 0$.

5

6 The proof of the second results (4.33) is similar to the proof of Equation (4.32).

7 To show (4.33), it is sufficient to prove that

$$\begin{aligned} B &= \frac{1}{4} \mathbb{E} \left(\sum_{\substack{v \subseteq u \\ |v|=1}}^{|u|} \sum_{\substack{v_1 \subseteq u \\ |v_1|=1}}^{|u|} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_v^{(3)})] [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_{v_1}^{(4)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_{v_1}^{(4)})]^T \right) \\ &= \frac{1}{2} \Sigma_u^{tot}. \end{aligned}$$

8 Indeed, as $\mathbb{E} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)}) \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_{v_1}^{(4)})^T] = \mathbb{E} [\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_v^{(3)}) \mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_{v_1}^{(4)})^T] = 0$,

9 $\mathbf{f}_u^{tot}(\mathbf{X}^{(1)}) = \sum_{\substack{v \subseteq u \\ |v|=1}}^{|u|} \mathbb{E}_{\mathbf{X}_v^{(3)}} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)})]$, and $\mathbf{f}_u^{tot}(\mathbf{X}^{(2)}) = \sum_{\substack{v_1 \subseteq u \\ |v_1|=1}}^{|u|} \mathbb{E}_{\mathbf{X}_{v_1}^{(4)}} [\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_{v_1}^{(4)})]$,

1 we have

$$\begin{aligned}
B &= \frac{1}{4} \sum_{\substack{v \subseteq u \\ |v|=1}}^{|u|} \sum_{\substack{v_1 \subseteq u \\ |v_1|=1}}^{|u|} \mathbb{E} \left([\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_v^{(3)})] [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_{v_1}^{(4)}) - \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_{v_1}^{(4)})]^T \right) \\
&= \frac{1}{4} \sum_{\substack{v \subseteq u \\ |v|=1}}^{|u|} \sum_{\substack{v_1 \subseteq u \\ |v_1|=1}}^{|u|} \mathbb{E} [\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)}) \mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_{v_1}^{(4)})^T] \\
&\quad + \frac{1}{4} \sum_{\substack{v \subseteq u \\ |v|=1}}^{|u|} \sum_{\substack{v_1 \subseteq u \\ |v_1|=1}}^{|u|} \mathbb{E} [\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_v^{(3)}) \mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_{v_1}^{(4)})^T] \\
&= \frac{1}{4} \sum_{\substack{v \subseteq u \\ |v|=1}}^{|u|} \sum_{\substack{v_1 \subseteq u \\ |v_1|=1}}^{|u|} \mathbb{E} \left[\mathbb{E}_{\mathbf{X}_v^{(3)}} (\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)})) \mathbb{E}_{\mathbf{X}_{v_1}^{(4)}} (\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_{v_1}^{(4)})^T) \right] \\
&\quad + \frac{1}{4} \sum_{\substack{v \subseteq u \\ |v|=1}}^{|u|} \sum_{\substack{v_1 \subseteq u \\ |v_1|=1}}^{|u|} \mathbb{E} \left[\mathbb{E}_{\mathbf{X}_v^{(3)}} (\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_v^{(3)})) \mathbb{E}_{\mathbf{X}_{v_1}^{(4)}} (\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_{v_1}^{(4)})^T) \right] \\
&= \frac{1}{4} \mathbb{E} \left[\sum_{\substack{v \subseteq u \\ |v|=1}}^{|u|} \mathbb{E}_{\mathbf{X}_v^{(3)}} (\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_v^{(3)})) \sum_{\substack{v_1 \subseteq u \\ |v_1|=1}}^{|u|} \mathbb{E}_{\mathbf{X}_{v_1}^{(3)}} (\mathbf{d}(\mathbf{X}^{(1)}, \mathbf{X}_{v_1}^{(3)})^T) \right] \\
&\quad + \frac{1}{4} \mathbb{E} \left[\sum_{\substack{v \subseteq u \\ |v|=1}}^{|u|} \mathbb{E}_{\mathbf{X}_v^{(3)}} (\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_v^{(3)})) \sum_{\substack{v_1 \subseteq u \\ |v_1|=1}}^{|u|} \mathbb{E}_{\mathbf{X}_{v_1}^{(3)}} (\mathbf{d}(\mathbf{X}^{(2)}, \mathbf{X}_{v_1}^{(3)})^T) \right] \\
&= \frac{1}{2} \Sigma_u^{tot}.
\end{aligned}$$

2 AppendixF. Proof of Theorem 3

3 Knowing that $\Sigma_u^{sup} = \mathbb{E}[\mathbf{K}^{sup}(\mathcal{X}, \mathcal{Y})]$, the corresponding U-statistic of two sam-
4 ple associated with the *kernel* $\mathbf{K}^{sup}(\mathcal{X}, \mathcal{Y})$ is $\widehat{\Sigma}_u^{sup}$.

5 It is obvious that $\text{Vec} \left[\widehat{\Sigma}_u^{sup} \right]$ is also an unbiased estimator of $\text{Vec}[\Sigma_u^{sup}]$, and it is
6 symmetric w.r.t the two types of inputs. Therefore, it follows from the theory of
7 U-statistics that $\text{Vec} \left[\widehat{\Sigma}_u^{sup} \right]$ has a minimum variance for the class of unbiased es-
8 timators that make use of the kernel $\mathbf{K}^{sup}(\mathcal{X}, \mathcal{Y})$ and have finite 4th moment (see
9 [1; 10] for more details). We also have the following results:

1

(a) unbiasedness

$$\mathbb{E} \left[\text{Vec} \left[\widehat{\Sigma}_u^{sup} \right] \right] = \text{Vec}[\Sigma_u^{sup}];$$

(b) consistency

$$\text{Vec} \left[\widehat{\Sigma}_u^{sup} \right] \xrightarrow{\mathcal{P}} \text{Vec}[\Sigma_u^{sup}];$$

(c) asymptotic normality

$$\sqrt{m} \left(\text{Vec} \left[\widehat{\Sigma}_u^{sup} \right] - \text{Vec}[\Sigma_u^{sup}] \right) \xrightarrow{\mathcal{D}} \mathcal{N} \left(\mathbf{0}, \mathbb{V}[\text{Vec}(\mathbf{K}^{sup})] \right).$$

2 Points (a), (b), and (d) are equivalent to Points (i), (ii), and (iii) of Theorem 3.

3 **AppendixG. Proof of Theorem 5**4 The results about the consistency are obvious bearing in mind Corollaries 1-2
5 and the Slutsky theorem.

6

For the asymptotic normality, first, bearing in mind Slutsky's theorem and using
Corollary 1, we can write

$$\sqrt{m} \left(\frac{\text{Tr}(\widehat{\Sigma}_u^{sup})}{\text{Tr}(\widehat{\Sigma})} - \frac{\text{Tr}(\Sigma_u^{sup})}{\text{Tr}(\widehat{\Sigma})} \right) \xrightarrow{\mathcal{D}} \mathcal{N} \left(0, \frac{\mathbb{V}(\text{Tr}[\mathbf{K}^{sup}(\mathcal{X}, \mathcal{Y})])}{[\text{Tr}(\Sigma)]^2} \right).$$

7 Second, we need to show that $\sqrt{m} \left(\frac{\text{Tr}(\widehat{\Sigma}_u^{sup})}{\text{Tr}(\widehat{\Sigma})} - \frac{\text{Tr}(\Sigma_u^{sup})}{\text{Tr}(\widehat{\Sigma})} \right)$ and $\sqrt{m} \left(\frac{\text{Tr}(\widehat{\Sigma}_u^{sup})}{\text{Tr}(\widehat{\Sigma})} - \frac{\text{Tr}(\Sigma_u^{sup})}{\text{Tr}(\Sigma)} \right)$
8 are asymptotically equivalent and therefore have the same asymptotic distribution.

9 Indeed,

$$10 \sqrt{m} \left(\frac{\text{Tr}(\widehat{\Sigma}_u^{sup})}{\text{Tr}(\widehat{\Sigma})} - \frac{\text{Tr}(\Sigma_u^{sup})}{\text{Tr}(\widehat{\Sigma})} \right) - \sqrt{m} \left(\frac{\text{Tr}(\widehat{\Sigma}_u^{sup})}{\text{Tr}(\widehat{\Sigma})} - \frac{\text{Tr}(\Sigma_u^{sup})}{\text{Tr}(\Sigma)} \right) = \sqrt{m} \text{Tr}(\Sigma_u^{sup}) \left(\frac{1}{\text{Tr}(\Sigma)} - \frac{1}{\text{Tr}(\widehat{\Sigma})} \right) \rightarrow 0,$$

1 because, the probability

$$\begin{aligned} \mathbb{P} \left[\sqrt{m} \left| \frac{1}{\widehat{\text{Tr}}(\Sigma)} - \frac{1}{\text{Tr}(\Sigma)} \right| \geq \epsilon \right] &\leq \frac{m}{\epsilon^2} \mathbb{E} \left[\left(\frac{1}{\widehat{\text{Tr}}(\Sigma)} - \frac{1}{\text{Tr}(\Sigma)} \right)^2 \right] \\ &\leq \frac{m}{\epsilon^2} \frac{C}{M} \xrightarrow{m, M \rightarrow +\infty} 0, \end{aligned}$$

- 2 as $\sqrt{M} \left(\frac{1}{\widehat{\text{Tr}}(\Sigma)} - \frac{1}{\text{Tr}(\Sigma)} \right) \xrightarrow{\mathcal{D}} \mathcal{N}(0, C)$, which is obtained by applying the delta method
- 3 to $\sqrt{M} \left(\widehat{\text{Tr}}(\Sigma) - \text{Tr}(\Sigma) \right) \xrightarrow{\mathcal{D}} \mathcal{N}(0, C_0)$.