



HAL
open science

Right hemisphere superiority for executive control of attention

Alfredo Spagna, Tae Hyeong Kim, Tingting Wu, Jin Fan

► **To cite this version:**

Alfredo Spagna, Tae Hyeong Kim, Tingting Wu, Jin Fan. Right hemisphere superiority for executive control of attention. *Cortex*, 2020, 122, pp.263 - 276. 10.1016/j.cortex.2018.12.012 . hal-03489504

HAL Id: hal-03489504

<https://hal.science/hal-03489504v1>

Submitted on 7 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

RUNNING HEAD: Right hemisphere advantage for executive control of attention

Right hemisphere superiority for executive control of attention

Alfredo Spagna^{1,2*}, Tae Hyeong Kim³, Tingting Wu³, Jin Fan^{3,4*}

¹Department of Psychology, Columbia University in the city of New York , NY, USA

²Institute du Cerveau et de la Moelle épinière, ICM, Inserm U 1127, CNRS UMR 7225,
Sorbonne Université, Paris, France

³Department of Psychology, Queens College, the City University of New York, Flushing, NY,
USA

Departments of ⁴Psychiatry and ⁵Neuroscience, Icahn School of Medicine at Mount Sinai, New
York, NY, USA

* Correspondence should be addressed to:

Alfredo Spagna, Ph. D.

Department of Psychology,

Columbia University in the city of New York , NY, USA

As5559@columbia.edu

Jin Fan, Ph.D.

Department of Psychology,

Queens College, CUNY

65-30 Kissena Blvd, Flushing, NY

jin.fan@qc.cuny.edu

Introduction

Executive control of attention plays a central role in human cognition as the mechanism that allows the selection and prioritization of the processing of goal-relevant information to reach consciousness, and is achieved via the interplay with the other two components of attention, the alerting function for the preparation and maintenance of a state of readiness, and the orienting function for the direction of attention towards relevant features of a stimulus (Fan et al., 2009; Petersen & Posner, 2012; Spagna, Mackie, & Fan, 2015). Evidence in favor of a right hemisphere dominance for orienting of attention came initially from studies of patients with unilateral hemispatial neglect (Bartolomeo, 2007; Bartolomeo & Chokron, 2002; Chica et al., 2012; Heilman & Van Den Abell, 1980; Kinsbourne, 1987; Lunven & Bartolomeo, 2016; Mesulam, 1999; Posner, Cohen, & Rafal, 1982; Posner, Walker, Friedrich, & Rafal, 1984), and were later supported by most of the behavioral patterns showing better performance in orienting to stimuli presented in the left visual field (processed by the right hemisphere, RH) compared to stimuli presented in the right visual field (processed by the left hemisphere, LH) in healthy individuals (De Schotten et al., 2011; Kincade, Abrams, Astafiev, Shulman, & Corbetta, 2005; Marotta, Lupianez, & Casagrande, 2012; Shulman et al., 2009; Smigasiewicz, Westphal, & Verleger, 2017; Vossel, Weidner, Driver, Friston, & Fink, 2012; Zago et al., 2017; Zuanazzi & Cattaneo, 2017). Because the purpose of orienting is to ultimately facilitate the processing of information and conflict resolution (Callejas, Lupianez, Funes, & Tudela, 2005; Callejas, Lupianez, & Tudela, 2004; Fan et al., 2009) by focusing more on the task-relevant feature or location of the imperative stimuli and therefore achieve cognitive control (Mackie, Van Dam, & Fan, 2013), theoretically there should be a RH superiority for the executive control of attention. However, there is inconsistent evidence regarding the right-lateralization of the other specific

Right hemisphere advantage for executive control

functions of attention, especially for the executive control (Asanowicz, Marzecova, Jaskowski, & Wolski, 2012; Greene et al., 2008; Konrad et al., 2005), with some studies showing better conflict resolution in the RH compared to the LH (Asanowicz et al., 2012; Garavan, Ross, & Stein, 1999; Mecklinger, von Cramon, Springer, & Matthes-von Cramon, 1999; Spielberg et al., 2011), but not in others (Greene et al., 2008; Konrad et al., 2005; Spagna, Martella, Fuentes, Marotta, & Casagrande, 2016; Wu, Weissman, Roberts, & Woldorff, 2007). Therefore, whether a RH advantage exists for executive control remains unclear, and determining the laterality of this function will further support the notion of RH dominance for attention (Heilman & Van Den Abell, 1980; Kinsbourne, 1987; Mesulam, 1981; Weintraub & Mesulam, 1987).

The pivotal role of the RH in the allocation of attentional resources, especially in the visuospatial environment, was initially (and consistently throughout the past 40 years) shown by studies in patients with unilateral hemispatial neglect, a neurological disorder that often follows a lesion to the right parietal cortex and / or underlying white matter which results in patients failing to direct their attention to the contra-lesional (i.e, left) side of the space (Bartolomeo & Chokron, 2002; Chica, Thiebaut de Schotten, Bartolomeo, & Paz-Alonso, 2018; Corbetta & Shulman, 2011; Heilman & Van Den Abell, 1980; Lunven & Bartolomeo, 2016; Mesulam, 1999; Posner et al., 1984; Rafal, 1994; Rastelli et al., 2013; Toba et al., 2018). This deficit in the ability to select information (in other terms “orient”) for goal-directed behavior following RH damage has been studied in a great deal of literature regarding the interaction between hemisphere and this function of attention (e.g., Benwell, Thut, Grant, & Harvey, 2014; Chica et al., 2012; Foxe, McCourt, & Javitt, 2003; Longo, Trippier, Vagnoni, & Lourenco, 2015; Marotta et al., 2012; Shulman et al., 2010; Spagna et al., 2016). Based on this evidence, the *hemispatial theory* (Heilman & Van Den Abell, 1980) and the *interhemispheric competition account* (Kinsbourne,

Right hemisphere advantage for executive control

1987) are two models proposing alternative mechanisms underlying the RH advantage of attention: while the former stated that the RH advantage is rather a “disadvantage” of the LH that is able to orient only towards the contralateral visual field (as opposed to the RH that can direct attention to both visual fields, hence the dominance), the latter proposed that each hemisphere has its own “contralateral vector of attention”, and that a lesion to one of the two hemispheres disrupts the normal balance and favors the orienting towards the ipsilesional visual field. Perhaps due to the involvement of both hemispheres in orienting, as shown in a wealth of neurophysiological studies (Corbetta, Patel, & Shulman, 2008; Corbetta & Shulman, 2002; Fan, McCandliss, Fossella, Flombaum, & Posner, 2005; Xuan et al., 2016), behavioral studies have revealed inconsistent results concerning this function of attention, with some studies revealing a RH dominance (exp. 1 of Asanowicz et al., 2012; Evert, McGlinchey-Berroth, Verfaellie, & Milberg, 2003; Greene et al., 2008; Kałamała, Drożdżowicz, Szewczyk, Marzecová, & Wodniecka, 2018; Poynter, Ingram, & Minor, 2010; Śmigasiewicz, Asanowicz, Westphal, & Verleger, 2014), while other studies failed to find a hemispheric asymmetry (exp. 2 of Greene et al., 2008; Spagna et al., 2016; Tao, Marzecová, Taft, Asanowicz, & Wodniecka, 2011). Controversies also exist regarding the RH superiority of the other two components, the alerting and executive control function. Behavioral studies have not found significant visual field difference for the alerting network in healthy individuals (Asanowicz et al., 2012; Greene et al., 2008; Marzecova, Asanowicz, KrivÁ, & Wodniecka, 2012; Spagna et al., 2016), possibly due to the simplicity of the testing procedure used, as stronger attentional asymmetries have been shown to come from more difficult tasks (Jonides, 1979; Verleger et al., 2009; Welcome & Chiarello, 2008). Nonetheless, a greater deficit in alerting has been found after RH damage to regions of the parietal lobe partially overlapping with those related to the hemispatial neglect

Right hemisphere advantage for executive control

discussed above (Fernandez-Duque & Posner, 2001; Petersen & Posner, 2012; see also Posner, 2008). For the executive control, behavioral evidence of an hemispherical asymmetry mostly derives from a lateralized version of the Stroop task (Stroop, 1935), a task that heavily relies on the well-known left-lateralized language function, and a RH advantage for conflict processing has been found in some studies (Asanowicz et al., 2012; Kałamała et al., 2018; Marzecova et al., 2012; Poynter et al., 2010; Weekes & Zaidel, 1996) but not in others (Belanger & Cimino, 2002; Greene et al., 2008; Konrad et al., 2005; Spagna et al., 2016). Overall, whether a RH dominance of attention exists across all three functions remains unclear, along with the mechanisms responsible for such unilateral advantage of attention.

A potential reason for the above-mentioned inconsistent results, especially regarding the executive control function, may reside in the neglected consideration of the interplay among the components (Badre, 2011) that leads to the unitary construct of attention. Although initially conceived as three independent components, both in terms of function (Fan, McCandliss, Sommer, Raz, & Posner, 2002) and associated brain structure (Fan et al., 2005), the interplay between the alerting, orienting, and executive control was soon after identified as a key component in the prioritization of mental computations (Callejas et al., 2005; Callejas et al., 2004; Chica et al., 2012; Fan et al., 2009; Liu, Bengson, Huang, Mangun, & Ding, 2016; Macaluso, 2010; Martella, Casagrande, & Lupianez, 2011; Spagna, Dong, et al., 2015; Spagna, Mackie, et al., 2015; Spagna et al., 2014). For example, efficient dynamics among these functions may ameliorate the inattentive symptoms associated with hemispatial neglect (Chica et al., 2012), which may open critical rehabilitation perspectives (Manly, Hawkins, Evans, Woldt, & Robertson, 2002; Robertson, Manly, Andrade, Baddeley, & Yiend, 1997). Evidence of modulatory effects involving the alerting function were first observed in studies showing the

Right hemisphere advantage for executive control

beneficial effect of an alerting cue on the orienting function (Fernandez-Duque & Posner, 1997; Fuentes & Campoy, 2008; Q. Li, Liu, Huang, & Huang, 2018; Mullane, Lawrence, Corkum, Klein, & McLaughlin, 2016; Wiegand, Petersen, Bundesen, & Habekost, 2017), and the detrimental effect of an alerting cue (whether visual or auditory) in conflict resolution (Asanowicz & Marzecová, 2017; Callejas et al., 2005; Callejas et al., 2004; Zani & Proverbio, 2017). The synergistic cooperation between the endogenous (i.e., voluntary) orienting and executive control function has also been shown in the form of a more precise selection of target information reducing the distracting effect of irrelevant information (Callejas et al., 2005; Callejas et al., 2004; Fan et al., 2009; Fuentes & Campoy, 2008; Spagna, Mackie, et al., 2015), while exogenous (i.e., automatic) orienting has been shown to impair conflict resolution (Trautwein, Singer, & Kanske, 2016). Executive control has been proposed to be located at the top of a hierarchical structure and acting irrespectively of sensory modalities (i.e., supramodal) (Donohue, Liotti, Perez, & Woldorff, 2012; Moris Fernandez, Macaluso, & Soto-Faraco, 2017; Roberts & Hall, 2008; Spagna et al., 2017; Spagna, Mackie, et al., 2015), with the alerting and orienting functions being located at a lower level and tied more to modality-specific mechanisms (Bushara et al., 1999; Langner et al., 2011; Salmi, Rinne, Degerman, Salonen, & Alho, 2007; Thiel & Fink, 2007; Ward, 1994; Yang & Mayer, 2013). Therefore, investigating the hemispherical asymmetries of the executive control under different states of alerting and orienting may provide useful insights on the dynamics from which attention emerges as a unitary cognitive function.

In this study we employed the visual field methodology (Bourne, 2006) to examine the efficiency and interactions of the attentional networks separately in the RH and LH. The hemispherical asymmetry of the executive control was investigated under different alerting and

Right hemisphere advantage for executive control

orienting conditions by using a lateralized version of the attentional network test (LANT-R). In this task, the hemispherical difference for executive control of attention was examined by presenting the target in the left or right visual field, under the conditions of congruent or incongruent flankers to generate the conflict effect, preceded by bilateral or unilateral visual cues to trigger the alerting and the orienting functions, respectively. We predicted a RH superiority in the executive control function, as shown by a reduced conflict effect, and that such superiority should be facilitated by the activation of alerting and orienting functions.

Method and Materials

Participants


Fifty students taking the Introduction to Psychology course at Queens College, the City University of New York (CUNY) participated in this study. Data from two participants were not included in the analyses due to low accuracy (approaching chance level) and an overall response time (RT) greater than two standard deviations (SD) from the group mean. The remaining participants ($n = 48$) consisted of 40 females and 8 males, with an average age of 20.8 years old ($SD = 3.22$), ranging from 18 to 25. All but one participant were right-handed, and all participants had normal or corrected-to-normal vision. Written informed consent approved by the Institutional Review Board of Queens College, CUNY was obtained from all participants prior to participation.

The Lateralized Attention Network Test - Revised (LANT-R)

The LANT-R is a modified version of the revised attention network test (Fan et al., 2009) to measure the hemispherical differences in the efficiency of the attentional functions (alerting, orienting, executive control) and their interactions. **Figure 1** illustrates the sequence of events of the LANT-R. On the screen with a gray background, there were two vertically aligned

Right hemisphere advantage for executive control

rectangular boxes with a black outline located to the left and right of a central fixation cross. For each trial, five arrows appeared in either the left or right box, with the center arrow (the target) pointing either up or down, and the other arrows located above or below the target pointing either toward the same direction (flanker congruent condition) or toward the opposite direction (flanker incongruent condition). Participants were required to indicate the direction of the target by pressing the corresponding button on the mouse. The target was cued under one of three cueing conditions: double cue (i.e., the outline of both boxes changing from black to white), spatial cue (the outline of one of the boxes changing from black to white), or no cue (no change in the outline of any of the boxes). The double cue was used as an alerting stimulus by providing temporal information about the impending target, regardless of location. The spatial cue was designed to validly or invalidly orient the participant's attention to either the left or the right side, thus providing both temporal and spatial information about the impending target.



Right hemisphere advantage for executive control

In each trial, participants were required to indicate the direction of a central arrow surrounded both above and below by two flanker arrows, either pointing in the same (congruent) or in the opposite (incongruent) direction as the target.

The size of the central fixation cross, located in the middle of the screen and visible throughout the entire task, was 1° of visual angle. The five black arrows that appeared on either side were $.56^\circ$ each. The arrows were separated by $.06^\circ$ of space, and the stimuli (target arrow and the flankers) subtended a total of 3.27° . Each box was located 5° from the central fixation cross. Preceding the target by three cue-to-target intervals (0 ms, 400 ms, 800 ms) equally likely to be presented, the change in the outline of the boxes from black to white lasted 100 ms, then the target and flankers were presented for 500 ms. Following the arrow presentation, there was a variable inter-trial interval ranging from 2000 to 12000 ms, with a mean interval of 4000 ms. Each trial had a mean duration of 5000 ms. The test consisted of 4 blocks, with each block containing 72 trials. On half of the trials (i.e., 144), the participant was shown a valid spatial cue, while the remaining trials were equally divided in 48 trials of a double cue condition, 48 trials of an invalid spatial cue condition, and 48 trials of a no cue condition. There was equal number of trials in the congruent and incongruent condition. In total, each block took 420 seconds to be completed, and the entire experiment took around 30 minutes to be completed. Prior to the beginning of the experimental session, the participant completed a short practice session consisting of 32 trials. During the practice session, participants were provided with feedback on accuracy and response time on each trial.

The practice session and the experimental session provided instructions in written form on the computer screen. It was emphasized that the participants fixate on the central fixation cross for the duration of each trial and press the mouse button that corresponded to the direction

Right hemisphere advantage for executive control

of the target arrow as quickly and accurately as possible. The mouse was tilted 90 degrees to the left, so that the right mouse button indicated “up”, and the left mouse button indicated “down”. The mouse was also aligned to the middle of the computer screen and participant’s midline of the body. To diminish the effects of hemispheric bias through motor control, the participant used both hands on the mouse to respond (Gable, Poole, & Cook, 2013): the right index finger was used to press the up-button while the left index finger was used to press the down-button. The task was displayed on a 17-inch LCD monitor, and the LANT-R was programmed in E-Prime (Psychology Software Tools, Pittsburgh, PA). Each participant completed the testing on a computer located in a silent and well-illuminated room.

Operational definition of the attentional effects and interactions

A $4 \times 2 \times 2$ factorial design was used in this experiment, with *cue condition* (no cue, double cue, valid cue, invalid cue), *conflict condition* (congruent, incongruent), and *visual field* (left, right) as within-subject factors. The same operational definitions for the attentional functions presented originally in (Fan et al., 2009) were used to estimate the attentional effects and are shown in **Table 1**. The *alerting effect* represents the performance benefit produced by the increased arousal compared to the no cue condition. The *orienting effect* is equivalent to *moving + engaging* operation defined in (Fan et al., 2009) and represents the performance benefit produced by a valid spatial information compared to the temporal information provided by an alerting cue. The *disengaging effect* represents the cost of disengaging from an invalid spatial cue. Combining the *disengaging* operation and the *moving + engaging* operation leads to the *validity effect*, which represents the extent to which a valid spatial cue condition benefits the participant’s performance compared to the cost in performance due to the invalid spatial cue condition. The *conflict effect* represents the cost in solving the conflict caused by the incongruent

Right hemisphere advantage for executive control

flankers, and a larger conflict effect corresponds to a less efficient executive control function (Fan et al., 2002). The interaction effects between these attentional functions were calculated by comparing the conflict effects under different cue conditions: *alerting by conflict interaction effect*, with a negative value indicating a negative impact of alerting on conflict processing. *Orienting by conflict interaction effect*, with a positive value indicating a more efficient conflict processing because of orienting. *Disengaging by conflict interaction effect*, with a positive value indicating a less efficient conflict processing because of invalid orienting compared to an alerting cue. *Validity by conflict interaction effect*, with a positive value indicating a less efficient conflict processing because of invalid orienting compared to a valid orienting cue. All the above-defined effects and interactions were computed separately for the RH and LH, according to the location where target and flankers were presented.

Data Analyses

Mean RT and error rate were calculated for each condition, with error trials (incorrect or missing responses) (less than 2% per subject) and RT outliers (above two standard deviations) (less than 2.23% per subject) being excluded from analyses on RT. Hemispherical differences in attentional effects, overall RT, and error rate, were tested using paired-sample *t* tests (one-tailed, hypothesizing a RH advantage). Effect sizes are reported as Cohen's *d*. Further, a $4 \times 2 \times 2$ ANOVA was conducted to test for the presence of *omnibus* effects and interactions among cue conditions, congruent conditions, and the visual fields. Planned comparisons were then conducted to analyze significant interactions between the conflict effect and the alerting, orienting, disengaging, and validity effects were tested using $2 \times 2 \times 2$ ANOVAs, with an alpha-level set to .05. Specifically, a 2 (*Hemisphere*: RH, LH) \times 2 (*Alerting*: no cue, double cue) \times 2 (*Conflict*: congruent, incongruent) was conducted to identify differences in the alerting \times conflict

Right hemisphere advantage for executive control

interaction in the two hemispheres. A 2 (*Hemisphere*: RH, LH) \times 2 (*Orienting*: double cue, valid cue) \times 2 (*Conflict*: congruent, incongruent) was conducted to examine the orienting \times conflict interaction separately in each hemisphere. A 2 (*Hemisphere*: RH, LH) \times 2 (*Disengaging*: invalid cue, double cue) \times 2 (*Conflict*: congruent, incongruent) was conducted to examine the disengaging \times conflict interaction separately in each hemisphere. A 2 (*Hemisphere*: RH, LH) \times 2 (*Validity*: invalid, valid cue) \times 2 (*Conflict*: congruent, incongruent) was conducted to examine the validity \times conflict interaction separately in each hemisphere. Effects sizes are reported as partial eta squared (η^2).

Results

Table 2 shows the RT and error rate for all conditions, separately for each hemisphere. The overall mean RT was 598 ms (SD = 95 ms), and the overall mean error rate was 9.12% (SD = 9.90). Participants were significantly faster at responding to stimuli presented to the RH (598 ± 93 ms) compared to the LH (602 ± 96 ms, $t_{(47)} = -2.71$, $p < .05$, $d = 0.04$), while this difference was not significant for the error rate (RH: $9.02 \pm 10.00\%$, LH: $9.22 \pm 9.82\%$, $t_{(47)} = -0.42$, $p = .97$, $d = 0.02$).

Hemispherical differences in the attentional effects

Figure 2 shows the alerting, orienting, disengaging, validity, and conflict effects separately for each hemisphere. For the alerting effect, the hemispherical difference was not significant in RT (RH: 44 ± 37 ms, LH: 33 ± 36 ms, $t_{(47)} = 1.42$, $p = .08$, $d = .29$), as well as in error rate (RH: $2.52 \pm 7.23\%$, LH: $1.30 \pm 6.50\%$, $t_{(47)} = 1.19$, $p = .12$, $d = .24$). For the orienting effect, the hemispherical difference was significant in RT ($t_{(47)} = 2.66$, $p < .01$, $d = .54$), indicating that the orienting effect was smaller when stimuli were presented to the RH (34 ± 28 ms) compared to the LH (53 ± 33 ms). This effect was also significant in error rate ($t_{(47)} = 2.92$,

Right hemisphere advantage for executive control

$p < .001$, $d = .59$), with a smaller effect when stimuli were presented to the RH ($0.14 \pm 5.89\%$) compared to the LH ($3.18 \pm 5.73\%$). For the disengaging, the hemispherical difference was significant in RT ($t(47) = 2.66$, $p < .01$, $d = .54$), indicating the disengaging effect was greater when stimuli were presented to the RH (62 ± 40 ms) compared to the LH (44 ± 34 ms). This effect was also significant in error rate ($t(47) = 2.93$, $p < .001$, $d = .60$), with a greater effect when stimuli were presented to the RH ($7.03 \pm 8.02\%$) compared to the LH ($2.34 \pm 7.03\%$). For the validity effect, the hemispherical difference was not significant in RT (RH: 96 ± 36 ms, LH: 99 ± 38 ms, $t(47) = 0.07$, $p = .94$, $d = .01$) as well as in error rate (RH: $7.17 \pm 7.81\%$, LH: $5.52 \pm 6.71\%$, $t(47) = 1.27$, $p = .21$). For the conflict effect (see also **Figure 3**), the hemispherical difference was significant in RT ($t(47) = .15$, $p < .01$, $d = .03$), showing a smaller conflict effect in the RH (106 ± 58 ms) compared to the LH (127 ± 56 ms), while this effect was not significant in error rate (RH: $12.17 \pm 10.17\%$, LH: $12.01 \pm 11.48\%$, $t(47) = 0.14$, $p = .89$, $d = .03$).

Right hemisphere advantage for executive control





Figure 2. Attentional effects and interactions in terms of reaction time (top panel), and error rate (lower panel). Error bars represent the standard error of the mean. ** = $p < .01$; *** = $p < .001$.



Right hemisphere advantage for executive control

Figure 3. Average reaction time (left panel) and error rate (right panel) for the congruent and incongruent conditions. Error bars represent the standard error of the mean. *** = $p < .001$.

Hemispherical differences in terms of main effects and interactions

For the analysis conducted in RT, (the top panel of **Figure 4**), the main effect of the factor *Cue* was significant, ($F(1, 47) = 233.61, p < .001, \eta^2 = .83$), indicating that participants were faster in the valid cue condition (548 ± 94 ms) compared to the double cue (592 ± 111 ms, $p < .001$), nocue (631 ± 114 ms, $p < .001$), and invalid cue (645 ± 121 ms, $p < .001$). Further, participants were faster in the double cue condition compared to the no cue condition ($p < .001$) and invalid cue condition ($p < .001$), and faster in the nocue compared to the invalid cue condition ($p < .05$). The main effect of the factor *Conflict* was significant ($F(1, 47) = 218.89, p < .001, \eta^2 = .82$), indicating that participants were faster in the congruent condition (541 ± 77 ms) compared to the incongruent condition (667 ± 116 ms, $p < .001$). The main effect of the factor *Hemisphere* was significant ($F(1, 47) = 7.32, p < .01, \eta^2 = .14$), indicating that participants were faster in response to stimuli presented to the RH (599 ± 114 ms) compared to stimuli presented to the LH (609 ± 119 ms, $p < .01$). The interaction *Cue by Conflict* was significant, ($F(3, 141) = 15.10, p < .001, \eta^2 = .24$), the interaction *Cue by Hemisphere* was significant ($F(3, 141) = 3.64, p < .05, \eta^2 = .07$), the interaction *Conflict by Hemisphere* was significant ($F(1, 47) = 13.14, p < .001, \eta^2 = .22$), and the three way interaction *Cue by Conflict by Hemisphere* was also significant ($F(3, 141) = 4.26, p < .01, \eta^2 = .08$). In line with the hypothesis driven nature of the current manuscript, planned comparisons were conducted to further analyze these interactions (see section below).

For the analysis conducted in Error rate, (the lower panel of **Figure 4**), the main effect of the factor *Cue* was significant, ($F(1, 47) = 24.95, p < .001, \eta^2 = .35$), indicating that participants

Right hemisphere advantage for executive control

committed fewer errors in the valid cue condition ($5.89 \pm 8.37\%$) compared to the nocue ($9.46 \pm 12.10\%$, $p < .001$), and invalid cue ($12.24 \pm 16.04\%$, $p < .001$), while the difference was not significant compared to the double cue ($7.55 \pm 11.67\%$, $p = .09$). Further, participants made fewer errors in the double cue condition compared to the invalid cue condition ($p < .001$) while this difference was not significant compared to the no cue condition ($p = .17$). The difference between nocue and invalid cue trials was significant ($p < .05$). The main effect of the factor *Conflict* was significant ($F(1, 47) = 68.58$, $p < .001$, $\eta^2 = .59$), indicating that participants made fewer errors in the congruent condition ($2.74 \pm 4.98\%$) compared to the incongruent condition ($14.83 \pm 14.74\%$, $p < .001$). The main effect of the factor *Hemisphere* was not significant ($F < 1$). The interaction *Cue by Conflict* ($F(3, 141) = 21.09$, $p < .001$, $\eta^2 = .31$) and the *Cue by Hemisphere* interactions ($F(3, 141) = 5.05$, $p < .01$, $\eta^2 = .10$) were significant, while the *Conflict by Hemisphere* ($F < 1$) and the three way interaction *Cue by Conflict by Hemisphere* ($F < 1$) were not. In line with the hypothesis driven nature of the current manuscript, planned comparisons were conducted to further analyze these interactions (see section below).

Right hemisphere advantage for executive control




Figure 4. Average reaction time (top panel) and error rate (lower panel) for the four cue conditions: a) no cue; b) double cue; c) valid cue; d) invalid cue, separately for the incongruent and congruent conditions and for stimuli presented in the RH or LH. Error bars represent the standard error of the mean.

Right hemisphere advantage for executive control

Hemispherical differences in terms of the interactions between attentional functions

For brevity, here we report only results of the three-way interactions between the conflict effect and the hemispheres under different cue conditions, which are the focus of the current paper. A full depiction of the results regarding the 2 (*Hemisphere*) \times 2 (*Cue conditions*) \times 2 (*Conflict*) ANOVAs can be found in **Figure 5**, and a full description can be found in **Table S1**.

The Alerting by Conflict interaction. For the 2 (*Hemisphere*: RH, LH) \times 2 (*Alerting*: no cue, double cue) \times 2 ANOVA (*Conflict*: congruent, incongruent) conducted on the RT (the top panel of **Figure 5a**), the three-way interaction was significant ($F(1, 47) = 4.03, p = .05, \eta^2 = .08$). The follow-up analysis conducted on the *Hemisphere* by *Conflict* interaction separately for the no cue and double cue conditions showed that for the no cue condition, showed that the two-way interaction was not significant ($F < 1$), while for the double cue condition the two-way interaction was significant ($F(1, 47) = 18.49, p < .001, \eta^2 = .28$), showing that participants were significantly faster at the incongruent condition when stimuli were presented to the RH (634 ± 111 ms) compared to the LH (675 ± 107 ms, $p < .001$). For the ANOVA conducted on error rate (the bottom panel of **Figure 5a**), the three-way interaction was not significant ($p \geq .24$).

The Orienting by Conflict interaction. For the 2 (*Hemisphere*: RH, LH) \times 2 (*Alerting*: double cue, valid cue) \times 2 ANOVA (*Conflict*: congruent, incongruent) conducted on the RT (the top panel of **Figure 5b**), the three-way interaction was not significant ($F(1, 47) = 1.33, p = .25, \eta^2 = .03$). For the ANOVA on error rate (the bottom panel of **Figure 5b**), the three-way interaction was also not significant ($p \geq .38$).

The Disengaging by Conflict interaction. For the 2 (*Hemisphere*: RH, LH) \times 2 (*Alerting*: double cue, invalid cue) \times 2 ANOVA (*Conflict*: congruent, incongruent) conducted on the RT (the top panel of **Figure 5c**), the three-way interaction was significant ($F(1, 47) = 15.38, p <$

Right hemisphere advantage for executive control

.001, $\eta^2 = .25$). The follow-up analysis conducted on the *Hemisphere* by *Conflict* interaction separately for the invalid cue and double cue conditions showed that for the invalid cue condition, the two-way interaction was not significant ($F < 1$), while it was significant for the double cue condition, ($F(1, 47) = 18.49, p < .001, \eta^2 = .28$), showing that participants were significantly faster at the incongruent condition when stimuli were presented to the RH (634 ± 111 ms) compared to the LH (675 ± 107 ms, $p < .001$), while this difference was not significant for the congruent condition (RH: 529 ± 70 ms, LH: 531 ± 68 ms, $p = .74$). For the ANOVA conducted on error rate (the bottom panel of **Figure 5c**), the three-way interaction was not significant ($p \geq .36$).

The Validity by Conflict interaction. For the 2 (*Hemisphere*: RH, LH) \times 2 (*Alerting*: double cue, invalid cue) \times 2 ANOVA (*Conflict*: congruent, incongruent) conducted on the RT (the top panel of **Figure 5d**), the three-way interaction was significant ($F(1, 47) = 7.27, p < .01, \eta^2 = .13$). The follow-up analysis conducted on the *Hemisphere* by *Conflict* interaction separately for the valid cue and invalid cue conditions showed that for the valid cue condition, the two-way interaction was significant ($F(1, 47) = 25.23, p < .001, \eta^2 = .35$), indicating that for the congruent condition participants were slower in the RH (503 ± 66 ms) compared to the LH (493 ± 61 ms), while in the incongruent condition participants were faster in the RH (590 ± 90 ms) compared to the LH (606 ± 95 ms). For the invalid cue condition, the two-way interaction was not significant ($F < 1$). For the ANOVA conducted on error rate (the bottom panel of **Figure 5c**), the three-way interaction was not significant ($p \geq .36$).

Right hemisphere advantage for executive control




Figure 5. Average reaction time (top panel) and error rate (lower panel) for the interactions between the task conditions: a) alerting by conflict; b) orienting by conflict; c) disengaging by conflict; d) validity by conflict. Error bars represent the standard error of the mean.

Discussion

In the present study, we used a lateralized version of the ANT-R and the visual field methodology to examine the efficiency and interactions of the attentional networks separately in the RH and LH, with a specific focus to the hemispherical asymmetry of the executive control under different alerting and orienting conditions. In summary, significant behavioral differences were found for the orienting, disengaging, and executive control of attention, and interactions between either the alerting and the orienting function with the executive control of attention, showing an advantage in all these functions for the RH. These result supports the long-standing proposal of a RH advantage for the orienting of attention and extend it to also include

Right hemisphere advantage for executive control

hemispherical asymmetries for executive control of attention. Specifically, the RH advantage in conflict resolution occurs under conditions of increased alertness or valid spatial orienting, suggesting a synergistic dynamic among the three functions. Nonetheless, our results showed no evidence for a hemispherical difference in the spatial cue conditions (i.e., no difference between left or right cues for either valid or invalid conditions). Therefore, we have no evidence for a “disadvantage” of the LH compared to the RH in the orienting of attention (a result that would have favored the hemispatial theory (Heilman & Van Den Abell, 1980), and while it is plausible that, in line with the interhemispheric competition account (Kinsbourne, 1987), each hemisphere has its own contralateral vector of attention, although the lack of lesion data makes our conclusion speculative.

Consistent with our (Fan et al., 2009; Mackie et al., 2013; Spagna, Mackie, et al., 2015) and other studies (Asanowicz & Marzecová, 2017; Callejas et al., 2005; Callejas et al., 2004; Chica et al., 2012; Marotta et al., 2012; Roca et al., 2012), a complex pattern of interactions was shown among the three attentional functions, further proving that the interplay among them is necessary for the selection and prioritization of the processing of goal-relevant information (Badre, 2011). For instance, we found that indicating the location where the target would be presented by means of a valid spatial cue reduced the conflict effect by decreasing the response time to a target flanked by incongruent information. This is consistent with previous evidence showing the beneficial effect of the endogenous orienting function on the executive control of attention (e.g., Callejas et al., 2005; Fan et al., 2009; Spagna, Mackie, et al., 2015; Xuan et al., 2016), and our results further extend this knowledge to the exogenous orienting function, by showing that the conflict resolution benefited more from a valid cue compared to an invalid cue when the process was conducted by the RH. Conversely, a wealth of studies showed that the

Right hemisphere advantage for executive control

presentation of an alerting cue has a cost on the conflict resolution (Asanowicz & Marzecová, 2017; Callejas et al., 2005; Callejas et al., 2004; Xuan et al., 2016), a pattern theoretically linked to a “U-shape” relationship (and justified by the norepinephrine dose-response relationship underlying the alerting function (Aston-Jones & Cohen, 2005)), with a phasic increase of the arousal state altering the efficiency of the executive control of attention (Thiele & Bellgrove, 2018). In this study, however, we show a pattern of facilitation produced by an alerting cue on the conflict resolution when stimuli were processed by the RH compared to the LH, indicating the superiority of this hemisphere in conflict resolution is benefited by the activation of the alerting function. At a closer look, our results and the evidence from the previous studies can coexist due to the following reasons: 1) we also show a greater conflict effect in the double cue condition compared to the no cue condition, when averaging performance across the two hemispheres; 2) however, if further broken down into the specific task conditions, the above mentioned effect reveals that participants showed overall better performance at the double cue compared to the no cue effect; and 3) the interaction between these two components of attention within each visual field shows that a double cue condition was significantly less beneficial to the conflict processing in the LH. While the most obvious explanation for this interaction effect is to state a RH advantage for both alerting, executive control, and their interaction, it may also be that the worsening of the conflict resolution in the LH under the alerting condition results from more limited attentional resources of this hemisphere that more quickly are exhausted when the two functions are contextually engaged.

Consensus exists regarding the neural substrate supporting the activation of interactive attentional networks (Xuan et al., 2016), showed as a complex pattern of interaction among cortical and subcortical areas (Petersen & Posner, 2012). Specifically, a bilateral fronto-parietal

Right hemisphere advantage for executive control

network (composed of the frontal eye fields, areas near and along the intraparietal sulcus, and the dorsolateral prefrontal cortex), a cingulo-opercular network (composed of the anterior cingulate cortex, and the anterior insular cortex), and subcortical areas such as the thalamus, superior colliculi, and the basal ganglia, were found to consistently interact during attentional tasks (Callejas, Shulman, & Corbetta, 2014; Corbetta et al., 2008; Corbetta & Shulman, 2002; Fan et al., 2005; Krauzlis, Lovejoy, & Zénon, 2013; Parlatini et al., 2017; Patel et al., 2015; Posner, 2012; Posner & Fan, 2008; Shulman et al., 2009; Wang et al., 2010; Xuan et al., 2016).

However, bilateral involvement and functional asymmetries are not mutually exclusive properties in our brain (Arbula et al., 2017; Bartolomeo & Thiebaut de Schotten, 2016; Corballis, 2009, 2017; A. Vallesi, Arbula, Capizzi, Causin, & D'Avella, 2015), and the existence of hemispherical differences for executive control of attention in terms of activation magnitude seems plausible. This is particularly true if we consider that the function of executive control of attention is to allow the selection and prioritization of the processing of goal-relevant information to reach consciousness (Fan, Fossella, Sommer, Wu, & Posner, 2003; Mackie & Fan, 2017; Mackie et al., 2013; Petersen & Posner, 2012; Spagna, Mackie, et al., 2015) and that this function inevitably interacts with the hemispherical specializations related to the specific type of information to be handled (Marotta & Casagrande, 2017; Spagna et al., 2016; Spagna et al., 2014). More specifically, hemispherical asymmetries were first identified in “split brain” patients (i.e., corpus callosotomy intervention as part of medical treatment) (Sperry, 1968), one of the earliest evidence for hemispheric dominance for verbal and language functions in the LH and for non-verbal and spatial functions in the RH (see Bartolomeo & Thiebaut de Schotten, 2016 for a review). In the present study, we used non-verbal information as imperative stimuli,

Right hemisphere advantage for executive control

an aspect that might have magnified the subtle interactions among the executive control of attention and the hemispheric dominance of the RH for this type of stimuli.

Alternatively, the RH advantage in conflict resolution may be explained by brain mechanisms revealed in neuroimaging studies. Neuroimaging evidence for a RH dominance associated with the executive control of attention exist at the functional level (Garavan et al., 1999), as shown by the association between participants' monitoring of task contingencies and the activation of prefrontal cortex in the RH (see Antonino Vallesi, 2012 for a review; Antonino Vallesi & Crescentini, 2011), response inhibition and the association with the right inferior frontal cortex (Aron, Robbins, & Poldrack, 2014; Cai, Ryali, Chen, Li, & Menon, 2014; Levy & Wagner, 2011), increased metabolic activity in the medial prefrontal cortex associated with monitor the auditory environment (Cohen, Semple, Gross, King, & Nordahl, 1992), and the anterior cingulate cortex of the RH for response selection (Turken & Swick, 1999), motor control (Paus, 2001) and attention shifting (Kondo, Osaka, & Osaka, 2004). Further, regions within the frontal and parietal lobes of the right hemisphere have been consistently shown to play a critical role in the more broad construct of attention, by modulating activity in the sensory cortices (Bressler, Tang, Sylvester, Shulman, & Corbetta, 2008; Ruff et al., 2009), and by dynamically adjusting the focus of attention (Malhotra, Coulthard, & Husain, 2009; Marshall, O'Shea, Jensen, & Bergmann, 2015; O'Shea, Muggleton, Cowey, & Walsh, 2004; Ronconi, Basso, Gori, & Facoetti, 2014). Consistently, in one of our previous studies using the ANT together with fMRI (Fan et al., 2005), in which the presentation of the stimuli was not lateralized, we showed a significant activation of the right temporo-parietal junction for the alerting function, together with clusters in the right parietal lobe for the orienting function, and a right-centered cluster of anterior cingulate cortex activation for executive control of attention. Together with the evidence

Right hemisphere advantage for executive control

regarding the necessity of the fronto-parietal and cingulo-opercular networks of the RH for both the alerting (J. Li et al., 2016; Q. Li et al., 2018; Perin, Godefroy, Fall, & de Marco, 2010; Posner, 2008) and orienting (Bartolomeo, 2014; Bartolomeo, Thiebaut de Schotten, & Chica, 2012; Chica et al., 2018; Chica et al., 2012; Mesulam, 1999), this neuroimaging evidence is in support of the findings from the present study regarding the executive control of attention.

One limitation of the present study must be, however, kept in mind. Here, we did not employ eye tracking techniques and therefore have no direct measure of the eye position of participants throughout the task. This is point may limit the conclusion of our studies, because eye movements must be avoided at all costs for the lateralized presentation to work. However, evidence exists that suggests carefully instructing participants before the beginning of the task regarding the relevance of maintaining central fixation can be effective in controlling eye movements (e.g., Moscovitch, 1986; Śmigasiewicz et al., 2014; Smigasiewicz et al., 2010; Verleger, Smigasiewicz, & Moller, 2011; Verleger et al., 2009). Additionally, a variety of behavioral studies investigating the hemispherical differences in the attentional functions using behavioral tasks with lateralized stimuli found interactions between the attentional functions and the visual field with and without using eye-tracker. It is worth noting that the lack of a control of eye movement may have a negative impact so that the hemispheric difference would not be significant. Here, we are not making conclusions about the negative findings (absence of hemispherical advantage) but on the positive findings (presence of hemispherical advantage of the RH). In other words, the significant difference between hemisphere should not be due to the presence of eye movement, but due to the true effect in terms of hemispherical difference. Overall, although the lack of control of participants' central fixation limits the validity of our

Right hemisphere advantage for executive control

conclusion regarding the orienting effect, it is not as relevant for our result concerning the executive control of attention, which was the focus of the study.

Our study is consistent with existing behavioral and neural evidence indicating a RH advantage for attention, and further provides novel knowledge regarding how this advantage also extends to the executive control of attention and its dynamics with the alerting and orienting function. These results may shed light on some of the inconsistencies shown in previous studies that did not find RH dominance for executive control and for the interplay between the three functions and pave the way for future studies by combining our task with neuroimaging techniques, aimed at characterizing the neural dynamics underlying the RH advantage for executive control of attention.

Right hemisphere advantage for executive control

Acknowledgements

This study was supported by the National Institute of Mental Health (NIMH) of the National Institutes of Health (NIH) under Award Number R01 MH094305.

Reference List

- Arbula, S., Pacella, V., De Pellegrin, S., Rossetto, M., Denaro, L., D'Avella, D., . . . Vallesi, A. (2017). Addressing the selective role of distinct prefrontal areas in response suppression: A study with brain tumor patients. *Neuropsychologia*, *100*, 120-130. doi: 10.1016/j.neuropsychologia.2017.04.018
- Aron, A. R., Robbins, T. W., & Poldrack, R. A. (2014). Inhibition and the right inferior frontal cortex: one decade on. *Trends Cogn Sci*, *18*(4), 177-185. doi: 10.1016/j.tics.2013.12.003
- Asanowicz, D., & Marzecová, A. (2017). Differential effects of phasic and tonic alerting on the efficiency of executive attention. *Acta psychologica*, *176*, 58-70.
- Asanowicz, D., Marzecova, A., Jaskowski, P., & Wolski, P. (2012). Hemispheric asymmetry in the efficiency of attentional networks. *Brain Cogn*, *79*(2), 117-128. doi: 10.1016/j.bandc.2012.02.014
- Aston-Jones, G., & Cohen, J. D. (2005). An integrative theory of locus coeruleus-norepinephrine function: adaptive gain and optimal performance. *Annu. Rev. Neurosci.*, *28*, 403-450. doi: 10.1146/annurev.neuro.28.061604.135709
- Badre, D. (2011). Defining an ontology of cognitive control requires attention to component interactions. *Top Cogn Sci*, *3*(2), 217-221. doi: 10.1111/j.1756-8765.2011.01141.x
- Bartolomeo, P. (2007). Visual neglect. *Curr Opin Neurol*, *20*(4), 381-386. doi: 10.1097/WCO.0b013e32816aa3a3
- Bartolomeo, P. (2014). The attention systems of the human brain *Attention Disorders After Right Brain Damage* (pp. 1-19): Springer London.
- Bartolomeo, P., & Chokron, S. (2002). Orienting of attention in left unilateral neglect. *Neuroscience & Biobehavioral Reviews*, *26*(2), 217-234.
- Bartolomeo, P., & Thiebaut de Schotten, M. (2016). Let thy left brain know what thy right brain doeth: Inter-hemispheric compensation of functional deficits after brain damage. *Neuropsychologia*, *93*(Pt B), 407-412. doi: 10.1016/j.neuropsychologia.2016.06.016
- Bartolomeo, P., Thiebaut de Schotten, M., & Chica, A. B. (2012). Brain networks of visuospatial attention and their disruption in visual neglect. *Front Hum Neurosci*, *6*, 110. doi: 10.3389/fnhum.2012.00110
- Belanger, H. G., & Cimino, C. R. (2002). The lateralized stroop: a meta-analysis and its implications for models of semantic processing. *Brain Lang*, *83*(3), 384-402.
- Benwell, C. S., Thut, G., Grant, A., & Harvey, M. (2014). A rightward shift in the visuospatial attention vector with healthy aging. *Frontiers in aging neuroscience*, *6*.
- Bourne, V. J. (2006). The divided visual field paradigm: Methodological considerations. *Laterality*, *11*(4), 373-393.
- Bressler, S. L., Tang, W., Sylvester, C. M., Shulman, G. L., & Corbetta, M. (2008). Top-down control of human visual cortex by frontal and parietal cortex in anticipatory visual spatial attention. *J Neurosci*, *28*(40), 10056-10061. doi: 10.1523/JNEUROSCI.1776-08.2008
- Bushara, K. O., Weeks, R. A., Ishii, K., Catalan, M.-J., Tian, B., Rauschecker, J. P., & Hallett, M. (1999). Modality-specific frontal and parietal areas for auditory and visual spatial localization in humans. *Nature neuroscience*, *2*(8), 759-766.
- Cai, W., Ryali, S., Chen, T., Li, C. S., & Menon, V. (2014). Dissociable roles of right inferior frontal cortex and anterior insula in inhibitory control: evidence from intrinsic and task-related functional parcellation, connectivity, and response profile analyses across multiple datasets. *J Neurosci*, *34*(44), 14652-14667. doi: 10.1523/jneurosci.3048-14.2014

Right hemisphere advantage for executive control

- Callejas, A., Lupianez, J., Funes, M. J., & Tudela, P. (2005). Modulations among the alerting, orienting and executive control networks. *Exp Brain Res*, 167(1), 27-37. doi: 10.1007/s00221-005-2365-z
- Callejas, A., Lupianez, J., & Tudela, P. (2004). The three attentional networks: on their independence and interactions. *Brain Cogn*, 54(3), 225-227. doi: 10.1016/j.bandc.2004.02.012
- Callejas, A., Shulman, G. L., & Corbetta, M. (2014). Dorsal and ventral attention systems underlie social and symbolic cueing. *Journal of cognitive neuroscience*, 26(1), 63-80.
- Chica, A. B., Thiebaut de Schotten, M., Bartolomeo, P., & Paz-Alonso, P. M. (2018). White matter microstructure of attentional networks predicts attention and consciousness functional interactions. *Brain Struct Funct*, 223(2), 653-668. doi: 10.1007/s00429-017-1511-2
- Chica, A. B., Thiebaut de Schotten, M., Toba, M., Malhotra, P., Lupianez, J., & Bartolomeo, P. (2012). Attention networks and their interactions after right-hemisphere damage. *Cortex*, 48(6), 654-663. doi: 10.1016/j.cortex.2011.01.009
- Cohen, R. M., Semple, W. E., Gross, M., King, A. C., & Nordahl, T. E. (1992). Metabolic brain pattern of sustained auditory discrimination. *Exp Brain Res*, 92(1), 165-172.
- Corballis, M. C. (2009). The evolution and genetics of cerebral asymmetry. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1519), 867-879.
- Corballis, M. C. (2017). The Evolution of Lateralized Brain Circuits. *Front Psychol*, 8, 1021. doi: 10.3389/fpsyg.2017.01021
- Corbetta, M., Patel, G., & Shulman, G. L. (2008). The reorienting system of the human brain: from environment to theory of mind. *Neuron*, 58(3), 306-324.
- Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in the brain. *Nat Rev Neurosci*, 3(3), 201-215. doi: 10.1038/nrn755
- Corbetta, M., & Shulman, G. L. (2011). Spatial neglect and attention networks. *Annual review of neuroscience*, 34, 569.
- De Schotten, M. T., Dell'Acqua, F., Forkel, S. J., Simmons, A., Vergani, F., Murphy, D. G., & Catani, M. (2011). A lateralized brain network for visuospatial attention. *Nature neuroscience*, 14(10), 1245-1246.
- Donohue, S. E., Liotti, M., Perez, R., 3rd, & Woldorff, M. G. (2012). Is conflict monitoring supramodal? Spatiotemporal dynamics of cognitive control processes in an auditory Stroop task. *Cogn Affect Behav Neurosci*, 12(1), 1-15. doi: 10.3758/s13415-011-0060-z
- Evert, D. L., McGlinchey-Berroth, R., Verfaellie, M., & Milberg, W. P. (2003). Hemispheric asymmetries for selective attention apparent only with increased task demands in healthy participants. *Brain and Cognition*, 53(1), 34-41.
- Fan, J., Fossella, J., Sommer, T., Wu, Y., & Posner, M. I. (2003). Mapping the genetic variation of executive attention onto brain activity. *Proc Natl Acad Sci U S A*, 100(12), 7406-7411. doi: 10.1073/pnas.0732088100
- Fan, J., Gu, X., Guise, K. G., Liu, X., Fossella, J., Wang, H., & Posner, M. I. (2009). Testing the behavioral interaction and integration of attentional networks. *Brain and Cognition*, 70(2), 209-220. doi: 10.1016/j.bandc.2009.02.002
- Fan, J., McCandliss, B. D., Fossella, J., Flombaum, J. I., & Posner, M. I. (2005). The activation of attentional networks. *Neuroimage*, 26(2), 471-479. doi: 10.1016/j.neuroimage.2005.02.004
- Fan, J., McCandliss, B. D., Sommer, T., Raz, A., & Posner, M. I. (2002). Testing the efficiency and independence of attentional networks. *J Cogn Neurosci*, 14(3), 340-347. doi: 10.1162/089892902317361886

Right hemisphere advantage for executive control

- Fernandez-Duque, D., & Posner, M. I. (1997). Relating the mechanisms of orienting and alerting. *Neuropsychologia*, *35*(4), 477-486.
- Fernandez-Duque, D., & Posner, M. I. (2001). Brain imaging of attentional networks in normal and pathological states. *Journal of Clinical and Experimental Neuropsychology*, *23*(1), 74-93.
- Foxe, J. J., McCourt, M. E., & Javitt, D. C. (2003). Right hemisphere control of visuospatial attention: line-bisection judgments evaluated with high-density electrical mapping and source analysis☆. *Neuroimage*, *19*(3), 710-726.
- Fuentes, L. J., & Campoy, G. (2008). The time course of alerting effect over orienting in the attention network test. *Exp Brain Res*, *185*(4), 667-672. doi: 10.1007/s00221-007-1193-8
- Gable, P. A., Poole, B. D., & Cook, M. S. (2013). Asymmetrical hemisphere activation enhances global-local processing. *Brain Cogn*, *83*(3), 337-341. doi: 10.1016/j.bandc.2013.09.012
- Garavan, H., Ross, T. J., & Stein, E. A. (1999). Right hemispheric dominance of inhibitory control: an event-related functional MRI study. *Proc Natl Acad Sci U S A*, *96*(14), 8301-8306.
- Greene, D. J., Barnea, A., Herzberg, K., Rassis, A., Neta, M., Raz, A., & Zaidel, E. (2008). Measuring attention in the hemispheres: the lateralized attention network test (LANT). *Brain Cogn*, *66*(1), 21-31. doi: 10.1016/j.bandc.2007.05.003
- Heilman, K. M., & Van Den Abell, T. (1980). Right hemisphere dominance for attention: the mechanism underlying hemispheric asymmetries of inattention (neglect). *Neurology*, *30*(3), 327-330.
- Jonides, J. (1979). Left and right visual field superiority for letter classification. *Q J Exp Psychol*, *31*(3), 423-439. doi: 10.1080/14640747908400737
- Kałamala, P., Drożdżowicz, A., Szewczyk, J., Marzecová, A., & Wodniecka, Z. (2018). Task strategy may contribute to performance differences between monolinguals and bilinguals in cognitive control tasks: ERP evidence. *Journal of Neurolinguistics*, *46*, 78-92.
- Kincade, J. M., Abrams, R. A., Astafiev, S. V., Shulman, G. L., & Corbetta, M. (2005). An event-related functional magnetic resonance imaging study of voluntary and stimulus-driven orienting of attention. *J Neurosci*, *25*(18), 4593-4604. doi: 10.1523/JNEUROSCI.0236-05.2005
- Kinsbourne, M. (1987). Mechanisms of unilateral neglect. *Advances in psychology*, *45*, 69-86.
- Kondo, H., Osaka, N., & Osaka, M. (2004). Cooperation of the anterior cingulate cortex and dorsolateral prefrontal cortex for attention shifting. *Neuroimage*, *23*(2), 670-679. doi: 10.1016/j.neuroimage.2004.06.014
- Konrad, K., Neufang, S., Thiel, C. M., Specht, K., Hanisch, C., Fan, J., . . . Fink, G. R. (2005). Development of attentional networks: an fMRI study with children and adults. *Neuroimage*, *28*(2), 429-439. doi: 10.1016/j.neuroimage.2005.06.065
- Krauzlis, R. J., Lovejoy, L. P., & Zénon, A. (2013). Superior colliculus and visual spatial attention. *Annual review of neuroscience*, *36*.
- Langner, R., Kellermann, T., Boers, F., Sturm, W., Willmes, K., & Eickhoff, S. B. (2011). Modality-specific perceptual expectations selectively modulate baseline activity in auditory, somatosensory, and visual cortices. *Cereb Cortex*, *21*(12), 2850-2862. doi: 10.1093/cercor/bhr083
- Levy, B. J., & Wagner, A. D. (2011). Cognitive control and right ventrolateral prefrontal cortex: reflexive reorienting, motor inhibition, and action updating. *Ann N Y Acad Sci*, *1224*, 40-62. doi: 10.1111/j.1749-6632.2011.05958.x
- Li, J., Chen, X., Ye, W., Jiang, W., Liu, H., & Zheng, J. (2016). Alteration of the alertness-related network in patients with right temporal lobe epilepsy: A resting state fMRI study. *Epilepsy Res*, *127*, 252-259. doi: 10.1016/j.eplepsyres.2016.09.013

Right hemisphere advantage for executive control

- Li, Q., Liu, P., Huang, S., & Huang, X. (2018). The effect of phasic alertness on temporal precision. *Atten Percept Psychophys*, *80*(1), 262-274. doi: 10.3758/s13414-017-1418-2
- Liu, Y., Bengson, J., Huang, H., Mangun, G. R., & Ding, M. (2016). Top-down modulation of neural activity in anticipatory visual attention: control mechanisms revealed by simultaneous EEG-fMRI. *Cerebral Cortex*, *26*(2), 517-529.
- Longo, M. R., Trippier, S., Vagnoni, E., & Lourenco, S. F. (2015). Right hemisphere control of visuospatial attention in near space. *Neuropsychologia*, *70*, 350-357.
- Lunven, M., & Bartolomeo, P. (2016). Attention and spatial cognition: Neural and anatomical substrates of visual neglect. *Annals of physical and rehabilitation medicine*.
- Macaluso, E. (2010). Orienting of spatial attention and the interplay between the senses. *Cortex*, *46*(3), 282-297. doi: 10.1016/j.cortex.2009.05.010
- Mackie, M.-A., & Fan, J. (2017). Functional Neuroimaging of Deficits in Cognitive Control. In E. Goldber (Ed.), *Executive Functions in Health and Disease* (Vol. 1, pp. 249-300): Academic Press.
- Mackie, M.-A., Van Dam, N. T., & Fan, J. (2013). Cognitive control and attentional functions. *Brain and Cognition*, *82*(3), 301-312. doi: 10.1016/j.bandc.2013.05.004
- Malhotra, P., Coulthard, E. J., & Husain, M. (2009). Role of right posterior parietal cortex in maintaining attention to spatial locations over time. *Brain*, *132*(Pt 3), 645-660. doi: 10.1093/brain/awn350
- Manly, T., Hawkins, K., Evans, J., Woldt, K., & Robertson, I. H. (2002). Rehabilitation of executive function: facilitation of effective goal management on complex tasks using periodic auditory alerts. *Neuropsychologia*, *40*(3), 271-281.
- Marotta, A., & Casagrande, M. (2017). Developmental differences in cognitive control of social information. *Infant and Child Development*, *26*(4), e2005.
- Marotta, A., Lupianez, J., & Casagrande, M. (2012). Investigating hemispheric lateralization of reflexive attention to gaze and arrow cues. *Brain Cogn*, *80*(3), 361-366. doi: 10.1016/j.bandc.2012.08.001
- Marshall, T. R., O'Shea, J., Jensen, O., & Bergmann, T. O. (2015). Frontal eye fields control attentional modulation of alpha and gamma oscillations in contralateral occipitoparietal cortex. *Journal of Neuroscience*, *35*(4), 1638-1647.
- Martella, D., Casagrande, M., & Lupianez, J. (2011). Alerting, orienting and executive control: the effects of sleep deprivation on attentional networks. *Exp Brain Res*, *210*(1), 81-89. doi: 10.1007/s00221-011-2605-3
- Marzecova, A., Asanowicz, D., KrivÁ, L. U., & Wodniecka, Z. (2012). The effects of bilingualism on efficiency and lateralization of attentional networks. *Bilingualism: Language and Cognition*, *16*(03), 608-623. doi: 10.1017/s1366728912000569
- Mecklinger, A. D., von Cramon, D. Y., Springer, A., & Matthes-von Cramon, G. (1999). Executive control functions in task switching: evidence from brain injured patients. *J Clin Exp Neuropsychol*, *21*(5), 606-619. doi: 10.1076/jcen.21.5.606.873
- Mesulam, M. M. (1981). A cortical network for directed attention and unilateral neglect. *Ann Neurol*, *10*(4), 309-325. doi: 10.1002/ana.410100402
- Mesulam, M. M. (1999). Spatial attention and neglect: parietal, frontal and cingulate contributions to the mental representation and attentional targeting of salient extrapersonal events. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, *354*(1387), 1325-1346.

Right hemisphere advantage for executive control

- Moris Fernandez, L., Macaluso, E., & Soto-Faraco, S. (2017). Audiovisual integration as conflict resolution: The conflict of the McGurk illusion. *Hum Brain Mapp*, *38*(11), 5691-5705. doi: 10.1002/hbm.23758
- Moscovitch, M. (1986). Afferent and efferent models of visual perceptual asymmetries: theoretical and empirical implications. *Neuropsychologia*, *24*(1), 91-114.
- Mullane, J. C., Lawrence, M. A., Corkum, P. V., Klein, R. M., & McLaughlin, E. N. (2016). The development of and interaction among alerting, orienting, and executive attention in children. *Child Neuropsychol*, *22*(2), 155-176. doi: 10.1080/09297049.2014.981252
- O'Shea, J., Muggleton, N. G., Cowey, A., & Walsh, V. (2004). Timing of target discrimination in human frontal eye fields. *J Cogn Neurosci*, *16*(6), 1060-1067. doi: 10.1162/0898929041502634
- Parlatini, V., Radua, J., Dell'Acqua, F., Leslie, A., Simmons, A., Murphy, D. G., . . . Thiebaut de Schotten, M. (2017). Functional segregation and integration within fronto-parietal networks. *Neuroimage*, *146*, 367-375. doi: 10.1016/j.neuroimage.2016.08.031
- Patel, G. H., Yang, D., Jamerson, E. C., Snyder, L. H., Corbetta, M., & Ferrera, V. P. (2015). Functional evolution of new and expanded attention networks in humans. *Proceedings of the National Academy of Sciences*, *112*(30), 9454-9459.
- Paus, T. (2001). Primate anterior cingulate cortex: where motor control, drive and cognition interface. *Nat Rev Neurosci*, *2*(6), 417-424. doi: 10.1038/35077500
- Perin, B., Godefroy, O., Fall, S., & de Marco, G. (2010). Alertness in young healthy subjects: an fMRI study of brain region interactivity enhanced by a warning signal. *Brain Cogn*, *72*(2), 271-281. doi: 10.1016/j.bandc.2009.09.010
- Petersen, S. E., & Posner, M. I. (2012). The attention system of the human brain: 20 years after. *Annu Rev Neurosci*, *35*, 73-89. doi: 10.1146/annurev-neuro-062111-150525
- Posner, M. I. (2008). Measuring alertness. *Ann N Y Acad Sci*, *1129*, 193-199. doi: 10.1196/annals.1417.011
- Posner, M. I. (2012). Imaging attention networks. *Neuroimage*, *61*(2), 450-456. doi: 10.1016/j.neuroimage.2011.12.040
- Posner, M. I., Cohen, Y., & Rafal, R. D. (1982). Neural systems control of spatial orienting. *Philos Trans R Soc Lond B Biol Sci*, *298*(1089), 187-198.
- Posner, M. I., & Fan, J. (2008). Attention as an organ system. *Topics in integrative neuroscience*, 31-61.
- Posner, M. I., Walker, J. A., Friedrich, F. J., & Rafal, R. D. (1984). Effects of parietal injury on covert orienting of attention. *The Journal of Neuroscience*, *4*(7), 1863-1874.
- Poynter, W., Ingram, P., & Minor, S. (2010). Visual field asymmetries in attention vary with self-reported attention deficits. *Brain Cogn*, *72*(3), 355-361. doi: 10.1016/j.bandc.2009.10.014
- Rafal, R. D. (1994). Neglect. *Curr Opin Neurobiol*, *4*(2), 231-236.
- Rastelli, F., Tallon-Baudry, C., Migliaccio, R., Toba, M. N., Ducorps, A., Pradat-Diehl, P., . . . Bartolomeo, P. (2013). Neural dynamics of neglected targets in patients with right hemisphere damage. *Cortex*, *49*(7), 1989-1996.
- Roberts, K. L., & Hall, D. A. (2008). Examining a supramodal network for conflict processing: a systematic review and novel functional magnetic resonance imaging data for related visual and auditory stroop tasks. *Journal of cognitive neuroscience*, *20*(6), 1063-1078.
- Robertson, I. H., Manly, T., Andrade, J., Baddeley, B. T., & Yiend, J. (1997). Oops!': performance correlates of everyday attentional failures in traumatic brain injured and normal subjects. *Neuropsychologia*, *35*(6), 747-758.

Right hemisphere advantage for executive control

- Roca, J., Fuentes, L. J., Marotta, A., Lopez-Ramon, M. F., Castro, C., Lupianez, J., & Martella, D. (2012). The effects of sleep deprivation on the attentional functions and vigilance. *Acta Psychol (Amst)*, *140*(2), 164-176. doi: 10.1016/j.actpsy.2012.03.007
- Ronconi, L., Basso, D., Gori, S., & Facoetti, A. (2014). TMS on right frontal eye fields induces an inflexible focus of attention. *Cerebral Cortex*, *24*(2), 396-402.
- Ruff, C. C., Blankenburg, F., Bjoertomt, O., Bestmann, S., Weiskopf, N., & Driver, J. (2009). Hemispheric differences in frontal and parietal influences on human occipital cortex: direct confirmation with concurrent TMS-fMRI. *J Cogn Neurosci*, *21*(6), 1146-1161. doi: 10.1162/jocn.2009.21097
- Salmi, J., Rinne, T., Degerman, A., Salonen, O., & Alho, K. (2007). Orienting and maintenance of spatial attention in audition and vision: multimodal and modality-specific brain activations. *Brain Struct Funct*, *212*(2), 181-194. doi: 10.1007/s00429-007-0152-2
- Shulman, G. L., Astafiev, S. V., Franke, D., Pope, D. L., Snyder, A. Z., McAvoy, M. P., & Corbetta, M. (2009). Interaction of stimulus-driven reorienting and expectation in ventral and dorsal frontoparietal and basal ganglia-cortical networks. *J Neurosci*, *29*(14), 4392-4407. doi: 10.1523/JNEUROSCI.5609-08.2009
- Shulman, G. L., Pope, D. L., Astafiev, S. V., McAvoy, M. P., Snyder, A. Z., & Corbetta, M. (2010). Right hemisphere dominance during spatial selective attention and target detection occurs outside the dorsal frontoparietal network. *J Neurosci*, *30*(10), 3640-3651. doi: 10.1523/JNEUROSCI.4085-09.2010
- Śmigasiewicz, K., Asanowicz, D., Westphal, N., & Verleger, R. (2014). Bias for the left visual field in rapid serial visual presentation: Effects of additional salient cues suggest a critical role of attention. *Journal of cognitive neuroscience*, *27*(2), 266-279.
- Smigasiewicz, K., Shalgi, S., Hsieh, S., Moller, F., Jaffe, S., Chang, C. C., & Verleger, R. (2010). Left visual-field advantage in the dual-stream RSVP task and reading-direction: a study in three nations. *Neuropsychologia*, *48*(10), 2852-2860. doi: 10.1016/j.neuropsychologia.2010.05.027
- Smigasiewicz, K., Westphal, N., & Verleger, R. (2017). Leftward bias in orienting to and disengaging attention from salient task-irrelevant events in rapid serial visual presentation. *Neuropsychologia*, *94*, 96-105. doi: 10.1016/j.neuropsychologia.2016.11.025
- Spagna, A., Dong, Y., Mackie, M.-A., Li, M., Harvey, P. D., Tian, Y., . . . Fan, J. (2015). Clozapine improves the orienting of attention in schizophrenia. *Schizophrenia Research*, *168*(1), 285-291. doi: 10.1016/j.schres.2015.08.009
- Spagna, A., He, G., Jin, S., Gao, L., Mackie, M. A., Tian, Y., . . . Fan, J. (2017). Deficit of supramodal executive control of attention in schizophrenia. *J Psychiatr Res*, *97*, 22-29. doi: 10.1016/j.jpsychires.2017.11.002
- Spagna, A., Mackie, M.-A., & Fan, J. (2015). Supramodal Executive Control of Attention. *Frontiers in Psychology*, *6*. doi: 10.3389/fpsyg.2015.00065
- Spagna, A., Martella, D., Fuentes, L., Marotta, A., & Casagrande, M. (2016). Hemispheric modulations of the attentional networks. *Brain and cognition*, *108*, 73.
- Spagna, A., Martella, D., Sebastiani, M., Maccari, L., Marotta, A., & Casagrande, M. (2014). Efficiency and interactions of alerting, orienting and executive networks: The impact of imperative stimulus type. *Acta Psychologica*, *148*, 209-215. doi: 10.1016/j.actpsy.2014.02.007
- Sperry, R. W. (1968). Hemisphere disconnection and unity in conscious awareness. *American Psychologist*, *23*(10), 723.

Right hemisphere advantage for executive control

- Spielberg, J. M., Miller, G. A., Engels, A. S., Herrington, J. D., Sutton, B. P., Banich, M. T., & Heller, W. (2011). Trait approach and avoidance motivation: lateralized neural activity associated with executive function. *Neuroimage*, *54*(1), 661-670. doi: 10.1016/j.neuroimage.2010.08.037
- Stroop, J. R. (1935). Studies of interference in serial verbal reactions. *Journal of experimental psychology*, *18*(6), 643.
- Tao, L., Marzecová, A., Taft, M., Asanowicz, D., & Wodniecka, Z. (2011). The efficiency of attentional networks in early and late bilinguals: the role of age of acquisition. *Frontiers in psychology*, *2*.
- Thiel, C. M., & Fink, G. R. (2007). Visual and auditory alertness: modality-specific and supramodal neural mechanisms and their modulation by nicotine. *J Neurophysiol*, *97*(4), 2758-2768. doi: 10.1152/jn.00017.2007
- Thiele, A., & Bellgrove, M. A. (2018). Neuromodulation of Attention. *Neuron*, *97*(4), 769-785. doi: 10.1016/j.neuron.2018.01.008
- Toba, M. N., Rabuffetti, M., Duret, C., Pradat-Diehl, P., Gainotti, G., & Bartolomeo, P. (2018). Component deficits of visual neglect: "Magnetic" attraction of attention vs. impaired spatial working memory. *Neuropsychologia*, *109*, 52-62. doi: 10.1016/j.neuropsychologia.2017.11.034
- Trautwein, F.-M., Singer, T., & Kanske, P. (2016). Stimulus-Driven Reorienting Impairs Executive Control of Attention: Evidence for a Common Bottleneck in Anterior Insula. *Cerebral Cortex*, *26*(11), 4136-4147.
- Turken, A. U., & Swick, D. (1999). Response selection in the human anterior cingulate cortex. *Nat Neurosci*, *2*(10), 920-924. doi: 10.1038/13224
- Vallesi, A. (2012). Organisation of executive functions: hemispheric asymmetries. *Journal of Cognitive Psychology*, *24*(4), 367-386.
- Vallesi, A., Arbula, S., Capizzi, M., Causin, F., & D'Avella, D. (2015). Domain-independent neural underpinning of task-switching: an fMRI investigation. *Cortex*, *65*, 173-183. doi: 10.1016/j.cortex.2015.01.016
- Vallesi, A., & Crescentini, C. (2011). Right fronto-parietal involvement in monitoring spatial trajectories. *Neuroimage*, *57*(2), 558-564. doi: 10.1016/j.neuroimage.2011.04.061
- Verleger, R., Smigasiewicz, K., & Moller, F. (2011). Mechanisms underlying the left visual-field advantage in the dual stream RSVP task: evidence from N2pc, P3, and distractor-evoked VEPs. *Psychophysiology*, *48*(8), 1096-1106. doi: 10.1111/j.1469-8986.2011.01176.x
- Verleger, R., Sprenger, A., Gebauer, S., Fritzmanna, M., Friedrich, M., Kraft, S., & Jaskowski, P. (2009). On why left events are the right ones: neural mechanisms underlying the left-hemifield advantage in rapid serial visual presentation. *J Cogn Neurosci*, *21*(3), 474-488. doi: 10.1162/jocn.2009.21038
- Vossel, S., Weidner, R., Driver, J., Friston, K. J., & Fink, G. R. (2012). Deconstructing the architecture of dorsal and ventral attention systems with dynamic causal modeling. *J Neurosci*, *32*(31), 10637-10648. doi: 10.1523/JNEUROSCI.0414-12.2012
- Wang, L., Liu, X., Guise, K. G., Knight, R. T., Ghajar, J., & Fan, J. (2010). Effective connectivity of the fronto-parietal network during attentional control. *Journal of Cognitive Neuroscience*, *22*(3), 543-553. doi: 10.1162/jocn.2009.21210
- Ward, L. M. (1994). Supramodal and modality-specific mechanisms for stimulus-driven shifts of auditory and visual attention. *Canadian Journal of Experimental Psychology*, *48*(2), 242-257.
- Weekes, N. Y., & Zaidel, E. (1996). The effects of procedural variations on lateralized Stroop effects. *Brain Cogn*, *31*(3), 308-330. doi: 10.1006/brcg.1996.0049

Right hemisphere advantage for executive control

- Weintraub, S., & Mesulam, M. M. (1987). Right cerebral dominance in spatial attention. Further evidence based on ipsilateral neglect. *Arch Neurol*, *44*(6), 621-625.
- Welcome, S. E., & Chiarello, C. (2008). How dynamic is interhemispheric interaction? Effects of task switching on the across-hemisphere advantage. *Brain Cogn*, *67*(1), 69-75. doi: 10.1016/j.bandc.2007.11.005
- Wiegand, I., Petersen, A., Bundesen, C., & Habekost, T. (2017). Phasic alerting increases visual attention capacity in younger but not in older individuals. *Visual Cognition*, *25*(1-3), 343-357.
- Wu, C. T., Weissman, D. H., Roberts, K. C., & Woldorff, M. G. (2007). The neural circuitry underlying the executive control of auditory spatial attention. *Brain research*, *1134*, 187-198.
- Xuan, B., Mackie, M.-A., Spagna, A., Wu, T., Tian, Y., Hof, P. R., & Fan, J. (2016). The activation of interactive attentional networks. *NeuroImage*, *129*, 308-319. doi: 10.1016/j.neuroimage.2016.01.017
- Yang, Z., & Mayer, A. R. (2013). An event-related fMRI study of exogenous orienting across vision and audition. *Hum Brain Mapp*. doi: 10.1002/hbm.22227
- Zago, L., Petit, L., Jobard, G., Hay, J., Mazoyer, B., Tzourio-Mazoyer, N., . . . Mellet, E. (2017). Pseudoneglect in line bisection judgement is associated with a modulation of right hemispheric spatial attention dominance in right-handers. *Neuropsychologia*, *94*, 75-83. doi: 10.1016/j.neuropsychologia.2016.11.024
- Zani, A., & Proverbio, A. M. (2017). How voluntary orienting of attention and alerting modulate costs of conflict processing. *Sci Rep*, *7*, 46701. doi: 10.1038/srep46701
- Zuanazzi, A., & Cattaneo, L. (2017). The right hemisphere is independent from the left hemisphere in allocating visuospatial attention. *Neuropsychologia*, *102*, 197-205. doi: 10.1016/j.neuropsychologia.2017.06.005

Right hemisphere advantage for executive control

Table 1. Operational definition of the attentional effects and interactions




Alerting effect	No cue	<i>minus</i>	Double cue
Orienting effect	Double cue	<i>minus</i>	Valid cue
Disengaging effect	Invalid cue	<i>minus</i>	Double cue
Validity effect	Invalid cue	<i>minus</i>	Valid cue
Conflict effect	Flanker incongruent	<i>minus</i>	Flanker congruent
Alerting by Conflict	no cue, incongruent <i>minus</i> no cue, congruent	<i>minus</i>	double cue, incongruent <i>minus</i> double cue, congruent
Orienting by Conflict	double cue, incongruent <i>minus</i> double cue, congruent	<i>minus</i>	valid cue, incongruent <i>minus</i> valid cue, congruent
Disengaging by Conflict	invalid cue, incongruent <i>minus</i> invalid cue, congruent	<i>minus</i>	double cue, incongruent <i>minus</i> double cue, congruent
Validity by Conflict	invalid cue, incongruent <i>minus</i> invalid cue, congruent	<i>minus</i>	valid cue, incongruent <i>minus</i> valid cue, congruent

RUNNING HEAD: Right hemisphere advantage for executive control of attention



Table 2. Mean reaction time in ms (\pm SD) and Error Rate in percentage (\pm SD) for each task condition and separately for the two hemispheres

RT	Congruent				Incongruent			
	nocue	double	valid	invalid	nocue	double	valid	invalid
R	563	529	504	568	687	634	590 (90)	719
H	(67)	(70)	(66)	(73)	(112)	(111)		(118)
L	570	531	493	573	702	675	606 (96)	720
H	(87)	(68)	(61)	(78)	(109)	(107)		(108)
ER								
R	2.95	1.56	2.43	3.82	14.93	11.28	10.13	23.09
H	(5.83)	(3.71)	(4.27)	(6.19)	(14.17)	(12.45)	(10.53)	(18.05)
L	4.17	2.95	1.80	2.26	15.80	14.41	9.20	19.79
H	(6.66)	(4.70)	(2.40)	(4.46)	(13.12)	(15.54)	(9.54)	(17.58)

Cue



Target




100 ms

0 – 800 ms
Mean = 400 ms

500 ms

2000 - 12000 ms
Mean = 4000 ms





no cue




double cue



spatial cue

valid


invalid









a. Alerting x Conflict



b. Orienting x Conflict



c. Disengaging x Conflict



d. Validity x Conflict

