



**HAL**  
open science

## On some open questions concerning determinantal inequalities

Mohammad M. Ghabries, Hassane Abbas, Bassam Mourad

► **To cite this version:**

Mohammad M. Ghabries, Hassane Abbas, Bassam Mourad. On some open questions concerning determinantal inequalities. *Linear Algebra and its Applications*, 2020, 596, pp.169 - 183. 10.1016/j.laa.2020.03.009 . hal-03489558

**HAL Id: hal-03489558**

**<https://hal.science/hal-03489558v1>**

Submitted on 20 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

## On some open questions concerning determinantal inequalities

Mohammad M. GHABRIES<sup>a,b,\*</sup>, Hassane ABBAS<sup>c</sup>, Bassam MOURAD<sup>c</sup>

<sup>a</sup>LAREMA, Faculte des Sciences-Departement de mathematiques, Angers, France

<sup>b</sup>KALMA, Department of Mathematics, Faculty of Science, Lebanese University, Beirut, Lebanon

<sup>c</sup>Department of Mathematics, Faculty of Science, Lebanese University, Beirut, Lebanon

---

### Abstract

In 2017, M. Lin formulated two conjectures concerning determinantal inequalities for positive semi-definite matrices  $A$  and  $B$ , and which can be stated as follows

$$\det(A^2 + |AB|^p) \geq \det(A^2 + |BA|^p) \quad \text{for } p \geq 0$$

and

$$\det(A^2 + |AB|^p) \geq \det(A^2 + A^p B^p) \quad \text{for } 0 \leq p \leq 2.$$

The main goal of this paper is to confirm the first conjecture in a slightly more general setting namely in the case when  $A$  and  $B$  are Hermitian, and also to prove the second conjecture when  $0 \leq p \leq \frac{4}{3}$ . Various related inequalities are then presented and we conclude with an open log-majorization question.

*Keywords:* Determinantal inequalities; Hermitian matrix; Positive semi-definite matrix; Log-majorization; Eigenvalues; Furuta inequality.

*2010 MSC:* 15A45, 15A60, 47A64

---

### 1. Introduction

Audenaert [2] proved the following determinantal inequality, for  $n \times n$  positive semi-definite matrices  $A$  and  $B$ ,

$$\det(A^2 + |BA|) \leq \det(A^2 + AB) \quad (1)$$

that answers a question arising in the study of interpolation methods for image processing in diffusion tensor imaging. Recently, Lin [6] generalized Audenaert's result by proving

$$\det(A^2 + |BA|^p) \leq \det(A^2 + A^p B^p), \quad 0 \leq p \leq 2. \quad (2)$$

In the same paper, he complemented (1) by proving that

$$\det(A^2 + |AB|) \geq \det(A^2 + AB). \quad (3)$$

Clearly, Inequalities (1) and (3) imply that

$$\det(A^2 + |AB|) \geq \det(A^2 + |BA|). \quad (4)$$

---

\*Corresponding author

URL: [mahdi.ghabries@gmail.com](mailto:mahdi.ghabries@gmail.com) (Mohammad M. GHABRIES)

8 In the same paper, it was asked whether it is possible to find a generalization of (4) and put forward the  
 9 following conjecture.

**Conjecture 1.1.** *Let  $A$  and  $B$  be positive semi-definite matrices. Then, for all  $p \geq 0$ , we have*

$$\det(A^2 + |AB|^p) \geq \det(A^2 + |BA|^p).$$

10 In addition, he introduced the following conjecture which is a generalization of (3) and a complementing  
 11 result for (2).

**Conjecture 1.2.** *Let  $A$  and  $B$  be  $n \times n$  positive semi-definite matrices. Then, for  $0 \leq p \leq 2$  it holds that*

$$\det(A^2 + |AB|^p) \geq \det(A^2 + A^p B^p).$$

12 Lin [6] was able to prove Conjecture 1.1 for  $p = 1$  and for all  $p$  positive even integers, and Conjecture  
 13 1.2 for  $p = 0, 1$  and  $2$ . All other cases for both conjectures remain unsolved.

14  
 15 The main purpose of this paper is to confirm Conjecture 1.1 in a slightly more general setting; namely in  
 16 the case where  $A$  and  $B$  are Hermitian matrices, and also to show that Conjecture 1.2 is valid for  $0 \leq p \leq \frac{4}{3}$ .  
 17 In addition, we shall prove that the determinantal inequality of Conjecture 1.2 is also true for all  $2 \leq p \leq 4$ ,  
 18 however it remains open for  $\frac{4}{3} < p < 2$ .

19 To proceed, we first fix some notation. Let  $M_n$  be the space of  $n \times n$  complex matrices where its  
 identity matrix is denoted by  $I_n$ . The modulus of a complex matrix  $X$  is defined as  $|X| = (X^* X)^{1/2}$ .  
 As usual, we shall write  $X \geq 0$  to indicate that  $X$  is positive semi-definite. Also, for Hermitian matrices  
 $X, Y \in M_n$ , we shall say that  $X \geq Y$  if  $X - Y$  is positive semi-definite matrix. Moreover, if the eigenvalues  
 $\lambda_1(X), \lambda_2(X), \dots, \lambda_n(X)$  of a matrix  $X$  are real, then we shall always assume that they are arranged in  
 decreasing order, that is

$$\lambda_1(X) \geq \lambda_2(X) \geq \dots \geq \lambda_n(X).$$

For a Hermitian matrix  $X \in M_n$ , we shall denote

$$\lambda(X) = (\lambda_1(X), \lambda_2(X), \dots, \lambda_n(X))^t$$

20 which is clearly a real vector of order  $n$ .

21  
 22 Majorization relations are great tools for deriving determinantal inequalities, see for example [9, Chapter  
 23 10] for more details on this subject. If  $\lambda(A), \lambda(B) \in \mathbb{R}_+^n$ , then by  $\lambda(A) \prec_{wlog} \lambda(B)$ , we mean that  $\lambda(A)$  is  
 24 *weakly log-majorized* by  $\lambda(B)$ , that is

$$\prod_{i=1}^k \lambda_i(A) \leq \prod_{i=1}^k \lambda_i(B) \quad \text{for all } k = 1, 2, \dots, n. \quad (5)$$

25 In addition, we shall write  $\lambda(A) \prec_{log} \lambda(B)$  and we will say that  $\lambda(A)$  is *log-majorized* by  $\lambda(B)$  if (5) is  
 26 true and equality holds for  $k = n$ .

27 **2. Comparing  $\det(A^2 + |BA|^p)$  with  $\det(A^2 + A^p B^p)$  when  $A, B \geq 0$ .**

The main purpose here is to study the relation between the two determinantal quantities

$$\det(A^2 + |BA|^p) \text{ and } \det(A^2 + A^p B^p) \text{ for } p \in [0, +\infty) \text{ and with } A, B \geq 0.$$

28 We shall start with the following lemma which is obtained from a result proved by A. Matsumoto, R.  
29 Nakamoto and M. Fujii [7] by using anti-symmetric tensor product argument.

30 **Lemma 2.1.** *Let  $X$  and  $Y$  be two positive semi-definite matrices. Then,*

- 31 1.  $\lambda(X^{\frac{k+t}{2}} Y^t X^{\frac{k+t}{2}}) \prec_{wlog} \lambda(X^{\frac{k}{2}} (X^{\frac{1}{2}} Y X^{\frac{1}{2}})^t X^{\frac{k}{2}})$  for all  $0 \leq t \leq 1$  and  $k \geq 0$ .  
32 2.  $\lambda(X^{\frac{k+t}{2}} Y^t X^{\frac{k+t}{2}}) \succ_{wlog} \lambda(X^{\frac{k}{2}} (X^{\frac{1}{2}} Y X^{\frac{1}{2}})^t X^{\frac{k}{2}})$  for all  $t \geq 1$  and  $0 \leq k \leq 1$ .

33 Next, we prove the following elementary lemma which constitutes one of the basis of our main results.  
34 In fact, it is a slight generalization<sup>1</sup> of Lemma A on page 129 of [5].

**Lemma 2.2.** *Let  $X$  and  $Y$  be two invertible matrices. Then, for all  $t \in \mathbb{R}$ ,*

$$(X^* Y^* Y X)^t = X^* Y^* (Y X X^* Y^*)^{t-1} Y X.$$

35 *Proof.* Let  $X^* Y^* = U |X^* Y^*|$  be the polar decomposition of the matrix  $X^* Y^*$ , where  $U$  is unitary. Then,  
36 clearly we obtain  $U = X^* Y^* |X^* Y^*|^{-1}$ ,  $U^* = |X^* Y^*|^{-1} Y X$ , and  $Y X = |X^* Y^*| U^*$ .

$$\begin{aligned} (X^* Y^* Y X)^t &= (U |X^* Y^*| |X^* Y^*| U^*)^t \\ &= (U |X^* Y^*|^2 U^*)^t \\ &= U |X^* Y^*|^{2t} U^* \\ &= X^* Y^* |X^* Y^*|^{-1} |X^* Y^*|^{2t} |X^* Y^*|^{-1} Y X \\ &= X^* Y^* (Y X X^* Y^*)^{t-1} Y X. \end{aligned}$$

37

□

38 The next lemma is also needed for our purposes and it shows a close connection between log-majorization  
39 and determinantal inequalities and can be found in [6, (P2)].

**Lemma 2.3.** *Let  $X$  and  $Y$  be two matrices in  $M_n$ . If  $\lambda(X), \lambda(Y)$  are in  $\mathbb{R}_+^n$  such that  $\lambda(X) \prec_{wlog} \lambda(Y)$ ,  
then*

$$\det(I_n + X) \leq \det(I_n + Y).$$

40 As a consequence, we have the following result.

41 **Theorem 2.1.** *Let  $A$  and  $B$  be two positive semi-definite matrices. Then*

- 42 1.  $\det(A^2 + |BA|^p) \geq \det(A^2 + A^p B^p)$  for  $2 \leq p \leq 4$ .  
43 2. If  $0 \leq p \leq 2$  or  $p \geq 4$ , then  $\det(A^2 + |BA|^p) \leq \det(A^2 + A^p B^p)$ .

---

<sup>1</sup>Thanks to the reviewer for pointing this out.

44 *Proof.* 1. Without loss of generality, we shall assume that  $A$  and  $B$  are positive definite matrices as the  
 45 general case can be then obtained by a continuity argument. For  $2 \leq p \leq 4$ , then we can write

$$\begin{aligned}
 \lambda(A^{-1}(AB^2A)^{\frac{p}{2}}A^{-1}) &= \lambda\left(A^{-1}(AB(BA^2B)^{\frac{p}{2}-1}BA)A^{-1}\right) && \text{(by Lemma 2.2)} \\
 &= \lambda\left(B(BA^2B)^{\frac{p}{2}-1}B\right) \\
 &= \lambda\left((B^2)^{\frac{1}{2}}\left((B^2)^{\frac{1}{2}}(A^2)(B^2)^{\frac{1}{2}}\right)^{\frac{p}{2}-1}(B^2)^{\frac{1}{2}}\right) \\
 &\succ_{wlog} \lambda\left((B^2)^{1+\frac{p}{2}-1}(A^2)^{\frac{p}{2}-1}\right) && \text{(by Lemma 2.1, Part 1)} \\
 &= \lambda(A^{p-2}B^p).
 \end{aligned}$$

46 So,

$$\lambda(A^{-1}(AB^2A)^{\frac{p}{2}}A^{-1}) \succ_{wlog} \lambda(A^{p-2}B^p). \quad (6)$$

47 Next, applying Lemma 2.3 on (6) gives

$$\det(I_n + A^{-1}(AB^2A)^{\frac{p}{2}}A^{-1}) \geq \det(I_n + A^{p-2}B^p). \quad (7)$$

48 Multiplying both sides of (7) by  $\det(A^2) > 0$  completes the proof of the first part.

49

50 2. The proof for the case where  $0 \leq p \leq 2$ , is due to [6], while the case where  $p \geq 4$  can be done in a  
 51 similar fashion as in the first case by making use this time of Part 2 of Lemma 2.1.  $\square$

### 52 3. Proof of Conjecture 1.1

53 The starting point here is a lemma dealing with a majorization inequality which appears in [8]. For the  
 54 sake of completeness, we reproduce the proof here.

**Lemma 3.1.** *Let  $Y$  be a positive semi-definite matrix and  $X$  be any Hermitian matrix. Then for all  $p, q \in [0, +\infty)$ , it holds that*

$$\lambda(XY^pXY^q) \prec_{wlog} \lambda(X^2Y^{p+q}).$$

*Proof.* By appealing to a standard argument (anti-symmetric product), then it suffices to prove that

$$\lambda_1(XY^pXY^q) \leq \lambda_1(X^2Y^{p+q}).$$

Without loss of generality, we shall assume that  $X$  is invertible as the general case can be done by continuity argument. In addition, we shall assume that  $q \leq p$  and  $\lambda_1(X^2Y^{p+q}) = 1$ . Now, obviously proving our claim is equivalent to showing that

$$\lambda_1(XY^pXY^q) \leq 1.$$

The fact that the largest eigenvalue of the matrix  $X^2Y^{p+q}$  is equal to 1, clearly implies that

$$\lambda_j(X^2Y^{p+q}) \leq 1 \text{ for all } 1 \leq j \leq n.$$

55 But this is equivalent to  $XY^{p+q}X \leq I_n$  which in turn gives

$$Y^{p+q} \leq (X^{-1})^2. \quad (8)$$

56 Next, applying Lowner-Heinz on (8) for a power  $0 \leq \frac{p}{p+q} \leq 1$ , yields

$$Y^p \leq ((X^{-1})^2)^{\frac{p}{p+q}}. \quad (9)$$

Again, taking a power  $0 \leq \frac{q}{p} \leq 1$  in both sides of (9), we get

$$Y^q \leq ((X^{-1})^2)^{\frac{q}{p+q}}.$$

57 Hence,

$$(X^2)^{\frac{q}{p+q}} \leq Y^{-q}. \quad (10)$$

58 Therefore,

$$\begin{aligned} \lambda_1(XY^pXY^q) &= \lambda_1\left(Y^{q/2}XY^pXY^{q/2}\right) \\ &\leq \lambda_1\left(Y^{q/2}X((X^{-1})^2)^{\frac{p}{p+q}}XY^{q/2}\right) && \text{(by using (9))} \\ &= \lambda_1\left(Y^{q/2}(X^2)^{\frac{q}{p+q}}Y^{q/2}\right) \\ &\leq \lambda_1\left(Y^{q/2}Y^{-q}Y^{q/2}\right) && \text{(by using (10))} \\ &= \lambda_1(I_n) \\ &= 1. \end{aligned}$$

59

□

60 As a result, we have the following theorem.

61 **Theorem 3.1.** *Let  $Y$  be a positive definite matrix and  $X$  be a Hermitian matrix. Then for all  $p, q \in [0, \infty)$*

$$\lambda(XY^pXY^{-q}) \succ_{wlog} \lambda(X^2Y^{p-q}). \quad (11)$$

62 *Proof.* The proof will be divided into three cases.

Case 1: Let  $q \geq 2p$ . By Schur's complement we have

$$M = \begin{bmatrix} Y^{-\frac{q}{2}}XY^pXY^{-\frac{q}{2}} & Y^{-\frac{q}{2}}X^2Y^{p-\frac{q}{2}} \\ Y^{p-\frac{q}{2}}X^2Y^{-\frac{q}{2}} & Y^{p-\frac{q}{2}}XY^{-p}XY^{p-\frac{q}{2}} \end{bmatrix} \geq 0.$$

Applying Theorem 10.20 in [9, p. 352] gives

$$\lambda_1\left(Y^{-\frac{q}{2}}XY^pXY^{-\frac{q}{2}}\right) \cdot \lambda_1\left(Y^{p-\frac{q}{2}}XY^{-p}XY^{p-\frac{q}{2}}\right) \geq \lambda_1\left(Y^{-\frac{q}{2}}X^2Y^{p-\frac{q}{2}}\right)^2.$$

63 That is,

$$\lambda_1(XY^pXY^{-q}) \cdot \lambda_1(XY^{2p-q}XY^{-p}) \geq \lambda_1(X^2Y^{p-q})^2. \quad (12)$$

64 In view of Lemma 3.1, it is worthy to observe that

$$\begin{aligned} \lambda(X(Y^{-1})^{q-2p}X(Y^{-1})^p) &\prec_{wlog} \lambda(X^2(Y^{-1})^{q-p}) \\ &= \lambda(X^2Y^{p-q}). \end{aligned}$$

65 Therefore,

$$\begin{aligned}\lambda_1(Y^{2p-q}XY^{-p}X) &= \lambda_1(X(Y^{-1})^{q-2p}X(Y^{-1})^p) \\ &\leq \lambda_1(X^2Y^{p-q}).\end{aligned}$$

Thus, from (12) we obtain

$$\lambda_1(XY^pXY^{-q}) \geq \lambda_1(X^2Y^{p-q}) \quad \text{for } q \geq 2p \geq 0.$$

66 By a standard anti-symmetric tensor product argument, inequality (11) is true for all  $q \geq 2p$ .

67

68 **Case 2:** Let  $p \leq q \leq 2p$ . The idea of the proof here depends on writing the interval  $[p, 2p] = \bigcup_{k=2}^{\infty} [\frac{k+1}{k}p, 2p]$   
69 and then proving (11) for each subinterval. We start with with case  $k = 2$  i.e. for  $2p \geq q \geq \frac{3p}{2}$ . Then,

$$\begin{aligned}\lambda(Y^pXY^{-q}X) &= \lambda\left(Y^{\frac{3p}{2}}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})Y^{p-q}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})Y^{\frac{p}{2}}\right) \\ &= \lambda\left(Y^{2p}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})Y^{-(q-p)}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})\right) \\ &= \lambda\left((Y^{-1})^{-2p}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})(Y^{-1})^{q-p}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})\right).\end{aligned}$$

70 Now considering this last expression and noticing that  $2p \geq 2(q-p) \geq 0$ , then in view of Case 1;  
71 replacing  $Y$  with  $Y^{-1}$ ,  $X$  with  $Y^{-\frac{p}{2}}XY^{-\frac{p}{2}}$ ,  $2p$  with  $q$  and lastly  $q-p$  with  $p$ , we obtain

$$\begin{aligned}\lambda(Y^pXY^{-q}X) &\succ_{wlog} \lambda\left(Y^{3p-q}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})^2\right) \\ &= \lambda(Y^{2p-q}XY^{-p}X) \\ &\succ_{wlog} \lambda(Y^{p-q}X^2) \quad (\text{again using Case 1 as } p \geq 2(2p-q) \geq 0).\end{aligned}$$

72 Next, using a similar argument, we prove (11) is true for  $k = 3$  i.e. for  $2p \geq q \geq \frac{4p}{3}$ . As earlier, we  
73 can write

$$\begin{aligned}\lambda(Y^pXY^{-q}X) &= \lambda\left(Y^{\frac{3p}{2}}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})Y^{p-q}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})Y^{\frac{p}{2}}\right) \\ &= \lambda\left(Y^{2p}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})Y^{-(q-p)}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})\right) \\ &= \lambda\left((Y^{-1})^{-2p}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})(Y^{-1})^{q-p}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})\right) \\ &\succ_{wlog} \lambda\left(Y^{3p-q}(Y^{-\frac{p}{2}}XY^{-\frac{p}{2}})^2\right) \quad (\text{in view of of Case 1 as } 2p \geq 2(q-p) \geq 0) \\ &= \lambda(Y^{2p-q}XY^{-p}X) \\ &\succ_{wlog} \lambda(Y^{p-q}X^2) \quad (\text{similarly in view of Case } k = 2 \text{ as } p \geq \frac{3}{2}(2p-q) \geq 0).\end{aligned}$$

74 Continuing this way, one can easily see (using induction) that for any positive integer  $k$ , inequality  
75 (11) is true for all  $q$  with  $2p \geq q \geq \frac{(k+1)p}{k}$ . Finally, the proof for this case can be achieved by letting  
76  $k$  tends to infinity.  
77

78 **Case 3:** Let  $q \leq p$ . To complete the proof of this case, it suffices to apply the preceding two cases on  $Y^{-1}$ .

79

□

80 As applications, we have the following result.

**Corollary 3.1.** *Let  $A$  and  $B$  be two Hermitian matrices. Then, for all  $p \in [0, \infty)$  we have*

$$\det(A^2 + |BA|^p) \leq \det(A^2 + |AB|^p).$$

81 *Proof.* As usual, we shall assume that  $A$  and  $B$  are invertible, the general case is by continuity argument.

82 Then, for all  $p \in [0, +\infty)$  we have

$$\begin{aligned} \lambda\left(A^{-1}(AB^2A)^{\frac{p}{2}}A^{-1}\right) &= \lambda\left(B(BA^2B)^{\frac{p}{2}-1}B\right) && \text{(Using Lemma 2.2)} \\ &= \lambda\left(B^2(BA^2B)^{\frac{p}{2}-1}\right) \\ &\prec_{\log} \lambda\left(B(BA^2B)^{\frac{p}{2}}B(BA^2B)^{-1}\right) && \text{(Using Theorem 3.1)} \\ &= \lambda\left(A^{-1}(BA^2B)^{\frac{p}{2}}A^{-1}\right). \end{aligned}$$

Next, applying Lemma 2.3 gives

$$\det(I_n + A^{-1}(AB^2A)^{\frac{p}{2}}A^{-1}) \leq \det(I_n + A^{-1}(BA^2B)^{\frac{p}{2}}A^{-1}).$$

Finally, multiplying both sides with  $\det(A^2) > 0$  yields

$$\det(A^2 + |BA|^p) \leq \det(A^2 + |AB|^p).$$

83

□

#### 84 **4. Conjecture 1.2**

85 In this section, our purpose is to find for what values of  $k$  and  $t$  the following majorization inequality

$$\lambda(A^{\frac{k}{2}-t}B^t) \prec_{\log} \lambda(A^{\frac{k}{4}}(B^{\frac{1}{2}}A^{-1}B^{\frac{1}{2}})^tA^{\frac{k}{4}}) \quad (13)$$

86 is valid, where  $A$  and  $B$  are positive definite matrices.

87

88 We shall start here with the following lemma which is needed for our purposes and is well known as  
89 Furuta's inequality [4].

**Lemma 4.1.** *Let  $A, B$  be two positive semi-definite matrices such that  $A \geq B$ . Then, for all  $p \geq 1, r \geq 0$ ,*

$$A^{(p+2r)/p} \geq (A^r B^p A^r)^{1/p}.$$

90 Now we are in a position to prove the next theorem which shows that (13) is valid for all  $0 \leq t \leq 1$  and  
91  $k \geq 4t$ .

**Theorem 4.1.** *Let  $A$  and  $B$  be two positive definite matrices. Then for all  $0 \leq t \leq 1$  and  $k \geq 4t$*

$$\lambda(A^{\frac{k}{2}-t}B^t) \prec_{\log} \lambda(A^{\frac{k}{4}}(B^{\frac{1}{2}}A^{-1}B^{\frac{1}{2}})^tA^{\frac{k}{4}}).$$



*Proof.* Let  $0 \leq t \leq 1$  and  $k \geq 4t$ . Using Schur's complement, we know that

$$M = \begin{bmatrix} A^{\frac{k}{4}-t} B^t (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^t B^t A^{\frac{k}{4}-t} & A^{\frac{k}{4}-t} B^t A^{\frac{k}{4}} \\ A^{\frac{k}{4}} B^t A^{\frac{k}{4}-t} & A^{\frac{k}{4}} (B^{\frac{1}{2}} A^{-1} B^{\frac{1}{2}})^t A^{\frac{k}{4}} \end{bmatrix} \geq 0.$$

92 Then

$$\left( \lambda_1(A^{\frac{k}{2}-t} B^t) \right)^2 \leq \lambda_1(A^{\frac{k}{4}} (B^{\frac{1}{2}} A^{-1} B^{\frac{1}{2}})^t A^{\frac{k}{4}}) \cdot \lambda_1(A^{\frac{k}{4}-t} B^t (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^t B^t A^{\frac{k}{4}-t}). \quad (14)$$

93 As mentioned earlier, in order to prove our claim, then it is enough to prove that

$$\lambda_1(A^{\frac{k}{2}-t} B^t) \geq \lambda_1(A^{\frac{k}{4}-t} B^t (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^t B^t A^{\frac{k}{4}-t}), \quad (15)$$

which is in turn equivalent to showing that

$$B^{\frac{t}{2}} A^{\frac{k}{2}-t} B^{\frac{t}{2}} \leq I_n \Rightarrow A^{\frac{k}{4}-t} B^t (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^t B^t A^{\frac{k}{4}-t} \leq I_n.$$

For this, let  $B^{\frac{t}{2}} A^{\frac{k}{2}-t} B^{\frac{t}{2}} \leq I_n$ , then  $A^{\frac{k}{2}-t} \leq B^{-t}$ . First, making use of Löwner-Heinz inequality for  $0 \leq \frac{t}{\frac{k}{2}-t} \leq 1$  gives

$$A^t \leq B^{-\frac{t^2}{\frac{k}{2}-t}}.$$

Next, applying Lemma 4.1 yields

$$\left( B^{-\frac{t^2}{\frac{k}{2}-t}} \right)^{\frac{p+2r}{p}} \geq \left[ \left( B^{-\frac{t^2}{\frac{k}{2}-t}} \right)^r (A^t)^p \left( B^{-\frac{t^2}{\frac{k}{2}-t}} \right)^r \right]^{\frac{1}{p}}.$$

94 Now, replacing  $p$  with  $\frac{1}{t} \geq 1$  and  $r$  with  $\frac{\frac{k}{2}-t}{2t^2} \geq 0$  gives

$$B^{-\frac{\frac{k}{2}-t}{\frac{k}{2}-t}} \geq (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^t.$$

95 Pre-post multiplying both sides by  $B^t > 0$  implies that

$$(B^t)^{\frac{\frac{k}{2}-2t}{\frac{k}{2}-t}} = B^t B^{-\frac{\frac{k}{2}-t}{\frac{k}{2}-t}} B^t \geq B^t (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^t B^t. \quad (16)$$

However,  $A^{\frac{k}{2}-t} \leq B^{-t}$ , so that  $A^{-(\frac{k}{2}-t)} \geq B^t$ . By again appealing to Löwner-Heinz inequality for  $0 \leq \frac{\frac{k}{2}-2t}{\frac{k}{2}-t} \leq 1$  we obtain

$$(A^{-(\frac{k}{2}-t)})^{\frac{\frac{k}{2}-2t}{\frac{k}{2}-t}} \geq (B^t)^{\frac{\frac{k}{2}-2t}{\frac{k}{2}-t}},$$

96 which gives

$$A^{-(\frac{k}{2}-2t)} \geq (B^t)^{\frac{\frac{k}{2}-2t}{\frac{k}{2}-t}}. \quad (17)$$

97 Now it is worthy to observe that inequalities (16) and (17) yield

$$A^{-(\frac{k}{2}-2t)} \geq B^t (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^t B^t.$$

98 Hence,  $A^{\frac{k}{4}-t}B^t(B^{-\frac{1}{2}}AB^{-\frac{1}{2}})^tB^tA^{\frac{k}{4}-t} \leq I_n$ , and therefore, (15) is true for all  $0 \leq t \leq 1$  and  $k \geq 4t$ .  
On the other hand, using (14) and (15) gives

$$\lambda_1(A^{\frac{k}{2}-t}B^t) \leq \lambda_1(A^{\frac{k}{4}}(B^{\frac{1}{2}}A^{-1}B^{\frac{1}{2}})^tA^{\frac{k}{4}}), \quad 0 \leq t \leq 1, \quad k \geq 4t.$$

Thus, by a standard anti-symmetric tensor product argument, we get

$$\lambda(A^{\frac{k}{2}-t}B^t) \prec_{w \log} \lambda(A^{\frac{k}{4}}(B^{\frac{1}{2}}A^{-1}B^{\frac{1}{2}})^tA^{\frac{k}{4}}), \quad 0 \leq t \leq 1, \quad k \geq 4t.$$

Finally, the proof is complete by making use of the fact that

$$\det(A^{\frac{k}{2}-t}B^t) = \det(A^{\frac{k}{4}}(B^{\frac{1}{2}}A^{-1}B^{\frac{1}{2}})^tA^{\frac{k}{4}}).$$

99

□

100 Our next goal is to show that (13) is also true for all  $\frac{1}{2} \leq t \leq 1$  and  $k \geq 6t - 2$ . First, we need the  
101 following lemma.

**Lemma 4.2.** *Let  $A$  and  $B$  be two positive definite matrices. Then, for all  $\frac{1}{2} \leq t \leq 1$  and  $k \geq 6t - 2$ , we have*

$$\lambda\left(A^{\frac{k+2t}{4}}\left(A^{-\frac{1}{2}}BA^{-\frac{1}{2}}\right)^{2t}A^{\frac{2t+k}{4}}\right) \succ_{\log} \lambda(A^{\frac{k}{2}-t}B^{2t}).$$

*Proof.* As in similar situations, it is enough to prove that for all  $\frac{1}{2} \leq t \leq 1$  and  $k \geq 6t - 2$

$$\lambda_1\left(A^{\frac{k+2t}{4}}\left(A^{-\frac{1}{2}}BA^{-\frac{1}{2}}\right)^{2t}A^{\frac{2t+k}{4}}\right) \geq \lambda_1(A^{\frac{k}{2}-t}B^{2t}).$$

First, in view of Lemma 2.2, we obtain the following equality:

$$A^{\frac{2t+k}{4}}\left(A^{-\frac{1}{2}}BA^{-\frac{1}{2}}\right)^{2t}A^{\frac{2t+k}{4}} = A^{\frac{2t+k-2}{4}}B^{\frac{1}{2}}\left(B^{\frac{1}{2}}A^{-1}B^{\frac{1}{2}}\right)^{2t-1}B^{\frac{1}{2}}A^{\frac{2t+k-2}{4}}.$$

102 For the sake of clarification, we shall use the following notation. For all  $\frac{1}{2} \leq t \leq 1$  and  $k \geq 6t - 2$ , let

$$\begin{aligned} X &= A^{\frac{2t+k-2}{4}}B^{\frac{1}{2}}\left(B^{\frac{1}{2}}A^{-1}B^{\frac{1}{2}}\right)^{2t-1}B^{\frac{1}{2}}A^{\frac{2t+k-2}{4}}, \\ Y &= A^{\frac{2t+k-2}{4}}B^{2t}A^{\frac{k-6t+2}{4}}, \quad \text{and} \\ Z &= A^{\frac{k-6t+2}{4}}B^{2t-\frac{1}{2}}\left(B^{-\frac{1}{2}}AB^{-\frac{1}{2}}\right)^{2t-1}B^{2t-\frac{1}{2}}A^{\frac{k-6t+2}{4}}. \end{aligned}$$

Now by making use of Schur complement, the following  $2 \times 2$  block matrix

$$M = \begin{bmatrix} X & Y \\ Y^* & Z \end{bmatrix}$$

is positive semi-definite and hence  $\lambda_1(X) \cdot \lambda_1(Z) \geq \lambda_1(Y)^2$ . Our next goal is to show that  $\lambda_1(Z) \leq \lambda_1(Y)$  which in turn gives  $\lambda_1(X) \geq \lambda_1(Y)$ . Noticing that  $\lambda_1(Y) = \lambda_1(B^tA^{\frac{k}{2}-t}B^t)$ , then in order to prove  $\lambda_1(Z) \leq \lambda_1(Y)$ , it suffices to show that

$$B^tA^{\frac{k}{2}-t}B^t \leq I_n \Rightarrow Z \leq I_n.$$

For this purpose, let  $B^t A^{\frac{k}{2}-t} B^t \leq I_n$ , then clearly  $A^{\frac{k}{2}-t} \leq B^{-2t}$ . Using Löwner-Heinz inequality for  $0 \leq \frac{2t-1}{\frac{k}{2}-t} \leq 1$  gives

$$A^{2t-1} \leq B^{-\frac{2t(2t-1)}{\frac{k}{2}-t}}.$$

Now, applying Lemma 4.1 yields

$$\left( B^{-\frac{2t(2t-1)}{\frac{k}{2}-t}} \right)^{\frac{p+2r}{p}} \geq \left[ \left( B^{-\frac{2t(2t-1)}{\frac{k}{2}-t}} \right)^r (A^{2t-1})^p \left( B^{-\frac{2t(2t-1)}{\frac{k}{2}-t}} \right)^r \right]^{\frac{1}{p}}.$$

Next, taking  $p = \frac{1}{2t-1} \geq 1$  and  $r = \frac{\frac{k}{2}-t}{4t(2t-1)} \geq 0$  implies that

$$B^{-\frac{(\frac{k}{2}+t)(2t-1)}{\frac{k}{2}-t}} \geq (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^{2t-1}.$$

Pre-post multiplying both sides with  $B^{2t-\frac{1}{2}} > 0$  gives

$$B^{2t-\frac{1}{2}} B^{-\frac{(\frac{k}{2}+t)(2t-1)}{\frac{k}{2}-t}} B^{2t-\frac{1}{2}} = (B^{2t})^{\frac{k-6t+2}{\frac{k}{2}-t}} \geq B^{2t-\frac{1}{2}} (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^{2t-1} B^{2t-\frac{1}{2}}.$$

Again, noticing that  $A^{\frac{k}{2}-t} \leq B^{-2t}$  implies that  $A^{-(\frac{k}{2}-t)} \geq B^{2t}$ , then by appealing to Löwner-Heinz for  $0 \leq \frac{\frac{k-6t+2}{2}}{\frac{k}{2}-t} \leq 1$  we obtain

$$(A^{-(\frac{k}{2}-t)})^{\frac{k-6t+2}{2}} \geq (B^{2t})^{\frac{k-6t+2}{\frac{k}{2}-t}}$$

which gives

$$A^{-(\frac{k-6t+2}{2})} \geq B^{2t-\frac{1}{2}} (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^{2t-1} B^{2t-\frac{1}{2}}.$$

Hence,

$$Z = A^{\frac{k-6t+2}{4}} B^{2t-\frac{1}{2}} (B^{-\frac{1}{2}} A B^{-\frac{1}{2}})^{2t-1} B^{2t-\frac{1}{2}} A^{\frac{k-6t+2}{4}} \leq I_n.$$

Therefore, for all  $\frac{1}{2} \leq t \leq 1$  and  $k \geq 6t - 2$ ,

$$\lambda_1(X) \geq \lambda_1(Y).$$

Thus, by an anti-symmetric tensor product argument and by the fact that  $\det(X) = \det(Y)$ , we get

$$\lambda(X) \succ_{\log} \lambda(Y).$$

103 Finally, noting that  $\lambda(X) = \lambda \left( A^{\frac{k+2t}{4}} \left( A^{-\frac{1}{2}} B A^{-\frac{1}{2}} \right)^{2t} A^{\frac{2t+k}{4}} \right)$  and  $\lambda(Y) = \lambda(A^{\frac{k}{2}-t} B^{2t})$ , the proof is  
104 complete.  $\square$

**Theorem 4.2.** *Let  $A$  and  $B$  be two positive definite matrices. Then, for all  $\frac{1}{2} \leq t \leq 1$  and  $k \geq 6t - 2$*

$$\lambda \left( A^{\frac{k}{4}} \left( B^{\frac{1}{2}} A^{-1} B^{\frac{1}{2}} \right)^t A^{\frac{k}{4}} \right) \succ_{\log} \lambda(A^{\frac{k}{2}-t} B^t).$$

105 *Proof.* Let  $\frac{1}{2} \leq t \leq 1$  and  $k \geq 6t - 2$ . Then, we can write

$$\begin{aligned}
\lambda \left( A^{\frac{k}{4}} \left( B^{\frac{1}{2}} A^{-1} B^{\frac{1}{2}} \right)^t A^{\frac{k}{4}} \right) &= \lambda \left( A^{\frac{k}{4}} \left( A^{\frac{1}{2}} (A^{-\frac{1}{2}} B^{\frac{1}{2}} A^{-1} B^{\frac{1}{2}} A^{-\frac{1}{2}}) A^{\frac{1}{2}} \right)^t A^{\frac{k}{4}} \right) \\
&\succ_{\log} \lambda \left( A^{\frac{k}{2}+t} \left( A^{-\frac{1}{2}} B^{\frac{1}{2}} A^{-1} B^{\frac{1}{2}} A^{-\frac{1}{2}} \right)^t \right) \quad (\text{by Part 1 of Lemma 2.1}) \\
&= \lambda \left( A^{\frac{k}{2}+t} \left( A^{-\frac{1}{2}} B^{\frac{1}{2}} A^{-\frac{1}{2}} \right)^{2t} \right) \\
&= \lambda \left( A^{\frac{k+2t}{4}} \left( A^{-\frac{1}{2}} B^{\frac{1}{2}} A^{-\frac{1}{2}} \right)^{2t} A^{-\frac{k+2t}{4}} \right) \\
&\succ_{\log} \lambda \left( A^{\frac{k}{2}-t} (B^{\frac{1}{2}})^{2t} \right) \quad (\text{by Lemma 4.2}) \\
&= \lambda(A^{\frac{k}{2}-t} B^t).
\end{aligned}$$

106

□

107 As a result of Theorem 4.1 and Theorem 4.2, we have the following corollary.

**Corollary 4.1.** *Let  $A$  and  $B$  be two positive semi-definite matrices. Then, for  $(0 \leq p \leq 2$  and  $k \geq 2p)$  or for  $(1 \leq p \leq 2$  and  $k \geq 3p - 2)$  we have*

$$\det(A^k + |AB|^p) \geq \det(A^k + A^p B^p).$$

108 *Proof.* Without loss of generality we may assume that  $A$  and  $B$  are positive definite matrices, the general  
109 case is by continuity argument. Suppose that  $0 \leq p \leq 2$  and  $k \geq 2p$ . Then, replacing  $A$  with  $A^{-2}$ ,  $B$  with  
110  $B^2$  and  $t$  with  $0 \leq \frac{p}{2} \leq 1$  in Theorem 4.1 gives

$$\lambda(A^{p-k} B^p) \prec_{\log} \lambda(A^{-\frac{k}{2}} (BA^2 B)^{\frac{p}{2}} A^{-\frac{k}{2}}). \quad (18)$$

Applying Lemma 2.3 on (18) yields

$$\det(I_n + A^{-\frac{k}{2}} (BA^2 B)^{\frac{p}{2}} A^{-\frac{k}{2}}) \geq \det(I_n + A^{p-k} B^p).$$

Next, multiplying both sides with  $\det(A^k) > 0$ , we obtain

$$\det(A^k + |AB|^p) \geq \det(A^k + A^p B^p), \quad 0 \leq p \leq 2, \quad k \geq 2p.$$

111 For the case when  $1 \leq p \leq 2$  and  $k \geq 3p - 2$ , the proof can be done in a similar fashion by making use  
112 this time of Theorem 4.2. □

113 As an analogue of Theorem 2.1, Corollary 3.1, and Corollary 4.1, we have the following result which  
114 gives a partial answer of Conjecture 1.2.

**Theorem 4.3.** *Let  $A$  and  $B$  be two positive semi-definite matrices. Then, for all  $0 \leq p \leq \frac{4}{3}$  or  $2 \leq p \leq 4$ , the following holds*

$$\det(A^2 + |AB|^p) \geq \det(A^2 + A^p B^p).$$

*Proof.* Corollary 4.1 is a general case of

$$\det(A^2 + |AB|^p) \geq \det(A^2 + A^p B^p), \quad 0 \leq p \leq \frac{4}{3}.$$

Combining Part 1 of Theorem 2.1 with Corollary 3.1 implies that for all  $2 \leq p \leq 4$

$$\det(A^2 + |AB|^p) \geq \det(A^2 + |BA|^p) \geq \det(A^2 + A^p B^p).$$

115

□

116 We conclude this paper with the following conjecture which is very much related to our work in this  
117 section, and whose validation for  $k = 2$  would imply Conjecture 1.2.

**Conjecture 4.1.** *Let  $A$  and  $B$  be two positive definite matrices. Then for all  $k \geq 0$  and  $0 \leq t \leq 1$ ,*

$$\lambda \left( A^{\frac{k}{4}} \left( B^{\frac{1}{2}} A^{-1} B^{\frac{1}{2}} \right)^t A^{\frac{k}{4}} \right) \succ_{\log} \lambda(A^{\frac{k}{2}-t} B^t).$$

## 118 Acknowledgements

119 The authors sincerely thank the reviewer for his valuable comments and encouragement. Also, thanks to  
120 the handling Editor for his useful suggestions. The authors acknowledge financial support from the Lebanese  
121 University research grants program.

## 122 Declaration of Competing Interest

123 The authors declare that they have no competing interests.

## 124 References

- 125 [1] H. Abbas and M.M. Ghabries, Some Generalizations and Complements of Determinantal Inequalities,  
126 Math. Inequal. Appl. v23, Number 1 (2020), 169–176.
- 127 [2] K.M.R. Audenaert, A determinantal inequality for the geometric mean with an application in diffusion  
128 tensor, 2015, arXiv:1502.06902v2.
- 129 [3] R. Bhatia, Matrix Analysis, GTM 169, Springer-Verlag, New York, 1997.
- 130 [4] T. Furuta,  $A \geq B \geq 0$  assures  $(B^r A^p B^r)^{1/q} \geq B^{(p+2r)/q}$  for  $r \geq 0, p \geq 0, q \geq 1$  with  $(1 + 2r)q \geq$   
131  $p + 2r$ , Proc. Amer. Math. Soc. 101 (1987) 85-88.
- 132 [5] T. Furuta, Invitation to Linear Operators: From matrices to bounded linear operators on a Hilbert space,  
133 CRC Press, Taylor Francis Group, 2001.
- 134 [6] M. Lin, On a determinantal inequality arising from diffusion tensor imaging, Commun. Contemp.  
135 Math. 19 (2017), 1650044, 6 pp.
- 136 [7] A. Matsumoto, R. Nakamoto, M. Fujii, Reverse of Bebiano-Lemos-Providencia inequality and Com-  
137 plementary Furuta inequality (Inequalities on Linear Operators and its Applications), Departmental  
138 Bulletin Paper, Kyoto University, 2008, pp.91-98.

- 139 [8] H. Abbas, M.M. Ghabries, B. Mourad, New determinantal inequalities concerning Hermitian and pos-  
140 itive semi-definite matrices, (2019) submitted.
- 141 [9] F. Zhang, Matrix Theory: Basic Results and Techniques, Springer, New York, 2nd ed., 2011.