



Biological Applications and Toxicity Minimization of Semiconductor Quantum Dots

Samira Filali, Fabrice Pirot, Pierre Miossec

► To cite this version:

Samira Filali, Fabrice Pirot, Pierre Miossec. Biological Applications and Toxicity Minimization of Semiconductor Quantum Dots. Trends in Biotechnology, 2020, 38, pp.163 - 177. <10.1016/j.tibtech.2019.07.013>. <hal-03489685>

HAL Id: hal-03489685

<https://hal.science/hal-03489685v1>

Submitted on 21 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

Biological applications and toxicity minimization of semiconductor quantum dots

Samira Filali^{a,b}, Fabrice Pirot^b, Pierre Miossec^{a*}

^a Immunogenomics and Inflammation Research Unit EA 4130, Department of Immunology and Rheumatology, Edouard Herriot Hospital, Hospices Civils de Lyon, University of Lyon, Lyon, France

^b Laboratory of Research and Development of Industrial Galenic Pharmacy and laboratory of Tissue Biology and Therapeutic Engineering UMR-CNRS 5305, Pharmacy Department, FRIPHARM Plateform, Edouard Herriot Hospital, Hospices Civils de Lyon, University of Lyon, Lyon, France, Fripharm.com

*Correspondence: samira.filali@chu-lyon.fr (Dr. Samira FILALI)

Keywords: quantum dots, stem cells, differentiated cells, organs, animal species

Abstract

The extraordinary potential of semiconductor quantum dots (QDs) has resulted in their widespread application in various fields, from engineering technology and development of laboratory techniques to biomedical imaging and therapeutic strategies. However, the toxicity of QDs remains a concern and has limited their applications in human health. Better understanding the behavior of QDs as it relates to their composition will enable exploring their limitations and developing a strategy to control their toxicity for potential therapeutic applications. Here, approaches to minimize their toxicities according to the specific cell type, organ, or animal species are described. This review summarizes recent promising works at the levels of cells, organs, and whole organisms.

Potential of QDs in the biological fields

The multitude of researchers exploring the remarkable potential of **quantum dots** (QDs) (see Glossary, Box 1 and 2) in biomedical research, from tools in medical imaging to the development of sophisticated theranostic agents, indicate the great potential of QDs in this field. Among the variety of nanoprobes (upconversion nanoparticles, fluorescent proteins, and graphene-based nanomaterials), semiconductor QDs have distinguished themselves in the field of imaging due to their size and their optical properties (stable fluorescence, brightness) and cost-effectiveness relative to conventional fluorescent dyes. Their extremely small size (order of magnitude of biological molecules, 1 -10 nm) makes these QDs ideal candidates for identification and tracking of biomolecules and organelles inside the cell and especially in the nucleus, unlike upconversion nanoparticles that have a larger size (20-50 nm) and cannot reach the nucleus[1]. Moreover, unlike carbon dots, red fluorescence emission for QDs semiconductors, which is the preferred color in biological applications due to cells autofluorescence phenomena in the 400 to 600 nm range (yellow, green, blue), is possible with semiconductor QDs. As a result, despite the toxicity conferred by semiconductor QDs (although recently controversial for so-called "biocompatible" nanoparticles), better understanding the occurrence of their toxicity and strategies to minimize their toxicity would enable their better use. In targeted imaging, QDs targeting tumors or biomarkers play a crucial role in the detection, treatment, monitoring, and prognostic evaluation of certain diseases. However, for purely diagnostic applications, in which cell damage must be avoided, the potential toxicity caused by the release of metal ions remains a major limitation to the use of QDs, thereby hindering their translation from preclinical studies to clinical studies in biomedical imaging. A number of reviews have been published to address various questions, including the recent progress made on QD surface modification, describing their structure-activity relationships (Figure 1), mechanisms of uptake in cells, and biomedical applications. However, few reviews

have specifically presented a body of knowledge that enables a detailed overview on the behavior of QDs including or not means to minimize toxicity according to the specific cell type, tissue/organ, or animal species. This review aims to collate information on recent advances in the use of QDs in biological fields, with the objective of describing the various effects of QDs reported thus far, and to elucidate their behavior in living organisms, notably for future applications in humans.

Application of QDs at the cellular level

Stem cells (see Glossary) are able to multiply indefinitely *in vitro* in an undifferentiated state, and can differentiate in the presence of adequate factors. QDs have proved to be of great value in cell imaging, particularly in the monitoring of cell differentiation, and in functional studies. However, the biological effect may differ depending on the type of stem cell or differentiated cell (Table 1).

Stem cells

The hemocompatibility of QDs was assessed in hematopoietic stem cells by combining Tat Peptide with CdSe/ZnS-QDs, which made it possible to monitor transplantation successfully [2]. However, in differentiated hematopoietic cells (human blood neutrophil granulocytes, monocytes, lymphocytes, and platelets), harmful effects were observed that were specific to both their morphology (maximum accumulation of QDs with monocytes and minimum with lymphocytes) and function (despite the presence of the shell, neutrophils underwent cell death after phagocytosis of CdSe/ZnS-QDs, and CdTe-QD-induced platelet aggregation in the absence of plasma) [3,4].

Concerning epithelial stem cells, depending on QD composition, the effects may be either deleterious or beneficial. Several studies have provided explanations for the cytotoxic effects

77 of QDs, which occur notably via interaction of CdTe-QDs with aromatic rings and amine
78 groups from trans-membrane proteins and the cytoskeleton[5], and the predomination of the
79 overall cellular response, which is mediated by the minor non-endocytic uptake of CdSe/ZnS-
80 QDs in BEAS-2B human bronchial epithelial cells [6]. Despite the use of multi-shell CdSe-
81 CdS/ZnS-QDs in the Calu-3 human lung cell monolayer, mechanical stress and activation of
82 Ca^{2+} influx were induced, affecting the lung barrier [7]. However, a different study showed that
83 these QDs did not elicit noticeable changes in the long-term **TEER** (see Glossary) after a
84 70 day-exposure, suggesting their potential use for drug delivery at the surface of the lung
85 epithelium for short periods [8]. Glass incorporation in CdSe-QDs in place of coating by greater
86 polymer-enabled reduction of the cytotoxicity on HaCaT keratinocytes. The higher rigidity and
87 density of glass relative to polymer enabled more efficient blocking of Cd and Se ions, making
88 these QDs safer for biological applications [9].

89 Nerve stem cells (NSCs) appear to be more resistant than other stem cells. After uptake, mainly
90 via microfilaments and microtubules, CdSe/ZnS-polymer-QDs and graphene-QDs had little
91 effect on the growth and proliferation of primary NSCs within a week[10,11]. CdSe/ZnS-QDs
92 did not affect either differentiation potential of NSCs or the protein expression of neuron- and
93 astrocyte-derived primary nerve stem cells [11]. In the GT1-7 differentiated neuronal cell line,
94 CdSe/ZnS-QDs were incorporated into only a limited fraction of the whole lysosomal
95 compartment, inducing a slight but significant reduction in cell survival and proliferation [12].

96 Mesenchymal stem cells are important for synthesizing and repairing the musculoskeletal
97 system (such as cartilage cells, bone cells, fat cells, synovial cells, or myoblasts). QDs have
98 been mainly used for the characterization of these cell types, with a majority of the QDs having
99 no negative impact on cell viability and differentiation [13,14]. The use of ZnS/CdSe-
100 streptavidin-QDs conjugated with a Fab fragment of an anti-myc antibody enabled the tracking
101 of GLUT4 movement in 3T3L1 adipocytes after stimulation by insulin or in an insulin-resistant

state [15]. The composition of CdSe/ZnS-polymer-QDs conjugated with a unique antibody against a heat shock chaperone (mortalin) allowed the successful tracking of mesenchymal stem cells labeled during tissue repair, following 26 weeks of allogeneic transplantation in a rabbit model of osteochondral defects [16]. The construction of a siRNA-coupled CdSe/ZnS-PEG-QDs has been successfully achieved, and has both demonstrated the involvement of *SOX9* in chondrogenesis and enabled the real-time monitoring of transfection [17]. CdTe-QDs have allowed the simultaneous characterization of two types of cadmium-induced cell death (apoptosis and autophagy) [18].

QD toxicity in male and female germ cells and during embryonic development has also been evaluated. At concentrations between 1 and 125 nM of CdSe-QDs, no toxicity was detected in germ cells (i.e., in terms of viability and motility of spermatozoa) and embryos. At 500 nM, toxic effects on both pre-and post-implantation embryo development and female germ cells (with decreased maturation and fertilization rates) were reported, which were reduced with the incorporation of a ZnS shell. The transfer across the placental barrier resulted in Cd accumulation in pups, which was reduced after coating with PEG or SiO₂ [19].

Differentiated cells

Overall, Cd²⁺-based QDs had deleterious effects in hepatic cells, including CdSe/ZnS-QDs-induced L02 hepatocyte pyroptosis, liver inflammation, and dysfunction[20], as well as CdTe-QD-induced mitochondrial damage in human hepatocellular carcinoma cells (HepG2), which were not solely attributable to cadmium released from QDs, as the observed effects were similar or greater to those of cadmium chloride[21]. However, graphene-QDs did not induce cytotoxicity in mouse primary hepatocytes [22].

An extensive investigation of the signaling mechanism involved in the induction of cytotoxicity of shell-less QDs has been performed in endothelial cells; notably, the pathways implicated in

127 CdTe-QDs-mediated mitochondrial-dependent apoptosis in human umbilical vein endothelial
128 cells have been studied in detail [23]. Nevertheless, the use of shells or an amorphous form [24]
129 seems sufficient to reduce their toxicity. Furthermore, additional applications may emerge
130 following the discovery of stronger accumulation of CdSe/ZnS-QDs in slow-flowing, small
131 caliber venous vessels than in fast-flowing, high caliber arterial vessels [25] and the large-scale
132 internalization of CdSe/ZnS-QDs conjugated to monoclonal antibodies targeting the murine
133 transferrin receptor (Ri7) by brain capillary endothelial cells [26].

134 The effects of unshelled QDs in nephrocytes have been discussed in detail. L-cysteine-capped
135 CdTe-QDs decreased cell viability and modified the structure and activity of copper–zinc
136 superoxide dismutase in mouse primary nephrocytes and hepatocytes. Notably, this effect was
137 abrogated by the addition of ascorbic acid [27]. The study of pig renal cells revealed the
138 involvement of the Wnt pathway in CdTe-QDs-induced damage [27]. The evaluation of the
139 nephrotoxicity in HK-2 human epithelial tubular cells using various metallic QDs with
140 differential solubility (TiO₂, ZnO, and CdS) revealed that soluble CdS- and ZnO-QDs elicited
141 cell death in a dose-dependent manner, and this effect was attributed to the release of metallic
142 cations. However, insoluble TiO₂-QDs had no cytotoxic effect [28]. Targeting bladder
143 urothelial cancer cells was performed successfully using CdSe/ZnS-QDs conjugated with the
144 prostate stem cell antigen [29].

145 Recently, promising applications in both diagnostics and therapy of microbial infections have
146 been reported. QD-specific antibodies designed for use as biosensors via the association of
147 rabbit anti-p38 antibody and CdSe/ZnS-QDs, were shown to successfully infiltrate porous
148 silicon whose pores contained immobilized Egp38 antigen. These QDs were used to detect
149 *Echinococcus granulosus* for early diagnosis of hydatid disease [30]. Photoexcited QDs have
150 been used to kill a wide range of multidrug-resistant bacterial clinical isolates, including
151 methicillin-resistant *Staphylococcus aureus*, carbapenem-resistant *Escherichia coli*, extended-

spectrum β -lactamase-producing *Klebsiella pneumoniae*, and *Salmonella typhimurium*, while leaving mammalian cells intact. Their proposed application in clinical phototherapy is suggested for the treatment of infections [31].

Application of QDs in organs

Ex vivo investigation would enable elucidation of QD-induced effects on a collection of specific cells in their environment. The different applications of QDs were grouped by organ system in this review (Table 2).

Integumentary system

Several studies have investigated QD penetration through the skin. In one of these studies, CdSe/ZnS- mPEG-5000-QDs were injected subcutaneously into CDF1 mice or deposited onto human skin. The results showed that QD diffusion in the dermis is limited by the basement membrane and dense connective tissue fibers; this resulted in negligible QD penetration into the epidermis, hair follicles, sebaceous and sweat glands, nerves, and blood vessels. The total penetration of QDs could only be achieved via a 10-minute massage of damaged or stripped stratum corneum. An increase in pH (8.3) of QD formulations enabled some degree of QD penetration of intact skin [32–34]. Furthermore, simultaneous use of cell therapy and graphene QDs accelerated the repair of skin lesions [35].

Brain system

Owing to their nanoscale size, which enables them to cross the blood–brain barrier, QDs represent a revolutionary theranostic tool for the brain system. CdSe/ZnS-streptavidin-QDs injected by stereotaxis into the mouse brain were able to specifically label the microglia in primary cortical cultures via macrophage scavenger receptors and mannose receptors present

on their surface, and did not induce toxicity (i.e. induction of inflammation) in the short term [36]. Recently, graphene-QDs were shown to successfully reduce symptoms in mice primed to develop Parkinson's disease by inhibiting fibrillization of α -syn and interacting directly with mature fibrils, thus triggering their disaggregation. These QDs were also to inhibit the aggregation of A β peptides which leads to Alzheimer's disease. These promising results constitute a considerable advance in the treatment of complex pathologies for which no effective therapies are available[37,38].

Digestive system

The digestive system is essential for the assimilation of nutrients, immune and endocrine defenses, and the metabolism of endogenous or exogenous molecules (e.g., drugs). A study using CdTe/CdS-QDs administrated by intravenous injection showed that QDs were retained in the sinusoids and selectively taken up by sinusoidal cells (Kupffer cells and liver sinusoidal endothelial cells) within 3 h, but not by hepatocytes[39]. However, the absorption of QDs at the intestinal level has been poorly studied. Previous studies focusing on nanoparticle absorption in general describe the requirement for well-defined characteristics to cross the mucus barrier and the intestinal epithelium lining the intestinal lumen. In nematodes (representative of intestine models), CdTe-QDs influenced lipid metabolism via alteration of the molecular basis of both synthesis and degradation of fatty acid, but not degradation of phospholipids, and increased in fat storage in intestine [40]. The spleens of rats were also exposed by CdSe/CdS-PEG-QDs coupled or not by folic acid with different doses(0–300 nM), and both were revealed to be non-toxic. Moreover, the addition of folic acid significantly improved the uptake of QDs in a dose-dependent manner, highlighting the importance of surface coating properties [41]. VEGFR2-CdSe/ZnS-QDs were used to successfully distinguish between diseased and non-diseased tissue in the colons of carcinogen-treated mice with high

sensitivity and specificity. However, the complete removal of the contrast agent by flushing is difficult to achieve [42,43].

Pulmonary system

The respiratory system is the preferred site for exchange of gas and airborne particles between the body and the external environment. The evaluation of the efficacy or toxicology of QDs in the pulmonary system is therefore essential. A short-term exposure to 240 µg CdO-QDs/m³ for seven days (3 h/day) caused pulmonary injury and tissue remodeling in mice, and altered immune function [44]. Intratracheal instillation of CdSe/ZnS-QDs or CdSe/ZnS-PEG-QDs persistently induced acute neutrophil infiltration, followed by interstitial lymphocyte infiltration, transiently reduced pulmonary function, and a granulomatous reaction on day 17. Although the granulomatous reaction disappeared for CdSe/ZnS-PEG-QDs at day 90, PEGylation was not sufficient to reduce the risk of side effects associated with QDs [45]. Upon further exploration, genetic predisposition factors and atopy (in this case dust mite allergy) were found to induce impaired pulmonary mechanics by exposure to QDs [46,47].

Urinary system

The urinary system is essential for the excretion of endogenous and exogenous substances that are either catabolized or intact. Studies of the kidneys of mice have shown that CdTe-QDs deplete glutathione levels, which reduce the ability of the liver and kidneys to eliminate hydroxyl and superoxide anion radicals, thus inducing oxidative damage to tissues. However, a partial recovery was observed after 28 days [48,49]. Free CdSe/ZnS-QDs and anti-CD47-CdSe/ZnS-QDs were intravesically instilled in mice. Examination of the *in vivo* biodistribution of anti-CD47-CdSe/ZnS-QDs revealed the absence of significant accumulation of QDs outside

of the bladder, and no acute toxicity up to 7 days post-administration. Mice showed extravesical biodistribution of QDs under some conditions; however, this was rare [50].

Cardiovascular system

The cardiovascular system transports blood from the heart to the peripheral regions of the body and vice versa, and carries a wide variety of substances. The ideal geometry and functionalization capabilities of QDs make them an excellent tool for visualizing the cardiovascular system. Accordingly, CdSe/ZnS-QDs have proved their utility in the study of sarcomere dynamics during excitation-contraction coupling in healthy and diseased cardiac muscle and in the whole heart [51]. By combining *in vivo* cryotechniques and the injection of glutathione-coated QDs into kidneys, QD distribution in the vascular system was achieved, primarily in glomerular blood capillaries for a few seconds and extending to peritubular blood capillaries by five seconds [52]. With regard to therapy, the amorphous form of SeQDs has proved to be effective in prevention of endothelial dysfunction and atherosclerosis [24].

Musculoskeletal system

The musculoskeletal system gives form, stability, and movement to the human body. The application of QDs on this system thus far has been confined to imaging. The use of CdSe/ZnS-streptavidin-QDs bound to biotin-phalloidin, which labels skeletal actin filaments, has enabled rapid, quantitative, and inexpensive imaging of sarcomeric movements, notably step-size measurement for low-duty cycle muscle myosin [51,53]. A recent study reported that QDs were mainly found in the synovial membrane, but significantly less on cartilage, one week after being intra-articularly injected into the joints of healthy or osteoarthritis horses [14]. Among the selected articles, no studies were conducted to evaluate the spread of QDs in the bone system, and QD accumulation in bone remains to be defined.

Applications of QDs in animals

Invertebrates

In order to evaluate the integrity of water ecosystems, two aquatic species, polyp *Hydra vulgaris* and coral *Stylophora pistillata*, representative of freshwater and sea habitats, respectively, were exposed to CdSe/ZnS-polymer-QDs and CdTe-QDs at concentrations exceeding the Cd²⁺ content in seawater. Despite the presence of a shell and polymer, CdSe/ZnS-polymer-QDs induced dramatic changes, including morphological and global transcriptional changes, albeit less than those induce by CdTe-QDs [54–57]. In order to track the visualization of cyclin E within cells of developing *Xenopus laevis* embryos in real time, CdSe/ZnS-QDs have been coupled to cell-cycle regulatory proteins (recombinant (His6)-cyclin E protein), and their accumulation in the nucleus at midblastula transition (6 h post-fertilization) has been confirmed [58].

Drosophila, which has high genetic homology with the human genome and has innate immunity signaling pathways that are similar to those of mammals, is an excellent model for genotoxicology and immunotoxicology. QDs were administered via the ingestion of QD-supplemented food (CdSe/ZnS- or InP/ZnS-QDs, 100 and 500 pM) and stress response (hsp70, hsp83), genotoxic stress (p53), and apoptotic cell death (Dredd) genes were observed following the administration of CdSe/ZnS-QDs. However, no significant response was observed following the administration of InP/ZnS-QDs [59].

Small vertebrates

Although alternatives to animal studies should be favored, the advantages provided by studies in small animals are not negligible (*i.e.*, transposition of whole biological systems). Various QDs (CdSe/ZnS-QDs, PbS/CdS/ZnS-QDs, CdTe/ZnS-QDs, CdTe-QDs, and QD innovative

structures) and modes of administration (intravenous, intraperitoneal, intratracheal, plantar, and oral) were evaluated in mice and rats (Table 3). CdTe-QDs affected steroid hormone and lysine biosyntheses and taurine metabolism [60]. CdSe/ZnS-QDs injected intravenously in mice at a concentration up to 10 mg/kg body weight (BW) did not induce significant toxic effects during four weeks of administration; however, a dose of 20 mg/kg BW caused death in two-thirds of mice shortly after the start of experiment [61]. Biodistribution kinetics of intravenously injected PbS/CdS/ZnS-QDs at 0.04 mg/mL revealed the presence of QDs in the lungs after short time periods (first 0-30 s) followed by accumulation in the liver (60 – 120 s) with complete disappearance of fluorescence at the injection site. After 25 days, elimination by the urinary system lead to the total elimination of the QDs [62,63]. Similarly, biodistribution studies of a single intravenous dose (0.2 μ mol/kg) of CdTe/ZnS-QDs defined an elimination half-life of 12 - 14 h [64]. After intraperitoneal injection of CdSe/CdS-QDs every 3 days for a period of 15 days, an increase in the levels of LDH, NADPH, and IL-6 pro-inflammatory cytokines was specifically detected in the plasma, liver, and spleen [65]. Globally, QD accumulation in the liver, kidney, and spleen was observed in the four previously cited studies, regardless of route of administration. The injection dose, particle size, and surface charge have been identified as factors affecting mean residence time, apparent volume of distribution, and clearance, particularly in the case of Ag₂S QDs [66]. The assessment of pulmonary toxicity following intratracheal injection of CdSe/ZnS-QDs revealed lung injury and inflammation in a dose-dependent manner, which peaked at days 7 and 14 post-exposure, with a holding time of QD fluorescence of 1 week. The built-up of Cd²⁺ occurred only in lung-associated lymph nodes and kidneys for 28 days [67]. An overdose (50 mg/kg) of phospholipid micelle-encapsulated CdSe-CdS/ZnS-PEG-QDs over 112 days (16 weeks) did not induce toxicity, despite the accumulation of QDs in the liver and spleen in Kunming mice [68]. The evaluation of orally administered 0.2 mg/kg and 0.4 mg/kg BW/day

of CdTe-QDs/quercetin complex for 3 weeks highlighted the utility of QDs as nano-carriers by reducing inflammation significantly and promoting cartilage regeneration [69]. Intraperitoneal injection (10 mg/kg) of graphene QDs effectively ameliorated autoimmune encephalomyelitis [70]. The evaluation of CdSe/ZnS-QDs (10, 20, and 40 mg/kg BW injection) on the male reproductive system at different stages of development was performed in 32 male mice (adult group) and 24 pregnant mice (embryo group) on day 8 of gestation, and no significant effect on fetal testis development and on adult groups at up to 20 mg/kg BW was observed. However, at 40 mg/kg BW, a decrease in lamina propria, destruction in interstitial tissue, deformation of seminiferous tubules, and a reduction in number of spermatogonia, spermatocytes, and spermatids was observed [71–73]. Embryotoxicity following the injection of CdTe-QDs on the 13th day of gestation in pregnant rats was assessed in 121 fetuses and via histological analysis of placentas. This study revealed dose-dependent survival rates of fetuses (Table 3) due to placental damage rather than QD penetration and accumulation in the fetuses [74], and an impairment on the first two generations of placenta growth [75].

Large vertebrates

Biomedical research in large animal models is rare, especially in non-human primates, which are unfortunately irreplaceable as models of certain diseases owing to their close similarity to humans.

In the cellular therapy field, assessment of cell migration and adhesion is of paramount importance. CdSe/ZnS-polymer-QDs were used to successfully label and assess the adhesion of goat adipose tissue-derived stem cells (g-ADSCs) after 30, 60, and 90 days of cell infusion in the goat mammary glands. The labeling did not affect cellularity and morphology, and was maintained long-term, even after freezing of liquid NO₂. The health status of goats following QD administration has not been specified. [76].

Both phospholipid micelle-encapsulated CdSe/CdS/ZnS-QDs (ph-QDs) [77] and F-127-encapsulated-silicon-QDs [78] were administrated to monkeys (25 and 200 mg/kg BW, respectively) via short intravenous transfusion (30 min). For both QDs, no signs of toxicity were clearly detected in terms of general conditions (body mass, physical, mental, and nutritional behavior), biological parameters (biochemical, no signs of infection or allergic or toxic reactions, liver, and renal). The histological analyses have been unchanged despite QD accumulation in kidney, liver, and spleen, indicating that the breakdown and QDs clearance is quite slow, unlike in mice in which there were signs of delayed and time-dependent increased liver damage (inflammation, proliferation of Kupffer cells, multifocal cholestasis, and spotty necrosis of hepatic cells). The fate of the two types of QDs could not be determined [77,78]. The injection of ph-QDs in pregnant monkeys at a gestational age of 100 days resulted in a miscarriage rate of 60% after having crossed the placenta from the mother to fetus and induced acute hepatocellular injury in the mother after a week of administration [79].

Concluding Remarks and Future Perspectives

The rapid expansion of the popularity of QDs has resulted in the publication of a large number of studies on the subject, with high variability in the structure and composition of the QDs studied, thus making synthesis and comparisons difficult (see Outstanding Questions). Nevertheless, 50% of the studies are based on CdSe/ZnS-polymer-QDs produced by the same supplier, which presents both an advantage (comparability of articles) and a limitation (fewer studies on the structure of other semiconductor QDs).

At the cellular level, overall toxic effects appear at a certain threshold concentration that differs according to the sensitivity of each cell type [109]. Stem cells appear to be more resistant under certain conditions than differentiated cells, and the presence of QDs does not seem to affect stem cell differentiation. Depending on the intended application (*in vitro* imaging, diagnostic,

or theranostic), it is essential to define the optimal concentration range to obtain desired effects and toxicity thresholds for each study, and not conclude, without due investigation, that QDs show global cytotoxicity. Interestingly, the results with innovative QD structures incorporating new materials are encouraging for safer use in medical imaging.

With regard to organs, the lungs, kidneys, and the liver are the most affected by toxicity at high concentrations due to QD accumulation. The involvement of these organs is globally observed for all nanoparticles, probably inherent to their size, rather than specifically for QDs [80]. As observed for a number of anticancer and antiretroviral drugs, accumulation in organs might be associated with potential toxicities, but preliminary precautions can be taken (improvement of hydration for nephrotoxic risk, addition of adjuvant protective organ) and does not necessarily constitute an obstacle to their use. Additional studies are needed to adjust QD concentrations according to their use. Surprisingly, the brain seems to be more resistant to the toxic effects of QDs; the same observations have been made at the cellular level, and revolutionary applications in neurodegenerative diseases are being explored.

Based on the observed negative effects on the ecosystem, QDs should be used with the same caution as chemotherapy treatments. The studies carried out on small and large animals are encouraging, taking into account the benefit/risk balance; however, animal studies on concrete cases of pathologies have yet to be conducted.

Regarding strategies for minimizing toxicity, in the case of stem cells, the addition of a shell reduces toxicity, especially by means of polymerization. Biocompatibility has been achieved by incorporating QDs in glass. For differentiated cells, the addition of a shell associated with polymerization is in some cases not sufficient; the use of QDs in amorphous form or the addition of adjuvant alleviates the effects. Other winning strategies such as functionalization or infiltration into silicone have minimized toxicity. Except for the pulmonary system, where the toxicity can be exacerbated due to factors such as genetic predisposition, the addition of shells

associated with polymerization is sufficient to limit the toxicity in the organs. In invertebrates, all cadmium-based QDs, even when combined with a shell, polymerization and functionalization, show toxicity. Only a change to another chemical element in another group reduces the toxicity. For small animals, the preferred means of toxicity minimization is the addition of a multishell. For large vertebrates, all minimization strategies have been successfully applied at once, and these QDs have been proven to be non-toxic, except in the case of pregnancy. Despite their toxicity, semiconductor QDs have been noted for their ability to respond to a wide range of applications. The toxicity of QDs can be exploited favorably to destroy cells that have lost their capacity to die (i.e., tumors) via a controlled toxicology strategy. A strategy of diffusion limitation can also be considered by playing on a galenic formulation without modifying the structure of the QD.

Before their potential is fully exploited in biological application, in-depth studies of pharmacokinetics, taking into account all routes of administration, are necessary and must be supplemented by regimens with repeated administrations. The fate of QDs in the body must be further clarified; particularly, the mechanism of elimination must be elucidated.

To reach clinical and industrial translation by establishing a pharmaceutical product for human clinical trials, close cooperation between different disciplines (chemist, physicists, biologists, pharmacists, pharmacologists, clinicians) and legal instances is paramount in order to take into account biological and environmental considerations and reglementary requirements (Figure 2). For this purpose, it is necessary to agree on a standardized and large-scale QDs preparation method with good and reproducible performance and to define a set of standard quality control for the characterization of QDs. However, between batches, there may always be a lack of reproducibility that should be overcome by additional controls and may be validated by functional tests. Another challenge is to ensure regulations to qualify the raw material in pharmaceutical grade; it is necessary to ensure the absence of heavy metals. However, heavy

401 metals are constituents of semiconductor QDs. Without demonstrating the effectiveness of the
402 toxicity minimization strategies or an argument in favor of controlled toxicology weighing the
403 benefits/risks for certain cases, accompanied by progress in legal regulation, the development
404 of a pharmaceutical preparation that is employable in the clinical setting cannot be envisaged.

405

406 Competing Financial Interests: The authors declare having no conflicts of interest related to this
407 paper.

408

409

410

Box 1. Structure-activity relationship of QDs

A QD is a nanocrystal made of semiconductor materials consisting of a few million atoms (but only a small number of free electrons, ≤ 100) that is small enough to exhibit quantum mechanical properties [81]. QDs can be composed of metallic atoms (e.g. Ni, Co, Pt, Au) or mostly of semiconductor materials, such as elements of periodic group II – VI (CdTe, CdSe, CdS, ZnSe, ZnS, PbS, PbSe, SnTe) [82] or from III – V group such as In and Ga [83]. In addition, QDs can surround by shell often to reduce ion release and therefore, its toxicity [84]. QDs can also be functionalized by adding a terminal function such as - NH₂ or - COOH allowing their hydrophilization [85], polymers to reduce their toxicity or improve their absorption [86], peptides [87] or antibodies [88] to specifically target a biological element (Figure 1).

Box 2. Characterization of quantum dots

Initially, their optical properties are evaluated by UV-visible [89] and photoluminescence spectroscopy [90,91], which is a fast and nondestructive technique allowing for the excitation, absorption, and emission spectra used for calculation of quantum yield to be determined [92]. The band gap studies were determined by optical diffuse reflectance spectra measurement [93,94]. The size and morphology of QDs can be estimated by scanning or transmission electron microscopy [95], dynamic light scattering (DLS) [96,97], or mathematic approaches from absorption edges using Henglein empirical curve [98]. To define the functionalization, Fourier transformed infrared (FTIR) spectroscopic measurements can be used [99]. The analysis of structure and elemental composition can be performed by X-ray diffraction [83,100] and energy dispersive X-ray analysis [101,102], respectively. Non-destructive techniques such as Raman spectroscopy explore the confined electronic structure of QDs [103–105]. QDs dosage suspension can be achieved by high pressure liquid chromatography via size exclusion chromatography with a fluorimetric detector [106]. The metal ion content in final QDs was also

436 assayed by inductively coupled plasma atomic emission spectrometry [107], and QD behavior
437 was studied by electrochemical methods [108].

438

439

440 **Glossary**

441 **Long-term transepithelial electrical resistance TEER:** commonly used *in vitro* for
442 the assessment of the integrity of the epithelial cell layers.

443

444 **Quantum dots:** The term quantum comes from these mechanical properties at the
445 atomic and subatomic scale with zero-dimensional as a fixed point.

446

447 **Stem cells:** consist of three forms: (i) adult stem cells (ASC) also called stem cells
448 specific to the tissues contained in the body (e.g., hematopoietic, epithelial, neuronal,
449 mesenchymal), (ii) embryonic stem cells or (iii) embryonic germ cells.

450

References

- 1 Himmelstoß, S.F. and Hirsch, T. (2019) A critical comparison of lanthanide based upconversion nanoparticles to fluorescent proteins, semiconductor quantum dots, and carbon dots for use in optical sensing and imaging. *Methods Appl Fluoresc* 7, 022002
- 2 Lei, Y. *et al.* (2012) Labeling of hematopoietic stem cells by Tat peptide conjugated quantum dots for cell tracking in mouse body. *J Nanosci Nanotechnol* 12, 6880–6886
- 3 Pleskova, S.N. *et al.* (2018) The interaction between human blood neutrophil granulocytes and quantum dots. *Micron* 105, 82–92
- 4 Samuel, S.P. *et al.* (2015) CdTe quantum dots induce activation of human platelets: implications for nanoparticle hemocompatibility. *Int J Nanomedicine* 10, 2723–2734
- 5 Cepeda-Pérez, E. *et al.* (2016) SERS and integrative imaging upon internalization of quantum dots into human oral epithelial cells. *Journal of Biophotonics* 9, 683–693
- 6 Zhao, X. *et al.* (2015) Effect of nonendocytic uptake of nanoparticles on human bronchial epithelial cells. *Anal. Chem.* 87, 3208–3215
- 7 Yin, H. *et al.* (2017) Quantum dots modulate intracellular Ca²⁺ level in lung epithelial cells. *Int J Nanomedicine* 12, 2781–2792
- 8 Turdalieva, A. *et al.* (2016) Bioelectric and Morphological Response of Liquid-Covered Human Airway Epithelial Calu-3 Cell Monolayer to Periodic Deposition of Colloidal 3-Mercaptopropionic-Acid Coated CdSe-CdS/ZnS Core-Multishell Quantum Dots. *PLoS ONE* 11, e0149915
- 9 Ando, M. *et al.* (2016) Cytotoxicity of CdSe-based quantum dots incorporated in glass nanoparticles evaluated using human keratinocyte HaCaT cells. *Biosci. Biotechnol. Biochem.* 80, 210–213
- 10 Shang, W. *et al.* (2014) The uptake mechanism and biocompatibility of graphene quantum dots with human neural stem cells. *Nanoscale* 6, 5799–5806
- 11 Zhang, J. *et al.* (2011) Labeling primary nerve stem cells with quantum dots. *J Nanosci Nanotechnol* 11, 9536–9542
- 12 Corazzari, I. *et al.* (2013) Localization of CdSe/ZnS quantum dots in the lysosomal acidic compartment of cultured neurons and its impact on viability: potential role of ion release. *Toxicol In Vitro* 27, 752–759
- 13 Kundrotas, G. *et al.* (2019) Uptake and distribution of carboxylated quantum dots in human mesenchymal stem cells: cell growing density matters. *J Nanobiotechnology* 17, 39

- 14 Grady, S.T. *et al.* (2019) Persistence of fluorescent nanoparticle-labelled bone marrow mesenchymal stem cells in vitro and after intra-articular injection. *J Tissue Eng Regen Med* 13, 191–202
- 15 Fujita, H. *et al.* (2010) Identification of Three Distinct Functional Sites of Insulin-mediated GLUT4 Trafficking in Adipocytes Using Quantitative Single Molecule Imaging. *Mol Biol Cell* 21, 2721–2731
- 16 Yoshioka, T. *et al.* Fate of bone marrow mesenchymal stem cells following the allogeneic transplantation of cartilaginous aggregates into osteochondral defects of rabbits. *J Tissue Eng Regen M.* 5, 437–443
- 17 Wu, Y. *et al.* (2016) Functional quantum dot-siRNA nanoplexes to regulate chondrogenic differentiation of mesenchymal stem cells. *Acta Biomater* 46, 165–176
- 18 Filali, S. *et al.* (2018) Live-stream characterization of cadmium-induced cell death using visible CdTe-QDs. *Sci. Rep.* 8, 12614
- 19 Hsieh, M.-S. *et al.* (2009) Cytotoxic Effects of CdSe Quantum Dots on Maturation of Mouse Oocytes, Fertilization, and Fetal Development. *Int J Mol Sci* 10, 2122–2135
- 20 Peng, L. *et al.* (2015) Metallomics Study of CdSe/ZnS Quantum Dots in HepG2 Cells. *ACS Nano* 9, 10324–10334
- 21 Nguyen, K.C. *et al.* (2015) Mitochondrial Toxicity of Cadmium Telluride Quantum Dot Nanoparticles in Mammalian Hepatocytes. *Toxicol. Sci.* 146, 31–42
- 22 Song, S.H. *et al.* (2015) Primary hepatocyte imaging by multiphoton luminescent graphene quantum dots. *Chem. Commun.* 51, 8041–8043
- 23 Yan, M. *et al.* (2016) Cytotoxicity of CdTe quantum dots in human umbilical vein endothelial cells: the involvement of cellular uptake and induction of pro-apoptotic endoplasmic reticulum stress. *Int J Nanomedicine* 11, 529–542
- 24 Zhu, M.-L. *et al.* (2019) Amorphous nano-selenium quantum dots improve endothelial dysfunction in rats and prevent atherosclerosis in mice through Na⁺/H⁺ exchanger 1 inhibition. *Vascul. Pharmacol.* DOI: 10.1016/j.vph.2019.01.005
- 25 Jiang, X.-Y. *et al.* (2017) Quantum dot interactions and flow effects in angiogenic zebrafish (*Danio rerio*) vessels and human endothelial cells. *Nanomedicine* 13, 999–1010
- 26 Paris-Robidas, S. *et al.* (2016) Internalization of targeted quantum dots by brain capillary endothelial cells in vivo. *J. Cereb. Blood Flow Metab.* 36, 731–742
- 27 Trabelsi, H. *et al.* (2013) Subacute toxicity of cadmium on hepatocytes and nephrocytes in the rat could be considered as a green biosynthesis of nanoparticles. *Int J Nanomedicine* 8, 1121–1128

- 28 Pujalte, I. *et al.* (2015) Cytotoxic effects and cellular oxidative mechanisms of metallic nanoparticles on renal tubular cells: impact of particle solubility. *Toxicol Res* 4, 409–422
- 29 Yuan, R. *et al.* (2018) Quantum dot-based fluorescent probes for targeted imaging of the EJ human bladder urothelial cancer cell line. *Exp Ther Med* 16, 4779–4783
- 30 Li, Y. *et al.* (2017) Detection of Echinococcus granulosus antigen by a quantum dot/porous silicon optical biosensor. *Biomed Opt Express* 8, 3458–3469
- 31 Courtney, C.M. *et al.* (2016) Photoexcited quantum dots for killing multidrug-resistant bacteria. *Nat. Mater* 15, 529–534
- 32 Kulvietis, V. *et al.* Distribution of polyethylene glycol coated quantum dots in mice skin. *Exp Dermatol.* 22, 157–159
- 33 Prow, T.W. *et al.* (2012) Quantum dot penetration into viable human skin. *Nanotoxicology* 6, 173–185
- 34 Gratieri, T. *et al.* (2010) Penetration of quantum dot particles through human skin. *J Biomed Nanotechnol* 6, 586–595
- 35 Haghshenas, M. *et al.* (2019) Use of embryonic fibroblasts associated with graphene quantum dots for burn wound healing in Wistar rats. *In Vitro Cell. Dev. Biol. Anim.* 55, 312–322
- 36 Minami, S.S. *et al.* (2012) Selective targeting of microglia by quantum dots. *J Neuroinflammation* 9, 22
- 37 Liu, Y. *et al.* (2015) Graphene quantum dots for the inhibition of β amyloid aggregation. *Nanoscale* 7, 19060–19065
- 38 Kim, D. *et al.* (2018) Graphene quantum dots prevent α -synucleinopathy in Parkinson's disease. *Nat. Nanotechnol.* 13, 812–818
- 39 Liang, X. *et al.* (2015) Intravital Multiphoton Imaging of the Selective Uptake of Water-Dispersible Quantum Dots into Sinusoidal Liver Cells. *Small* 11, 1711–1720
- 40 Wu, Q. *et al.* (2016) Quantum dots increased fat storage in intestine of Caenorhabditis elegans by influencing molecular basis for fatty acid metabolism. *Nanomed-nanotechnol.* 12, 1175–1184
- 41 Haque, M.M. *et al.* (2013) Effects of folic acid and polyethylene glycol coated quantum dots on toxicity and tissue uptake to precision-cut spleen slices of rats. *J Pharm Investig* 43, 375–383
- 42 Carbary-Ganz, J.L. *et al.* (2015) In vivo molecular imaging of colorectal cancer using quantum dots targeted to vascular endothelial growth factor receptor 2 and optical coherence tomography/laser-induced fluorescence dual-modality imaging. *J Biomed Opt* 20, 096015

- 43 Carbary-Ganz, J.L. *et al.* (2014) Quantum dots targeted to vascular endothelial growth factor receptor 2 as a contrast agent for the detection of colorectal cancer. *J Biomed Opt* 19, 086003
- 44 Blum, J.L. *et al.* (2014) Short-term inhalation of cadmium oxide nanoparticles alters pulmonary dynamics associated with lung injury, inflammation, and repair in a mouse model. *Inhal Toxicol* 26, 48–58
- 45 Wu, T. and Tang, M. (2014) Toxicity of quantum dots on respiratory system. *Inhal Toxicol* 26, 128–139
- 46 Scoville, D.K. *et al.* (2018) Quantum dot induced acute changes in lung mechanics are mouse strain dependent. *Inhal Toxicol* 30, 397–403
- 47 Scoville, D.K. *et al.* (2019) Quantum dots and mouse strain influence house dust mite-induced allergic airway disease. *Toxicol. Appl. Pharmacol.* 368, 55–62
- 48 Wang, J. *et al.* (2017) Dose and time effect of CdTe quantum dots on antioxidant capacities of the liver and kidneys in mice. *Int J Nanomedicine* 12, 6425–6435
- 49 Zhao, L. *et al.* (2019) Kidney Toxicity and Response of Selenium Containing Protein-glutathione Peroxidase (Gpx3) to CdTe QDs on Different Levels. *Toxicol. Sci.* 168, 201–208
- 50 Pan, Y. *et al.* (2017) In vivo biodistribution and toxicity of intravesical administration of quantum dots for optical molecular imaging of bladder cancer. *Sci Rep* 7,
- 51 Kobirumaki-Shimozawa, F. *et al.* (2012) Sarcomere Imaging by Quantum Dots for the Study of Cardiac Muscle Physiology. *J Biomed Biotechnol* 2012,
- 52 Terada, N. *et al.* (2016) Application of “In Vivo Cryotechnique” to Visualization of Microvascular Blood Flow in Mouse Kidney by Quantum Dot Injection. In *In Vivo Cryotechnique in Biomedical Research and Application for Bioimaging of Living Animal Organs* pp. 219–221, Springer, Tokyo
- 53 Wang, Y. *et al.* (2013) The Qdot-labeled actin super-resolution motility assay measures low-duty cycle muscle myosin step size. *Biochemistry* 52, 1611–1621
- 54 Ambrosone, A. *et al.* (2017) Dissecting common and divergent molecular pathways elicited by CdSe/ZnS quantum dots in freshwater and marine sentinel invertebrates. *Nanotoxicology* 11, 289–303
- 55 Liu, T. *et al.* (2014) Hematopoiesis toxicity induced by CdTe quantum dots determined in an invertebrate model organism. *Biomaterials* 35, 2942–2951
- 56 Yan, S.-Q. *et al.* (2016) Reproductive toxicity and gender differences induced by cadmium telluride quantum dots in an invertebrate model organism. *Sci Rep* 6, 34182

- 57 Xing, R. *et al.* (2016) Targeting and retention enhancement of quantum dots decorated with amino acids in an invertebrate model organism. *Sci Rep* 6, 19802
- 58 Brandt, Y.I. *et al.* (2015) Quantum dot assisted tracking of the intracellular protein Cyclin E in *Xenopus laevis* embryos. *J Nanobiotechnol* 13, 31
- 59 Alaraby, M. *et al.* (2015) Assessing potential harmful effects of CdSe quantum dots by using *Drosophila melanogaster* as in vivo model. *Sci. Total Environ.* 530–531, 66–75
- 60 Khoshkam, M. *et al.* (2018) Synthesis, characterization and in vivo evaluation of cadmium telluride quantum dots toxicity in mice by toxicometabolomics approach. *Toxicol. Mech. Methods* 28, 539–546
- 61 Bozrova, S.V. *et al.* (2017) Semiconductor quantum dot toxicity in a mouse in vivo model. *J. Phys.: Conf. Ser.* 784, 012013
- 62 Benayas, A. *et al.* (2015) PbS/CdS/ZnS Quantum Dots: A Multifunctional Platform for In Vivo Near-Infrared Low-Dose Fluorescence Imaging. *Adv. Funct. Mater.* 25, 6650–6659
- 63 Yaghini, E. *et al.* (2018) In vivo biodistribution and toxicology studies of cadmium-free indium-based quantum dot nanoparticles in a rat model. *Nanomedicine* 14, 2644–2655
- 64 Liu, N. *et al.* (2013) Degradation of aqueous synthesized CdTe/ZnS quantum dots in mice: differential blood kinetics and biodistribution of cadmium and tellurium. *Part Fibre Toxicol* 10, 37
- 65 Haque, M.M. *et al.* (2013) Acute toxicity and tissue distribution of CdSe/CdS-MPA quantum dots after repeated intraperitoneal injection to mice. *J Appl Toxicol* 33, 940–950
- 66 Javidi, J. *et al.* (2019) Pharmacokinetics, Tissue Distribution and Excretion of Ag₂S Quantum Dots in Mice and Rats: the Effects of Injection Dose, Particle Size and Surface Charge. *Pharm. Res.* 36, 46
- 67 Roberts, J.R. *et al.* (2013) Lung toxicity and biodistribution of Cd/Se-ZnS quantum dots with different surface functional groups after pulmonary exposure in rats. *Part Fibre Toxicol* 10, 5
- 68 Liu, J. *et al.* (2013) Toxicity assessment of phospholipid micelle-encapsulated cadmium-based quantum dots using Kunming mice. *RSC Adv.* 3, 1768–1773
- 69 Jeyadevi, R. *et al.* (2013) Enhancement of anti arthritic effect of quercetin using thioglycolic acid-capped cadmium telluride quantum dots as nanocarrier in adjuvant induced arthritic Wistar rats. *Colloids Surf B Biointerfaces* 112, 255–263
- 70 Tomic, J. *et al.* (2019) Graphene quantum dots inhibit T cell-mediated neuroinflammation in rats. *Neuropharmacology* 146, 95–108

617 71 Amiri, G. *et al.* (2016) Comparison of Toxicity of CdSe: ZnS Quantum Dots on Male
618 Reproductive System in Different Stages of Development in Mice. *Int J Fertil Steril* 9, 512–
619 520

620 72 Xu, G. *et al.* (2016) The Reproductive Toxicity of CdSe/ZnS Quantum Dots on the in
621 vivo Ovarian Function and in vitro Fertilization. *Sci Rep* 6, 37677

622 73 Valipoor, A. *et al.* (2015) A comparative study about toxicity of CdSe quantum dots on
623 reproductive system development of mice and controlling this toxicity by ZnS coverage.
624 *Nanomedicine Journal* 2, 261–268

625 74 Zalgeviciene, V. *et al.* (2017) Quantum dots mediated embryotoxicity via placental
626 damage. *Reprod Toxicol* 73, 222–231

627 75 Hong, W. *et al.* (2019) CdSe/ZnS Quantum Dots Impaired the First Two Generations of
628 Placenta Growth in an Animal Model, Based on the Shh Signaling Pathway. *Nanomaterials*
629 (*Basel*) 9,

630 76 Costa, C.R.M. *et al.* (2017) Labeling of adipose-derived stem cells with quantum dots
631 provides stable and long-term fluorescent signal for ex vivo cell tracking. *In Vitro*
632 *Cell.Dev.Biol.-Animal* 53, 363–370

633 77 Ye, L. *et al.* (2012) A pilot study in non-human primates shows no adverse response to
634 intravenous injection of quantum dots. *Nat. Nanotechnol.* 7, 453–458

635 78 Liu, J. *et al.* (2013) Assessing clinical prospects of silicon quantum dots: studies in mice
636 and monkeys. *ACS Nano* 7, 7303–7310

637 79 Ye, L. *et al.* (2019) Comparing Semiconductor Nanocrystal Toxicity in Pregnant Mice
638 and Non-Human Primates. *Nanotheranostics* 3, 54–65

639 80 Zhang, Y.-N. *et al.* (2016) Nanoparticle–liver interactions: Cellular uptake and
640 hepatobiliary elimination. *J Control Release* 240, 332–348

641 81 Sain, S. *et al.* (2014) Microstructure and photoluminescence properties of ternary
642 Cd_{0.2}Zn_{0.8}S quantum dots synthesized by mechanical alloying. *J Nanopart Res* 16, 2673

643 82 Azpiroz, J.M. *et al.* (2014) Benchmark Assessment of Density Functional Methods on
644 Group II–VI MX (M = Zn, Cd; X = S, Se, Te) Quantum Dots. *J Chem Theory Comput* 10,
645 76–89

646 83 Daly, A.B. *et al.* (2017) Optical and structural properties in type-II InAlAs/AlGaAs
647 quantum dots observed by photoluminescence, X-ray diffraction and transmission electron
648 microscopy. *Superlattices Microstruct.* 110, 1–9

649 84 Abbasi, S. *et al.* (2017) CdSe and CdSe/CdS core-shell QDs: New approach for
650 synthesis, investigating optical properties and application in pollutant degradation.
651 *Luminescence* 32, 1137–1144

652 85 Spirin, M.G. *et al.* (2013) Hydrophilization of CdSe quantum dots with surfactants.
653 *Colloid J* 75, 427–432

654 86 Bobrovsky, A. *et al.* (2016) Quantum dot–polymer composites based on nanoporous
655 polypropylene films with different draw ratios. *Eur Polym J* 82, 93–101

656 87 Thovhogi, N. *et al.* 27-Apr-(2018) Peptide-functionalized quantum dots for potential
657 applications in the imaging and treatment of obesity. *Int J Nanomedicine* 13, 2551-2559

658 88 Brazhnik, K. *et al.* (2014) Advanced Procedure for Oriented Conjugation of Full-Size
659 Antibodies with Quantum Dots. In *Quantum Dots: Applications in Biology* pp. 55–66,
660 Humana Press, New York, NY

661 89 Mansur, H.S. and Mansur, A.A.P. (2011) CdSe quantum dots stabilized by carboxylic-
662 functionalized PVA: Synthesis and UV–vis spectroscopy characterization. *Mater Chem Phys*
663 125, 709–717

664 90 Alemu, Y.A. *et al.* (2018) Enhanced photoluminescence from CuInS₂/ZnS quantum
665 dots: Organic superacid passivation. *Mater. Lett.* 219, 178–181

666 91 Ahia, C.C. *et al.* (2018) Photoluminescence and structural properties of unintentional
667 single and double InGaSb/GaSb quantum wells grown by MOVPE. *Physica B: Condensed*
668 *Matter* 535, 13–19

669 92 Grabolle, M. *et al.* (2009) Determination of the Fluorescence Quantum Yield of
670 Quantum Dots: Suitable Procedures and Achievable Uncertainties. *Anal. Chem.* 81, 6285–
671 6294

672 93 Borah, P. *et al.* (2018) Quantum confinement induced shift in energy band edges and
673 band gap of a spherical quantum dot. *Physica B: Condensed Matter* 530, 208–214

674 94 Marotti, R.E. *et al.* (2006) Crystallite size dependence of band gap energy for
675 electrodeposited ZnO grown at different temperatures. *Sol. Energy Mater Sol.* 90, 2356–
676 2361

677 95 Sinclair, R. *et al.* (2014) The Stanford Nanocharacterization Laboratory (SNL) and
678 Recent Applications of an Aberration-Corrected Environmental Transmission Electron
679 Microscope. *Adv Eng Mater* 16, 476–481

680 96 Ramírez-García, G. *et al.* (2015) Functionalization and characterization of persistent
681 luminescence nanoparticles by dynamic light scattering, laser Doppler and capillary
682 electrophoresis. *Colloids Surf B Biointerfaces* 136, 272–281

- 97 Ramírez-García, G. *et al.* (2017) Characterization of phthalocyanine functionalized
quantum dots by dynamic light scattering, laser Doppler, and capillary electrophoresis. *Anal*
Bioanal Chem 409, 1707–1715
- 98 Henglein, A. (1989) Small-particle research: physicochemical properties of extremely
small colloidal metal and semiconductor particles. *Chem. Rev.* 89, 1861–1873
- 99 Shim, M. *et al.* (2000) Long-Lived Delocalized Electron States in Quantum Dots: A
Step-Scan Fourier Transform Infrared Study. *J. Phys. Chem. B* 104, 1494–1496
- 100 Bonu, V. and Das, A. (2013) Size Distribution of SnO₂ Quantum Dots Studied by UV–Visible, Transmission Electron Microscopy and X-Ray
Diffraction. *Mapan* 28, 259–262
- 101 Choi, Y.J. *et al.* (2012) Cyto-/genotoxic effect of CdSe/ZnS quantum dots in human
lung adenocarcinoma cells for potential photodynamic UV therapy applications. *J Nanosci*
Nanotechnol 12, 2160–2168
- 102 Takahashi, C. *et al.* (2017) Imaging of intracellular behavior of polymeric nanoparticles
in Staphylococcus epidermidis biofilms by slit-scanning confocal Raman microscopy and
scanning electron microscopy with energy-dispersive X-ray spectroscopy. *Mater Sci Eng C*
76, 1066–1074
- 103 Cepeda-Pérez, E. *et al.* (2016) Interaction of TGA@CdTe Quantum Dots with an
Extracellular Matrix of Haematococcus pluvialis Microalgae Detected Using Surface-
Enhanced Raman Spectroscopy (SERS). *Appl Spectrosc* 70, 1561–1572
- 104 Kostić, R. *et al.* (2017) Off-Resonant Raman Spectroscopy of ZnS Quantum Dots. In
Proceedings of the IV Advanced Ceramics and Applications Conference pp. 203–215,
Atlantis Press, Paris
- 105 Biermann, A. *et al.* (2017) Interface formation during silica encapsulation of colloidal
CdSe/CdS quantum dots observed by in situ Raman spectroscopy. *J Chem Phys* 146, 134708
- 106 Pitkänen, L. and Striegel, A.M. (2016) Size-exclusion chromatography of metal
nanoparticles and quantum dots. *Trends Analyt Chem* 80, 311–320
- 107 Hao, C.L. (2012) , In Vivo Stability and Biodistribution of Quantum Dots by Inductively
Coupled Plasma-Atomic Emission Spectrometry. , in *Advanced Materials Research*, 412,
pp. 449–452
- 108 Gondim, C.S. *et al.* (2018) Development and Validation of an Electrochemical
Screening Methodology for Sulfonamide Residue Control in Milk Samples Using a
Graphene Quantum Dots@Nafion Modified Glassy Carbon Electrode. *Food Anal. Methods*
11, 1711–1721

717 109 Bilal, M. et al. (2019) Bayesian Network Resource for Meta-Analysis: Cellular
718 Toxicity of Quantum Dots. Small , 1900510.
719
720
721

722 Figure legends

723

724 Figure 1: Schematic illustration of the structure-activity relationship of QDs. Since the
725 inorganic core is biologically inactive, a hydrophilization step is necessary for its use in
726 biological applications. In order to minimize core-induced toxicity, surface modification
727 strategies may be envisaged (i.e., a capping, a polymerization and a functionalization).

728

729 Figure 2: Impact of QDs toxicity including biological and environmental considerations and
730 reglementary requirements for QD use in human.

731

732

733 Table 1: Biological effects depending on the structure of QDs and the type of stem cells or differentiated cells.

Core material	Shell material	Functionalization/Encapsulation	Type of cell	Outcome	Reference
<i>Stem cells</i>					
CdTe-QDs	-	-	Epithelial	Toxicity	[6]
	-	-	Synoviocytes	Monitoring, toxicity	[18]
CdSe-QDs	-	-	Embryonic	Concentration dependent toxicity	[19]
	ZnS	-	Epithelial	Toxicity	[6]
		-	Embryonic	↘ toxicity	[19]
		-	Neuronal	Slight toxicity	[11]
		Anti-myc antibody	Adipocytes	Monitoring	[15]
		Anti-mortalin antibody	Chondrocytes	Monitoring	[16]
		Tat Peptide	Hematopoietic	Monitoring	[2]
	CdS/ZnS	-	Epithelial	± toxicity	[7,8]
	ZnS - polymer	-	Embryonic	↘ toxicity	[19]
		-	Neuronal	No toxicity under threshold detection	[10,11]
		SiSOX9	Chondrocytes	Monitoring	[17]
	-incorporated glass	-	Epithelial	No toxicity under threshold detection	[9]
Graphene-QDs	-	-	Neuronal	No toxicity under threshold detection	[10]
<i>Differentiated cells</i>					
CdTe-QDs	-	-	Endothelial	Toxicity	[21]
	-	-	Platelets	Toxicity	[4]
	-	-	Renal	Toxicity	[28]
	-	L-cysteine	Renal	Toxicity	[27]
CdSe-QDs	ZnS	-	Monocytes, neutrophils, granulocytes	Toxicity	[3]
		-	Lymphocytes	No toxicity under threshold detection	[3]
		-	Hepatocytes	Toxicity	[20]
		-	Urothelial	No toxicity under threshold detection	[29]
		Photoexcited	Multidrug-resistant bacteria	Bactericidal	[31]
		Anti-Ri7 antibody	Endothelial	Built-up	[25]
		Anti-p38 antibody infiltrated in porous silicon pores immobilized Egp38 antigen	Parasites	Detection	[30]
	ZnS - PEG	-	Endothelial	Built-up	[24]
Graphene-QDs	-	-	Hepatocytes	No toxicity under threshold detection	[22]

734 Table 2: Biological effects depending on the structure of QDs and the organs.

Core material	Shell material	Functionalization	Type of organ	Outcome	Reference
CdTe-QDs	-	-	kidney	Toxicity	[48,49]
	-	-	Intestine	Toxicity	[40]
	CdS	-	Liver	Monitoring	[39]
CdO-QDs	-	-	Lung	Toxicity	[44]
CdSe-QDs	ZnS	-	Brain	No toxicity under threshold detection, visualization	[36]
		-	Heart	Monitoring	[51]
		-	Bladder	No toxicity under threshold detection, no diffusion	[46]
		-	Lung	Toxicity	[45]
		-	Joint	No toxicity under threshold detection	[14]
		GSH	Vascular	Monitoring	[52]
		VEGFR2	Tumor	Detection	[42,43]
	ZnS - polymer	Biotinylated phalloidin	Muscle	Monitoring	[51,53]
		Anti-CD47 antibody	Bladder	No toxicity under threshold detection, no diffusion	[50]
		-	Skin	No penetration	[32-34]
		-	Lung	Toxicity	[41]
		-	Spleen	No toxicity under threshold detection	[41]
		Folic acid	Spleen	Built-up	[41]
Graphene-QDs	-	-	Skin	Therapy	[35]
	-	-	Brain	Therapy	[37-38]

735

736

737 Table 3: Biological effects depending on the structure of QDs and the animal species.

Core material	Shell material	Functionalization /Encapsulation	Animal species	Outcome	Reference
CdTe-QDs	-	-	Mice	Toxicity	[60]
	-	-	Pregnant rat	Dose-dependent embryotoxicity	[74-75]
	-	Quercetin complex	Rat	Therapy	[69]
CdSe-QDs	ZnS	-	Mice	Half-life = 12 – 14h	[64]
	ZnS or CdS	-	Drosophila	Toxicity	[59]
		-	Mice	Dose-dependent toxicity	[61]
		-	Rat	Toxicity	[65]
		-	Male reproductive system in mice	Dose-dependent toxicity	[71-73]
		(His6)-Cyclin E protein	Aquatic species	Monitoring	[58]
	ZnS - polymer	-	Aquatic species	Toxicity	[54-57]
		-	Goat	Monitoring	[76]
	PbS/ZnS	Phospholipid micelle-encapsulated	Mice	No toxicity under threshold detection	[68]
		-	Mice	No toxicity under threshold detection	[62;63]
		Phospholipid micelle-encapsulated	Primate	No toxicity under threshold detection	[77]
			Pregnant primate	Toxicity	[79]
InP-QDs	ZnS	-	Drosophila	No toxicity under threshold detection	[59]
Graphene-QDs	-	-	Rat	Therapy	[70]
Silicon-QDs	- polymer	-	Primate	No toxicity under threshold detection	[78]

738

739

Hydrophilization


↗ water solubility
Biologically compatible



Aqueous coating




Inorganic core




Organic ligands

Biologically incompatible

Inorganic shell



Capping




↗ Photostability
↗ Brightness
↘ Photo-oxidation
↗ Solubility
↗ durability of core

Functionalization

Targeted therapy
Biomarker
Detection & Specific diagnosis
Molecular imaging

Polymerization

Biocompatibility
Stability
Amphiphilic
Facilitated conjugaison



Antibody

Receptors

Biotinylated peptides

Drugs


Biological considerations

Cytotoxicity

Absorption, metabolism, distribution or elimination

Inflammation

Quantum dots



Pharmaceutical grade raw material (permitted heavy metal threshold excess)

Lack of regulation for the pharmaceutical development of specific nanoparticles

Absence of clinical trials in humans

Optical, sizes and analytics differences between batches

Regulatory requirements

Environmental considerations

Non-ecological QD synthesis reagents

Waste management of QDs