



Treatment of calcinosis cutis in systemic sclerosis and dermatomyositis: A review of the literature

Hélène Traineau, Rohit Aggarwal, Jean-Benoît Monfort, Patricia Senet, Chester V. Oddis, Carlo Chizzolini, Annick Barbaud, Camille Francès, Laurent Arnaud, François Chasset

► To cite this version:

Hélène Traineau, Rohit Aggarwal, Jean-Benoît Monfort, Patricia Senet, Chester V. Oddis, et al.. Treatment of calcinosis cutis in systemic sclerosis and dermatomyositis: A review of the literature. Journal of The American Academy of Dermatology, 2020, 82 (2), pp.317 - 325. 10.1016/j.jaad.2019.07.006 . hal-03489704

HAL Id: hal-03489704

<https://hal.science/hal-03489704>

Submitted on 21 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Treatment of calcinosis cutis in systemic sclerosis and dermatomyositis: a review of the literature

Hélène Traineau¹, MD, Rohit Aggarwal², MD, Jean-Benoît Monfort¹, MD, Patricia Senet¹, MD, Chester V. Oddis², MD, Carlo Chizzolini^{3, 4}, MD, PhD, Annick Barbaud¹ MD, PhD, Camille Francès¹ MD, Laurent Arnaud⁵, MD, PhD, François Chasset¹ MD

¹Sorbonne Université, Faculté de Médecine Sorbonne Université, AP-HP, Service de Dermatologie et Allergologie, Hôpital Tenon, F-75020 Paris, France

² Division of Rheumatology and Clinical Immunology, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA, USA

³ Department of Immunology & Allergy, University Hospital and School of Medicine, Geneva, Switzerland

⁴ Department of Pathology & Immunology, School of Medicine, Geneva, Switzerland.

⁵ Service de rhumatologie, Hôpitaux Universitaires de Strasbourg, Laboratoire d'ImmunoRhumatologie Moléculaire, Centre National de Référence des Maladies Systémiques Auto-immunes Rares Est Sud-Ouest (RESO), INSERM UMR_S1109, Université de Strasbourg, F-67000 Strasbourg, France

Corresponding author & reprint requests:

François Chasset, MD, Sorbonne université, AP-HP, Service de Dermatologie et d'Allergologie, Hôpital Tenon, 4 rue de la Chine 75970 Paris CEDEX 20, France

Phone number: (+33156 01 75 47). Fax number: (+331 56 01 72 32)

Email: francois.chasset@aphp.fr

Funding sources: none

Conflict of interest: Laurent ARNAUD has received honoraria from Roche-Chugai, Grifols, LFB, Pfizer, UCB, Carlo CHIZZOLINI has received travel support from Roche-Chugai

IRB status: not necessary

Statement of any prior presentation: none

Abstract words count: 199, **Capsule words count:** 53, **Text word count:** 2386, **Figures count:** 1, **Table count:** 4

Abstract:

Background: We have limited data on the treatment of calcinosis cutis associated with systemic sclerosis (SSc) and dermatomyositis (DM).

Objective: To assess the efficacy and tolerance of available treatments for calcinosis cutis based on previously published studies.

Method: We performed a systematic review of studies published in MEDLINE, Embase, and the Cochrane library between 1980 and July 2018. The strength of clinical data was graded according to the modified Oxford Centre for Evidence-Based Medicine Levels of Evidence.

Results: In all, 30 studies (288 patients) were included. Eleven therapeutic classes, surgery and physical treatments were identified as potential treatments for calcinosis cutis. From the results of a small randomized controlled trial and 4 retrospective studies, low-dose warfarin should not be used for calcinosis cutis (Level IB evidence). Several studies suggested the use of diltiazem and bisphosphonates (Level IV). Considering biologic therapies, rituximab has shown interesting results in both DM and SSc, whereas TNF inhibitors may be useful in juvenile DM (Level IV). Intralesional sodium thiosulfate may be a promising alternative (Level IV).

Limitations: Few included studies had a high level of evidence.

Conclusion: This study highlights the efficacy and tolerance profiles of available treatments for calcinosis cutis, with a focus on level of evidence.

Key words: calcinosis cutis, systemic sclerosis, dermatomyositis, level of evidence

INTRODUCTION

Calcinosis cutis is defined by the deposition of insoluble calcium in the skin and subcutaneous tissues.¹ DM and SSc are the most frequent autoimmune connective tissue disorders associated with calcinosis cutis.² Indeed, calcinosis cutis develops in about 30% of adult DM patients³ and 30% to 70% of juvenile DM patients.⁴⁻⁷ Moreover, the prevalence of calcinosis ranges from 18% to 49% in SSc patients.⁸⁻¹² Significant advances in understanding the SSc and DM pathogenesis, newer classification criteria and advances in disease management have resulted in improved survival in SSc and DM.^{4,5,13} Nevertheless, long-term morbidity remains a major issue.^{14,15}

Dystrophic calcinosis is associated with considerably impaired quality of life due to ulceration and secondary infections, both resulting in extreme debilitation.^{7,16} Although early aggressive intervention may prevent calcinosis cutis development,¹⁷⁻¹⁹ treatment remains challenging. Furthermore, few randomized controlled trials (RCTs) have been performed²⁰ and we lack specific guidelines for calcinosis cutis management in DM or SSc. However, several case series or prospective cohort studies focusing on calcinosis cutis treatment have been published.²¹

To better define an evidence-based treatment approach and to provide the best available evidence for physicians, we performed a systematic review of case series and cohort studies investigating the management of calcinosis cutis in patients with DM or SSc.

MATERIAL AND METHODS

This systematic review was performed according to the 2009 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist.²²

Literature search and information sources:

We performed a systematic review of MEDLINE/PubMed, Embase and the Cochrane database between January 1980 to July 2018, with no restriction on language. The search strategy combined free text search, exploded MESH/EMTREE terms and all synonyms of the following Medical Subject Headings terms: systemic sclerosis, dermatomyositis, calcinosis cutis. The grey literature was also explored to avoid publication bias. We also searched for additional articles from the reference lists of relevant papers.

Study selection and eligibility criteria

Observational studies or RCTs were considered if 1) they included patients with DM or SSc; 2) the number of patients with calcinosis cutis was available; 3) patients received a specific treatment for calcinosis cutis or for the connective tissue disease with a specific assessment of calcinosis cutis outcomes; and 4) the number of patients treated and the number of responders were available. Given the rarity of calcinosis cutis but to avoid publication bias based on single case reports, we excluded case series of fewer than 3 patients along with reviews, editorials and guidelines (**Figure 1**). The quality of studies was assessed by the Newcastle-Ottawa Assessment Scale²³ for observational studies and the Cochrane collaboration Risk of Bias tool²⁴ for RCTs.

Data extraction and assessment of calcinosis cutis outcomes

The post-treatment calcinosis cutis response was reported as complete or partial. Complete response was defined as the complete disappearance of calcinosis cutis, and partial response was any improvement according to the study protocol, which included reduction in the size of calcinotic deposits and healing of ulcerations. When only pain reduction was reported, the treatment was considered a failure. Adverse events were recorded. The relapse rate was defined as the reappearance of lesions after a complete or partial response. All data were extracted independently by 2 investigators. In the tables, when data were available, adult and pediatric patients are presented separately.

Levels of evidence and treatment recommendation:

The strength of clinical data and subsequent treatment recommendations were graded according to the modified Oxford Centre for Evidence-Based Medicine Levels of Evidence and Grades of Recommendation.²⁵

RESULTS

Literature search and characteristics of included studies

Our literature search identified 3032 citations; reports for 30 studies^{10,20,26–53} were included in this systematic review (**Figure 1**), including 2 RCTs,^{20,30} 11 prospective cohort studies^{31,32,36,43–48,51,53} and 17 retrospective studies.^{10,26–29,33–35,37–42,49,50,52} The sample size ranged from 3 to 78 patients, for a total of 288 patients (SSc=108, adult DM=66, juvenile DM=90, and DM or SSc but unspecified diagnosis=24).

Overall, the methodological quality of included studies was low, with 26 cohort studies graded as poor quality and only 2 as fair quality.

Warfarin: Warfarin, a vitamin-K antagonist, was used in one RCT²⁰ and 4 retrospective studies^{26–29}, with a total of 19 patients (DM=10, SSc=6, unspecified=3) (**Table 1**). A dosage of 1 mg/day was most commonly prescribed. The mean calcinosis cutis duration was from 4 months²⁷ to 10 years.²⁶ A small placebo-controlled trial²⁶ found no clinical improvement in calcinosis cutis in 5 patients who received warfarin 1 mg/day. From the results of 5 studies, the partial response rate ranged from 0%^{20,26,29} to 2/3 (66%)^{28,54}, with no complete response observed in all but one study (2/3; 66%).⁵⁴ No relapse was observed after 2 years' follow-up in one study.²⁷ Adverse events were not reported.

Diltiazem: Diltiazem, a calcium channel blocker, was used in 3 retrospective cohort studies in 38 patients (DM=12, SSc=12, unspecified =14)^{10,28,29} (**Table 1**). The dosage ranged from 60 mg 3 times daily to 480 mg/day. The partial response rate ranged from 0/12 (0%)²⁹ to 9/14 (64%).²⁸ No complete responses were reported in the 3 studies and no adverse events were reported in the study of Vayssairat et al.¹⁰

Rituximab: Rituximab was used in 7 studies (SSc=18, adult DM=9, juvenile DM=32) including 3 prospective studies,^{31–33} 3 retrospective studies^{29,34,35} and one RCT versus placebo

focusing on the efficacy of rituximab on skin lesions including calcinosis cutis.³⁰ Study characteristics are summarized in **Table 2**. The mean calcinosis cutis duration ranged from 3.4 to 12 years.^{33,34} From the results of 6 studies, the partial response rate ranged from 0/6 (0%)³⁴ to 100% in 3 studies including 3 to 9 patients.^{31,32,35} The complete response rate ranged from 0%^{29,33,34} to 100% in a study of 3 SSc patients.³¹ In 2 studies, the relapse rate was 0% with a follow-up of 12 to 60 months.^{31,35} Adverse events were assessed in 3 studies and included localized bacterial infection of the calcinosis (n=2), moderate acute infusion-related events (n=1), and intestinal perforation 2 weeks after a combination of rituximab and pulse methylprednisolone infusions (n=1).

Other biologic agents: Tumor necrosis factor α (TNF- α) inhibitors and abatacept (CTLA4-Ig) were used in 4 studies^{29,36–38}, with a total of 30 DM patients (mostly juvenile DM). The results of 3 infliximab studies^{29,36,38} showed a partial response rate ranging from 0% (0/2)²⁹ to 80% in a prospective study of 5 juvenile DM patients³⁶. No complete responses were noted and adverse events were not reported.

Bisphosphonates: Four retrospective cohort studies of 17 individuals (DM=15, unspecified =2)^{28,39–41} assessed the efficacy of bisphosphonates for calcinosis cutis (**Table 3**). Specific bisphosphonates and the therapeutic regimens were heterogeneous. The partial response ranged from 3/6 (50%)⁴⁰ to 3/3 (100%)³⁹ and the complete response rate from 0% to 33%.^{28,39–41} In 2 patients receiving pamidronate with complete response, one relapse was noted after 4 years; in this case, alendronate was then used, which resulted in a second complete response. No adverse events were recorded in the only study assessing safety.⁴¹

Intravenous immunoglobulins: Two retrospective studies including 15 DM patients^{29,42} assessed the efficacy of intravenous immunoglobulins on calcinosis cutis. Galimberti et al.⁴² reported a partial response in 5/8 (62%), with no complete responders. Conversely, no

objective response was observed in the study of Fredi et al.²⁹ including 7 DM patients, with only one reporting pain improvement. Adverse events were not reported in these studies.

Sodium thiosulfate (STS): Five studies^{29,48–51} assessing different regimens of STS or its metabolites involved 20 patients (SSc=9, adult DM=10, juvenile DM=1). Two studies^{48,49} of 7 patients assessed the efficacy of intravenous STS. No objective improvement was found, and pain improvement was noted in only one patient.⁴⁹ Topical STS conferred no improvement in 5 DM patients.²⁹ Partial response was achieved with topical sodium metabisulfite (a metabolite of STS) in 3/3 patients (2 DM and 1 SSc), without relapse,⁵⁰ including one with complete response. Finally, in a prospective study, partial response was achieved in 5/5 SSc patients, including complete response in 2 receiving intralesional STS injection.⁵¹ Adverse events of intralesional STS included transient pain (n=2/5) and local infection (n=1/5).⁵¹

Minocycline: Minocycline (50-200 mg/day) was used in 2 studies of 12 patients (SSc=9, unspecified=3).^{28,52} Robertson et al.⁵² reported a partial response in 8/9 patients and Balin et al.²⁸ a partial response in 1/3. Adverse events included nausea (n=1), dizziness (n=1) and the conversion of calcinosis cutis deposits to a blue/black color.⁵²

Colchicine: Balin et al.²⁸ found a partial response in 3/7 patients (43%), including 1/7 with complete response using colchicine doses < 1.2 mg/day. Fredi et al.²⁹ reported only 1/9 partial response with colchicine.

Cyclophosphamide: One prospective study used cyclophosphamide for treating refractory or severe juvenile DM, including 14 patients with calcinosis cutis.⁵³ Complete response of calcinosis cutis was noted in 9/14 (64%), with a follow-up between 12 and 24 months.

Surgery and physical therapies: Five prospective studies and one retrospective study (n=55 patients: SSc=26, DM=1, unspecified=28) assessed surgical intervention and physical therapy

for calcinosis cutis improvement^{28,43-47} (**Table 3**). Surgical excision in 2 studies led to 80% improvement.⁴⁷ Balin et al.²⁸ reported a partial response of 27/28 (96%), including 22/28 (79%) with complete response. Two small prospective cohort studies of 7 patients noted a partial response rate of 33% to 100% with extracorporeal shock-wave therapy, with no complete response.^{44,45} The relapse rate was not reported. Adverse events included transient pain with extrusion of calcific debris. Another prospective study of 6 SSc patients reported a partial improvement rate of 83% with a carbon dioxide laser.⁴³ Adverse events included poor wound healing (n=5), hyperkeratosis (n=4) and postoperative infections (n=2). Relapse was observed in 2/6 patients within 3 to 4 months.⁴³

DISCUSSION

In this systematic review, we identified 30 studies (288 patients) focusing on the treatment of calcinosis cutis associated with SSc and DM. **Table 4** summarizes the available treatments for calcinosis cutis with a focus on underlying diseases and levels of evidence.

Currently, we lack specific guidelines for managing calcinosis cutis in autoimmune connective tissue disorders. In the recent consensus-based recommendations for the management juvenile DM, an intensification of immunosuppressive therapy was suggested, but no specific treatment was recommended.¹⁹ Moreover, treatment of calcinosis cutis is not included in the updated EULAR recommendations for treating SSc.⁵⁵

From our systematic review, several drugs used to treat calcinosis cutis have potential therapeutic interest. Because of the small number of patients and the treatment heterogeneity among included studies, pooled response rates were not calculated. However, several important findings may be underlined. From the results of a small RCT, warfarin conferred no improvement in calcinosis cutis⁵⁶, and no partial response was observed in most included studies.^{20,26,29} Therefore, warfarin should not be considered for treating calcinosis cutis. This suggestion is further supported by the fact that warfarin could promote ectopic calcification via under-carboxylated matrix gla protein.⁵⁷

Despite no complete response noted in the 3 studies of diltiazem, partial response was observed in some, particularly in Balin et al., in 9/14 (64%) patients. Some data support the use of calcium channel blockers for calcinosis cutis in SSc. Indeed, digital ischemia was strongly related to the occurrence of calcinosis cutis in a large study of 1300 SSc patients, and the use of calcium channel blockers was inversely associated with the presence of calcinosis cutis in this study.¹² Moreover, the use of calcium channel blockers is recommended for treating Raynaud phenomenon in SSc.⁵⁵ Therefore, although no formal

curative effect of diltiazem on calcinosis cutis could be demonstrated, a potential preventive effect cannot be ruled out, and therefore diltiazem may be considered for treating digital calcinosis cutis associated with SSc.

Rituximab has been increasingly used in DM and SSc because of favorable outcomes in DM skin lesions³⁰ and SSc skin sclerosis and lung function.⁵⁸ From the results of 6 studies, rituximab may be considered for treating calcinosis cutis both in DM and SSc. Indeed, although one study of 6 juvenile DM patients did not reported improvement,³⁴ most studies showed at least partial response, including 3 with 100% partial response.^{31,32,35} Moreover, 3 studies reported at least one patient with complete response.^{30,31,35} TNF- α inhibitors, particularly infliximab, may have a beneficial effect on calcinosis cutis in juvenile DM, but their use should be carefully scrutinized in SSc because of reports of severe exacerbation of pulmonary fibrosis associated with their use.⁵⁹

Bisphosphonates remain a therapeutic option mostly in DM, with at least partial response noted in 4 studies.^{28,39-41} Nevertheless, a lack of substantive data precludes recommending a specific regimen of bisphosphonates.

Intravenous sodium thiosulfate seemed ineffective,⁴⁹ but intralesional treatment could be a promising alternative.⁵¹

Several other treatments, such as intravenous immunoglobulin,⁴² minocycline,^{28,52} colchicine,^{28,29} and cyclophosphamide⁵³, improved calcinosis cutis in small case series but with limited level of evidence.

Surgery and physical therapies should be considered in calcinosis cutis, both in DM and SSc: several studies^{28,47} reported response rates higher than 80%. However, the surgical management of digital calcinosis cutis in SSc may lead to skin necrosis and limited range of

249 motion.⁶⁰ Less invasive procedures such as carbon dioxide laser⁴³ or extracorporeal shock-
250 wave therapy^{44,45} may be useful, but the level of evidence is weak.

251 Among the limitations of this systematic review is a possible publication bias. Given the
252 rarity of publications related to calcinosis cutis treatment, we included case series with at
253 least 3 patients which may have affected our results. However, single case reports were
254 excluded, as were case series of only 2 patients, in order to reduce the bias, as previously
255 described in systematic reviews of rare diseases.⁶¹ Moreover, increasing the minimum
256 number of patients to be considered for inclusion would have led to the exclusion of several
257 treatments with high potential interest. Another limitation is the low levels of evidence of the
258 reviewed studies. To date, only 2 small RCTs assessing improvement of calcinosis cutis have
259 been performed,^{20,30} including one with calcinosis cutis as the secondary outcome,³⁰ Other
260 studies were mainly low-quality cohort studies or case series with Level IV
261 recommendations.²⁵ The severity and the size and duration of calcinosis cutis disease are
262 important confounding factors that may affect the therapeutic response. Only calcinosis cutis
263 duration was reported in some studies, and data were inadequate to perform any subgroup
264 analyses.

265 The treatment of calcinosis cutis is a major unmet need. Identifying patients at high risk of
266 developing calcinosis cutis and its early treatment is recommended.¹⁹ On the basis of 30
267 studies including 288 SSc and DM patients, this systematic review provides evidence-based
268 guidance for practitioners.

269 **Acknowledgements:** We deeply thank Laura Smales (BioMedEditing) for English-language
270 editing of the manuscript.

271 **Conflict of interests:** Laurent ARNAUD has received honoraria from Roche-Chugai,
272 Grifols, LFB, Pfizer, UCB; Carlo CHIZZOLINI has received travel support from Roche.

273 **Abbreviation and acronym list:**

274 Dermatomyositis: DM

275 Systemic sclerosis: SSc

276 Randomized controlled trial: RCT

277 Tumor necrosis factor: TNF

278 Sodium thiosulfate: STS

279

References

1. Chander S, Gordon P. Soft tissue and subcutaneous calcification in connective tissue diseases. *Curr Opin Rheumatol*. 2012;24(2):158-164. doi:10.1097/BOR.0b013e32834ff5cd.
2. Walsh JS, Fairley JA. Calcifying disorders of the skin. *J Am Acad Dermatol*. 1995;33(5 Pt 1):693-706; quiz 707-710.
3. Sallum AME, Pivato FCMM, Doria-Filho U, et al. Risk factors associated with calcinosis of juvenile dermatomyositis. *J Pediatr (Rio J)*. 2007;0(0). doi:10.2223/JPED.1746
4. Ramanan AV, Feldman BM. Clinical features and outcomes of juvenile dermatomyositis and other childhood onset myositis syndromes. *Rheum Dis Clin North Am*. 2002;28(4):833-857.
5. McCann LJ, Juggins AD, Maillard SM, et al. The Juvenile Dermatomyositis National Registry and Repository (UK and Ireland)--clinical characteristics of children recruited within the first 5 yr. *Rheumatology*. 2006;45(10):1255-1260. doi:10.1093/rheumatology/kei099
6. Clemente G, Piotto DGP, Barbosa C, et al. High frequency of calcinosis in juvenile dermatomyositis: a risk factor study. *Rev Bras Reumatol*. 2012;52(4):549-553.
7. Robinson AB, Hoeltzel MF, Wahezi DM, et al. Clinical Characteristics of Children With Juvenile Dermatomyositis: The Childhood Arthritis and Rheumatology Research Alliance Registry: Investigating Juvenile DM Through a National Multicenter Registry. *Arthritis Care Res*. 2014;66(3):404-410. doi:10.1002/acr.22142
8. Morgan ND, Shah AA, Mayes MD, et al. Clinical and serological features of systemic sclerosis in a multicenter African American cohort: Analysis of the genome research in African American scleroderma patients clinical database. *Medicine (Baltimore)*. 2017;96(51):e8980. doi:10.1097/MD.00000000000008980
9. Cruz-Domínguez MP, García-Collinot G, Saavedra MA, et al. Clinical, biochemical, and radiological characterization of the calcinosis in a cohort of Mexican patients with systemic sclerosis. *Clin Rheumatol*. 2017;36(1):111-117. doi:10.1007/s10067-016-3412-9
10. Vayssairat M, Hidouche D, Abdoucheli-Baudot N, Gaitz JP. Clinical significance of subcutaneous calcinosis in patients with systemic sclerosis. Does diltiazem induce its regression? *Ann Rheum Dis*. 1998;57(4):252-254.
11. Belloli L, Ughi N, Massarotti M, Marasini B, Biondi ML, Brambilla G. Role of Fetuin-A in Systemic Sclerosis-associated Calcinosis. *J Rheumatol*. 2010;37(12):2638-2639. doi:10.3899/jrheum.100627
12. Baron M, Pope J, Robinson D, et al. Calcinosis is associated with digital ischaemia in systemic sclerosis—a longitudinal study. *Rheumatology*. 2016;55(12):2148-2155. doi:10.1093/rheumatology/kew313
13. Allanore Y, Avouac J, Kahan A. Systemic sclerosis: an update in 2008. *Joint Bone Spine*. 2008;75(6):650-655. doi:10.1016/j.jbspin.2008.07.003
14. Marie I. Morbidity and Mortality in Adult Polymyositis and Dermatomyositis. *Curr Rheumatol Rep*. 2012;14(3):275-285. doi:10.1007/s11926-012-0249-3

- 318 15. Panopoulos S, Bournia V-K, Konstantonis G, Fragiadaki K, Sfrikakis PP, Tektonidou MG.
319 Predictors of morbidity and mortality in early systemic sclerosis: Long-term follow-up data
320 from a single-centre inception cohort. *Autoimmun Rev.* 2018;17(8):816-820.
321 doi:10.1016/j.autrev.2018.02.008
- 322 16. Wu JJ, Metz BJ. Calcinosis Cutis of Juvenile Dermatomyositis Treated with Incision and
323 Drainage. *Dermatol Surg.* 2008;34(4):575-577. doi:10.1111/j.1524-4725.2007.34106.x
- 324 17. Ravelli A, Trail L, Ferrari C, et al. Long-term outcome and prognostic factors of juvenile
325 dermatomyositis: A multinational, multicenter study of 490 patients. *Arthritis Care Res.*
326 2010;62(1):63-72. doi:10.1002/acr.20015
- 327 18. Kim S, El-Hallak M, Dedeoglu F, Zurakowski D, Fuhlbrigge RC, Sundel RP. Complete and
328 sustained remission of juvenile dermatomyositis resulting from aggressive treatment. *Arthritis*
329 *Rheum.* 2009;60(6):1825-1830. doi:10.1002/art.24571
- 330 19. Enders FB, Bader-Meunier B, Baildam E, et al. Consensus-based recommendations for the
331 management of juvenile dermatomyositis. *Ann Rheum Dis.* 2017;76(2):329-340.
332 doi:10.1136/annrheumdis-2016-209247
- 333 20. Berger RG, Featherstone GL, Raasch RH, McCartney WH, Hadler NM. Treatment of calcinosis
334 universalis with low-dose warfarin. *Am J Med.* 1987;83(1):72-76.
- 335 21. Valenzuela A, Song P, Chung L. Calcinosis in scleroderma. *Curr Opin Rheumatol.*
336 2018;30(6):554-561. doi:10.1097/BOR.0000000000000539
- 337 22. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA Statement for Reporting Systematic
338 Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions: Explanation
339 and Elaboration. *PLoS Med.* 2009;6(7):e1000100. doi:10.1371/journal.pmed.1000100
- 340 23. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of
341 nonrandomized studies in meta-analyses. *Eur J Epidemiol.* 2010;25(9):603-605.
342 doi:10.1007/s10654-010-9491-z
- 343 24.
344 [Http://Handbook.Cochrane.Org/Chapter_8/8_5_the_cochrane_collaborations_tool_for_asse](http://Handbook.Cochrane.Org/Chapter_8/8_5_the_cochrane_collaborations_tool_for_assessing_risk_of_bias.Htm)
345 [ssing_risk_of_bias.Htm](http://Handbook.Cochrane.Org/Chapter_8/8_5_the_cochrane_collaborations_tool_for_assessing_risk_of_bias.Htm).
- 346 25. Centre for Evidence-based Medicine website. Levels of evidence (March 2009). Available at:
347 <http://www.cebm.net/oxford-centre-evidence-based-medicine-levels-evidence-march-2009/>.
- 348 26. Lassoued K, Saiag P, Anglade MC, Roujeau JC, Touraine RL. Failure of warfarin in treatment of
349 calcinosis universalis. *Am J Med.* 1988;84(4):795-796.
- 350 27. Cukierman T. Low dose warfarin treatment for calcinosis in patients with systemic sclerosis.
351 *Ann Rheum Dis.* 2004;63(10):1341-1343. doi:10.1136/ard.2003.014431
- 352 28. Balin SJ, Wetter DA, Andersen LK, Davis MDP. Calcinosis cutis occurring in association with
353 autoimmune connective tissue disease: the Mayo Clinic experience with 78 patients, 1996-
354 2009. *Arch Dermatol.* 2012;148(4):455-462. doi:10.1001/archdermatol.2011.2052

- 355 29. Fredi M, Bartoli F, Cavazzana I, et al. SAT0469 Calcinosis Cutis in Poly-Dermatomyositis: Clinical
356 and Therapeutic Study. *Ann Rheum Dis*. 2015;74(Suppl 2):830.2-831.
357 doi:10.1136/annrheumdis-2015-eular.4163
- 358 30. Aggarwal R, Loganathan P, Koontz D, Qi Z, Reed AM, Oddis CV. Cutaneous improvement in
359 refractory adult and juvenile dermatomyositis after treatment with rituximab. *Rheumatology*.
360 2017;56(2):247-254. doi:10.1093/rheumatology/kew396
- 361 31. Moazedi-Fuerst FC, Kielhauser SM, Bodo K, Graninger WB. Dosage of rituximab in systemic
362 sclerosis: 2-year results of five cases. *Clin Exp Dermatol*. 2015;40(2):211-212.
363 doi:10.1111/ced.12450
- 364 32. Narváez J., Sancho J.J.A., Castellvi I., Herrera S., Molina M.M., Castillo D., De La Morena Barrio
365 I., Villarino M.R., Ferrer A.M., García D.Y., Pascual E.V., Llobet J.M., Latorre F.G., Nolla J.M.
366 Long-term efficacy of rituximab in systemic sclerosis,. *Arthritis and Rheumatology (2014) 66*
367 *SUPPL. 10 (S737)*.
- 368 33. Giuggioli D, Lumetti F, Colaci M, Fallahi P, Antonelli A, Ferri C. Rituximab in the treatment of
369 patients with systemic sclerosis. Our experience and review of the literature. *Autoimmun Rev*.
370 2015;14(11):1072-1078. doi:10.1016/j.autrev.2015.07.008
- 371 34. Bader-Meunier B, Decaluwe H, Barnerias C, et al. Safety and Efficacy of Rituximab in Severe
372 Juvenile Dermatomyositis: Results from 9 Patients from the French Autoimmunity and
373 Rituximab Registry. *J Rheumatol*. 2011;38(7):1436-1440. doi:10.3899/jrheum.101321
- 374 35. Alhemairi M., Muzaffer M. Effectiveness of rituximab therapy on severe calcinosis in 4 children
375 with JDM,Pediatric Rheumatology (2017) 15 Supplement 1.
- 376 36. Riley P, McCann LJ, Maillard SM, Woo P, Murray KJ, Pilkington CA. Effectiveness of infliximab in
377 the treatment of refractory juvenile dermatomyositis with calcinosis. *Rheumatology*.
378 2008;47(6):877-880. doi:10.1093/rheumatology/ken074
- 379 37. DeGuzman M., Singla S., Mizesko M., Sagcal-Gironella A.C. Abatacept as adjunct therapy for
380 the calcinosis of juvenile dermatomyositis: A single-center experience. *Arthritis and*
381 *Rheumatology (2017) 69 Supplement 4 (137-138)*. Date of Publication: 1 Apr 2017.
- 382 38. Boulter E, Beard L, Ryder C, Pilkington C, UK Juvenile Dermatomyositis Research Group (JDRG).
383 Effectiveness of anti-TNF- α agents in the treatment of refractory juvenile dermatomyositis.
384 *Pediatr Rheumatol*. 2011;9(Suppl 1):O29. doi:10.1186/1546-0096-9-S1-O29
- 385 39. Marco Puche A, Calvo Penades I, Lopez Montesinos B. Effectiveness of the treatment with
386 intravenous pamidronate in calcinosis in juvenile dermatomyositis. *Clin Exp Rheumatol*.
387 2010;28(1):135-140.
- 388 40. Tayfur AC, Topaloglu R, Gulhan B, Bilginer Y. Bisphosphonates in juvenile dermatomyositis with
389 dystrophic calcinosis. *Mod Rheumatol*. 2015;25(4):615-620.
390 doi:10.3109/14397595.2014.988197
- 391 41. Saini I, Kalaivani M, Kabra SK. Calcinosis in juvenile dermatomyositis: frequency, risk factors
392 and outcome. *Rheumatol Int*. 2016;36(7):961-965. doi:10.1007/s00296-016-3467-6

- 393 42. Galimberti F, Li Y, Fernandez AP. Intravenous immunoglobulin for treatment of
394 dermatomyositis-associated dystrophic calcinosis. *J Am Acad Dermatol*. 2015;73(1):174-176.
395 doi:10.1016/j.jaad.2015.03.047
- 396 43. Bottomley WW, Goodfield MJ, Sheehan-Dare RA. Digital calcification in systemic sclerosis:
397 effective treatment with good tissue preservation using the carbon dioxide laser. *Br J*
398 *Dermatol*. 1996;135(2):302-304.
- 399 44. Blumhardt S, Frey DP, Toniolo M, Alkadhi H, Held U, Distler O. Safety and efficacy of
400 extracorporeal shock wave therapy (ESWT) in calcinosis cutis associated with systemic
401 sclerosis. *Clin Exp Rheumatol*. 2016;34 Suppl 100(5):177-180.
- 402 45. Sultan-Bichat N, Menard J, Perceau G, Staerman F, Bernard P, Reguia Z. Treatment of calcinosis
403 cutis by extracorporeal shock-wave lithotripsy. *J Am Acad Dermatol*. 2012;66(3):424-429.
404 doi:10.1016/j.jaad.2010.12.035
- 405 46. Shetty S. A pilot study of acetic acid iontophoresis and ultrasound in the treatment of systemic
406 sclerosis-related calcinosis. *Rheumatology*. 2005;44(4):536-538.
407 doi:10.1093/rheumatology/keh536
- 408 47. Fahmy FS, Evans DM, Devaraj VS. Microdrilling of digital calcinosis. *Eur J Plast Surg*.
409 1998;21(7):378-380. doi:10.1007/s002380050122
- 410 48. Trysberg E., Werna S., Sakiniene E. Effect of sodium thiosulfate on calcinosis cutis associated
411 with connective tissue disease,. *Annals of the Rheumatic Diseases (2014) S73*.
- 412 49. Mageau A, Guignon V, Ratzimbasafy V, et al. Intravenous sodium thiosulfate for treating
413 tumoral calcinosis associated with systemic disorders: Report of four cases. *Joint Bone Spine*.
414 2017;84(3):341-344. doi:10.1016/j.jbspin.2016.10.009
- 415 50. del Barrio-Díaz P, Moll-Manzur C, Álvarez-Veliz S, Vera-Kellet C. Topical sodium metabisulfite
416 for the treatment of calcinosis cutis: a promising new therapy. *Br J Dermatol*. 2016;175(3):608-
417 611. doi:10.1111/bjd.14412
- 418 51. Baumgartner-Nielsen J, Olesen A. Treatment of Skin Calcifications with Intra-lesional Injection
419 of Sodium Thiosulphate: A Case Series. *Acta Derm Venereol*. 2016;96(2):257-258.
420 doi:10.2340/00015555-2206
- 421 52. Robertson LP, Marshall RW, Hickling P. Treatment of cutaneous calcinosis in limited systemic
422 sclerosis with minocycline. *Ann Rheum Dis*. 2003;62(3):267-269.
- 423 53. Moraitis E, Arnold K, Wedderburn L, Pilkington C, JDRG. PRoS-FINAL-2130-A: Effectiveness of
424 intravenous cyclophosphamide in severe or refractory juvenile dermatomyositis - a national
425 cohort study UK and Ireland. *Pediatr Rheumatol*. 2013;11(Suppl 2):P143. doi:10.1186/1546-
426 0096-11-S2-P143
- 427 54. Cukierman T, Elinav E, Korem M, Chajek-Shaul T. Low dose warfarin treatment for calcinosis in
428 patients with systemic sclerosis. *Ann Rheum Dis*. 2004;63(10):1341-1343.
429 doi:10.1136/ard.2003.014431
- 430 55. Kowal-Bielecka O, Fransen J, Avouac J, et al. Update of EULAR recommendations for the
431 treatment of systemic sclerosis. *Ann Rheum Dis*. 2017;76(8):1327-1339.
432 doi:10.1136/annrheumdis-2016-209909

- 433 56. Berger RG, Featherstone GL, Raasch RH, McCartney WH, Hadler NM. Treatment of calcinosis
434 universalis with low-dose warfarin. *Am J Med.* 1987;83(1):72-76.
- 435 57. Palaniswamy C, Sekhri A, Aronow WS, et al. Association of warfarin use with valvular and
436 vascular calcification: a review. *Clin Cardiol* 2011; 34:74–81.
- 437 58. Thiebaut M, Launay D, Rivière S, et al. Efficacy and safety of rituximab in systemic sclerosis:
438 French retrospective study and literature review. *Autoimmun Rev.* 2018;17(6):582-587.
439 doi:10.1016/j.autrev.2017.12.010
- 440 59. Allanore Y. Fatal exacerbation of fibrosing alveolitis associated with systemic sclerosis in a
441 patient treated with adalimumab. *Ann Rheum Dis.* 2006;65(6):834-835.
442 doi:10.1136/ard.2005.044453
- 443 60. Bogoch ER, Gross DK. Surgery of the hand in patients with systemic sclerosis: outcomes and
444 considerations. *J Rheumatol.* 2005;32(4):642-648.
- 445 61. Micieli R, Alavi A. Treatment for Livedoid Vasculopathy: A Systematic Review. *JAMA Dermatol.*
446 2018;154(2):193-202. doi:10.1001/jamadermatol.2017.4374

447

448

449 **Figure Legends:**

450 **Figure 1. Flow-charts for study selection.** DM dermatomyositis, SSc: systemic sclerosis

451

452 **Table 1. Characteristics of included studies evaluating the effect of warfarin and diltiazem on calcinosis cutis improvement**

Name/year	Disease (N)	Study design	Calcinosis duration at inclusion, years	Dose regimen	Partial response, N (%)	Complete response, N (%)	Follow-up, mean (range), months	Level of evidence, grade*
Warfarin								
<i>Adults</i>								
Berger, 1987	SSc (2)	RCT	8 (6-10)	1 mg/day	0 (0%)	0 (0%)	18	IB
Lassoued, 1988	SSc (1) DM (5)	R	10 (2-25)	1 mg/day	0 (0%)	0 (0%)	14.6 (7-28)	IV
Cukierman, 2004	SSc (3)	R	0.4	1 mg/day	2 (66%)	2 (66%)	20 (12-24)	IV
Balin, 2012	NA (3)	R	NA	NA	2(66%)	0 (0%)	104 (1-696)	IV
Fredi, 2015	DM (2)	R	NA	NA	0 (0%)	0 (0%)	201.8	IV
<i>Children</i>								
Berger, 1987	DM/SSc (1) DM (2)	RCT	5.3 (3-9)	1 mg/day	0 (0%)	0 (0%)	18	IB
Diltiazem								
<i>Adults</i>								
Vayssairat, 1998	SSc (12)	R	11.5	60mg x3/day	3 (25 %)	0 (0%)	78 (12-180)	IV
Balin, 2012	NA (14)	R	NA	≤480 mg/day	9 (64%)	0 (0%)	104	IV
Fredi, 2015	DM (12)	R	NA	NA	0 (0%)	0 (0%)	201.8	IV

453 RCT: randomized controlled trial, R: retrospective, SSc: systemic sclerosis, DM: dermatomyositis, NA: not available *according to modified Oxford Centre
 454 for Evidence-Based Medicine Levels of Evidence and Grades of Recommendation

455
 456
 457
 458
 459
 460
 461
 462
 463

464 **Table 2. Characteristics of included studies evaluating the effect of rituximab on calcinosis cutis improvement**

Name/year	Disease (N)	Study design	Dose regimen	Partial response, N (%)	Complete response, N (%)	Follow-up, mean (range), months	Level of evidence, grade*
Adult							
Aggarwal, 2016³⁰	DM (7)	RCT	0.575-1g/m2 at weeks 0/1	NA	1 (14%)	1.46	IB
Moazedi-fuerst, 2015³¹	SSc (3)	P	500 mg/m2 at weeks 0/2 then every 3 months	3 (100%)	3(100%)	NA (12-24)	IV
Narvaez, 2014³²	SSc (9)	P	NA	9 (100%)	NA	NA	IV
Giuggioli, 2015³³	SSc (6)	P	375 mg/m2 at weeks 0/1/2/3	3 (50%)	0 (0%)	30 (18-48)	IV
Fredi, 2015²⁹	DM (2)	R	NA	1 (50)	0 (0%)	201.8	IV
Children							
Aggarwal, 2016³⁰	DM (22)	RCT	0.575-1g/m2 at weeks 0/1	NA	1 (4%)	1.46	IB
Bader-Meunier, 2011³⁴	DM (6)	R	2x500mg/m2 (n=3) 4x375 mg/m2 (n=3)	0 (0%)	0 (0%)	NA (20.2-36)	IV
Alhemairi, 2017³⁵	DM (4)	R	NA	4 (100%)	1 (25%)	NA (36-60)	IV

465 DM: dermatomyositis, SSc: systemic sclerosis RCT: randomized controlled trial, P: prospective, R: retrospective, NA: not available

466 *according to modified Oxford Centre for Evidence-Based Medicine Levels of Evidence and Grades of Recommendation

467

468

469

470

471 **Table 3. Characteristics of included studies evaluating the effect of bisphosphonates, surgery and physical therapies on calcinosis cutis improvement**

Name/year	Disease (N)	Study design	Drug	Dose regimen	Partial response, N (%)	Complete response, N (%)	Follow-up, mean (range), months	Level of evidence, grade*
Biphosphonates								
<i>Adult</i>								
Balin, 2012 ²⁸	NA (2)	R	Etidronate	NA	1 (50%)	0 (0%)	NA	IV
Children								
Marco Puche, 2009 ³⁹	DM (3)	R	Pamidronate	IV 1mg/kg/day at day 1, 2, 3 every 3 months	3 (100%)	1 (33%)	42	IV
Tayfur, 2015 ⁴⁰	DM (3)	R	Pamidronate	IV 1mg/kg/day every 3 months	2 (66%)	1 (33%)	84	IV
	DM (1)	R	Risedronate	PO 1.25 mg/day	1 (100%)	1 (100%)	84	IV
	DM (2)	R	Alendronate	PO 70mg/week	0 (0%)	0 (0%)	84	IV
Saini, 2016 ⁴¹	DM (6)	R	Alendronate	NA	4 (66%)	0 (0%)	22.32 (4.9-27.7)	IV
Surgery and physical therapies								
<i>Adult</i>								
Bottomley, 1996	SSc (6)	P	Carbon dioxide laser	Range of power between 7-5 and 10 W	5 (83%)	NS	> 6	IV
Balin, 2012	NA (28)	R	Surgical excision		27 (96%)	22 (79%)	104	IV
	NA (1)	R	Low frequency ultrasound	NS	1 (100%)	0 (0%)	NA	IV
Blumhardt, 2016	SSc (4)	P	ESWT	1 day/week 3 weeks	4 (100%)	0 (0%)	3	IV
Sultant-Bichat, 2012	DM (1) SSc (2)	P	ESWT	1 day/3 weeks, 3 times	1 (33%)	0 (0%)	8	IV
Shetty, 2005	SSc (3)	P	Iontophoresis of acetic acid + ultrasound	3 d/ week, 3 weeks	0 (0%)	0 (0%)	0.75	IIB
Fahmy, 1998	SSc (15)**	P	Surgical excision (microdrilling)		12 (80%)	NA	0.1	IV

472 R: retrospective, IV: intravenously, PO: Per mouth, NA: not available, DM: dermatomyositis, SSc: systemic sclerosis; ESWT: extracorporeal shock-wave therapy;
473 *according to modified Oxford Centre for Evidence-Based Medicine Levels of Evidence and Grades of Recommendation. **Improvement in 12/15 digits in 10 treated
474 patients.

475 **Table 4. Summary of available treatments for treating calcinosis cutis in DM and SSc**
 476 **patients**

Drug/therapeutic class ^{ref}	Number of DM/SSc patients treated	Overall response rate (%)	Comments ^{ref} (level of evidence)	Grade of recommendation
Warfarin ^{20,26–29}	6/10 (NA=3)	0-66%	Should not be recommended (IB)	B
Diltiazem ^{10,28,29}	12/12 (NA=14)	0-64%	-May have a preventive effect ¹² -Should be discussed in SSc patients and DM with Raynaud's phenomenon (IV)	C
Infliximab ^{29,36,38}	20/0	0-80%	May be discussed in DM patients (IV)	C
Abatacept ³⁷	4/0	100%	May be discussed in DM patients (IV)	C
Rituximab ^{29–35}	41/18	0-100%	May be discussed in DM and SSc patients	C
Biphosphonates ^{28,39–41}	15/0 (NA=2)	0-100%	May be discussed in DM patients (IV)	C
Intravenous immunoglobulins ^{29,42}	15/0	0-62%	May be discussed in DM patients (IV)	C
Minocyclin ^{28,52}	0/9 (NA=3)	33-88%	May be discussed in SSc patients (IV)	C
Colchicine ^{28,29}	(NA=16)	11-43%	May be discussed in DM and SSc patients (IV)	C
Cyclophosphamide ⁵³	14/0	69%	May be discussed in DM patients (IV)	C
Intravenous sodium thiosulfate ^{48,49}	4/3	0%	Should not be recommended (IV)	C
Topical* or intralesional sodium thiosulfate ^{29,50,51}	7/6	0-100%	May be discussed in DM and SSc patients (IV)	C
Surgical excision and physical therapies ^{28,47}	0/15 (NA=28)	80-96%	-May be discussed in DM and SSc patients (IV) -Carbon dioxide laser ⁴³ , low frequency ultrasound ²⁸ and ESWT ^{44,45} may be alternative treatments to surgery (IV) -Iontophoresis of acetic acid + ultrasound ⁴⁶ seems ineffective (IIB)	C C C

477 DM: dermatomyositis; SSc: systemic sclerosis; NA: DM or SSc patients but unspecified
 478 diagnosis; * including sodium metabisulfite

479

480

