



HAL
open science

Critical Evaluation of Whole-Body Cryostimulation Protocol in Race Horses

Fabien Bogard, Bastien Bouchet, Sébastien Murer, Jean Robert Filliard,
Fabien Beaumont, Guillaume Polidori

► **To cite this version:**

Fabien Bogard, Bastien Bouchet, Sébastien Murer, Jean Robert Filliard, Fabien Beaumont, et al.. Critical Evaluation of Whole-Body Cryostimulation Protocol in Race Horses. Journal of Equine Veterinary Science, 2020, 88, pp.102944 -. 10.1016/j.jevs.2020.102944 . hal-03489832

HAL Id: hal-03489832

<https://hal.science/hal-03489832>

Submitted on 22 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Critical evaluation of whole body cryostimulation protocol in race horses

F. Bogard^a, B. Bouchet^{b,c}, S. Murer^a, JR Filliard^{c,d}, F. Beaumont^a,
G. Polidori^{a,c}

^a GRESPI EA4694, University of Reims Champagne-Ardenne, Reims, France

^b Cryotera, Bezannes, France

^c French Society of Whole Body Cryotherapy (SFCCE), Paris, France

^d Medical Department, National Institute of Sport, Expertise and Performance (INSEP), Paris, France

* Corresponding author: fabien.bogard@univ-reims.fr

Abstract

Cold therapy is commonly used to relieve pain and inflammation and to aid in muscle recovery after exercise in human medicine. A number of applications have also been observed in veterinary practice. In this article, a critical evaluation of equine protocol applied with a new commercial concept of equine Whole Body Cryostimulation (WBC) was made. With this new concept of WBC, the protocol usually utilized for relieving pain and discomfort in humans, has been extended to horses. The investigations described herein focus on the reduction of horse skin temperature when applying human WBC protocols. Based on infrared thermography measurements, results show that exposing a horse for 3 minutes to a temperature of -140°C, which are conventional parameters used for humans, does not induce sufficient skin thermal gradients in horses. Consequently, beneficial cold reflexes such as vasomotor, neuroconduction and biochemical reactions, cannot be triggered. Further investigations should therefore be carried out to design an adequate protocol specifically aimed at horses.

Keywords

Cryotherapy, mares, thermal mapping, Whole Body Cryostimulation protocol, cold reflexes

1. Introduction

Localized cold, whether by means of ice, snow or cold water, has been used in humans for over 2000 years mostly because of its analgesic effects. It also helps reduce muscular spasms and provides muscle relaxation by blocking nerve conduction [1], [2]. This applies to both nociceptive and sensory fibers, thus the decrease in sensory influx would lead to a decrease in muscle spasms contributing to analgesia in the affected area [3]. It is highly prized by athletes in their muscle recovery process [4], [5] and explains why the practice of WBC as a complementary therapy or a natural recovery strategy in humans has become widespread in recent years [6], [7]. Indeed, numerous studies [8], [9], [10] have accumulated scientific evidence supporting the beneficial effects of WBC in the human medical domain as an alternate treatment or rehabilitation technique. Similar findings have been made in the sporting realm, especially in the context of post-exercise recovery [1], [11]. The principle commonly implemented is to cause a thermal stress in a confined space at a temperature generally comprised between -160°C and -110°C, for a limited time of about 3 minutes.

Applications of cold to mammals are few [12] and only horses actually undergo cryotherapy sessions in local treatment [13], [14], [15]. Several articles in the literature recommend the use of cold, and provide advice related to methods and local duration of action on horses [16], [17], [18]. Used by veterinarians since the early 2000s, cryotherapy techniques for horses consist in ice applied through a continuous cold boot, sprayed liquid nitrogen or the use of a cryotherapy probe and gas cryotherapy [17], [18], [1]. It has been shown that these localized treatment techniques can bring

56 real benefits in post-traumatic treatment [19], [20], [21]. However, reaching the same temperature
57 in the subcutaneous tissue as on the skin involves the onset of a thermal shock. If not, the
58 temperature of deeper tissues will only decrease a few minutes after the application of cold; the
59 cool-down being gradual and therefore the magnitude of cooling will be less than at the skin level
60 [22]. Analgesia is believed to result from slowed nerve conduction, inhibiting the transmission of pain
61 signals to the dorsal horn of the spinal cord [23], [24], [25]. This phenomenon is observed in humans
62 when muscle temperature drops down to a threshold of 27°C [18].
63

64 Today, in the absence of any scientific guidelines for horses, experimental commercial WBC systems
65 for horses are based on the same concept as for humans. More precisely, the whole horse body
66 undergoes a thermal shock at -140°C, caused by exposure to gaseous nitrogen in a semi-closed space
67 over a limited time frame (typically 3 minutes). In humans, this duration is required to trigger cold
68 reflexes, such as vasoconstriction, slowdown or even blocking of nerve conduction, ultimately
69 leading to pain relief [26].
70

71 The purpose of this study was to test the validity of this human WBC technique transposed to horses,
72 by measuring the level of the skin thermal variations thus obtained. To this end, the change in
73 cutaneous temperature was measured over the whole body of the horse after exposure to an
74 extremely cold atmosphere generated by WBC embedded in a new commercial device.
75 Measurements were performed using InfraRed Thermography (IRT) to acquire skin temperature, its
76 response being sensitive to extreme cold. The potential role of IRT in the diagnosis and treatment
77 monitoring is mentioned in veterinary applications [22], especially in equine medicine [27], [28],
78 [29], [30], [31], [32], [33]. This powerful, noninvasive and easy-to-handle technique has a broad range
79 of application. It is used in horses to diagnose symptoms in the fields of equine orthopedics and
80 lameness [34], palmar or plantar digital neurectomy [35], detection of jugular venipuncture [36] and
81 design of training protocols [37], [38]. Then, the critical evaluation of the WBC protocol used in
82 humans at -140°C and extended to race horses, as it is used today, will be carried out by infrared
83 thermography.
84

85 2. Materials and methods

87 2.1. Thermal behavior and dissipation in horses and humans

88 Horses and humans share thermal characteristics. First, they are both homeothermic mammals [25],
89 with a core temperature independent of the surrounding environment and tending to constancy at
90 rest due to neurophysiologic mechanisms balancing heat production and heat losses. This balance is
91 maintained at rest in an environmental temperature range called thermoneutral zone, estimated in
92 horses in the range 5-25°C [39], [40] and in the range 25°C-30°C for a naked man. Core temperatures
93 are very similar, namely 38°C [41] for horses and 37°C for humans. Same goes for the average skin
94 temperature, namely 33°C for horses and humans. The thermal balance is therefore governed by the
95 relationship:

$$96 \text{Heat}_{\text{production}} = \text{Heat}_{\text{dissipation}} \pm \text{Heat}_{\text{storage}} \quad (1)$$

97
98
99 In humans, it has been shown that the internal temperature does not vary during WBC sessions [42].
100 The reason is that the exposure to cold is too short to induce a decrease in internal temperature. If a
101 net internal heat loss had initiated, heat storage would have been negative and the body
102 temperature would have fallen ($S < 0$) [43]. If the internal temperature is constant, it can be assumed
103 that the rate of heat storage ($\text{Heat}_{\text{storage}}$) is zero ($S = 0$). This assumption in humans can reasonably be
104 transposed to horses.

105 Major differences between horses and humans may lie in the way heat dissipates even if for both
106 species, convective (including respiration) and radiative thermal exchanges are the main heat
107 transfer modes:

$$108 \varphi_{\text{heat losses (W/m}^2)} = (\varphi_{\text{conv}} + \varphi_{\text{resp}}) + \varphi_{\text{rad}} \quad (2)$$

109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159

Convective and radiative exchanges φ_{conv} and φ_{rad} are expressed using Newton and Boltzmann laws respectively:

$$\varphi_{\text{heat losses}} (W/m^2) = h_c(T_S - T_\alpha) + \varphi_{\text{resp}} + \sigma \varepsilon (T_S^4 - T_\alpha^4) \quad (3)$$

where h_c is the free convective heat transfer coefficient, σ is the Stefan-Boltzmann constant ($\sigma = 5.67 \times 10^{-8} Wm^{-2}K^{-4}$) and ε is the skin radiative emissivity. Parameters T_S and T_α stand for the skin and ambient temperatures respectively, expressed in K . In horses, the heat loss related to the convective respiratory mechanism φ_{resp} is estimated between 15 and 25% of total loss [42], while it lies around only 10.5% in humans [44].

2.2. Horses studied

Five healthy thoroughbred mares (3 years \pm 3 months of age, 448 \pm 18kg) from a racing stable in Chantilly (France) were recruited for the study. All were followed by a veterinarian who declared them sound, with no musculoskeletal disease history and no clinical symptoms of illness. In the opinion of the horse trainer, no external sign of estrus was observed at the time of the trials, which took place at the end of October, outside the breeding season [50]. Prior to the experiments, early in the morning, the horses performed a horse's standard exercise protocol composed of 800 m at promenade allure ($\approx 7 \text{ km}\cdot\text{h}^{-1}$), 1200 m trotting ($\approx 14 \text{ km}\cdot\text{h}^{-1}$), 1400 m hunting canter ($\approx 20 \text{ km}\cdot\text{h}^{-1}$) and 1500 m quick canter ($\approx 50 \text{ km}\cdot\text{h}^{-1}$). They stayed in the stable for 3 hours to cool down afterwards. The owners of the horses as well as the horse trainer who owns the race stable gave their consent for this study.

2.3. Whole Body Cryostimulation chamber for horses

The development of this cryotherapy device for horses was inspired by the mobile units for athletes that enable cryotherapy devices to be brought directly onto sporting events or race sites. In the present case, the truck is replaced by a two-horses van (Figures 1A and 1B). The technology is based on the decompression of liquid nitrogen to reach extreme temperatures very quickly inside the cryostimulation chamber. For safety considerations, the head of the horses remained outside the cooled zone to avoid any nitrogen respiration (Figure 1B). This principle is found in cryo-cabins for humans [1], where the top face of the cylinder constituting the cryotherapy chamber is open such that the free surface of nitrogen lies at the neck level. Moreover, horses were equipped with earplugs to avoid possible stress due to noises emitted by the device. It should be noted that all the mares were familiar with this kind of cryostimulation and none of them showed any resistance to entering the chamber during the experiments, nor appeared to be under any stress according to the horse trainer feedback. However, it is worth reminding that naïve horses must be gradually acquainted with entering the mobile chamber to minimize stress.

For all horses, the selected protocol was identical to the one conventionally used for humans and commonly adopted in cryo-cabins, i.e. an exposure of 3 minutes at -140°C . Each horse was thermally photographed (instantaneous capture) on its left side before the trainer lead it into the van (Fig. 1A) for the cryostimulation session. This picture is the reference thermal level. After the cryostimulation session, immediately after leaving the van, the horse is thermally photographed again on the same side. This last picture makes it possible to evaluate the influence of thermal shock on the horse's skin temperature in comparison with the reference image. Although all five horses were evaluated in the same way, the experiments were not repeated over several days, which may be a limitation of the study.



Fig. 1. Mobile cryostimulation chamber (A); Horse's head out of the cryotherapy device (B).

2.4. Infrared thermography and environmental conditions

Skin temperatures were recorded using a ThermoVision SC620 thermal imaging camera (Flir Systems, Wilsonville, OR) in accordance with the standard protocol of infrared imaging in medicine [51]. For each horse, two infrared thermography pictures were acquired, the first one just before entering the cryotherapy chamber, and the second just after leaving it.

It is worth noting that healthy horses display thermal symmetry similarly to humans. Indeed, healthy humans have a high degree of thermal symmetry in terms of both magnitude and pattern [52]. The results can easily be extended to mammals and more specifically to horses. This is the reason why only the left laterality of the horses was thermally treated.

Since the experiments took place outdoors, knowledge of the atmospheric conditions is necessary to qualify dissipative heat transfers. Measurements of outdoor temperature, atmospheric pressure, humidity and wind speed were performed throughout the experiment, and are summarized in Table 1. Experiments were conducted in a shaded area away from the sun.


Outdoor temperature	Relative humidity	Outdoor pressure	Wind speed	Weather
12.7 °C	69.35 %	1025 hPa	0.7 m.s ⁻¹	Sunny

Table 1. Atmospheric working conditions during IRT measurements.

For accurate measurement by infrared thermography, the knowledge of skin emissivity is key, since it reflects the effectiveness of the emission of infrared radiation by the skin. In published works on equine thermography, the emissivity has not been accurately defined, ranging from 0.95 to 1. An estimated emissivity value of 0.98, already adopted in the literature [52], was selected.

It should be noted that in race horses, the use of infrared thermography to access skin temperatures comes without generating measurement bias since these horses are characterized by a coat that is often short [53].


Quantitatively, thermal imaging was analyzed according to regions of interest that we identified as being some of the areas where muscle tensions appear in the horse's trunk [54]. In detail, nine areas have been identified, corresponding respectively to the subcutaneous positions of the trapezius muscles over the shoulder, latissimus dorsi muscle of the back, medium and superficial buttock muscles, semi-tendinosus and gluteo-femoral muscles of the hindlimb, latissimus dorsi muscle and brachiocephalic muscle of the neck. These areas are depicted in Figure 2. We hypothesize that in horses, as in humans [55], there is a relationship between surface temperature and intramuscular temperature. Thus, measuring the surface temperatures over the 9 identified areas must make it possible to obtain information on the corresponding deep tissues, which are ordinarily the target tissues of cryotherapy applications.



201
202
203 **Fig.2.** Areas of muscle tension in the horse’s trunk [54] selected for the IRT imaging.

204
205
206 **3. Results**


207
208 Figure 3 presents the infrared thermography mappings of the same mare just before and
209 immediately after the cryostimulation session, at -140°C for a 3 minutes’ duration, with identical
210 color-coded scales. It clearly appears in Figure 3A, at rest and before the cryostimulation session,
211 that the skin temperature is not uniform over the body surface, the maximum amplitude measured
212 being 10.5°C (maximum at the throat latch: 32.4°C and minimum at the fetlock of the hind leg:
213 21.9°C). Higher skin temperatures are generally obtained in the vicinity of vital organs that must be
214 thermally preserved. Indeed, in these areas, greater heat production (to preserve vital organs)
215 induces greater heat dissipation to maintain the thermal balance, resulting in an increase in skin
216 temperature.



218
219
220 **Fig. 3.** « Red-Whisper » (3 years old, 466 kg) just before the cryo-session (A),
221 immediately post-session (B)

222
223 It should be noted that for each mare and each area investigated, the measurement obtained is not a
224 one-point measurement but an average measurement over an area of approximately 100 cm².
225 Figure 4 represents the statistically averaged measured temperature over all mares before and
226 immediately post cryostimulation treatment, for the different body regions of interest. The *
227 indicates a significant skin temperature difference between pre and post treatment ($P \leq 0.05$), β

228 indicates a significant skin temperature difference between pre and post treatment ($P \leq 0.001$).
 229 Muscular area numbers are presented in Figure 2.
 230 Standard deviation is also reported. The observation that can be drawn from these results seems to
 231 be that temperature decreases too slightly between pre and post-cryostimulation. For a -140°C
 232 cryostimulation during 3 minutes, corresponding to human protocol, the induced skin cooling varies
 233 from -1.12°C for the large dorsal to -2.80°C for medium buttock in horses.
 234
 235



236
 237
 238
 239 **Fig.4.** Average skin temperature measurements and standard deviation obtained through IRT.

240
 241
 242 **4. Discussion**

243
 244 It is reminded that as a consequence of the dissimilar morphologies between humans and horses, all
 245 the comparisons herein are questionable. Nevertheless, it can be estimated from the results that skin
 246 cooling level observed in horses is 5 to 6 times lower than that of humans, when the same
 247 cryostimulation protocol at -140°C for a duration of 3 minutes is applied.

248 For comparison purposes about Figure 3, we can refer to a recent study in which 10 human subjects
 249 (50% female; means \pm S.D.: age 45.8 ± 5.5 years, height 168.7 ± 9.3 cm, weight 75.3 ± 13.1 kg, body
 250 fat percentage 19.3 ± 9.8) participated in cryostimulation sessions with the same protocol [58].
 251 Thermal imaging was quantitatively analyzed according to given regions of interest, namely chest,
 252 upper back, arms, stomach, lower back and legs. In humans, the induced skin cooling was found to
 253 vary from -6.30°C for the upper back to -14.35°C for the legs.

254 This difference could be attributed to several factors. First, contrary to radiative dissipative heat
 255 transfer which is independent of the body shape, convective heat transfer is quite different between
 256 horses and humans. Their body shapes are actually quite dissimilar; the human body can roughly be
 257 assimilated in shape to a volumetric heat source embedded in a thin vertical cylinder while horses
 258 can be considered as a large horizontal one. Convective heat transfer stemming from these simplified
 259 geometries are therefore different [45]. Subsequently, at an ambient temperature between -3°C and
 260 37°C , horses undergo a total heat loss [40] corresponding to a dissipative heat flux density of about
 261 142 W/m^2 (without evaporating losses), while humans at rest present a value close to 50 W/m^2 . This
 262 order of magnitude may be calculated by considering a key physiological parameter in cryotherapy,
 263 which is the ratio between the mass (kg) and body surface area (m^2) of humans and horses alike. It
 264 can reasonably be estimated at $80 \text{ kg}/1.8 \text{ m}^2$ for an average male human, and $500 \text{ kg}/5 \text{ m}^2$ for horses
 265 [47], which shows that horses need to dissipate 2.5 to 3 times as much heat through each square
 266 meter of body surface area than men. Therefore, thermal balance is ensured by a higher heat
 267 production in horses than in humans in the same proportion. When facing extreme cold conditions,
 268 the lower the body mass relative to body surface, the higher the heat dissipation compared to basal

269 metabolic rate; this explains in part why humans are more subject to hypothermia than horses, and
270 why horses may tolerate temperatures as low as -40°C [48]. Another key parameter when it comes to
271 facing cold is the tissue thermal insulance. This value, which is rarely studied in animals, is
272 nevertheless fundamental in the way the body reacts to cold. The reference value of tissue thermal
273 insulance in horses ($0.10 \text{ m}^2\text{K/W}$ [40]) is twice that of humans ($0.05 \text{ m}^2\text{K/W}$ – derived from [42]),
274 which also explains why the latter cannot withstand temperatures as low as horses do. One of the
275 major reasons why the discrepancies between tissue thermal insulance in horses and humans lies in
276 their skin structures, which provides a physiological barrier between the external and internal
277 environments as well as higher thermal inertia [59] than humans. Horses and humans strongly differ
278 in this matter, even though skin is the largest organ in both these species and constitutes a
279 temperature regulator as well as the key factor when it comes to heat loss issued from the body. Its
280 thickness is estimated on average to be about 3.8 mm in horses [49], while it is commonly accepted
281 that the average value in humans lies around 0.6 mm. For obvious reasons, the thickness and
282 structure (presence of leather) of the skin may affect the thermal insulance of tissues. These
283 different thermal characteristics account for the discrepancies in the reactions of horses and humans
284 to extreme cold. In the end, these differences prevented the cold from penetrating rapidly into the
285 horse's tissues during the reduced time frame of the cryostimulation session.

286 A number of potential limitations need to be considered. First, the thermal mapping depicted in
287 Figure 3 and obtained before cryostimulation indicates values lower than the average skin
288 temperature in horses mentioned in the literature, i.e. around 33°C . One of the reasons is that
289 experiments were conducted in a real outdoor situation in October in northern France at a
290 temperature of 12.7°C and a light breeze of $0.7 \text{ m}\cdot\text{s}^{-1}$. Under these climatic conditions, the
291 characteristics of dissipative heat transfers are obviously different from those that could be obtained
292 under laboratory conditions. This would support in part an assumption found in the literature, which
293 states that wind speed has an influence on the amplitude of skin temperature variation [57]. Another
294 limitation may lie in the reduced number of horses involved in the study.

295 In spite of the discrepancies in thermal characteristics mentioned above, the principle of equine
296 thermoregulation in the face of extreme cold remains similar to that of humans. Thus, one may
297 reasonably think that WBC can also produce beneficial effects in horses. Given the current state of
298 knowledge, transferring human protocols to horses produces insufficient benefits in terms of skin
299 temperature decrease.

300 One possible solution would be to select colder treatment temperatures, but nitrogen-based WBC
301 technologies do not allow it [60]. Therefore, only the extension of the exposure time may be
302 considered. To define the optimal duration of exposure to cold, a test campaign related to the skin
303 cooling kinetics of the horse should be performed by measuring thermal fields at increasing time
304 intervals of exposure and on a larger number of animals. This would make it possible to estimate the
305 thermoregulatory behavior of the horse in WBC, which remains unknown to date, unlike that of
306 humans [61]. Knowledge of this behavior will provide essential information in the definition of a WBC
307 protocol adapted to the horse's particularities.

308
309

310 **5. Conclusion**

311

312 This experimental study focused on a critical evaluation of whole body cryostimulation protocol in
313 race horses. For this purpose, the reaction to extreme cold of five race horses was analyzed by means
314 of infrared thermography. The benefits of WBC on muscle recovery in equestrian sport seem illusory
315 at the current state when protocols are transposed from humans (3 min at 140°C) to horses. Indeed,
316 the recorded skin variations seem far too small to be able to induce interesting therapeutic or muscle
317 recovery effects. Future studies appear necessary and should therefore focus on analyzing cold
318 exposure duration to define an appropriate protocol for race horses.

319
320

321 **Ethics Statement**

322 Based on European directive 2010/63 transposed into French law (decree 2013-118), no ethical issue
323 arises from this study as experimental data is only gathered from observational field studies.
324 Observations were made in the frame of normal cryosessions scheduled by the trainer, under his full
325 responsibility. The horses were familiar with these sessions: every effort was made to ensure their
326 welfare by minimizing disturbance caused by researchers and the horses showed no sign of stress.

327 **Conflict of interest**

328 The authors have no conflict of interest to declare.

329

330 **Acknowledgments**

331 The authors would like to thank Mikel Delzangles's team and owners of race horses for making this
332 experimental work possible. They also gratefully acknowledge the French Society of Whole Body
333 Cryotherapy (SFCCE) for their interest in this study.

334

335

336 **References**

- 337 [1] Bouzigon R, Grappe F, Ravier G, Dugue B. Whole- and partial-body cryostimulation/cryotherapy:
338 Current technologies and practical applications. *J Therm Biol* 2016;61:67-81.
- 339 [2] Herrera E, Sandoval MC, Camargo DM, Salvini TF. Motor and sensory nerve conduction are
340 affected differently by ice pack, ice massage, and cold-water immersion. *Phys Ther.*
341 2010;(90):581-591.
- 342 [3] Commandre FA, Fish F. Douleur, Cryothérapie et sport. *Médecine du sport.* 1997;71(1):5-11.
- 343 [4] Ziemann E, Olek RA, Kujach S, Grzywacz T, Antosiewicz J, Garsztka T, et al. Five-day whole-body
344 cryostimulation, blood inflammatory markers, and performance in high-ranking professional
345 tennis players. *J Athl Train.* 2012;47(6):664–672.
- 346 [5] Lombardi G, Lanteri P, Porcelli S, Mauri C, Colombini A, Grasso D, et al. Hematological profile
347 and martial status in Rugby players during whole body Cryostimulation. *PLoS One*
348 2013;8(2):e55803.
- 349 [6] Hausswirth C, Louis J, Bieuzen F, Pournot H, Fournier J, Filliard JR, Brisswalter J. Effects of whole-
350 body cryotherapy vs. far-infrared vs. passive modalities on recovery from exercise-induced
351 muscle damage in highly-trained runners. *PLoS One.* 2011;(6):e27749.
- 352 [7] Bleakley C, Bieuzen F, Davison Gareth W, Costello J. Whole-body cryotherapy: empirical
353 evidence and theoretical perspectives. *Open Access J. Sports Med.* 2014;(5):25-36.
- 354 [8] Vitenet M, Tubez F, Marreiro I, Polidori G, Taiar R, Legrand F, Boyer F. Effect of Whole Body
355 Cryotherapy interventions on health-related quality of life in fibromyalgia patients: a
356 randomized controlled trial. *Comp Therap in Med.* 2018;36:6–8.
- 357 [9] Bouzigon R, Ravier G, Dugue B, Grappe F. The use of whole-body cryostimulation to improve the
358 quality of sleep in athletes during high level standard competitions. *Br J Sports Med.*
359 2014;(48):572.
- 360 [10] Stanek A, Cholewka A, Galuda J, Drzazga Z, Sieron A, Sieron-Stoltny K. Can whole-body
361 cryotherapy with subsequent kinesiotherapy procedures in closed type cryogenic chamber
362 improve BASDAI, BASFI, and some spine mobility parameters and decrease pain intensity in
363 patients with ankylosing spondylitis? *BioMed Res Int.* 2015;(2015):11p.
- 364 [11] Rose C, Edwards KM, Siegler J, Graham K, Caillaud C. Whole-body cryotherapy as a recovery
365 technique after exercise: a review of the literature. *Int J Sports Med.* 2017;38(14):1049–1060.
- 366 [12] Richman AW, Kirby AL, Rosenkrantz W, Muse R. Persistent papilloma treated with cryotherapy
367 in three dogs. *Vet. Dermatol* 2017;28:625-e154.
- 368 [13] Hunt ER. Response of twenty-seven horses with lower leg injuries to cold spa bath
369 hydrotherapy, *J of Eq Vet Sc* 2001;21(4):188-193.
- 370 [14] Petrov R, MacDonald M, Tesch A, Van Hoogmoed L. Influence of topically applied cold treatment
371 on core temperature and cell viability in equine superficial digital flexor tendons. *Am J Vet Res*
372 2003;64(7):835-844.
- 373 [15] Ramey DW. Cold therapy in the horse. *Equine Practice* 1999;21(1):19-22.

- 374 [16] Desbrosse F. Les propriétés analgésiques de la cryothérapie au CO2 hyperbare. *Pratique*
375 *Vétérinaire Equine* 2003;35:97-106.
- 376 [17] Pollitt CC, Van Eps AW. Prolonged, continuous distal limb cryotherapy in the horse. *J Equine Vet*
377 *Sci* 2004;36(3):216-220.
- 378 [18] Roszkowska K, Witkowska-Pilaszewicz O, Przewozny M, Cywinska A. Whole body and partial
379 body cryotherapies – lessons from human practice and possible application for horses. *BMC Vet*
380 *Res.* 2018;14:394.
- 381 [19] van Eps AW, Orsini JA. A comparison of seven methods for continuous therapeutic cooling of the
382 equine digit. *Equine Vet J.* 2016;48(1):120–124.
- 383 [20] van Eps AW, Pollitt CC. Equine laminitis model: cryotherapy reduces the severity of lesions
384 evaluated seven days after induction with oligofructose. *Equine Vet J.* 2009;41(8):741–746.
- 385 [21] Kullmann A, Holcombe SJ, Hurcombe SD, Roessner HA, Hauptman JG, Geor RJ, et al. Prophylactic
386 digital cryotherapy is associated with decreased incidence of laminitis in horses diagnosed with
387 colitis. *Equine Vet J.* 2014;46(5):554–559.
- 388 [22] Delongea M. Utilisation des piscines, marcheurs aquatiques et spas dans la rééducation et
389 l'entraînement du cheval, Thèse de l'Université Claude-Bernard. Lyon1, France. 2008.
- 390 [23] Galletier A. *Le Pur-Sang Arabe.* Editions Atlas. 2011:125p.
- 391 [24] Langlois B. Inter breed variation in the horse with regard to cold adaptation: a review. *Livestock*
392 *Production Sci* 1994;40:1-7.
- 393 [25] Piccione G, Giannetto C, Marafioti C, Casella S, Assenza A, Fazio F. Comparison of daily rhythm of
394 rectal and auricular temperatures in horses kept under a natural photoperiod and constant
395 darkness. *J of Thermal Biology* 2011;36(4):245-249.
- 396 [26] Stocks JM, Taylor NAS, Tipton MJ, Greenleaf JE. Human physiological responses to cold
397 exposure. *Aviat Space Environ Med* 2004;75:444-57.
- 398 [27] Soroko M, Howell K, Dudek K, Wilk I, Zastrzeżyńska M, Janczarek I. A Pilot Study Into the Utility
399 of Dynamic Infrared Thermography for Measuring Body Surface Temperature Changes During
400 Treadmill Exercise in Horses. *J of Equine Vet Sc* 2018;62:44-46.
- 401 [28] Rekant S., Lyons MA, Pacheco JM, Arzt J., Rodriguez, LL., Veterinary applications of infrared
402 thermography. *Am J Vet Res* 2016;77(1):98-107.
- 403 [29] Soroko M, Howell K. Infrared Thermography: Current Applications in Equine Medicine. *J of*
404 *Equine Vet Sc* 2018;60:90–96.
- 405 [30] Redaelli V, Bergero D, Zucca E, Ferrucci F, Nanni Costa L, Crosta L, Luzi F. Use of Thermography
406 Techniques in Equines: Principles and Applications. *J of Equine Vet Sc* 2014;34(3):345-350.
- 407 [31] Eddy AL, Van Hoogmoed LM, Snyder JR. The role of thermography in the management of equine
408 lameness. *The Vet J* 2001;162:172-181.
- 409 [32] Simon EL, Gaughan EM, EPP T, Spire M. Influence of exercise on thermographically determined
410 surface temperatures of thoracic and pelvic limbs in horses, *J Am Vet Med Assoc*
411 2006;229(12):1940-4.
- 412 [33] Loving NS, Johnston AN. Therapy for leg swelling In: *Veterinary Manual for the Performance*
413 *Horse.* Blackwell Science, London: 1995, p.238-242.
- 414 [34] Soroko M, Henklewski R, Filipowski H, Jodkowska E. The Effectiveness of Thermographic Analysis
415 in Equine Orthopedics. *J of Equine Vet Sc* 2013;33:760-762.
- 416 [35] Van Hoogmoed LM, Snyder JR. Use of infrared thermography to detect injections and palmar
417 digital neurectomy in horses. *The Vet Journal.* 2002;164:129-141.
- 418 [36] DGLISH J, le Jeune S, Pypendo BH, Millares Ramirez E., Turner TA. Use of Infrared Thermography
419 to Detect Jugular Venipuncture in the Horse. *J Equine Vet Sc.* 2017;59:1-6.
- 420 [37] Yarnella K, Fleming J, Stratton TD, Brassington R. Monitoring changes in skin temperature
421 associated with exercise in horses on a water treadmill by use of infrared thermography. *J of*
422 *Thermal Biology* 2014;45:110-116.
- 423 [38] Zalewski P, Klawe JJ, Pawlak J, Tafil-Klawe M, Newton J. Thermal and hemodynamic response to
424 whole-body cryostimulation in healthy subjects. *Cryobiology.* 2013;66(3):295–302.
- 425 [39] Zalewski P, Buszko K, Zawadka-Kunikowska M, Slomko J, Szrajda J, Klawe JJ, et al. Cardiovascular
426 and autonomic responses to whole-body cryostimulation in essential hypertension. *Cryobiology.*
427 2014;69(2):249–255.

- 428 [40] Morgan K. Thermoneutral zone and critical temperatures of horses. *J. Therm Biol* 1998;23(1):59-
429 61.
- 430 [41] Green AR, Gates RS, Lawrence LM. Measurement of horse core body temperature. *J of Thermal*
431 *Biology*. 2005;30(5):370-377.
- 432 [42] Polidori G, Cuttelli S, Hammond L, Langdon D, Legrand F, Taiar R, Boyer FC, Costello JT. Should
433 whole body cryotherapy sessions be differentiated between women and men? A preliminary
434 study on the role of the body thermal resistance. *Med Hyp* 2018;120:60-64.
- 435 [43] Parsons KC. *Human Thermal Environments: the effects of hot, moderate, and cold environments*
436 *on human health, comfort and performance* 2nd Ed. Taylor & Francis, London and New York.
437 2003.
- 438 [44] Hinchcliff KW, Geor RJ, Kaneps AJ. *Equine exercise physiology. The Science of Exercise in the*
439 *Athletic Horse*. Saunders Ltd Ed. Philadelphia 2008:464p.
- 440 [45] Kakaç S, Yener Y, Pramuanjaroenkij A. *Convective Heat Transfer*. 3rd Edition. CRC Press. Boca
441 Raton 2014:622 p.
- 442 [46] Morgan K, Ehrlemark A, Sällvik K. Dissipation of heat from standing horses exposed to ambient
443 temperatures between -3°C and 37°C . *J Therm Biol* 1997;22(3):177-186.
- 444 [47] Martin D, Nankervis K. *Equine Exercise Physiology*. London, Blackwell Publishing 2002:296p.
- 445 [48] Cymbaluk NF, Christison GI. Environmental effects on thermoregulation and nutrition of horses.
446 *Vet Clin North Am Equine Pract* 1990;6(2):355-72.
- 447 [49] Wakuri H, Mutoh K, Ichikawa H, Liu B. Microscopic anatomy of the equine skin with special
448 reference to the dermis. *Okajimas Folia Anat. Jpn* 1995;72(2-3);177-184.
- 449 [50] Cuervo-Arango J, Clark A. The first ovulation of the breeding season in the mare: the effect of
450 progesterone priming on pregnancy rate and breeding management (hCG response rate and
451 number of services per cycle and mare). *Anim Reprod Sci* 2010;118(2-4):265-9.
- 452 [51] Ring E, Ammer K. The technique of infrared imaging in medicine. *Thermol Intern* 2000;10:7-14.
- 453 [52] Jiang LJ, Ng EY, Yeo AC et al. A perspective on medical infrared imaging. *J Med Eng Technom*.
454 2005;29(6):257-267.
- 455 [53] Westermann S, Buchner HHF, Schramel JP, Tichy A, Stanek C. Effects of infrared camera angle
456 and distance on measurement and reproducibility of thermographically determined
457 temperatures of the distolateral aspects of the forelimbs in horses. *J Am Vet Med Assoc*
458 2013;242: 388-395.
- 459 [54] Turner TA, Purohit RC, Fessler JF. Thermography: a review in equine medicine. *Comp Cont Educ*
460 *Pract Vet* 1986;8:855-861.
- 461 [55] Denoix JM, Pailloux JP. *Approche de la kinésithérapie du cheval*. Editions Maloine. Paris.
462 1997:291p.
- 463 [56] Hardaker NJ, Moss AD, Richards J, Jarvis S, McEwan I, Selfe J. Relationship between
464 intramuscular temperature and skin surface temperature as measured by thermal imaging
465 camera, *Thermol Int* 2007;17(2):45-50.
- 466 [57] Westermann S, Stanek C, Schramel JO, Ion A, Buchner HF. The effect of airflow on
467 thermographically determined temperature of the distal forelimb of the horse. *Eq Vet Journal*
468 2013;45:637-641.
- 469 [58] Polidori G, Taiar R, Legrand F, Beaumont F, Murer S, Bogard F, Boyer FC. Infrared thermography
470 for assessing skin temperature differences between Partial Body Cryotherapy and Whole Body
471 Cryotherapy devices at -140°C . *Inf. Phys. and Tech* 2018;93:158-161.
- 472 [59] Hodgson DR, Davis RE, McConaghy FF. Thermoregulation in the horse in response to exercise. *Br*
473 *Vet J* 1994;150:219-234.
- 474 [60] Baranov A, Pakhomov O, Fedorov A, Ivanov V, Zaitsev A, Polyakov R. *Technique and Technology*
475 *of Whole-Body Cryotherapy (WBC)*. IntechOpen; 2019.
- 476 [61] Costello JT, Culligan K, Selfe J, Donnelly AE. Muscle, Skin and Core Temperature after -110°C
477 Cold Air and 8°C Water Treatment. *Plos One*. 2012;7(11): e48190.