

**E-health. Patterns of use and perceived benefits and barriers among people living with HIV  
and their physicians. Part 2: health apps and smart devices.**

C JACOMET<sup>1</sup>, R OLOGEANU-TADDEI<sup>2</sup>, J PROUTEAU<sup>3</sup>, C LAMBERT<sup>4</sup>, F LINARD<sup>5</sup>, P BASTIANI<sup>6</sup> and  
P DELLAMONICA<sup>7</sup>

<sup>1</sup>Service des Maladies Infectieuses et Tropicales, CHU Clermont-Ferrand, France.  
cjacomet@chu-clermontferrand.fr

<sup>2</sup>Systèmes d'Information–Montpellier Recherche en Management & Polytech Montpellier,  
Université de Montpellier, France. roxana.ologeanu-taddei@umontpellier.fr

<sup>3</sup>Service des Maladies Infectieuses et Tropicales, CHU Clermont-Ferrand. France  
jprouteau@chu-clermontferrand.fr

<sup>4</sup>Délégation Recherche Clinique & Innovation, CHU Clermont-Ferrand, France.  
clambert@chu-clermontferrand.fr

<sup>5</sup>Services des Maladies Infectieuses et Tropicales, CHU Tenon, CHU Hôtel Dieu, APHP-  
Sorbonne Université, France. Francoise.linard@outlook.com

<sup>6</sup>AC SIDA, Nice. France pbastia06@gmail.com

<sup>7</sup>Distinguished Lecturer, Service des Maladies Infectieuses et Tropicales, Université de la  
Cote d'Azur, France. dellamopierre@yahoo.fr

**Keywords:** HIV; e-health; mobile applications

## **Abstract**

**Objectives.** To evaluate patterns of use and perceived benefits and barriers to health/wellness applications (apps) and smart devices among people living with HIV (PLHIV) and their physicians.

**Methods.** Online multicenter observational survey (October 15-19, 2018).

**Results.** Study participation was accepted by 229 physicians and 838/1,377 PLHIV followed in 46 centers, of which 325 (39%) responded online. Overall, 83/288 (29%) PLHIV had already downloaded at least one app: these 'downloaders' were younger (OR 0.96  $\pm$ 0.01,  $p=0.004$ ), educated to at least university entry level (OR 2.27  $\pm$ 0.86,  $p=0.03$ ), and more frequently used geolocation-based dating websites (OR 3.00  $\pm$ 1.09,  $p=0.002$ ). However, 227/314 (72%) PLHIV claimed they were ready to use an app recommended by a physician. For the 60/83 PLHIV who answered, the ideal app would be a vaccination tracker (76%) to better communicate with their physician (68%). However, 96/277 (42%) physicians were unable to answer this question and for 94/227 (41%) of them, the ideal patient app would be for schedule management. Although PLHIV used smart devices, 231/306 (75%) would want to report the data to their physicians and 137/225 (61%) of physicians would welcome this exchange. The main physician-side barrier to this exchange was concerns over data security.

**Conclusion.** mHealth apps and smart devices have failed to garner adoption by PLHIV. There is a case for good-quality health data sharing and exchange if PLHIV are provided with appropriately secure tools and physicians are backed up by adapted legislation.

## Introduction

Digital technologies have invested every aspect of our lives. A survey conducted by specialists pollsters Opinion Way to coincide with the Distree #Connect 2017 smart tech forum found that 45% of French people see smart devices as “a revolution, much like the Internet a few years back” [1]. A recent BVA survey found that the French put health ahead of safety: 64% of people surveyed claimed that smart device development research should focus first on health applications, against 60% for safety [2]. Digital health industry analysts and strategy consultants Research 2 Guidance reported an estimated 259,000 health apps in 2016 against 100,000 just a year before [3]. eHealth, particularly through mHealth-enabling smart devices, poses a number of challenges for prevention and management of HIV infection. The emergent cultural phenomenon of ‘quantified self’ goes in tandem with the spread of increasingly ubiquitous wearable smart devices (sensors, wristbands, watches plus mobile apps). The movement extends and revisits ways to measure body, mind, and activity patterns [4]. This first ‘quantified self’ step teaches practitioners better self-analysis and self-surveillance of their own health indicators. It is claimed as an easy, inclusive, and readily accessible step towards better preventive care or better-coordinated care pathways [5]. It may lead to a second phase called ‘modified self’ where practitioners learn to modify their behaviors and routines and better manage the ageing process, which is a major issue for PLHIV [6]. There is already an array of practical apps, like the HIV prevention apps for men who have sex with men that have been rolled out by the European Center for Disease Prevention and Control (ECDC), or like the Canadian mHealth app HIVSmart that guides people through the process of getting HIV-tested, with links to healthcare facilities and counsellors [7-8]. Others like Hiv/Aidstestapp and Seronet/L’info en + have been created by a French nonprofit called AIDES [9-11]. The systematic reviews that identified HIV-related

apps have also shown that they register few downloads [12-13]. Ethical questions have emerged and, over time, mistrust, misinformation, and concern over unwelcome surveillance and unsafe data protection have surfaced. In France, it was only in 2016 that the French National Authority for Health (French acronym HAS) finally issued long-overdue guidelines on information to users, health-related content, technical media, security/reliability, and uses/utilization [14]. Furthermore, caregivers today rarely make use of the data collected by smart devices and mobile apps. Swendeman *et al.* studied HIV-clinic healthcare providers' attitudes to apps, focusing on their goal to deliver better quality-of-life, and identified a number of barriers, chiefly organizational barriers [15].

We report a survey conducted in 2018 on PLHIV and their physicians that set out to identify patterns of use and perceived benefits and barriers to eHealth in France. The question addressed is whether mHealth, defined as “the practice of medicine and public health supported by mobile devices such as mobile phones, patient monitoring systems, personal digital assistants, and other wireless-enabled devices” serves PLHIV and their physicians as part of a wider self-managed care offer?

## **Methods**

We carried out an online multicenter observational ‘random-week’ survey on all HIV-positive patients referred for consultation at short-stay outpatient clinics from October 15-19, 2018 via regional coordinated care organizations (COREVIH care pathway coordinators), as described and reported elsewhere [16].

## Results

The survey was led at 51 clinics throughout France, and 255 physicians who had seen 1,377 PLHIV attending consultation during the study period were surveyed [16]. Of these 1,377 patients, 144 were ineligible, 395 refused to participate, and 838 were given login details to complete the survey. A total of 325 patients attending 46 of the centers, including 191 in-hospital outpatients (59%), completed the online questionnaire. The Ile-de-France contingent counted 117 PLHIV (36% of the sample). There was a significant difference in survey participants vs non-participants on gender make-up (72% men vs 65%, respectively;  $p=0.03$ ) and mean age ( $52.6 \pm 11.9$  years vs  $50.6 \pm 12.1$ , respectively;  $p=0.01$ ).

Table 1 reports the sociodemographic and medical characteristics of patients who took part in the survey. The subset of 287/325 respondents who completed these items had a mean age of  $53 \pm 12$  years and a majority were men, including 77% born in France. Half were living with their long-term partner. Nearly 66% were educated to university entry level or had been through higher education, 46% were in stable employment, and 46% were in precarious employment. They had been HIV-positive for  $17 \pm 10$  years and had been on antiretroviral treatment for  $14 \pm 8$  years, with 92% having undetectable viral load ( $<50$  copies/mL) and a mean CD4 T-cell count of  $620 \pm 375/\text{mm}^3$ . Furthermore, 45% were also on other associated treatments. Most saw their primary care physician one, two, or three times a year and their HIV specialist twice a year, and only 29% did not also see other specialist physicians. At the time the survey was completed, mean patient fitness status self-reported on a 0–100 scale was  $77 \pm 20$ .

Mean age of the physicians who took part in the survey was  $48 \pm 10$  years, and 57% were women. A large majority (72%) worked full-time at hospital, with 71% working in an infectious disease clinic.

## Questionnaire surveying PLHIV

### 1. Apps

A total of 9 out of 10 PLHIV (*i.e.*, 279/314; 89%) owned a smartphone and a little less than half (132/314; 42%) owned a tablet computer.

Stated opinions on mHealth apps by the 311 respondents were as follows: three-quarters (232; 75%) were comfortable with technology, 188 (60%) trusted the technology, 212 (68%) found the time to use the apps, and half (156; 50%) trusted their scientific value credentials, while 137 (44%) stated that using the apps helped them feel in tune with the times and a third (103; 33%) stated that the apps gave grounds to talk technology with friends. However, the dominant concerns were that their data would be used for marketing purposes (250; 81%) or by unauthorized people (241; 78%), and 74% did not want to share their data. Other concerns voiced by PLHIV were that the apps may become overly intrusive (173; 56%), may have hidden costs (119; 39%), or may become addictive (80; 26%).

Nearly a third of respondents (83/288; 29%) had already downloaded 'mHealth/wellness' apps (Tables 2 and 3). Compared to non-downloaders, this 'downloaders' subset was significantly younger ( $p < 0.001$ ) and more educated ( $p = 0.002$ ), spent more time in sex bars ( $p = 0.01$ ) and on geolocation-based dates ( $p < 0.001$ ), were active users of recreational drugs ( $p = 0.02$ ), and had spent less time HIV-positive ( $p = 0.002$ ) and on antiretroviral treatment ( $p = 0.02$ ). Multivariate analysis found that these downloaders were younger (OR  $0.96 \pm 0.01$ ,

$p=0.004$ ), educated to at least university entry level (OR 2.27  $\pm$ 0.86,  $p=0.03$ ), and more frequently used geolocation-based dating apps (OR 3.00  $\pm$ 1.09,  $p=0.002$ ) (Table 4).

At the time of the survey, only 10% (30/288) used wellness apps and 18% (52/288) used fitness tracking apps. Only 12 patients (4%) used health-related apps (HIV and/or others) to feel more confident in their health and have more firmly documented discussions with their physician (67%), or to improve their general health and be more self-managing (58%).

Only 9/314 (3%) PLHIV had already been advised on mobile apps by a healthcare professional, but the app functions and features failed to meet the needs of one third of them. However, 227/314 (72%) claimed they were ready to use an app recommended by a physician, far more than by a pharmacist (74/314; 24%) or a nonprofit organization (57/314; 18%).

A total of 60 PLHIV answered the question investigating the role that apps play or may play: 75% responded better tracking of their bodyweight, fitness and sleep, 68% responded better communication with their physician, 67% responded better self-surveillance of comorbidity, and 65% responded as a disease prevention tool, whereas 35% saw using apps as a way to reduce the number of appointments needed with physicians. Note that 46/60 (77%) were against sharing this data, while those who were for sharing this data would prefer to share with their physician (79%), pharmacist (50%), or a nurse (29%).

For around three quarters of the 310 respondents, the ideal app would serve primarily for monitoring vaccinations (235/310; 76%), general all-round health (234/310; 75%), and adverse drug effects (232/310; 75%), as well as for keeping track of blood test appointments (221/310; 71%) and as a record of treatments (210/310; 68%).

Respondents attached very little value to buying drugs and medication online — only 8% (24/310) of PLHIV had already bought over-the-counter products through online channels.

## 2. Smart devices

Although 236/306 PLHIV respondents had already heard about smart devices, only 66 (22%) owned one — in most cases a step counter, smart watch, and sleep quality tracker. Most had owned their device for more than a year, and 11% continued to use them today. These 66 respondents were younger ( $p=0.009$ ), had spent less time HIV-positive ( $p=0.003$ ), and more often used geolocation-based dating apps ( $p=0.003$ ) than the other 224 respondents. Of the 34 respondents that used these smart devices today, 65% reported that the devices helped them stay fit and be more efficient in their efforts to get fitter, but only 26% reported that they served for better-informed discussions with their physicians. The majority (88%) stated that they will continue using them unless they are paid-subscription services. Only 12% consulted their physician less often than recommended, and 70% of them would like to be able to send their data to their HIV specialist and/or primary care physician, against only 12% to their pharmacist, 15% to their basic health insurance, and 8% to their supplemental health insurance.

If they had to get mHealth-equipped, 201/306 (66%) PLHIV would need technical support and 215/306 (70%) would need medical support: either from their HIV specialist (204/233;88%) or their primary care physician (155/233; 67%) ahead of their pharmacist (70/233; 30%), a nurse (42/233; 18%), a purpose-dedicated website (25/233; 11%), or the health insurance system (26/233; 11%).

## Questionnaire surveying physicians

The physicians had ubiquitously adopted smartphones — nearly all of them (216/227; 95%) owned one. Practically all of these physicians used their smartphone to browse the Internet (214; 99%), and 212 of them (93%) had downloaded apps: 77 (34%) used wellness apps and 92 (41%) used fitness tracking apps. Half of them in total (117/227; 52%) used these apps, and this subsample had no differentiating sociodemographic characteristics.

Ninety-four (41%) physicians thought their patients used mHealth apps, but only 41 (18%) could give an idea of how their patients used the apps, and 24 of these 41 (56%) thought that less than 10% of their patients were actually health-app users.

According to the physicians' opinion, what patients look for most in an HIV support app is schedule management (94/227; 41%), social connectivity (65/227; 28%), peer testimonies (53/227; 23%), psychological well-being (30/227; 13%), and keeping track of adverse events (30/227; 13%), blood test results (23/227; 10%), and physical and psychological HIV-related parameters. A little short of half (96/277; 42%) were unable to answer this question.

However, for 165/227 (73%) physicians, apps were adjunctive tools, and for 127/227 (56%), apps empowered and informed patients to self-manage their infection. Only 18% thought apps could help strengthen the physician–patient relationship and only 24% thought apps could make patient encounters more efficient. Note that 209/227 (92%) physicians stated that the law was too lax on the collection and use of this data.

For 176/227 (75%) physicians, really useful apps would empower patients with better information and therapeutic education: for 40%, the apps would serve for clinical decision support and for 37% they would enable better monitoring, but only 46 (20%) thought it feasible to allow online data entry in the patient's health record.

A majority of the physicians (150/226; 66%) did not know whether their patients used smart sensors. Among just 24 who did know, the majority asserted that no more than 5% of patients used them. However, 137/225 (61%) physicians would welcome mainstream use of smart devices, as they ready their patients for the digital health revolution (71%) and educate their patients on how to sensibly use emerging technologies (53%). Those physicians who were against essentially cited data security issues (57/88; 65%) and information overload (49%) rather than issues tied to data confidentiality (40%), medical liability (26%), inadequate data (26%), or trustworthiness (17%).

## **Discussion**

Nearly a third of the 325 PLHIV respondents had already downloaded mHealth/wellness apps — including just 3% on the advice of their physician — but only 10% reported using the apps at the time of the survey. These active users were younger and more highly educated, and they already used apps socially. This 10% mHealth app user rate is the same as the rate estimated by the few physicians who claimed they knew the app. Only 22% of PLHIV surveyed owned at least one smart device, with no differences in sociodemographic characteristics to active app users, and 11% continued using them today. This converges with the numbers estimated by the 24/299 physicians who stated that they knew their patients' patterns of mHealth device use. On top of the gap between patient-reported practices and physician-reported perceptions, there is another gap between patient-reported needs and physician-reported needs: 76% of the 310 PLHIV stated that the ideal app would be a general health and vaccinations tracker for the purpose of better

communicating with their physician, whereas 42% of the 227 physicians surveyed said they had no idea how PLHIV would find use for an app.

A French survey led by Odoxa in 2015 reported that 29% of people regularly used mainstream smart devices and only 5% of patients had been recommended a connected technology solution by their physician [17]. We observed that PLHIV are timid mHealth users: they make disappointingly few downloads of HIV-related apps and are even turning away from using them. The early hope and promise of new health information technologies has given way to mistrust due to insecurity surrounding data confidentiality and personal data protection. The ‘innovative biotechnology packages’ that were designed to improve quality of life across the board had offered a ‘framework’ and ‘horizon’ for hope and a less uncertain future [18]. These researches fueled belief that life was improvable and promised to perform ‘miracles’ in the field of medicine — at a cost of desubjectivizing the body and disembodiment of the conscious subject. However, the surrounding economy selling the promise of a new ‘biotechnological regime’ is ultimately a ‘commerce and industry-driven’ vision grounded in a triangle of ‘market forces, self-determination and public health’ [18]. Connected medical devices (CMD) come with a legal obligation to protect all personal health data, but note that not all connected mobile apps and smart devices are qualified as CMD — the others are smart devices that have no stated medical purpose, for which France has nevertheless issued good-practice guidance to industry to ensure that personal data remains secured and reliable [14]. Furthermore, the sponsors of technoscience-driven promises are arguably in the business of selling hope and may well be overselling the possibilities of solving psychosocial-medical problems with technology solutions [19]. In this sense, their promises equate to speculative bubbles.

Are these app tools overly intrusive, as claimed by 56% of the PLHIV surveyed? On the one hand, without a firm command of the information produced, knowing the state of one's body is liable to generate stress and anxiety [20]; yet on the other hand, app-based 'self-healthcare' may create an illusion of some kind of body-health control. This self-led health assessment may give the illusion of escaping avoidable morbidity and premature death [21]. This biometric data could open the door to intrusive body-health control by service providers or even health insurance companies, which would have acute social and psychological repercussions [22]. Mathieu-Fritz & Guillot [23], in a paper on the 'patient work system', showed that the 'price to pay' for using self-care tools entails more work, more uncertainty, and — crucially — an 'experimental mindset' geared to testing the value and reliability of new devices.

The landmark market approval for the world's first digital pill — called Abilify, indicated to treat chronic psychosis — to track medication adherence raises tricky ethical questions [24]. On one side of the equation, carer responsibility dictates tight control of perfect medication adherence for the good of the patient. On the other side, the patient becomes dispossessed of their disease, with efficacy overriding patient autonomy since wider community principles dictate the provision of care for all and care to keep scarce health resources cost-effective, and so this digital system ultimately raises the issue of individual freedom.

General disinterest for HIV mHealth may also be induced by inadequate follow-up, gaps in counselling or gaps in personalized status analysis, due to a lack of appropriate health education. Our findings point to distortion in the end-purpose of mHealth apps. Note that the 310 PLHIV respondents stated that the ideal app would be a general health and vaccinations tracker for the purpose of better communicating with their physician, whereas

42% of the physicians surveyed said they had no idea how PLHIV would find use for an mHealth app. The other half of the physicians surveyed thought that their patients see utility in schedule management and for creating social exchange among peers. Less than one in five physicians thought an mHealth platform could strengthen the physician–patient relationship, as suggested in the EmERGE project [25]. Physicians today are routinely struggling with tight time schedules and poor working conditions, especially in hospitals, which puts the physician–patient relationship under strain [26-27]. These new connected devices thus need to find room and space in an already-crowded organization/ecosystem. Furthermore, published findings on the use of digital technologies in medical practice were long a source of controversy [28]. Many of the papers were not fit for analysis as they failed to meet robust methodological evaluation standards [29], and a majority of them failed to demonstrate any real medical service to patients, except for CMDs which enabled ‘live’ telemonitoring that brings significant benefits in heart failure prevention and diabetes management, chiefly through appropriate professional organizational-level service delivery and appropriately-trained professionals [30-32]. The disorganized use of smart health tech and the lack of cooperation between industry, patients, and healthcare providers to ensure data remains secure and reliable have since been addressed and trial methodologies have since been adapted, with encouraging results [33]. There are now some apps purpose-designed for PLHIV outreach, to support engagement in prevention and medication adherence, or to support continuity of care [34-36]. mHealth solutions could also address inequity in access to care for larger segments of the HIV-positive population in low-income countries. A study in South Africa found that mHealth could address the lack of communication in sharing patient health information between clinics, community health workers, and patients, as well as improve health information transfer to government officials

to better allocate healthcare resources, and maybe even enable better patient identification [37], yet nation-wide adoption of mHealth has still not occurred [38].

PLHIV used smart devices that were useful adjunctive tools for managing their care, but three quarters of them would like to send their data to their physician. This fits with the 61% of physicians who would welcome routine use of smart devices, which they see as a vector for delivering health education or a way to empower patient self-management, in line with the conclusions of the national evaluator of health apps and smart devices that certifies medical devices [13]. The main barriers to adoption cited by physicians arise from concerns over data security and information overload, as already signaled by Dr J Melchior from Louvain University in Belgium [39]. In terms of data sharing and confidentiality with 'health/wellness' apps that do not strictly host health data, a recent study surveying PLHIV captured the situation well [40]. The authors found that more and more community-member outreach workers are using in-community mobile apps and websites, which raises a number of ethical challenges on four fronts: i) managing personal and professional boundaries with clients; ii) disclosing personal or identifiable information to professionals; iii) maintaining client confidentiality and anonymity; iv) security and data storage measures for online information. Discussions under the EmERGE project surrounding network exchange platforms converged towards a critical need to implement the highest level of data security/privacy, as there is still strong stigmatization around HIV-positive people, and unlike when a bank account gets hacked, the social damage done is impossible to repair [25].

To conclude, mHealth apps and smart devices have failed to garner adoption by PLHIV. The reason appears to be a combination of marketing-related mistrust raising data security, confidentiality, and privacy fears, and an absence of adequate personalized monitoring and medical analysis — which physicians lack the time and perhaps even the motivation to

deliver. These barriers to uptake of mHealth apps and smart devices could be removed if physicians, with legislative backing and better hospital-side organization, engaged in promoting them. If app certification could become one of the roles and responsibilities of scholarly societies working in concert with patient advocacy groups, and if the lead 'eHealth' specialists could produce the requisite technical assistance, then data sharing and exchange could become adopted within a new conception of the HIV care pathway.

### **Disclosure of interest**

The authors declare no conflict of interests.

### **Acknowledgements**

CJ, PB, FL, ROT, and PD drafted the protocol, JP put the two surveys online, CL performed the statistical analysis, CJ interpreted the findings and drafted the article, and FL, ROT, and PD reread and reviewed the draft.

The surveys were organized, coordinated, and delivered on-site with help from the following physicians and CRTs:

C. Faudon, T. Allegre **Aix-en-Provence**; M. Lombard, JL. Schmit **Amiens**; S. Rehaïem, V. Rabier **Angers**; A. Valran, E. Piet **Annecy**; J. Gerbe, F. Caby **Argenteuil**; E. Chevalier, A. Foltzer **Besancon**; L. Traore, O. Bouchaud **Bobigny**; JC. Duthe, S. Ansart, L. de Saint Martin **Brest**; P. Goubin, R. Verdon **Caen**; J. Prouteau, C. Jacomet **Clermont-Ferrand**; C. Chesnel, L. Richier, S. Gallien, B. Elharrar **Créteil**; I. Calmont, A. Cabié **Fort-de-France**; M. Marcou, P. de Truchis **Garches**; C. Sautron, MP. Moiton **La Réunion Saint-Denis**; L. Lainé, S. Leautez-Nainville **La**

**Roche-sur-Yon**; P. Camps, M. Roncato-Saberan **La Rochelle**; D. Bornarel, A Greder-Belan **Le Chesnay**; AS. Keita, A. Cheret **Le Kremlin Bicêtre**; F. Dangeul-Potier, H. Hitoto **Le Mans**; B. Montoya, JJ. Laurichesse **Mantes-la-Jolie**; S. Tassi, E. Frogel **Marne-la-Vallée**; A. Suzel-Ritleng, O. Zaegel-Faucher **Marseille Sainte Marguerite**; C. Fernandez, J. Reynes **Montpellier**; M. Benomar, G. Beck-Wirth **Mulhouse**; A. Soria, F. Raffi **Nantes**; I. Touitou, P. Pugliese **Nice**; P. Gougeon, S. Sunder **Niort**; MJ. Dulucq, L. Slama **Paris Hôtel-Dieu**; C. Louisin, C. Duvivier, PH. Consigny **Paris Necker Institut Pasteur**; M. Pauchard, D. Beniken, C. Katlama, A. Simon **Paris Pitié-Salpêtrière**; C. Tran, D. Bollens **Paris Saint-Antoine**; A. Adda Lievin, G. Thibaut, P. Pialoux **Paris Tenon**; L. Pinheiro, H. Aumaitre **Perpignan**; L. Gerard, H. Masson **Poissy**; D. Plainchamp, G. Lemoal **Poitiers**; JC. Duthe, P. Perfezou **Quimper**; I. Kmiec, F. Bani-Sadr **Reims**; C. Guennoun, C. Arvieux **Rennes**; G. Unal, A. Lesourd **Rouen**; V. Ronat, A. Fresard **Saint-Etienne**; B. Montoya, Y. Welker **Saint-Germain-en-Laye**; C. Grand Courault, C. Michau **Saint-Nazaire**; P. Fisher, D. Rey **Strasbourg**; V. Lambry, A. Lafeuillade **Toulon**; M. Digumber, O. Robineau **Tourcoing**; E. Racamier, H. Champagne **Valence**; L. Richier, S. Dellion **Villeneuve-Saint-Georges**.

**Funding:** none

## References

1. OpinionWay. Les Français et les objets connectés. Distree Connect. Mars 2017. 16 pages. <https://www.opinion-way.com/fr/sondage-d-opinion/sondages-publies/marketing/opinionway-pour-distreeconnect-2017-les-francais-et-les-objets-econnectes-mars-2017.html> (accessed on March 31, 2020)
2. Leclerc E, Vanryp B. Baromètre de l'innovation. Sondage BVA pour Syntec numérique. 12 mars 2014. 17 pages . [https://syntec-numerique.fr/sites/default/files/Documents/barometre\\_de\\_linnovation\\_sn\\_bva\\_cloud\\_-\\_moral\\_editeurs\\_mars2014\\_0.pdf](https://syntec-numerique.fr/sites/default/files/Documents/barometre_de_linnovation_sn_bva_cloud_-_moral_editeurs_mars2014_0.pdf) (accessed on March 31, 2020)
3. Research 2 Guidance. Current Status and Trends of the mHealth App Market. mHealth Economics 2016. 28 pages. <https://research2guidance.com/product/mhealth-app-developer-economics-2016/> (accessed on March 31, 2020)
4. "Quantified Self". Revue Réseaux, Éditions La Découverte. 2019, 4 (216): 276 pages.
5. Jacomet C, Berland P, Guiguet M, Simon A, Rey D, Arvieux C, Pugliese P, Gerbaud L. Impact of age on care pathways of people living with HIV followed up in hospital. *AIDS Care*. 2017;29(1):105-111.
6. Morlat P. Prise en charge médicale des personnes vivant avec le VIH – Recommandations du groupe d'experts. 6 aout 2019. <https://cns.sante.fr/actualites/prise-en-charge-du-vih-recommandations-du-groupe-dexperts/> (accessed on March 31, 2020)
7. Use of smartphone application advertising for HIV prevention among men who have sex with men in the European Union/European Economic Area. ECDC 2017. <https://www.ecdc.europa.eu/sites/default/files/documents/Smartphone-apps-HIV-prevention-2017.pdf> (accessed on March 31, 2020)
8. Pant pai N. HIVSmart! An award-winning smartphone application that allows self-testing for HIV. <http://nitikapantpai.com/hivsmart/> (accessed on March 31, 2020)
9. Home HIV Test APK. Amazing finger home HIV test prank to find about HIV symptoms. <https://www.apktouch.com/apps/com.RaholjiFingerprintApps.FingerHomeHIVTestPrank.HIVSymptoms.EarlySymptomsOfHIV> (accessed on March 31, 2020)
10. HIV/ AIDS test. Apps on google Play. [https://play.google.com/store/apps/details?id=yakovenko.hivsymptoms&hl=en\\_US](https://play.google.com/store/apps/details?id=yakovenko.hivsymptoms&hl=en_US) (accessed on March 31, 2020)
11. Seronet. L'info en +. <https://www.seronet.info/> (accessed on March 31, 2020)

12. Muessig KE, Nekkanti M, Bauermesister J, Bull S, Hightow-Weidam LB. A systematic review of recent smartphone, Internet and Web 2.0 intervention to address the HIV continuum of care. *Curr HIV/AIDS Rep* 2015; 12(1): 173-190
13. Santo K, Richtering SS, Chalmers J, Thiagalingam A, Chow CK, Redfern J. Mobile phone apps to improve medication adherence: a systematic stepwise process to identify high quality apps. *JMIR Mhealth Uhealth* 2016; 4 (4): e132
14. Haute Autorité de santé. Référentiel de bonnes pratiques sur les applications et les objets connectés en santé (MobileHealth ou mHealth). Evaluation et amélioration des pratiques. Octobre 2016. 60 p.  
[https://www.has-sante.fr/upload/docs/application/pdf/2016-11/has\\_ref\\_apps\\_oc.pdf](https://www.has-sante.fr/upload/docs/application/pdf/2016-11/has_ref_apps_oc.pdf)  
(accessed on May 3, 2019).
15. Swendeman D, Farmer S, Mindry D, Lee S.J, Medich M. HIV care providers' attitudes regarding mobile phone applications and web-based dashboards to support patient self-management and care coordination: results from a qualitative feasibility study. *J HIV AIDS* 2016; 2(4).
16. Jacomet C, Bastiani P, Prouteau J, Lambert C, Linard F, Ologeanu-Taddei R, et al. E-santé. E-health. Patterns of use and perceived benefits and barriers among people living with HIV and their doctors. Part 1 : Internet for information retrieval and social network. *Med Mal Inf.* In press.
17. Sliman G. Baromètre santé 360. La santé connectée. Odoxa. Janvier 2015. 20p.  
<http://www.odoxa.fr/sondage/barometre-sante-360-sante-connectee/>
18. Annette Leibing, Virginie Tournay (dir.), *Les technologies de l'espoir. La fabrique d'une histoire à accomplir*, Presses de l'Université Laval, coll. « Sociétés, cultures et santé », 2010, 302 p., EAN : 9782763789958
19. Joly, P.B. (2015). Le régime des promesses technoscientifiques. In Audétat, M. *Pourquoi tant de promesses*, Paris : Hermann, pp.31-48
20. Association française des correspondants à la protection des données à caractère personnel (AFCDP). Quantified self connecté et Informatique et Libertés 2015.  
[https://www.researchgate.net/publication/288825232\\_Quantified\\_Self\\_connecte\\_et\\_Informatique\\_Libertes](https://www.researchgate.net/publication/288825232_Quantified_Self_connecte_et_Informatique_Libertes) (accessed on May 3, 2019)
21. Andrieu B. L'osmose technique avec son corps viv@ant : une auto-santé connectée du patient immersif. *L'évolution psychiatrique* 2016; 8: 1253-65.

22. Vallancien G. la médecine sans médecin? Le numérique au service du malade. Paris: Gallimard 2015.
23. Mathieu-Fritz A, Guillot C. Les dispositifs d'autosurveillance du diabète et les transformations du « travail du patient ». Les nouvelles formes de temporalité, de réflexivité et de connaissance de soi liées à l'expérience de la maladie chronique. *Revue d'anthropologie des connaissances*. 2017; 4 (11): 641-675.
24. Cosgrove L, Cristea IA, Shaughnessy AF , Mintzes B, naudet F. Digital aripiprazole or digital evergreening? A systematic review of the evidence and its dissemination in the scientific literature and in the media. *BMJ Evid Based Med* 2019 Dec 24 (6): 231-238.
25. Marent B, henwood F, Darking M, EmERGE Consortium. Development of an mHealth platform for HIV Care : gathering user perspectives through co-design workshops and interviews. *JMIRmhealth and Uhealth* 2018 ;6 (10) : e184.
26. Raveyre M, Ughetto P. « On est toujours dans l'urgence » : surcroît ou défaut d'organisation dans le sentiment d'intensification du travail ? Colloque «Organisation, intensité du travail, qualité du travail ». Centre d'études de l'emploi, CEPREMAP et LATTTS Paris, 21-23 novembre 2002.
27. Gheorghiu M D, Moatty F, L'hôpital en mouvement. Changements organisationnels et conditions de travail, Éditions Liaisons, coll. « Liaisons Sociales », 2013, 303 p., ISBN : 978-2-87880-976-3.
28. P. Simon. Les leçons apprises des principales études sur les objets connectés en télémédecine et santé mobile. *European Research in Telemedecine* 2017; 6: 67-77.
29. Steventon A, Bardsley M, Billings J, Dixon J, Doll H, Hirani S et al. Whole System Demonstrator Evaluation Team. Effects of telehealth on use of secondary care and mortality: findings from the WSD cluster randomized trial. *BMJ* 2012; 344: 3874-92
30. Ahmed S, Ernst P, Bartlett SJ, Valois MF, Zaihra T, Paré G et al. The effectiveness of web-based asthma self-management system. My Asthma portal (MAP): a pilot randomized controlled trial. *J med Internet Res* 2016; 18 (12): e13
31. Kitsiou S, Pare G, Jaan M. Effects of home telemonitoring interventions on patients with chronic heart failure: an overview of systematic reviews. *J Med Internet Res* 2015; 17(3): e63
32. Charpentier G, Benhamou PY, Dardari D, Clergeot A, Franc S, Schaepepelynk-Belicar P et al, TeleDiab Study Group. The diabeo software enabling individualized insulin dose adjustments combined with telemedicine support improves HBA1C in poorly controlled

- type 1 diabetic patients: a 6-month, randomized, open label, a parallel-group, multicenter trial (ReleDiab1 Study). *Diabetes Care* 2011; 34(3): 533-9.
33. Denis F, Basch E, Septans AL, Bennouna J, Urban T, Dueck AC, et al. Two-year survival comparing web-based symptom monitoring vs routine surveillance following treatment for lung cancer. *JAMA*. 2019;321(3) :306–7.
  34. Milam J, Jain S, Daar E, Dube M, Seefreid E, Ellorin E, et al. Controlled Trial of an Internet-Based Risk Reduction Intervention in HIV+ Men Who Have Sex With Men. Conference on Retroviruses and Opportunistic Infections (CROI); March 3–6, 2014; Boston, MA. Poster #943, [https://www.croiconference.org/sites/default/files/uploads/CROI2014\\_Final\\_Abstracts.pdf](https://www.croiconference.org/sites/default/files/uploads/CROI2014_Final_Abstracts.pdf) (accessed on May 3, 2019)
  35. Miranda J, Cote J, Godin G, Blais M, Otis J, Gueheneuc YG, et al. An Internet-Based Intervention (Condom-Him) to Increase Condom Use Among HIV-Positive Men Who Have Sex with Men: Protocol for a Randomized Controlled Trial. *JMIR Res Protoc*. 2013; 2: e39
  36. Hightow-Weidman L, Muessig K, Knudtson K, Srivatsa M, Lawrence H, LeGrand S, et al. A gamified smartphone app to support engagement in care and medication adherence of HIV positive young men who have sex with men (Allyquest): development and pilot study. *JMIR Public Health Survey* 2018; 4 (2): e34.
  37. Van Heerden A, Harris M D, Van Rooyen H, Barnabas R, Ramanathan N, Ngcobo N, et al. Perceived mHealth barriers and benefit for home-based HIV testing and counseling and other care: qualitative findings from health officials, community health workers and persons living with HIV in South Africa. *Soc Sci Med* 2017; 183: 97-105.
  38. Krah EF, de Kruijff JG. Exploring the ambivalence evidence base of mobile health (mhealth): A systematic literature review on the use of mobile phones for the improvement of community health in Africa. *Digital health*. 2016; 2: 1-20.
  39. J Melchior, F Ndiemo. De la médecine 1.0 à la médecine 3.0 : le patient connecté. Décembre 2016. <https://www.louvainmedical.be/fr/article/de-la-medecine-10-la-medecine-30-le-patient-connecte> (accessed on May 3, 2019)
  40. Fantus S, Souleymanov R, Lachowsky NJ, Brennan DJ. The emergence of ethical issues in the provision of online sexual health outreach for gay, bisexual, two-spirit and other

men who have sex with men: perspectives of online outreach workers. BMC Med Ethics [Internet] 2017; 18

**Table 1.** Sociodemographic and medical characteristics of patients (N=287)

<b>Variable</b>		<b>n (%)</b>  <b>Mean ± SD</b>
Age	Years	53 ±12
Gender	Male	204 (71)
	Female	82 (29)
	Transgender	1 (0.4)
Life situation	Long-term relationship	145 (51)
Sexual orientation	Heterosexual	130 (45)
	Homosexual	123 (43)
	Other	12 (4)
Children	At least one	121 (42)
Country of birth	Born in France	220 (77)
Department of birth <sup>1</sup>	Ile-de-France	55 (25)
Department of residency	Ile-de-France	91 (32)
Educational attainment	Baccalaureate (university entry qualification) or higher	190 (66)

Occupational status	Stable employment	133 (46)
	Retired	61 (21)
	Incapacity	36 (13)
	Jobseeker	32 (11)
Precarious employment	EPICES score	31 ±23
	Non-precarius	153 (54)
Meeting places	Bars-clubs (non-sex-oriented)	81 (28)
	Sex clubs	40 (14)
	Online	61 (21)
Last HIV viral load measure	Undetectable viral load	262 (92)
Last CD4 cell count	/mm <sup>3</sup>	620±375
Time to HIV test	Years	17 ±10
Time on antiretroviral treatment	Years	14 ±8
Smoking	Yes	78 (28)
	Ex-smoker	64 (22)
Alcohol use	More than once a week	137 (48)
Recreational drug use	Yes	39 (14)
	Ex-user	17 (6)
Lipodystrophy	Presence	58 (20)

Other associated treatments	Presence	129 (45)
	Antihypertensive	59 (21)
	Psychiatric help	45 (16)
	Cardiovascular	29 (10)
	Antidiabetic	26 (9)
	Hyperlipidemia	16 (6)
	Osteoarticular	16 (6)
	Neurological	13 (5)
	Hepatitis B or C	9 (3)
	Renal	8 (3)
	Cancer	5 (2)
Monitoring	In-hospital	246 (86)
Primary care physician	None	40 (14)
	One, two, or three	168 (59)
	Four or more	78 (27)
Consultation with an HIV specialist	One or two over the year	165 (57)
	Three or more	122 (43)
Consultations with other specialists	None	82 (29)
	One, two, or three	155 (54)
	Four or more	50 (17)
Self-reported patient fitness status <sup>2</sup>	0–100	77 ±20

<sup>1</sup> 220 respondents

<sup>2</sup> 325 respondents

**Table 2.** Sociodemographic characteristics of people living with HIV who have already downloaded health/wellbeing apps (N=83) versus others (N=205)

<b>Variables</b>		<b>n (%)</b> <b>Mean ±SD</b>	<b>Downloaders</b>	<b>Non-downloaders</b>	<b>p-value</b>
Age <sup>1</sup>	Years	52 ±12	48 ±13	54 ±11	<0.001
Gender <sup>1</sup>	Male	188 (71)	55 (73)	133 (72)	0.38
	Female	73 (28)	20 (26)	53 (28)	
	Transgender	1 (1)	1 (1)	0	
Life situation <sup>1</sup>	Long-term relationship	133 (51)	38 (50)	95 (51)	0.87
Sexual orientation <sup>1</sup>	Heterosexual	118 (45)	28 (37)	90 (48)	0.16
	Homosexual	114 (44)	40 (53)	74 (40)	
	Other/Refuse to identify	30 (11)	8 (10)	22 (12)	
Number of children <sup>2</sup>	At least one child	108 (42)	26 (35)	82 (44)	0.15
Country of birth <sup>1</sup>	Born in France	200 (76)	56 (74)	144 (77)	0.52
Department of birth <sup>1</sup>	Ile-de-France	50 (25)	17 (30)	33 (23)	0.28
Department of residency <sup>1</sup>	Ile-de-France	85 (32)	29 (38)	56 (30)	0.21

Educational attainment <sup>1</sup>	Baccalaureate (university entry qualification) or higher	177 (68)	62 (82)	115 (62)	0.02
Occupational status <sup>1</sup>	Stable employment	124 (47)	41 (54)	83 (45)	0.20
	Retired	54 (21)	13 (17)	41 (22)	
	Incapacity	32 (12)	5 (7)	27 (15)	
	Jobseeker	28 (11)	11 (14)	17 (9)	
Precarious existence <sup>3</sup>	EPICES score	31 ±23	28 ±24	32 ±22	0.14
	Non-precarious	143 (55)	48 (63)	95 (51)	0.08
Meeting places <sup>1</sup>	Bars-clubs (non-sex-oriented)	77 (29)	28 (37)	49 (26)	0.09
	Sex clubs	39 (15)	18 (24)	21 (11)	0.01
	Online	59 (23)	32 (42)	27 (15)	<0.001

<sup>1</sup> 262 respondents (76 Downloaders, 186 Non-downloaders)

<sup>2</sup> 260 respondents (75 Downloaders, 185 Non-downloaders)

<sup>3</sup> 261 respondents (76 Downloaders, 185 Non-downloaders)

**Table 3.** Medical characteristics of people living with HIV who have already downloaded health/wellbeing apps (N=83) versus others (N=205)

Variables		n (%) Mean $\pm$ SD	Downloaders	Non-downloaders	p-value
Last HIV viral load measure <sup>3</sup>	Undetectable viral load	238 (91)	70 (92)	168 (91)	0.74
Last CD4 cell count <sup>4</sup>	/mm <sup>3</sup>	613 $\pm$ 366	620 $\pm$ 352	610 $\pm$ 373	0.28
Time to HIV test <sup>5</sup>	Years	17 $\pm$ 10	14 $\pm$ 11	18 $\pm$ 10	0.002
Time on antiretroviral treatment <sup>5</sup>	Years	14 $\pm$ 8	11 $\pm$ 8	15 $\pm$ 8	0.002
Smoking <sup>1</sup>	Yes	70 (27)	23 (30)	47 (25)	0.54
	Ex-smoker	61 (23)	19 (25)	42 (26)	
Alcohol use <sup>1</sup>	Once or more a week	126 (48)	36 (47)	90 (48)	0.88
Recreational drug use <sup>1</sup>	Yes	37 (14)	15 (20)	22 (12)	0.02
	Ex-user	15 (6)	8 (11)	7 (4)	
Lipodystrophy <sup>5</sup>	Presence	55 (21)	17 (23)	38 (21)	0.76
	Presence	111 (42)	29 (38)	82 (44)	0.38

Other associated treatments <sup>1</sup>	Antihypertensive	50 (19)	12 (16)	38 (20)	0.39
	Psychiatric help	38 (15)	13 (17)	25 (13)	0.45
	Cardiovascular	23 (9)	4 (5)	19 (10)	0.20
	Antidiabetic	22 (8)	6 (8)	16 (9)	0.85
	Hyperlipidemia	13 (5)	4 (5)	9 (5)	1.00
	Osteoarticular	13 (5)	3 (4)	10 (5)	0.76
	Neurological	11 (4)	3 (4)	8 (4)	0.90
	Hepatitis B or C	8 (3)	4 (5)	4 (2)	0.24
	Renal	6 (2)	1 (1)	5 (3)	0.68
	Cancer	5 (2)	1 (1)	4 (2)	1.00
Monitoring <sup>1</sup>	In-hospital	226 (86)	64 (84)	162 (87)	0.54
Primary care physician <sup>1</sup>	None	36 (14)	6 (8)	30 (16)	0.10
	One, two, or three	158 (60)	45 (59)	113 (61)	
	Four or more	68 (26)	25 (33)	43 (23)	
Consultations with an HIV physician <sup>1</sup>	One or two over the year	152 (58)	47 (62)	105 (56)	0.42
	Three or more	110 (42)	29 (38)	81 (44)	
Consultations with other specialists <sup>1</sup>	None	76 (29)	25 (33)	51 (27)	0.22
	One, two, or three	142 (54)	35 (46)	107 (58)	
	Four or more	44 (17)	16 (21)	28 (15)	

<sup>1</sup> 262 respondents (76 Downloaders, 186 Non-downloaders)

<sup>3</sup> 261 respondents (76 Downloaders, 185 Non-downloaders)

<sup>4</sup> 257 respondents (76 Downloaders, 181 Non-downloaders)

<sup>5</sup> 260 respondents (76 Downloaders, 184 Non-downloaders)

**Table 4.** Characteristics of people living with HIV who have already downloaded health/wellbeing apps (multivariate analysis)

Patients who have already downloaded mHealth/wellness apps	OR [95% CI]	p-value
Homosexuality (vs heterosexuality)	1.14 ±0.48 [0.51–2.58]	NS
Refuses to identify their sexual orientation (vs heterosexuality)	1.32 ±0.75 [0.43–4.01]	NS
Age	0.96 ±0.01 [0.94–0.99]	0.004
At least one child	1.35 ±0.53 [0.63–2.90]	NS
Resident of the Ile-de-France	0.71 ±0.24 [0.37–1.38]	NS
Baccalaureate (university qualification) or higher	2.27 ±0.86 [1.08–4.77]	0.031
Precarious employment	0.60 ±0.19	NS

	[0.32–1.12]	
Use of geolocation-based dating apps	3.00 ±1.09 [1.48–6.12]	0.002
One, two, or three consultations with their primary care physician (vs none)	1.99 ±1.07 [0.69–5.73]	NS
Four consultations with their primary care physician (vs none)	3.86 ±2.23 [1.24–11.99]	0.019