



HAL
open science

Comparative analysis of predictive values of the kinetics of 11 circulating miRNAs and of CA125 in ovarian cancer during first line treatment (a GINECO study)

Patrick Robelin, Michel Tod, Olivier Colomban, Joel Lachuer, Isabelle Ray-Coquard, Gaëtan De Rauglaudre, Florence Joly, Annick Chevalier-Place, Pierre Combe, Alain Lortholary, et al.

► To cite this version:

Patrick Robelin, Michel Tod, Olivier Colomban, Joel Lachuer, Isabelle Ray-Coquard, et al.. Comparative analysis of predictive values of the kinetics of 11 circulating miRNAs and of CA125 in ovarian cancer during first line treatment (a GINECO study). *Gynecologic Oncology*, 2020, 159, pp.256 - 263. 10.1016/j.ygyno.2020.07.021 . hal-03491450

HAL Id: hal-03491450

<https://hal.science/hal-03491450v1>

Submitted on 17 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Comparative analysis of predictive values of the kinetics of 11 circulating miRNAs and**
2 **of CA125 in ovarian cancer during first line treatment (a GINECO study).**

3

4 Patrick ROBELIN^{1,2}, Michel TOD^{1,3}, Olivier COLOMBAN¹, Joel LACHUER⁴, Isabelle
5 RAY-COQUARD⁵, Gaëtan DE RAUGLAUDRE⁶, Florence JOLY⁷, Annick CHEVALIER-
6 PLACE⁸, Pierre COMBE⁹, Alain LORTHOLARY¹⁰, Salima HAMIZI¹¹, Nadia RABAN¹²,
7 Gwénaél FERRON¹³, Jérôme MEUNIER¹⁴, Dominique BERTON-RIGAUD¹⁵, Jérôme
8 ALEXANDRE¹⁶, Marie-Christine KAMINSKY¹⁷, Coraline DUBOT¹⁸, Alexandra LEARY¹⁹,
9 Emmanuelle MALAURIE²⁰, Benoit YOU^{1,2}

10 1. EMR3738, Ciblage Thérapeutique en Oncologie, Faculté de Médecine et de
11 Maïeutique Lyon-Sud Charles Mérieux, Université Claude Bernard Lyon 1,
12 Oullins, France ; Université de Lyon, Lyon, France.

13 2. Service d'Oncologie Médicale, CITOHL, Centre Hospitalo-Universitaire Lyon-
14 Sud, Institut de Cancérologie des Hospices Civils de Lyon (IC-HCL), Pierre-
15 Bénite, France ; GINECO, Paris, France

16 3. Pharmacie, Hôpital de la Croix Rousse, Hospices civils de Lyon, Lyon, France.

17 4. ProfileXpert, SFR-Est, CNRS UMR-S3453, INSERM US7, Lyon, France.

18 5. Centre Léon Bérard, Cancérologie médicale - 28 Prom. Léa et Napoléon
19 Bullukian, 69008 Lyon Université Claude Bernard Lyon 1, GINECO, Paris,
20 France

21 6. Institut Sainte-Catherine, Oncologie radiothérapie, 250 Chemin De Baigne Pieds
22 Cs80005, 84918 Avignon ; GINECO, Paris, France

23 7. Centre François Baclesse, Oncologie Médicale, 13 Rue du Général Harris
24 14000 Caen ; Université Unicaen, GINECO, Paris, France

25 8. Centre Oscar Lambret, cancérologie gynécologique, 3 Frédéric COMBEMALE
26 59020 Lille ; GINECO, Paris, France

27 9. Hôpital Européen Georges Pompidou, Oncologie Médicale, 20 Rue Leblanc,
28 75015 Paris ; GINECO, Paris, France

- 29 10. Hôpital Privé du Confluent, Oncologie médicale, 4 rue Eric Tabarly 44200 Nantes
30 ; GINECO, Paris, France
- 31 11. Centre Hospitalier Lyon Sud, Oncologie médicale, 165 Chemin Du Grand Revoyet
32 (CH LYON SUD), 69310 Pierre Benite ; GINECO, Paris, France
- 33 12. Hôpital de la Milétrie, Centre Hospitalier Universitaire de Poitiers, Oncologie
34 médicale, 2 Rue De La Miletrie 86000 Poitiers ; GINECO, Paris, France
- 35 13. Institut Claudius Regaud, Service de chirurgie oncologique; 1 avenue Irène Joliot-
36 Curie; 31059 Toulouse CEDEX 09; GINECO, Paris, France
- 37 14. Centre Hospitalier Régional d'Orléans, Oncologie médicale ,14 Avenue De
38 L'Hôpital, 45100 Orléans; GINECO, Paris, France
- 39 15. ICO site René Gauducheau, Oncologie médicale, Boulevard Jacques Monod,
40 44800 Saint Herblain; GINECO, France
- 41 16. Hôpital Cochin, Cancérologie oncologie, 27, Rue du Faubourg Saint Jacques
42 75014 Paris ; GINECO, Paris, France
- 43 17. Institut de cancérologie de lorraine, Alexis Vautrin, Oncologie médicale, 6 avenue
44 de bourgogne 54500 Vandœuvre-lès-Nancy; GINECO, Paris, France
- 45 18. Centre RENE HUGUENIN INSTITUT CURIE, Oncologie médicale, 35 Rue
46 Dailly, 92210 Saint Cloud; GINECO, Paris, France
- 47 19. Institut Gustave Roussy, Département de médecine, 114 rue Edouard Vaillant
48 94805 VILLEJUIF ; GINECO, Paris, France
- 49 20. Centre Hospitalier Intercommunal de Créteil, Oncologie Médicale et
50 Radiothérapie, 40 Avenue de Verdun, 94000 Créteil ; GINECO, Paris, France

51 The authors declare no potential conflicts of interest.

52 **Running title:** Kinetics of miRNAs and of CA125 in ovarian cancer

53 **Corresponding author:**

54 Dr Patrick Robelin,
55 Service d'Oncologie Médicale, Centre Hospitalo-Universitaire Lyon-Sud
56 EMR UCBL/HCL 3738,
57 Chemin du Grand Revoyet, 69495 Pierre-Benite, France.
58 Tel: +33 4 78 86 43 18; E-mail: patrick.robelin@chu-lyon.fr

59

60 ***Electronic word count:***

61 Abstract: 243 words

62 Manuscript (excluding title page, abstract, translational relevance, references, tables, legends
63 to figures): 3270 words

64 2 Figures, 4 Tables, 7 Supplementary Figures

65 Number of references: 36

66 Abstract

67 **Objective:** MicroRNAs (miRNAs) are promising biomarkers in ovarian cancer. Their kinetics
68 during treatment might be useful for monitoring disease burden, and guiding treatments in
69 patients treated with peri-operative chemotherapy and interval debulking surgery (IDS).

70 **Methods:** Serial blood samples of patients enrolled in the randomized phase II CHIVA trial,
71 comparing first line carboplatin-paclitaxel +/- nintedanib (NCT01583322) and IDS, were
72 investigated to assess the kinetics of 11 relevant miRNAs. Their prognostic/predictive values
73 regarding the likelihood of complete IDS, and the patient survival, were assessed and
74 compared to those of CA125 kinetics. The selection of the miRNAs (miR-15b-5p, miR-16-5p,
75 miR-20a-5p, miR-21-5p, miR-93-5p, miR-122-5p, miR-150-5p, miR-195-5p, miR-200b-3p,
76 miR-148b-5p and miR-34a-5p) was based on the expression levels found with a large
77 explorative panel, and on the literature data.

78 **Results:** 756 serial blood samples from 119 patients were analyzed for a total of 8172 miRNA
79 assays, and 1299 CA125 values. The longitudinal kinetics of the miRNA expressions were
80 highly inconsistent, and were not related to CA125 dynamics. The miRNA changes during
81 neoadjuvant treatment were not found associated with RECIST tumor response or IDS
82 outcomes. Decreases of miR-34a-5p and miR-93-5p were associated with PFS benefit
83 ($p=0.009$) and OS benefits ($p<0.001$), respectively, using univariate tests.

84 **Conclusions:** The longitudinal kinetics of miRNA expressions during neo-adjuvant treatment
85 in ovarian cancer patients were inconsistent, and were not found to be associated with tumor
86 burden changes. Although some prognostic value could be discussed, no predictive value
87 regarding tumor responses or IDS quality could be identified.

88 Introduction

89 Ovarian cancer ranks fifth for mortality linked to female cancer in western countries [1]. In
90 2011, the Gynecologic Cancer Intergroup (GCIG) defined CA125 response as a $\geq 50\%$
91 reduction in CA125 levels from baseline, on a minimum 28 day period [2]. However, the
92 relevance of this specific criterion, has recently been questioned, since it was not found to
93 exhibit prognostic or predictive values regarding progression free survival (PFS) and overall
94 survival (OS) in several studies [3–6].

95 In first line setting, stage III or IV ovarian cancer patients with diseases considered to be not
96 amenable to primary cytoreductive surgery are usually treated with neoadjuvant
97 chemotherapy and interval debulking surgery (IDS) [7]. One of the main prognostic factors in
98 patients treated with such a strategy is the completeness of the cytoreductive surgery
99 (complete CC0 versus incomplete CC1-CC2) [8]. In this context, there are no validated
100 biomarkers prone to help predict the likelihood of complete IDS, or patient survival.

101 MicroRNAs (miRNAs) are small non-coding RNA molecules that bind specifically to their
102 target mRNAs for inhibiting the expression of genes after transcription [9]. Several studies
103 reported promising outcomes about the diagnostic [10–14], prognostic [10,12,13,15] or
104 predictive values [16–18] of miRNAs in ovarian cancer patients. Some data suggested that the
105 tumor burden changes observed during treatment correlated with the kinetics of some
106 circulating miRNAs in ovarian cancers patients [17,18].

107 We hypothesized that the kinetics of some miRNAs of interest may be more helpful than
108 CA125 for monitoring disease burden and guiding clinicians in decision making for ovarian
109 cancers patients. The aim of the present study was designed to assess the kinetics of 11
110 relevant miRNAs and of CA125 in first line treatment, as a way of comparing their prognostic
111 and predictive values regarding the likelihood of complete IDS.

112

113 Materials and methods

114 Patient population

115 Clinical and biological data were derived from the randomized phase II trial CHIVA
116 (NCT01583322). In this trial, patients with International Federation of Gynecology and
117 Obstetrics (FIGO) stage IIC or IV ovarian carcinomas planned to be treated with peri-
118 operative chemotherapy and IDS, were randomly allocated to 1) the standard arm carboplatin
119 AUC 5 or 6 and paclitaxel 175mg/m² every 3 weeks with placebo, or 2) the experimental arm
120 with the same regimen combined to nintedanib at 200mg twice daily. The enrolled patients
121 received 3 to 4 cycles before cytoreductive surgery, and then 3 to 4 cycles after surgery,
122 followed by a maintenance treatment with nintedanib/placebo for up to 2 years [19].

123 Nintedanib (BIBF 1120, VARGATEFTM) is a receptor tyrosine kinase (RTK) inhibitor with
124 potential antiangiogenic and antineoplastic activities. The multitargeted tyrosine kinase
125 inhibitor BIBF 1120 selectively binds to and inhibits vascular endothelial growth factor
126 receptor (VEGFR), fibroblast growth factor receptor (FGFR) and platelet-derived growth
127 factor receptor (PDGFR) tyrosine kinases, which may result in the induction of endothelial
128 cell apoptosis; a reduction in tumor vasculature; and the inhibition of tumor cell proliferation
129 and migration.

130 Blood samples for miRNA, along with CA125 concentrations, were assayed at the 8
131 following times during the study: at screening, cycles 2, and 3; before IDS; the three cycles
132 after IDS; and at progression, if any. Moreover, the CA125 values were assayed every 3
133 months after the end of chemotherapy for follow-up. The following parameters were collected
134 for the present study: age, weight, height, pathology type, grade, FIGO stage, treatment arm,
135 post-operative disease residuals as judged by the surgeon (no macroscopic residual, tumor

136 residuals < 0.25cm, tumor residuals 0.25 to 2.5 cm, tumor residuals > 2.5 cm), and the
137 radiological tumor objective response to treatment according to RECIST V1.1 criteria
138 (complete response (CR), partial response (PR), stable disease (SD) or progressive disease
139 (PD)).

140 The protocol was approved by an ethic committee and health authorities in 2012. Written
141 informed consent was obtained from all patients before study entry.

142 Serum miRNAs quantification and analysis

143 The plasma samples were centrifuged, and stored at -80°C. MiRNAs were extracted with the
144 Qiagen miRNeasy microkit. The targets miRNAs were quantified by qRT-PCR using the
145 Qiagen kit custom miScript miRNA PCR assay. After performing the Qiazol lysis step, *C.
146 elegans* miR-39 miRNA (cel-miR-39) was added as an external control for extraction
147 efficiency and normalization.

148 Analysis were performed using the $2^{-\Delta\Delta Ct}$ method [20]. The ΔCt is the difference of expression
149 (expressed as cycle threshold (Ct)) between the target miRNA (miRNA being tested) and a
150 housekeeping miRNA (external control, miRNA cel-miR-39) for each sample. The difference
151 between the ΔCt of the target sample and the ΔCt of the reference sample (calibrator sample:
152 first sample of the first patient) is the $\Delta\Delta Ct$. By calculating the $2^{-\Delta\Delta Ct}$, the relative
153 quantification (RQ) of the target gene expression sample was compared to the reference
154 sample. RQs greater or less than 1 were considered as over-expression or under-expressions,
155 respectively. Samples with expressions lower than the limit of quantification (Ct result > 35
156 cycles) were excluded from the analyses and the graphics, because they were considered as
157 non-significant expressions of miRNAs.

158 MiRNAs selection

159 More than 1000 cancer miRNAs have been reported in the literature [21]. Eleven miRNAs
160 were selected for the present study. This number resulted from a selection performed in 2
161 steps. First, a large panel of 84 human circulating miRNAs (MIHS-106ZA, QIAGEN) were
162 tested in 8 randomly selected patients in order to identify the 8 miRNAs which were the most
163 relevant for our kinetic analysis with the following criteria: sufficient expression for
164 replication (at least 75% of Ct lower than 35 cycles), and kinetics considered to be consistent
165 during treatment (in the sense that the miRNA level should increase or decrease during
166 treatment periods). Moreover, the selection was guided by the literature analysis with the
167 requirement that these miRNAs had to be involved in the ovarian carcinogenesis, and they
168 were shown to exhibit differential expressions in between ovarian cancer patients and control
169 groups. The eight miRNAs selected from this step were: hsa-miR-15b-5p, hsa-miR-16-5p,
170 hsa-miR-20a-5p, hsa-miR-21-5p, hsa-miR-93-5p, hsa-miR-122-5p, hsa-miR-150-5p and hsa-
171 miR-195-5p (Supplementary Figure S1). In addition, three additional miRNAs were selected
172 based on publications reporting relevant kinetics of miRNAs during chemotherapy: hsa-miR-
173 200b-3p, hsa-miR-148b-5p, and hsa-miR-34a-5p [17,18,22].

174 The eleven selected miRNAs were quantified in the remaining 111 patients.

175 Kinetic analyses of longitudinal miRNAs and CA125 changes during CHIVA trial

176 The time changes of miRNA titers and of CA125 were assessed on the following time
177 windows: 1) screening to pre-surgery timepoint; 2) pre-surgery timepoint to first cycle after
178 surgery (cycle 4); and 3) third cycle after surgery to progression, if any. The differences in
179 between plasma miRNA RQs and CA125 concentrations on these time periods were assessed
180 using Wilcoxon signed-rank tests. The longitudinal changes in miRNAs RQ and in CA125
181 levels were graphically assessed for each patient, and for the global population.

182 Prognostic and predictive values of miRNA and of CA125 longitudinal time changes during
183 neo-adjuvant chemotherapy

184 As previously done for CA125 [2] and other serum biomarkers [23], the time changes of
185 miRNA RQs and of CA125 concentrations during neoadjuvant chemotherapy were calculated
186 using their percentage decreases from the baseline timepoint (screening value) to the pre-
187 operative timepoint (before IDS).

188 Receiver operating characteristic (ROC) curves were used to evaluate the discriminative
189 predictive ability of miRNA & CA125 percentage decreases regarding the likelihood of
190 complete CC0 IDS, and of radiological tumor response (assessed using RECIST V1.1
191 criteria).

192 PFS and OS were calculated as the time intervals from treatment start to the date of
193 progression or death, whichever occurred first, for PFS; and to the death only for OS.

194 The 5 following thresholds were arbitrarily chosen: the cutoffs maximizing 1) the prediction
195 ability of CC0 IDS (CC0 vs CC1-CC2); 2) the prediction ability of tumor response (complete
196 response and partial response (CR & PR) versus stable disease and progressive disease (SD &
197 PD), using the Youden index statistic criterion with ROC curves; along with 3) the 1st
198 quartiles of the biomarker percentage decrease distributions; 4) the cutoffs associated with a
199 decline of miRNA or CA125 by 25%; 5) the cutoffs associated with a decline of miRNA or
200 CA125 by 50%.

201 The predictive values of every miRNA and CA125 percentage variations, dichotomized by
202 the above thresholds, regarding PFS and OS were first investigated using c-index tests [24].
203 The prognostic values of CA125 and miRNA percentage variations found to be the most
204 significant with the c-index analyses, were subsequently assessed using Kaplan-Meier method
205 and log-rank tests.

206 All tests were two-sided with 5% alfa risks. All statistical analyses were performed using R
207 3.3.2 software with “survival”, "ROC" and "pec" packages.

208

209 Results

210 Patients characteristics

211 188 patients were included in the CHIVA trial from May 2013 to July 2015. The data from
212 119 of them (63%) could be analyzed for the present study, comprising 8 patients chosen for
213 the selection of 8 miRNAs from the large panel, and 111 remaining patients for the kinetic
214 analyses of 11 miRNAs (Supplementary Figure S2). The characteristics of the included
215 patients are presented in Table 1 and were similar for the 2 groups.

216 Total 8172 miRNAs concentrations were assayed. The concentrations were above the limits
217 of quantification for 6674 samples (82%), ranging from 23% to 98% depending of the
218 miRNA. All the 1299 CA125 measured values were above the limits of quantification (Table
219 2).

220 Kinetic analysis of miRNAs and CA125 longitudinal time changes in CHIVA trial

221 The kinetics of miRNAs were poorly assessable. No statistically significant changes of the
222 miRNAs expressions during treatment were found using Wilcoxon signed-rank tests.
223 However, statistically significant increases of miRNAs expressions were identified at disease
224 progressions for 6 mi-RNA (Table 2).

225 The comparative analyses of miRNA and CA125 kinetics during the disease managements
226 showed that the time-changes of CA125 were more homogeneous and rational, than those of
227 miRNAs (Table 2, Figure 1 and Supplementary Figures S3 to S7). Declines of CA125 by
228 50% or more during chemotherapy were found in 98% of patients (GCIG CA125 response

229 criterion), while such analyses were not assessable for the miRNAs, due to their hectic and
230 irrational kinetics, with high inter-individual variability.

231 Predictive value of miRNAs regarding the tumor response rate or interval debulking
232 cytoreductive surgery score

233 The predictive abilities of every miRNA expression percentage variation during neoadjuvant
234 chemotherapy regarding the likelihood of complete IDS (yes vs no) using ROC area under the
235 curve (AUC) are presented in Table 3. The ROC AUCs ranged from 0.51 to 0.88; none of
236 them being statistically significant, except for the hsa-miR-200b-3p. The same lack of
237 significant predictive value was found for the likelihood of tumor response (CR-PR vs SD-
238 PD) with ROC AUCs ranging from 0.50 to 0.86; all of them being non-significant, except for
239 hsa-miR-200b-3p (Table 3).

240 The same analyses done with CA125 percentage decrease showed a ROC AUC at 0.58 (95%
241 CI 0.45-0.71) for prediction of CC0 IDS, and at 0.64 (95% CI 0.51-0.76) for prediction of
242 tumor response (Table 3).

243 The values of the different thresholds maximizing the likelihood of complete surgery and of
244 tumor response for miRNA and CA125 percentage decreases are presented in Table 3.

245 Predictive value of miRNA kinetics for survivals

246 The predictive values of miRNA and CA125 variations, dichotomized by the different cutoffs,
247 regarding PFS and OS are shown in Table 4. The c-index values of miRNA time changes
248 ranged from 0.47 to 0.67 for PFS; and from 0.44 to 0.81 for OS. The highest significant
249 predictive values regarding PFS were observed with the hsa-miR-15b-5p (categorized by the
250 threshold optimizing the tumor response likelihood) at 0.63 (95% CI 0.57-0.69), and with the
251 hsa-miR-34a-5p (categorized by the cutoff associated with a decline by 25%) at 0.64 (95% CI
252 0.52-0.76).

253 For OS, the highest significant predictive values were obtained with the has-miR-15b-5p
254 (categorized by the threshold optimizing the tumor response likelihood) at 0.63 (95% CI 0.53-
255 0.73), and with the hsa-miR-93-5p (categorized by the threshold optimizing the prediction of
256 complete IDS) at 0.66 (95% CI 0.56-0.76). The similar analyses done with the CA125
257 kinetics showed c-index values ranging from 0.50 to 0.54 for PFS, and 0.50 to 0.56 for OS.

258 The outcomes regarding the prognostic values of the most significant miRNAs using Kaplan-
259 Meier survival curves and log-rank tests are presented in Figure 2. Hsa-miR-15b-5p and hsa-
260 miR-34a-5p were associated with better PFS (median PFS = 13.2 vs 20.8 months and 13.7 vs
261 24.8 months, respectively, $p < 0.01$), whilst hsa-miR-15b-5p and hsa-miR-93-5p were
262 associated with better OS (median OS = 34.2 vs 52.0 and 34.3 vs NR months, respectively,
263 $p < 0.01$). CA125 kinetics assessed with the similar methodology were not significant,
264 including with the 50% decrease cutoff recognized as the GCIG CA125 response criterion.

265

266 Discussion

267 This present study represents the first analysis on the actual predictive and prognostic values
268 of miRNA longitudinal kinetics in cancer patients managed with medical-surgical treatments.
269 MiRNAs have been recently reported as major actors in the tumorigenesis [25]. The miRNA
270 expression time changes were expected to correlate with the tumor burden changes, thereby
271 representing promising predictive biomarkers [17,18,26,27]. The ovarian cancer population
272 was particularly adequate, since there is a need for reliable serum biomarkers to overcome the
273 limitation of imaging, and also because miRNA kinetic value could be assessed with respect
274 to the validated serum marker CA125 [2,28].

275 Several authors reported the prognostic or predictive values of miRNA expressions (e.g. miR-
276 34a-5p level before treatment associated with CC0 cytoreductive surgery in 56 patients, ROC

277 AUC = 0.78) [22], or miRNA expression changes during treatments (e.g. miR-200b change
278 from baseline to post-operative period regarding PFS in 33 patients) [17].

279 The innovative design of the randomized phase II trial CHIVA, in which serial blood samples
280 were regularly performed during disease management, enabled us to assess the kinetics of 11
281 miRNAs measured in a properly conducted prospective trial. This miRNA selection was
282 based on initial explorative analyses with a large panel of miRNAs in 8 patients, and based on
283 the literature data. Our selection appeared adequate since the miRNAs were highly expressed
284 in 82% of samples.

285 The kinetic analysis of miRNA demonstrates that no rational longitudinal profiles could be
286 identified for any of them throughout the treatment. The large intra-individual and inter-
287 individual variability of miRNA time-changes made any longitudinal assessment impossible
288 with mathematical modeling. As a consequence, it was not adequate to search for
289 relationships between tumor burden alterations and miRNA kinetics using mathematical
290 modeling, contrarily to what was done with CA125 [3,29–32].

291 Our outcomes in favor a lack of predictive value of miRNA are not consistent with the data
292 reported in the literature earlier [17,18,22,26,27,33]. This discrepancy could be understood by
293 the methodology used by the other teams so far. In those studies, the kinetic analyses were
294 performed on maximum two or three time points (contrarily to our approach based on 8
295 timepoints per patient), and it is possible that the limited numbers of time points would have
296 induced biases in the outcomes given the high inter-individual variability of miRNA
297 expressions. Moreover, the large inconsistency in the types of miRNAs reported in the
298 literature assumed to exhibit prognostic or predictive values highlights the difficulty to
299 identify miRNAs with reproducible prognostic/predictive values.

300 The present study present limitations prone to reduce the relevance of data. First, it was not
301 possible to assess all miRNAs, and we had to select an affordable panel. We selected 11
302 miRNAs that appeared rational, based on an explorative analysis including a large panel of 84
303 miRNAs, and based on the literature. Our conclusions about the lack of assessable
304 longitudinal prognostic or predictive kinetic profiles may not necessarily be transposable to
305 other miRNAs. The selection of the best cut-offs for analyzing miRNAs kinetics in a way that
306 would make the analyses helpful for clinical purpose was another difficulty. We chose to
307 implement common statistical approaches, such as ROC curve tests and quartiles, but the
308 relevance is questionable. None of the cutoffs that we assessed appeared superior. Moreover
309 the expressions of the assessed miRNAs were below the lower limit of quantification in 18%
310 of measurements, thereby making the kinetics analysis even more complicated, especially for
311 miR-200b-3p. Furthermore, the impact of the tumor histological subtypes or BRCA status on
312 miRNAs kinetics could not be assessed. Of note, recent data on the kinetics of CA125
313 assessed using mathematical modeling suggest that the modeled kinetic parameter KELIM
314 could be a promising predictor of complete IDS likelihood, or of survival, in ovarian cancer
315 patients [34–36]. However this parameter was not assessed here, because we thought that
316 miRNA and CA125 kinetics had to be investigated with the same methodologies.

317 The present study is the largest work including serial assessment of a high numbers of
318 miRNA, in cancer patients managed with multi-modal treatment in a properly conducted
319 phase II trial. The outcomes indicate miRNA may not be adequate biomarkers for longitudinal
320 kinetic analyses during treatment. Consistently with the inconsistent data of the literature on
321 the prognostic or the predictive values of miRNA titers or time changes, we could not identify
322 any miRNA prone to be reproducible predictors of IDS score, tumor response, or survivals in
323 ovarian cancer patients.

324 Acknowledgements

325 We thank all the patients and their families, the investigators and study teams, Pharmacists,
326 Pathologists, Biologists, ARCAGY GINECO Project Manager and study teams and
327 ARCAGY GINECO biological resource center for their support of the study.

328 Conflict of interest statement

329 The authors declare no potential conflicts of interest.

330 Funding

331 This study was supported by the ARCAGY GINECO group.

332

333

- 335 [1] Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. *CA Cancer J Clin* 2017;67:7–
336 30. <https://doi.org/10.3322/caac.21387>.
- 337 [2] Rustin GJS, Vergote I, Eisenhauer E, Pujade-Lauraine E, Quinn M, Thigpen T, et al.
338 Definitions for response and progression in ovarian cancer clinical trials incorporating
339 RECIST 1.1 and CA 125 agreed by the Gynecological Cancer Intergroup (GCIG). *Int J*
340 *Gynecol Cancer Off J Int Gynecol Cancer Soc* 2011;21:419–23.
341 <https://doi.org/10.1097/IGC.0b013e3182070f17>.
- 342 [3] You B, Colomban O, Freyer G, Leary A, Ray-Coquard IL, Lortholary A, et al. Two
343 prognostic populations of ovarian cancer patients defined by CA125 modeled kinetic
344 parameter KELIM (AGO-OVAR 7 & 9; ICON 7 AGO/GINECO/ MRC CTU/GCIG
345 trials). *J Clin Oncol* 2017;35:5554–5554.
346 https://doi.org/10.1200/JCO.2017.35.15_suppl.5554.
- 347 [4] Azad NS, Annunziata CM, Steinberg SM, Minasian L, Premkumar A, Chow C, et al.
348 Lack of reliability of CA125 response criteria with anti-VEGF molecularly targeted
349 therapy. *Cancer* 2008;112:1726–32. <https://doi.org/10.1002/cncr.23374>.
- 350 [5] Rustin GJS, van der Burg MEL, Griffin CL, Guthrie D, Lamont A, Jayson GC, et al.
351 Early versus delayed treatment of relapsed ovarian cancer (MRC OV05/EORTC 55955):
352 a randomised trial. *Lancet Lond Engl* 2010;376:1155–63. [https://doi.org/10.1016/S0140-6736\(10\)61268-8](https://doi.org/10.1016/S0140-6736(10)61268-8).
- 354 [6] Pujade-Lauraine E, Wagner U, Aavall-Lundqvist E, Gebiski V, Heywood M, Vasey PA,
355 et al. Pegylated liposomal Doxorubicin and Carboplatin compared with Paclitaxel and
356 Carboplatin for patients with platinum-sensitive ovarian cancer in late relapse. *J Clin*
357 *Oncol Off J Am Soc Clin Oncol* 2010;28:3323–9.
358 <https://doi.org/10.1200/JCO.2009.25.7519>.
- 359 [7] Wright AA, Bohlke K, Armstrong DK, Bookman MA, Cliby WA, Coleman RL, et al.
360 Neoadjuvant Chemotherapy for Newly Diagnosed, Advanced Ovarian Cancer: Society
361 of Gynecologic Oncology and American Society of Clinical Oncology Clinical Practice
362 Guideline. *J Clin Oncol Off J Am Soc Clin Oncol* 2016;34:3460–73.
363 <https://doi.org/10.1200/JCO.2016.68.6907>.
- 364 [8] du Bois A, Reuss A, Pujade-Lauraine E, Harter P, Ray-Coquard I, Pfisterer J. Role of
365 surgical outcome as prognostic factor in advanced epithelial ovarian cancer: a combined
366 exploratory analysis of 3 prospectively randomized phase 3 multicenter trials: by the
367 Arbeitsgemeinschaft Gynaekologische Onkologie Studiengruppe Ovarialkarzinom
368 (AGO-OVAR) and the Groupe d'Investigateurs Nationaux Pour les Etudes des Cancers
369 de l'Ovaire (GINECO). *Cancer* 2009;115:1234–44. <https://doi.org/10.1002/cncr.24149>.
- 370 [9] Shukla GC, Singh J, Barik S. MicroRNAs: Processing, Maturation, Target Recognition
371 and Regulatory Functions. *Mol Cell Pharmacol* 2011;3:83–92.
- 372 [10] Meng X, Müller V, Milde-Langosch K, Trillsch F, Pantel K, Schwarzenbach H.
373 Diagnostic and prognostic relevance of circulating exosomal miR-373, miR-200a, miR-
374 200b and miR-200c in patients with epithelial ovarian cancer. *Oncotarget* 2016;7:16923–
375 35. <https://doi.org/10.18632/oncotarget.7850>.
- 376 [11] Liang H, Jiang Z, Xie G, Lu Y. Serum microRNA-145 as a novel biomarker in human
377 ovarian cancer. *Tumour Biol J Int Soc Oncodevelopmental Biol Med* 2015;36:5305–13.
378 <https://doi.org/10.1007/s13277-015-3191-y>.
- 379 [12] Meng X, Joosse SA, Müller V, Trillsch F, Milde-Langosch K, Mahner S, et al.
380 Diagnostic and prognostic potential of serum miR-7, miR-16, miR-25, miR-93, miR-
381 182, miR-376a and miR-429 in ovarian cancer patients. *Br J Cancer* 2015;113:1358–66.
382 <https://doi.org/10.1038/bjc.2015.340>.

- 383 [13] Shapira I, Oswald M, Lovecchio J, Khalili H, Menzin A, Whyte J, et al. Circulating
384 biomarkers for detection of ovarian cancer and predicting cancer outcomes. *Br J Cancer*
385 2014;110:976–83. <https://doi.org/10.1038/bjc.2013.795>.
- 386 [14] Langhe R, Norris L, Saadeh FA, Blackshields G, Varley R, Harrison A, et al. A novel
387 serum microRNA panel to discriminate benign from malignant ovarian disease. *Cancer*
388 *Lett* 2015;356:628–36. <https://doi.org/10.1016/j.canlet.2014.10.010>.
- 389 [15] Gong L, Wang C, Gao Y, Wang J. Decreased expression of microRNA-148a predicts
390 poor prognosis in ovarian cancer and associates with tumor growth and metastasis.
391 *Biomed Pharmacother Biomedecine Pharmacother* 2016;83:58–63.
392 <https://doi.org/10.1016/j.biopha.2016.05.049>.
- 393 [16] Liu W, Wang S, Zhou S, Yang F, Jiang W, Zhang Q, et al. A systems biology approach
394 to identify microRNAs contributing to cisplatin resistance in human ovarian cancer cells.
395 *Mol Biosyst* 2017;13:2268–76. <https://doi.org/10.1039/c7mb00362e>.
- 396 [17] Kapetanakis N-I, Uzan C, Jimenez-Pailhes A-S, Gouy S, Bentivegna E, Morice P, et al.
397 Plasma miR-200b in ovarian carcinoma patients: distinct pattern of pre/post-treatment
398 variation compared to CA-125 and potential for prediction of progression-free survival.
399 *Oncotarget* 2015;6:36815–24.
- 400 [18] Benson EA, Skaar TC, Liu Y, Nephew KP, Matei D. Carboplatin with Decitabine
401 Therapy, in Recurrent Platinum Resistant Ovarian Cancer, Alters Circulating miRNAs
402 Concentrations: A Pilot Study. *PLoS ONE* 2015;10.
403 <https://doi.org/10.1371/journal.pone.0141279>.
- 404 [19] Vargatef in Addition to First Line Chemotherapy With Interval Debulking Surgery in
405 Patients With Ovarian Cancer - Full Text View - ClinicalTrials.gov n.d.
406 <https://clinicaltrials.gov/ct2/show/NCT01583322> (accessed September 11, 2018).
- 407 [20] Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T)
408 method. *Nat Protoc* 2008;3:1101–8.
- 409 [21] Kozomara A, Griffiths-Jones S. miRBase: annotating high confidence microRNAs using
410 deep sequencing data. *Nucleic Acids Res* 2014;42:D68-73.
411 <https://doi.org/10.1093/nar/gkt1181>.
- 412 [22] Shah JS, Gard GB, Yang J, Maidens J, Valmadre S, Soon PS, et al. Combining serum
413 microRNA and CA-125 as prognostic indicators of preoperative surgical outcome in
414 women with high-grade serous ovarian cancer. *Gynecol Oncol* 2018;148:181–8.
415 <https://doi.org/10.1016/j.ygyno.2017.11.005>.
- 416 [23] Almufti R, Wilboux M, Oza A, Henin E, Freyer G, Tod M, et al. A critical review of the
417 analytical approaches for circulating tumor biomarker kinetics during treatment. *Ann*
418 *Oncol Off J Eur Soc Med Oncol* 2014;25:41–56. <https://doi.org/10.1093/annonc/mdt382>.
- 419 [24] Gerds TA, Kattan MW, Schumacher M, Yu C. Estimating a time-dependent
420 concordance index for survival prediction models with covariate dependent censoring.
421 *Stat Med* 2013;32:2173–84. <https://doi.org/10.1002/sim.5681>.
- 422 [25] Deb B, Uddin A, Chakraborty S. miRNAs and ovarian cancer: An overview. *J Cell*
423 *Physiol* 2017. <https://doi.org/10.1002/jcp.26095>.
- 424 [26] Kovacova J, Juracek J, Poprach A, Buchler T, Kopecky J, Fiala O, et al. Candidate
425 MicroRNA Biomarkers of Therapeutic Response to Sunitinib in Metastatic Renal Cell
426 Carcinoma: A Validation Study in Patients with Extremely Good and Poor Response.
427 *Anticancer Res* 2018;38:2961–5. <https://doi.org/10.21873/anticancer.12546>.
- 428 [27] Frères P, Josse C, Bovy N, Boukerroucha M, Struman I, Bours V, et al. Neoadjuvant
429 Chemotherapy in Breast Cancer Patients Induces miR-34a and miR-122 Expression. *J*
430 *Cell Physiol* 2015;230:473–81. <https://doi.org/10.1002/jcp.24730>.

- 431 [28] Tuxen MK, Sölétormos G, Dombernowsky P. Serum tumour marker CA 125 in
432 monitoring of ovarian cancer during first-line chemotherapy. *Br J Cancer* 2001;84:1301–
433 7. <https://doi.org/10.1054/bjoc.2001.1787>.
- 434 [29] You B, Colomban O, Heywood M, Lee C, Davy M, Reed N, et al. The strong prognostic
435 value of KELIM, a model-based parameter from CA 125 kinetics in ovarian cancer: data
436 from CALYPSO trial (a GINECO-GCIG study). *Gynecol Oncol* 2013;130:289–94.
437 <https://doi.org/10.1016/j.ygyno.2013.05.013>.
- 438 [30] Ducoulombier S, Golfier F, Colomban O, Benayoun D, Bolze P-A, Tod M, et al.
439 Modeling CA-125 During Neoadjuvant Chemotherapy for Predicting Optimal
440 Cytoreduction and Relapse Risk in Ovarian Cancer. *Anticancer Res* 2017;37:6879–86.
441 <https://doi.org/10.21873/anticancer.12150>.
- 442 [31] Wilbaux M, Hénin E, Oza A, Colomban O, Pujade-Lauraine E, Freyer G, et al. Dynamic
443 modeling in ovarian cancer: an original approach linking early changes in modeled
444 longitudinal CA-125 kinetics and survival to help decisions in early drug development.
445 *Gynecol Oncol* 2014;133:460–6. <https://doi.org/10.1016/j.ygyno.2014.04.003>.
- 446 [32] Wilbaux M, Hénin E, Oza A, Colomban O, Pujade-Lauraine E, Freyer G, et al.
447 Prediction of tumour response induced by chemotherapy using modelling of CA-125
448 kinetics in recurrent ovarian cancer patients. *Br J Cancer* 2014;110:1517–24.
449 <https://doi.org/10.1038/bjc.2014.75>.
- 450 [33] Zhu W, Liu M, Fan Y, Ma F, Xu N, Xu B. Dynamics of circulating microRNAs as a
451 novel indicator of clinical response to neoadjuvant chemotherapy in breast cancer.
452 *Cancer Med* 2018. <https://doi.org/10.1002/cam4.1723>.
- 453 [34] Colomban O, Tod M, Leary A, Ray-Coquard I, Lortholary A, Hardy-Bessard AC, et al.
454 Early Modeled Longitudinal CA-125 Kinetics and Survival of Ovarian Cancer Patients:
455 A GINECO AGO MRC CTU Study. *Clin Cancer Res Off J Am Assoc Cancer Res* 2019.
456 <https://doi.org/10.1158/1078-0432.CCR-18-3335>.
- 457 [35] Robelin P, Tod M, Colomban O, Louvet C, Lotz J-P, Abadie Lacourtoisie S, et al.
458 Modeled CA-125 kinetics during neoadjuvant chemotherapy for predicting the
459 likelihood of optimal interval debulking surgery in ovarian cancer patients: Data from
460 CHIVA trial (a GINECO study). *J Clin Oncol* 2019;37:5546–5546.
461 https://doi.org/10.1200/JCO.2019.37.15_suppl.5546.
- 462 [36] You B, Robelin P, Tod M, Louvet C, Lotz J-P, Abadie-Lacourtoisie S, et al. CA-125
463 ELIMination rate constant K (KELIM) Is A Marker Of Chemosensitivity In Patients
464 With Ovarian Cancer: Results from the Phase II CHIVA trial. *Clin Cancer Res Off J Am*
465 *Assoc Cancer Res* 2020. <https://doi.org/10.1158/1078-0432.CCR-20-0054>.
- 466

467 **Figures Legends**

468 **Figure 1. Longitudinal time changes of the miR-195 expression and of CA125**
469 **concentrations.** (A) Individual patient profiles of miR-195 and CA125 kinetics according to
470 completeness of cytoreduction (green line: complete cytoreductive surgery; red: incomplete
471 cytoreductive surgery); (B) Changes in populations values of blood miR-195 expressions and
472 CA125 concentrations during management steps and at progression

473 Abbreviations: S: Screening; C: Pre-surgery; P: Progression.

474

475 **Figure 2. Prognostic values of the expression changes of the most significant miRNAs**
476 **and of CA125 regarding PFS and OS using Kaplan-Meier survival curves of patients.**

477 (A) PFS with miR-15b categorized by the threshold optimizing the tumor response likelihood
478 and miR-34a categorized by the 25% decline threshold; (B) OS with miR-15b categorized by
479 the threshold optimizing the tumor response and miR-93 categorized by the threshold
480 optimizing the complete IDS likelihood; (C) PFS with CA125 categorized by the first quartile
481 of the RV and OS with CA125 categorized the threshold maximizing the complete IDS
482 likelihood.

483 IDS: interval debulking surgery; PFS: progression free survival; OS: overall survival.

484

485 **Tables**

486 **Table 1. Patient characteristics at inclusion**

487

488 **Table 2. Expressions of the 11 miRNA and of CA125. Statistical significances of the**
489 **comparisons of expressions on 3 different time windows using Wilcoxon signed-rank**
490 **tests: 1) from screening (baseline) to IDS; 2) from IDS to cycle 4; 3) from cycle 6 to**
491 **disease progression.**

492

493 **Table 3. Outcomes of the ROC curve AUC analyses on 1) Predictive ability of miRNA**
494 **and CA125 percentage decreases during neoadjuvant chemotherapy regarding the**
495 **likelihood of complete cytoreductive surgery (complete cytoreductive surgery with no**
496 **macroscopic residuals vs incomplete debulking with residuals), or of tumor response (CR +**
497 **PR vs SD + PD); 2) Identification of the optimal cutoff for predictions of the likelihood of**
498 **complete cytoreductive surgery, or of tumor response using Youden criterion.**

499

500 **Table 4. C-index values regarding the predictive values of miRNA and CA125**
501 **percentage decreases, categorized by the 5 different cutoffs for PFS and OS.**

502




Figure 1.









Figure 2.

Table 1. Patient characteristics at inclusion

Characteristics	MiRNA selection	
	patients	Test patients
Patients number, <i>n</i>	8	111
Age, years, median (range)	65 (31-71)	64 (31-79)
ECOG PS, <i>n</i> (%)		
0-1	7 (88%)	98 (88%)
2	1 (12%)	12 (11%)
NA	0 (0%)	1 (1%)
FIGO stage, <i>n</i> (%)		
IIIc	7 (88%)	85 (77%)
IV	1 (12%)	26 (23%)
Histological type, <i>n</i> (%)		
Serous/papillary	8 (100%)	98 (88%)
Endometrioid	0 (0%)	1 (1%)
Mucinous	0 (0%)	1 (1%)
Clear cells	0 (0%)	1 (1%)
Undifferentiated	0 (0%)	3 (3%)
NA	0 (0%)	7 (6%)
Histological grade, <i>n</i> (%)		
Grade 1	0 (0%)	3 (3%)
Grade 2	1 (12%)	10 (9%)
Grade 3	5 (63%)	79 (71%)
NA	2 (25%)	19 (17%)
CA125 at inclusion (xULN), median (range)	24.4 (2.6-250.4)	26.9 (0.5-784.3)
IDS performed, <i>n</i> (%)	8 (100%)	86 (77%)
Completeness of cytoreduction, <i>n</i> (%)		
No macroscopic residuals	5 (63%)	70 (81%)
Tumoral residuals < 0,25 cm	0 (0%)	6 (7%)
Tumoral residuals 0,25 < R < 2.5 cm	3 (37%)	7 (8%)
Tumoral residuals > 2.5 cm	0 (0%)	2 (2%)
NA	0 (0%)	1 (1%)
Treatment with nintedanib, <i>n</i> (%)	5 (63%)	68 (61%)
Response to neo-adjuvant chemotherapy, <i>n</i> (%)		
CR	1 (12%)	2 (2%)
PR	3 (37%)	45 (41%)
SD	4 (50%)	57 (51%)
PD	0 (0%)	2 (2%)
NA	0 (0%)	5 (5%)
Progression free survival in months, median (95% CI)	15.6 (11.8-NR)	16.2 (14.4-18.4)
Overall survival in months, median (95% CI)	NR (38.6-NR)	40.1 (36.2-NR)
Death, <i>n</i> (%)	3 (37%)	55 (50%)

ECOG PS: ECOG Performance Status; IDS: Interval debulking surgery; NA: Non-available;

ULN: Upper limit of normal; CR: Complete response; PR: Partial response; SD: Stable disease; PD: Progressive disease CI: Confident interval; NR: Not reached.

Table 2. Expressions of the 11 miRNA and of CA125. Statistical significances of the comparisons of expressions on 3 different time windows using Wilcoxon signed-rank tests: 1) from screening (baseline) to IDS; 2) from IDS to cycle 4; 3) from cycle 6 to disease progression.

Serum biomarkers	% of samples with RQ greater than LoQ	Significances of expression differences on 3 time windows		
		1) Screening to IDS <i>p</i> value	2) IDS to Cycle 4 <i>p</i> value	3) Cycle 6 to Progression <i>p</i> value
hsa-miR-15b-5p	97%	0.986	0.184	0.001
hsa-miR-16-5p	98%	0.397	0.454	0.001
hsa-miR-20a-5p	92%	0.886	0.441	0.013
hsa-miR-21-5p	98%	0.405	0.920	0.005
hsa-miR-93-5p	90%	0.974	0.441	0.026
hsa-miR-122-5p	96%	0.296	0.309	0.437
hsa-miR-150-5p	90%	0.182	0.012	0.081
hsa-miR-195-5p	96%	0.660	0.186	0.007
hsa-miR-34a-5p	57%	0.915	0.080	0.117
hsa-miR-148b-5p	54%	0.379	0.624	0.181
hsa-miR-200b-3p	23%	0.375	0.461	0.500
CA125	100%	<0.001	0.109	<0.001

IDS: Interval debulking surgery; RQ: Relative quantification; LoQ: limit of quantification;

Table 3. Outcomes of the ROC curve AUC analyses on 1) Predictive ability of miRNA and CA125 percentage decreases during neoadjuvant chemotherapy regarding the likelihood of complete cytoreductive surgery (complete cytoreductive surgery with no macroscopic residuals vs incomplete debulking with residuals), or of tumor response (CR + PR vs SD + PD); 2) Identification of the optimal cutoff for predictions of the likelihood of complete cytoreductive surgery, or of tumor response using Youden criterion.

Serum marker	Predictive value of miRNA and CA125 kinetics using ROC curve AUCs		Optimal threshold of the percentage decline maximizing the prediction ability of	
	Prediction of complete cytoreductive surgery (95% CI)	Prediction of tumor response (95% CI)	Complete cytoreductive surgery	Tumor response
hsa-miR-15b-5p	0.55 (0.41-0.70)	0.50 (0.36-0.63)	-6%	+27%
hsa-miR-16-5p	0.55 (0.40-0.69)	0.64 (0.50-0.76)	-23%	-11%
hsa-miR-20a-5p	0.53 (0.38-0.68)	0.57 (0.43-0.72)	+11%	+29%
hsa-miR-21-5p	0.52 (0.37-0.66)	0.53 (0.40-0.66)	-32%	+9%
hsa-miR-93-5p	0.52 (0.37-0.67)	0.67 (0.54-0.80)	+15%	+111%
hsa-miR-122-5p	0.58 (0.43-0.73)	0.58 (0.45-0.72)	+8%	+18%
hsa-miR-150-5p	0.56 (0.41-0.72)	0.51 (0.37-0.65)	-32%	+5%
hsa-miR-195-5p	0.52 (0.37-0.67)	0.60 (0.46-0.73)	-13%	-33%
hsa-miR-34a-5p	0.54 (0.27-0.80)	0.53 (0.32-0.74)	-41%	+124%
hsa-miR-148b-5p	0.63 (0.40-0.86)	0.58 (0.37-0.79)	-30%	+168%
hsa-miR-200b-3p	0.88 (0.65-1.00)	0.86 (0.60-1.00)	-10%	-3%
CA125	0.57 (0.43-0.70)	0.63 (0.49-0.76)	-99%	-97%

ROC: Receiver operating characteristic; AUC: Area under the curve; IDS: Interval debulking surgery; CI: Confident interval.

Table 4. C-index values regarding the predictive values of miRNA and CA125 percentage decreases, categorized by the 5 different cutoffs for PFS and OS.

Serum marker	Optimal threshold for predicting complete cytoreductive surgery (SD)	Optimal threshold for predicting tumor response (SD)	1 st Quartile of percentage decrease distribution (SD)	Percentage decrease of 25% (SD)	Percentage decrease of 50% (SD)
Analyses for PFS					
hsa-miR-15b-5p	0.61 (0.04)	0.63 (0.03)	0.60 (0.03)	0.60 (0.03)	0.58 (0.03)
hsa-miR-16-5p	0.55 (0.04)	0.57 (0.04)	0.55 (0.03)	0.53 (0.03)	0.53 (0.03)
hsa-miR-20a-5p	0.57 (0.04)	0.51 (0.04)	0.50 (0.03)	0.52 (0.04)	0.51 (0.04)
hsa-miR-21-5p	0.59 (0.03)	0.60 (0.03)	0.58 (0.03)	0.61 (0.04)	0.60 (0.03)
hsa-miR-93-5p	0.60 (0.04)	0.57 (0.03)	0.58 (0.04)	0.58 (0.04)	0.57 (0.03)
hsa-miR-122-5p	0.54 (0.04)	0.56 (0.03)	0.55 (0.03)	0.57 (0.04)	0.56 (0.03)
hsa-miR-150-5p	0.59 (0.04)	0.57 (0.04)	0.59 (0.04)	0.58 (0.04)	0.56 (0.03)
hsa-miR-195-5p	0.60 (0.04)	0.56 (0.03)	0.57 (0.03)	0.58 (0.03)	0.53 (0.03)
hsa-miR-34a-5p	0.60 (0.05)	0.56 (0.04)	0.57 (0.05)	0.64 (0.06)	0.55 (0.05)
hsa-miR-148b-5p	0.49 (0.06)	0.54 (0.04)	0.47 (0.05)	0.55 (0.05)	0.51 (0.05)
hsa-miR-200b-3p	0.58 (0.11)	0.67 (0.11)	0.64 (0.11)	0.64 (0.11)	0.59 (0.08)
CA125	0.51 (0.03)	0.51 (0.04)	0.52 (0.03)	0.50 (0)	0.50 (0.02)
Analyses for OS					
hsa-miR-15b-5p	0.60 (0.05)	0.63 (0.05)	0.61 (0.04)	0.61 (0.04)	0.62 (0.04)
hsa-miR-16-5p	0.57 (0.05)	0.58 (0.05)	0.56 (0.04)	0.55 (0.04)	0.55 (0.04)
hsa-miR-20a-5p	0.57 (0.05)	0.53 (0.05)	0.55 (0.04)	0.55 (0.05)	0.48 (0.05)
hsa-miR-21-5p	0.59 (0.04)	0.61 (0.04)	0.59 (0.04)	0.60 (0.05)	0.60 (0.04)
hsa-miR-93-5p	0.66 (0.05)	0.60 (0.05)	0.59 (0.05)	0.64 (0.05)	0.57 (0.04)
hsa-miR-122-5p	0.54 (0.05)	0.56 (0.04)	0.54 (0.04)	0.59 (0.05)	0.55 (0.04)
hsa-miR-150-5p	0.58 (0.04)	0.55 (0.05)	0.58 (0.04)	0.58 (0.04)	0.59 (0.04)
hsa-miR-195-5p	0.63 (0.05)	0.58 (0.04)	0.59 (0.04)	0.60 (0.04)	0.55 (0.04)
hsa-miR-34a-5p	0.53 (0.07)	0.52 (0.06)	0.49 (0.07)	0.51 (0.07)	0.51 (0.06)
hsa-miR-148b-5p	0.53 (0.07)	0.52 (0.05)	0.54 (0.06)	0.55 (0.07)	0.50 (0.06)
hsa-miR-200b-3p	0.81 (0.18)	0.75 (0.19)	0.44 (0.18)	0.44 (0.18)	0.58 (0.12)
CA125	0.50 (0.03)	0.54 (0.04)	0.51 (0.04)	0.50 (0)	0.52 (0.02)

Abbreviations: SD: Standard deviation; IDS: Interval debulking surgery; PFS: Progression

free survival, OS: Overall survival, CI: Confident interval; NR: Not Reached.