



Meta-trial of awake prone positioning with nasal high flow therapy: Invitation to join a pandemic collaborative research effort

Jie Li, Ivan Pavlov, John G. Laffey, Oriol Roca, Sara Mirza, Yonatan Perez, Bairbre Mcnicholas, David Cosgrave, David Vines, Elsa Tavernier, et al.

► To cite this version:

Jie Li, Ivan Pavlov, John G. Laffey, Oriol Roca, Sara Mirza, et al.. Meta-trial of awake prone positioning with nasal high flow therapy: Invitation to join a pandemic collaborative research effort. *Journal of Critical Care*, 2020, 60, pp.140 - 142. <10.1016/j.jcrc.2020.07.020>. <hal-03492426>

HAL Id: hal-03492426

<https://hal.science/hal-03492426v1>

Submitted on 22 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

Meta-trial of awake prone positioning with nasal high flow therapy: invitation to join a pandemic collaborative research effort

Jie Li¹, Ivan Pavlov², John G. Laffey³, Oriol Roca⁴, Sara Mirza¹, Yonatan Perez⁵,
Bairbre McNicholas³, David Cosgrave³, David Vines¹, Elsa Tavernier⁶, Stephan
Ehrmann⁷

1: Department of Cardiopulmonary Sciences, Division of Respiratory Care, Rush
University Medical Center, Chicago, IL

2: Department of Emergency Medicine, Hôpital de Verdun, Montréal, Québec,
Canada

3: Anaesthesia and Intensive Care Medicine, School of Medicine, National
University of Ireland Galway; Department of Anaesthesia, University Hospital
Galway, Saolta Hospital Group, Ireland

4: Servei de Medicina Intensiva, Hospital Universitari Vall d'Hebron, Institut de
Recerca Vall d'Hebron, Barcelona, Spain ; Departament de Medicina, Universitat
Autònoma de Barcelona, Bellaterra, Spain ; Ciber Enfermedades Respiratorias
(CIBERES), Instituto de Salud Carlos III, Madrid, Spain

5: CHRU Tours, Médecine Intensive Réanimation, CIC INSERM 1415, REVA and
CRICS-TriggerSEP research networks, Tours, France

6: Universities of Tours and Nantes, INSERM 1246-SPHERE, Tours, France;
Clinical Investigation Center, INSERM 1415, CHRU Tours, Tours, France

7: CHRU Tours, Médecine Intensive Réanimation, CIC INSERM 1415, REVA and CRICS-TriggerSEP research networks, Tours, France ; and INSERM, Centre d'étude des pathologies respiratoires, U1100, Université de Tours, Tours, France

Corresponding author: Stephan Ehrmann

- stephanehrmann@gmail.com
- Médecine Intensive Réanimation. CHRU de Tours. 2, Bd Tonnellé. 37044 Tours cedex 9. France
- telephone: +33 (0)2 47 47 28 55
- fax: +33 (0)2 47 39 65 36

Authors contributions: ET, JL and SE initially designed the project, SE drafted the manuscript, all authors contributed significantly to study design and concept, all authors reviewed the manuscript for important intellectual content.

Financial disclosure: This work was supported by OpenAI. (3180 18th St, San Francisco, CA 94110, United States), Fisher & Paykel (15 Maurice Paykel Place, East Tamaki, Auckland 2013 New Zealand) and the French ministry of health (Appel d'offre PHRC inter-régional GIRCI grand-ouest).

Conflict of interest statement: JL discloses research support from Fisher & Paykel Healthcare for another research project. IP has been a speaker for Fisher & Paykel Healthcare within the last 3 years. All compensation was paid to the charitable foundation at the hospital where he works, and he did not personally receive any compensation. OR provides consultancy to Hamilton Medical but he did not received any personal fee. All compensations were received by the Institute

of Research of his Institution. He received speaker fees by Air Liquide. SE received unrestricted research grants, travel fee reimbursements and speaker fees from Fisher & Paykel Healthcare, consulting fees from La Diffusion Technique Française, consulting fees and unrestricted research grants from Aerogen Ltd, and an unrestricted research grant from Hamilton medical. Other authors have no conflict of interest to declare.

Word count: 928

Key words: Research design [MeSH], COVID-19 [MeSH], Meta-analysis [MeSH], Pandemics [MeSH], Acute respiratory distress syndrome

The ongoing coronavirus disease pandemic (COVID-19) poses a great challenge to healthcare systems worldwide. Beyond challenges for direct patient care, optimal conduct of research under the specific conditions of the pandemic is a matter of concern and discussion. We present the *meta-trial* concept as a scientifically, clinically, ethically and socially sound method to carry out optimal clinical research in the setting of a pandemic. We take the example of such a *meta-trial*, set up to investigate prone positioning among awake patients undergoing nasal high flow therapy and invite journal readers to join this collaborative research effort.

COVID-19 constraints to clinical research: the pandemic has placed the research community under great pressure with the urgent need for results, given the lack of knowledge concerning optimal management of patients suffering this new disease. Public pressure is high due to the lack of specific effective therapy regarding this major threat to public health; the traditional pace of clinical research being considered as not adapted by most stakeholders. Worldwide funding agencies and regulatory bodies changed their procedures in order to speed up the research process. This effort has led to the launch of a high number of clinical studies within a short period of time to an unprecedented extent. As of 2020 June 15th, 2138 COVID-19 trials were registered in clinicaltrials.gov.

Pros / Cons of national independent trials vs international trials: Numerous such trials launched simultaneously across hospitals in various countries address similar research questions. E.g. the search terms of “COVID AND Prone” to retrieve studies evaluating patient prone positioning on clinicaltrials.gov yielded 6,

31 and 44 results on the 5th of April, 5th of May and 5th of June respectively. This poses a major risk of redundant work, poor research resource allocation and incompleteness of some trials -- this happened all too often during previous epidemics (1-3). Researchers may waste time writing protocols from scratch while others already obtained funding or regulatory approvals. Although data generated by these numerous trials may ultimately be meta-analyzed, the time required for these numerous individual trials to publish and then compile data may be incompatible with the pressure of the epidemic. Several stakeholders called for a coordinated research effort, which should ideally take place at the international level (1).

However, setting up an international trial requires tremendous resources and time to finalize a unique protocol translated in various languages, coordinate all regulatory and ethical approvals, and conduct trials and data quality assessment in each country (4, 5). Such efforts have been conducted successfully for observational studies with the support of international research networks and scientific societies (6). However, the hurdles to set up a large scale international interventional randomized controlled trial are incompatible with the resources of an academic sponsor and are thus often restricted to pharmaceutical companies. When evaluating non-pharmacological interventions such as prone positioning, which has been proven to reduce mortality in mechanically ventilated patients with ARDS (7), the lack of foreseeable return on investment precludes such funding.

The *meta-trial* as a pragmatic solution for efficient pandemic clinical

research: in order to combine the benefits of international research with the fast setup of national trials, we propose to coordinate multiple national investigator-initiated trials in the form of a prospective meta-analysis. This so-called “*meta-trial*” consists of aggregating data from various national trials during the data collection (8). To study prone positioning in awake patients with COVID-19 pneumonia, 2 such trials were registered on clincaltrials.gov (one in the USA, the other in France) in mid-march 2020, and the investigators got in contact with 3 other groups planning trials with very similar inclusion and outcome criteria in Canada, Ireland and Spain, who all joined the *meta-trial* project. Each trial could be set up within a few weeks given the accelerated procedures in place during the COVID-19 outbreak. Planned sample sizes of individual trials ranged from 198 to 346 patients with the total planned inclusion of 1386 patients across 5 countries. Investigators and methodologists of all groups organized several web meetings to harmonize inclusion criteria and primary and secondary outcomes of the *meta-trial*. Given the planned sample sizes, an interim analysis plan was developed at the *meta-trial* level analyzing aggregated data every 200 patients. A memorandum of understanding and data sharing agreement were drafted (available upon request).

The *meta-trial* concept enables researchers to combine the agility of smaller national trials into a much larger international project in a short period of time (Table 1). *Meta-trial* interim analysis enables to detect a positive or negative response to the scientific question as soon as an adequate sample size is reached across several

countries, thus potentially speeding up the research process dramatically (9, 10). Adherence to methodological standards of individual trials represents a guarantee of a high level of overall final quality. Furthermore, by estimating the treatment effect across the various trials upfront, the *meta-trial* may provide stronger evidence in favor of external validity and replicability of the individual trials.

To the best of our knowledge, the *meta-trial* concept has never been experienced in real life across several countries, and feasibility uncertainties do exist. The present project may serve as a guidance for future research projects set up in a pandemic context.

How to join: the *meta-trial* is a living project in the sense that other groups can join as long as they adhere to the general principle and abide by ethical regulations in their country. Protocols and clinical record files are made available upon request by the core investigator group of the *meta-trial*. Readers are invited to contact authors for any additional information and to join the project (Awake.Prone.Meta.Trial@gmail.com).

REFERENCES :

- 1: Phua J, Weng L, Ling L, Egi M, Lim CM, Divatia JV, Shrestha BR, Arabi YM, Ng J, Gomersall CD, Nishimura M, Koh Y, Du B. Intensive care management of coronarvirus disease 2019 (COVID-19): Challenges and Recommendations. *Lancet Respir Med* 2020;8:506-517
- 2: Baden LR, Rubin EJ, Morrissey S, Farrar JJ, Drazen JM. We can do better — improving outcomes in the midst of an emergency. *N Engl J Med* 2017;377:1482-1484
- 3: Ivie RMJ, Vail EA, Wunsch H, Goldklang MP, Fowler R, Moitra VK. Patient eligibility for randomized controlled trials in critical care medicine: an international two-center observational study. *Crit Care Med* 2017;45:216-224
- 4: Rishu AH, Marinoff N, Julien L, Dumitrascu M, Marten N, Eggertson S, Willems S, Ruddell S, Lane D, Light B, Stelfox HT, Jouvett P, Hall R, Reynolds S, Daneman N, Fowler RA. Time required to initiate outbreak and pandemic observational research. *J Crit Care* 2017;40:7-10
- 5: Burns KEA, Rizvi L, Tan W, Marshall JC, Pope K. Participation of ICUs in critical care pandemic research: a province wide, cross-sectional survey. *Crit Care Med* 2013;41:1009-1016
- 6: Bellani G, Laffey JG, Pham T, Fan E, Brochard L, Esteban A, Gattinoni L, van Haren F, Larsson A, McAuley DF, Ranieri M, Rubenfeld G, Thompson BT, Wriggle H, Slutsky AS, Pesenti A. Epidemiology, patterns of care, and mortality for

patients with acute respiratory distress syndrome in intensive care units in 50 countries. JAMA 2016;315:788-800

7: Munshi L, Del Sorbo L, Adhikari NKJ, Hodgson CL, Wunsch H, Meade MO, Uleryk E, Mancebo J, Pesenti A, Ranieri M, Fan E. Prone position for acute respiratory distress syndrome. A systematic review and meta-analysis. Ann Am Thorac Soc 2017;14:S280-S288

8: Tavernier E, Trinquart L, Giraudeau B. Finding alternatives to the dogma of power based sample size calculation: is a fixed sample size prospective meta-experiment a potential alternative? Plos One 2016;11:e0158604

9: Simonsen L, Higgs E, Taylor RJ. Clinical research networks are key to accurate and timely assessment of pandemic clinical severity. Lancet Glob Health 2018;6:e956-e957

10: Memoli MJ. Pandemic research in the ICU: always be prepared. Crit Care Med 2013;41:1147-1148

Table 1: Comparison of the meta-trial concept to alternative designs			
	Individual trials followed by a retrospective meta-analysis	International single trial	<i>Meta-trial</i> : prospective international meta-analysis
Eligibility criteria for participants	Heterogenous between trials	Uniform within the trial	Similar between trials (may have some heterogeneity within clinical relevance)
Baseline data	Heterogenous between trials	Uniform within the trial	Common set of variables in data sharing agreement
Intervention details and how they were administered	Heterogenous between trials	Uniform within the trial	Uniform between trials: agreement between individual investigators to deliver same intervention
Pre-specified primary and secondary outcome measures	Heterogenous between trials	Uniform within the trial	Uniform between trials (investigators agree on a common set of outcomes)
Samples size and Interim analysis	Heterogenous between trials, interim analyses impossible at the meta-level	One sample size calculation for the trial, interim analyses possible	Meta-trial design transcends original sample size calculation, interim analyses possible at the meta-level
Randomization- sequence generation, stratification, allocation sequence, concealment and blinding	Heterogenous between trials	Centralized randomization	May differ for each site but fundamental randomization principles adhered to
Statistical methods	Heterogenous original analyses, meta-analysis on effect sizes to compute a summary effect	Uniform within the trial, adjustments possible	Uniform within the trials, meta-analysis on individual participant data, adjustments possible
Analysis populations: intention to treat, Per protocol, subgroups	Heterogenous between studies	Uniform within the trial	Uniform between the trials (agreement on uniform analysis population)
Data quality and safety monitoring	Each trialist is responsible for his or her trial	Centralized data monitoring	Each trialist is responsible for his or her trial
Funding	Multiple funding	Centralized funding	Multiple funding
Set-up time	Short	Long	Short
Time to completion	Long	Short	Short
Protocols	Multiple original protocols	One	Multiple original protocols followed by a meta-trial protocol
Ethics	Each trialist is responsible for his or her trial	Centralized submission process	Each trialist is responsible for his or her trial