



**HAL**  
open science

## Voiding Dysfunction after Colorectal Surgery for Endometriosis: A Systematic Review and Meta-analysis

Elie Vesale, Horace Roman, Gaby Moawad, Louise Benoit, Cyril Touboul, Emile Darai, Sofiane Bendifallah

► **To cite this version:**

Elie Vesale, Horace Roman, Gaby Moawad, Louise Benoit, Cyril Touboul, et al.. Voiding Dysfunction after Colorectal Surgery for Endometriosis: A Systematic Review and Meta-analysis. *Journal of Minimally Invasive Gynecology*, 2020, 27, pp.1490 - 1502.e3. 10.1016/j.jmig.2020.07.019 . hal-03493014

**HAL Id: hal-03493014**

**<https://hal.science/hal-03493014v1>**

Submitted on 7 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **VOIDING DYSFUNCTION AFTER COLORECTAL SURGERY FOR ENDOMETRIOSIS: A**  
2 **SYSTEMATIC REVIEW AND META-ANALYSIS**

3  
4 Elie Vesale (M.D)<sup>1,2,6</sup>, Horace Roman (M.D, PhD)<sup>4</sup>, Gaby Moawad (M.D)<sup>5</sup>, Louise Benoit  
5 (M.D)<sup>1,3</sup>, Cyril Touboul (M.D, PhD)<sup>1,2,3</sup>, Emile Darai(M.D, PhD)<sup>1,2,3</sup>, Sofiane Bendifallah(M.D,  
6 PhD)<sup>1,2,3</sup>

7 1. Department of Gynecology and Obstetrics, Tenon University Hospital, Assistance  
8 Publique des Hôpitaux de Paris (AP-HP), Sorbonne University, Paris, France.

9 2. Groupe de Recherche Clinique 6 (GRC6-UPMC) : Centre Expert En Endométriase  
10 (C3E)

11 3. UMR\_S938 Sorbonne University, Paris, France

12 4. Clinique Tivoli-Ducos, Bordeaux, France.

13 5. Department of Obstetrics and Gynecology, George Washington University School of  
14 Medicine and Health Sciences, Washington, D.C., USA.

15 6. Department of Gynecology and Obstetrics, Medical Center of the Sud-Francilien,  
16 Corbeil-Essonne, France

17 **Corresponding Author:**

18 Louise BENOIT (MD)

19 Department of Gynecology and Obstetrics, Tenon University Hospital

20 Adress: 4 rue de la Chine, 75020 Paris, France

21 Phone: + 33 1 56 01 54 10

22 Fax: + 33 1 56 01 73 17

23 E-mail: [louise.am.benoit@gmail.com](mailto:louise.am.benoit@gmail.com)

24

25 **Funding:** None

26 **Declaration of interest:** None

27 This manuscript has not been submitted elsewhere.

28 **Disclosure statement:** The authors declare that they have no conflicts of interest and  
29 nothing to disclose.

30 **Word count :** 2807 (<3000)

31

32 **Precis**

33 In this meta-analysis, we found that, rectal shaving causes less postoperative voiding  
34 dysfunction than discoid excision or segmental resection in colorectal surgery for  
35 endometriosis.

36 **Abstract**

37 Objective: Surgical management of deep endometriosis is associated with a high  
38 incidence of lower urinary tract dysfunction. The aim of the current systematic review and  
39 meta-analysis was to assess the rates of voiding dysfunction according to colorectal shaving,  
40 discoid excision and segmental resection for deep endometriosis.

41 Data sources: We performed a systematic review using bibliographic citations from  
42 PubMed, Clinical Trials.gov, Embase, Cochrane Library, and Web of Science databases.  
43 MeSH terms for colorectal endometriosis and voiding dysfunction were combined and  
44 restricted to the French and English languages. The final search was performed on August  
45 28, 2019. The outcome measured was the occurrence of postoperative voiding dysfunction.

46 Methods of study selection: Study Quality Assessment Tools were used to assess the  
47 quality of included studies. Studies rated as good and fair were included. Two reviewers  
48 independently assessed the quality of each included study, discrepancies were discussed  
49 and, if consensus was not reached, a third reviewer was consulted.

50 Tabulation, Integration and results: Out of 201 relevant published reports, 51 studies  
51 were ultimately reviewed systematically and 13 were included in the meta-analysis. Rectal  
52 shaving was statistically less associated with postoperative voiding dysfunction than  
53 segmental colorectal resection (OR 0.34, 95% CI 0.18-0.63,  $I^2= 0\%$ ,  $p=0.0006$ ) or discoid  
54 excision (OR 0.22, 95% CI 0.09-0.51,  $I^2 = 0\%$ ,  $p=0.0004$ ). No significant difference was  
55 noted when comparing discoid excision and segmental colorectal resection (OR 0.74, 95%  
56 CI 0.32-1.69,  $I^2= 29\%$ ,  $p=0.47$ ). Similarly, rectal shaving was associated with a lower risk of  
57 self-catheterization >1 month compared with segmental colorectal resection (OR 0.3, 95% CI  
58 0.14-0.66,  $I^2= 0\%$ ,  $p=0.003$ ). This outcome was no longer significant when comparing  
59 discoid excision and segmental colorectal resection (OR 0.72, 95% CI 0.4-1.31,  $I^2 = 63\%$ ,  
60  $p=0.28$ ).

61

62            Conclusion: Colorectal surgery for endometriosis has a significant impact on urinary  
63 function regardless of the technique. However, rectal shaving causes less postoperative  
64 voiding dysfunction than discoid excision or segmental resection.

65 **Key words**

66 Endometriosis; bowel involvement; meta-analysis; voiding dysfunction, gastrointestinal

67 endometriosis.

68 **Introduction**

69           Deep endometriosis (DE) surgery can involve the uterosacral ligaments, rectovaginal  
70 septum, vagina, pouch of Douglas, rectum, rectosigmoid junction, and parametrium. It can  
71 induce voiding dysfunction, especially in the context of bowel resection (segmental, discoid  
72 resection or shaving).

73           In this setting, the occurrence of postoperative voiding dysfunction ranges from 0-  
74 40% from one study to another (1) and is mainly due to iatrogenic injury of the pelvic  
75 anatomic nerves. Indeed, the bladder is innervated by the inferior hypogastric plexus  
76 originating from the superior hypogastric plexus (autonomous), the pelvic nerves  
77 (parasympathetic) and the hypogastric nerve (sympathetic). The ureter's sphincter is  
78 innervated by the pudendal nerve (somatic). The inferior hypogastric plexus can be injured  
79 at two different locations during surgery for colo-rectal endometriosis: during the dissection  
80 of the utero-sacral ligaments (inferior hypogastric nerve) or during the opening of the para-  
81 rectal space (hypogastric nerve). Although it may resolve spontaneously after one to three  
82 weeks, it can also persist for months, years or for a lifetime (2). Its occurrence has been  
83 reported to affect the quality of life in 20-30% of patients after DE surgery (3)(4) and there  
84 are few therapeutic options: mainly intermittent self-catheterization and neuromodulation  
85 (5).

86           For colorectal endometriosis, three main surgical approaches have been reported  
87 (6)(7): (i) a radical procedure allowing the complete removal of the endometriosis location by  
88 segmental resection (8)(9), and conservative surgery based on (ii) discoid resection or (iii)  
89 rectal shaving. Currently, pelvic surgeons agree that a nerve-sparing (NS) technique reduces  
90 the risk of persistent urinary retention due to iatrogenic injury of the pelvic autonomic nerves  
91 compared with the conventional (non-NS) technique (5). However, despite the NS technique,  
92 colorectal endometriosis surgery is still associated with a high rate of voiding dysfunction.

93           Therefore, the purpose of this systematic review and meta-analysis was to assess  
94 the occurrence of voiding dysfunction after surgery for DE with colorectal involvement



95 according to the surgical approach, i.e. rectal shaving, discoid excision or colorectal  
96 segmental resection. The impact of NS techniques were also assessed.

97 **Material and methods**

98 We performed a systematic review and meta-analysis following the  
99 recommendations from the Cochrane Collaboration (10) and in accordance with the  
100 PRISMA (Preferred Reported Items for Systematic Reviews and Meta-analyses)  
101 guidelines (11) . The protocol was approved beforehand (International Prospective  
102 Register of Systematic Reviews PROSPERO ID: 3528920).

103 *Electronic search*

104 The following databases were searched: PubMed, Clinical Trials.gov, Embase,  
105 Cochrane Library, and Web of Science databases. The major MeSH search terms used were  
106 (“endometriosis” or “deep infiltrating endometriosis” or “deep endometriosis” or “deep pelvic  
107 endometriosis” or “colorectal endometriosis”) and (“surgery for endometriosis” or  
108 “conservative management” or “radical management” or “colorectal resection” or “rectal  
109 shaving” or “full thickness resection” or “discoid excision”) AND “treatment”, “outcomes”,  
110 “short- and long-term results”, and “voiding dysfunction”. Studies in French or English  
111 published before April 1, 2019 were included. Abstracts were excluded if the authors did not  
112 provide adequate information when contacted. The final search of each database was  
113 performed on August 28, 2019.

114 Two reviewers (EV and SB) independently assessed the quality of each included  
115 study. In case of discrepancies a third reviewer was consulted (ED, HR). The percentages,  
116 odd ratios (ORs) and confidence intervals (CIs) of each study were obtained or calculated  
117 from existing data (12)(13).

118 *Outcome Measures*

119 The primary outcome was the occurrence of postoperative voiding dysfunction. As  
120 the voiding dysfunction rate varies according to the definition, we decided to first analyze the  
121 risk of voiding dysfunction regardless its definition and then to take a closer look at the risk of

122 self-catheterization lasting more than 1 month. Moreover, we analyzed the rate of voiding  
123 dysfunction according to the surgical management of the colorectum as this outcome varies  
124 greatly depending on conservative management or radical colorectal resection.

#### 125 *Risk of Bias*

126 The risk of bias was assessed using clarity of the statements, the eligibility criteria, size  
127 of the population, ethical considerations, representativeness, response rate, data collection,  
128 objective, definition of the objective, and consistency between research question and data  
129 reported.

130 The quality of the included studies was evaluated with the Study Quality Assessment  
131 Tools(14). Studies were considered as "satisfactory" when at least 70% of 12-14 assessment  
132 criteria were met, "fair" when 50% were respected, and "poor" when fewer than 50% of the  
133 criteria were fulfilled.

#### 134 *Statistical Analysis*

135 ORs and their CIs were derived from each study as the primary endpoint. Pooled  
136 response means, defined as the estimated overall mean difference (with 95% CI), were  
137 described using forest plots. For pooled response means, a p-value <0.05 was considered  
138 significant. If no observed events was reported by the study for a given outcome, a 0.001  
139 integer continuity correction was determined.

140 Statistical heterogeneity among the studies was assessed using Cochran's Q test and  
141 the  $I^2$  index. A  $I^2$  <50% or p-values of <0.1 signified that there was no significant  
142 heterogeneity. If no heterogeneity was observed, the fixed-effects model was adopted. In  
143 case of heterogeneity, the random-effects model was used for pooled estimates. If  $I^2$  > 50%  
144 (high heterogeneity), subgroup analyses were carried out when possible. If sufficient data  
145 were available, a predefined subgroup analyses for the primary outcome was undertaken.

146 All statistical analysis was performed on Review Manager (RevMan, IOS, version (5.3),  
147 Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2011). For the  
148 comparison of the nerve sparing techniques, the number of patients and percentages are  
149 presented. Comparison was performed using Fisher's exact test (qualitative variable) and  $p <$   
150 0.05 was considered statistically significant. For this analysis, statistical analysis was  
151 performed using R 3.6.0 software (<http://lib.stat.cmu.edu/R/CRAN/>).

152 **Results**

153 *Study selection*

154 A total of 201 studies fulfilling our electronic search criteria were identified. After  
155 screening by title and abstract and removing duplicate papers, 148 full text articles were  
156 assessed. Of these, one systemic review was excluded. From the 147 remaining papers, 79  
157 were excluded and among the 68 remaining papers, 17 were not eligible for the review  
158 (Figure 1). Among the 51 papers included in the review, 38 were not included in the meta-  
159 analysis because these studies did not compare different surgical techniques. The final result  
160 for the descriptive analysis included a total of 51 studies, 13 articles for the global meta-  
161 analysis (regardless of the definition) and five articles for the meta-analysis of self-  
162 catheterization > 1 month (Table 1).

163

164 *Study characteristics – Descriptive analysis*

165 Twenty-four prospective studies were identified; two randomized controlled trials, one  
166 case control study and 24 retrospective studies with a total 5962 patients undergoing surgery  
167 for DE involving the colorectum. Among them, 1734 (29.1%) underwent rectal shaving, 573  
168 (9.6%) discoid excision, and 3655 (61.3%) segmental resection. Voiding dysfunction was  
169 observed in 452 (7.6%) patients with 41 (2.4%), 43 (7.5%) and 368 (10.1%) in the rectal  
170 shaving, the discoid excision and the segmental resection groups, respectively (Table 1).

171 Eight studies (15)(16)(17)(18)(19)(20)(21)(22) included 1202 patients and defined  
172 voiding dysfunction as self-catheterization lasting at least one month after surgery. One  
173 (1.2%) patient was included in the shaving group, 0 in the discoid excision group, and 143  
174 (13.6%) in the segmental resection group (Supplementary Table 1).

175 Thirteen studies (23)(24)(25)(15)(26)(16)(27)(28)(19)(29)(30)(31)(32) including 1694  
176 patients (643 rectal shavings, 357 discoid excisions and 728 segmental resections) reported  
177 at least two of the three types of colorectal surgery and were thus included in the meta-  
178 analysis (Supplementary Table 2). Five studies (23)(15)(16)(28)(19) among them defined  
179 voiding dysfunction as self-catheterization > 21 days (Supplementary Table 3).

180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206

*Postoperative voiding dysfunction regardless of the definition*

When analyzing the occurrence of postoperative voiding dysfunction regardless of the definition, rectal shaving was statistically less associated with postoperative voiding dysfunction than segmental colorectal resection (OR 0.34, 95% CI 0.18-0.63,  $I^2= 0\%$ ,  $p=0.0006$ ) (Figure 2). This difference was no longer significant when comparing discoid excision and segmental colorectal resection (OR 0.74, 95% CI 0.32-1.69,  $I^2= 29\%$ ,  $p=0.47$ ) (Figure 2). Rectal shaving resulted in less voiding dysfunction than discoid excision (OR 0.22, 95% CI 0.09-0.51,  $I^2 = 0\%$ ,  $p=0.0004$ ) (Figure 2).

*Postoperative voiding dysfunction defined as self-catheterization > 21 days*

Rectal shaving was associated with a lower risk of prolonged self-catheterization (i.e. lasting for more than 21 days) compared with segmental colorectal resection (OR 0.3, 95% CI 0.14-0.66,  $I^2= 0\%$ ,  $p=0.003$ ) (Figure 3). This outcome was no longer significant when comparing discoid excision and segmental colorectal resection (OR 0.72, 95% CI 0.4-1.31,  $I^2 = 63\%$ ,  $p=0.28$ ) (Figure 3). As no studies included both shaving and discoid excision and defined voiding dysfunction as self-catheterization time >21 days, no comparison was evaluable for these techniques.

*Postoperative voiding dysfunction according to conservative versus radical surgery*

A comparison between conservative surgery (i.e. rectal shaving and discoid excision) and radical surgery (i.e. segmental colorectal resection) demonstrated that conservative strategy had a lower risk of postoperative voiding dysfunction (Risk Difference -0.04, 95% CI -0.07; -0.02,  $I^2= 23\%$ ,  $p=0.001$ ). (Figure 4)

*Nerve-sparing technique*

207 Twenty-six studies including 3120 patients (1402 conservative and 1718 radical  
208 surgeries) clearly reported an NS technique while 25 studies including 2843 patients (1001  
209 conservative and 1842 radical surgeries) did not specifically mention an NS technique (Table  
210 2).

211 Sixty-two (4.4%) and 18 (1.8%) patients, respectively, had postoperative voiding  
212 dysfunction after conservative surgery with an NS and with a non-NS technique ( $p=0.03$ ). For  
213 radical surgery, 222 (12.9%) patients and 208 (11.3%) patients had postoperative voiding  
214 dysfunction after an NS and a non-NS technique, respectively ( $p=0.15$ ).

215

216

#### 217 *Duration of postoperative self-catheterization*

218 The mean time of self-catheterization could not be estimated for each group as not all  
219 the studies reported the exact catheterization time. The median time of self-catheterization  
220 after segmental colorectal resection according to eight studies  
221 (33)(34)(35)(36)(37)(21)(38)(39) was  $31.5 \pm 6.1$  days.

222

223 **Discussion**

224 We report a systematic review and meta-analysis focusing on the occurrence of  
225 voiding dysfunction after surgery for deep endometriosis (DE) with colorectal involvement  
226 comparing three surgical approaches. The estimated overall rate of voiding dysfunction was  
227 7.6% with a prevalence of 2.4%, 7.5% and 10.1% after rectal shaving, discoid excision and  
228 segmental colorectal resection, respectively. Self-catheterization > 1 month mainly occurred  
229 after segmental colorectal resection (13.6%). Evaluating urinary function preoperatively is  
230 therefore of paramount importance in digestive endometriosis.

231 The assessment of voiding dysfunction before and after colorectal resection for  
232 endometriosis surgery is a major issue in daily practice. However, the current evidence-  
233 based literature is marked by a major weakness due to the lack of consensus in the definition  
234 used. For instance, voiding dysfunction may correspond to a variety of clinical conditions  
235 such as urinary retention, bladder atonia, incomplete urinary bladder emptying requiring  
236 intermittent catheterization after postoperative day 7 or with a post-voiding bladder volume of  
237 more than 100ml (15)(2)(40). Likewise, some authors define the duration of voiding  
238 dysfunction as transient when self-catheterization lasts under 30 days (2) or up to 6 weeks,  
239 while others extend this definition to 6 months (33)(41). All these definitions which should be  
240 reporting the same outcome render interpretation and comparison of clinical data somewhat  
241 difficult and heterogeneous. Moreover, few reports distinguish preoperative from *de novo*  
242 voiding dysfunction which can occur in 7.8% of their patients with colorectal  
243 endometriosis. To overcome such a limitation, we decided to present the results considering  
244 (i) the voiding dysfunction whatever the definition used (tables 2,3,4), (ii) the voiding  
245 dysfunction based on the definition of self-catheterization > 1 month, and also (iii) the use of  
246 an NS technique.

247 Regardless of the definition, rectal shaving appears to result in a lower risk of  
248 voiding dysfunction than discoid excision (OR 0.22, 95% CI 0.09-0.51,  $I^2 = 0\%$ ,  $p=0.0004$ )  
249 and segmental colorectal resection (OR 0.34, 95% CI 0.18-0.63,  $I^2 = 0\%$ ,  $p= 0.0006$ ).



250 Interestingly, this difference was no longer significant when comparing discoid excision  
251 and segmental resection, although considerable heterogeneity was observed between the  
252 studies (OR 0.74, 95% CI 0.32-1.69,  $I^2= 29\%$ ,  $p=0.47$ ). Self-catheterization >1 month  
253 occurred in 13.6% of the patients and almost only after segmental colorectal resection.  
254 Considering this definition, we observed that the risk was significantly lower after rectal  
255 shaving compared to segmental resection (OR 0.3, 95% CI 0.14-0.66,  $I^2= 0\%$ ,  $p=0.003$ ).  
256 The difference was not significant when discoid resection was compared to segmental  
257 resection although a high heterogeneity among studies was observed (OR 0.72, 95% CI  
258 0.4-1.31,  $I^2 = 63\%$ ,  $P=0.28$ ). The most frequent type of surgery was segmental resection  
259 (61.3% of patients) since this technique was most historical surgical technics and can be  
260 proposed no matter the lesion size, the circumference involved and the presence of  
261 multifocal lesions.

262 Colorectal endometriosis is a source of urinary dysfunction in itself due to infiltration  
263 of the hypogastric plexus. Urinary function should thus be systematically evaluated  
264 preoperatively to distinguish preexisting from *de novo* voiding dysfunction subsequent to  
265 surgery (17)(42)(35). Considering the high heterogeneity of the rates of voiding  
266 dysfunction reported in the various studies (ranging from 0 to 40%) (Table 1), future  
267 studies should include evaluation of preoperative urinary symptoms by urodynamic testing  
268 and electrophysiology exam to determine whether intraoperative nerve damage of the  
269 pelvic plexus is the cause. In this setting, Ballester et al. conducted a comparative study  
270 between a group of patients with DE and a control group (35) based on a validated  
271 questionnaire (International Prostate Symptom Score). They showed a significantly higher  
272 incidence of preoperative voiding dysfunction in the DE group.

273 Several authors hypothesized (43) the need of systematic preoperative urodynamic  
274 investigation in the context of DE as the rate of preoperative voiding dysfunction can be  
275 more than 25%. However, identification of patients who might benefit from preoperative  
276 urodynamic testing remain debatable. The majority of published studies in the current

277 systematic review did not include preoperative assessment of lower urinary tract  
278 dysfunction therefore we were unable to refine the assessment of voiding dysfunction  
279 rates. Moreover, Bonneau et al reported that half of their patients with preoperative voiding  
280 dysfunction had normal urodynamic test values (2).

281 NS surgery for DE with colorectal involvement has been shown to decrease the  
282 incidence of urinary side effects (44)(45). De Resende Junior et al. recently conducted a  
283 meta-analysis (41) including four studies that strictly compared NS and non-NS techniques  
284 (33)(46)(47)(48). The authors found a common OR of 0.19 [95%CI: 0.03–1.17; (I  
285 21/450.20%; P1/40.09)] for the need of self-catheterization at discharge in the NS group  
286 compared to the conventional non-NS technique.

287 In the current review, 26 studies clearly mentioned an NS technique compared to 25  
288 studies where it was unclear whether this technique was used. In contrast to De Resende  
289 Junior et al.'s review, we found no difference in postoperative voiding dysfunction  
290 according to the use of an NS technique. However, the interpretation of these results is  
291 compromised because: i) the studies did not aim to compare the techniques; ii) a study  
292 which does not mention an NS technique does not necessarily exclude NS surgery; iii)  
293 surgery with NS are usually more invasive procedures with a high extent of resection; iv)  
294 there is a lack of objective and uniform criteria to define postoperative voiding dysfunction  
295 and; v) surgical skills may have improved over time.

296 Resende et al (41) could not detect significant benefits of NS compared with  
297 conventional non-NS surgery in terms of relative risk (RR) of urinary retention at discharge.  
298 Nevertheless, they found a statistically significant advantage of NS with respect to RR of  
299 persistent urinary retention by the need for self-catheterization for 3 months or more even  
300 if no clear criteria (e.g., minimum residual volume) was presented to define the need for  
301 self-catheterization after surgery. Although several NS techniques have been reported for  
302 delayed functional complications (e.g. urinary retention, constipation, and sexual  
303 dysfunction) due to autosomal nerve damage of the bowel, they have not been 100%  
304 successful (45). The surgeon should consider both of these complications when choosing

305 the most appropriate surgical procedure and provide information on functional outcomes to  
306 patients treated for colorectal endometriosis (29). Indeed, identifying and sparing the pelvic  
307 nerves should be standard practice in 2020 as it is accepted that the more nerves are  
308 preserved, the fewer sexual, rectal and bladder disorders you cause (49).

309 The strengths of this review include the use of an exhaustive search strategy applied  
310 to five different databases. In addition, it seems to be the first to report the risk of  
311 postoperative voiding dysfunction comparing three surgical approaches for DE with  
312 colorectal involvement. Methodological limitations such as differences between criteria  
313 used for defining the voiding dysfunction may limit the quality of the available evidence.  
314 Furthermore, the absence of taking preoperative voiding dysfunction into account should  
315 be mentioned as a limitation. However, most of the articles included in this review lacked  
316 data on preoperative voiding dysfunction and we were thus unable to know if the  
317 postoperative dysfunction was due to operative damage. In addition, the heterogeneity  
318 between some studies may limit the accuracy and the validity of the results. Finally, for the  
319 sake of completeness, some comparisons, in particular that of NS procedures, may be  
320 rough estimates rather than accurate comparisons because of the lack of studies  
321 comparing NS and classical surgery in the context of colorectal resection for DE.

322 **Conclusion**

323 Bowel surgery for DE has a significant impact on urinary function regardless of the  
324 technique and patients should be clearly informed pre-operatively. Rectal shaving causes  
325 less postoperative voiding dysfunction than discoid excision or segmental resection. It is  
326 therefore essential to detect preoperative urinary symptoms and to define a high-risk group  
327 for which urodynamic testing could be helpful. Moreover, a consensual definition of  
328 postoperative voiding dysfunction as well as the indication for self-catheterization should be  
329 used to develop a predictive tool which would be useful when counseling patients and to  
330 determine clinical trial eligibility.

331 **References**

- 332 1. [Urinary functional disorders bound to deep endometriosis and to its treatment: review of  
333 the literature]. - PubMed - NCBI [Internet]. [cited 2020 Mar 29]. Available from:  
334 <https://www.ncbi.nlm.nih.gov/pubmed/?term=24831568>
- 335 2. Bonneau C, Zilberman S, Ballester M, Thomin A, Thomassin-Naggara I, Bazot M, et al.  
336 Incidence of pre- and postoperative urinary dysfunction associated with deep infiltrating  
337 endometriosis: relevance of urodynamic tests and therapeutic implications. *Minerva*  
338 *Ginecol.* 2013 Aug;65(4):385–405.
- 339 3. Byrne D, Curnow T, Smith P, Cutner A, Saridogan E, Clark TJ, et al. Laparoscopic  
340 excision of deep rectovaginal endometriosis in BSGE endometriosis centres: a  
341 multicentre prospective cohort study. *BMJ Open.* 2018 09;8(4):e018924.
- 342 4. Roman H, Chanavaz-Lacheray I, Ballester M, Bendifallah S, Touleimat S, Tuech J-J, et  
343 al. High postoperative fertility rate following surgical management of colorectal  
344 endometriosis. *Hum Reprod Oxf Engl.* 2018 01;33(9):1669–76.
- 345 5. Nyangoh Timoh K, Canlorbe G, Verollet D, Peyrat L, Ballester M, Amarenco G, et al.  
346 Contribution of sacral neuromodulation to manage persistent voiding dysfunction after  
347 surgery for deep infiltrating endometriosis with colorectal involvement: preliminary  
348 results. *Eur J Obstet Gynecol Reprod Biol.* 2015 Jul;190:31–5.
- 349 6. Abrão MS, Petraglia F, Falcone T, Keckstein J, Osuga Y, Chapron C. Deep  
350 endometriosis infiltrating the recto-sigmoid: critical factors to consider before  
351 management. *Hum Reprod Update.* 2015 Jun;21(3):329–39.
- 352 7. Donnez J, Nisolle M, Gillerot S, Smets M, Bassil S, Casanas-Roux F. Rectovaginal  
353 septum adenomyotic nodules: a series of 500 cases. *Br J Obstet Gynaecol.* 1997  
354 Sep;104(9):1014–8.
- 355 8. Nezhat C, Nezhat F, Ambroze W, Pennington E. Laparoscopic repair of small bowel and  
356 colon. A report of 26 cases. *Surg Endosc.* 1993 Apr;7(2):88–9.
- 357 9. Darwish B, Roman H. Surgical treatment of deep infiltrating rectal endometriosis: in favor  
358 of less aggressive surgery. *Am J Obstet Gynecol.* 2016;215(2):195–200.
- 359 10. idostatistics. Higgins, J. P. T., & Green, S. (Eds.) (2008). *Cochrane handbook for*  
360 *systematic reviews of interventions* [Internet]. IDoStatistics. 2016 [cited 2019 May 4].

- 361 Available from: <https://idostatistics.com/higgins-green-2008-cochrane-handbook->  
362 [systematic-reviews-interventions/](https://idostatistics.com/higgins-green-2008-cochrane-handbook-systematic-reviews-interventions/)
- 363 11. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The  
364 PRISMA statement for reporting systematic reviews and meta-analyses of studies that  
365 evaluate healthcare interventions: explanation and elaboration. *BMJ*. 2009 Jul  
366 21;339:b2700.
- 367 12. Luo D, Wan X, Liu J, Tong T. Optimally estimating the sample mean from the sample  
368 size, median, mid-range, and/or mid-quartile range. *Stat Methods Med Res*.  
369 2018;27(6):1785–805.
- 370 13. Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median,  
371 range, and the size of a sample. *BMC Med Res Methodol*. 2005 Apr 20;5:13.
- 372 14. Study Quality Assessment Tools | National Heart, Lung, and Blood Institute (NHLBI)  
373 [Internet]. [cited 2020 Jun 29]. Available from: [https://www.nhlbi.nih.gov/health-](https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools)  
374 [topics/study-quality-assessment-tools](https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools)
- 375 15. Jayot A, Nyangoh Timoh K, Bendifallah S, Ballester M, Darai E. Comparison of  
376 Laparoscopic Discoid Resection and Segmental Resection for Colorectal Endometriosis  
377 Using a Propensity Score Matching Analysis. *J Minim Invasive Gynecol*. 2018  
378 Apr;25(3):440–6.
- 379 16. Roman H, Milles M, Vassilieff M, Resch B, Tuech J-J, Huet E, et al. Long-term functional  
380 outcomes following colorectal resection versus shaving for rectal endometriosis. *Am J*  
381 *Obstet Gynecol*. 2016 Dec;215(6):762.e1-762.e9.
- 382 17. Ballester M, Belghiti J, Zilberman S, Thomin A, Bonneau C, Bazot M, et al. Surgical and  
383 clinical impact of extraserosal pelvic fascia removal in segmental colorectal resection for  
384 endometriosis. *J Minim Invasive Gynecol*. 2014 Dec;21(6):1041–8.
- 385 18. Ruffo G, Scopelliti F, Scioscia M, Ceccaroni M, Mainardi P, Minelli L. Laparoscopic  
386 colorectal resection for deep infiltrating endometriosis: analysis of 436 cases. *Surg*  
387 *Endosc*. 2010 Jan;24(1):63–7.
- 388 19. Stepniewska A, Pomini P, Guerriero M, Scioscia M, Ruffo G, Minelli L. Colorectal  
389 endometriosis: benefits of long-term follow-up in patients who underwent laparoscopic  
390 surgery. *Fertil Steril*. 2010 May 1;93(7):2444–6.

- 391 20. Ferrero S, Anserini P, Abbamonte LH, Ragni N, Camerini G, Remorgida V. Fertility after  
392 bowel resection for endometriosis. *Fertil Steril*. 2009 Jul;92(1):41–6.
- 393 21. Ghezzi F, Cromi A, Ciravolo G, Rampinelli F, Braga M, Boni L. A new laparoscopic-  
394 transvaginal technique for rectosigmoid resection in patients with endometriosis. *Fertil*  
395 *Steril*. 2008 Nov;90(5):1964–8.
- 396 22. Mereu L, Ruffo G, Landi S, Barbieri F, Zaccoletti R, Fiaccavento A, et al. Laparoscopic  
397 treatment of deep endometriosis with segmental colorectal resection: short-term  
398 morbidity. *J Minim Invasive Gynecol*. 2007 Aug;14(4):463–9.
- 399 23. Abo C, Moatassim S, Marty N, Saint Ghislain M, Huet E, Bridoux V, et al. Postoperative  
400 complications after bowel endometriosis surgery by shaving, disc excision, or segmental  
401 resection: a three-arm comparative analysis of 364 consecutive cases. *Fertil Steril*.  
402 2018;109(1):172-178.e1.
- 403 24. Hudelist G, Aas-Eng MK, Birsan T, Berger F, Sevelde U, Kirchner L, et al. Pain and  
404 fertility outcomes of nerve-sparing, full-thickness disk or segmental bowel resection for  
405 deep infiltrating endometriosis-A prospective cohort study. *Acta Obstet Gynecol Scand*.  
406 2018 Dec;97(12):1438–46.
- 407 25. Bourdel N, Comptour A, Bouchet P, Gremeau A-S, Pouly J-L, Slim K, et al. Long-term  
408 evaluation of painful symptoms and fertility after surgery for large rectovaginal  
409 endometriosis nodule: a retrospective study. *Acta Obstet Gynecol Scand*. 2018  
410 Feb;97(2):158–67.
- 411 26. Segmental and Discoid Resection are Preferential to Bowel Shaving for Medium-Term  
412 Symptomatic Relief in Patients With Bowel Endometriosis. - PubMed - NCBI [Internet].  
413 [cited 2020 Mar 29]. Available from:  
414 <https://www.ncbi.nlm.nih.gov/pubmed/?term=27544881>
- 415 27. Diguisto C, Hébert T, Paternotte J, Kellal I, Marret H, Ouldamer L, et al. [Robotic assisted  
416 laparoscopy: comparison of segmentary colorectal resection and shaving for colorectal  
417 endometriosis]. *Gynecol Obstet Fertil*. 2015 Apr;43(4):266–70.
- 418 28. Fanfani F, Fagotti A, Gagliardi ML, Ruffo G, Ceccaroni M, Scambia G, et al. Discoid or  
419 segmental rectosigmoid resection for deep infiltrating endometriosis: a case-control  
420 study. *Fertil Steril*. 2010 Jul;94(2):444–9.

- 421 29. Roman H, Loisel C, Resch B, Tuech JJ, Hochain P, Leroi AM, et al. Delayed functional  
422 outcomes associated with surgical management of deep rectovaginal endometriosis with  
423 rectal involvement: giving patients an informed choice. *Hum Reprod Oxf Engl*. 2010  
424 Apr;25(4):890–9.
- 425 30. Brouwer R, Woods RJ. Rectal endometriosis: results of radical excision and review of  
426 published work. *ANZ J Surg*. 2007 Jul;77(7):562–71.
- 427 31. Jatan AK, Solomon MJ, Young J, Cooper M, Pathma-Nathan N. Laparoscopic  
428 management of rectal endometriosis. *Dis Colon Rectum*. 2006 Feb;49(2):169–74.
- 429 32. Mohr C, Nezhat FR, Nezhat CH, Seidman DS, Nezhat CR. Fertility considerations in  
430 laparoscopic treatment of infiltrative bowel endometriosis. *JSLs*. 2005 Mar;9(1):16–24.
- 431 33. Che X, Huang X, Zhang J, Xu H, Zhang X. Is nerve-sparing surgery suitable for deeply  
432 infiltrating endometriosis? *Eur J Obstet Gynecol Reprod Biol*. 2014 Apr;175:87–91.
- 433 34. Boileau L, Laporte S, Bourgaux J-F, Rouanet J-P, Filleron T, Mares P, et al.  
434 [Laparoscopic colorectal resection for deep pelvic endometriosis: Evaluation of post-  
435 operative outcome]. *J Gynecol Obstet Biol Reprod (Paris)*. 2012 Apr;41(2):128–35.
- 436 35. Ballester M, Chereau E, Dubernard G, Coutant C, Bazot M, Daraï E. Urinary dysfunction  
437 after colorectal resection for endometriosis: results of a prospective randomized trial  
438 comparing laparoscopy to open surgery. *Am J Obstet Gynecol*. 2011 Apr;204(4):303.e1-  
439 6.
- 440 36. Dousset B, Leconte M, Borghese B, Millischer A-E, Roseau G, Arkwright S, et al.  
441 Complete surgery for low rectal endometriosis: long-term results of a 100-case  
442 prospective study. *Ann Surg*. 2010 May;251(5):887–95.
- 443 37. Dubernard G, Rouzier R, David-Montefiore E, Bazot M, Daraï E. Urinary complications  
444 after surgery for posterior deep infiltrating endometriosis are related to the extent of  
445 dissection and to uterosacral ligaments resection. *J Minim Invasive Gynecol*. 2008  
446 Apr;15(2):235–40.
- 447 38. Seracchioli R, Poggioli G, Pierangeli F, Manuzzi L, Gualerzi B, Savelli L, et al. Surgical  
448 outcome and long-term follow up after laparoscopic rectosigmoid resection in women  
449 with deep infiltrating endometriosis. *BJOG Int J Obstet Gynaecol*. 2007 Jul;114(7):889–  
450 95.



- 451 39. Darai E, Thomassin I, Barranger E, Detchev R, Cortez A, Houry S, et al. Feasibility and  
452 clinical outcome of laparoscopic colorectal resection for endometriosis. *Am J Obstet*  
453 *Gynecol.* 2005 Feb;192(2):394–400.
- 454 40. Darai E, Dubernard G, Coutant C, Frey C, Rouzier R, Ballester M. Randomized trial of  
455 laparoscopically assisted versus open colorectal resection for endometriosis: morbidity,  
456 symptoms, quality of life, and fertility. *Ann Surg.* 2010 Jun;251(6):1018–23.
- 457 41. de Resende JAD, Cavalini LT, Crispi CP, de Freitas Fonseca M. Risk of urinary retention  
458 after nerve-sparing surgery for deep infiltrating endometriosis: A systematic review and  
459 meta-analysis. *Neurourol Urodyn.* 2017;36(1):57–61.
- 460 42. Kondo W, Bourdel N, Tamburro S, Cavoli D, Jardon K, Rabischong B, et al.  
461 Complications after surgery for deeply infiltrating pelvic endometriosis. *BJOG Int J Obstet*  
462 *Gynaecol.* 2011 Feb;118(3):292–8.
- 463 43. [Urinary functional and urodynamic preoperative evaluation of patients with deep pelvic  
464 surgical endometriosis: about 12 cases]. - PubMed - NCBI [Internet]. [cited 2020 Mar 29].  
465 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/?term=18494148>
- 466 44. The LANN technique to reduce postoperative functional morbidity in laparoscopic radical  
467 pelvic surgery. - PubMed - NCBI [Internet]. [cited 2020 Mar 29]. Available from:  
468 <https://www.ncbi.nlm.nih.gov/pubmed/?term=16310695>
- 469 45. Landi S, Ceccaroni M, Perutelli A, Allodi C, Barbieri F, Fiaccavento A, et al. Laparoscopic  
470 nerve-sparing complete excision of deep endometriosis: is it feasible? *Hum Reprod Oxf*  
471 *Engl.* 2006 Mar;21(3):774–81.
- 472 46. Kavallaris A, Mebes I, Evagyelinos D, Dafopoulos A, Beyer DA. Follow-up of  
473 dysfunctional bladder and rectum after surgery of a deep infiltrating rectovaginal  
474 endometriosis. *Arch Gynecol Obstet.* 2011 May;283(5):1021–6.
- 475 47. Ceccaroni M, Clarizia R, Bruni F, D'Urso E, Gagliardi ML, Roviglione G, et al. Nerve-  
476 sparing laparoscopic eradication of deep endometriosis with segmental rectal and  
477 parametrial resection: the Negrar method. A single-center, prospective, clinical trial. *Surg*  
478 *Endosc.* 2012 Jul;26(7):2029–45.
- 479 48. Volpi E, Ferrero A, Sismondi P. Laparoscopic identification of pelvic nerves in patients  
480 with deep infiltrating endometriosis. *Surg Endosc.* 2004 Jul;18(7):1109–12.

481 49. Uccella S, Gisone B, Serati M, Biasoli S, Marconi N, Angeretti G, et al. Functional  
482 outcomes of nerve-sparing laparoscopic eradication of deep infiltrating endometriosis: a  
483 prospective analysis using validated questionnaires. Arch Gynecol Obstet.  
484 2018;298(3):639–47.

485

486

487 Tables

488 Table 1: Studies reporting voiding dysfunction rate after colorectal surgery for deep

489 | endometriosis

490 Table 2: Nerve sparing versus non-nerve sparing technique in colorectal surgery for deep

491 endometriosis

492 Figure legends

493 Figure 1: Flow chart of the eligibility of studies for inclusion in the meta-analysis

494

495 Figure 2. Meta-analysis of voiding dysfunction, no matter the definition after colorectal  
496 surgery for deep endometriosis.

497 1.1. Shaving versus segmental resection no matter the definition of voiding dysfunction

498 1.2. Disc excision versus segmental resection no matter the definition of voiding dysfunction

499 1.3. Shaving versus disc excision no matter the definition of voiding dysfunction

500

501 Figure 3. Meta-analysis of voiding dysfunction defined by self-catheterization at 3 weeks after  
502 colorectal surgery for deep endometriosis.

503 2.1. Shaving versus segmental resection

504 2.2. Disc excision versus segmental resection


505




506 Figure 4. Meta-analysis of voiding dysfunction after radical versus conservative colorectal  
507 surgery for deep endometriosis



508 3.1. Voiding dysfunction, no matter the definition



509 4.1. Voiding dysfunction, defined by self-catheterization at 3 weeks

Figure 1: Eligibility of studies for inclusion in the meta-analysis











Study Author (year)	Type	n analyzed	n Shaving	Voiding dysfunction if shaving n(%)	n Disc excision	Voiding dysfunction if disc excision n(%)	n conservative surgery	Voiding dysfunction if conservative surgery excision n(%)	n Segmental resection	Voiding dysfunction if segmental resection n(%)	Definition of voiding dysfunction	Self-catherization time	Inclusion period	Nerve sparing surgery
<b>ABO 2018 * 29307394</b>	Retrospective	364	145	8 (5.5)	80	16(20)	225	24(10.7)	139	20(4.3)	Transitory bladder atony requiring auto-catheterization	3 weeks to 6 months	June 2009 - December 2015	Yes
<b>ERDEM 2018 29664797</b>	Retrospective	51	NC	NC	NC	NC	NC	NC	51	2(3)	Transient urinary retention	NR	June 2002 – May 2011	Yes
<b>HUDELIST 2018 * 30080244</b>	Prospective	134	NC	NC	32	3 (9.4)	32	3(9.4)	102	6(5.9)	urinary retention	NR	March 2011 - August 2016	Yes
<b>BOURDEL 2017* 29143306</b>	Retrospective	195	172	6(3.5)	NC	NC	172	6(3.5)	23	1(4.3)	urinary retention	NR	January 2000 - June 2013	NR
<b>JAYOT 2017 * 28987649</b>	Prospective	98	NC	NC	31	0(0)	31	0(0)	31	7(22)	persistent voiding dysfunction	>30 days	January 2015 - September 2016	Yes
<b>RENNER 2017 28374101</b>	Retrospective	113	NC	NC	NC	NC	NC	NC	112	15(15.3)	Postoperative residual urine >100 mL	NR	2001–2010	NR
<b>ROMAN 2017 28139235</b>	Prospective	111	NC	NC	111	10(9)	111	10(9)	NC	NC	Transitory bladder atony requiring auto-catheterization	21 to 180 days	June 2009 - June 2016	NR
<b>MILLOCHAU 2017 28893658</b>	Retrospective	20	NC	NC	20	4(19)	20	4(19)	NC	NC	Transitory bladder atony requiring self-catheterization	>7 days	March 2011 - December 2016	Yes
<b>MARTY 2017 28673871</b>	Retrospective	110	110	2(1.8)	NC	NC	110	2(1.8)	NC	NC	bladder atony requiring daily self-catheterization	>3 weeks	December 2012 - December 2016	NR
<b>AFORS 2016 * 27544881</b>	Retrospective	92	57	1(1.75)	15	2(13.3)	72	3(4.2)	30	2(6.6)	urinary retention which resolved spontaneously within 15 days	NC	January 2010 - September 2012	Yes
<b>ROMAN 2016 * 27393269</b>	Retrospective	70	46	1(2.2)	NC	NC	46	1(2.2)	25	3(12)	Bladder atony requiring daily catheterization	> 1 month	January 2005 - January 2010	Yes
<b>ROMAN 2016 26474734</b>	Prospective	103	NC	NC	NC	NC	NC	NC	103	5(4.9)	transitory urinary dysfunction requiring systematic daily catheterization	NR	June 2009 - November 2014	Yes
<b>ROMAN 2016 27565263</b>	Retrospective	122	122	8(6.6)	NC	NC	122	8(6.6)	NC	NC	bladder postvoiding volume >100 mL on	6 +/- 4 weeks	June 2009 - September 2014	Yes

											the day of discharge from the hospital			
<b>AKLADIOS 2015 25109779</b>	Retrospective	41	NC	NC	NC	NC	NC	NC	41	4(9.7)	transient urinary retention requiring self-catheterization	NR	2004 – 2011	Yes
<b>DIGUISTO 2015 * 25813434</b>	Retrospective	28	22	3 (13.6)	NC	NC	22	3 (13.6)	6	1(16.7)	Self-catheterization at discharge from the hospital	NR	December 2011 - June 2014	NR
<b>ROMAN 2015 26347968</b>	Prospective	50	NC	NC	50	8(16)	50	8(16)	NC	NC	Bladder atony requiring self-catheterization	3 weeks to 6 months	June 2009 - November 2014	Yes
<b>MEULEMAN 2014 23579578</b>	Prospective	76	NC	NC	NC	NC	NC	NC	76	2(2.6)	bladder atony	NR	September 2006 - October 2008	NR
<b>FLEISCH 2014 24407554</b>	Retrospective	4	NC	NC	NC	NC	NC	NC	4	2(50)	transient postoperative urinary retention	NR	2008 – 2010	Yes
<b>RUFFO 2014 24877097</b>	Retrospective	900	NC	NC	NC	NC	NC	NC	900	114(14.7)	Dysuria	NR	January 2002 - December 2010	NR
<b>BALLESTER 2014 24858943</b>	Prospective	227	NC	NC	NC	NC	NC	NC	227	45(22)	Voiding dysfunction requiring self-catheterization	>1 month	2001-2011	Yes
<b>Che 2014 24576484</b>	Prospective	21	NC	NC	NC	NC	NC	NC	21	9(4.3)	patient requiring self-catheterization postoperatively	56 days (range: 25 days to eight months)	September 2010 - July 2012	Yes
<b>MANGLER 2014 24726619</b>	Prospective	71	NC	NC	NC	NC	NC	NC	71	2(2.8)	residual urine at discharge	NR	September 2004 - December 2012	Yes
<b>BOILEAU 2012 22071018</b>	Retrospective	23	NC	NC	NC	NC	NC	NC	23	2(8.9)	De novo dysuria requiring self-catheterization	15 days	2005 – 2011	NR
<b>MEULEMAN 2011 22729087</b>	Retrospective	45	NC	NC	NC	NC	NC	NC	45	1(2.2)	Atonic neurogenic urinary bladder necessitating intermittent urethral catheterization	first 10 weeks after the operation	September 2004 - September 2006	Yes
<b>WOLTHUIS 2011 21427115</b>	Prospective	21	NC	NC	NC	NC	NC	NC	21	1(4.8)	atonic neurogenic urinary bladder necessitating intermittent urethral catheterization	NR	September 2009 - September 2010	NR
<b>BALLESTER 2011 21256472</b>	RCT	52	NC	NC	NC	NC	NC	NC	52	15(29)	Self-catheterization or Dysuria	3 months	January 2006 - December 2008	Yes
<b>DONNEZ 2010 20547557</b>	Prospective	500	500	4(0.8)	NC	NC	500	4(0.8)	NC	NC	urinary retention	NR	2004-2008	Yes
<b>FANFANI 2010 * 19394600</b>	Case-control	136	NC	NC	48	0(0)	48	0(0)	88	13(14.7)	Urinary retention after 30 days	NR	January 2003 - December 2007	NR

<b>DOUSSET 2010</b> <b>20395847</b>	Prospective	100	NC	NC	NC	NC	NC	NC	100	16(16)	A transient peripheral neurogenic bladder defined when intermittent self-catheterization was required after hospital discharge within 15 days	6 weeks (range: 3–18 weeks)	1995-2003	Yes
<b>RUFFO 2010</b> <b>19466489</b>	Retrospective	436	NC	NC	NC	NC	NC	NC	436	71(19.9)	urinary retention with need for self-catheterization at 1 month	>1 month	January 2002 - March 2008	Yes
<b>STEPNIEWSKA 2010 *</b> <b>19836731</b>	Prospective	100	40	0(0)	NC	NC	40	0(0)	60	3(5)	urine retention for >1 month	>1 month	2000 – 2005	Yes
<b>ROMAN 2010 *</b> <b>20106836</b>	Retrospective	41	NC	NC	16	0(0)	16	0(0)	25	2(8)	Bladder dysfunction requiring systematic or occasional catheterization .12 months after surgery	NR	January 2005 - June 2008	Yes
<b>FERRERO 2009</b> <b>18684451</b>	Prospective	46	NC	NC	NC	NC	NC	NC	46	2(4.3)	Urinary retention after 30 d from surgery	>1 month	May 1999 - May 2006	Yes
<b>LANDI 2008</b> <b>18656048</b>	Retrospective	35	NC	NC	35	0(0)	35	0(0)	NC	NC	NR	NC	January 2003 - September 2006	Yes
<b>BENBARA 2008</b> <b>19019719</b>	Retrospective	50	NC	NC	NC	NC	NC	NC	50	10(30.4)	Post-operative dysuria spontaneously disappeared between 8 days and 1 year	NR	1998 – 2005	No
<b>DUBERNAND 2008</b> <b>18313000</b>	Retrospective	52	NC	NC	NC	NC	NC	NC	52	12(23)	acute retention requiring self-catheterization of bladder	85 days (range: 10–420)	April 2001 - April 2004	No
<b>KOSSI 2008</b> <b>19438891</b>	Retrospective	31	NC	NC	NC	NC	NC	NC	31	2(6.4)	Urinary retention	NR	January 2004 - May 2007	No
<b>BROUWER 2007 *</b> <b>17610695</b>	Prospective	163	18	0(0)	58	0(0)	76	0(0)	137	2(1.46)	urinary symptoms requiring an indwelling catheter in the postoperative period	NR	1995 - 2005	No
<b>SLACK 2007</b> <b>17877680</b>	Retrospective	128	128	5(1.4)	NC	NC	128	5(1.4)	NC	NC	urinary catheterization	<7 days	May 1999 - September 2006	No
<b>GHEZZI 2007</b> <b>18163992</b>	Prospective	33	NC	NC	NC	NC	NC	NC	33	3(9.1)	urinary retention that required intermittent self-catheterization	>1 month	October 2005 - June 2007	No
<b>MEREU 2007</b> <b>17630164</b>	Prospective	192	NC	NC	NC	NC	NC	NC	192	9(4.7)	Urinary retention after 30 days	>1 month	January 2003 - December 2005	Yes
<b>SERACHIOLLI 2007</b> <b>17501958</b>	Prospective	22	NC	NC	NC	NC	NC	NC	22	5(22.7)	Transient urinary retention requiring	2-12 weeks	January 2001 - January 2003	No

											self-catheterization			
<b>JATAN 2006 *</b> <b>16322961</b>	Retrospective	95	43	0(0)	38	0(0)	81	0(0)	14	1(7.1)	urinary retention	NR	April 1996 - August 2004	No
<b>MOHR2005 *</b> <b>15791964</b>	Retrospective	178	100	0(0)	39	0(0)	139	0(0)	48	1(2)	urinary retention	NR	March 1990 - March 1997	No
<b>DARAI2005</b> <b>15695977</b>	Prospective	40	NC	NC	NC	NC	NC	NC	40	7(17.5)	Urinary retention required bladder catherization	2 weeks (range: 2-4 weeks)	March 2001 - March 2003	No
<b>LANDI2005</b> <b>16449312</b>	Prospective	65	NC	NC	NC	NC	NC	NC	65	4(6.2)	bladder self- catheterization for >50 days	>50 days	December 2003 - September 2004	No
<b>TOUBOUL2005</b> <b>25294552</b>	RCT	40	NC	NC	NC	NC	NC	NC	40	5(12.5)	self-catheterization more than 6 months	>6 months	January 2006 - December 2008	No
<b>CAMPAGNACCI</b> <b>2005</b> <b>15759190</b>	Prospective	7	NC	NC	NC	NC	NC	NC	7	1(14)	temporary urinary retention	NR	March 1995 - March 2003	No
<b>MARPEAU2004</b> <b>15550878</b>	Prospective	32	NC	NC	NC	NC	NC	NC	32	6(18.8)	self-catheterization	3 weeks	From March 2001 - March 2003	No
<b>PASSOVER 2000</b> <b>10960302</b>	Prospective	34	NC	NC	NC	NC	NC	NC	34	0(0)	NR	NC	NR	No
<b>DONNEZ 1995</b> <b>7782444</b>	Prospective	231	231	3(1.3)	NC	NC	231	3(1.3)	NC	NC	urinary retention	NR	NR	No
<b>Total</b>		5962	1734	41(2.4)	573	43(7.5)	2307	84(3.6)	3655	368(10.1)				

NC= non-concerned

NR= non-reported

### endometriosis

\*Studies reporting at least two groups of colorectal surgery.

<b>Nerve sparing technique</b>	<b>Inclusion period</b>	<b>n conservative surgery</b>	<b>Voiding dysfunction if conservative surgery n(%)</b>	<b>n Radical surgery</b>	<b>Voiding dysfunction if radical surgery n(%)</b>	<b>Non-Nerve sparing or not clearly reported</b>	<b>Inclusion period</b>	<b>n conservative surgery</b>	<b>Voiding dysfunction if conservative surgery n(%)</b>	<b>n Radical surgery</b>	<b>Voiding dysfunction if radical surgery n(%)</b>
<b>ABO 2018 29307394</b>	June 2009 - December 2015	225	24(10.7)	139	20(4.3)	<b>BOURDEL 2017 29143306</b>	January 2000 - June 2013	172	6(3.5)	23	1(4.3)
<b>ERDEM 2018 29664797</b>	June 2002 – May 2011	NC	NC	51	2(3)	<b>RENNER 2017 28374101</b>	2001–2010	NC	NC	112	15(15.3)
<b>HUDELIST 2018 30080244</b>	March 2011 - August 2016	172	6(3.5)	23	1(4.3)	<b>MARTY 2017 28673871</b>	December 2012 - December 2016	110	2(1.8)	NC	NC
<b>JAYOT 2017 28987649</b>	January 2015 - September 2016	31	0(0)	31	7(22)	<b>DIGUISTO 2015 25813434</b>	December 2011 - June 2014	22	3 (13.6)	6	1(16.7)
<b>ROMAN 2017 28139235</b>	June 2009 - June 2016	111	10(9)	NC	NC	<b>MEULEMAN 2014 23579578</b>	September 2006 - October 2008	NC	NC	76	2(2.6)
<b>MILLOCHAU 2017 28893658</b>	March 2011 - December 2016	20	4(19)	NC	NC	<b>RUFFO 2014 24877097</b>	January 2002 - December 2010	NC	NC	900	114(14.7)
<b>AFORS 2016 27544881</b>	January 2010 - September 2012	72	3(4.2)	30	2(6.6)	<b>BOILEAU 2012 22071018</b>	2005 – 2011	NC	NC	23	2(8.9)
<b>ROMAN 2016 27393269</b>	January 2005 - January 2010	46	1(2.2)	25	3(12)	<b>WOLTHUIS 2011 21427115</b>	September 2009 - September 2010	NC	NC	21	1(4.8)
<b>ROMAN 2016 26474734</b>	June 2009 - November 2014	NC	NC	103	5(4.9)	<b>FANFANI 2010 19394600</b>	January 2003 - December 2007	48	0(0)	88	13(14.7)
<b>ROMAN 2016 27565263</b>	June 2009 - September 2014	122	8(6.6)	NC	NC	<b>BENBARA 2008 19019719</b>	1998 – 2005	NC	NC	50	10(30.4)
<b>AKLADIOS 2015 25109779</b>	2004 – 2011	NC	NC	41	4(9.7)	<b>DUBERNAN D 2008 18313000</b>	April 2001 - April 2004	NC	NC	52	12(23)
<b>ROMAN 2015 26347968</b>	June 2009 - November 2014	22	3 (13.6)	6	1(16.7)	<b>KOSSI 2008 19438891</b>	January 2004 - May 2007	NC	NC	31	2(6.4)
<b>FLEISCH 2014 24407554</b>	2008 – 2010	NC	NC	4	2(50)	<b>BROUWER 2007 17610695</b>	1995 - 2005	76	0(0)	137	2(1.46)
<b>BALLESTER 2014 24858943</b>	2001-2011	NC	NC	227	45(22)	<b>SLACK 2007 17877680</b>	May 1999 - September 2006	128	5(1.4)	NC	NC
<b>Che 2014 24576484</b>	September 2010 - July 2012	NC	NC	21	9(4.3)	<b>GHEZZI 2007</b>	October 2005 - June 2007	NC	NC	33	3(9.1)

							<b>18163992</b>					
<b>MANGLER 2014</b> <b>24726619</b>	September 2004 - December 2012	NC	NC	71	2(2.8)		<b>SERACHIO LLI 2007</b> <b>17501958</b>	January 2001 - January 2003	NC	NC	22	5(22.7)
<b>MEULEMAN 2011</b> <b>22729087</b>	September 2004 - September 2006	NC	NC	45	1(2.2)		<b>JATAN2006</b> <b>16322961</b>	April 1996 - August 2004	81	0(0)	14	1(7.1)
<b>BALLESTER 2011</b> <b>21256472</b>	January 2006 - December 2008	NC	NC	52	15(29)		<b>MOHR2005</b> <b>15791964</b>	March 1990 - March 1997	139	0(0)	48	1(2)
<b>DONNEZ 2010</b> <b>20547557</b>	2004-2008	500	4(0.8)	NC	NC		<b>DARAI2005</b> <b>15695977</b>	March 2001 -March 2003	NC	NC	40	7(17.5)
<b>DOUSSET 2010</b> <b>20395847</b>	1995-2003	NC	NC	100	16(16)		<b>LANDI2005</b> <b>16449312</b>	December 2003 - September 2004	NC	NC	65	4(6.2)
<b>RUFFO 2010</b> <b>19466489</b>	January 2002 - March 2008	NC	NC	436	71(19.9)		<b>TOUBOUL 2005</b> <b>25294552</b>	January 2006 - December 2008	NC	NC	40	5(12.5)
<b>STEPNIEWSKA 2010</b> <b>19836731</b>	2000 – 2005	40	0(0)	60	3(5)		<b>CAMPAGN ACCI 2005</b> <b>15759190</b>	March 1995 - March 2003	NC	NC	7	1(14)
<b>ROMAN 2010</b> <b>20106836</b>	January 2005 - June 2008	16	0(0)	25	2(8)		<b>MARPEAU 2004</b> <b>15550878</b>	From March 2001 - March 2003	NC	NC	32	6(18.8)
<b>FERRERO 2009</b> <b>18684451</b>	May 1999 - May 2006	NC	NC	46	2(4.3)		<b>PASSOVER 2000</b> <b>10960302</b>	NR	NC	NC	34	0(0)
<b>LANDI 2008</b> <b>18656048</b>	January 2003 - September 2006	35	0(0)	NC	NC		<b>DONNEZ 1995</b> <b>7782444</b>	NR	231	3(1.3)	NC	NC
<b>MEREU 2007</b> <b>17630164</b>	January 2003 - December 2005	NC	NC	192	9(4.7)							
<b>26 studies</b>	1999 - 2016	1402	62(4.4)	1718	222(12.9)		<b>25 studies</b>	1995-2013	1001	18(1.8)	1842	208(11.3)

**Table 2: Nerve sparing versus non-nerve sparing technique in colo-rectal surgery for deep endometriosis**