



**HAL**  
open science

## Temperature and fuel availability control fire size/severity in the boreal forest of central Northwest Territories, Canada

Dorian M. Gaboriau, Cécile C. Remy, Martin P. Girardin, Hugo Asselin, Christelle Hély, Yves Bergeron, Adam A. Ali

### ► To cite this version:

Dorian M. Gaboriau, Cécile C. Remy, Martin P. Girardin, Hugo Asselin, Christelle Hély, et al.. Temperature and fuel availability control fire size/severity in the boreal forest of central Northwest Territories, Canada. *Quaternary Science Reviews*, 2020, 250, pp.106697 -. 10.1016/j.quascirev.2020.106697 . hal-03493397

**HAL Id: hal-03493397**

**<https://hal.science/hal-03493397>**

Submitted on 21 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Temperature and fuel availability control fire size/severity in the boreal forest of**  
2 **central Northwest Territories, Canada**

3 Dorian M. Gaboriau<sup>1,2</sup>, Cécile C. Remy<sup>3</sup>, Martin P. Girardin<sup>4,5</sup>, Hugo Asselin<sup>1,4</sup>, Christelle Hély<sup>2,6,7</sup>,  
4 Yves Bergeron<sup>4,7</sup>, Adam A. Ali<sup>2,7</sup>

5 <sup>1</sup>School of Indigenous Studies, Université du Québec en Abitibi-Témiscamingue, 445 Boulevard de  
6 l'Université, Rouyn-Noranda, QC, J9X 5E4, Canada

7 <sup>2</sup>Institut des Sciences de l'Evolution, UMR 5554 CNRS-IRD-Université Montpellier-EPHE, 2 place  
8 Eugène Bataillon, 34095 Montpellier, France

9 <sup>3</sup>Department of Biology, University of New Mexico, Albuquerque, United States

10 <sup>4</sup>Centre for Forest Research, Université du Québec à Montréal, P.O. Box 8888, Stn. Centre-ville,  
11 Montréal, QC, H3C 3P8, Canada

12 <sup>5</sup>Natural Resources Canada, Canadian Forest Service, Laurentian Forestry Centre, 1055 rue du PEPS,  
13 P.O. Box 10380, Stn. Sainte-Foy, Québec, QC, G1V 4C7, Canada

14 <sup>6</sup>École Pratique des Hautes Études, PSL University, Paris, France

15 <sup>7</sup>Forest Research Institute, Université du Québec en Abitibi-Témiscamingue, 445 Boulevard de  
16 l'Université, Rouyn-Noranda, QC, J9X 5E4, Canada

17 **Corresponding author**

18 dorian.gaboriau@uqat.ca

19

20

21 **Abstract**

22 The north-central Canadian boreal forest experienced increased occurrence of large and severe wildfires  
23 caused by unusually warm temperatures and drought events during the last decade. It is, however,  
24 difficult to assess the exceptional nature of this recent wildfire activity, as few long-term records are  
25 available in the area. We analyzed macroscopic sedimentary charcoal from four lakes and pollen grains  
26 from one of those lakes to reconstruct long-term fire regimes and vegetation histories in the boreal forest  
27 of central Northwest Territories. We used regional estimates of past temperature and hydrological  
28 changes to identify the climatic drivers of fire activity over the past 10,000 years. Fires were larger and  
29 more severe during warm periods (before *ca.* 5,000 cal. yrs. BP and during the last 500 years) and when  
30 the forest landscape was characterized by high fuel abundance, especially fire-prone spruce. In contrast,  
31 colder conditions combined with landscape opening (i.e., lower fuel abundance) during the Neoglacial  
32 (after *ca.* 5,000 cal. yrs. BP) were related with a decline in fire size and severity. Fire size and severity  
33 increased during the last five centuries, but remained within the Holocene range of variability.  
34 According to climatic projections, fire size and severity will likely continue to increase in central  
35 Northwest Territories in response to warmer conditions, but precipitation variability, combined with  
36 increased abundance of deciduous species or opening of the landscape, could limit fire risk in the future.

37

38 **Keywords**

39 large wildfires; charcoal; pollen; fire size; lake sediments; vegetation dynamics; climate change;  
40 Holocene

41

42

43

## 44 **1. Introduction**

45 The global climate is experiencing rapid warming conducive to increased occurrence of extreme  
46 weather events like heatwaves and droughts (IPCC, 2014; Mann et al., 2017). Such conditions may  
47 increase the occurrence of exceptionally large and severe wildfires due to the strong positive correlation  
48 between temperature and area burned (Abatzoglou and Kolden, 2013; Ali et al., 2012; Duffy et al., 2005;  
49 Gaboriau et al., under review; Kasischke and Turetsky, 2006; Turco et al., 2017). Particularly high  
50 temperatures and recurrent droughts during the last decade have led to severe wildfire seasons in forest  
51 ecosystems worldwide, for example in Australia (Boer et al., 2020; Nolan et al., 2020), Sweden (SFOR,  
52 2018), Chile (De la Barrera et al., 2018), Greece and the USA (Smith et al., 2019), as well as in the  
53 Northwest Territories (NWT) in Canada where about 3.4 million ha burned during the summer of 2014  
54 (Kochtubajda et al., 2019; NTENR, 2015).

55 Large wildfires in the NWT mainly occur from June to September. Lightning-induced fires are  
56 predominant and fire frequency and area burned are among the highest in the North American boreal  
57 forest (Erni et al., 2020; Veraverbeke et al., 2017). However, the 2014 wildfire season was exceptional  
58 in comparison with observations of the last decade (3.4 million ha burned in 2014, compared to a mean  
59 of 622 thousand hectares burned annually from 2009 to 2019; Gaboriau et al., under review). More than  
60 380 fires were observed, some of which having burned more than 150,000 hectares. Moreover,  
61 firefighting costs reached more than CAD 56 million during this exceptional year, compared to an  
62 annual mean of CAD 7.5 million in the previous 20 years (NTENR, 2015).

63 Few remnant trees are left behind in areas burned during large and severe wildfires (Erni et al.,  
64 2017; Stephens et al., 2014). Recolonization is dominated by fast-growing pioneer deciduous tree  
65 species whose seeds disperse over long distances (Chapin et al., 2010; Walker et al., 2018). Large  
66 wildfires also reduce access to ecosystem services (Adams, 2013), cause health issues in human

67 populations (Dodd et al., 2018), and decrease the capacity of Indigenous people to maintain traditional  
68 activities on the land (Berkes and Davidson-Hunt, 2006).

69 Since the 1950s, mean annual temperature increased by over 1.5°C in Canada, leading to an  
70 earlier onset of snowmelt across much of the boreal forest (Zhang et al., 2011). According to climate  
71 projections, temperature will continue to rise in the next decades, particularly in the north-central  
72 Canadian boreal forest (Price et al., 2013; Wang et al., 2014). Summer and fall precipitation are  
73 projected to increase in the boreal forest of Canada and annual precipitation could increase up to 20% by  
74 2100. However, these conditions will not necessarily lead to increased soil and vegetation moisture, as  
75 projected warmer temperatures will lead to increased evapotranspiration (IPCC, 2014; Meehl et al.,  
76 2007). Hence, the drying effect of higher temperatures in the future will not necessarily be compensated  
77 by increased precipitation. Boreal forest landscapes will consequently be altered by a longer growing  
78 season (Jain et al., 2018), increased fire activity (Flannigan et al., 2013; Hassol, 2005; Wang et al., 2017;  
79 Wotton et al., 2017), higher proportion of deciduous tree species and more open forest canopies than  
80 currently (Baltzer et al., submitted; Boulanger et al., 2017; Chaste et al., 2019; Mekonnen et al., 2019).  
81 However, the robustness of the predictions of climate change impacts on wildfire regimes, particularly  
82 large wildfires, is limited by our lack of knowledge of the long-term interactions between climate,  
83 vegetation and fire (Ali et al., 2012; Conedera et al., 2009; Coogan et al., 2019). While the main drivers  
84 of fire activity have been documented in the NWT for the recent past (1965-2016; Gaboriau et al.,  
85 submitted), few palaeoecological data are available at multi-millennial timescales.

86 To get insights into the long-term relationships between climate, vegetation and fire, we used  
87 high-resolution sedimentary proxies (pollen grains and charcoal particles) to reconstruct vegetation and  
88 fire histories for the past 10,000 years in the boreal forest of central NWT. We compared the  
89 reconstructed vegetation and fire histories with previous studies based on regional climatic

90 reconstructions. Our objectives were (1) to determine whether the recent wildfire regime (last 500 years)  
91 is within or outside the Holocene range of variability and (2) to decipher the respective roles of climate  
92 and vegetation controls on past fire activity. We expected that the recent fire regime would be  
93 characterized by larger and/or more severe wildfires compared to the past 10,000 years in response to  
94 ongoing climate warming and recurrent development of severe drought episodes.

## 95 **2. Material and methods**

### 96 *2.1. Study area and present-day vegetation*

97 We sampled sediments from lakes *Emile*, *Izaak*, *Paradis* and *Saxon* (unofficial names) located north  
98 of Yellowknife, in central NWT, and 5-100 km distant from each other (Fig. 1). The sampled lakes have  
99 a mean elevation of 382 m (Table 1). According to the Canadian National Fire Database (CNFDB,  
100 <http://cwfis.cfs.nrcan.gc.ca/ha/nfdb>), time since the last fire varied between 4 and 39 years in the  
101 watersheds of the studied lakes (Table 1). The regional climate is dry continental, with long cold winters  
102 and short warm summers (Environnement Canada, 2017). The study area is located close to the treeline  
103 (Timoney et al., 2019) and is part of the Taiga Boreal Shield West ecoregion, characterized by rock  
104 outcrops and discontinuous permafrost (Olson et al., 2001). Vegetation is currently dominated by conifer  
105 tree species, mostly black spruce (*Picea mariana* (Mill.) B.S.P.). White spruce (*Picea glauca* (Moench)  
106 Voss.) is mainly found in the southwestern part of the study area, and jack pine (*Pinus banksiana*  
107 Lamb.) mostly occupies the southern part of the NWT, *i.e.* in the Taiga Plains. Broadleaf tree species are  
108 also present in younger stands having regenerated after fire, mostly paper birch (*Betula papyrifera*  
109 Marsh) and trembling aspen (*Populus tremuloides* Michx.), such as observed at lake *Emile* (Appendix  
110 S1A). The shrub layer is mainly composed of *Betula glandulosa* Michx., *Alnus alnobetula* (Ehrh.) K.  
111 Köch, *Andromeda polifolia* L., *Ledum palustre* L., and *Chamaedaphne calyculata* (L.) Moench. The  
112 ground layer is mostly composed of *Vaccinium vitis-idaea* L., *Rubus chamaemorus* L., *Kalmia*

113 *angustifolia* L. and *Lycopodium annotinum* L., with abundant mosses (mostly *Sphagnum* spp.), while  
114 *Cladonia* lichens are found on xeric rock outcrops.

## 115 2.2. Past regional climate and vegetation

116 Previous studies have documented long-term regional climatic and vegetation changes in the  
117 northwestern and north-central Canadian boreal forest, which in turn had impacts on wildfire regimes.  
118 Before *ca.* 11,000 cal. yrs. BP (calibrated years before present), high solar radiation and summer  
119 insolation increased temperatures in the northwestern Canada (Viau et al., 2006), reduced snow cover,  
120 and led to the retreat of the Laurentide ice sheet (Dyke, 2005). Deglaciation occurred between *ca.*  
121 10,000 – 9,000 cal. yrs. BP in the study area (Dalton et al., 2020) and left behind numerous lakes and  
122 swamps scattered across the landscape (Latifovic et al., 2017; MacDonald, 1995). Following ice retreat,  
123 the land was quickly colonized by tundra vegetation soon followed by trees once temperatures warmed  
124 during the early Holocene (Dyke, 2005). The Holocene Thermal Maximum (HTM) occurred in northern  
125 continental Canada during the mid-Holocene (between 7,300 and 4,300 cal. yrs. BP; Kaufman et al.,  
126 2004). During this period, major vegetation reorganization occurred in response to warmer temperature,  
127 such as treeline migration to the north and increased abundance of fire-adapted species (i.e. jack pine  
128 and black spruce), as observed in the NWT and southwestern Yukon (Gajewski et al., 2014; Pienitz et  
129 al., 1999; Sulphur et al., 2016). The decrease in temperature after 4,300 cal. yrs. BP, characterizing the  
130 Neoglacial period, favored coniferous species such as spruce as well as deciduous shade-tolerant tree  
131 species such as paper birch in permafrost-free areas (Moser and MacDonald, 1990; Sulphur et al., 2016).  
132 However, interactions between climate, vegetation and fire depend on regional atmospheric and oceanic  
133 oscillations, as observed in the eastern Canadian boreal forest where conifer species also increased in  
134 density after 3,500 cal. yrs. BP (Fr chet te et al., 2018; Remy et al., 2017).

135       2.3. *Sampling*

136       We sampled lake sediments in June 2018 from the deepest point in each of the four lakes. The  
137       vegetation of the watersheds was mainly composed of recently burned conifer tree species and/or shrub  
138       birch having grown after the last fire (Appendix S1). The four lakes were small, deep, circular, and  
139       characterized by the absence of connections to surrounding watercourses (Table 1). We used a Kajak-  
140       Brinkhurst (KB) gravity corer to collect the most recently deposited material at the water-sediment  
141       interface (Glew 1991). We subsampled surface sediments on site in 0.5-cm thick sections stocked in  
142       plastic bags. We collected deeper sediments in 1-m cores using a 5-cm diameter Russian corer; we  
143       wrapped these cores in aluminum foil and placed them in hemicylindrical tubes for protection. We sliced  
144       the sediment cores in the laboratory into continuous 1-cm thick subsamples that we stocked in plastic  
145       bags and kept refrigerated at 4°C until analysis. Surface sediments were sampled at a shorter interval  
146       (0.5cm) than deeper sediments (1 cm) to ease comparison with recent (last 500 years) reconstructions of  
147       vegetation and fire history.

148       2.4. *Radiocarbon dating and age-depth models*

149       The basal 10 cm of the deepest sediment cores were mostly clay, corresponding to the beginning  
150       of lake sedimentation following deglaciation in the four sites, whereas the upper sediment was  
151       composed of a gyttja richer in organic matter. We constructed core chronologies based on radiocarbon  
152       dating of bulk gyttja samples by <sup>14</sup>C accelerator mass spectrometry (AMS). We used the ‘*WinBacon*’  
153       v.2.3.7 R package (Blaauw and Christen, 2011) to reconstruct Bayesian sediment accumulation histories  
154       and calibrate age-depth models constructed from seven dates for lakes *Emile*, *Paradis* and *Saxon* and  
155       nine dates for lake *Izaak* (Table S1). We used the IntCal13.14C calibration curve for terrestrial northern  
156       hemisphere material (Reimer et al., 2013). We interpolated ages at contiguous 0.5-cm depth intervals  
157       and all dates were expressed in calibrated years before present (cal. yrs. BP). We used linear



158 interpolation between  $^{14}\text{C}$  ages rather than best-fit curves, assuming that both methods yield equivalent  
159 chronologies (Blaauw, 2010; Trachsel and Telford, 2017).

### 160 *2.5. Subsample chemical preparation and charcoal analysis*

161 To distinguish charcoal from other biological material, we applied a common charcoal extraction  
162 protocol to each subsample using the ‘chemical digestion’ method (Winkler, 1985), based on charcoal  
163 resistance to chemicals (Mooney and Tinner, 2011). We removed a 1.4-cm<sup>3</sup> subsample from each liquid  
164 subsample of surface sediment (KB) and 1-cm<sup>3</sup> for the rest of the core (solid part), following Mustaphi  
165 et al. (2015). We took sediments in the central part of each core slice to limit the risk of contamination  
166 with modern material. We deflocculated and bleached the subsamples by placing them in a potassium  
167 hydroxide (KOH) solution combined with bleach and sodium hexametaphosphate – (NaPO<sub>3</sub>)<sub>6</sub> – on a  
168 stirring table for 24 hours at room temperature; which later allowed us to differentiate black charcoal  
169 from bleached organic matter (Braadbaart and Poole, 2008; Swain, 1973). We wet-sieved the solution  
170 through a 160 μm mesh to collect larger charcoal particles produced by fire events having mostly  
171 occurred 0-3 km from the lakeshore (Oris et al., 2014). We sorted charcoal particles in a Petri dish, and  
172 measured their area using a camera (D-Moticam 1080, Motic Images, 2018) mounted on a binocular  
173 microscope and connected to an image-analysis software (MOTICAM IMAGE Plus 3.0).

### 174 *2.6. Fire history reconstruction*

175 We reconstructed burned biomass (hereafter *BB*; unitless) at each study site by measuring the  
176 cumulative area of charcoal particles per cm<sup>3</sup> for each subsample. Based on numerical age-depth  
177 models, we transformed this measure into charcoal accumulation rate (hereafter CHAR, i.e. mm<sup>2</sup> cm<sup>-2</sup>  
178 yr<sup>-1</sup>), using the R package ‘*paleofire*’ v.1.2.3 (Blarquez et al., 2014). We pooled and smoothed these  
179 series (using a 500-year window) by (1) rescaling initial CHAR values using min-max transformation,  
180 (2) homogenizing the variance using Box-Cox transformation, and (3) rescaling the values to Z-scores

181 (Power et al., 2008). The resulting values (cumulative charcoal area) are interpreted as representing the  
182 regional signature of fires having occurred in the watersheds of the studied lakes, and thus the regional  
183 burned biomass (hereafter *RegBB*; unitless).

184 We detected past fire events (Appendix S2) within each individual CHAR series using the  
185 CharAnalysis v.1.1 software (Higuera, 2009; available at <https://github.com/phiguera/CharAnalysis>).  
186 Following Brossier et al. (2014), we used the narrowest time window allowing us to obtain a median  
187 Signal-to-Noise Index greater than 3.0 (Table 1). We obtained the past regional fire frequency (hereafter  
188 *RegFF*; fire.year<sup>-1</sup>) by combining the smoothed series using the R package ‘*paleofire*’ v.1.2.3 (Blarquez  
189 et al., 2014). We assessed the significance of changes in *RegFF* and *RegBB* by using a bootstrap  
190 procedure with 999 iterations (BCI; 90%).

191 Following Ali et al. (2012), we added a constant equal to 1 to *RegBB* and *RegFF* and we  
192 computed the ratio (*FS* index), which we interpreted as indicating fire size and/or severity. The *FS* index  
193 corresponds to mean burned biomass per fire calculated for 500-year intervals, and thus, reflects fire  
194 size. Considering that large fires in boreal North America are known to cause high vegetation mortality,  
195 the *FS* index could also represent fire severity (Ali et al., 2012; Cansler and McKenzie, 2014; Hély et al.,  
196 2020; Hennebelle et al., 2020; Kelly et al., 2013). We assumed that high *FS* index values corresponded  
197 to periods when the study area experienced larger and more severe wildfires. Following Power et al.  
198 (2008), we used a reference period, here corresponding to 500-0 cal. yrs. BP (by convention, 0 cal. yr.  
199 BP corresponding to AD 1950), to calculate *RegBB*, *RegFF* and *FS* index anomalies.

## 200 2.7. Vegetation history reconstruction

201 We reconstructed the Holocene vegetation history using palynological analysis of subsamples of  
202 bulk sediment (1 cm<sup>3</sup>) taken at regular 4 cm intervals along the core of lake *Emile*. We counted and  
203 determined pollen and spores in sediment subsamples using standard techniques described by Faegri and

204 Iversen, (1989). We counted a minimum of 300 pollen grains of terrestrial taxa for each subsample  
205 (Djamali and Cilleros, 2020), under a microscope with a  $\times 200$  or  $\times 400$  magnification factor. We  
206 identified pollen grains based on the Pollen Atlas of Arctic and Boreal Canada (Williams, 2006) and  
207 pollen keys from (Vincent, 1973) and Richard (1970). We identified tree pollen to the genus level. We  
208 also identified the green algae *Pediastrum* and spores of aquatic plants. We added exotic marker pollen  
209 (*Eucalyptus*) to each subsample to estimate pollen concentration (grains  $\text{cm}^3$ ) and pollen accumulation  
210 rate ( $n$  grains  $\text{cm}^{-2} \text{yr}^{-1}$ ) for taxa with an average percentage greater than 0.1% (Appendix S3), using the  
211 R package 'rioja' (Juggins, 2017). We measured the total pollen accumulation rate (PAR), based on all  
212 terrestrial taxa with percentages greater than 0.1%.

### 213 2.8. Temperature and drought

214 We used quantitative temperature reconstructions from four different proxies (described in Appendix  
215 S4) and originating from northwestern and north-central Canada (Lecavalier et al., 2017; Porter et al.,  
216 2019; Upiter et al., 2014; Viau et al., 2006). The reconstructions included (i) mean air temperature in  
217 July from 6,000 cal. yrs. BP to present estimated from chironomid assemblages sampled in lake  
218 sediments of the central Northwest Territories (Upiter et al., 2014), (ii) mean summer temperature  
219 anomalies relative to the 1961-1990 period in central Yukon, from 10,000 cal. yrs. BP to present,  
220 estimated using precipitation isotopes in syngenetic permafrost (Porter et al., 2019), (iii) mean Arctic  
221 summer air temperature anomalies relative to the preindustrial period reconstructed from ice melt in the  
222 Agassiz ice cap (Ellesmere Island, Northwest Territories) (Lecavalier et al., 2017), and (iv) mean July  
223 temperature estimated from pollen records for the past 10,000 years across northwestern North America  
224 (Viau et al., 2006). We assumed that these reconstructions captured temperature fluctuations during the  
225 summer months, i.e. the period of warm and dry weather conducive to wildfire activity in the boreal  
226 forest. For each reconstruction, we interpolated and standardized values for each year between 10,000

227 cal. yrs. BP and present, converting the data in anomalies relative to the mean for the entire period. We  
228 then averaged the four reconstructions. Then, we assessed the significance of temporal changes in  
229 temperature by bootstrapping the pooled means 999 times (BCI; 90%). We present temperature  
230 anomalies relative to the reference period 500-0 cal. yrs. BP.

231 To quantify the relationships among the fire, vegetation and temperature time series reconstructed at  
232 different temporal resolutions, and to avoid interpolation when time series were not sampled at identical  
233 points, we measured the Pearson correlation coefficient with 95% confidence interval between the  
234 automatically binned undetrended time series following Mudelsee (2013) and using the R package  
235 'BINCOR' (Polanco-Martinez et al., 2019).

236 We used three indicators of hydrological variability to define Holocene dryness periods assumed to  
237 correspond to low vegetation productivity (inferred from pollen and plant macrofossil assemblages) and  
238 low lake water levels (inferred from diatoms) (Lauriol et al., 2009; Pienitz et al., 1999; Viau and  
239 Gajewski, 2009). We compared these records providing independent evidence of dryness periods with  
240 our fire reconstructions to detect possible temporal overlap of specific fire regimes and dryness periods.

### 241 **3. Results**

#### 242 *3.1. Age-depth models*

243 Core length varied from 359 cm at lake *Izaak* to 456 cm at lake *Emile* (Table S1). According to the  
244 Bayesian age-depth models, basal sediments were dated between *ca.* 10,000 cal. yrs. BP at lake *Saxon*  
245 and *ca.* 9,530 cal. yrs. BP at lake *Izaak* (Fig. 2). We used 9,530 cal. yrs. BP as the earliest date for  
246 wildfire regime reconstructions, as it was included in all four reconstructions. The four age-depth  
247 models had similar sedimentation rates, varying between 0.036 cm and 0.049 cm year<sup>-1</sup> (Table 1),  
248 similar to the results a previous study of other boreal lakes in the NWT (Crann et al., 2015).

249       3.2. Regional fire history

250       The regional fire activity remained fairly constant throughout the Holocene with approximately 5  
251 fires per millennium (+2 anomaly; Fig. 3A). Mean *RegFF* very slightly increased from *ca.* 9,500 cal.  
252 yrs. BP to *ca.* 500 cal. yrs. BP, before decreasing during the reference period (500-0 cal. BP). *RegBB*  
253 steadily increased starting from *ca.* 9,500 cal. yrs. BP, peaked between *ca.* 7,000-5,000 cal. yrs. BP,  
254 before gradually decreasing to present values, the lowest of the entire series (Fig. 3B). Higher *FS* index  
255 values were recorded before 4,000 cal. yrs. BP (Fig. 3C), and, although to a lesser extent, during the  
256 reference period (500-0 cal. yrs. BP).

257       3.3. Vegetation history

258       We identified a total of 41 pollen taxa in the whole sequence extracted at lake *Emile*, covering the  
259 period from *ca.* 9,700 cal. yrs. BP to present. Eighteen taxa had a mean pollen percentage > 0.1% over  
260 the entire study period (Appendix S3). The PAR diagram is dominated by few tree and shrub taxa  
261 throughout the sequence (*Picea*, *Betula*, *Pinus* and *Alnus*; Fig. 4). *Artemisia*, Cyperaceae, Ericaceae,  
262 Poaceae, *Juniperus*, *Larix* and *Salix* were also present, but in low numbers.

263       PAR was low from *ca.* 9,700 to 7,800 cal. yrs. BP, concomitant with a stable and relatively low  
264 sedimentation rate (Fig. 4). Most trees and shrubs were relatively low, except *Populus*, *Myrica* and  
265 *Lycopodium*, which recorded their highest values at that time. A marked increase of *Picea* (likely black  
266 spruce, MacDonald et al., 1993) occurred *ca.* 7,800 cal. yrs. BP, along with a simultaneous expansion of  
267 *Alnus alnobetula* subsp. *crispa*. At the same time, the lake recorded a major decrease in *Populus*, *Myrica*  
268 and *Lycopodium*. PAR increased markedly, while the sedimentation rate remained stable. PAR, *Picea*  
269 and *Alnus alnobetula* subsp. *crispa* remained high until *ca.* 6,000 cal. yrs. BP, before decreasing,  
270 concomitant with an increase of the sedimentation rate *ca.* 5,700 cal. yrs. BP. *Pinus* (likely jack pine;  
271 Sulphur et al., 2016), *Betula* and *Alnus incana* subsp. *rugosa* increased somewhat later, while the

272 sedimentation rate remained high. After *ca.* 4,000 cal. yrs. BP, the sedimentation rate decreased and  
273 remained stable for the last two millennia of the record, while *Pinus* remained relatively high, especially  
274 after *ca.* 2,500 cal. yrs. BP. A short-duration PAR peak occurred between *ca.* 3,000 and 2,200 cal. yrs.  
275 BP, although the sedimentation rate remained stable. *Pediastrum* and *Nuphar* increased simultaneously  
276 *ca.* 2,200 cal. yrs. BP. During the last two millennia, *Picea*, *Pinus*, *Betula* and *Alnus* sp. decreased, while  
277 Ericaceae and Cyperaceae remained stable or even slightly increased, such as for *Juniperus*. The  
278 sedimentation rate decreased after 500 cal. yrs. BP, following a slight increase between *ca.* 1,500 and  
279 500 cal. yrs. BP.

#### 280 *3.4. Temperature and drought*

281 We identified three main climatic periods over the Holocene, based on pooled summer temperature  
282 reconstructions (Fig. 5). Larger confidence intervals reveal more uncertainty in temperature estimations  
283 from 10,000 to 6,500 cal. yrs. BP, corresponding to the early Holocene period. Overall, maximum  
284 temperature values were recorded between *ca.* 6,500 and 4,500 cal. yrs. BP, before decreasing until *ca.*  
285 1,500 cal. yrs. BP, and increasing again drastically during the last few centuries. According to inferred  
286 hydrological conditions (Fig. 6; Lauriol et al., 2009; Pienitz et al., 1999; Viau and Gajewski, 2009), the  
287 period between *ca.* 7,000-3,000 cal. yrs. BP was characterized by wetter conditions in central NWT and  
288 northern Yukon, compared to the rest of the study period. A major change in diatom-inferred dissolved  
289 organic carbon at *ca.* 5,800 cal. yrs. BP suggests a warm and humid interval during the mid-Holocene  
290 (Fig. 6a; Pienitz et al., 1999), resulting in high lake productivity (Fig. 6c; Lauriol et al., 2009).  
291 Conversely, high annual precipitation were recorded before *ca.* 7,500 cal. yrs. BP and around *ca.* 1,000  
292 cal. yrs. BP (Fig. 6b; Viau and Gajewski, 2009).

293        *3.5. Relationships between fire, temperature and vegetation*

294        Warm periods (*ca.* 10,000 to 5,000 cal. yrs. BP and last 500 years) coincided with low *RegFF*, high  
295 *FS* index, and relatively high *RegBB* (Table 2). Higher temperatures corresponded with higher *Populus*,  
296 *Myrica* and *Lycopodium* abundance. A cooler period (*ca.* 5,000 to 500 cal. yrs. BP) was characterized by  
297 a low *FS* index and a landscape richer in *Pinus*, *Larix*, *Juniperus*, Poaceae, *Pediastrum* and *Nuphar*  
298 (Table 2). PAR, interpreted as fuel availability, coincided with high *RegFF* and high *RegBB*, especially  
299 for *Picea*, *Betula* and *Alnus alnobetula* subsp. *crispa* (Table 2).

300        **4. Discussion**

301        Our fire history reconstructions based on macroscopic sedimentary charcoal provide evidence that  
302 fire size and/or severity were higher during warmer periods (i.e. before *ca.* 5,000 cal. yrs. BP and during  
303 the last 500 years) than during the Neoglacial (after *ca.* 5,000 cal. yrs. BP). During the Holocene, the  
304 *RegBB* metric was positively correlated with the *FS* index because *RegFF* was relatively constant from  
305 the early to the late Holocene.

306        *4.1. Fire, climate and vegetation interactions*

307        *Early Holocene (10,000-6,500 cal. yrs. BP)*

308        The spatial variability of the climate data used in air temperature reconstructions shown by the large  
309 confidence interval around summer temperature during the early Holocene (i.e. before 6,500 cal. yrs.  
310 BP; Figure 5) is due to differences in time of deglaciation across northern Canada (Dyke, 2005;  
311 Kaufman et al., 2004). Summers were warmer in far northern NWT and in central Yukon (Lecavalier et  
312 al., 2017; Porter et al., 2019) but cooler in northwestern Canada (Viau et al., 2006), probably due to a  
313 later time of deglaciation. Conditions were also dry at this time in north-central Canada but not in  
314 northwestern Canada (i.e. Mackenzie region; Fig. 6), confirming the presence of the Laurentide ice sheet  
315 in the study area during the early Holocene. PAR indicates that the first stage of vegetation colonization

316 following ice retreat was characterized by an open woodland, as observed in previous studies for the  
317 same period and in the same area (Conedera et al., 2009; Macumber et al., 2011; Sulphur et al., 2016).  
318 The landscape was dominated by pioneer taxa such as *Populus* and *Betula* (likely dwarf birch, Andrews  
319 et al., 1980). The relatively high percentages of *Picea* spp. (~ 40-50 %) was likely due to long-distance  
320 transport of pollen by wind, from populations located to the southwest of the study area (Campbell et al.,  
321 1999). Low tree abundance before 8,000 cal. yrs. BP can be explained by the time required for  
322 northward migration following deglaciation (Gajewski et al., 1993; Moser and MacDonald, 1990;  
323 Ritchie, 1985). Dry conditions also could have limited tree cover. During the early Holocene, deciduous  
324 shrubs, less inflammable than conifer trees, and low fuel abundance likely limited fire ignition and  
325 spread, but might have promoted large and/or severe fires in shrub areas during dry years.

#### 326 *Mid-Holocene (6,500-5,000 cal. yrs. BP)*

327 Warmer and wetter climate during the mid-Holocene in northern continental Canada (Kaufman et al.,  
328 2004; Porter et al., 2019; Ritchie et al., 1983) favored tree growth, leading to gradual densification of the  
329 vegetation cover as suggested by the increase in PAR values despite a relatively stable sedimentation  
330 rate. This result is in line with previous reconstructions in southwestern Yukon (Cwynar and Spear,  
331 1995; Gajewski et al., 2014), Alaska (Tinner et al., 2006) and Northwest Territories (Sulphur et al.,  
332 2016). Warm temperature and wet conditions of the mid-Holocene especially favored dense conifer  
333 forests (*Picea* and *Pinus* spp.), causing fuel accumulation conducive to large and severe fires, as  
334 previously observed in Alaska (Hoecker et al., 2020). During this period, wet conditions limited the  
335 increase in fire frequency (*RegFF*), which remained stable, while biomass burning (*RegBB*) was higher.  
336 Because the period was wet, it suggests that these non-frequent large wildfires occurred during episodic  
337 drought that dried the fuel and favored fire spread. High fire activity during the mid-Holocene promoted



338 fire-prone coniferous species (*Picea* and *Pinus*) and pioneer trees and shrubs (*Betula* and *Alnus*), as  
339 previously observed in the study area (Parisien et al., 2020).

340

341 *Neoglacial (5,000-1,000 cal. yrs. BP)*

342 The Neoglacial was characterized by a gradual decrease in temperature, reaching an all-time low *ca.*  
343 1,500 cal. yrs. BP. Cooling favored the expansion of *Pediastrum* and *Nuphar ca.* 3,000 cal. yrs. BP, as  
344 observed in Alaska (Edwards et al., 2000). The Neoglacial was also characterized by landscape opening,  
345 as shown by a decrease in *Picea* and increase in Poaceae, also observed in previous studies (MacDonald,  
346 1995; Pienitz et al., 1999). Lower fuel abundance, likely due to cooler and drier conditions less  
347 favorable to shrub and tree productivity, hindered fire spread leading to lower biomass burned.

348

349 *Last millennium (1,000-0 cal. yrs. BP)*

350 During the last 1,000 years, temperature varied but remained low until *ca.* 500 cal. yrs. BP before a  
351 marked increase, especially during the last century. Vegetation density and fire frequency were as low as  
352 during the early Holocene, while *RegBB* reached its lowest levels for the entire time series. Contrary to  
353 our expectations, the recent fire size/severity (i.e. over the last 500 years) is below the maximum values  
354 observed during the mid-Holocene.

355

356 *4.2 Implications for future fire risk*

357 We provide evidence that both climate conditions and vegetation dynamics played a key role in  
358 shaping the wildfire regime over the past 10,000 years in central NWT. Fire size and/or severity were  
359 higher under the warmer and wetter climate of the mid-Holocene (7,000 to 5,000 cal. yrs. BP), which  
360 favored fuel availability, corroborating recent observations on large wildfires during the last decades in

361 the study area (Gaboriau et al., under review). Our results can be used to anticipate future fire risk and to  
362 elaborate risk mitigation strategies including fuel management.

363 While temperature is expected to continue increasing in western Canada over the 21<sup>st</sup> century,  
364 dryness periods could be more severe than in the past (Price et al., 2013). Hence, fire frequency might  
365 increase (Wotton et al., 2017), but not necessarily biomass burning. Indeed, temperature increase could  
366 lead to the conversion of coniferous to deciduous forests (Hansen et al., in press; Mekonnen et al.,  
367 2019), or to a more open landscape (Asselin and Payette, 2005; Baltzer et al., submitted), which would  
368 negatively feedback on fire ignition and spread by reducing fuel flammability, combustibility, and/or  
369 connectivity. Hence, for large wildfires to be frequent in the future, the warming-induced increase in  
370 evapotranspiration will have to be compensated by increased precipitation to produce sufficient fuel  
371 (Flannigan et al., 2016).

## 372 **5. Acknowledgements**

373 We thank Benoît Brossier, David Gervais, Julia Morarin, Laure Paradis and David Pretorius for their  
374 assistance in the field. We are also grateful to Christine Simard for her help in identifying charcoal  
375 fragments, Jordan Paillard for pollen counting and identification and Pierre J. H. Richard for suggestions  
376 on an earlier draft. This work was supported by Polar Knowledge Canada (Grant # NST-1718-0014), the  
377 Natural Sciences and Engineering Research Council of Canada, the Canadian Forest Service, the  
378 National Geography Society (Grant # EC-386R-18), and the French University Institute (IUF).

379 Data and R codes used in this manuscript are available [here](#).

380

## 381 **6. References**

- 382 Abatzoglou, J.T., Kolden, C.A., 2013. Relationships between climate and macroscale area burned in the  
383 western United States. *International Journal of Wildland Fire* 22, 1003–1020.
- 384 Adams, M.A., 2013. Mega-fires, tipping points and ecosystem services: managing forests and  
385 woodlands in an uncertain future. *Forest Ecology and Management* 294, 250–261.
- 386 Ali, A.A., Blarquez, O., Girardin, M.P., Hély, C., Tinquaut, F., El Guellab, A., Valsecchi, V., Terrier,  
387 A., Bremond, L., Genries, A., 2012. Control of the multimillennial wildfire size in boreal North  
388 America by spring climatic conditions. *Proceedings of the National Academy of Sciences* 109,  
389 20966–20970.
- 390 Andrews, J.T., Mode, W.N., Webber, P.J., Miller, G.H., Jacobs, J.D., 1980. Report on the distribution of  
391 dwarf birches and present pollen rain, Baffin Island, NWT, Canada. *Arctic* 33, 50–58.
- 392 Asselin, H., Payette, S., 2005. Late Holocene opening of the forest tundra landscape in northern Québec,  
393 Canada. *Global Ecology and Biogeography* 14, 307–313.
- 394 Baltzer, J.L., Nicola, D., Walker, X.J., Greene, D.F., Mack, M.C., submitted. Fire and the decline of fire-  
395 adapted black spruce in the boreal forest. *Science*.
- 396 Berkes, F., Davidson-Hunt, I.J., 2006. Biodiversity, traditional management systems, and cultural  
397 landscapes: examples from the boreal forest of Canada. *International Social Science Journal* 58,  
398 35–47.
- 399 Blaauw, M., 2010. Methods and code for ‘classical’ age-modelling of radiocarbon sequences. *Quaternary*  
400 *Geochronology* 5, 512–518.
- 401 Blaauw, M., Christen, J.A., 2011. Flexible paleoclimate age-depth models using an autoregressive  
402 gamma process. *Bayesian analysis* 6, 457–474.
- 403 Blarquez, O., Vannière, B., Marlon, J.R., Daniau, A.-L., Power, M.J., Brewer, S., Bartlein, P.J., 2014.  
404 paleofire: an R package to analyse sedimentary charcoal records from the Global Charcoal  
405 Database to reconstruct past biomass burning. *Computers & Geosciences* 72, 255–261.
- 406 Boer, M.M., de Dios, V.R., Bradstock, R.A., 2020. Unprecedented burn area of Australian mega forest  
407 fires. *Nature Climate Change* 10, 171–172.
- 408 Boulanger, Y., Taylor, A.R., Price, D.T., Cyr, D., McGarrigle, E., Rammer, W., Sainte-Marie, G.,  
409 Beaudoin, A., Guindon, L., Mansuy, N., 2017. Climate change impacts on forest landscapes  
410 along the Canadian southern boreal forest transition zone. *Landscape Ecology* 32, 1415–1431.
- 411 Braadbaart, F., Poole, I., 2008. Morphological, chemical and physical changes during charcoalification  
412 of wood and its relevance to archaeological contexts. *Journal of Archaeological Science* 35,  
413 2434–2445.
- 414 Brossier, B., Oris, F., Finsinger, W., Asselin, H., Bergeron, Y., Ali, A.A., 2014. Using tree-ring records  
415 to calibrate peak detection in fire reconstructions based on sedimentary charcoal records.  
416 *Holocene* 24, 635–645.
- 417 Campbell, I.D., McDonald, K., Flannigan, M.D., Kringayark, J., 1999. Long-distance transport of pollen  
418 into the Arctic. *Nature* 399, 29–30.
- 419 Cansler, C.A., McKenzie, D., 2014. Climate, fire size, and biophysical setting control fire severity and  
420 spatial pattern in the northern Cascade Range, USA. *Ecological Applications* 24, 1037–1056.
- 421 Carcaillet, C., Bouvier, M., Fréchette, B., Larouche, A.C., Richard, P.J., 2001. Comparison of pollen-  
422 slide and sieving methods in lacustrine charcoal analyses for local and regional fire history.  
423 *Holocene* 11, 467–476.
- 424 Chapin, F.S., McGuire, A.D., Ruess, R.W., Hollingsworth, T.N., Mack, M.C., Johnstone, J.F.,  
425 Kasischke, E.S., Euskirchen, E.S., Jones, J.B., Jorgenson, M.T., 2010. Resilience of Alaska’s  
426 boreal forest to climatic change. *Canadian Journal of Forest Research* 40, 1360–1370.

- 427 Chaste, E., Girardin, M.P., Kaplan, J.O., Bergeron, Y., Hély, C., 2019. Increases in heat-induced tree  
428 mortality could drive reductions of biomass resources in Canada's managed boreal forest.  
429 *Landscape Ecology* 34, 403–426.
- 430 Conedera, M., Tinner, W., Neff, C., Meurer, M., Dickens, A.F., Krebs, P., 2009. Reconstructing past fire  
431 regimes: methods, applications, and relevance to fire management and conservation. *Quaternary*  
432 *Science Reviews* 28, 555–576.
- 433 Coogan, S.C., Robinne, F.-N., Jain, P., Flannigan, M.D., 2019. Scientists' warning on wildfire—a  
434 Canadian perspective. *Canadian Journal of Forest Research* 49, 1015–1023.
- 435 Crann, C.A., Patterson, R.T., Macumber, A.L., Galloway, J.M., Roe, H.M., Blaauw, M., Swindles, G.T.,  
436 Falck, H., 2015. Sediment accumulation rates in subarctic lakes: insights into age-depth  
437 modeling from 22 dated lake records from the Northwest Territories, Canada. *Quaternary*  
438 *Geochronology* 27, 131–144.
- 439 Cwynar, L., Spear, R., 1995. Paleovegetation and paleoclimatic changes in the Yukon at 6 ka BP.  
440 *Géographie physique et Quaternaire* 49, 29–35.
- 441 Dalton, A.S., Margold, M., Stokes, C.R., Tarasov, L., Dyke, A.S., Adams, R.S., Allard, S., Arends, H.E.,  
442 Atkinson, N., Attig, J.W., 2020. An updated radiocarbon-based ice margin chronology for the  
443 last deglaciation of the North American Ice Sheet Complex. *Quaternary Science Reviews* 234,  
444 106223.
- 445 De la Barrera, F., Barraza, F., Favier, P., Ruiz, V., Quense, J., 2018. Megafires in Chile 2017:  
446 Monitoring multiscale environmental impacts of burned ecosystems. *Science of the Total*  
447 *Environment* 637–638, 1526–1536. <https://doi.org/10.1016/j.scitotenv.2018.05.119>
- 448 Djamali, M., Cilleros, K., 2020. Statistically significant minimum pollen count in Quaternary pollen  
449 analysis; the case of pollen-rich lake sediments. *Review of Palaeobotany and Palynology* 275,  
450 104156.
- 451 Dodd, W., Howard, C., Rose, C., Scott, C., Scott, P., Cunsolo, A., Orbinski, J., 2018. The summer of  
452 smoke: ecosocial and health impacts of a record wildfire season in the Northwest Territories,  
453 Canada. *The Lancet Global Health* 6, S30.
- 454 Duffy, P.A., Walsh, J.E., Graham, J.M., Mann, D.H., Rupp, T.S., 2005. Impacts of large-scale  
455 atmospheric–ocean variability on Alaskan fire season severity. *Ecological Applications* 15,  
456 1317–1330.
- 457 Dyke, A., 2005. Late Quaternary vegetation history of northern North America based on pollen,  
458 macrofossil, and faunal remains. *Géographie physique et Quaternaire* 59, 211–262.
- 459 Edwards, M.E., Bigelow, N.H., Finney, B.P., Eisner, W.R., 2000. Records of aquatic pollen and  
460 sediment properties as indicators of late-Quaternary Alaskan lake levels. *Journal of*  
461 *Paleolimnology* 24, 55–68.
- 462 Environnement Canada, 2017. Canadian Climate Normals 1971-2000.
- 463 Erni, S., Arseneault, D., Parisien, M.-A., Bégin, Y., 2017. Spatial and temporal dimensions of fire  
464 activity in the fire-prone eastern Canadian taiga. *Global Change Biology* 23, 1152–1166.
- 465 Erni, S., Wang, X., Taylor, S., Boulanger, Y., Swystun, T., Flannigan, M., Parisien, M.-A., 2020.  
466 Developing a two-level fire regime zonation system for Canada. *Canadian Journal of Forest*  
467 *Research* 50, 259–273.
- 468 Faegri, K., Iversen, J., 1989. *Textbook of Pollen Analysis*, 4th ed. John Wiley & Sons Ltd., Chichester.
- 469 Flannigan, M., Cantin, A.S., De Groot, W.J., Wotton, M., Newbery, A., Gowman, L.M., 2013. Global  
470 wildland fire season severity in the 21st century. *Forest Ecology and Management* 294, 54–61.

- 471 Flannigan, M.D., Wotton, B.M., Marshall, G.A., De Groot, W.J., Johnston, J., Jurko, N., Cantin, A.S.,  
472 2016. Fuel moisture sensitivity to temperature and precipitation: climate change implications.  
473 *Climatic Change* 134, 59–71.
- 474 Fréchette, B., Richard, P.J., Grondin, P., Lavoie, M., Larouche, A.C., 2018. Histoire postglaciaire de la  
475 végétation et du climat des pessières et des sapinières de l'ouest du Québec, Mémoire de  
476 Recherche Forestière. Gouvernement du Québec, ministère des Forêts, de la Faune et des Parcs,  
477 Direction de la recherche forestière.
- 478 Gaboriau, D.M., Asselin, H., Ali, A.A., Hély, C., Girardin, M.P., under review. Main drivers of recent  
479 extreme wildfire years and identification of climatic thresholds on the territory of the Tłı̨ch̨ First  
480 Nation, northwestern Canada. *International Journal of Wildland Fire*.
- 481 Gajewski, K., Bunbury, J., Vetter, M., Kroeker, N., Khan, A.H., 2014. Paleoenvironmental studies in  
482 southwestern Yukon. *Arctic* 67, 58–70.
- 483 Gajewski, K., Payette, S., Ritchie, J.C., 1993. Holocene vegetation history at the boreal-forest–shrub-  
484 tundra transition in north-western Quebec. *Journal of Ecology* 81, 433–443.
- 485 Glew, J.R., 1991. Miniature gravity corer for recovering short sediment cores. *Journal of*  
486 *Paleolimnology* 5, 285–287.
- 487 Hansen, W.D., Fitzsimmons, R., Olnes, J., Williams, A.P., in press. An alternate vegetation type proves  
488 resilient and persists for decades following forest conversion in the North American boreal  
489 biome. *Journal of Ecology*. <https://doi.org/10.1111/1365-2745.13446>
- 490 Hassol, S.J., 2005. Arctic Climate Impact Assessment - Scientific Report. Cambridge University Press.
- 491 Hély, C., Chaste, E., Girardin, M.P., Remy, C., Blarquez, O., Bergeron, Y., Ali, A.A., 2020. A Holocene  
492 perspective of vegetation controls on seasonal boreal wildfire sizes using numerical  
493 paleoecology. *Frontiers in Forests and Global Change* 3, 106.
- 494 Hennebelle, A., Aleman, J.C., Ali, A.A., Bergeron, Y., Carcaillet, C., Grondin, P., Landry, J., Blarquez,  
495 O., 2020. The reconstruction of burned area and fire severity using charcoal from boreal lake  
496 sediments. *Holocene*.
- 497 Higuera, P., 2009. CharAnalysis 0.9: diagnostic and analytical tools for sediment charcoal analysis.  
498 User's Guide, Montana State University, Bozeman, MT.
- 499 Hoecker, T.J., Higuera, P.E., Kelly, R., Hu, F.S., 2020. Arctic and boreal paleofire records reveal drivers  
500 of fire activity and departures from Holocene variability. *Ecology* 0, e03096.
- 501 IPCC, 2014. Climate Change 2014: Synthesis Report Summary for Policymakers. Contribution of  
502 Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on  
503 Climate Change.
- 504 Jain, P., Wang, X., Flannigan, M.D., 2018. Trend analysis of fire season length and extreme fire weather  
505 in North America between 1979 and 2015. *International Journal of Wildland Fire* 26, 1009–  
506 1020.
- 507 Juggins, S., 2017. Rioja: analysis of Quaternary science data. R package version (0.9–21).
- 508 Kasischke, E.S., Turetsky, M.R., 2006. Recent changes in the fire regime across the North American  
509 boreal region—Spatial and temporal patterns of burning across Canada and Alaska. *Geophysical*  
510 *Research Letters* 33, L09703.
- 511 Kaufman, D.S., Ager, T.A., Anderson, N.J., Anderson, P.M., Andrews, J.T., Bartlein, P.J., Brubaker,  
512 L.B., Coats, L.L., Cwynar, L.C., Duvall, M.L., 2004. Holocene thermal maximum in the western  
513 Arctic (0–180 W). *Quaternary Science Reviews* 23, 529–560.
- 514 Kelly, R., Chipman, M.L., Higuera, P.E., Stefanova, I., Brubaker, L.B., Hu, F.S., 2013. Recent burning  
515 of boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of the National*  
516 *Academy of Sciences* 110, 13055–13060.

- 517 Kobashi, T., Menviel, L., Jeltsch-Thömmes, A., Vinther, B.M., Box, J.E., Muscheler, R., Nakaegawa,  
518 T., Pfister, P.L., Döring, M., Leuenberger, M., 2017. Volcanic influence on centennial to  
519 millennial Holocene Greenland temperature change. *Scientific Reports* 7.
- 520 Kochtubajda, B., Stewart, R.E., Flannigan, M.D., Bonsal, B.R., Cuell, C., Mooney, C.J., 2019. An  
521 assessment of surface and atmospheric conditions associated with the extreme 2014 wildfire  
522 season in Canada's Northwest Territories. *Atmosphere-Ocean* 57, 73–90.
- 523 Latifovic, R., Pouliot, D., Olthof, I., 2017. Circa 2010 land cover of Canada: Local optimization  
524 methodology and product development. *Remote Sensing* 9, 1098.
- 525 Lauriol, B., Lacelle, D., Labrecque, S., Duguay, C.R., Telka, A., 2009. Holocene evolution of lakes in  
526 the Bluefish Basin, northern Yukon, Canada. *Arctic* 62, 212–224.
- 527 Lecavalier, B.S., Fisher, D.A., Milne, G.A., Vinther, B.M., Tarasov, L., Huybrechts, P., Lacelle, D.,  
528 Main, B., Zheng, J., Bourgeois, J., 2017. High Arctic Holocene temperature record from the  
529 Agassiz ice cap and Greenland ice sheet evolution. *Proceedings of the National Academy of  
530 Sciences* 114, 5952–5957.
- 531 MacDonald, G., 1995. Vegetation of the continental Northwest Territories at 6 ka BP. *Géographie  
532 physique et Quaternaire* 49, 37–43.
- 533 MacDonald, G.M., Edwards, T.W., Moser, K.A., Pienitz, R., Smol, J.P., 1993. Rapid response of  
534 treeline vegetation and lakes to past climate warming. *Nature* 361, 243–246.
- 535 Macumber, A.L., Neville, L.A., Galloway, J.M., Patterson, R.T., Falck, H., Swindles, G., Crann, C.,  
536 Clark, I., Gammon, P., Madsen, E., 2011. Paleoclimatological Assessment of the Northwest  
537 Territories and Implications for the Long-Term Viability of the Tibbitt to Contwoyto Winter  
538 Road, Part II: March 2010 Field Season Results. Open report 2010-010. Northwest Territories.  
539 Geoscience office, NT.
- 540 Mann, M.E., Rahmstorf, S., Kornhuber, K., Steinman, B.A., Miller, S.K., Coumou, D., 2017. Influence  
541 of anthropogenic climate change on planetary wave resonance and extreme weather events.  
542 *Scientific Reports* 7.
- 543 Meehl, G.A., Stocker, T.F., Collins, W.D., Friedlingstein, P., Gaye, T., Gregory, J.M., Kitoh, A., Knutti,  
544 R., Murphy, J.M., Noda, A., 2007. Global climate projections.
- 545 Mekonnen, Z.A., Riley, W.J., Randerson, J.T., Grant, R.F., Rogers, B.M., 2019. Expansion of high-  
546 latitude deciduous forests driven by interactions between climate warming and fire. *Nature Plants*  
547 5, 952–958.
- 548 Mooney, S.D., Tinner, W., 2011. The analysis of charcoal in peat and organic sediments. *Mires and Peat*  
549 7, 1–18.
- 550 Moser, K.A., MacDonald, G.M., 1990. Holocene vegetation change at treeline north of Yellowknife,  
551 Northwest Territories, Canada. *Quaternary Research* 34, 227–239.
- 552 Motic Images, 2018. Motic Images Plus 3.0 ML Software.
- 553 Mudelsee, M., 2013. Climate time series analysis, classical statistical and bootstrap methods. Springer,  
554 Dordrecht.
- 555 Mustaphi, C.J.C., Davis, E.L., Perreault, J.T., Pisaric, M.F., 2015. Spatial variability of recent  
556 macroscopic charcoal deposition in a small montane lake and implications for reconstruction of  
557 watershed-scale fire regimes. *Journal of Paleolimnology* 54, 71–86.
- 558 Nolan, R.H., Boer, M.M., Collins, L., Resco de Dios, V., Clarke, H., Jenkins, M., Kenny, B., Bradstock,  
559 R.A., 2020. Causes and consequences of eastern Australia's 2019-20 season of mega-fires.  
560 *Global Change Biology* 26, 3756–3758. <https://doi.org/10.1111/gcb.14987>
- 561 NTENR, 2015. 2014 Fire season review report (No. 249– 17(5)). Legislative assembly of the Northwest  
562 Territories [https://www.assembly.gov.nt.ca/sites/default/files/td\\_249-175.pdf](https://www.assembly.gov.nt.ca/sites/default/files/td_249-175.pdf).

- 563 Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V., Underwood, E.C.,  
564 D'amico, J.A., Itoua, I., Strand, H.E., Morrison, J.C., 2001. Terrestrial ecoregions of the world: a  
565 new map of life on Earth. *BioScience* 51, 933–938.
- 566 Oris, F., Ali, A.A., Asselin, H., Paradis, L., Bergeron, Y., Finsinger, W., 2014. Charcoal dispersion and  
567 deposition in boreal lakes from 3 years of monitoring: differences between local and regional  
568 fires. *Geophysical Research Letters* 41, 6743–6752.
- 569 Parisien, M.-A., Barber, Q.E., Hirsch, K.G., Stockdale, C.A., Erni, S., Wang, X., Arseneault, D., Parks,  
570 S.A., 2020. Fire deficit increases wildfire risk for many communities in the Canadian boreal  
571 forest. *Nature Communications* 11, 2121.
- 572 Pienitz, R., Smol, J.P., MacDonald, G.M., 1999. Paleolimnological reconstruction of Holocene climatic  
573 trends from two boreal treeline lakes, Northwest Territories, Canada. *Arctic, Antarctic, and  
574 Alpine Research* 31, 82–93.
- 575 Polanco-Martinez, J.M., Medina-Elizalde, M.A., Goni, M.F.S., Mudelsee, M., 2019. BINCOR: An R  
576 package for Estimating the Correlation between Two Unevenly Spaced Time Series. *R Journal*  
577 11, 170–184.
- 578 Porter, T.J., Schoenemann, S.W., Davies, L.J., Steig, E.J., Bandara, S., Froese, D.G., 2019. Recent  
579 summer warming in northwestern Canada exceeds the Holocene thermal maximum. *Nature  
580 Communications* 10, 1631.
- 581 Power, M.J., Marlon, J., Ortiz, N., Bartlein, P.J., Harrison, S.P., Mayle, F.E., Ballouche, A., Bradshaw,  
582 R.H., Carcaillet, C., Cordova, C., 2008. Changes in fire regimes since the Last Glacial  
583 Maximum: an assessment based on a global synthesis and analysis of charcoal data. *Climate  
584 Dynamics* 30, 887–907.
- 585 Price, D.T., Alfaro, R.I., Brown, K.J., Flannigan, M.D., Fleming, R.A., Hogg, E.H., Girardin, M.P.,  
586 Lakusta, T., Johnston, M., McKenney, D.W., 2013. Anticipating the consequences of climate  
587 change for Canada's boreal forest ecosystems. *Environmental Reviews* 21, 322–365.
- 588 Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H.,  
589 Edwards, R.L., Friedrich, M., 2013. IntCal13 and Marine13 radiocarbon age calibration curves  
590 0–50,000 years cal. BP. *Radiocarbon* 55, 1869–1887.
- 591 Remy, C.C., Lavoie, M., Girardin, M.P., Hély, C., Bergeron, Y., Grondin, P., Oris, F., Asselin, H., Ali,  
592 A.A., 2017. Wildfire size alters long-term vegetation trajectories in boreal forests of eastern  
593 North America. *Journal of Biogeography* 44, 1268–1279.
- 594 Richard, P., 1970. Atlas pollinique des arbres et de quelques arbustes indigènes du Québec. *Naturaliste  
595 Canadien* 97, 1–34; 97-161; 241–306.
- 596 Ritchie, J.C., 1985. Late-Quaternary Climatic and Vegetational Change in the Lower Mackenzie Basin,  
597 Northwest Canada. *Ecology* 66, 612–621.
- 598 Ritchie, J.C., Cwynar, L.C., Spear, R.W., 1983. Evidence from north-west Canada for an early Holocene  
599 Milankovitch thermal maximum. *Nature* 305, 126–128.
- 600 SFOR, 2018. News and views: the burning hot summer of 2018. *Scandinavian Journal of Forest  
601 Research* 33, 724–727.
- 602 Smith, A., Schismenos, S., Stevens, G., Hutton, L., Chalaris, M., Emmanouloudis, D., 2019.  
603 Understanding large-scale fire events: megafires in Attica, Greece and California, USA. *United  
604 Nations Major Group for Children and Youth, USA*, pp. 29–34.
- 605 Stephens, S.L., Burrows, N., Buyantuyev, A., Gray, R.W., Keane, R.E., Kubian, R., Liu, S., Seijo, F.,  
606 Shu, L., Tolhurst, K.G., 2014. Temperate and boreal forest mega-fires: characteristics and  
607 challenges. *Frontiers in Ecology and the Environment* 12, 115–122.  
608 <https://doi.org/10.1890/120332>

- 609 Sulphur, K.C., Goldsmith, S.A., Galloway, J.M., Macumber, A., Griffith, F., Swindles, G.T., Patterson,  
610 R.T., Falck, H., Clark, I.D., 2016. Holocene fire regimes and treeline migration rates in sub-  
611 arctic Canada. *Global and Planetary Change* 145, 42–56.
- 612 Swain, A.M., 1973. A history of fire and vegetation in Northeastern Minnesota as recorded in lake  
613 sediments. *Quaternary Research* 3, 383–390.
- 614 Timoney, K.P., Mamet, S.D., Cheng, R., Lee, P., Robinson, A.L., Downing, D., Wein, R.W., 2019. Tree  
615 cover response to climate change in the forest-tundra of north-central Canada: fire-driven  
616 decline, not northward advance. *Ecoscience* 26, 133–148.
- 617 Tinner, W., Hu, F.S., Beer, R., Kaltenrieder, P., Scheurer, B., Krähenbühl, U., 2006. Postglacial  
618 vegetational and fire history: pollen, plant macrofossil and charcoal records from two Alaskan  
619 lakes. *Vegetation History and Archaeobotany* 15, 279–293.
- 620 Trachsel, M., Telford, R.J., 2017. All age–depth models are wrong, but are getting better. *Holocene* 27,  
621 860–869.
- 622 Turco, M., von Hardenberg, J., AghaKouchak, A., Llasat, M.C., Provenzale, A., Trigo, R.M., 2017. On  
623 the key role of droughts in the dynamics of summer fires in Mediterranean Europe. *Scientific*  
624 *Reports* 7, 81.
- 625 Uptier, L.M., Vermaire, J.C., Patterson, R.T., Crann, C.A., Galloway, J.M., Macumber, A.L., Neville,  
626 L.A., Swindles, G.T., Falck, H., Roe, H.M., 2014. Middle to late Holocene chironomid-inferred  
627 July temperatures for the central Northwest Territories, Canada. *Journal of Paleolimnology* 52,  
628 11–26.
- 629 Veraverbeke, S., Rogers, B.M., Goulden, M.L., Jandt, R.R., Miller, C.E., Wiggins, E.B., Randerson,  
630 J.T., 2017. Lightning as a major driver of recent large fire years in North American boreal  
631 forests. *Nature Climate Change* 7, 529–534.
- 632 Viau, A.E., Gajewski, K., 2009. Reconstructing millennial-scale, regional paleoclimates of boreal  
633 Canada during the Holocene. *Journal of Climate* 22, 316–330.
- 634 Viau, A.E., Gajewski, K., Sawada, M.C., Fines, P., 2006. Millennial-scale temperature variations in  
635 North America during the Holocene. *Journal of Geophysical Research: Atmospheres* 111,  
636 D09102.
- 637 Vincent, J.-S., 1973. A palynological study for the Little Clay Belt, northwestern Quebec. *Naturaliste*  
638 *Canadien* 100, 59–70.
- 639 Walker, X.J., Baltzer, J.L., Cumming, S.G., Day, N.J., Johnstone, J.F., Rogers, B.M., Solvik, K.,  
640 Turetsky, M.R., Mack, M.C., 2018. Soil organic layer combustion in boreal black spruce and  
641 jack pine stands of the Northwest Territories, Canada. *International Journal of Wildland Fire* 27,  
642 125–134.
- 643 Wang, X., Parisien, M.-A., Taylor, S.W., Candau, J.-N., Stralberg, D., Marshall, G.A., Little, J.M.,  
644 Flannigan, M.D., 2017. Projected changes in daily fire spread across Canada over the next  
645 century. *Environmental Research Letters* 12, 025005.
- 646 Wang, Y., Hogg, E.H., Price, D.T., Edwards, J., Williamson, T., 2014. Past and projected future changes  
647 in moisture conditions in the Canadian boreal forest. *Forestry Chronicle* 90, 678–691.
- 648 Williams, J.W., 2006. An atlas of pollen-vegetation-climate relationships for the United States and  
649 Canada. American Association of Stratigraphic Palynologists Foundation, Dallas, Texas.
- 650 Winkler, M.G., 1985. Charcoal analysis for paleoenvironmental interpretation: a chemical assay.  
651 *Quaternary Research* 23, 313–326.
- 652 Wotton, B.M., Flannigan, M.D., Marshall, G.A., 2017. Potential climate change impacts on fire intensity  
653 and key wildfire suppression thresholds in Canada. *Environmental Research Letters* 12, 095003.



654 Zhang, X., Brown, R., Vincent, L., Skinner, W., Feng, Y., Mekis, E., 2011. Canadian climate trends,  
655 1950–2007, in: Canadian Biodiversity: Ecosystem Status and Trends 2010, Technical Thematic  
656 Report No. 5. Canadian Councils of Resource Ministers, Ottawa, ON.

**Table 1.** Characteristics of the four studied lakes and sediment records.

<i>Lakes</i>	<b>Emile</b>	<b>Izaak</b>	<b>Paradis</b>	<b>Saxon</b>
--------------	--------------	--------------	----------------	--------------

---

*Characteristics*

Latitude (°N)	64°03'16.64"	64°05'56.21"	64°00'23.54"	63°48'26.06"
Longitude (°W)	114°06'21.67"	114°10'33.08"	114°59'19.24"	114°58'33.29"
Surface area (ha)	1.4	1.5	1.8	0.9
Elevation (m above sea level)	393	394	347	392
Maximum water depth (cm)	150	115	235	540
Maximum length (m)	203	162	271	177
Lakeshores (Flat / Abrupt)	F	F	F	A
Fluvial input (Present / Absent)	A	A	A	A
Date of the last local fire (CNFD)	2014	2014	1979	1998


*Sediment records*

Total length of sediment (cm)	456	359	450	395
Depth of the basal <sup>14</sup> C date (cm)	448	340	441	350
Sedimentation rate (cm year <sup>-1</sup> )	0.049	0.036	0.046	0.036
Median signal-to-noise index	5.4	4.2	4.1	3.7
Total number of significant fire events	57	49	55	46


---

**Table 2.** BINCOR Pearson correlation coefficient (with 95% confidence interval) between fire metrics, temperature, total pollen accumulation rate (PAR), and accumulation rates of taxa with an average percentage greater than 0.1% over the entire study period. \*\*\* p < 0.001, \*\* p < 0.01 and \* p < 0.05.


	<i>RegFF</i>	<i>RegBB</i>	<i>FS index</i>	Temperature
<b>Temperature</b>	-0.53 [-0.63, -0.42]***	0.21 [0.07, 0.35]**	0.66 [0.58, 0.74]***	-
<b>PAR</b>	0.50 [0.35, 0.62]***	0.58 [0.45, 0.68]***	0.14 [-0.04, 0.31]	-0.13 [-0.30, 0.04]
<i>Picea</i>	0.37 [0.21, 0.51]***	0.64 [0.53, 0.74]***	0.29 [0.12, 0.44]***	0.00 [-0.17, 0.17]
<i>Pinus</i>	0.63 [0.52, 0.72]***	-0.16 [-0.32, 0.00]	-0.56 [-0.67, -0.44]***	-0.52 [-0.63, -0.39]***
<i>Betula</i>	0.27 [0.12, 0.42]***	0.31 [0.16, 0.45]***	0.09 [-0.08, 0.25]	-0.06 [-0.22, 0.1]
<i>Populus</i>	-0.34 [-0.47, -0.18]***	0.04 [-0.13, 0.20]	0.28 [0.12, 0.43]***	0.18 [0.02, 0.34]*
<i>Alnus alnobetula</i> subsp. <i>crispa</i>	0.40 [0.25, 0.53]***	0.42 [0.28, 0.55]***	0.09 [-0.08, 0.25]	-0.02 [-0.18, 0.14]
<i>Alnus incana</i> subsp. <i>rugosa</i>	0.23 [0.07, 0.38]**	0.16 [0.00, 0.32]*	-0.02 [-0.18, 0.15]	-0.14 [-0.30, 0.02]
<i>Juniperus</i>	0.09 [-0.08, 0.25]	-0.30 [-0.45, -0.15]***	-0.33 [-0.47, -0.17]***	-0.38 [-0.51, -0.23]***
<i>Salix</i>	0.21 [0.05, 0.36]*	0.10 [-0.06, 0.26]	-0.04 [-0.21, 0.12]	-0.17 [-0.32, -0.01]*
<i>Larix</i>	0.21 [0.05, 0.36]*	-0.23 [-0.38, -0.07]**	-0.34 [-0.48, -0.19]***	-0.26 [-0.41, -0.1]**
<b>Cyperaceae</b>	0.20 [0.03, 0.35]*	-0.01 [-0.17, 0.15]	-0.13 [-0.29, 0.03]	-0.19 [-0.34, -0.03]*
<b>Ericaceae</b>	0.14 [-0.03, 0.29]	0.11 [-0.06, 0.27]	0.00 [-0.16, 0.17]	0.01 [-0.15, 0.17]
<i>Artemisia</i>	-0.05 [-0.22, 0.11]	0.10 [-0.06, 0.26]	0.13 [-0.04, 0.29]	0.15 [-0.01, 0.31]
<i>Myrica</i>	-0.46 [-0.58, -0.32]***	-0.01 [-0.17, 0.16]	0.33 [0.17, 0.47]***	0.38 [0.23, 0.51]***
<i>Lycopodium</i>	-0.31 [-0.45, -0.15]***	-0.05 [-0.21, 0.12]	0.19 [0.03, 0.35]*	0.2 [0.04, 0.35]*
<b>Poaceae</b>	0.23 [0.07, 0.38]**	-0.04 [-0.21, 0.12]	-0.19 [-0.34, -0.02]*	-0.24 [-0.39, -0.08]**
<i>Pediastrum</i>	0.28 [0.12, 0.43]***	-0.23 [-0.38, -0.07]**	-0.37 [-0.5, -0.22]***	-0.23 [-0.38, -0.07]**
<i>Potamogeton</i>	-0.14 [-0.30, 0.02]	0.14 [-0.02, 0.30]	0.22 [0.06, 0.37]**	0.10 [-0.06, 0.26]
<i>Nuphar</i>	0.38 [0.23, 0.51]***	0.01 [-0.15, 0.18]	-0.24 [-0.39, -0.08]**	-0.24 [-0.39, -0.08]**




**Figure 1.** Locations of the studied lakes in the Northwest Territories (north-central Canada), showing ecozones (adapted from Olson et al., 2001) and treeline (adapted from Timoney et al., 2019).



**Figure 2.** WinBacon outputs for lakes *Emile*, *Izaak*, *Paradis* and *Saxon*. Upper left panels describe the MCMC (Markov Chain Monte Carlo) iterations (the distribution is stationary with little structure among neighbouring iterations). Upper middle panels show the distribution of sediment accumulation rates. Upper right panels show the memory corresponding to the variation of sediment accumulation rate in time. Main panels show the calibrated <sup>14</sup>C dates (see Table S1 for full details on the chronology) and age-depth models (with 95% confidence intervals).




**Figure 3.** Holocene fire activity anomalies interpolated using a 500-year bandwidth smoothing, relative to the 500-0 cal. yrs. BP reference period (black horizontal line) for individual and regional fire-history reconstructions based on sediment charcoal records from lakes *Emile*, *Izaac*, *Paradis* and *Saxon*: (A) Regional Fire Frequency (*RegFF*), (B) Regional Biomass Burning (*RegBB*) and (C) Fire Size/Severity (*FS* index) based on regional biomass burning and fire frequency. Yellow shaded areas in each panel represent the 90% bootstrap confidence intervals.




**Figure 4.** Sedimentation ( $\text{cm}^{-1} \text{ year}^{-1}$ ) and pollen accumulation rates ( $\text{grains cm}^{-2} \text{ year}^{-1}$ ) at lake *Emile* for total terrestrial pollen (PAR, in red) and for the main taxa: trees (in brown), shrubs (in green), herbaceous plants (in yellow) and green algae (*Coenobium*) and aquatic plants (in blue) all having an average percentage greater than 0.1% over the entire study period. Pale areas represent  $\times 5$  exaggeration. Note that the scale differs for each pollen or spore taxon.





**Figure 5.** Mean summer temperature anomalies over the Holocene, relative to the 500-0 cal. yrs. BP reference period, obtained from the calculation of the means of standardized independent temperature datasets described in Appendix S4. The yellow shaded area indicates the 90% bootstrap confidence interval.



**Figure 6.** Dryness periods inferred from hydrological conditions from different locations in north-central and northwestern Canada. Distance from the study area increases from top to bottom. (a) Central Northwest Territories (Pienitz et al., 1999; dissolved organic carbon (DOC) inferred from diatom assemblages). (b) Mackenzie region (50°-70°N, 120°-140°W; Viau and Gajewski, 2009; annual precipitation inferred from pollen assemblages). (c) Northern Yukon (Lauriol et al., 2009; lake-level inferred from plant macrofossil analysis).