



## **Push-pull strategy combined with net houses for controlling cowpea insect pests and enhancing crop yields**

Seydou Diabate, Thibaud Martin, Lucy Kananu Murungi, Komi K.M. Fiaboe, John Wesonga, John M. Kimani, Emilie Deletre

### **► To cite this version:**

Seydou Diabate, Thibaud Martin, Lucy Kananu Murungi, Komi K.M. Fiaboe, John Wesonga, et al.. Push-pull strategy combined with net houses for controlling cowpea insect pests and enhancing crop yields. *Crop Protection*, 2021, 141, pp.105480 -. <10.1016/j.cropro.2020.105480>. <hal-03493614>

**HAL Id: hal-03493614**

**<https://hal.science/hal-03493614v1>**

Submitted on 15 Dec 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY-NC 4.0 - Attribution - Non-commercial use - International License

**Push-pull strategy combined with net houses for controlling cowpea insect pests and enhancing crop yields**

Seydou Diabate<sup>1,2,3</sup>, Thibaud Martin<sup>1,3,4</sup>, Lucy Kananu Murungi<sup>2</sup>, Komi K. M. Fiaboe<sup>5</sup>, John Wesonga<sup>2</sup>, John M. Kimani<sup>6</sup>, Emilie Deletre<sup>1,3,4</sup>

<sup>1</sup>International Centre of Insect Physiology and Ecology, P.O. Box 30772-00100, Nairobi, Kenya

<sup>2</sup>Jomo Kenyatta University of Agriculture and Technology, P.O Box 62000-00200, Nairobi, Kenya

<sup>3</sup>CIRAD UPR HortSys, 34398 Montpellier, France

<sup>4</sup>Université de Montpellier, 34090 Montpellier, France

<sup>5</sup>IITA-Cameroon, BP. 2008 (Messa), IRAD Main Road, Nkolbisson, Yaoundé, Cameroon

<sup>6</sup>Kenya Agricultural and Livestock Research Organization, Industrial Crops Research Institute Mtwapa, P.O. Box 16-80109, Mtwapa, Kenya

Corresponding author: [seydoudiabate01@gmail.com](mailto:seydoudiabate01@gmail.com)

## Abstract

Net houses can be used in tropical environments to protect crops such as cowpea against large insect pests, thereby avoiding pesticide treatments while sustainably mitigating the effects of climate change. We investigated a push-pull strategy to prevent small insect pest outbreaks in a net house. The push component consisted of two stimulus plants, i.e. *Cymbopogon citratus* and *Tagetes minuta*, and the pull stimuli consisted of visual cues from blue and yellow sticky traps. Field experiments were set up in central Kenya and conducted during a rainy and a dry season, involving an open field control treatment, and three management treatments consisting of (1) an open field push-pull treatment, (2) a net house treatment and (3) a combined net house + push-pull treatment. *Trialeurodes vaporariorum* infestations were lower in the net house and net house + push-pull treatments than in the two open field treatments during the dry period or in the control treatment during the rainy period. *Aphis craccivora* infestations were higher in the net house and net house + push-pull treatments than in the control and open field push-pull treatments during the dry period, while no differences were observed among treatments during the rainy period. *Megalurothrips sjostedti* infestations did not vary among treatments in both periods. Among the larger insect pests, *Clavigralla tomentosicollis* infestations were lower in the net house and net house + push-pull treatments than in the open field treatments during the dry period, while *Maruca vitrata* infestations were lower in the net house treatment than in the control. During the rainy period, *C. tomentosicollis* infestations were higher in the net house + push-pull treatment than in the net house treatment, whereas *M. vitrata* infestations did not vary among treatments. Compared to the control, *Empoasca* sp. infestations were lower in the net house and net house + push-pull treatments in both periods, and in the open field push-pull treatment in the rainy period. Cowpea pod and grain yield and quality were higher in the net house and net house + push-pull treatments than in the control irrespective of the period. Although the treatments 1 reduced some of the pests, the net house and net house + push-pull treatments were effective in protecting cowpeas against most of the pests while improving pod yields in both periods.

Keywords: Net house, integrated pest management, push-pull, sticky traps, insect pests

## 1. Introduction

Population growth in developing countries has increased the demand for safe nutritional food (Alexandratos and Bruinsma, 2012). Vegetable production plays an important role in feeding the growing urban population in sub-Saharan regions (OECD and FAO, 2016). Farmers have been resorting to intensive chemical pesticide treatments to reduce crop pest damage and thus help meet the rising food demand (de Bon et al., 2014). Chemical pesticides are effective for pest control but they often have concomitant negative impacts on beneficial arthropods, the environment and human health (de Bon et al., 2014). Netting technology provides an alternative for reducing chemical pesticide treatments (Martin et al., 2006, 2015) by protecting different vegetable crops against a wide range of pests, boosting soil moisture, stabilizing the air temperature and increasing crop yields and quality (Martin et al., 2006; Gogo et al., 2012; Saidi et al., 2013; Simon et al., 2014; Gogo et al., 2014b; Nordey et al., 2017). Knitted nets with a 30-40 mesh size (0.9-0.4 mm diameter) have proven effective as a barrier against large size insect pests (body length > 2 mm) such as lepidopterans and dipterans, while providing enough ventilation to mitigate the adverse climate conditions that prevail particularly in the Kenyan highlands (Martin et al., 2015). However, net houses with such netting do not provide an effective barrier to small insects (body length < 2 mm) as they can pass through the mesh.

We investigated a push-pull crop protection strategy based on insect cues combined with the use of netting technology to curb outbreaks of small insect pests in net houses. In the push-pull strategy, attractant and repellent stimuli from different sources are simultaneously used to control the spatial distribution of insect pests and thus reduce their abundance on the target crop (Cook et al., 2006). Attractant stimuli divert insects from the target crop by means of taste, egg-laying stimulants, volatile plant attractants, sexual pheromones, and visual stimulants. Repellent stimuli prevent insect pests from finding or accepting its host via the emission of repellent volatiles, alarm pheromones, anti-feeding agents, visual distractions, egg-laying repellents and irritants (Cook et al., 2006; Eigenbrode et al., 2016).

Cowpea, (*Vigna unguiculata* L. Walp) is an African indigenous vegetable (AIV) that can reduce food insecurity and malnutrition in Africa (Muniu, 2017) since about 200 million people consume cowpea grains or leaves (Popelka et al., 2006). For example, cowpea is the top ranking AIV cropped in Kenya, with a total of 65 million t produced over a 24,431 ha area (HCDA, 2014). Netting technology and the push-pull strategy could be combined to provide an

alternative solution for controlling large and small insect pests of cowpea, thus reducing the need for pesticide treatments.

The pod borer (*Maruca vitrata* Fabricius) (Lepidoptera: Crambidae), and the pod bug (*Clavigralla tomentosicollis* Stål) (Hemiptera: Coreidae), are two large insect pests of cowpea that can cause 80% yield losses if no pesticides are applied (Ekesi et al., 2002; OECD, 2015). *C. tomentosicollis* feeds extensively on fresh pods (Koono et al., 2002), whereas *M. vitrata* feeds on both flowers and pods (Singh and Jackai, 1988). The bean flower thrips (*Megalurothrips sjostedti* Trybom) (Thysanoptera: Thripidae), and the black legume aphid (*Aphis craccivora* Koch) (Homoptera: Aphididae), are two small insect pests of cowpea that cause yield losses of 20–100%, depending on the outbreak severity (Ekesi et al., 1999; Obopile, 2006; Abtew et al., 2016; Mweke et al., 2018). The feeding activity of *M. sjostedti* larvae and adults can lead to flower abortion and shedding, thereby resulting in reduced crop yields, while heavy *A. craccivora* infestations may cause plant stunting and delayed flowering (Obopile and Ositile, 2010; Moritz et al., 2013). In addition, *A. craccivora* may transmit viruses to plants (Stoetzel and Miller, 2001; Borowiak-Sobkowiak et al., 2017). A previous study showed that *M. sjostedti* is attracted by blue colours (Muvea et al., 2014), while *A. craccivora* is attracted to yellow sticky traps (Webb et al., 1994)—we thus used sticky traps as a pull stimulus in our study. Our previous experiment using olfactory tests showed that Mexican marigold (*Tagetes minuta* L.), and lemongrass (*Cymbopogon citratus* DC. Stapf), plants in the vegetative stage were repellent to female *M. sjostedti* (Diabate et al., 2019a). Freshly cut lemongrass leaves produce citral, an organic volatile which is repellent to *M. sjostedti*. Mexican marigold and lemongrass are either repellent or insecticidal to *A. craccivora* (Morillo-Rejesus and Decena, 1982; Ofuya and Okuku, 1994). In this study, we combined these repellent push plants with the use of a net house. Insect control using a push-pull strategy can avoid the need for chemical insecticides, thereby increasing the opportunity of natural enemies to reduce insect pest populations (Khan et al., 1997).

We hypothesised that nets could effectively protect cowpea crops against large insect pests but not small insect pests. The aim of the present study was to evaluate the effectiveness of three control strategies against cowpea pests, by combining (a) use of an insect net house to provide a physical barrier, (b) repellent companion plants to provide an olfactory barrier, and (c) a visual trap in the form of coloured sticky strips as a pest attractant to increase cowpea grain yields and quality.

## 2. Materials and Methods

## 2.1. Site

The field trial was conducted at Kenya Agricultural and Livestock Research Organization (KALRO), Industrial Crops Research Centre (ICRC) Mwea (0°37'09.0"S 37°22'09.4"E) in Kirinyaga County, central Kenya. Mwea is one of the main areas in Kenya where vegetables, including cowpea, are grown (Musebe et al., 2005). The four seasons in Kenya include a short rainy season (October-November), a hot dry season (January-March), a long rainy season (March-June), and a cold dry season (July-August) (Hassan, 1998; Foba et al., 2015). The experiment was conducted during two periods: in the dry period from 3 October 2017 to 16 January 2018, and in the rainy period from 30 January 2018 to 22 May 2018. The dry period included one rainy month (10 October to 7 November 2017) and three dry months (14 November to 9 January 2018), with 358.8 mm total rainfall and 22.71°C mean temperature. The rainy period included one dry month (30 January 2018 to 27 February 2018) followed by three rainy months (27 February to 15 May 2018), with 679.5 mm total rainfall and 22.50°C mean temperature. Rainfall and temperature data were collected at the KALRO ICRC Mwea Kirogo Research Farm.

## 2.2. Treatments and experimental design

The net house was a locally manufactured high iron framed tunnel covered with a transparent knitted polyethylene AgroNet 0.4 (A to Z Textile Mills, Arusha, Tanzania) with a 40 mesh size (0.4 mm diameter hole size). The high tunnel was 5 m wide, 10 m long and 2.5 m high (flat on top) in the middle and 2 m high at the sides with a double door. Rain drops seeped through the top of the net house. Coloured (blue, yellow) sticky traps were purchased from the RealIPM Company, Thika, Kenya. Cowpea, *V. unguiculata* var. Machakos 66 (M66), seeds were purchased from Dryland Seed Ltd., and lemongrass seedlings were from Simlaw Seeds Company Ltd., in Nairobi, Kenya. Mexican marigold seeds were field collected at KALRO-ICRC Mwea. About 40 kg of cow manure was spread per plot (10 m × 5 m) 2 weeks before sowing the cowpea seeds. Two cowpea seeds were sown in two hills on each side of the micro-irrigation emitters at a distance of 25 cm with 1 m inter-row spacing. Each experimental plot measured 5 m × 10 m and 60 cowpea plants were planted per row, for a total of 300 plants (60 plants × 5 rows) per plot.

The experiment involved a total of sixteen plots, with four replicate plots for each of the four treatments, including: an open field control treatment (open field), and three management treatments consisting of (1) an open field push-

pull treatment, (2) a net house treatment and (3) a combined net house + push-pull strategy treatment. Plots were laid out in a Latin square design with four replicates (Figure 1). The treatment 1 plots included two repellent plants, lemongrass and Mexican marigold as push components (Calumpang et al., 2013) and coloured sticky traps (blue and yellow) as pull components (Webb et al., 1994; Muvea et al., 2014). Mexican marigold was planted at 20 cm intervals and lemongrass at 2 m intervals around the plots. Mexican marigold served as a repellent plant via its continuous release of repellent volatiles. Moreover, on the basis of previous findings that volatiles emitted by freshly cut lemongrass leaves were effective in repelling *M. sjostedti* for 24 h (Diabate et al., 2019a), freshly cut lemongrass leaf pieces (about 5 cm of each leaf cut with scissors once a week) were also applied between the rows as organic mulch to repel *M. sjostedti* and *A. craccivora*. In the treatment 3 plots, Mexican marigold was planted outside the net house to discourage insects from passing through the net and lemongrass was planted inside the net house along the sides to repel insects that managed to enter. The experimental plots were separated from each other by 2-m buffer strips of bare soil. Blue sticky traps were used to catch *M. sjostedti* and yellow sticky traps served to capture other insects including whiteflies and aphids. Four strips of sticky tape (15 cm × 10 m) per push-pull plot were placed at 1 m intervals between the rows, 1 m above the soil and attached to sticks fixed to the ground. The traps were removed and replaced with new ones monthly over the 4-month cowpea cropping period. Insects on 10×10 cm samples of each replaced trap were then counted in the laboratory. No chemical insecticide treatments were carried out during in the field experiment, but a biopesticide—a mixture of 5 ml of liquid soap and 5 g of pepper powder in 20 l of water—was applied when a threshold of 1,000 aphids per plant was reached on three plants/plot in net house treatments. This biopesticide was sprayed twice in all treatments (on 17 and 24 November 2017) to slightly reduce the aphid populations, since high *A. craccivora* infestation of cowpea plants can cause high plant mortality (Annan et al., 1994). Biopesticide sprays were conducted during the dry period to avoid infested plant loss during the field trial, but this was not done during the rainy period. This soap/pepper solution provides a safe means to reduce aphid populations (Poswal and Akpa, 1991; Pahla et al., 2014), but it is less effective than synthetic insecticides in open field conditions (Ahmed et al., 2019; Smaili et al., 2014). To the best of our knowledge, there are no reports of this biopesticide impacting any of the other insects sampled in the study.

### 2.3. Sampling methods of insects and yield

The insect pests were counted 2 weeks after sowing and thereafter on a weekly basis for 14 and 15 weeks during the dry and rainy periods, respectively. Two plants per row were randomly selected and 10 individual plants were monitored in 5 rows per plot, for a total of 160 plants monitored per sampling date. The first and last plants of each row were not used to avoid border effects. The insects were counted between 9 am and 3 pm in all plots. Adult and immature insects were visually counted on different parts of each cowpea plant—large insect pests such as leafhoppers, *Empoasca* sp., brown pod-sucking bugs, *C. tomentosicollis*, and larval bean pod borers, *M. vitrata*, were counted on whole plants. Small adult insects such as bean flower thrips, *M. sjostedti*, black legume aphids, *A. craccivora*, and greenhouse whiteflies, *Trialeurodes vaporariorum* Westwood, were counted on one old lower leaf and one young upper leaf per cowpea plant. *A. craccivora* and *M. sjostedti* were also counted on two picked open flowers, and on two pods per plant. *A. craccivora* was counted on a 5 cm portion of the plant stem (the number of insects counted on 1 cm were multiplied by 5 cm representing the portion of the plant stem).

The number of flowers and pods were counted on 10 plants randomly selected per plot. The harvested pods, undamaged pods or pods damaged by insects were counted and weighed to evaluate the quantity of marketable pods. Cowpea grains harvested on each plot were placed in bags and weighed on a mechanical kitchen scale.

## **2.4. Data analysis**

The data were systematically log transformed ( $\log(x+0.5)$ ) to ensure homogeneity of variance and conformation to normality before using a linear mixed model with random intercepts. Treatments were considered as fixed factors while plots were random factors. The model was fitted using the *lmer* function from the *lme4* package (Bates et al., 2015), and the fixed factor effects were tested using Type II Wald F tests. When the tests revealed a significant treatment effect, means were separated by least squares means (lsmeans) adjusted using the Tukey method. Between-period comparisons of insects counted on sticky traps and treatments were performed via Student's t tests after transformation. R version 3.3.2 software (R Core Team, 2016) was used for all analyses.

## **3. Results**

### **3.1. Colonization of plants by small insect pests in both dry and rainy periods**

In the dry period, cowpea colonisation by insects with piercing-sucking mouthparts such as *T. vaporariorum* and *A. craccivora* started in the control and treatment 1 experiments (open field) 2 and 3 weeks after sowing, respectively



(Figure 2a, 2b). *A. craccivora* infestations were low in the open field treatments compared to the treatments 2 and 3. *T. vaporariorum* was observed on leaves while *A. craccivora* was noted on leaves and stems during cowpea seedling stage and on flowers and pods during the reproductive stage. Peak *T. vaporariorum* infestations were detected 6 weeks after sowing in the control plots, whereas *A. craccivora* outbreaks peaked 8 and 9 weeks after sowing in the treatment 3 and 2 during the flowering and podding stages, respectively. *M. sjostedti* infestations started during the cowpea flowering stage 8 weeks after sowing in both open field treatments (Figure 2c).

In the rainy period, *T. vaporariorum*, *A. craccivora* and *M. sjostedti* infestations began in periods similar to those noted in the dry season treatments (Figure 3a, b, c). *T. vaporariorum* infestations peaked 6 weeks after sowing in the control plots, while *A. craccivora* infestations peaked 7 weeks after sowing in the control plots. Three *M. sjostedti* infestation peaks occurred 13, 14 and 15 weeks after sowing in the treatments 2, 1 and control plots, respectively.

### 3.2. Effects of dry and rainy periods on small insect pests

The between-period comparison showed that in the rainy period small insect pest populations were lower than or equal to those in the dry period (Table 1). In all treatments, the *M. sjostedti* population did not differ the dry and rainy periods. However, *A. craccivora* numbers were higher in the treatments 2 and 3 in the dry period than in the rainy period (treatment 2:  $t = 2.186$ ,  $df = 114$ ,  $p = 0.030$ ; treatment 3:  $t = 4.172$ ,  $df = 114$ ,  $p < 0.001$ ). Conversely, the *A. craccivora* population was higher in the control and treatments 1 in the rainy period than in the dry period (control  $t = -1.899$ ,  $df = 114$ ,  $p = 0.059$ ; treatment 1:  $t = -1.986$ ,  $df = 114$ ,  $p = 0.049$ ). In the control treatment ( $t = 2.846$ ,  $df = 114$ ,  $p = 0.005$ ) and treatment 1 ( $t = 4.010$ ,  $df = 114$ ,  $p < 0.001$ ), *T. vaporariorum* whitefly populations were higher in the dry period than in the rainy period.

### 3.3. Effects of each treatment on small insect pests

#### 3.3.1. *Megalurothrips sjostedti* and *Aphis craccivora*

*M. sjostedti* populations did not vary among treatments in both periods (Table 1). *A. craccivora* infestations were higher in the treatments 2 and 3 than in control and treatment 1 during the dry period. Otherwise, *A. craccivora* infestations did not vary among treatments in the rainy period.

#### 3.3.2. *Trialeurodes vaporariorum*

*T. vaporariorum* numbers were lower in the treatments 2 and 3 than in the open field treatments during the dry period. *T. vaporariorum* populations were higher in the control than in the treatments 2 and 3 in the rainy period.

### **3.4. Effects of dry and rainy periods on large insect pests**

Infestations of the large insect pest *M. vitrata* were low infestations in all treatments, but they were significantly higher in the control during dry period than in the rainy period ( $t = 2.810$ ,  $df = 114$ ,  $p = 0.005$ ). *Empoasca* sp. numbers were found to be highest during the rainy period. *Empoasca* sp. infestations were higher in the open field treatments in the rainy period than in the dry period (control:  $t = -5.449$ ,  $df = 114$ ,  $p < 0.001$ ; treatment 1:  $t = -4.273$ ,  $df = 114$ ,  $p < 0.001$ ). *C. tomentosicollis* infestations were higher in the treatment 2 and 3 during the rainy period than during dry period (treatment 2:  $t = -2.457$ ,  $df = 114$ ,  $p = 0.015$ ; treatment 3:  $t = -4.922$ ,  $df = 114$ ,  $p < 0.001$ ).

### **3.5. Effects of each treatment on large insect pests**

#### *3.5.1. Maruca vitrata*

*M. vitrata* infestations were lower in the treatment 2 than in the control during the dry period, but no differences were observed among treatments during the rainy period.

#### *3.5.2. Empoasca* sp.

*Empoasca* sp. infestations were lower in the treatments 2 and 3 than in both open field treatments during the dry period. *Empoasca* sp. numbers were higher in the control than in all other treatments during the rainy period. In addition, *Empoasca* sp. numbers in the treatment 1 were higher than in the treatments 2 and 3 but lower than in the control.

#### *3.5.3. Clavigralla tomentosicollis*

*C. tomentosicollis* infestations were lower in the treatments 2 and 3 than in control and treatment 1 during the dry period (Table 1). However, *C. tomentosicollis* numbers were higher in the treatment 3 than in the treatment 2 in the rainy period. *C. tomentosicollis* infestations in control and treatment 1 were not significantly different.

### **3.6. Pests caught by coloured sticky traps with the push-pull strategy**

The yellow sticky traps caught higher *T. vaporariorum* numbers in the treatment 1 than in the treatment 3, but no differences were observed between these two treatments with blue sticky traps (Table 2). The blue sticky traps caught more *M. sjostedti* in the treatment 1 than in the treatment 3. No differences were observed in *M. sjostedti* numbers caught on the yellow sticky traps when comparing the treatments 1 with treatments 3. *A. craccivora* numbers caught on yellow and blue sticky traps did not differ between the treatments 1 and 3. Only winged aphids were caught on the traps, whereas mostly nonwinged aphids were detected on cowpea plants.

### **3.7. Influence of the planting period on yield parameters**

There were no significant differences in the number of open flowers when comparing the same treatment in the two periods (Table 3). All yield parameters observed in the present study were lower in both of the open field treatments in the rainy period than in the dry period, including the total harvested pod number (control:  $t = 4.486$ ,  $df = 114$ ,  $p < 0.001$ ; treatment 1:  $t = 4.441$ ,  $df = 114$ ,  $p < 0.001$ ), the total harvested pod weight (control:  $t = 4.469$ ,  $df = 114$ ,  $p < 0.001$ ; treatment 1:  $t = 4.649$ ,  $df = 114$ ,  $p < 0.001$ ), marketable harvested pods (control:  $t = 4.577$ ,  $df = 114$ ,  $p < 0.001$ ; treatment 1:  $t = 4.518$ ,  $df = 114$ ,  $p < 0.001$ ) and the marketable harvested pod weight (control:  $t = 4.528$ ,  $df = 114$ ,  $p < 0.001$ ; treatment 1:  $t = 4.757$ ,  $df = 114$ ,  $p < 0.001$ ). On average, the harvested pod number in the open field treatments was 220-fold (control) and 55-fold (treatment 1) greater during the dry period compared to rainy period. No differences in yield parameters were observed between periods in both of the net house treatments.

### **3.8. Effects of treatments on crop yield**

The open flower numbers did not differ among the treatments in both periods and nor any of the pod parameters differ among the treatments in the dry period (Table 3). However, the total harvested pods and their weight were 2-fold greater in the treatment 2 than in the control during the dry period. In addition, marketable harvested pods were 2-fold greater in the treatments 2 than in both of the open field treatments, while the marketable pod weights were 2-fold greater in the treatments 2 and 3 than in the control during the dry period. During the rainy period, the total harvested pods, harvested pod weight, marketable harvested pods and marketable pod and grain weights were greater in the treatments 2 and 3 than in the control and treatment 1. However, no differences were observed between the control and treatment 1 and between the treatment 2 and the treatment 3 during the rainy period. The grain quality in the plots was only assessed during the rainy period and the treatments 2 and 3 grain was better than that harvested in the open field treatments in the rainy period (Supplementary data).

#### 4. Discussion

The present study showed that net house and net house + push-pull treatments were effective in protecting cowpea crops against most of the major large insect pests present. Previous studies revealed effective protection of vegetable crops grown in high or low tunnels under insect netting (Martin et al., 2015; Simon et al., 2014). For example, in Benin, netting reduced diamond back moth, *Plutella xylostella* L., large cabbage white, *Pieris brassicae* L., cabbage webworm, *Hellula undalis* Fabricius, and cotton bollworm *Helicoverpa armigera* Hübner, populations on cabbage crops (Simon et al., 2014).

Here we found that *T. vaporariorum* and *M. sjostedti* populations on cowpea crops in the net house and net house + push-pull treatments were always lower than in the control. The net house could work as a screen which disturbs the visual cues used by these insects to locate their host plants. Gogo et al. (2014b) suggested that the bright white colour of the nets could act as a visual barrier. Like many insects, whiteflies and thrips orient their movements visually (Teulon et al., 1999; Antignus et al., 2001). Net houses reduce UV light penetration, which may in turn interfere with insects' vision and dispersion under the net house (Raviv et al., 2004; Ben-Yakir et al., 2014). By contrast, we found that *A. craccivora* infestations were higher in the net house and net house + push-pull treatments than in open fields, particularly in the dry period. The aphid outbreaks that occurred in the net house in the dry period suggest that the temperature and humidity conditions under this shelter were suitable for aphids. Net houses are known to provide conditions conducive to aphid outbreaks, as reported with regard to *Lipaphis erysimi* (Kaltenbach) infestations on cabbage crops in Benin (Simon et al., 2014), *Myzus persica* (Sulzer) and *Aphis gossypii* (Glover) infestations on sweet pepper crops (Singh et al., 2004) grown under netting. Insect nets can also protect aphid populations from parasitoids and predators (Martin et al., 2013). The absence of natural enemies to regulate aphid populations in the net house and net house + push-pull treatments may explain the high *A. craccivora* numbers observed in our study. Conversely, the presence of natural enemies in open field treatments could explain the reduced *A. craccivora* numbers in the dry period—many adult and larval coccinellid beetles were actually observed on plants in open field treatments. Spraying a soap/pepper biopesticide solution on the cowpea crops reduced aphid populations in our study, in line with the findings of previous studies (Poswal and Akpa, 1991; Pahla et al., 2014). Releasing predators or parasitoids inside the net house is another way to reduce *A. craccivora* infestations. Yang et al. (2014) reported that the release of *Harmonia axyridis* (Pallas) natural enemies in the greenhouse was an effective

way to reduce *Aphis gossypii* Glover populations on strawberry and cucumber. The severe aphid infestations we noted on cowpea flowers and pods could explain why cowpea yields were not as high as expected in the net house and net house + push-pull treatments. The best cowpea yields were nevertheless obtained in the net house treatment compared to other treatments despite the *A. craccivora* outbreak. The M66 cowpea cultivar used in this study may be tolerant to *A. craccivora*, since this cowpea genotype was an improved variety from the Kenya Agricultural Livestock Research Organization (KALRO) which is partly tolerant to aphid damage (Kenya Agricultural Research Institute (KARI), 2008; Kimutai, 2017).

We also hypothesised that the combined use of netting and the push-pull strategy (treatment 3) could help control small insect pests, particularly thrips on bean flowers. In our study, lemongrass (*C. citratus*) and Mexican marigold (*T. minuta*) were used to repel (push) insects from the cowpea crop and sticky traps were used as attractants (pull) to catch the flying insects. In laboratory assays, *M. sjostedti* was shown to be repelled by fresh cut lemongrass leaves but not by old cut leaves (Diabate et al., 2019a), but *M. sjostedti* numbers did not vary among any of the treatments in our study. Lemongrass leaves were cut once a week and spread between the rows to repel thrips. In addition, *T. minuta* likely continuously released repellent volatiles that prevented *M. sjostedti* from locating the cowpea plants. The repellent effect of volatiles could have been of short duration or the cut lemongrass leaves were perhaps not sufficiently abundant to repel *M. sjostedti*. The fact that there was a greater abundance of cowpea plants relative to lemongrass and Mexican marigold plants in the field could mean that there was an overwhelming presence of attractive volatiles, thus decreasing repellent effect of the companion plants. The efficacy of repellent volatiles could be enhanced in the field by sowing companion plants that produce greater quantities of repellent volatiles, or otherwise repellent essential oils could be provided through dispensers or injected in the irrigation system. We previously identified repellent citral volatiles in lemongrass and repellent blends of dihydrotagetone, (Z)-3-hexenyl acetate, limonene and (Z)- $\beta$ -ocimene volatiles in marigold (Diabate et al., 2019a). These compounds could be supplemented via dispensers to lengthen the repellence period in the field. Although the blue sticky traps were effective in catching *M. sjostedti* in the open field push-pull treatment, the number of thrips on cowpea plants in this treatment did not differ from that in the control. *M. sjostedti* was more attracted to cowpea flowers than to the blue traps. Frey et al. (1994) reported that the plant growth stage may have an impact on the efficacy of colour traps. For example, the number of *F. occidentalis* thrips caught on blue sticky traps in ornamental greenhouses was reported to be lower during flowering than during the vegetative stage (Frey et al., 1994).

By contrast, our findings showed that the *Empoasca* sp. population was lower in the open field push-pull treatment than in the control during the rainy period. Volatiles released by Mexican marigold and lemongrass could have been involved in the reduction of *Empoasca* sp. populations in the open field push-pull treatment. Calumpang et al. (2013) observed a reduction in green leafhopper, *Amrasca biguttula* (Ishida), populations when lemongrass was intercropped with eggplant. It was also reported that using Mexican marigold as an intercrop and maize as a border crop reduced *A. biguttula biguttula* leafhopper populations on eggplant crops in the field (Sujayanand et al., 2015). On the other hand, yellow sticky traps may contribute to reducing *Empoasca* sp. numbers. The attraction of sharp-nosed leafhoppers, *Scaphytopius magdalensis* (Provancher), to yellow traps was also demonstrated by Rodriguez-Saona et al. (2012). Combining visual and olfactory cues in a push-pull strategy could prove effective in controlling *Empoasca* sp. populations.

The open field push-pull treatment did not have an impact on insect populations except *Empoasca* sp. and therefore did not increase the pest control efficacy of the net house + push-pull treatment in the present study. Several authors have demonstrated the efficacy of non-host plant volatiles in reducing insect populations in the field (Parolin et al., 2012), whereas many other studies failed to reveal their efficacy in the same conditions (Held et al., 2003; Webster and Cardé, 2016). For example, intercropping French marigold, a non-host plant, with host potato plants did not reduce Colorado potato beetle, *Leptinotarsa decemlineata* (Say), populations but conversely it increased the incidence of *L. decemlineata* attacks (Moreau et al., 2006). In our study, we observed no such perverse effects of the push-pull strategy on any cowpea pests.

We observed that cowpea yields and insect populations varied with the period. Populations of large insect pests were higher in the rainy period than in the dry period. Conversely, small insect pests such as *M. sjostedti*, *T. vaporariorum* and *A. craccivora* were much lower in the rainy period than in the dry period. In Kenya, the heavy rainfall between March and June 2018 may have been responsible for the reduction in insect populations via egg destruction and partial larval mortality. Nyasani et al. (2013) reported that heavy rain destroyed western flower thrips, *F. occidentalis*, larvae in the field and hence reduced their population density. In our study, *A. craccivora* populations were reduced in the net house and net house + push-pull treatments in the rainy period compared to the dry period. The top of net house was flat and rain penetrated through the mesh. Rainfall is considered to regulate aphid population densities (Kaakeh and Dutcher, 1993). The lower aphid numbers in the net house and net house +

push-pull treatments suggest that the microclimate was unsuitable for these pests during the rainy period. Rainfall increased the relative humidity under the net, which could have a negative impact on *A. craccivora* populations by enhancing the environmental conditions for entomopathogenic fungi. Kataria and Kumar, (2017) reported that *A. craccivora* population patterns were negative correlated with minimum temperature, relative humidity and rainfall. In addition, *A. craccivora* propagate parthenogenetically in Africa (Irwin, 1980). This asexual reproduction enables aphids to multiply and spread rapidly under optimal conditions (Iluz, 2011). However, conditions might be different in the future and the lack of gene combinations could be a disadvantage to the offspring, thus curbing the growth populations of this pest. Cowpea pod yields in the open field treatments were very poor during the rainy period compared to the dry period. The reduction in pod yield may have been due to excessive rainfall in the rainy period, as already observed by Parwada (2016). In addition, cowpea plants in the open field treatments were heavily damaged by fungi and diseases in the rainy period. Edema et al. (1997) reported a higher incidence and severity of viral diseases, anthracnose and scab on cowpea crops in the rainy period than in the dry period. Fungi such as cercospora leaf spot may affect cowpea pod production and quality. Our study showed that net houses could help farmers to produce cowpea intensively. Despite the fact the rainy period was longer than usual during our study, cowpea production was always higher in the net house and net house + push-pull treatments than in the open field conditions. Net technology thus seems to be an efficient tool to help farmers grow vegetables in unpredictable weather conditions.

During both climate periods, all yield parameters evaluated were higher in the net house and net house + push-pull treatment than in the open field. The net house protected cowpeas against large insect pests or had no special effect on some of them, except in the rainy period when a *C. tomentosicollis* infestation was observed in the net house + push-pull treatment. Our findings did not enable us to explain why the *C. tomentosicollis* population was high in the net house + push-pull treatment in the rainy period, but the high density of Mexican marigold plants around the net house might have attracted or protected them. Further investigations are needed to elucidate this issue. Despite, the abundance of *Empoasca* sp. in the open fields during the rainy period, the net houses (used in treatments net house and net house + push-pull) prevented this pest from infesting the cowpea crops in the rainy period. The use of net houses in horticulture usually has a major impact on crop yields by reducing populations of large insect pests. *M. vitrata* and *C. tomentosicollis* can cause cowpea yield losses of up to 80% in Africa (Ekesi et al., 2002; OECD, 2015). The reduction of large insect pests could thus partially explain the increased crop yield in the net house and

net house + push-pull treatments, whereas the modified microclimate in the net house could boost plant development. Saidi et al. (2013) reported that the constant high temperature and soil moisture in net houses improved plant growth and yield. Moreover, we noticed that the quality of the grains produced in the net house and net house + push-pull treatments were higher. In Kenya, Gogo et al. (2014) also reported faster development, higher pod yield, and better quality of green beans, another leguminous plant that was grown under low netting covered tunnels as compared to uncovered plants.

In our study, the cowpea pod numbers were higher in the dry period than in the rainy period. Conditions in the dry period were better for cowpea pod production compared to the rainy period. Cowpea is an herbaceous warm-season annual and many authors have reported its good adaptation to high temperatures and drought (Ehlers and Hall, 1997; Timko et al., 2007). Moreover, the reduction of large insect pests in the dry period consequently reduced cowpea flower and pod damage due to low feeding activity. Conversely, small insect pests were more abundant in the dry period. Populations of *M. sjostedti* thrips and *T. vaporariorum* whiteflies on cowpea plants in the net house and net house + push-pull treatments were always lower than the open field populations, which increased very rapidly, particularly in the dry period. *M. sjostedti* outbreaks were linked to the flower emergence. *M. sjostedti* preferentially feeds on flowers, causing them to fall (Moritz et al., 2013). However, there was no difference in the total number of pods and flowers among the different treatments in the dry period. The high *A. craccivora* infestation of cowpea vegetative parts, flowers and pods in the net house and net house + push-pull treatments did not decrease yields in the dry period. Small insect pests therefore seem to have a limited impact on cowpea production, contrary to large insect pests.

## 5. Conclusion

The findings of this study showed that net house and net house + push-pull treatments considerably reduced pest infestations on cowpeas and that the damage was mainly incurred by large insect pests such as *Empoasca* sp., *M. vitrata* and *C. tomentosicollis* along with smaller insect pests such as *M. sjostedti* and *T. vaporariorum*. We also noted that net house grown cowpeas could be subject to high *A. craccivora* infestation. Aphid outbreaks seemed to have a limited impact on cowpea yields, so populations of this pest could be reduced by biopesticide treatment. The net house + push-pull treatment did not improve *T. vaporariorum*, *A. craccivora* and *M. sjostedti* control. However, the open field push-pull treatment reduced *Empoasca* sp. infestations during the rainy period. The study also showed



that cowpea production was much higher in the net house and net house + push-pull treatments than in the open field. In addition, cowpea pod yields were higher in all treatments during the dry period than in the rainy period. The large insect pest populations were low during the dry period and high in the rainy period, contrary to small insect pest populations. The study showed that cowpea production was impacted by the period and large insect pest infestations. This study showed that netting technology provides an effective alternative to chemical pesticide treatments for vegetable growing in the tropics regardless of the season. We showed that netting technology could be slightly improved to control sucking insect pests using the push-pull strategy. Cowpea yields were highly affected by heavy rains in both open field treatments but not in the net house and net house + push-pull treatments. Future research should be focused on investigating the effect of microclimate and insect pests on cowpea yield in net houses. In Mwea (Kenya), the rainy season is not the best period to grow cowpeas in the open fields despite the low insect pest pressure. But we found that the cowpea yield and quality in the net house treatment was still high, thus confirming the need for netting to produce off-season vegetable crops when prices may be higher due to the low market supply. This study confirmed that netting is an efficient tool for protecting vegetable crops against insect pests and producing higher yields of safe vegetables.

**Acknowledgments:** We are grateful to the German Federal Ministry for Economic Cooperation and Development (BMZ), the French Agricultural Research Centre for International Development (CIRAD) and Kenya Agricultural and Livestock Research Organization (KALRO) financial support. We thank UK Aid, the Swedish International Development Cooperation Agency (SIDA), the Swiss Agency for Development and Cooperation (SDC) and the Kenyan Government for their core support for *icipe* research. The first author was supported by a German Academic Exchange Service (DAAD) In-Region Postgraduate Scholarship through *icipe*.

## References

- Ahmed, M.T., Miah, M.R.U., Amin, M.R., Hossain, M.M., Suh, S.J., Kwon, Y.J., 2019. Plant material as an alternative tool for management of aphid in country bean field. *Int. J. Pest Manag.* 65, 171–176. <https://doi.org/10.1080/09670874.2018.1494864>
- Alexandratos, N., Bruinsma, J., 2012. World agriculture towards 2030/2050: the 2012 revision, ESA Working Paper No. 12-03 (FAO, Rome, 2012). [https://doi.org/10.1016/S0264-8377\(03\)00047-4](https://doi.org/10.1016/S0264-8377(03)00047-4)
- Annan, I.B., Saxena, K.N., Schaefer, G.A., Tingey, W.M., 1994. Effects of infestation by cowpea aphid (Homoptera: Aphididae) on different growth stages of resistant and susceptible cowpea cultivars. *Int. J. Trop. Insect Sci.* 15, 401–410. <https://doi.org/10.1017/s1742758400015733>
- Antignus, Y., Nestel, D., Cohen, S., Lapidot, M., 2001. Ultraviolet-deficient greenhouse environment affects whitefly attraction and flight-behavior. *Environ. Entomol.* 30, 394–399. <https://doi.org/10.1603/0046-225x-30.2.394>
- Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. lme4: Linear mixed-effects models using Eigen and S4. R package version 1.1-8. Retrieved from <http://CRAN.R-project.org/package=lme4>
- Ben-Yakir, D., Antignus, Y., Offir, Y., Shahak, Y., 2014. Photosensitive nets and screens can reduce insect pests and diseases in agricultural crops. *Acta Hort.* 1015, 95–102. <https://doi.org/10.17660/ActaHortic.2014.1015.10>
- Borowiak-Sobkowiak, B., Durak, R., & Wilkaniec, B. (2017). Morphology, biology and behavioral aspects of *Aphis craccivora* (Hemiptera:Aphididae) on *Robinia pseudoacacia*. *Acta Sci Pol-Hortoru.* 16, 39–49.
- Calumpang, S.M.F., Bayot, R.G., Vargas, D.G., Ebuenga, D.M., Gonzales, P.G., 2013. Impact of intercropping lemon grass (*Cymbopogon citratus* Stapf.) on infestation of Eggplant Fruit and Shoot Borer (*Leucinodes*

443        *orbonalis Guenee*) in Eggplant (*Solanum melongena* L.). Siliman J. 54, 114–130

444        Cook, S.M., Khan, Z.R., Pickett, J.A., 2006. The use of push-pull strategies in integrated pest management. Annu.  
445        Rev. Entomol. 52, 375–400. <https://doi.org/10.1146/annurev.ento.52.110405.091407>

446        de Bon, H., Huat, J., Parrot, L., Sinzogan, A., Martin, T., Malézieux, E., Vayssières, J.F., 2014. Pesticide risks from  
447        fruit and vegetable pest management by small farmers in sub-Saharan Africa. a review. Agron. Sustain. Dev.  
448        34, 723–736. <https://doi.org/10.1007/s13593-014-0216-7>

449        Diabate, S., Deletre, E., Murungi, L.K., Fiaboe, K.K.M., Subramanian, S., Wesonga, J., Martin, T., 2019b.  
450        Behavioural responses of bean flower thrips (*Megalurothrips sjostedti*) to vegetative and floral volatiles from  
451        different cowpea cultivars. Chemoecology 29, 73–88. <https://doi.org/10.1007/s00049-019-00278-0>

452        Diabate, S., Martin, T., Murungi, L.K., Fiaboe, K.K.M., Subramaniana, S., Wesonga, J., Deletre, E., 2019a.  
453        Repellent activity of *Cymbopogon citratus* and *Tagetes minuta* and their specific volatiles against  
454        *Megalurothrips sjostedti*. J. Appl. Entomol. 00, 1–12. <https://doi.org/https://doi.org/10.1111/jen.12651>

455        Edema, R., Adipala, E., Florini, D.A., 1997. Influence of season and cropping system on occurrence of cowpea  
456        diseases in Uganda. Plant Dis. 81, 465–468. <https://doi.org/10.1094/PDIS.1997.81.5.465>

457        Ehlers, J.D., Hall, A.E., 1997. Cowpea (*Vigna unguiculata* L. Walp.). F. Crop. Res. 53, 187–204.  
458        [https://doi.org/10.1016/S0378-4290\(97\)00031-2](https://doi.org/10.1016/S0378-4290(97)00031-2)

459        Eigenbrode, S.D., Birch, A.N.E., Lindzey, S., Meadow, R., Snyder, W.E., 2016. A mechanistic framework to  
460        improve understanding and applications of push-pull systems in pest management. J. Appl. Ecol. 53, 202–212.  
461        <https://doi.org/10.1111/1365-2664.12556>

462        Ekesi, S., Adamu, R.S., Maniania, N.K., 2002. Ovicidal activity of entomopathogenic hyphomycetes to the legume  
463        pod borer , *Maruca vitrata* and the pod sucking bug , *Clavigralla tomentosicollis*. Crop Prot. 21, 589–595.

464        Foba, C.N., Salifu, D., Lagat, Z.O., Gitonga, L.M., Akutse, K.S., Fiaboe, K.K.M., 2015. Species composition,  
465        distribution, and seasonal abundance of *Liriomyza* leafminers (Diptera: Agromyzidae) under different  
466        vegetable production systems and agroecological zones in Kenya. Environ. Entomol. 44, 223–232.  
467        <https://doi.org/10.1093/ee/nvu065>

468 Frey, J.E., Cortada, R.V., Helbling, H., 1994. The potential of flower odours for use in population monitoring of  
 469 western flower thrips *Frankliniella occidentalis* Perg. (Thysanoptera: Thripidae). *Biocontrol Sci Technol.* 4,  
 470 177–186.

471 Gogo, E.O., Saidi, M., Itulya, F.M., Martin, T., Ngouajio, M., 2012. Microclimate modification using eco-friendly  
 472 nets for high-quality tomato transplant production by small-scale farmers in East Africa. *Horttechnology* 22,  
 473 292–298.

474 Gogo, E., Saidi, M., Itulya, F., Martin, T., Ngouajio, M., 2014a. Eco-friendly nets and floating row covers reduce  
 475 pest infestation and improve tomato (*Solanum lycopersicon* L.) yields for smallholder farmers in Kenya.  
 476 *Agronomy* 4, 1–12. <https://doi.org/10.3390/agronomy4010001>

477 Gogo, E., Saidi, M., Ochieng, J.M., Martin, T., Baird, V., Ngouajio, M., 2014b. Microclimate modification and  
 478 insect pest exclusion using agronet improve pod yield and quality of French bean. *HortScience* 49, 1298–  
 479 1304.

480 Hassan, R.M., 1998. Maize technology development and transfer. A GIS application for research planning in  
 481 Kenya., CAB International, Walford, UK. CIMMYT and Kenya Agricultural Research Institute, pp 227

482 HCDA, 2014. Horticulture Validated report- Ministry of Agriculture & Horticultural Crops Development Authority.

483 Held, D.W., Gonsiska, P., Potter, D.A., 2003. Evaluating companion planting and non-host masking odors for  
 484 protecting roses from the Japanese beetle (Coleoptera: Scarabaeidae). *J. Econ. Entomol.* 96, 81–87.  
 485 <https://doi.org/10.1093/jee/96.1.81>

486 Iluz, D., 2011. The Plant–Aphid Universe, in: Dubinsky, J.S. and Z. (Ed.), *All Flesh Is Grass. Plant-Animal*  
 487 *Interrelationships.* pp. 91–118. <https://doi.org/10.1093/aob/mcr214>

488 Irwin, M.E., 1980. Sampling Methods in Soybean Entomology, in: Kogan, M., Herzog, D.C. (Eds.), *Sampling*  
 489 *Methods in Soybean Entomology, Springer Series in Experimental Entomology.* Springer New York, New  
 490 York, NY, pp. 239–259. <https://doi.org/10.1007/978-1-4612-9998-1>

491 Kaakeh, W., Dutcher, J., 1993. Effect of rainfall on population abundance of aphids (Homoptera : Aphididae ) on  
 492 Pecan. *J. Entomol. Sci.* 28, 283–286. <https://doi.org/10.18474/0749-8004-28.3.283>

493 Kataria, R., Kumar, D., 2017. Population dynamics of *Aphis craccivora* (Koch) and its natural enemies on bean crop  
 494 in relation to weather parameters in Vadodara, Gujarat, India. Legum. Res. 40, 571–579.  
 495 <https://doi.org/10.18805/lr.v0iOF.10282>

496 Kenya Agricultural Research Institute (KARI), 2008. Growing cow peas in dry areas.  
 497 [http://www.kalro.org/fileadmin/publications/brochuresII/Growing\\_cow\\_peas\\_in\\_dry\\_areas.pdf](http://www.kalro.org/fileadmin/publications/brochuresII/Growing_cow_peas_in_dry_areas.pdf). (accessed 16  
 498 April 2018)

499 Khan, Z.R., Ampong-Nyarko, K., Chiliswa, P., Hassanali, A., Kimani, S., Lwande, W., Overholt, W.A., Pickett,  
 500 J.A., Smart, L.E., Wadhams, L.J., Woodcock, C.M., 1997. Intercropping increases parasitism of pests. Nature  
 501 388, 631–632. <https://doi.org/10.1038/41681>

502 Kimutai, W.J., 2017. Effect of selected bradyrhizobia and nutrients on cowpea biomass, biological nitrogen and  
 503 yield in Kilifi and Mbeere regions of Kenya. Masters Thesis, Kenyatta University (Kenya). p 69

504 Koona, P., Osisanya, E.O., Jackai, L., Tamo, M., Markham, R.H., 2002. Resistance in accessions of cowpea to the  
 505 coreid pod-bug *Clavigralla tomentosicollis* ( Hemiptera : Coreidae ). J. Econ. Entomol. 95, 1281–1288.

506 Martin, T., Assogba-Komlan, F., Houndete, T., Hougard, J.M., Chandre, A.F., 2006. Efficacy of mosquito netting  
 507 for sustainable small holders' cabbage production in Africa. J. Econ. Entomol 99, 450–454.  
 508 <https://doi.org/10.1603/0022-0493-99.2.450>

509 Martin, T., Palix, R., Kamal, A., Delétré, E., Bonafos, R., Simon, S., Ngouajio, M., 2013. A repellent net as a new  
 510 technology to protect cabbage crops. J. Econ. Entomol. 106, 1699–1706. <https://doi.org/10.1603/EC13004>

511 Martin, T., Simon, S., Parrot, L., Komlan, F.A., Vidogbena, F., Adegbidi, A., Baird, V., Saidi, M., Kasina, M.,  
 512 Wasilwa, L.A., Subramanian, S., Ngouajio, M., 2015. Eco-friendly nets to improve vegetable production and  
 513 quality in sub-Saharan Africa. Acta Hort. 1105, 221–227. <https://doi.org/10.17660/ActaHortic.2015.1105.31>

514 Morallo-Rejesus, B., Decena, A., 1982. The activity, isolation, purification and identification of the insecticidal  
 515 principles from Tagetes. Philipp. J. Crop Sci. 7, 31–36

516 Moreau, T.L., Warman, P.R., Hoyle, J., 2006. An evaluation of companion planting and botanical extracts as  
 517 alternative pest controls for the colorado potato beetle. Biol. Agric. Hortic. 23, 351–370.

518 <https://doi.org/10.1080/01448765.2006.9755336>

519 Moritz, G., Brandt, S., Triapitsyn, S., Subramanian, S., 2013. Pest thrips in East Africa: identification and  
520 information tools. QAAFI Biological Information Technology (QBIT), The University of Queensland,  
521 Brisbane. <https://doi.org/ISBN 978-1-74272-067-8>

522 Muniu, F.K., 2017. Characterization and evaluation of local cowpea accessions and their response to organic and  
523 inorganic nitrogen fertilizers in Coastal Kenya. B.S.c Thesis. University of Nairobi. pp 72

524 Musebe, R., Dorward, P., Karanja, D., 2005. Socio-economic report for project ( R8296 / ZA0568 ), promotion of  
525 sustainable approaches for the management of root-knot nematodes on vegetables in Kenya.

526 Muvea, A.M., Waiganjo, M.M., Kutima, H.L., Osiemo, Z., Nyasani, J.O., Subramanian, S., 2014. Attraction of pest  
527 thrips (Thysanoptera: Thripidae) infesting French beans to coloured sticky traps with Lurem-TR and its utility  
528 for monitoring thrips populations. Int. J. Trop. Insect Sci. 34, 197–206.  
529 <https://doi.org/10.1017/S174275841400040X>

530 Mweke, A., Ulrichs, C., Nana, P., Akutse, K.S., Fiaboe, K.K.M., Maniania, N.K., Ekesi, S., 2018. Evaluation of the  
531 Entomopathogenic Fungi *Metarhizium anisopliae*, *Beauveria bassiana* and *Isaria sp.* for the Management of  
532 *Aphis craccivora* (Hemiptera: Aphididae). J. Econ. Entomol. 111, 1587–1594.  
533 <https://doi.org/10.1093/jee/toy135>

534 Nordey, T., Basset-Mens, C., De Bon, H., Martin, T., Déletré, E., Simon, S., Parrot, L., Desprez, H., Huat, J., Biard,  
535 Y., Dubois, T., Malézieux, E., 2017. Protected cultivation of vegetable crops in sub-Saharan Africa: limits and  
536 prospects for smallholders. a review. Agron. Sustain. Dev. 37, 53. <https://doi.org/10.1007/s13593-017-0460-8>

537 Nyasani, J.O., Meyhöfer, R., Subramanian, S., Poehling, H.M., 2013. Seasonal abundance of western flower thrips  
538 and its natural enemies in different French bean agroecosystems in Kenya. J. Pest Sci. (2004). 86, 515–523.  
539 <https://doi.org/10.1007/s10340-013-0491-0>

540 Obopile, M., 2006. Economic threshold and injury levels for control of cowpea aphid, *Aphis crassivora* Linnaeus  
541 (Homoptera: Aphididae), on cowpea. Afr. Plant Prot. 12, 111–115. <https://hdl.handle.net/10520/EJC87791>

542 Obopile, M., Ositile, B., 2010. Life table and population parameters of cowpea aphid, *Aphis craccivora* Koch  
 543 (Homoptera: Aphididae) on five cowpea *Vigna unguiculata* (L. Walp.) varieties. J. Pest Sci. (2004). 83, 9–14.  
 544 <https://doi.org/10.1007/s10340-009-0262-0>

545 OECD, 2015. Consensus document on the biology of cowpea (*Vigna unguiculata* (L.) Walp.), Series on  
 546 Harmonisation of Regulatory Oversight in Biotechnology No. 60. [https://doi.org/ENV/JM/MONO\(2015\)48](https://doi.org/ENV/JM/MONO(2015)48)

547 OECD, FAO, 2016. Agriculture in Sub-Saharan Africa: Prospects and challenges for the next decade, in: OECD-  
 548 FAO Agricultural Outlook 2016-2025.

549 Ofuya, T.I., Okuku, I.E., 1994. Insecticidal effect of some plant extracts on the cowpea aphid *Aphis craccivora*  
 550 Koch (Homoptera: Aphididae). Anzeiger für Schädlingskd. Pflanzenschutz Umweltschutz 67, 127–129.  
 551 <https://doi.org/10.1007/BF01909033>

552 Pahla, I., Moyo, M., Muzemu, S., Muziri, T., 2014. Evaluating the effectiveness of botanical sprays in controlling  
 553 Aphids (*Brevicoryne brassicae*) on rape (*Brassica napus* L). Int. J. Agron. Agric. Res. 5, 1–6.

554 Painter, R.H., 1951. Insect resistance in crop plants, vol. 72, No. 6, 481. LWW

555 Parolin, P., Bresch, C., Desneux, N., Brun, R., Bout, A., Boll, R., Poncet, C., 2012. Secondary plants used in  
 556 biological control: a review. Int. J. Pest Manag. 58, 91–100. <https://doi.org/10.1080/09670874.2012.659229>

557 Parwada, C., 2016. Evaluation of seasonal rainfall changes on the growth performance of *Vigna unguiculata*  
 558 (cowpeas) in Zimbabwe. Acad. J. Agric. Res. 4, 067–071. <https://doi.org/10.15413/ajar.2015.0188>

559 Popelka, J.C., Gollasch, S., Moore, A., Molvig, L., Higgins, T.J.V., 2006. Genetic transformation of cowpea (*Vigna*  
 560 *unguiculata* L.) and stable transmission of the transgenes to progeny. Plant Cell Rep. 25, 304–312.  
 561 <https://doi.org/10.1007/s00299-005-0053-x>

562 Poswal, M.A.T., Akpa, A.D., 1991. Current trends in the use of traditional and organic methods for the control of  
 563 crop pests and diseases in Nigeria. Trop. Pest Manag. 37, 329–333.  
 564 <https://doi.org/10.1080/09670879109371609>

565 Raviv, M., Antignus, Y., Yishay, R., 2004. Invited review UV radiation effects on pathogens and insect pests of

566 greenhouse-grown crops. Photochem. Photobiol. 79, 219–226.

567 Rodriguez-Saona, C.R., Byers, J.A., Schiffhauer, D., 2012. Effect of trap color and height on captures of blunt-  
568 nosed and sharp-nosed leafhoppers (Hemiptera: Cicadellidae) and non-target arthropods in cranberry bogs.  
569 Crop Prot. 40, 132-144.

570 Saidi, M., Gogo, E.O., Itulya, F.M., Martin, T., Ngouajio, M., 2013. Microclimate modification using eco-friendly  
571 nets and floating row covers improves tomato (*Lycopersicon esculentum* ) yield and quality for small holder  
572 farmers in East Africa. Agric. Sci. 4, 577–584.

573 Saroch, V.K., 2000. biology and management of bean aphid, *Aphis craccivora* (Koch) on mung crop. M.Sc thesis.  
574 Sher-E-Kashmir, University of Agricultural Sciences and Technology.pp 55

575 Simon, S., Komlan, F.A., Adjaïto, L., Mensah, a., Coffi, H.K., Ngouajio, M., Martin, T., 2014. Efficacy of insect  
576 nets for cabbage production and pest management depending on the net removal frequency and microclimate.  
577 Int. J. Pest Manag. 60, 208–216. <https://doi.org/10.1080/09670874.2014.956844>

578 Singh, D., Kaur, S., Dhillon, T.S., 2004. Protected cultivation of sweet pepper hybrids under net-house in indian  
579 conditions. Acta Hortic. 659, 515–521.

580 Singh, S.R., Jackai, L.E.N., 1988. The legume pod-borer, *Maruca Testulalis* ( Geyer ): past , present and future  
581 research. Insect Sci. its Appl. 9, 1–5.

582 Smaili, M.C., El Ghadraoui, L., Gaboun, F., Benkirane, R., Blenzar, A., 2014. Impact of some alternative methods  
583 to chemical control in controlling aphids (Hemiptera: Sternorrhyncha) and their side effects on natural  
584 enemies on young Moroccan citrus groves. Phytoparasitica 42, 421–436. [https://doi.org/10.1007/s12600-013-](https://doi.org/10.1007/s12600-013-0379-9)  
585 0379-9

586 Stoetzel, M.B., Miller Gary, L. (2001). Aerial feeding aphids of corn in the united states with reference to the root-  
587 feeding *Aphis maidiradicis* (Homoptera: Aphididae). Florida Entomol. 84, 83–98.  
588 <https://doi.org/10.2307/3496667>



589 Sujayanand, G.K., Sharma, R.K., Shankarganesh, K., Saha, S., Tomar, R.S., 2015. Crop diversification for  
590 sustainable insect pest management in Eggplant (Solanales: Solanaceae). *Florida Entomol.* 98, 305–314.  
591 <https://doi.org/10.1653/024.098.0149>

592 Teulon, D.A.J., Hollister, B., Butler, R.C., Cameron, E.A., 1999. Colour and odour responses of flying western  
593 flower thrips: Wind tunnel and greenhouse experiments. *Entomol. Exp. Appl.* 93, 9–19.  
594 <https://doi.org/10.1023/A:1003893905051>

595 Togola, A., Boukar, O., Belko, N., Chamarthi, S.K., Fatokun, C., Tamo, M., Oigiangbe, N., 2017. Host plant  
596 resistance to insect pests of cowpea (*Vigna unguiculata* L. Walp.): achievements and future prospects.  
597 *Euphytica* 213, 1–16. <https://doi.org/10.1007/s10681-017-2030-1>

598 Timko, M.P., Ehlers, J.D., Roberts, P. A., 2007. Cowpea, in: *Genome Mapping and Molecular Breeding in Plants:*  
599 *Pulses, Sugar and Tuber Crops.* pp. 49–67.

600 Webb, S.E., Kok-Yokomi, M.L., Voegtlin, D.J., 1994. Effect of trap color on species composition of alate aphids  
601 (Homoptera : Aphididae ) caught over watermelon plants. *Florida Entomol.* 77, 146–154.  
602 <https://doi.org/10.2307/3495881>

603 Webster, B., Cardé, R.T., 2016. Use of habitat odour by host-seeking insects. *Biol. Rev.* 92, 1241–1249.  
604 <https://doi.org/10.1111/brev.12281>

605 Yang, N.W., Zang, L.S., Wang, S., Guo, J.Y., Xu, H.X., Zhang, F., Wan, F.H., 2014. Biological pest management  
606 by predators and parasitoids in the greenhouse vegetables in China. *Biol. Control* 68, 92–102.  
607 <https://doi.org/10.1016/j.biocontrol.2013.06.012>

608

### Figure captions


**Figure 1:** Schematic representation of the experimental field set up in a 4x4 Latin square design with a total of 16 plots and randomized rows and columns. 300 plants were grown per plot and plots were separated by 2-m bare soil buffer strips. Four strips of sticky tape (15 cm x 10 m) per push-pull plot were placed at 1 m intervals between the rows.

**Figure 2:** Effects of treatments on adult *Trialeurodes vaporariorum* (a), adult and larval *Aphis craccivora* (b), adult and larval *Megalurothrips sjostedti* (c) population fluctuations in the dry period at KALRO-Mwea (Kenya).




Two applications of a mixed pepper/soap solution in all treatments on 17 and 24 November 2017


**Figure 3:** Effects of treatments on adult *Trialeurodes vaporariorum*, adult and larval *Aphis craccivora*, adult and larval *Megalurothrips sjostedti* population fluctuations in the rainy period at KALRO-Mwea (Kenya).






Plots



A push-pull treatment plot





**Table 1:** Mean ( $\pm$  SE) and abundance of insects observed per plant among the treatments in the dry period<sup>1</sup> and rainy period<sup>2</sup> at KALRO- Mwea (Kenya).

|                    |                                    | Dry period         |                   |                       |                      |       |           | Rainy period        |                    |                    |                   |       |           |
|--------------------|------------------------------------|--------------------|-------------------|-----------------------|----------------------|-------|-----------|---------------------|--------------------|--------------------|-------------------|-------|-----------|
|                    | Insect pests                       | Control            | Treatment 1       | Treatment 2           | Treatment 3          | F     | P-value   | Control             | Treatment 1        | Treatment 2        | Treatment 3       | F     | P-value   |
| <i>small pests</i> | <i>Megalurothrips sjostedti</i>    | 51.0 $\pm$ 12.6 a  | 42.6 $\pm$ 11.2 a | 16.3 $\pm$ 5.8 a      | 9.9 $\pm$ 3.7 a      | 2.54  | 0.10      | 31.4 $\pm$ 7.2 A    | 31.3 $\pm$ 7.1 A   | 18.8 $\pm$ 5.1 A   | 7.1 $\pm$ 1.5 A   | 1.67  | 0.22      |
|                    | <i>Aphis craccivora</i>            | 24.6 $\pm$ 21.5 a* | 5.10 $\pm$ 2.8 a* | 406.5 $\pm$ 213.08 b* | 518.8 $\pm$ 191.5 b* | 17.33 | P < 0.001 | 101.3 $\pm$ 39.0 A* | 43.0 $\pm$ 16.5 A* | 67.7 $\pm$ 32.9 A* | 12.5 $\pm$ 6.0 A* | 1.20  | 0.34      |
|                    | <i>Trialeurodes vaporariorum</i>   | 14.0 $\pm$ 4.2 a*  | 5.6 $\pm$ 1.7 a*  | 3.8 $\pm$ 2.7 b       | 0.1 $\pm$ 0.1b       | 18.54 | P < 0.001 | 3.7 $\pm$ 1.2 A*    | 1.0 $\pm$ 0.3 AB*  | 0.3 $\pm$ 0.1 B    | 0.1 $\pm$ 0.1 B   | 9.37  | 0.0018    |
| <i>large pests</i> | <i>Maruca vitrata</i>              | 0.3 $\pm$ 0.1 a*   | 0.2 $\pm$ 0.1 ab  | 0 b                   | 0.0 $\pm$ 0.0 ab     | 4.43  | 0.025     | 0 A*                | 0.0 $\pm$ 0.0 A    | 0A                 | 0A                | 2.45  | 0.11      |
|                    | <i>Empoasca</i> sp.                | 0.4 $\pm$ 0.1 a*   | 0.3 $\pm$ 0.1 a*  | 0 b                   | 0.0 $\pm$ 0.0 b      | 8.77  | 0.002     | 8.0 $\pm$ 1.6 A*    | 3.1 $\pm$ 0.7 B*   | 0.1 $\pm$ 0.1 C    | 0 $\pm$ 0 C       | 35.62 | P < 0.001 |
|                    | <i>Clavigralla tomentosicollis</i> | 7.5 $\pm$ 3.0 a    | 5.6 $\pm$ 2.5 a   | 0.2 $\pm$ 0.2 b*      | 0 b*                 | 9.41  | 0.001     | 2.7 $\pm$ 0.8 AB    | 4.2 $\pm$ 1.2 AB   | 2.4 $\pm$ 1.3 A*   | 10.0 $\pm$ 2.5 B* | 3.68  | 0.043     |

<sup>1</sup> From 3 October 2017 to 9 January 2018; <sup>2</sup> From 30 January 2018 to 15 May 2018, SE = Standard error, numerator and denominator degrees of freedom (df = 3,

12). Treatment 1= Push-pull strategy; Treatment 2 = Net house; Treatment 3 = Combined net house + push-pull strategy. Small letters correspond to the comparison of treatments in the dry period and capital letters refer to the comparison of treatments in the rainy period using a linear mixed model (F, p-value). If significant, means were separated by least squares means (lsmeans) adjusted using the Tukey method. \*indicates significant differences between the same treatment in the two periods according to the Student's t test. The same letter in the same row means not different.

**Table 2:** Mean ( $\pm$  SE) number of insects: *A. craccivora*, *M. sjostedti* and *T. vaporariorum* caught on the coloured sticky traps during the two periods. Dry period from 3 October 2017 to 9 January 2018. Rainy period from 30 January 2018 to 15 May 2018.

| Insects                |                | Blue trap          | Yellow trap          |
|------------------------|----------------|--------------------|----------------------|
| <i>T. vaporariorum</i> | Treatment 1    | 10.6 $\pm$ 12.8a   | 2315.3 $\pm$ 1771.5a |
|                        | Treatment 3    | 1.0 $\pm$ 2.0a     | 91.2 $\pm$ 100.0b    |
|                        | <i>t-test</i>  | 1.438              | 2.583                |
|                        | <i>P value</i> | 0.172              | 0.021                |
| <i>M. sjostedti</i>    | Treatment 1    | 265.4 $\pm$ 131.7a | 4.4 $\pm$ 4.5a       |
|                        | Treatment 3    | 27.0 $\pm$ 14.2b   | 2.1 $\pm$ 1.4a       |
|                        | <i>t-test</i>  | 3.487              | 0.241                |
|                        | <i>P value</i> | 0.003              | 0.812                |
| <i>A. craccivora</i>   | Treatment 1    | 0a                 | 64.6 $\pm$ 99.9a     |
|                        | Treatment 3    | 0.25 $\pm$ 0.32a   | 93.9 $\pm$ 88.6a     |
|                        | <i>t-test</i>  | -1.527             | -0.993               |
|                        | <i>P value</i> | 0.148              | 0.337                |

SE = Standard error, df= 14. Treatment 1 = Push-pull strategy; Treatment 3 = Combined net house + push-pull strategy. Small letters refer to the comparison of two treatments per insect in each column using the Student's t test at  $p < 0.05$ . The same letter means not significantly different

**Table 3:** Mean ( $\pm$  SE) flowers per plant, the total and weight of harvested pods, marketable weight of harvested pods and number of grains per plot among the treatments in the dry period<sup>1</sup> and rainy period<sup>2</sup> at KALRO- Mwea (Kenya).

|                           | Dry period          |                     |                     |                     |      |         | Rainy period     |                  |                    |                    |       |           |
|---------------------------|---------------------|---------------------|---------------------|---------------------|------|---------|------------------|------------------|--------------------|--------------------|-------|-----------|
|                           | Control             | Treatment 1         | Treatment 2         | Treatment 3         | F    | P-value | Control          | Treatment 1      | Treatment 2        | Treatment 3        | F     | P-value   |
| Flowers                   | 2.9 $\pm$ 0.7 a     | 5.8 $\pm$ 2.0 a     | 3.7 $\pm$ 0.9 a     | 3.2 $\pm$ 0.9 a     | 0.13 | 0.93    | 1.7 $\pm$ 0.4 A  | 2.5 $\pm$ 0.5 A  | 3.3 $\pm$ 0.7 A    | 3.1 $\pm$ 0.5 A    | 1.02  | 0.41      |
| Total harvested pods      | 110.3 $\pm$ 30.3 a* | 120.8 $\pm$ 31.7 a* | 181.60 $\pm$ 49.6 a | 149.4 $\pm$ 39.1 a  | 0.18 | 0.90    | 0.5 $\pm$ 0.5 A* | 2.2 $\pm$ 1.7 A* | 105.3 $\pm$ 26.3 B | 73.9 $\pm$ 17.8 B  | 11.63 | P < 0.001 |
| Total weight of pods (g)  | 282.4 $\pm$ 78.4 a* | 288.9 $\pm$ 73.7 a* | 478.2 $\pm$ 131.7 a | 403.9 $\pm$ 101.6 a | 0.16 | 0.91    | 1.3 $\pm$ 1.3 A* | 3.0 $\pm$ 2.7 A* | 276.6 $\pm$ 70.4 B | 187.2 $\pm$ 44.7 B | 11.84 | P < 0.001 |
| Marketable harvested pods | 70.6 $\pm$ 19.1 a*  | 75.4 $\pm$ 20.4 a*  | 156.8 $\pm$ 43.4 a  | 124.3 $\pm$ 32.1 a  | 0.31 | 0.81    | 0 A*             | 1.0 $\pm$ 0.7 A* | 104.9 $\pm$ 26.0 B | 69.4 $\pm$ 16.4 B  | 13.32 | P < 0.001 |
| Marketable weight pod (g) | 198.2 $\pm$ 53.6 a* | 201.1 $\pm$ 51.4 a* | 431.1 $\pm$ 119.9 a | 348.9 $\pm$ 86.2 a  | 0.29 | 0.83    | 0 A*             | 0.3 $\pm$ 0.3 A* | 272.3 $\pm$ 68.7 B | 181.7 $\pm$ 43.5 B | 13.62 | P < 0.001 |
| Grains                    |                     |                     |                     |                     |      |         | 0.7 $\pm$ 0.7 A  | 0.8 $\pm$ 0.7 A  | 227.5 $\pm$ 57.3 B | 136.2 $\pm$ 33.1 B | 12.28 | P < 0.001 |

<sup>1</sup> From 3 October 2017 to 9 January 2018; <sup>2</sup>From 30 January 2018 to 15 May 2018, SE = Standard error, numerator and denominator degrees of freedom (df = 3,

12). Treatment 1 = Push-pull strategy; Treatment 2 = Net house; Treatment 3 = Combined net house + push-pull strategy. Small letters refer to the comparison of treatments in the dry period and capital letters to comparison of treatments in the rainy period using a linear mixed model (F, p-value). If significant, means were separated by least squares means (lsmeans) adjusted using the Tukey method. \*indicates significant differences between the same treatment in the two periods with the Student's t test. The same letter in the same row means not different