



HAL
open science

Protecting Biodiversity (in All Its Complexity): New Models and Methods

Laura J. Pollock, Louise M.J. O'connor, Karel Mokany, Dan F. Rosauer,
Matthew V. Talluto, Wilfried Thuiller

► **To cite this version:**

Laura J. Pollock, Louise M.J. O'connor, Karel Mokany, Dan F. Rosauer, Matthew V. Talluto, et al..
Protecting Biodiversity (in All Its Complexity): New Models and Methods. Trends in Ecology &
Evolution, 2020, 35, pp.1119 - 1128. 10.1016/j.tree.2020.08.015 . hal-03493786

HAL Id: hal-03493786

<https://hal.science/hal-03493786>

Submitted on 21 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

1 **Protecting biodiversity (in all its complexity): new models and methods**

2

3 Laura J. Pollock^{1,2*}, Louise M. J. O'Connor², Karel Mokany³, Dan F. Rosauer⁴, Matthew V.

4 Talluto^{5,6} and Wilfried Thuiller^{2*}

5

6

7 ¹Department of Biology, McGill University, 1205 Dr. Penfield, Montréal, Québec H3A 1B1,

8 Canada.

9 ²Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LECA, Laboratoire d'Écologie

10 Alpine, F-38000 Grenoble, France

11 ³CSIRO, PO Box 1700, Canberra, Australian Capital Territory 2601, Australia

12 ⁴Research School of Biology, Australian National University, Acton, Australian Capital

13 Territory, 2601, Australia.

14 ⁵Department of Ecohydrology, Leibniz Institute for Freshwater Ecology and Inland Fisheries

15 Müggelseedamm 310, 12587 Berlin, Germany.

16 ⁶Department of Ecology, University of Innsbruck, Innrain 52, AT-6020 Innsbruck, Austria

17

18 *Corresponding Author: L. Pollock, lajosy@gmail.com, +33 6 1041 3058, @lajosy0

19

20 **Keywords**

21 Conservation prioritization, multi-trophic diversity, species distribution models, phylogenetic

22 diversity, functional diversity

23

24

25 **Abstract**

26 We are facing a biodiversity crisis at the same time as we are acquiring an unprecedented view of the
27 world's biodiversity. Vast new datasets (e.g. species distributions, traits, phylogenies, and interaction
28 networks) hold knowledge to better comprehend the depths of biodiversity change, reliably anticipate
29 these changes, and inform conservation actions. To harness this information for conservation, we need to
30 integrate the largely independent fields of biodiversity modelling and conservation. We highlight new
31 developments in each respective field, early examples of how they are being brought together, and ideas
32 for a future synthesis so conservation decisions can be made with a fuller awareness of the biodiversity at
33 stake.

34

35 **Main Text**

36

37 ***We need greater diversity in biodiversity conservation***

38 Aside from a small fraction of well-known species with established conservation status, most of
39 the world's biodiversity is poorly integrated into conservation efforts. Even for iconic clades
40 such as birds and mammals, only a subset of species are considered in conservation despite a
41 continuously refined understanding of the ecology and evolution of these clades[1,2]. Lesser
42 known taxa are nearly always overlooked[3] even those most at risk of extinction[4]. First, we
43 need to expand the focus of biodiversity in conservation programs to consider more of the tree of
44 life, even taxa without names[5,6], especially as increasing evidence suggests 'invisible'
45 diversity is important to ecosystem functioning [7]. Second, we must consider the role of species
46 in ecosystems[8]. Biodiversity is not simply the mere sum of species, but includes complex
47 interaction networks with fluxes of energy and nutrients[9,10]. Even conservation efforts for

48 focal charismatic species would benefit from considering the other species enabling their
49 existence. Biodiversity losses impact all facets of biodiversity (species, genetic diversity,
50 **functional diversity, phylogenetic diversity**, see Glossary) and spatial extents ranging from
51 local assemblages to the global pool of biodiversity. Considering these dimensions is essential to
52 prevent widespread biodiversity loss.

53

54 *Conservation evaluation often falls short for under-represented biodiversity*

55 *Simplified metrics likely miss many types of biodiversity*

56 Despite scientific recognition of the importance of biodiversity across multiple facets, most
57 planning and on-the-ground conservation still focus on a simplified representation of biodiversity
58 such as the amount or quality of habitat. Habitat-based metrics are used more often in
59 conservation than in ecology[11], and are often the basis of conservation offsetting, wetland
60 mitigation, conservation easements, conservation and management of listed species (alongside
61 population abundances, connectivity, and threat information when available), and are used at a
62 regional extent in coordinated reserve designs (e.g. Natura 2000,
63 https://ec.europa.eu/environment/nature/natura2000/index_en.htm). Reliance on habitat metrics
64 can overlook important biodiversity that falls outside of commonly held notions of high-quality
65 habitat[12], and these metrics do not necessarily represent population persistence[11]. While the
66 use of habitat metrics certainly has follow-on benefits to some diversity, its surrogacy value for
67 even well-known species is unclear[13], and unknown for many taxa and for different
68 **biodiversity facets.**

69

70 In cases where multiple species are considered together in applied conservation, they are most
71 often represented as species richness (the count of species in areas)[11]. The use of species
72 richness cannot address the known biases in extinction risk towards certain clades (e.g.
73 amphibians[14]) and functional types (e.g. top predators). While species richness and other
74 species-based metrics can be an effective surrogate for phylogenetic diversity[10] or functional
75 diversity[15], its surrogacy value depends on the conservation scenario[16] and tends to decrease
76 in scenarios where resources are limited (i.e. few sites available for protection)[17]. We have
77 very limited evidence for surrogacy value of different biodiversity facets outside of iconic animal
78 groups (e.g. does aboveground animal richness relate to soil microbial diversity?). Finally, how
79 species interact within ecosystems is almost never considered in applied conservation despite
80 links to the important conservation goals of ecosystem functioning and resilience[9].

81

82 *Independent conservation efforts fail to protect the world's biodiversity*

83 Safeguarding multiple facets of biodiversity—especially for the purposes of preventing
84 extinction, protecting our biological heritage, and retaining future options for society—requires
85 an understanding of the global biodiversity pool, arguably more than for other conservation aims
86 (e.g. managing local ecosystems or protecting ecosystem services)[18]. But conservation actions
87 are typically local or regional, which can lead to unintended outcomes for global
88 biodiversity[19]. First, species are declining globally[20], but not necessarily locally[21–23],
89 leading to a homogenization of biodiversity where different areas have increasingly similar
90 biodiversity[20]. Second, conservation priorities established for local species assemblages[16] or
91 even for species at a national scale[19] differ from global biodiversity priorities[16,19]. Lastly,
92 the emphasis on protecting large patches and connected habitats (e.g. species corridor
93 initiatives), which is based on research on metapopulations and conservation genetics, risks

94 overlooking important biodiversity (e.g. endemic species, highly functional communities) that
95 occur in small and isolated habitat patches [24]. A more spatially-coordinated approach is needed
96 that recognizes how local conservation efforts impact the regional and global biodiversity pool.

97

98 *The potential to bring together biodiversity modelling and conservation*

99 The fields of ecology and conservation planning already consider the spatial extents necessary
100 for this coordination, but they do so in slightly different ways. In systematic conservation
101 planning (SCP), biodiversity features (e.g. species distributions) are considered in planning units.
102 Planning units are compared across the entire area of interest using the principles of spatial
103 **complementarity** and **irreplaceability**. In ecology, a similar hierarchy can be found in the
104 delineation of α -(within-site or within-ecosystem diversity), β -(differentiation of sites in a
105 region; e.g. compositional turnover), and γ -**diversity** (diversity of the entire region). We refer to
106 γ -**diversity** as the total set of biodiversity in the area of interest, which could have a regional or
107 global extent, to better align with conservation planning. Both fields are progressing
108 independently toward a multi-scale, multi-faceted view of biodiversity (Figure 1). Now is the
109 time to integrate these fields to rapidly expand the amount of biodiversity data that reaches
110 conservation practice (Figure 2).

111

112 **Advances in biodiversity models**

113 While steady progress has been made on addressing the major biodiversity shortfalls with
114 increased data collection (e.g. citizen science[25], environmental DNA) and global initiatives to
115 compile data on species occurrences (e.g. GEO-BON initiative; geobon.org, GBIF; gbif.org),
116 traits (e.g. TRY;www.try-db.org), genetic and phylogenetic data (e.g. GenBank), and species

117 interaction data (e.g. GloBI; globalbioticinteractions.org), major shortfalls remain[26].
118 Biodiversity models are essential tools to rapidly overcome these shortfalls. We consider
119 **biodiversity models** to be any model that makes predictions for biodiversity (including species
120 occurrences or abundances, traits, phylogenetic placement, or species interactions). Here, we
121 highlight two model classes: (1) **imputation models** for missing biodiversity data and (2) **spatial**
122 **biodiversity models** for predicting taxa or other biodiversity facets for unsampled locations,
123 which often serve as an input for conservation planning (Figure 1).

124

125 *Imputation models*

126 Imputation models are needed to fill in the many gaps in biodiversity datasets. For example, they
127 can generate missing trait values for taxa[27], which can be estimated from phylogenies,
128 assuming traits have a general pattern of conserved evolution[28] or through multivariate
129 imputation that does not rely on phylogenies[29]. Increasingly sophisticated approaches are also
130 being used to generate phylogenies when some gene sequences are missing[30], and to combine
131 multiple datasets into aggregated phylogenies either based on the phylogenies themselves ('super
132 trees' or preferably from a combined gene matrix[31]. There is also growing interest in
133 predicting interactions between species based on their trait attributes (e.g. predator-prey trait
134 matching[32]) and spatial co-occurrence data[33]. Predicting species interactions across many
135 species or at multiple locations is difficult with severely limited data, but can be addressed by
136 combining empirical data and expert knowledge[34].

137

138 While imputation models are greatly expanding the coverage of biodiversity datasets, challenges
139 remain. Data collection is systematically geographically and taxonomically biased, which could

140 bias predictions, especially for rare taxa. We are increasingly relying on the same datasets for
141 different types of analyses, so there is also the danger of circularity if those analyses are later
142 combined (e.g. using the phylogeny to predict trait values and later combining phylogeny and
143 traits in spatial models). We need more robust tests for the impact of these biases, and to ensure
144 that uncertainty is propagated from the original data sources to all subsequent analyses. A partial
145 solution is to integrate methods and create a transparent and documented pipeline of analyses
146 (Figure 2), but we still must address the underlying causes of bias and use this knowledge to
147 guide future collection.

148

149

150 *Spatial Biodiversity models*

151

152 The Wallacean shortfall (inadequate knowledge of geographic distributions) greatly hinders
153 progress in identifying conservation priorities[26]. This shortfall is particularly problematic at
154 local scales—it is easier to define a regional or national set of species than to pinpoint the precise
155 distributions of species at fine spatial scales[26]. Biodiversity models are essential tools to
156 address this shortfall. In many cases, they can provide continuous coverage of species
157 distributions at a resolution relevant for conservation purposes[35]. The most common examples
158 are single species distribution models (i-SDMs; Fig. I), which have been and will continue to be
159 invaluable tools for conservation applications [35]. Advanced models can additionally
160 accommodate multiple facets (e.g. traits and phylogenies) and integrate elements of α - and β -
161 diversity (Box 1). A notable advance is the ability to combine a top-down approach focused on
162 an entire species assemblage with a bottom-up approach focused on single species (**‘sideways’**
163 **biodiversity models**; Box 1, Box 2). One such model, Joint Species Distribution Models (j-
164 SDMs, reviewed in [36]), offers potential improvements by modeling all species simultaneously

165 [37,38], enabling conditional predictions [39–41], and has a flexible hierarchy amenable to
166 adding other information such as traits[37], phylogenies[42], geographic space[43], temporal
167 data[44], and detection bias[45], which impact multifaceted biodiversity estimates[46]. Models
168 that can account for species interactions[47] are an exciting arena for future developments.

169

170 A key advantage of these advanced biodiversity models is the ability to process highly
171 multidimensional datasets (e.g. many taxa, sites, and environmental variables). Combining
172 multiple, large biodiversity datasets can quickly result in a computationally intractable problem.
173 Solutions include dimensionality reduction in a Bayesian framework[36,48] or via machine
174 learning[49]. However, what these approaches gain in flexibility may come at a cost in terms of
175 generality, and the relationships derived between explanatory variables and diversity metrics are
176 often empirical with little to no grounding in ecological theory. An important future research
177 avenue will be to rigorously test new biodiversity models to evaluate their capacity for aiding
178 ecological inference and for predicting to new situations (e.g. novel climates or altered
179 ecosystems).

180

181 **Advances in conservation evaluation of multifaceted biodiversity**

182 While the methods described above could offer a richer representation of biodiversity and
183 improved predictive abilities, identifying conservation solutions requires more than just
184 biodiversity predictions. These predictions must be translated to meaningful outputs that can be
185 used to meet defined objectives (e.g. protected area targets) or provide forecasts for decision
186 makers. In systematic conservation planning (SCP)[50], priorities are established that efficiently
187 meet goals for representation of biodiversity under a range of spatial or other constraints (e.g. in
188 a **spatial prioritization**). Targets can be set on diversity (e.g. 10% of each species range

189 protected[17]) or approaches without targets such as algorithms based on **weighted**
190 **endemism**[16,19]. While i-SDMs are commonly used as inputs to spatial prioritization, more
191 advanced biodiversity models are rarely used (see Box 2 for exceptions). Multiple biodiversity
192 facets are also rarely considered in conservation despite decades of recognized conservation
193 benefit of *e.g.* phylogenetic diversity[51]. However, in recent years methodological
194 advances[52–54] have enabled the use of phylogenies and functional diversity [16,53] in
195 conservation planning studies at regional[6,52] and global[16,17] extents, and initiatives such as
196 EDGE (evolutionarily distinct and globally endangered; edgeofexistence.org) are promoting the
197 conservation of multiple biodiversity facets. Species interactions are even more rarely
198 considered, although emerging examples show how interactions can be used to evaluate
199 ecological network collapse[55], and set spatial priorities[56] and management scenarios that
200 account for predators and their prey[57]

201

202 **The way forward: integrating biodiversity models and conservation evaluation**

203 Modeling has a critical, yet under-appreciated role in conservation planning and practice,
204 especially with respect to setting and achieving global conservation targets (Aichi targets for
205 threatened species, habitat loss, and protected areas) in the Strategic Plan for Biodiversity 2011-
206 2020 and the post-2020 global biodiversity framework[58]. Models that predict the distribution
207 of species in space and time are crucial to these global initiatives, and i-SDMs are already widely
208 used in conservation[35]. We believe the more advanced biodiversity models have vast potential
209 for representing not just species but entire communities, clades, or functional roles (see early
210 examples in Box 2), but they first must be better-integrated into conservation efforts.

211

212 The advantages of this integration are many. First, a combined approach retains the unique
213 strengths of each individual approach. Conservation planning would benefit from having access
214 to the most advanced statistical machinery including model-calibration, model validation, and
215 the propagation of uncertainty. Such models can also incorporate more flexible hierarchical
216 structures that account for biases in sampling biodiversity, different sources of inputs (e.g.
217 museum location data with community-based eDNA sampling), different extents (α - and β -
218 diversity), and different resolutions of input data. They can take advantage of new data types
219 (e.g. tracking data[59]) as they become available. Conversely, the field of biodiversity modelling
220 would benefit from tailoring predictions for their use in tangible conservation scenarios and
221 targets.

222

223 Second, an integrated approach reduces any redundancies that arise from using similar datasets
224 and similar approaches in an unsynchronized manner. The potential for redundancy becomes
225 greater the larger and the more complex the representation of biodiversity becomes. Redundancy
226 could be reduced on two fronts—the raw data inputs and the methodological similarities (e.g.
227 combining modelling and conservation optimization algorithms as discussed below).

228

229 Finally, this synthesis would make the pipeline of data-implementation more efficient and likely
230 to result in a better representation of under-represented biodiversity in conservation. Recent calls
231 to streamline analytical pipelines in ecology[60–62] point to increased efficiency [63] of readily
232 available technological solutions, such as user-friendly interfaces (e.g. R Shiny, Dash).

233 Importantly, the integration of conservation goals, targets, and scenarios into modelling would
234 require stakeholder input into the modelling decisions[64], which could lead to increased

235 coordination, trust and uptake of these models, decreasing the ‘black box’ problem of
236 biodiversity models[60]. It would engage biodiversity scientists beyond academia to understand
237 the needs of stakeholders and provide more tailored conservation guidance.

238

239 **Steps forward**

240 We recommend three related areas of focus to speed integration of biodiversity models and
241 conservation: (1) adapting biodiversity models to accommodate conservation goals and
242 scenarios, (2) combining biodiversity models and conservation optimization algorithms, and (3)
243 re-evaluating model validation in light of conservation goals and scenarios. Recent examples
244 show this integration is indeed possible (Box 2).

245

246 In many cases, statistical methods already exist to adapt biodiversity models for use in
247 conservation. For example, management actions (or management history) could be included as
248 model predictors, which could then be used to make predictions for different future actions.

249 More complex methods of scenario-building are also possible through altering various model
250 parameters [65,66](Box 2) or even adding sub-models of ecological processes (e.g. population
251 viability). Previously collected ‘prior’ information (information on species threat or response to
252 management action) can be combined with newly collected data to refine model estimates[67].

253 The challenge will be to find the best data and models for the conservation question at hand[68],
254 requiring close collaboration between modelers and conservation practitioners[65].

255

256 Combining conservation optimization algorithms with biodiversity models is another important
257 step (Figure 2) that will enable relevant ecological processes to be directly considered in the

258 conservation alternatives[65]. This integration will require some method developments, but will
259 benefit from the similarities in data types (Figure 1), especially when definitions are
260 synchronized (e.g. α -diversity is equivalent to diversity in planning units, β -diversity to
261 complementarity[69], γ -diversity to the total set of biodiversity). In particular, β -diversity could
262 be used in ecological models with scenarios (Box 2) or within optimization algorithms,
263 streamlining methods and opening the possibility of evaluating conservation value of
264 biodiversity not represented in a single-species approach (e.g. ecological interaction
265 networks[70]).

266

267 Finally, model evaluation—while always a critical step in biodiversity modelling—is especially
268 important with conservation scenarios that are often extrapolations to new situations (e.g.
269 different environmental conditions, alternative restoration scenarios, poorly known taxa). Model
270 evaluation will help address questions such as which type of model to use. For some
271 conservation applications, a highly flexible, accurate model is likely appropriate (e.g. predicting
272 the current distribution of a threatened species). For others, it might be preferable to have models
273 more closely aligned with ecological theory (e.g. when predicting to novel environments such as
274 climate change) or when the conservation question depends on a particular variable (e.g.
275 properties of a connectivity corridor). While there are examples that evaluate the ability of
276 advanced biodiversity models to predict rare species[71] and their interactions[72], very few
277 studies have tested how well models extrapolate to new conditions with independent datasets, or
278 whether they predict realistic conservation outcomes. Much more research is needed to
279 understand how biodiversity models will perform in an applied setting.

280

281

282 ***Concluding Remarks***

283 We urgently need to understand how biodiversity change impacts critical ecosystem functioning
284 and prunes the tree of life. We propose an integration of advanced biodiversity models with
285 conservation goals, targets, assessments, and practice. This integration would improve the ability
286 to rapidly evaluate biodiversity data, make predictions and recommend conservation action for
287 taxa, communities, and ecosystems. The backbone of this integration already exists with recent
288 developments in biodiversity modelling and conservation. Early examples show this integration
289 is possible, could be extended to a number of other conservation efforts and used to address
290 unresolved questions of how to best protect the world's biodiversity (see Outstanding
291 Questions).

292

293 **Acknowledgements**

294 LJP was supported with Marie Curie Fellowship 'Conserving the Legacy of Evolution into the
295 Future) AMD-659422-1. This work received funding from the ERA-Net BiodivERsA -
296 Belmont Forum, with the national funder Agence National pour la Recherche (FutureWeb:
297 ANR-18-EBI4-0009).

298

299

300 **Glossary Box**

301

302 **Alpha (α) diversity:** quantifies local diversity, usually species richness or effective number of
303 species (Shannon exponent)

304

305 **Beta (β) diversity:** quantifies the dissimilarity or turnover between locations in terms of
306 composition. Can be pairwise between locations, global (average dissimilarity across the region)
307 or the distinctiveness of a location compared to all others in the region.

308

309 **Biodiversity facets:** categories of biodiversity that describe relevant taxonomic or ecological
310 information to support biodiversity evaluation, and that apply to and can be (relatively) easily
311 measured for all or most taxa. Examples include taxonomic diversity, phylogenetic diversity,
312 genetic diversity, trait/functional diversity and network diversity.

313

314 **Biodiversity models:** statistical or process-based models that are used to make inferences and
315 predictions about the effect of the environment on biodiversity, accounting for ecological
316 processes explicitly or implicitly. Biodiversity can be represented in units ranging from
317 individuals to entire communities and additionally represent attributes of those taxa (e.g.
318 abundances, functional traits, phylogenetic position, threat).

319

320 **Complementarity (conservation planning):** quantifies the difference between locations in
321 terms of species or features represented within them (two locations are fully complementary if
322 their pairwise β -diversity is equal to 1).

323

324 **Complementarity (ecology):** two species are complementary if they fill different roles in an
325 ecosystem or use resources differently.

326

327 **Functional diversity (FD):** the diversity of functional forms in a species set (or community)
328 measured by a variety of metrics that use dendrograms or representations in multidimensional
329 space.

330

331 **Gamma (γ) diversity:** quantifies the total diversity across locations for an entire area of interest,
332 which could be at regional or global scale depending on the study.

333

334 **Imputation model:** a model that estimates values for missing data points in biodiversity datasets
335 (e.g. trait values for a particular taxa).

336

337 **Irreplaceability:** the extent to which a location or species is distinct from all others (opposite of
338 redundancy). For example, a location where an endemic species occurs is irreplaceable; an
339 irreplaceable species has a unique position in its community (in terms of function or interactions,
340 i.e. distinctive) or in the phylogeny.

341

342 **Macro-ecological model:** a biodiversity model that uses a top-down approach to model α - or β -
343 diversity directly rather than modeling the distributions of the component taxa.

344

345 **Phylogenetic diversity (PD):** The evolutionary diversity represented by sets of taxa, with the
346 most common metric (Faith's PD) being the branch length of the minimum spanning tree
347 connecting a set of species on a phylogeny[51] (sets of species can be from a single area or
348 multiple areas combined).

349

350 **'Sideways' biodiversity models:** models that predict the distribution of biodiversity with a
351 combination of bottom-up (i.e. single-taxon predictions) and top-down (i.e. models of the
352 properties of an assemblage or community of taxa) approaches.

353

354 **Spatial biodiversity models:** biodiversity models that are explicitly spatial, in which inferences
355 and predictions are made for biodiversity in certain locations (e.g. planning units or grid cells).

356

357 **Spatial prioritization:** A form of systematic conservation planning (SCP) that selects a set of
358 areas that maximizes conservation value given other constraints (e.g. cost, protected areas,
359 feasibility).

360

361 **Weighted endemism:** the ratio between a species local range of occurrence and its total range.
362 Note: this is different than the definition of endemism which is the extent to which a species
363 range is restricted to a particular location.

364

365

366 **Figure 1.** How biodiversity facets, different scales of biodiversity and conservation concerns
367 factor into biodiversity modelling and conservation evaluation. Most modelling and evaluation
368 involve particular target species rather than other facets of diversity (e.g. evolutionary history,
369 species interactions). Most biodiversity models focus on ecological processes rather than
370 explicitly considering conservation concerns, although many models are built for threatened
371 species or species of conservation interest resulting in a mostly unidirectional flow of
372 information from models to evaluation. In the case of spatial conservation planning, both models
373 and evaluation use similar data inputs and spatial organization of biodiversity. Both have local

374 scale (or planning units) and a regional or global scale. In particular, the use of metrics or
375 algorithms to differentiate sites from one another (beta diversity in modelling, complementarity
376 in conservation planning) are treated similarly in modelling and planning.

377

378 **Figure 2.** The steps and information flow in a typical data-implementation pipeline (left) and a
379 streamlined version (right) that combines models and evaluation in one step. We expect that
380 more biodiversity information would make it through to the application stage (orange arrows)
381 with a unified rather than disconnected method of evaluating biodiversity from the data
382 products to their use in conservation applications, and that the increasing transparency and
383 additional communication between biodiversity scientists and stakeholders is required by the
384 combined approach.

385

386 **Figure I.** (Embedded in Box 1). The dimensions of recent biodiversity models. Models can have
387 different types of response objectives: within-site models, between-site models and models of
388 individual species. These response objectives fall along two axes: (1) top-down approaches
389 where α -diversity or other diversity metrics are the focus versus bottom-up where individual
390 species are the focus (vertical axis), and (2) whether the focus is on predicting biodiversity
391 within sites or between sites (horizontal axis). DynamicFOAM: Dynamic Framework for
392 Occurrence Allocation in Metacommunities, GDM: Generalized Dissimilarity Modelling, MEM:
393 macro-ecological model, i-SDM: individual or single-species distribution model, s-SDM:
394 stacked species distribution model, j-SDM: joint species distribution model, SESAM: Spatially
395 explicit species assemblage modelling.

396

397 **Box 1. Modelling biodiversity from the top-down, bottom-up and sideways**

398 Biodiversity models can be defined by their ‘response objectives’ and internal structure
399 (particularly the response variable—what is being predicted). These response objectives can be
400 represented with axes that describe whether the objective is a top-down versus bottom-up or a
401 within- versus between-site depiction of biodiversity (Fig. I). Current modelling approaches can
402 be mapped onto this space, with potentially the most information rich approaches (grey circle in
403 Fig. I) cross-linking biodiversity facets (traits and phylogenies) while jointly predicting
404 community composition (e.g. species abundance or occurrence), within-site community
405 diversity, and between-site turnover.

406

407 In the top-down approach, known as a **macro-ecological model** (MEM), α -diversity is itself is
408 the response (e.g. species richness or phylogenetic diversity) and the input data is a site by
409 diversity matrix. β -diversity models (e.g. GDM; Generalized Dissimilarity Modelling[73] or
410 MBM; multi-faceted biodiversity models[49]) estimate turnover in species or phylogenetic
411 diversity[74] between sites as a function of differences in environmental and geographic
412 distances.

413

414 Bottom-up approaches, where individual species are the response variables (single-species
415 distribution model, i-SDM), are by far the most common type of biodiversity model used. Note
416 that α -diversity can also be derived from aggregating models of individual species together in a
417 stacked species distribution model (s-SDM). This feature is important for conservation planning,
418 because both the species pool of a set of sites and diversity metrics for entire species
419 assemblages can be calculated (e.g. endemism-based metrics). However, resulting diversity

420 metrics are simply the sum of the parts—there is no information on species associations or
421 community-level attributes.

422 ‘Sideways’ methods are methods that combine top-down and bottom-up approaches: Joint
423 Species Distribution Models (j-SDMs), which are the multi-species version of i-SDMs that
424 model effects of individual species and communities[39,40], and can include shared responses
425 between species and can include other information such as phylogeny and traits in a hierarchical
426 framework, and spatial and temporal dynamics[66]; Dynamic Framework for Occurrence
427 Allocation in Metacommunities (DynamicFOAM)[75], which balances richness and turnover to
428 predict occurrences for all species, SESAM: Spatially explicit species assemblage modelling[76],
429 which adjusts combined s-SDM predictions based on those of macro-ecological models.

430

431 **Box 2. Hybrid approaches for integrating advanced biodiversity models and conservation**
432 **decisions**

433 We have emphasized the need for models that join different spatial and taxonomic dimensions of
434 biodiversity that can be combined with conservation evaluation. Here are examples from two
435 classes of advanced biodiversity models (Box 1) that directly address applied conservation
436 questions. There is much scope for building upon and extending these early examples.

437

438 *1. Macro-ecological models*

439 For taxa for which there is no abundance or time-series data, models based on α - and β -diversity
440 (Box1) can also be used to directly design, predict, and compare conservation scenarios. For
441 example, β -diversity models can summarise the expected effects on future biodiversity of
442 changes to habitat condition, extent or management, as change in effective habitat area[77]. With
443 sophisticated models of environmental or land-use change for example, new protected areas,

444 habitat loss[78] or degradation and climate change[79], scenarios can be planned for the most
445 beneficial effect on biodiversity even at global extents[77], and can also be used with
446 phylogenetic[74] or functional[49] β -diversity.

447 2. 'Sideways' biodiversity models

448 Multi-species models that contain a hierarchy for a species-level and community (or ecosystem)-
449 level (Box 1) are widely relevant to conservation applications because they can simultaneously
450 estimate ecological processes relevant for individual species and for larger-scale attributes (e.g.
451 ecosystem properties). Perhaps the best example of this approach has been for evaluating
452 management strategies in fisheries, in which models of fish species (or sets of species) are
453 combined with ecosystem level attributes[64]. In this way, important ecological and ecosystem
454 constraints can be considered (e.g. population dynamics in response to habitat or predators)
455 along with management scenarios (harvest controls)[57]. Recent versions allow for evaluation of
456 harvest impacts, survey designs, and protection scenarios for finescale habitats in a changing
457 climate[66]. While these more elaborate designs are only possible for a subset of well-studied
458 taxa, this will be increasingly possible with ever-improving sampling and imputation models.
459

460 **References**

- 461 1 Jetz, W. *et al.* (2012) The global diversity of birds in space and time. *Nature* 491, 444–448
462 2 Mazel, F. *et al.* (2017) Global patterns of β -diversity along the phylogenetic time-scale: The
463 role of climate and plate tectonics. *Glob. Ecol. Biogeogr.* 26, 1211–1221
464 3 Eisenhauer, N. *et al.* (2019) Recognizing the quiet extinction of invertebrates. *Nat. Commun.*
465 10, 1–3

- 466 4 Davies, T. *et al.* (2018) Popular interest in vertebrates does not reflect extinction risk and is
467 associated with bias in conservation investment. *PLOS ONE* 13, e0203694
- 468 5 Asmyhr, M.G. *et al.* (2014) Systematic Conservation Planning for Groundwater Ecosystems
469 Using Phylogenetic Diversity. *PLoS ONE* 9, e115132
- 470 6 Rosauer, D.F. *et al.* (2018) Real-world conservation planning for evolutionary diversity in
471 the Kimberley, Australia, sidesteps uncertain taxonomy. *Conserv. Lett.* 11, e12438
- 472 7 Delgado-Baquerizo, M. *et al.* (2016) Microbial diversity drives multifunctionality in
473 terrestrial ecosystems. *Nat. Commun.* 7, 10541
- 474 8 Violle, C. *et al.* (2017) Functional Rarity: The Ecology of Outliers. *Trends Ecol. Evol.* 32,
475 356–367
- 476 9 Harvey, E. *et al.* (2017) Bridging ecology and conservation: from ecological networks to
477 ecosystem function. *J. Appl. Ecol.* 54, 371–379
- 478 10 Thompson, R.M. *et al.* (2012) Food webs: reconciling the structure and function of
479 biodiversity. *Trends Ecol. Evol.* 27, 689–697
- 480 11 Marshall, E. *et al.* (2020) What are we measuring? A review of metrics used to describe
481 biodiversity in offsets exchanges. *Biol. Conserv.* 241, 108250
- 482 12 Hobbs, R.J. (2016) Degraded or just different? Perceptions and value judgements in
483 restoration decisions. *Restor. Ecol.* 24, 153–158
- 484 13 Beier, P. *et al.* (2015) A review of selection-based tests of abiotic surrogates for species
485 representation. *Conserv. Biol.* 29, 668–679
- 486 14 González-del-Pliego, P. *et al.* (2019) Phylogenetic and Trait-Based Prediction of Extinction
487 Risk for Data-Deficient Amphibians. *Curr. Biol.* 29, 1557-1563.e3

- 488 15 Rapacciuolo, G. *et al.* (2019) Species diversity as a surrogate for conservation of
489 phylogenetic and functional diversity in terrestrial vertebrates across the Americas. *Nat.*
490 *Ecol. Evol.* 3, 53–61
- 491 16 Pollock, L.J. *et al.* (2017) Large conservation gains possible for global biodiversity facets.
492 *Nature* 546, 141–144
- 493 17 Rosauer, D.F. *et al.* (2017) Phylogenetically informed spatial planning is required to
494 conserve the mammalian tree of life. *Proc R Soc B* 284, 20170627
- 495 18 Pearson, R.G. (2016) Reasons to Conserve Nature. *Trends Ecol. Evol.* 31, 366–371
- 496 19 Pouzols, F.M. *et al.* (2014) Global protected area expansion is compromised by projected
497 land-use and parochialism. *Nature* 516, 383–386
- 498 20 Di Marco, M. *et al.* (2018) Changes in human footprint drive changes in species extinction
499 risk. *Nat. Commun.* 9, 1–9
- 500 21 Sax, D.F. and Gaines, S.D. (2003) Species diversity: from global decreases to local
501 increases. *Trends Ecol. Evol.* 18, 561–566
- 502 22 Vellend, M. *et al.* (2013) Global meta-analysis reveals no net change in local-scale plant
503 biodiversity over time. *Proc. Natl. Acad. Sci. U. S. A.* 110, 19456–19459
- 504 23 Cardinale, B.J. *et al.* (2018) Is local biodiversity declining or not? A summary of the debate
505 over analysis of species richness time trends. *Biol. Conserv.* 219, 175–183
- 506 24 Wintle, B. *et al.* (2018) Global synthesis of conservation studies reveals the importance of
507 small habitat patches for biodiversity. *Proc. Natl. Acad. Sci.* 116, 201813051
- 508 25 Kobori, H. *et al.* (2016) Citizen science: a new approach to advance ecology, education, and
509 conservation. *Ecol. Res.* 31, 1–19

- 510 26 Hortal, J. *et al.* (2015) Seven Shortfalls that Beset Large-Scale Knowledge of Biodiversity.
511 *Annu. Rev. Ecol. Evol. Syst.* 46, 523–549
- 512 27 Wilman, H. *et al.* (2014) EltonTraits 1.0: Species-level foraging attributes of the world's
513 birds and mammals. *Ecology* 95, 2027–2027
- 514 28 Swenson, N.G. (2014) Phylogenetic imputation of plant functional trait databases.
515 *Ecography* 37, 105–110
- 516 29 Penone, C. *et al.* (2014) Imputation of missing data in life-history trait datasets: which
517 approach performs the best? *Methods Ecol. Evol.* 5, 961–970
- 518 30 Bhattacharjee, A. and Bayzid, Md.S. (2020) Machine learning based imputation techniques
519 for estimating phylogenetic trees from incomplete distance matrices. *BMC Genomics* 21, 497
- 520 31 von Haeseler, A. (2012) Do we still need supertrees? *BMC Biol.* 10, 13
- 521 32 Gravel, D. *et al.* (2013) Inferring food web structure from predator–prey body size
522 relationships. *Methods Ecol. Evol.* 4, 1083–1090
- 523 33 Morales-Castilla, I. *et al.* (2015) Inferring biotic interactions from proxies. *Trends Ecol.*
524 *Evol.* 30, 347–356
- 525 34 Braga, J. *et al.* (2019) Spatial analyses of multi-trophic terrestrial vertebrate assemblages in
526 Europe. *Glob. Ecol. Biogeogr.* 28, 1636–1648
- 527 35 Guisan, A. *et al.* (2013) Predicting species distributions for conservation decisions. *Ecol.*
528 *Lett.* 16, 1424–1435
- 529 36 Warton, D.I. *et al.* (2015) So Many Variables: Joint Modeling in Community Ecology.
530 *Trends Ecol. Evol.* 30, 766–779
- 531 37 Pollock, L.J. *et al.* (2012) The role of functional traits in species distributions revealed
532 through a hierarchical model. *Ecography* 35, 716–725

- 533 38 Ovaskainen, O. and Soininen, J. (2011) Making more out of sparse data: hierarchical
534 modeling of species communities. *Ecology* 92, 289–295
- 535 39 Clark, J.S. *et al.* (2014) More than the sum of the parts: forest climate response from joint
536 species distribution models. *Ecol. Appl.* 24, 990–999
- 537 40 Pollock, L.J. *et al.* (2014) Understanding co-occurrence by modelling species simultaneously
538 with a Joint Species Distribution Model (JSDM). *Methods Ecol. Evol.* 5, 397–406
- 539 41 Ovaskainen, O. *et al.* (2010) Modeling species co-occurrence by multivariate logistic
540 regression generates new hypotheses on fungal interactions. *Ecology* 91, 2514–2521
- 541 42 Ives, A.R. and Helmus, M.R. (2011) Generalized linear mixed models for phylogenetic
542 analyses of community structure. *Ecol. Monogr.* 81, 511–525
- 543 43 Ovaskainen, O. *et al.* (2016) Uncovering hidden spatial structure in species communities
544 with spatially explicit joint species distribution models. *Methods Ecol. Evol.* 7, 428–436
- 545 44 Schliep, E.M. *et al.* (2018) Joint species distribution modelling for spatio-temporal
546 occurrence and ordinal abundance data. *Glob. Ecol. Biogeogr.* 27, 142–155
- 547 45 Tobler, M.W. *et al.* (2019) Joint species distribution models with species correlations and
548 imperfect detection. *Ecology* 100, e02754
- 549 46 Jarzyna, M.A. and Jetz, W. (2016) Detecting the Multiple Facets of Biodiversity. *Trends*
550 *Ecol. Evol.* 31, 527–538
- 551 47 Staniczenko, P.P.A. *et al.* (2017) Linking macroecology and community ecology: refining
552 predictions of species distributions using biotic interaction networks. *Ecol. Lett.* 20, 693–707
- 553 48 Taylor-Rodríguez, D. *et al.* (2017) Joint Species Distribution Modeling: Dimension
554 Reduction Using Dirichlet Processes. *Bayesian Anal.* 12, 939–967

555 49 Talluto, M.V. *et al.* (2018) Multifaceted biodiversity modelling at macroecological scales
556 using Gaussian processes. *Divers. Distrib.* 24, 1492–1502

557 50 Margules, C.R. and Pressey, R.L. (2000) Systematic conservation planning. *Nature* 405,
558 243–252

559 51 Faith, D.P. (1992) Conservation evaluation and phylogenetic diversity. *Biol. Conserv.* 61, 1–
560 10

561 52 Pollock, L.J. *et al.* (2015) Phylogenetic diversity meets conservation policy: small areas are
562 key to preserving eucalypt lineages. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 370, 20140007

563 53 Strecker, A.L. *et al.* (2011) Defining conservation priorities for freshwater fishes according
564 to taxonomic, functional, and phylogenetic diversity. *Ecol. Appl.* 21, 3002–3013

565 54 Strimas-Mackey, M. *prioritizr: Systematic Conservation Prioritization in R*,

566 55 McDonald-Madden, E. *et al.* (2016) Using food-web theory to conserve ecosystems. *Nat.*
567 *Commun.* 7, 1–8

568 56 Decker, E. *et al.* (2017) Incorporating ecological functions in conservation decision making.
569 *Ecol. Evol.* 7, 8273–8281

570 57 Punt, A.E. *et al.* (2016) Exploring the implications of the harvest control rule for Pacific
571 sardine, accounting for predator dynamics: A MICE model. *Ecol. Model.* 337, 79–95


572 58 Nicholson, E. *et al.* (2019) Scenarios and Models to Support Global Conservation Targets.
573 *Trends Ecol. Evol.* 34, 57–68


574 59 Hays, G.C. *et al.* (2019) Translating Marine Animal Tracking Data into Conservation Policy
575 and Management. *Trends Ecol. Evol.* 34, 459–473

576 60 Rapacciuolo, G. (2019) Strengthening the contribution of macroecological models to
577 conservation practice. *Glob. Ecol. Biogeogr.* 28, 54–60

- 578 61 Golding, N. *et al.* (2018) The zoon r package for reproducible and shareable species
579 distribution modelling. *Methods Ecol. Evol.* 9, 260–268
- 580 62 White, E.P. *et al.* (2019) Developing an automated iterative near-term forecasting system for
581 an ecological study. *Methods Ecol. Evol.* 10, 332–344
- 582 63 Lowndes, J.S.S. *et al.* (2017) Our path to better science in less time using open data science
583 tools. *Nat. Ecol. Evol.* 1, 0160
- 584 64 Plagányi, É.E. *et al.* (2014) Multispecies fisheries management and conservation: tactical
585 applications using models of intermediate complexity. *Fish Fish.* 15, 1–22
- 586 65 Scroggie, M.P. *et al.* (2019) Optimizing habitat management for amphibians: From simple
587 models to complex decisions. *Biol. Conserv.* 236, 60–69
- 588 66 Thorson, J.T. *et al.* (2019) Spatio-temporal models of intermediate complexity for ecosystem
589 assessments: A new tool for spatial fisheries management. *Fish Fish.* 20, 1083–1099
- 590 67 Morris, W.K. *et al.* (2015) The neglected tool in the Bayesian ecologist’s shed: a case study
591 testing informative priors’ effect on model accuracy. *Ecol. Evol.* 5, 102–108
- 592 68 Guillera-Aroita, G. *et al.* (2015) Is my species distribution model fit for purpose? Matching
593 data and models to applications. *Glob. Ecol. Biogeogr.* 24, 276–292
- 594 69 Bush, A. *et al.* (2016) Current Uses of Beta-Diversity in Biodiversity Conservation: A
595 response to Socolar *et al.* *Trends Ecol. Evol.* 31, 337–338
- 596 70 Ohlmann, M. *et al.* (2019) Diversity indices for ecological networks: a unifying framework
597 using Hill numbers. *Ecol. Lett.* 22, 737–747
- 598 71 Zhang, C. *et al.* (2020) Improving prediction of rare species’ distribution from community
599 data. *Sci. Rep.* 10, 12230

- 600 72 Flores-Tolentino, M. *et al.* (2020) Distribution and conservation of species is misestimated if
601 biotic interactions are ignored: the case of the orchid *Laelia speciosa*. *Sci. Rep.* 10, 9542
- 602 73 Ferrier, S. *et al.* (2007) Using generalized dissimilarity modelling to analyse and predict
603 patterns of beta diversity in regional biodiversity assessment. *Divers. Distrib.* 13, 252–264
- 604 74 Rosauer, D.F. *et al.* (2014) Phylogenetic generalised dissimilarity modelling: a new approach
605 to analysing and predicting spatial turnover in the phylogenetic composition of communities.
606 *Ecography* 37, 21–32
- 607 75 Mokany, K. *et al.* (2011) Combining α - and β -diversity models to fill gaps in our
608 knowledge of biodiversity: Filling gaps in biodiversity knowledge. *Ecol. Lett.* 14, 1043–1051
- 609 76 Guisan, A. and Rahbek, C. (2011) SESAM – a new framework integrating macroecological
610 and species distribution models for predicting spatio-temporal patterns of species
611 assemblages. *J. Biogeogr.* 38, 1433–1444
- 612 77 Hoskins, A.J. *et al.* (2019) Supporting global biodiversity assessment through high-
613 resolution macroecological modelling: Methodological underpinnings of the BILBI
614 framework. *Environ. Model. Softw.* 132, 104806
- 615 78 Allnutt, T.F. *et al.* (2008) A method for quantifying biodiversity loss and its application to a
616 50-year record of deforestation across Madagascar. *Conserv. Lett.* 1, 173–181
- 617 79 Prober, S.M. *et al.* (2012) Combining community-level spatial modelling and expert
618 knowledge to inform climate adaptation in temperate grassy eucalypt woodlands and related
619 grasslands. *Biodivers. Conserv.* 21, 1627–1650
- 620





MODEL OBJECTIVE SPACE

