



HAL
open science

MAIT Cell Development and Functions: the Microbial Connection

François Legoux, Marion Salou, Olivier Lantz

► **To cite this version:**

François Legoux, Marion Salou, Olivier Lantz. MAIT Cell Development and Functions: the Microbial Connection. *Immunity*, 2020, 53, pp.710 - 723. 10.1016/j.immuni.2020.09.009 . hal-03493882

HAL Id: hal-03493882

<https://hal.science/hal-03493882>

Submitted on 17 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

MAIT cell development and functions: the microbial connection

François Legoux^{1*}, Marion Salou^{1*} and Olivier Lantz^{1-3#}

- 1) INSERM U932, PSL University, Institut Curie, Paris, 75005, France.
- 2) Laboratoire d'immunologie clinique, Institut Curie, Paris, 75005, France.
- 3) Centre d'investigation Clinique en Biothérapie, Institut Curie (CIC-BT1428), Paris, 75005, France.

* equal contribution

Correspondence to:

Olivier Lantz: Laboratoire d'Immunologie and Inserm U932, Institut Curie, 26 rue d'Ulm, 75005 Paris, France.

Phone: +33 144324218

Fax: + 33 1 53 10 26 52

Email: Olivier.lantz@curie.fr

Short title: MAIT cells and bacteria

Abbreviations

5-A-RU: 5-amino-ribityl uracil

5-OE-RU: 5-(2-oxoethylideneamino)-6-D-ribitylaminouracil

5-OP-RU: 5-(2-oxopropylideneamino)-6-D-ribitylaminouracil

Ac-6-FP: acetyl-6-formyl-pterin

Areg: Amphiregulin

BM: Bone Marrow

CDR3: Complementary-Determining Region 3

DP: Double Positive

iNKT: invariant Natural Killer T

MAIT: Mucosal Associated Invariant T

MR1: MHC class I related molecule

PLZF: Promyelocytic Leukaemia Zinc Finger or Zbtb16: Zinc finger and BTB domain-containing protein 16

SAP: Slam Adaptor Protein

SPF: Specific Pathogen Free

TECs: Thymic Epithelial Cells

Tregs: regulatory T cells

Abstract (143 words)

Mucosal Associated Invariant T (MAIT) cells are an evolutionarily conserved T cell subset, which reacts to most bacteria through TCR-mediated recognition of metabolites derived from the vitamin B2 biosynthetic pathway. Microbiota-derived signals impact all stages of MAIT cell biology including intra-thymic development, peripheral expansion and functions in specific organs. In tissues, MAIT cells can integrate multiple signals and display effector functions involved in the defense against infectious pathogens. In addition to anti-bacterial activity, MAIT cells improve wound healing in the skin, suggesting a role in epithelium homeostasis through bi-directional interactions with the local microbiota. In humans, blood MAIT cell frequency is modified during several auto-immune diseases, which are often associated with microbiota dysbiosis further emphasizing the potential interplay of MAIT cells with the microbiota. Herein we will review how microbes interact with MAIT cells, from initial intra-thymic development to tissue colonization and functions.

(7961 words)

Introduction

Mucosal Associated Invariant T (MAIT) cells represent the most abundant T cell subset recognizing bacterial compounds (Franciszkiwicz et al., 2016). MAIT cell frequencies are very high in humans in the blood (1-10% of T cells), the liver (20-40%) and the lung and gut lamina propria (Dusseaux et al., 2011; Martin et al., 2009). They are much less abundant in mice (Rahimpour et al., 2015; Tilloy et al., 1999). They have been discovered almost 30 years ago following identification of an invariant T cell receptor- α (TCR α) chain (V α 7.2-J α 33) in CD4⁺CD8⁻ T cells from human blood (Porcelli et al., 1993). A T cell subset expressing a homologous TCR α chain has been then evidenced in mice and cattle (Tilloy et al., 1999), indicating evolutionary conservation and therefore important function(s). Selection by a non-polymorphic MHC class Ib molecule has also been demonstrated early on (Tilloy et al., 1999). This molecule has been identified as the evolutionarily-conserved MHC class I related molecule (MR1), which expression on hematopoietic cells is required for MAIT cell development (Treiner et al., 2003). MAIT cells have been rapidly characterized as T cells displaying immediate effector activities related to the expression of the master transcription factor PLZF (Zinc finger and BTB domain-containing protein 16, encoded by *Zbtb16*), also expressed by invariant natural killer T (iNKT) cells (Savage et al., 2008). MAIT cell dependency on commensal microbiota has been demonstrated as MAIT cells are absent from the periphery of germ-free mice (Treiner et al., 2003), indicating an intimate relationship between MAIT cells and bacteria. In the early 2010's, the ligand activating MAIT cells has been identified, first as a soluble microbial compound (Le Bourhis et al., 2010) and then as unstable vitamin B2 (riboflavin) precursor derivatives (Corbett et al., 2014; Kjer-Nielsen et al., 2012; Tastan et al., 2018). These molecules are only found in riboflavin-synthesizing bacteria and

yeasts (Mondot et al., 2016), conferring a broad anti-microbial reactivity to MAIT cells. This exquisite microbial specificity associated with the decrease of MAIT cell blood numbers during infections in humans (Gold et al., 2010; Grimaldi et al., 2014; Le Bourhis et al., 2013; Le Bourhis et al., 2010) and the protective effect of MAIT cells in several microbial infection experimental models ((Cui et al., 2015; Le Bourhis et al., 2013; Le Bourhis et al., 2010; Meierovics et al., 2013; Meierovics and Cowley, 2016; Wang et al., 2018), reviewed in (Godfrey et al., 2019; Salou et al., 2017)) support an anti-microbial function of MAIT cells.

Vitamin B2 metabolites produced by the microbiota travel through the body and control thymic development of MAIT cells (Legoux et al., 2019a). Microbial ligands may also be implicated in epithelial homeostasis, since MAIT cells promote tissue repair in response to TCR triggering (Constantinides et al., 2019; Hinks et al., 2019; Lamichhane et al., 2019; Leng et al., 2019). Thus, MAIT cells represent a direct way for the mammalian immune system to sense the microbiota, and to provide mucosal barrier protection. The role of vitamin B2 metabolites in tissue homeostasis is reminiscent of the way short chain fatty acids induce the expression of the Foxp3 transcription factor in T cells, with the acquisition of tissue-repair functions (Smith et al., 2013; Uchimura et al., 2018). Similarly, fMet peptides from some commensals expand H2-M3 major histocompatibility molecule-restricted T cells with both anti-microbial and tissue-repair functions (Harrison et al., 2019; Linehan et al., 2018). Although described only in primates, V γ 9 δ 2 γ δ T cells represent another subset of abundant, effector T cells with specificity for phospho-antigen metabolites, produced by microbes or stressed cells (Fournie and Bonneville, 1996). Thus, mammalian T cells have evolved several mechanisms for sensing the production of metabolites by the microbiota and for mounting a tissue-repair response in return.

Herein, we will first review the unique features of MR1 and MR1 ligands. We will then summarize the role of bacterial metabolites in the ontogeny and anti-microbial functions of MAIT cells. Finally, we will discuss the contribution of MAIT cells to tissue remodeling upon injury and how MAIT cell ligands together with other microbial signals may be involved.

1. MR1 and its ligands have unusual properties.

In contrast with classical MHC or CD1 molecules that present peptides or glycolipids, respectively, the only known MHC protein presenting metabolites to $\alpha\beta$ T cells is MR1. The *MR1* gene is absent from fish and birds, and appeared about 170 million years ago in the common ancestors of living marsupials and eutherians (Boudinot et al., 2016; Tsukamoto et al., 2013). *MR1* has been subsequently lost in several species, including *Lagomorphs* and all members of the *Carnivora* family (Boudinot et al., 2016). The MR1 protein shows remarkably high homology, with signs of purifying selection, indicating evolutionary pressure to maintain both the ligand binding groove and the external structure of the MR1 molecule (Boudinot et al., 2016; Huang et al., 2009). Associated with the absence of MR1 polymorphism, this suggests that the function of MR1 is to bind and present a limited number of ligands. Accordingly, only two molecules have

been described as strong agonists for MAIT cells, and both are derived from the vitamin B2 precursor 5-amino-ribityl uracil (5-A-RU) (Corbett et al., 2014; Soudais et al., 2015). The unstable 5-A-RU reacts with glyoxal or methylglyoxal, generated in the intermediary metabolism, and give rise to the short-lived adducts 5-(2-oxoethylideneamino)-6-D-ribitylaminouracil (5-OE-RU) and 5-(2-oxopropylideneamino)-6-D-ribitylaminouracil (5-OP-RU), respectively. Crystallographic studies reveal that both 5-OE-RU and 5-OP-RU establish a Schiff base with Lysine 43 of the MR1 groove, stabilizing these compounds (Corbett et al., 2014). Unless bound to MR1, 5-OP- and -OE-RU spontaneously cyclize and, though being still able to bind MR1, become weak agonists for MAIT cells (Kjer-Nielsen et al., 2012; Mak et al., 2017).

The MAIT cell TCR recognizes MR1:5-OP-RU complexes using a docking mode similar to mainstream T cells on classical MHC molecules, with a key role for the complementary-determining region (CDR) 3 of the TCR α chain in contacting the antigen (Awad et al., 2020; Gherardin et al., 2016; Reantragoon et al., 2012). CDR3 α sequences are composed of the junction of TRAV and TRAJ gene segments, with the stochastic addition or deletion of nucleotides in the TCR rearrangement process, resulting in extremely diverse sequences. Remarkably though, contact with 5-OP-RU is always mediated by the evolutionarily conserved Tyrosine 95 from the CDR3 of the TCR α chain. Indeed, in humans, cattle, pig (Xiao et al., 2019) and mice (Tilloy et al., 1999), the TCR α chain of MAIT cells is made of the evolutionarily conserved TRAV1 (official nomenclature of human V α 7.2) and TRAJ33 (and also TRAJ12 and TRAJ20 in humans) genes, associated with a CDR3 of fixed length (Reantragoon et al., 2013; Tilloy et al., 1999). Across mammals, the species which have lost the TRAV1 gene do not have a functional MR1 gene, which is either absent or pseudogenized (Boudinot et al., 2016). Conversely, when the *TRAV1* gene is present, the *MR1* gene is always functional. The CDR1 and CDR2 sequences of TRAV1 are also highly conserved when compared with other TRAV genes. Thus, strong selective pressure maintains TRAV1 and MR1 in a phylogenetic locked interaction, which is released when the TRAV1 locus is lost, as a consequence of the high variability of the TCR α locus during evolution (Mondot et al., 2016).

MR1 also binds a number of other small molecules, including other bacterial metabolites (Harriff et al., 2018), synthetic drugs (Keller et al., 2017a; Salio et al., 2020) and metabolites derived from human cell lysates (Lepore et al., 2017). Some MR1-bound metabolites can activate MAIT cells, but only at high concentrations. Of all the identified MR1 ligands, 5-OE-RU and 5-OP-RU exhibit a 100- to 1,000-fold higher potency than other MR1-binding molecules (Corbett et al., 2014; Harriff et al., 2018; Kjer-Nielsen et al., 2012).

In humans, very rare (< 1/1000 of MAIT cells) MR1:5-OP-RU reactive TRAV1^{neg} T cells also express PLZF (Zbtb16), hence their classification as MAIT-like cells (Koay et al., 2019a). In the four donors studied, these cells presented oligoclonal expansions and used TRAV36 associated with TRAJ34 or 37, and a CDR3 of constant length. Together with the use of a

predominant TCR β chain (TRBV28-TRBJ2.5), this suggests strong structural constraints to recognize MR1:5-OP-RU. However, no TRAV36 ortholog exists in mice indicating that this subset is not under strong selective pressure. In addition, a few TRAV1^{neg} mainstream T cells (i.e. not expressing PLZF), also called non-MAIT-like, recognize bacterial metabolites presented by MR1 with high affinity (Gherardin et al., 2016; Harriff et al., 2018; Meermeier et al., 2016). These cells are mostly CD8⁺, display a naive phenotype associated with a diverse repertoire. Finally, another TRAV1^{neg} TCR, identified in 2 individuals, recognizes a tumor-associated ligand presented by MR1 (Crowther et al., 2020). These T cells are neither activated by bacteria nor labelled with MR1:5-OP-RU tetramers. Interestingly, addition of the MR1-binding but non-agonist acetyl-6-formyl-pterin (Ac-6-FP) decreases their activation, suggesting the presence of an endogenous ligand (specific to tumor cells), with lower affinity than Ac-6-FP for MR1. In the absence of evolutionary conservation of a TCR shared between species, the physiological relevance of these TRAV1^{neg} MR1-restricted T cells is unclear.

Interestingly, a human TCR $\gamma\delta$ T cell subset can bind the alpha 3 domain of MR1 underneath the antigen binding groove (Le Nours et al., 2019). Although very rare (less than 1% of the $\gamma\delta$ T cells and 100-fold less frequent than MAIT cells), MR1-specific $\gamma\delta$ T cells may sense any MR1 ligand able to induce cell-surface expression of MR1, thereby potentially playing important roles in immune responses. Thus, MR1-specific $\gamma\delta$ T cells likely survey a much broader repertoire of metabolic ligands than MAIT cells, which are only activated by vitamin B2-derived metabolites. However, co-evolution of MR1 with TRAV1 strongly suggests that the main function of MR1 is to present antigens to TRAV1⁺ TCR $\alpha\beta$ MAIT cells. It is unclear whether MR1-specific $\gamma\delta$ T cells exist in other species.

The nature of MAIT antigens (i.e. small metabolites) likely affects the biology of MAIT cells. Indeed, in contrast with classical peptide or glycolipid antigens, 5-OP-RU applied onto the intact skin or delivered by oral gavage can be detected shortly after in the thymus (Legoux et al., 2019a). Hence, this metabolite can cross mucosal barriers, travel through the body and react with MR1 in the thymus (**Figure 1**). These results are consistent with known properties of other microbiota-derived metabolites, as microbial short chain fatty acids and tryptophan metabolites can reach distant host tissues such as the brain, thymus and foetus (Gomez de Agüero et al., 2016; Uchimura et al., 2018). Antibodies facilitate stability or transport of microbial metabolites (such as tryptophan metabolites) through the host (Gomez de Agüero et al., 2016). However, given that MAIT cell development requires 5-OP-RU to reach the thymus, and that MAIT cells develop normally in mice lacking B cells (Legoux et al., 2019a), antibodies are likely dispensable for 5-OP-RU transport. Whether 5-OP-RU requires a carrier in blood is unclear. The dynamic properties of 5-OP-RU *in vivo* may also affect the initiation and maintenance of MAIT cell responses (see below).

2. MAIT cell development is controlled by host intrinsic signals

MAIT cells develop in the thymus (Koay et al., 2016; Tilloy et al., 1999) in a process recently reviewed in details (Pellicci et al., 2020). Following positive selection, MAIT cells undergo intra-thymic differentiation and, contrary to mainstream T cells, acquire immediate effector functions associated with the related phenotype (Koay et al., 2016; Rahimpour et al., 2015) and the ability to migrate into non-lymphoid tissues (Salou et al., 2019). These original characteristics, shared with iNKT cells, are related to the expression of the master transcription factor PLZF (*Zbtb16*) during thymic development (Koay et al., 2016; Savage et al., 2008). PLZF (*Zbtb16*) is a known suppressor of the naïve T cell program and inducer of effector genes in iNKT cells (Mao et al., 2016; Savage et al., 2008). MAIT and iNKT subsets display highly similar transcriptomic programs once split according to expression of T-bet or ROR γ t, which define the iNKT1, MAIT1 and iNKT17, MAIT17 effector subsets, respectively (Salou et al., 2019). Accordingly, as for iNKT cells, positive selection of MAIT cells in mice strictly depends on the expression of the Signaling lymphocytic activation molecule (SLAM) Adaptor Protein (SAP), which signals downstream of SLAM molecules (Koay et al., 2019b; Legoux et al., 2019b). SLAM signaling can result in both inhibitory and stimulatory effects on the development of iNKT cells, but the molecular pathways involved are unclear (Lu et al., 2019; Tuttle et al., 2018; Zhao et al., 2012). Hence, positive selection of effector MAIT cells in mice requires signaling from the TCR and concomitant engagement of SLAM molecules at the surface of hematopoietic thymocytes (Legoux et al., 2019b) (**Figure 2**).

Bone marrow (BM) chimera experiments demonstrated that TRAV1⁺ T cells recognizing MR1:5-OP-RU complexes can be selected by MR1 expressed on thymic epithelial cells (TECs) (Legoux et al., 2019b). TECs do not express SLAM molecules, and thymocytes undergoing positive selection on TECs differentiate into naïve CD4⁺ T cells, which patrol secondary lymphoid organs but do not migrate into non-lymphoid tissues (**Figure 2**). In mice, we estimated that about half of the immature thymocytes specific for MR1:5-OP-RU are selected on SLAM⁺ hematopoietic cells, the other half being selected on Slam⁻ TECs (Legoux et al., 2019b).

Following positive selection on hematopoietic cells (Seach et al., 2013), MAIT precursors down-regulate CD24a and up-regulate *Klf2*, *Sell* and *S1pr1* (Legoux et al., 2019b), in a process similar to differentiation of mainstream T cells preparing for thymus egress (Weinreich and Hogquist, 2008). Some of these MAIT precursors may exit the thymus at this stage, as proposed for iNKT cells (Wang and Hogquist, 2018). Early thymic egress of MAIT cells is also suggested by the presence of MR1:5-OP-RU-specific T cells with remaining GFP expression in the spleen and lymph nodes of RAG-GFP mice (Winter et al., 2019). Following CD24a down-regulation, MAIT precursors begin to express the transcription factor PLZF (*Zbtb16*), with acquisition of high amounts of CD44 and subsequent effector differentiation into T-bet- or ROR γ t-expressing cells subsets (Legoux et al., 2019b; Salou et al., 2019). Effector differentiation requires *Satb1* and *Cxcr6* (Koay et al., 2019b). Although the sequence of molecular events associated with MAIT effector (PLZF (*Zbtb16*)⁺) differentiation is emerging (Koay et al., 2019b; Legoux et al., 2019b),

the mechanisms controlling MAIT1 versus MAIT17 lineage choice remain obscure. TCR avidity for the selecting ligand appears involved in iNKT1, 2 and 17 commitment, with stronger signaling for iNKT2 and iNKT17 in comparison with iNKT1 subsets (Tuttle et al., 2018). Given the similarity between iNKT and MAIT cell development, TCR avidity for the selecting antigen probably determines MAIT1 versus MAIT17 choice. Consistent with a role of TCR signaling during MAIT cell development, MAIT cell numbers are decreased in *miR181^{-/-}* (Winter et al., 2019) and in *Zap70*-hypomorphic (SKG) mice (Koay et al., 2019b). Studying MAIT1 and MAIT17 proportions in these mice could be informative on the role of TCR avidity in MAIT subset commitment since the iNKT2 subset, which requires stronger TCR signal for selection, is absent (Tuttle et al., 2018). The role of TCR strength is also supported by the lower proportion of MAIT17 cells in the thymus of *Ccr7^{-/-}* mice, which exhibit reduced TCR signaling (Davalos-Misslitz et al., 2007; Koay et al., 2019b).

Whether the developmental process of MAIT cells is conserved between mice and humans is unclear. Although some orthologous transcription factors are induced in both species upon thymic development (i.e. *ZBTB16*, *RORC*, *TBX21*), human MAIT cells exit the thymus as T-bet⁺RORγ⁺ cells while mouse MAIT cells express either transcription factors. In addition, SAP-deficient patients have a normal frequency of blood MAIT cells, and these cells express normal amounts of PLZF (*ZBTB16*) (Martin et al., 2009) and Legoux and Lantz, unpublished) suggesting additional mechanisms by which MAIT cells may differentiate into effector cells in humans.

3. MAIT cell development is controlled by the microbiota

The absence of MAIT cells in the mesenteric lymph nodes of germ-free mice has been recognized early on using RT-qPCR for the MAIT iTCRα chain mRNA (Treiner et al., 2003). Reduced MAIT cell frequency has been then reported in the thymus of germ-free mice using MR1:5-OP-RU tetramers (Koay et al., 2016). As parabiotic experiments demonstrate that MAIT cells do not circulate from the periphery to the thymus (Legoux et al., 2019a), the effect of microbiota on MAIT cells is likely happening directly in the thymus. Both the number of mature MAIT17 cells and of CD24⁺ immature MR1:5-OP-RU tetramer-labelled T cells are reduced in germ-free mice. The residual tetramer-positive cells have reduced TCR signaling and proliferation as compared with specific pathogen free (SPF) mice, suggesting that the microbiota provides a TCR ligand for thymic MAIT cell selection and proliferation. Experiments with germ-free mice mono-colonized with *E. coli* strains mutated in the vitamin B2 pathway either up-stream (*RibD*-deficient) or downstream (*RibE*-deficient) of MAIT antigen production establish that thymic MAIT cell development is directly dependent upon *RibD* gene expression and 5-A-RU production by bacteria (Legoux et al., 2019a). Both immature CD24⁺ precursors and mature MAIT17 cells are restored upon microbial colonization, which suggests *de novo* positive selection of MAIT cells by a microbial antigen. Alternatively, given that residual MAIT cells exist in germ-free mice (discussed below), the microbial ligand might impact MAIT cell development mostly through

expansion of MAIT17 cells selected beforehand on a self-antigen or even on empty MR1 molecules. The 5-A-RU-derived 5-OP-RU antigen rapidly travels from mucosal surfaces (skin or stomach) to the thymus, where it induces expansion of MAIT cells. A variety of thymic cells can capture exogenous 5-OP-RU, including TECs, dendritic cells and CD4⁺CD8⁺ double positive (DP) thymocytes, but the latter cells express the highest amounts of MR1 (Legoux et al., 2019a; Seach et al., 2013). Interestingly, only MAIT17 cells - but not MAIT1 cells - respond to microbial colonization and expand in the thymus (**Figure 2**).

Importantly, positive selection of MAIT cells takes place in germ-free mice, as indicated by the presence of small numbers of mature, cytokine-producing MAIT1 and MAIT17 thymocytes in these mice (Legoux et al., 2019a). By contrast, *Mr1*^{-/-} mice are completely devoid of mature MAIT cells, which indicates that MR1, either unloaded or loaded with unidentified self-ligand(s), mediates positive selection of MAIT cells. MR1-restricted self-metabolites in the thymus may be identical, or distinct, to the MR1 ligands suggested but not yet identified in human cell lysates (Lepore et al., 2017) or involved in the human MR1-dependent recognition of tumour cells by a T cell clone (Crowther et al., 2020).

Intriguingly, microbiota colonization of adult mice restores thymic MAIT cell development, but newly generated MAIT cells do not populate peripheral tissues such as lung (Legoux et al., 2019a) and skin (Constantinides et al., 2019). MAIT and iNKT subset frequencies in mice are highly correlated in different tissues (Salou et al., 2019), as are iNKT and MAIT cell frequencies in human cord blood (Ben Youssef et al., 2018). This suggests a competition for peripheral niches between T cell subsets expressing similar effector programs, such as iNKT or $\gamma\delta$ T cells. The increased MAIT cell frequency in the skin of $\gamma\delta$ T cell-deficient mice or in the spleen of *Cd1d*^{-/-} mice supports this hypothesis (Constantinides et al., 2019; Koay et al., 2016). Alternatively, the tissues or its endothelium may be permissive for colonization only for a short period after birth, as previously suggested for conventional T cells (Alferink et al., 1998). Finally, early life colonization may produce MAIT cells with different tissue-homing properties. Interestingly, 5-OP-RU alone is not sufficient for MAIT cell development when administered to adult mice (Constantinides et al., 2019; Legoux et al., 2019a), but enables MAIT cell development and skin homing in neonates (Constantinides et al., 2019). Thus, in addition to 5-A-RU-derived antigens, microbiota-induced signals are necessary to support MAIT cell maturation in the thymus of adults, but not in neonates. MAIT cell development is not affected in MyD88-deficient mice (Legoux et al., 2019a), which excludes a role for TLR signals and for IL-1, IL-18 and IL-33. IL-23 alone is not sufficient to induce thymic MAIT cell development (Constantinides et al., 2019), but may act in conjunction with 5-OP-RU to control MAIT cell development in adult SPF mice.

In humans, mature T cell development starts in utero, and V α 7.2⁺CD161⁺ T cells are found in the thymus at early age. As 5-OP-RU cross mucosal barriers in mice and microbial metabolites

can influence the development of innate lymphoid cells in utero (Gomez de Agüero et al., 2016), vitamin B2 metabolites produced by the microbiota of the mother might be able to reach the fetal thymus and contribute to MAIT cell selection and expansion. MR1:5-OP-RU-specific thymocytes identified with tetramer express high amounts of PLZF (*Zbtb16*), CD161 and IL-18R α (Koay et al., 2016), demonstrating intra-thymic acquisition of PLZF (*Zbtb16*). Contrary to mouse MAIT cells that become fully mature and memory in the thymus, only the most mature MAIT cells in the human thymus express CD161 and IL-18R α (Martin et al., 2009) as well as PLZF (*Zbtb16*) (Koay et al., 2016; Koay et al., 2019b; Martin et al., 2009). Accordingly, thymic MAIT cells exhibit many features of adult MAIT cells with expression of tissue homing chemokine receptors such as CCR6, and CXCR6. However, thymic mature and cord blood MAIT cells remain negative for the memory marker CD45RO (Ben Youssef et al., 2018; Martin et al., 2009). Still, thymic mature MAIT cells are able to secrete TNF- α and IFN- γ after PMA+ionomycin stimulation, but to a lesser extent than adult blood MAIT cells (Koay et al., 2016). According to RNAseq analysis, the cytotoxic capacity seems to be similarly acquired in the thymus (Koay et al., 2019b). This suggests that additional events occurring in the periphery, such as microbiota encounter, might be pivotal for the final maturation steps.

In cord blood, about 1% of T cells express high amounts of CD161 and IL-18R α as well as *Zbtb16* and *Rorc* (Walker et al., 2012), which are strongly associated to the MAIT differentiation program. However, only a minority uses the V α 7.2 TCR α chain among which less than 10% is labelled with MR1:5-OP-RU tetramers (Ben Youssef et al., 2018) suggesting that a large set of T cells with various specificities acquires a MAIT-like differentiation program in the thymus during fetal life, possibly through selection on DP thymocytes expressing various endogenous ligands (Lantz and Legoux, 2018). Notably, only the MR1:5-OP-RU tetramer-labelled T cells acquire a memory phenotype in the few weeks following birth and subsequently expand in 5 to 6 years to reach adult frequencies (Ben Youssef et al., 2018; Gherardin et al., 2018b; Koay et al., 2016). Among CD161^{hi} T cells, only iNKT and MAIT cells are found in adult blood (Ben Youssef et al., 2018). Associated with the oligoclonality of adult MAIT cells (Lepore et al., 2014; Tilloy et al., 1999), these data suggest that only T cells with the highest avidity for 5-OP-RU expand, which depends on their particular TCR β chain (Awad et al., 2020; Eckle et al., 2014). The other CD161^{hi} T cells which do not recognize microbial ligands presented in the right context would disappear or be diluted out by MAIT cells. Thus, like for mice in the thymus, the microbial ligands and maybe other microbiota-derived signals are key for the development of human MAIT cells in the periphery.

We hypothesize that the different locations (thymus versus periphery) where the microbial ligands impact MAIT cell development in mice versus humans is related to body size. As microbiota-produced 5-OP-RU seems to freely diffuse in the body, and is highly unstable unless

bound to MR1, its concentration in fluids is directly related to the rate of diffusion from the microbiota. This rate is proportional to mucosal surface, but the resulting concentration is also inversely proportional to the volume of distribution. Thus, in the absence of specific transporters or epithelial layer disruption, 5-OP-RU concentration should be inversely correlated with body size: $\text{concentration} = k \times \text{surface (size}^2) \div \text{volume (size}^3) = k \div \text{size}$ where k is the production rate per surface unit. Thus, the ability of 5-OP-RU to reach the thymus is probably higher in mice than in humans, which would rely more on endogenous ligands. The analysis of MAIT cell development in the thymus of phylogenetically close species with different body size may resolve this issue.

4. MAIT cells reside in peripheral tissues at steady state

MAIT cells are present in non-lymphoid tissues in both humans and mice, with highly variable percentages between individuals. The microbiota seems to be a strong determinant of MAIT cell numbers at steady state as the percentage of skin and lung MAIT cells is strongly cage-dependent in mice (Constantinides et al., 2019). While MAIT1 cells localize preferentially to the spleen, lymph nodes and liver (Rahimpour et al., 2015; Salou et al., 2019), MAIT17 cells also home to barrier tissues such as lung, skin, and gut (**Figure 3**), probably in relation with the differentiation program associated with ROR γ t vs T-bet expression.

Tissue tropism seems to be an intrinsic feature of the MAIT effector program. Indeed, in mice, both MAIT1 and MAIT17 thymocytes express gene signatures associated with tissue residency, rather than blood circulation (Salou et al., 2019). Thymic MAIT cells express the *Runx3*-related transcription program (Milner et al., 2017; Salou et al., 2019) controlling the establishment of conventional tissue-resident memory CD8⁺ T cells. Accordingly, as compared to single-positive thymocytes, thymic MAIT cells preferentially seed non-lymphoid organs such as lung and liver upon adoptive transfer (Salou et al., 2019). In B6 mice, MAIT cells represent only 0.03% of T cells in the thymus, but up to 1-3% of T cells in the liver, lung and gut lamina propria (Rahimpour et al., 2015). The MAIT program alone may be sufficient for tissue colonization as antigen recognition in lymph-nodes is dispensable for the recruitment of conventional memory CD8⁺ T cells into non-lymphoid tissues (Mackay et al., 2012). Accordingly, MAIT cells expand normally following *S. epidermidis* association in mice devoided of peripheral lymph nodes (*Lta*^{-/-}) (Constantinides et al., 2019). Still, BM chimera using *Mr1*⁺ donors into *Mr1*^{-/-} recipients results in decreased numbers of MAIT cells in the lungs (Wang et al., 2019), but this could also be explained by impaired thymic exit in *Mr1*^{-/-} recipients. Similarly to iNKT cells (Thomas et al., 2011), very few MAIT cells found in spleen, liver and lung (except lung MAIT1) recirculate between 5-week long parabiotic pairs (Salou et al., 2019), which indicates that MAIT cells are long-term resident cells in these tissues. In spleen and liver, MAIT1 cell residency relies on LFA1 and ICAM1 interactions (Salou et al., 2019). In spleen and lung, MAIT17 cells are located in the

extravascular space and do not rely on these interactions. In skin, while IL-23 is necessary for MAIT17 development or maintenance (Constantinides et al., 2019), the mechanisms of tissue retention are unknown.

In humans, the liver MAIT transcriptome is enriched in a residency signature (Milner et al., 2017) as compared to blood, suggesting that the tissue resident property is shared with mouse MAIT cells (Salou et al., 2019). In the buccal mucosa, MAIT cells express the residency markers CD103 and CD69 (Lu et al., 2020; Sobkowiak et al., 2019). However, MAIT cells from matched thoracic duct lymph and blood samples have a shared TCR repertoire, suggesting recirculation between compartments (Voillet et al., 2018). As MAIT cells in the blood lack CCR7 expression, the more likely scenario is migration from blood to tissues and then to the lymph and back to the blood, similarly to classical effector memory T cells (Beura and Masopust, 2014). As in mice, human MAIT cells appear targeted to tissues during thymic development, since tissue-homing molecules such as *CCR6*, *CCR5* and *CXCR6* are expressed in thymic MAIT cells (Lantz and Legoux, 2019; Salou et al., 2019). *RUNX3* and *CEBPD* (see below) are also upregulated in the more mature human thymic MAIT cells (Koay et al., 2019b). Accordingly, human MAIT cells are abundant in the lung (2-4%), liver (20-50%) and intestine (up to 60% of CD4⁺ T cells in the jejunum) (Dusseaux et al., 2011; Kurioka et al., 2016). Thus, as for mouse MAIT cells, the tissue-homing phenotype of human MAIT cells is acquired in the thymus.

Whether MAIT cells are homogeneous or not can be debated, as studied for other resident cells with similar effector functions such as innate lymphoid cells (Gury-BenAri et al., 2016). In mice, two single cell RNA-seq datasets suggest minimum heterogeneity in the thymus, once MAIT1 and MAIT17 subsets are separated (Koay et al., 2019b; Legoux et al., 2019b). In humans, blood MAIT cells appear very homogenous with high expression of IL-18R α , CD161 and CD26 allowing their identification by cytometry (Dusseaux et al., 2011; Martin et al., 2009). Together with anti V α 7.2 staining, these markers faithfully identify MR1:5-OP-RU tetramer-positive MAIT cells except in a few cases (Gherardin et al., 2018a). In tissue, single-cell RNAseq on broncho-alveolar lavage MAIT cells from pneumonia patients identifies three MAIT subpopulations with varying expression of *RORC*, *TBX21* and *GATA3*, but still transcriptionally very close (Lu et al., 2020). In the absence of comparison to reference cell populations such as naive and memory mainstream T cells, the biological meaning of distances on the UMAP graph is unclear (Kiselev et al., 2019). Similarly, cytometry analysis of 332 immune receptors on human blood MAIT cells revealed limited heterogeneity despite bimodal expression of few markers such as CD56, CD84 and CD94. High expression of these markers is associated with higher lymphokine secretion after IL-12 and IL-18 stimulation (Dias et al., 2017).

Altogether, MAIT cells from mice and humans are programmed in the thymus to seed and reside in barrier tissues. Once there, local cues probably modulate the MAIT transcription

program, as MAIT cells isolated from different organs express distinct effector gene sets ((Salou et al., 2019) and Salou and Lantz, unpublished observations). Thus, MAIT cells seem to display specific properties according to the organ, suggesting that their function is tailored to the tissue, as previously proposed for mainstream T cells (Matzinger and Kamala, 2011).

5. MAIT cells are protective during various infectious diseases

The strong evolutionary conservation of the MAIT-MR1 axis indicates important functions for MAIT cells. These functions could be related to particular properties of MAIT cells: secretion of specific effector molecules, antigen specificity, positioning in specific tissues or ability to be directly activated by certain cytokines. Rapid production of IFN- γ , TNF- α , IL-17, GM-CSF and cytotoxic molecules is a key characteristic of MAIT cells, shared with mainstream memory T cells (reviewed in (Franciszkiewicz et al., 2016; Godfrey et al., 2019; Salou et al., 2017)) (**Figure 4**). Rapid response may rely on preformed RNAs instead of de novo transcription, as shown by PMA-Ionomycin-induced IFN- γ secretion experiments in which transcription or translation is inhibited (Gutierrez-Arcelus et al., 2019). MAIT cells are able to detect the presence of bacteria through their TCR and they also express various cytokine receptors such as receptors for IL-18, IL-12, IL-7, IL-1, IL-15 and IL-23 (Franciszkiewicz et al., 2016). The integration of these signals modulates the pattern of secreted effector molecules, resulting in context-dependent functions (**Figure 4**). For instance, IL-23R-deficient mice have reduced expansion of MAIT cells upon lung bacterial infection (Wang et al., 2019). Recently, type 1 IFN and the gut-derived TL1A have been added to the list of cytokines modulating MAIT cell functions (Lamichhane et al., 2020; Leng et al., 2019).

Because of their antigen specificity, the involvement of MAIT cells in infections by 5-A-RU-producing bacteria has been extensively investigated, both in mice and humans. Briefly, the absence of MAIT cells results in defective or delayed clearance of several intra-cellular bacteria such as *Mycobacterium abscessus* (Le Bourhis et al., 2010), *M. bovis* BCG (Chua et al., 2012), *Francisella tularensis* (Meierovics et al., 2013; Meierovics and Cowley, 2016) and *Legionella longbeachae* (Wang et al., 2018).. MAIT cells are also important for the clearance of urinary tract infection (Cui et al., 2015) or during *E. coli* peritonitis (Georgel et al., 2011). In humans, the frequency of MAIT cells in the blood is decreased in several bacterial infections (Grimaldi et al., 2014; Le Bourhis et al., 2010) with concomitant increase at the infected sites in few reports (reviewed in (Godfrey et al., 2019; Salou et al., 2017)). Longitudinal follow-up of human blood samples after *Salmonella enterica* Typhi infection or a *Shigella* strain vaccine challenge support the hypothesis of a migration towards inflammatory sites, as blood MAIT cells decrease in numbers, appear activated, and exhibit clonal expansion (Grimaldi et al., 2014; Howson et al., 2018; Le Bourhis et al., 2010; Salerno-Goncalves et al., 2017). MAIT activation is also associated with increased CCR6 and CCR9 expression, which favor homing to inflamed tissues and gut, respectively. Moreover, C/EBP δ , important for transmigration into inflamed tissues (Lee et al., 2018), is highly expressed by human MAIT cells. However, whether the main

function of MAIT cells is the defense against microbial pathogens remains unclear. Indeed, no susceptibility to a particular bacterial pathogen has yet been ascribed to a genetic deficiency of the MAIT-MR1 axis, besides a polymorphism of the *Mr1* gene in a Vietnamese cohort that has been associated to disseminated tuberculosis susceptibility (Seshadri et al., 2017).

Interestingly, MAIT cells seem to be involved in the defense against viruses. MAIT cell frequencies are selectively decreased in the blood of patients during several viral infections ((Loh et al., 2016; van Wilgenburg et al., 2016) reviewed in (Godfrey et al., 2019; Ussher et al., 2018)) and MAIT cells protect from experimental influenza infection, with an increased mortality in *Mr1*^{-/-} animals without, however, modifications in the viral load (Wilgenburg et al., 2018). As viruses do not produce 5-A-RU, TCR-independent stimulation of MAIT cells is probably pivotal in these settings. Accordingly, MAIT cells are activated *in vitro* by the IL-12 and IL-18 secreted by influenza-infected monocytes (Loh et al., 2016).

The mechanisms of MAIT cell anti-infectious immunity are still unclear. *In vitro*, MAIT cells kill infected cells through granzymes and perforin (Kurioka et al., 2015; Le Bourhis et al., 2013). MAIT cells produce IFN- γ , TNF- α , IL-17 and (G)M-CSF (Franciszkievicz et al., 2016; Godfrey et al., 2019; Salou et al., 2017), which likely stimulates anti-infectious responses through myeloid cells. MAIT cells also secrete various proinflammatory chemokines (Hinks et al., 2019; Lamichhane et al., 2019), suggesting a role in orchestrating the immune response. The loss of protection imparted by IFN- γ -deficient MAIT cells transferred into MAIT-deficient animals suggests that IFN- γ is crucial for anti-*Legionella* and -influenza immunity (Wang et al., 2018; Wilgenburg et al., 2018). However, in these studies, MAIT cells were expanded through *Salmonella* infection of the donor animals, to obtain the large numbers of MAIT cells required for adoptive transfers. This procedure induces a ROR γ ⁺T-bet⁺ profile (Chen et al., 2017), distinct from the predominant ROR γ ⁺T-bet^{neg} profile at steady state but more similar to the phenotype of human MAIT cells. In *F. tularensis* infection, MAIT cells improve bacterial clearance and mainstream T cell recruitment (Meierovics and Cowley, 2016). Differentiation of monocyte-derived macrophages is delayed in *Mr1*^{-/-} mice, probably due to the absence of early GM-CSF production by MAIT cell. Transfer of monocyte-derived macrophages in *Mr1*^{-/-} animals restored CD4⁺ T cell recruitment (Meierovics and Cowley, 2016). Importantly, when fighting bacteria, MAIT cells release inflammatory mediators able to induce tissues damage, such as Granzyme B and K, implicated in killing cells infected by intracellular bacteria (Kurioka et al., 2015; Ussher et al., 2014; Ussher et al., 2016), and IL-4i (leading to the formation of H₂O₂) (Turtle et al., 2011).

Thus, the exact mechanisms, direct or indirect, leading to MAIT cell-dependent protection against infectious agents remain to be fully dissected. Recent analysis of MAIT transcriptome following various activation strategies unraveled a previously unappreciated tissue repair function (see below).

6. MAIT cells are involved in both anti-microbial immunity and tissue repair

When compared to cytokine-activated or resting conditions, *in vitro* TCR-mediated activation of human blood MAIT cells results in an up-regulation tissue repair mediators, including *IL22*, *AREG*, *VEGFA* and *TGFA*, together with inflammatory molecules (Hinks et al., 2019; Lamichhane et al., 2019; Leng et al., 2019). Accordingly, supernatant from human MAIT cells cultured with *E. coli*-infected THP1 cells promotes closure in an *in vitro* scratch assay (Leng et al., 2019). Stimulation of human blood MAIT cells with IL-2, IL-7, mitogen and CD28 for a long period (8 to 10 days) induces high amounts of IL-13 and IL-5 (Kelly et al., 2019), two cytokines involved in tissue repair (Van Dyken and Locksley, 2013). A tissue-repair signature is also enriched in mouse lung MAIT cells following *Legionella* infection (Hinks et al., 2019) and in skin MAIT cells from *S. epidermidis*-associated animals (Constantinides et al., 2019). A role for MAIT cells in wound healing has been directly demonstrated by measuring the epidermal tongue length 5 days after a punch biopsy of the skin, with faster wound healing in C57Bl/6 mice pretreated topically with 5-OP-RU (Constantinides et al., 2019). The same beneficial effect of MAIT cells has been observed in mice associated with *S. epidermidis* prior to wounding, but only in $\gamma\delta$ T cell-deficient mice, suggesting redundancy between $\gamma\delta$ T cells and MAIT cells in the skin (Constantinides et al., 2019). At steady state, MAIT cells in the lung express more tissue repair mediators than in liver or spleen (Salou et al., 2019). This is also true for human MAIT cells in liver as compared to blood (Salou et al., 2019). MAIT cells may also regulate intestinal epithelium tissue integrity, as gut permeability to FITC-Dextran after gavage is increased in NOD mice deficient for MR1 (Rouxel et al., 2017).

Apart from these cytokines, and the putative mediators of tissue repair described for commensal-specific H2-M3-restricted CD8⁺ T cells (Linehan et al., 2018), how MAIT cells repair tissues remains unknown. One important effector molecule in several tissue-repair models is amphiregulin (Areg). Areg is involved in the tissue repair function of Foxp3⁺ regulatory T cells (Tregs) following influenza infection or muscle injury (Arpaia et al., 2015; Burzyn et al., 2013). MAIT cells may influence various pathways to promote repair including mediators regulating angiogenesis or tissue repair according to a MAIT transcriptome analysis (Hinks et al., 2019).

Consistent with the production of pro-angiogenesis molecules and epithelial growth factors, MAIT cells may have a pro-tumoral role (Yan et al., 2020). While the described mechanism is impairment of IFN- γ secretion by NK and CD8⁺ T cell, the MAIT tissue repair function is also likely involved. This has been previously described for Treg cells (Green et al., 2017) or $\gamma\delta$ T cells (Jin et al., 2019). In three different models of transplanted tumors, *Mr1*^{-/-} mice develop reduced numbers of metastases, a phenotype reversed by MAIT cell adoptive transfer (Yan et al., 2020). The pro-tumoral effect of MAIT cells requires MR1 expression on the tumor cells. *In vivo*, the nature and the origin (endogenous or bacterial) of the MAIT ligand presented by tumor cells remains unknown, although tumoral colonization by bacteria is a possibility (Nejman et al., 2020). Similarly, exacerbated or uncontrolled tissue-repair function could have unwanted effects such as the

fibrotic gastritis induced by *H. pylori* infection once MAIT cells have been expanded with 5-OP-RU and TLR ligands (D'Souza et al., 2018). MAIT cells could also promote liver fibrosis (Bottcher et al., 2018; Hegde et al., 2018), maybe through direct activation by bacterial products reaching the liver by the portal vein (Riva et al., 2017). Similarly, MAIT cells have been implicated in kidney fibrosis (Law et al., 2019).

Thus, MAIT cells are able to exert various effector functions (**Figure 4**), consistent with the co-expression of several master transcription factors such as ROR γ t, T-bet, Helios and Eomes in humans (Leeansyah et al., 2015). The expression of these usually mutually exclusive transcription factors is reminiscent of the commensal-specific H2-M3-restricted CD8⁺ T cells populating the mouse skin following *S. epidermidis* colonization (Harrison et al., 2019; Linehan et al., 2018). In MAIT cells, the factors triggering one effector function over others are beginning to be unraveled as described in the next section.

7. MR1 ligands activate MAIT cells at steady state and during diseases

Recent *in vitro* evidence suggests a pivotal role of MAIT TCR signaling in eliciting tissue-repair function (Hinks et al., 2019; Lamichhane et al., 2019; Leng et al., 2019). Given that most microbes produce 5-A-RU (Mondot et al., 2016), and as microbiota-derived 5-OP-RU travels rapidly into the organism (Legoux et al., 2019a), tissue repair functions are likely constantly triggered (**Figure 4**). Therefore, MAIT cells would be involved both in tissue homeostasis, and recovery of tissue function after infection with 5-A-RU-producing pathogens. Infections with 5-A-RU-deficient pathogens might still activate MAIT cells via the commensal microbiota-derived 5-OP-RU. In thymic MAIT cells, microbiota-derived 5-OP-RU triggers MAIT17- but not MAIT1-proliferation, suggesting that MAIT17 may be the main actors of reparation.

When and in which circumstances the MAIT TCR is triggered *in vivo* is unclear. MR1 expression might not be high enough for MAIT cell activation in peripheral tissues, or MR1 may be loaded with non-agonist ligands. Indeed, DP thymocytes express 2-3 times more *Mr1* RNA than the alveolar macrophages and 5-6 times more than other cells in the body (Heng et al., 2008). MR1 is mostly intra-cellular at steady state and ligand availability seems to be the limiting factor for cell surface translocation (McWilliam et al., 2016). In HSV-1 infection, MR1 expression is suppressed and the protein is targeted for proteosomal degradation, preventing MR1 surface expression (McSharry et al., 2020). Similar mechanisms might occur at steady state or during other infections to prevent MAIT cell activation. Another layer of regulation may lie in the presence of other MR1 ligands, derived or not from bacteria, and binding MR1 without activating MAIT cells (Awad et al., 2020; Eckle et al., 2014; Harriff et al., 2018; Keller et al., 2017b; Soudais et al., 2015). Poor or null agonists, such as 5-OP-RU derived lumazines, may also regulate MAIT cell activation by preventing MR1 binding to stronger MAIT agonists. Since the relative proportion of these agonists or antagonists may vary according to microbiota composition and metabolic status (Schmaler et al., 2018), this would allow MAIT cells to sense microbiota ecology.

Whether TCR triggering is required for MAIT functions in pathological settings remains unclear as MAIT cells can be activated independently of TCR stimulation, through cytokines, either alone or in combination as reviewed elsewhere (Franciszkiewicz et al., 2016; Godfrey et al., 2019). In iNKT cells, TCR expression is dispensable for IFN- γ secretion in response to LPS stimulation *in vivo* (Vahl et al., 2013). TCR expression is also dispensable to maintain the iNKT cell lineage identity, including PLZF (*Zbtb16*) expression, in peripheral tissues. As MAIT and iNKT subsets have an almost identical transcriptional program (Salou et al., 2019), MAIT cell IFN- γ production may similarly be independent of TCR triggering. Moreover, for the Treg cells implicated in lung repair following influenza infection, Areg secretion is TCR-independent (Arpaia et al., 2015). In contrast, TCR signaling induces MAIT responses in several settings. Blocking MR1 decreases the number of metastasis, as well as NK and CD8 T cell secretion of IFN- γ (Yan et al., 2020). TCR triggering seems necessary but not sufficient for MAIT cell proliferation: 5-OP-RU inhalation increases MAIT cell numbers in the lungs only when associated with TLR 2, 3, 6 or 9 ligands (Chen et al., 2017). 5-OP-RU, together with IL-23, drives MAIT cell expansion and improves protection against legionella infection (Wang et al., 2019). Similarly, 5-A-RU-deficient salmonella does not induce MAIT cell proliferation, while 5-OP-RU addition does (Chen et al., 2017). Finally, deletion of MR1 before *S. epidermidis* association prevents MAIT cell increase in the skin (Constantinides et al., 2019). In this model, IL-18 blockade demonstrates an additional IL-18-dependency for the proliferation, while IL-17 secretion required IL-1 (Constantinides et al., 2019). Thus, TCR triggering seems important for proliferation following infection and for the protumoral effect of MAIT cells. However, as MAIT cells can be triggered either through their TCR or by microbial-induced lymphokines, the need for bacterial ligands is still unclear for MAIT cell control of bacterial infection.

This issue is also complicated by the possible difference in TCR or lymphokine dependency of MAIT1 versus MAIT17 subsets as most of the *in vivo* data were generated in barrier organs in which MAIT17 cells are the great majority. MAIT1 vs MAIT17 activation requirements and subsequent effector functions are likely very different as MAIT1 cells express more NK receptors and cytotoxic molecules than MAIT17 cells. In the thymus, triggering MAIT1 TCR results in lower Nur77 expression in comparison to MAIT17 cells (Legoux et al., 2019a; Legoux et al., 2019b). Thus, in contrast to MAIT17 cells that mainly express tissue repair mediators following TCR stimulation, MAIT1 cells may preferentially respond to danger signals and produce cytotoxic molecules. However, in a pathological context in which IL-1, IL-12, IL-18 as well as MAIT ligands are present, MAIT17 cells upregulate T-bet, and secrete type-1 effector molecules (IFN- γ or TNF- α) together with tissue repair mediators (Chen et al., 2017; Hinks et al., 2019; Wang et al., 2018).

Thus, it is unclear how MAIT cells arbitrate between anti-bacterial functions with the associated cost of inflicting tissue damage and tissue repair functions (Salou and Lantz, 2019) (**Figure 4**). Anti-infectious immunity and tissue repair are tightly interweaved processes as

infection and the subsequent immune response result in tissue damage, the resolution of which is key to maintain tissue integrity and functions. MAIT cells are likely involved in both processes through the variety of effector molecules they produce (Hinks et al., 2019; Lamichhane et al., 2019; Leng et al., 2019). Moreover, physiologic (such as stomach acidity) or physical stress (due to breathing in the lungs for instance (Solis et al., 2019), or little traumatic disruptions of barrier tissues) may put in contact the organism with usually excluded microbiota. As seen above, MAIT cells can be activated through TCR-dependent or -independent pathways, and a tight integration of these signals is pivotal in triggering one function versus another. In mice at steady-state, both functions appear segregated in two distinct subsets expressing either T-bet or ROR γ t. However, upon bacterial infection, single MAIT cells can co-express T-bet and ROR γ t. In such MAIT cells, as in humans, both effector functions may be achieved by the same cell sequentially or according to the lymphokine context.

8. Concluding remarks

Besides infectious diseases, the frequency of blood MAIT cells is modified during several auto-immune diseases: diabetes (Magalhaes et al., 2015; Rouxel et al., 2017), multiple sclerosis (Annibali et al., 2010; Mekinian et al., 2017; Mexhitaj et al., 2019; Miyazaki et al., 2011; Treiner and Liblau, 2015; Willing et al., 2018; Willing et al., 2014), inflammatory bowel disease (Serriari et al., 2014) as well as obesity (Magalhaes et al., 2015). All these diseases are associated with important dysbiosis (Honda and Littman, 2016). Interestingly, bariatric surgery leads to a rapid (3 months) recovery of MAIT cell blood numbers (Magalhaes et al., 2015). One of the first changes observed after bariatric surgery is microbiota modifications (Liu et al., 2017) although this is controversial (Aron-Wisnewsky et al., 2019). Thus, MAIT cells could sense the changes in microbiota ecology and the modifications of their blood frequency would reflect the dysbiosis which is common in these diseases.

Tissue location, antigen specificity and context-dependent effector functions of MAIT cells explain their versatile properties and their involvement in various settings such as anti-bacterial defense, tissue repair and epithelium homeostasis. Bacterial infections can inflict considerable damage to host tissues as pathogen clearance by immune cells involves the release of mediators which are also associated with cell death and loss of tissue integrity. Following pathogen elimination, tissue repair processes enable the reconstruction of the vasculature, intercellular matrix, epithelium and stromal cell networks. Through the expression of both anti-microbial and wound-healing effector molecules, MAIT cells are well equipped to contribute to both of these overlapping phases of immune responses to infections according to the lymphokine milieu and ligand availability. Future studies will elucidate the dynamics of vitamin B2-derived metabolites *in vivo*, at steady state and along the course of infections and wound healing processes. A better understanding of MAIT cell triggers, together with the development of stable agonists, will pave the way for harnessing the clinical potential of these abundant T cells.

Acknowledgments

This work was supported by the Institut National de la Santé et de la Recherche Médicale, Institut Curie, Agence Nationale de la Recherche (ANR) (Generic call (MAIT, NEOMAIT and DIABMAIT) and Labex DCBIOL) and by INCA.

Author contribution

All authors contributed to the writing.

Conflict of interest

The authors declare no competing interests

References

- Alferink, J., Tafuri, A., Vestweber, D., Hallmann, R., Hammerling, G.J., and Arnold, B. (1998). Control of neonatal tolerance to tissue antigens by peripheral T cell trafficking. *Science* *282*, 1338-1341.
- Annibaldi, V., Ristori, G., Angelini, D.F., Serafini, B., Mechelli, R., Cannoni, S., Romano, S., Paolillo, A., Abderrahim, H., Diamantini, A., *et al.* (2010). CD161^{high}CD8⁺T cells bear pathogenetic potential in multiple sclerosis. *Brain* *134*, 542-554.
- Aron-Wisnewsky, J., Prifti, E., Belda, E., Ichou, F., Kayser, B.D., Dao, M.C., Verger, E.O., Hedjazi, L., Bouillot, J.L., Chevallier, J.M., *et al.* (2019). Major microbiota dysbiosis in severe obesity: fate after bariatric surgery. *Gut* *68*, 70-82.
- Arpaia, N., Green, J.A., Moltedo, B., Arvey, A., Hemmers, S., Yuan, S., Treuting, P.M., and Rudensky, A.Y. (2015). A Distinct Function of Regulatory T Cells in Tissue Protection. *Cell* *162*, 1078-1089.
- Awad, W., Ler, G.J.M., Xu, W., Keller, A.N., Mak, J.Y.W., Lim, X.Y., Liu, L., Eckle, S.B.G., Le Nours, J., McCluskey, J., *et al.* (2020). The molecular basis underpinning the potency and specificity of MAIT cell antigens. *Nat Immunol* *21*, 400-411.
- Ben Youssef, G., Tourret, M., Salou, M., Ghazarian, L., Houdouin, V., Mondot, S., Mburu, Y., Lambert, M., Azarnoush, S., Diana, J.S., *et al.* (2018). Ontogeny of human mucosal-associated invariant T cells and related T cell subsets. *J Exp Med* *215*, 459-479.
- Beura, L.K., and Masopust, D. (2014). SnapShot: resident memory T cells. *Cell* *157*, 1488-1488 e1481.
- Bottcher, K., Rombouts, K., Saffioti, F., Roccarina, D., Rosselli, M., Hall, A., Luong, T., Tsochatzis, E.A., Thorburn, D., and Pinzani, M. (2018). MAIT cells are chronically activated in patients with autoimmune liver disease and promote pro-fibrogenic hepatic stellate cell activation. *Hepatology*.
- Boudinot, P., Mondot, S., Jouneau, L., Teyton, L., Lefranc, M.P., and Lantz, O. (2016). Restricting nonclassical MHC genes coevolve with TRAV genes used by innate-like T cells in mammals. *Proc Natl Acad Sci U S A* *113*, E2983-2992.
- Burzyn, D., Kuswanto, W., Kolodin, D., Shadrach, J.L., Cerletti, M., Jang, Y., Sefik, E., Tan, T.G., Wagers, A.J., Benoist, C., and Mathis, D. (2013). A special population of regulatory T cells potentiates muscle repair. *Cell* *155*, 1282-1295.
- Chen, Z., Wang, H., D'Souza, C., Sun, S., Kostenko, L., Eckle, S.B., Meehan, B.S., Jackson, D.C., Strugnell, R.A., Cao, H., *et al.* (2017). Mucosal-associated invariant T-cell activation and accumulation after in vivo infection depends on microbial riboflavin synthesis and co-stimulatory signals. *Mucosal immunology* *10*, 58-68.

- Chua, W.J., Truscott, S.M., Eickhoff, C.S., Blazevic, A., Hoft, D.F., and Hansen, T.H. (2012). Polyclonal mucosa-associated invariant T cells have unique innate functions in bacterial infection. *Infect Immun* *80*, 3256-3267.
- Constantinides, M.G., Link, V.M., Tamoutounour, S., Wong, A.C., Perez-Chaparro, P.J., Han, S.J., Chen, Y.E., Li, K., Farhat, S., Weckel, A., *et al.* (2019). MAIT cells are imprinted by the microbiota in early life and promote tissue repair. *Science* *366*.
- Corbett, A.J., Eckle, S.B., Birkinshaw, R.W., Liu, L., Patel, O., Mahony, J., Chen, Z., Reantragoon, R., Meehan, B., Cao, H., *et al.* (2014). T-cell activation by transitory neoantigens derived from distinct microbial pathways. *Nature* *509*, 361-365.
- Crowther, M.D., Dolton, G., Legut, M., Caillaud, M.E., Lloyd, A., Attaf, M., Galloway, S.A.E., Rius, C., Farrell, C.P., Szomolay, B., *et al.* (2020). Genome-wide CRISPR-Cas9 screening reveals ubiquitous T cell cancer targeting via the monomorphic MHC class I-related protein MR1. *Nat Immunol* *21*, 178-185.
- Cui, Y., Franciszkiewicz, K., Mburu, Y.K., Mondot, S., Le Bourhis, L., Premel, V., Martin, E., Kachaner, A., Duban, L., Ingersoll, M.A., *et al.* (2015). Mucosal-associated invariant T cell-rich congenic mouse strain allows functional evaluation. *J Clin Invest* *125*, 4171-4185.
- D'Souza, C., Pediongco, T., Wang, H., Scheerlinck, J.Y., Kostenko, L., Esterbauer, R., Stent, A.W., Eckle, S.B.G., Meehan, B.S., Strugnell, R.A., *et al.* (2018). Mucosal-Associated Invariant T Cells Augment Immunopathology and Gastritis in Chronic *Helicobacter pylori* Infection. *J Immunol*.
- Davalos-Misslitz, A.C., Worbs, T., Willenzon, S., Bernhardt, G., and Forster, R. (2007). Impaired responsiveness to T-cell receptor stimulation and defective negative selection of thymocytes in CCR7-deficient mice. *Blood* *110*, 4351-4359.
- Dias, J., Leeansyah, E., and Sandberg, J.K. (2017). Multiple layers of heterogeneity and subset diversity in human MAIT cell responses to distinct microorganisms and to innate cytokines. *Proc Natl Acad Sci U S A* *114*, E5434-E5443.
- Dusseaux, M., Martin, E., Serriari, N., Peguillet, I., Premel, V., Louis, D., Milder, M., Le Bourhis, L., Soudais, C., Treiner, E., and Lantz, O. (2011). Human MAIT cells are xenobiotic resistant, tissue-targeted, CD161hi IL-17 secreting T cells. *Blood* *117*, 1250-1259.
- Eckle, S.B., Birkinshaw, R.W., Kostenko, L., Corbett, A.J., McWilliam, H.E., Reantragoon, R., Chen, Z., Gherardin, N.A., Beddoe, T., Liu, L., *et al.* (2014). A molecular basis underpinning the T cell receptor heterogeneity of mucosal-associated invariant T cells. *J Exp Med* *211*, 1585-1600.
- Fournie, J.J., and Bonneville, M. (1996). Stimulation of gamma delta T cells by phosphoantigens. *Res Immunol* *147*, 338-347.
- Franciszkiewicz, K., Salou, M., Legoux, F., Zhou, Q., Cui, Y., Bessoles, S., and Lantz, O. (2016). MHC class I-related molecule, MR1, and mucosal-associated invariant T cells. *Immunol Rev* *272*, 120-138.
- Georgel, P., Radosavljevic, M., Macquin, C., and Bahram, S. (2011). The non-conventional MHC class I MR1 molecule controls infection by *Klebsiella pneumoniae* in mice. *Mol Immunol* *48*, 769-775.
- Gherardin, N.A., Keller, A.N., Woolley, R.E., Le Nours, J., Ritchie, D.S., Neeson, P.J., Birkinshaw, R.W., Eckle, S.B., Waddington, J.N., Liu, L., *et al.* (2016). Diversity of T

Cells Restricted by the MHC Class I-Related Molecule MR1 Facilitates Differential Antigen Recognition. *Immunity* 44, 32-45.

Gherardin, N.A., Souter, M.N., Koay, H.F., Mangas, K.M., Seemann, T., Stinear, T.P., Eckle, S.B., Berzins, S.P., d'Udekem, Y., Konstantinov, I.E., *et al.* (2018a). Human blood MAIT cell subsets defined using MR1 tetramers. *Immunol Cell Biol* 96, 507-525.

Gherardin, N.A., Souter, M.N.T., Koay, H.F., Mangas, K.M., Seemann, T., Stinear, T.P., Eckle, S.B.G., Berzins, S.P., d'Udekem, Y., Konstantinov, I.E., *et al.* (2018b). Human blood MAIT cell subsets defined using MR1 tetramers. *Immunol Cell Biol*.

Godfrey, D.I., Koay, H.F., McCluskey, J., and Gherardin, N.A. (2019). The biology and functional importance of MAIT cells. *Nat Immunol* 20, 1110-1128.

Gold, M.C., Cerri, S., Smyk-Pearson, S., Cansler, M.E., Vogt, T.M., Delepine, J., Winata, E., Swarbrick, G.M., Chua, W.J., Yu, Y.Y., *et al.* (2010). Human mucosal associated invariant T cells detect bacterially infected cells. *PLoS Biol* 8, e1000407.

Gomez de Agüero, M., Ganal-Vonarburg, S.C., Fuhrer, T., Rupp, S., Uchimura, Y., Li, H., Steinert, A., Heikenwalder, M., Hapfelmeier, S., Sauer, U., *et al.* (2016). The maternal microbiota drives early postnatal innate immune development. *Science* 351, 1296-1302.

Green, J.A., Arpaia, N., Schizas, M., Dobrin, A., and Rudensky, A.Y. (2017). A nonimmune function of T cells in promoting lung tumor progression. *J Exp Med* 214, 3565-3575.

Grimaldi, D., Le Bourhis, L., Sauneuf, B., Dechartres, A., Rousseau, C., Ouaz, F., Milder, M., Louis, D., Chiche, J.D., Mira, J.P., *et al.* (2014). Specific MAIT cell behaviour among innate-like T lymphocytes in critically ill patients with severe infections. *Intensive care medicine* 40, 192-201.

Gury-BenAri, M., Thaïss, C.A., Serafini, N., Winter, D.R., Giladi, A., Lara-Astiaso, D., Levy, M., Salame, T.M., Weiner, A., David, E., *et al.* (2016). The Spectrum and Regulatory Landscape of Intestinal Innate Lymphoid Cells Are Shaped by the Microbiome. *Cell* 166, 1231-1246 e1213.

Gutierrez-Arcelus, M., Teslovich, N., Mola, A.R., Polidoro, R.B., Nathan, A., Kim, H., Hannes, S., Slowikowski, K., Watts, G.F.M., Korsunsky, I., *et al.* (2019). Lymphocyte innateness defined by transcriptional states reflects a balance between proliferation and effector functions. *Nature communications* 10, 687.

Harriff, M.J., McMurtrey, C., Froyd, C.A., Jin, H., Cansler, M., Null, M., Worley, A., Meermeier, E.W., Swarbrick, G., Nilsen, A., *et al.* (2018). MR1 displays the microbial metabolome driving selective MR1-restricted T cell receptor usage. *Sci Immunol* 3.

Harrison, O.J., Linehan, J.L., Shih, H.Y., Bouladoux, N., Han, S.J., Smelkinson, M., Sen, S.K., Byrd, A.L., Enamorado, M., Yao, C., *et al.* (2019). Commensal-specific T cell plasticity promotes rapid tissue adaptation to injury. *Science* 363.

Hegde, P., Weiss, E., Paradis, V., Wan, J., Mabire, M., Sukriti, S., Rautou, P.E., Albuquerque, M., Picq, O., Gupta, A.C., *et al.* (2018). Mucosal-associated invariant T cells are a profibrogenic immune cell population in the liver. *Nature communications* 9, 2146.

Heng, T.S., Painter, M.W., and Immunological Genome Project, C. (2008). The Immunological Genome Project: networks of gene expression in immune cells. *Nat Immunol* 9, 1091-1094.

- Hinks, T.S.C., Marchi, E., Jabeen, M., Olshansky, M., Kurioka, A., Pediongco, T.J., Meehan, B.S., Kostenko, L., Turner, S.J., Corbett, A.J., *et al.* (2019). Activation and In Vivo Evolution of the MAIT Cell Transcriptome in Mice and Humans Reveals Tissue Repair Functionality. *Cell Rep* 28, 3249-3262 e3245.
- Honda, K., and Littman, D.R. (2016). The microbiota in adaptive immune homeostasis and disease. *Nature* 535, 75-84.
- Howson, L.J., Napolitani, G., Shepherd, D., Ghadbane, H., Kurupati, P., Preciado-Llanes, L., Rei, M., Dobinson, H.C., Gibani, M.M., Teng, K.W.W., *et al.* (2018). MAIT cell clonal expansion and TCR repertoire shaping in human volunteers challenged with *Salmonella Paratyphi A*. *Nature communications* 9, 253.
- Huang, S., Martin, E., Kim, S., Yu, L., Soudais, C., Fremont, D.H., Lantz, O., and Hansen, T.H. (2009). MR1 antigen presentation to mucosal-associated invariant T cells was highly conserved in evolution. *Proc Natl Acad Sci U S A* 106, 8290-8295.
- Jin, C., Lagoudas, G.K., Zhao, C., Bullman, S., Bhutkar, A., Hu, B., Ameh, S., Sandel, D., Liang, X.S., Mazzilli, S., *et al.* (2019). Commensal Microbiota Promote Lung Cancer Development via $\gamma\delta$ T Cells. *Cell* 176, 998-1013 e1016.
- Keller, A.N., Eckle, S.B., Xu, W., Liu, L., Hughes, V.A., Mak, J.Y., Meehan, B.S., Pediongco, T., Birkinshaw, R.W., Chen, Z., *et al.* (2017a). Drugs and drug-like molecules can modulate the function of mucosal-associated invariant T cells. *Nat Immunol* 18, 402-411.
- Keller, A.N., Eckle, S.B., Xu, W., Liu, L., Hughes, V.A., Mak, J.Y., Meehan, B.S., Pediongco, T., Birkinshaw, R.W., Chen, Z., *et al.* (2017b). Drugs and drug-like molecules can modulate the function of mucosal-associated invariant T cells. *Nat Immunol*.
- Kelly, J., Minoda, Y., Meredith, T., Cameron, G., Philipp, M.S., Pellicci, D.G., Corbett, A.J., Kurts, C., Gray, D.H., Godfrey, D.I., *et al.* (2019). Chronically stimulated human MAIT cells are unexpectedly potent IL-13 producers. *Immunol Cell Biol* 97, 689-699.
- Kiselev, V.Y., Andrews, T.S., and Hemberg, M. (2019). Challenges in unsupervised clustering of single-cell RNA-seq data. *Nat Rev Genet* 20, 273-282.
- Kjer-Nielsen, L., Patel, O., Corbett, A.J., Le Nours, J., Meehan, B., Liu, L., Bhati, M., Chen, Z., Kostenko, L., Reantragoon, R., *et al.* (2012). MR1 presents microbial vitamin B metabolites to MAIT cells. *Nature* 491, 717-723.
- Koay, H.F., Gherardin, N.A., Enders, A., Loh, L., Mackay, L.K., Almeida, C.F., Russ, B.E., Nold-Petry, C.A., Nold, M.F., Bedoui, S., *et al.* (2016). A three-stage intrathymic development pathway for the mucosal-associated invariant T cell lineage. *Nat Immunol* 17, 1300-1311.
- Koay, H.F., Gherardin, N.A., Xu, C., Seneviratna, R., Zhao, Z., Chen, Z., Fairlie, D.P., McCluskey, J., Pellicci, D.G., Uldrich, A.P., and Godfrey, D.I. (2019a). Diverse MR1-restricted T cells in mice and humans. *Nature communications* 10, 2243.
- Koay, H.F., Su, S., Amann-Zalcenstein, D., Daley, S.R., Comerford, I., Miosge, L., Whyte, C.E., Konstantinov, I.E., d'Udekem, Y., Baldwin, T., *et al.* (2019b). A divergent transcriptional landscape underpins the development and functional branching of MAIT cells. *Sci Immunol* 4.
- Kurioka, A., Ussher, J.E., Cosgrove, C., Clough, C., Fergusson, J.R., Smith, K., Kang, Y.H., Walker, L.J., Hansen, T.H., Willberg, C.B., and Klenerman, P. (2015). MAIT cells are licensed through granzyme exchange to kill bacterially sensitized targets. *Mucosal immunology* 8, 429-440.

- Kurioka, A., Walker, L.J., Klenerman, P., and Willberg, C.B. (2016). MAIT cells: new guardians of the liver. *Clin Transl Immunology* 5, e98.
- Lamichhane, R., Galvin, H., Hannaway, R.F., de la Harpe, S.M., Munro, F., Tyndall, J.D., Vernall, A.J., McCall, J.L., Husain, M., and Ussher, J.E. (2020). Type I interferons are important co-stimulatory signals during T cell receptor mediated human MAIT cell activation. *Eur J Immunol* 50, 178-191.
- Lamichhane, R., Schneider, M., de la Harpe, S.M., Harrop, T.W.R., Hannaway, R.F., Dearden, P.K., Kirman, J.R., Tyndall, J.D.A., Vernall, A.J., and Ussher, J.E. (2019). TCR- or Cytokine-Activated CD8(+) Mucosal-Associated Invariant T Cells Are Rapid Polyfunctional Effectors That Can Coordinate Immune Responses. *Cell Rep* 28, 3061-3076 e3065.
- Lantz, O., and Legoux, F. (2018). MAIT cells: an historical and evolutionary perspective. *Immunol Cell Biol* 96, 564-572.
- Lantz, O., and Legoux, F. (2019). MAIT cells: programmed in the thymus to mediate immunity within tissues. *Curr Opin Immunol* 58, 75-82.
- Law, B.M.P., Wilkinson, R., Wang, X., Kildey, K., Giuliani, K., Beagley, K.W., Ungerer, J., Healy, H., and Kassianos, A.J. (2019). Human Tissue-Resident Mucosal-Associated Invariant T (MAIT) Cells in Renal Fibrosis and CKD. *J Am Soc Nephrol* 30, 1322-1335.
- Le Bourhis, L., Dusseaux, M., Bohineust, A., Bessoles, S., Martin, E., Premel, V., Core, M., Sleurs, D., Serriari, N.E., Treiner, E., *et al.* (2013). MAIT cells detect and efficiently lyse bacterially-infected epithelial cells. *PLoS pathogens* 9, e1003681.
- Le Bourhis, L., Martin, E., Peguillet, I., Guihot, A., Froux, N., Core, M., Levy, E., Dusseaux, M., Meyssonier, V., Premel, V., *et al.* (2010). Antimicrobial activity of mucosal-associated invariant T cells. *Nat Immunol* 11, 701-708.
- Le Nours, J., Gherardin, N.A., Ramarathinam, S.H., Awad, W., Wiede, F., Gully, B.S., Khandokar, Y., Praveena, T., Wubben, J.M., Sandow, J.J., *et al.* (2019). A class of gammadelta T cell receptors recognize the underside of the antigen-presenting molecule MR1. *Science* 366, 1522-1527.
- Lee, C.H., Zhang, H.H., Singh, S.P., Koo, L., Kabat, J., Tsang, H., Singh, T.P., and Farber, J.M. (2018). C/EBPdelta drives interactions between human MAIT cells and endothelial cells that are important for extravasation. *Elife* 7, e32532.
- Leeansyah, E., Svard, J., Dias, J., Buggert, M., Nystrom, J., Quigley, M.F., Moll, M., Sonnerborg, A., Nowak, P., and Sandberg, J.K. (2015). Arming of MAIT Cell Cytolytic Antimicrobial Activity Is Induced by IL-7 and Defective in HIV-1 Infection. *PLoS pathogens* 11, e1005072.
- Legoux, F., Bellet, D., Daviaud, C., El Morr, Y., Darbois, A., Niort, K., Procopio, E., Salou, M., Gilet, J., Ryffel, B., *et al.* (2019a). Microbial metabolites control the thymic development of mucosal-associated invariant T cells. *Science* 366, 494-499.
- Legoux, F., Gilet, J., Procopio, E., Echasserieau, K., Bernardeau, K., and Lantz, O. (2019b). Molecular mechanisms of lineage decisions in metabolite-specific T cells. *Nat Immunol* 20, 1244-1255.
- Leng, T., Akther, H.D., Hackstein, C.P., Powell, K., King, T., Friedrich, M., Christoforidou, Z., McCuaig, S., Neyazi, M., Arancibia-Carcamo, C.V., *et al.* (2019). TCR and Inflammatory Signals Tune Human MAIT Cells to Exert Specific Tissue Repair and Effector Functions. *Cell Rep* 28, 3077-3091 e3075.

- Lepore, M., Kalinichenko, A., Calogero, S., Kumar, P., Paleja, B., Schmalzer, M., Narang, V., Zolezzi, F., Poidinger, M., Mori, L., and De Libero, G. (2017). Functionally diverse human T cells recognize non-microbial antigens presented by MR1. *Elife* 6.
- Lepore, M., Kalinichenko, A., Colone, A., Paleja, B., Singhal, A., Tschumi, A., Lee, B., Poidinger, M., Zolezzi, F., Quagliata, L., *et al.* (2014). Parallel T-cell cloning and deep sequencing of human MAIT cells reveal stable oligoclonal TCRbeta repertoire. *Nature communications* 5, 3866.
- Linehan, J.L., Harrison, O.J., Han, S.J., Byrd, A.L., Vujkovic-Cvijin, I., Villarino, A.V., Sen, S.K., Shaik, J., Smelkinson, M., Tamoutounour, S., *et al.* (2018). Non-classical Immunity Controls Microbiota Impact on Skin Immunity and Tissue Repair. *Cell* 172, 784-796 e718.
- Liu, R., Hong, J., Xu, X., Feng, Q., Zhang, D., Gu, Y., Shi, J., Zhao, S., Liu, W., Wang, X., *et al.* (2017). Gut microbiome and serum metabolome alterations in obesity and after weight-loss intervention. *Nat Med* 23, 859-868.
- Loh, L., Wang, Z., Sant, S., Koutsakos, M., Jegaskanda, S., Corbett, A.J., Liu, L., Fairlie, D.P., Crowe, J., Rossjohn, J., *et al.* (2016). Human mucosal-associated invariant T cells contribute to antiviral influenza immunity via IL-18-dependent activation. *Proc Natl Acad Sci U S A* 113, 10133-10138.
- Lu, B., Liu, M., Wang, J., Fan, H., Yang, D., Zhang, L., Gu, X., Nie, J., Chen, Z., Corbett, A.J., *et al.* (2020). IL-17 production by tissue-resident MAIT cells is locally induced in children with pneumonia. *Mucosal immunology*.
- Lu, Y., Zhong, M.C., Qian, J., Calderon, V., Cruz Tleugabulova, M., Mallevaey, T., and Veillette, A. (2019). SLAM receptors foster iNKT cell development by reducing TCR signal strength after positive selection. *Nat Immunol* 20, 447-457.
- Mackay, L.K., Stock, A.T., Ma, J.Z., Jones, C.M., Kent, S.J., Mueller, S.N., Heath, W.R., Carbone, F.R., and Gebhardt, T. (2012). Long-lived epithelial immunity by tissue-resident memory T (TRM) cells in the absence of persisting local antigen presentation. *Proc Natl Acad Sci U S A* 109, 7037-7042.
- Magalhaes, I., Pingris, K., Poitou, C., Bessoles, S., Venteclef, N., Kiaf, B., Beaudoin, L., Da Silva, J., Allatif, O., Rossjohn, J., *et al.* (2015). Mucosal-associated invariant T cell alterations in obese and type 2 diabetic patients. *J Clin Invest* 125, 1752-1762.
- Mak, J.Y., Xu, W., Reid, R.C., Corbett, A.J., Meehan, B.S., Wang, H., Chen, Z., Rossjohn, J., McCluskey, J., Liu, L., and Fairlie, D.P. (2017). Stabilizing short-lived Schiff base derivatives of 5-aminouracils that activate mucosal-associated invariant T cells. *Nature communications* 8, 14599.
- Mao, A.P., Constantinides, M.G., Mathew, R., Zuo, Z., Chen, X., Weirauch, M.T., and Bendelac, A. (2016). Multiple layers of transcriptional regulation by PLZF in NKT-cell development. *Proc Natl Acad Sci U S A* 113, 7602-7607.
- Martin, E., Treiner, E., Duban, L., Guerri, L., Laude, H., Toly, C., Premel, V., Devys, A., Moura, I.C., Tilloy, F., *et al.* (2009). Stepwise development of MAIT cells in mouse and human. *PLoS Biol* 7, e54.
- Matzinger, P., and Kamala, T. (2011). Tissue-based class control: the other side of tolerance. *Nat Rev Immunol* 11, 221-230.
- McSharry, B.P., Samer, C., McWilliam, H.E.G., Ashley, C.L., Yee, M.B., Steain, M., Liu, L., Fairlie, D.P., Kinchington, P.R., McCluskey, J., *et al.* (2020). Virus-Mediated

Suppression of the Antigen Presentation Molecule MR1. *Cell Rep* 30, 2948-2962 e2944.

McWilliam, H.E., Eckle, S.B., Theodossis, A., Liu, L., Chen, Z., Wubben, J.M., Fairlie, D.P., Strugnell, R.A., Mintern, J.D., McCluskey, J., *et al.* (2016). The intracellular pathway for the presentation of vitamin B-related antigens by the antigen-presenting molecule MR1. *Nat Immunol* 17, 531-537.

Meermeier, E.W., Laugel, B.F., Sewell, A.K., Corbett, A.J., Rossjohn, J., McCluskey, J., Harriff, M.J., Franks, T., Gold, M.C., and Lewinsohn, D.M. (2016). Human TRAV1-2-negative MR1-restricted T cells detect *S. pyogenes* and alternatives to MAIT riboflavin-based antigens. *Nature communications* 7, 12506.

Meierovics, A., Yankelevich, W.J., and Cowley, S.C. (2013). MAIT cells are critical for optimal mucosal immune responses during in vivo pulmonary bacterial infection. *Proc Natl Acad Sci U S A* 110, E3119-3128.

Meierovics, A.I., and Cowley, S.C. (2016). MAIT cells promote inflammatory monocyte differentiation into dendritic cells during pulmonary intracellular infection. *J Exp Med* 213, 2793-2809.

Mekinian, A., Mahevas, T., Mohty, M., Jachiet, V., Riviere, S., Fain, O., and Gaugler, B. (2017). Mucosal-associated Invariant Cells are Deficient in Systemic Sclerosis. *Scand J Immunol* 86, 216-220.

Mexhitaj, I., Nyirenda, M.H., Li, R., O'Mahony, J., Rezk, A., Rozenberg, A., Moore, C.S., Johnson, T., Sadovnick, D., Collins, D.L., *et al.* (2019). Abnormal effector and regulatory T cell subsets in paediatric-onset multiple sclerosis. *Brain* 142, 617-632.

Milner, J.J., Toma, C., Yu, B., Zhang, K., Omilusik, K., Phan, A.T., Wang, D., Getzler, A.J., Nguyen, T., Crotty, S., *et al.* (2017). Runx3 programs CD8(+) T cell residency in non-lymphoid tissues and tumours. *Nature* 552, 253-257.

Miyazaki, Y., Miyake, S., Chiba, A., Lantz, O., and Yamamura, T. (2011). Mucosal-associated invariant T cells regulate Th1 response in multiple sclerosis. *Int Immunol* 23, 529-535.

Mondot, S., Boudinot, P., and Lantz, O. (2016). MAIT, MR1, microbes and riboflavin: a paradigm for the co-evolution of invariant TCRs and restricting MHCII-like molecules? *Immunogenetics* 68, 537-548.

Nejman, D., Livyatan, I., Fuks, G., Gavert, N., Zwang, Y., Geller, L.T., Rotter-Maskowitz, A., Weiser, R., Mallel, G., Gigi, E., *et al.* (2020). The human tumor microbiome is composed of tumor type-specific intracellular bacteria. *Science* 368, 973-980.

Pellicci, D.G., Koay, H.F., and Berzins, S.P. (2020). Thymic development of unconventional T cells: how NKT cells, MAIT cells and gammadelta T cells emerge. *Nat Rev Immunol*.

Porcelli, S., Yockey, C.E., Brenner, M.B., and Balk, S.P. (1993). Analysis of T cell antigen receptor (TCR) expression by human peripheral blood CD4-8- alpha/beta T cells demonstrates preferential use of several V beta genes and an invariant TCR alpha chain. *J Exp Med* 178, 1-16.

Rahimpour, A., Koay, H.F., Enders, A., Clanchy, R., Eckle, S.B., Meehan, B., Chen, Z., Whittle, B., Liu, L., Fairlie, D.P., *et al.* (2015). Identification of phenotypically and functionally heterogeneous mouse mucosal-associated invariant T cells using MR1 tetramers. *J Exp Med* 212, 1095-1108.

- Reantragoon, R., Corbett, A.J., Sakala, I.G., Gherardin, N.A., Furness, J.B., Chen, Z., Eckle, S.B., Uldrich, A.P., Birkinshaw, R.W., Patel, O., *et al.* (2013). Antigen-loaded MR1 tetramers define T cell receptor heterogeneity in mucosal-associated invariant T cells. *J Exp Med* *210*, 2305-2320.
- Reantragoon, R., Kjer-Nielsen, L., Patel, O., Chen, Z., Illing, P.T., Bhati, M., Kostenko, L., Bharadwaj, M., Meehan, B., Hansen, T.H., *et al.* (2012). Structural insight into MR1-mediated recognition of the mucosal associated invariant T cell receptor. *J Exp Med* *209*, 761-774.
- Riva, A., Patel, V., Kurioka, A., Jeffery, H.C., Wright, G., Tarff, S., Shawcross, D., Ryan, J.M., Evans, A., Azarian, S., *et al.* (2017). Mucosa-associated invariant T cells link intestinal immunity with antibacterial immune defects in alcoholic liver disease. *Gut*.
- Rouxel, O., Da Silva, J., Beaudoin, L., Nel, I., Tard, C., Cagninacci, L., Kiaf, B., Oshima, M., Diedisheim, M., Salou, M., *et al.* (2017). Cytotoxic and regulatory roles of mucosal-associated invariant T cells in type 1 diabetes. *Nat Immunol* *18*, 1321-1331.
- Salerno-Goncalves, R., Luo, D., Fresnay, S., Magder, L., Darton, T.C., Jones, C., Waddington, C.S., Blohmke, C.J., Angus, B., Levine, M.M., *et al.* (2017). Challenge of Humans with Wild-type *Salmonella enterica* Serovar Typhi Elicits Changes in the Activation and Homing Characteristics of Mucosal-Associated Invariant T Cells. *Front Immunol* *8*, 398.
- Salio, M., Awad, W., Veerapen, N., Gonzalez-Lopez, C., Kulicke, C., Waithe, D., Martens, A.W.J., Lewinsohn, D.M., Hobrath, J.V., Cox, L.R., *et al.* (2020). Ligand-dependent downregulation of MR1 cell surface expression. *Proc Natl Acad Sci U S A*.
- Salou, M., Franciszkiewicz, K., and Lantz, O. (2017). MAIT cells in infectious diseases. *Curr Opin Immunol* *48*, 7-14.
- Salou, M., and Lantz, O. (2019). A TCR-Dependent Tissue Repair Potential of MAIT Cells. *Trends Immunol*.
- Salou, M., Legoux, F., Gilet, J., Darbois, A., du Halgouet, A., Alonso, R., Richer, W., Goubet, A.G., Daviaud, C., Menger, L., *et al.* (2019). A common transcriptomic program acquired in the thymus defines tissue residency of MAIT and NKT subsets. *J Exp Med* *216*, 133-151.
- Savage, A.K., Constantinides, M.G., Han, J., Picard, D., Martin, E., Li, B., Lantz, O., and Bendelac, A. (2008). The Transcription Factor PLZF Directs the Effector Program of the NKT Cell Lineage. *Immunity* *29*, 391-403.
- Schmaler, M., Colone, A., Spagnuolo, J., Zimmermann, M., Lepore, M., Kalinichenko, A., Bhatia, S., Cottier, F., Rutishauser, T., Pavelka, N., *et al.* (2018). Modulation of bacterial metabolism by the microenvironment controls MAIT cell stimulation. *Mucosal immunology* *11*, 1060-1070.
- Seach, N., Guerri, L., Le Bourhis, L., Mburu, Y., Cui, Y., Bessoles, S., Soudais, C., and Lantz, O. (2013). Double-positive thymocytes select mucosal-associated invariant T cells. *J Immunol* *191*, 6002-6009.
- Serriari, N.E., Eoche, M., Lamotte, L., Fumery, M., Marcelo, P., Chatelain, D., Barre, A., Nguyen-Khac, E., Lantz, O., Dupas, J.L., and Treiner, E. (2014). Innate Mucosal-Associated Invariant T (MAIT) cells are activated in Inflammatory Bowel Diseases. *Clinical and experimental immunology* *191*, 6002-6009.
- Seshadri, C., Thuong, N.T., Mai, N.T., Bang, N.D., Chau, T.T., Lewinsohn, D.M., Thwaites, G.E., Dunstan, S.J., and Hawn, T.R. (2017). A polymorphism in human MR1 is

associated with mRNA expression and susceptibility to tuberculosis. *Genes Immun* 18, 8-14.

Smith, P.M., Howitt, M.R., Panikov, N., Michaud, M., Gallini, C.A., Bohlooly, Y.M., Glickman, J.N., and Garrett, W.S. (2013). The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell homeostasis. *Science* 341, 569-573.

Sobkowiak, M.J., Davanian, H., Heymann, R., Gibbs, A., Emgard, J., Dias, J., Aleman, S., Kruger-Weiner, C., Moll, M., Tjernlund, A., *et al.* (2019). Tissue-resident MAIT cell populations in human oral mucosa exhibit an activated profile and produce IL-17. *Eur J Immunol* 49, 133-143.

Solis, A.G., Bielecki, P., Steach, H.R., Sharma, L., Harman, C.C.D., Yun, S., de Zoete, M.R., Warnock, J.N., To, S.D.F., York, A.G., *et al.* (2019). Mechanosensation of cyclical force by PIEZO1 is essential for innate immunity. *Nature* 573, 69-74.

Soudais, C., Samassa, F., Sarkis, M., Le Bourhis, L., Bessoles, S., Blanot, D., Herve, M., Schmidt, F., Mengin-Lecreulx, D., and Lantz, O. (2015). In Vitro and In Vivo Analysis of the Gram-Negative Bacteria-Derived Riboflavin Precursor Derivatives Activating Mouse MAIT Cells. *J Immunol* 194, 4641-4649.

Tastan, C., Karhan, E., Zhou, W., Fleming, E., Voigt, A.Y., Yao, X., Wang, L., Horne, M., Placek, L., Kozhaya, L., *et al.* (2018). Tuning of human MAIT cell activation by commensal bacteria species and MR1-dependent T-cell presentation. *Mucosal immunology*.

Thomas, S.Y., Scanlon, S.T., Griewank, K.G., Constantinides, M.G., Savage, A.K., Barr, K.A., Meng, F., Luster, A.D., and Bendelac, A. (2011). PLZF induces an intravascular surveillance program mediated by long-lived LFA-1-ICAM-1 interactions. *J Exp Med* 208, 1179-1188.

Tilloy, F., Treiner, E., Park, S.H., Garcia, C., Lemonnier, F., de la Salle, H., Bendelac, A., Bonneville, M., and Lantz, O. (1999). An invariant T cell receptor alpha chain defines a novel TAP-independent major histocompatibility complex class Ib-restricted alpha/beta T cell subpopulation in mammals. *J Exp Med* 189, 1907-1921.

Treiner, E., Duban, L., Bahram, S., Radosavljevic, M., Wanner, V., Tilloy, F., Affaticati, P., Gilfillan, S., and Lantz, O. (2003). Selection of evolutionarily conserved mucosal-associated invariant T cells by MR1. *Nature* 422, 164-169.

Treiner, E., and Liblau, R.S. (2015). Mucosal-Associated Invariant T Cells in Multiple Sclerosis: The Jury is Still Out. *Front Immunol* 6, 503.

Tsukamoto, K., Deakin, J.E., Graves, J.A., and Hashimoto, K. (2013). Exceptionally high conservation of the MHC class I-related gene, MR1, among mammals. *Immunogenetics* 65, 115-124.

Turtle, C.J., Delrow, J., Joslyn, R.C., Swanson, H.M., Basom, R., Tabellini, L., Delaney, C., Heimfeld, S., Hansen, J.A., and Riddell, S.R. (2011). Innate signals overcome acquired TCR signaling pathway regulation and govern the fate of human CD161(hi) CD8alpha semi-invariant T cells. *Blood* 118, 2752-2762.

Tuttle, K.D., Krovi, S.H., Zhang, J., Bedel, R., Harmacek, L., Peterson, L.K., Dragone, L.L., Lefferts, A., Halluszczak, C., Riemondy, K., *et al.* (2018). TCR signal strength controls thymic differentiation of iNKT cell subsets. *Nature communications* 9, 2650.

Uchimura, Y., Fuhrer, T., Li, H., Lawson, M.A., Zimmermann, M., Yilmaz, B., Zindel, J., Ronchi, F., Sorribas, M., Hapfelmeier, S., *et al.* (2018). Antibodies Set Boundaries

- Limiting Microbial Metabolite Penetration and the Resultant Mammalian Host Response. *Immunity* 49, 545-559 e545.
- Ussher, J.E., Bilton, M., Attwod, E., Shadwell, J., Richardson, R., de Lara, C., Mettke, E., Kurioka, A., Hansen, T.H., Klenerman, P., and Willberg, C.B. (2014). CD161⁺⁺ CD8⁺ T cells, including the MAIT cell subset, are specifically activated by IL-12+IL-18 in a TCR-independent manner. *Eur J Immunol* 44, 195-203.
- Ussher, J.E., van Wilgenburg, B., Hannaway, R.F., Ruustal, K., Phalora, P., Kurioka, A., Hansen, T.H., Willberg, C.B., Phillips, R.E., and Klenerman, P. (2016). TLR signaling in human antigen-presenting cells regulates MR1-dependent activation of MAIT cells. *Eur J Immunol* 46, 1600-1614.
- Ussher, J.E., Willberg, C.B., and Klenerman, P. (2018). MAIT cells and viruses. *Immunol Cell Biol*.
- Vahl, J.C., Heger, K., Knies, N., Hein, M.Y., Boon, L., Yagita, H., Polic, B., and Schmidt-Supprian, M. (2013). NKT cell-TCR expression activates conventional T cells in vivo, but is largely dispensable for mature NKT cell biology. *PLoS Biol* 11, e1001589.
- Van Dyken, S.J., and Locksley, R.M. (2013). Interleukin-4- and interleukin-13-mediated alternatively activated macrophages: roles in homeostasis and disease. *Annu Rev Immunol* 31, 317-343.
- van Wilgenburg, B., Scherwitzl, I., Hutchinson, E.C., Leng, T., Kurioka, A., Kulicke, C., de Lara, C., Cole, S., Vasanawathana, S., Limpitikul, W., *et al.* (2016). MAIT cells are activated during human viral infections. *Nature communications* 7, 11653.
- Voillet, V., Buggert, M., Slichter, C.K., Berkson, J.D., Mair, F., Addison, M.M., Dori, Y., Nadolski, G., Itkin, M.G., Gottardo, R., *et al.* (2018). Human MAIT cells exit peripheral tissues and recirculate via lymph in steady state conditions. *JCI Insight* 3.
- Walker, L.J., Kang, Y.H., Smith, M.O., Tharmalingham, H., Ramamurthy, N., Fleming, V.M., Sahgal, N., Leslie, A., Oo, Y., Geremia, A., *et al.* (2012). Human MAIT and CD8 α cells develop from a pool of type-17 precommitted CD8⁺ T cells. *Blood* 119, 422-433.
- Wang, H., D'Souza, C., Lim, X.Y., Kostenko, L., Pediongco, T.J., Eckle, S.B.G., Meehan, B.S., Shi, M., Wang, N., Li, S., *et al.* (2018). MAIT cells protect against pulmonary *Legionella longbeachae* infection. *Nature communications* 9, 3350.
- Wang, H., and Hogquist, K.A. (2018). CCR7 defines a precursor for murine iNKT cells in thymus and periphery. *Elife* 7.
- Wang, H., Kjer-Nielsen, L., Shi, M., D'Souza, C., Pediongco, T.J., Cao, H., Kostenko, L., Lim, X.Y., Eckle, S.B.G., Meehan, B.S., *et al.* (2019). IL-23 costimulates antigen-specific MAIT cell activation and enables vaccination against bacterial infection. *Sci Immunol* 4.
- Weinreich, M.A., and Hogquist, K.A. (2008). Thymic emigration: when and how T cells leave home. *J Immunol* 181, 2265-2270.
- Wilgenburg, B.V., Loh, L., Chen, Z., Pediongco, T.J., Wang, H., Shi, M., Zhao, Z., Koutsakos, M., Nussing, S., Sant, S., *et al.* (2018). MAIT cells contribute to protection against lethal influenza infection in vivo. *Nature communications* 9, 4706.
- Willing, A., Jager, J., Reinhardt, S., Kursawe, N., and Friese, M.A. (2018). Production of IL-17 by MAIT Cells Is Increased in Multiple Sclerosis and Is Associated with IL-7 Receptor Expression. *J Immunol* 200, 974-982.

Willing, A., Leach, O.A., Ufer, F., Attfield, K.E., Steinbach, K., Kursawe, N., Piedavent, M., and Friese, M.A. (2014). CD8(+) MAIT cells infiltrate into the CNS and alterations in their blood frequencies correlate with IL-18 serum levels in multiple sclerosis. *Eur J Immunol* *44*, 3119-3128.

Winter, S.J., Kunze-Schumacher, H., Imelmann, E., Grewers, Z., Osthues, T., and Krueger, A. (2019). MicroRNA miR-181a/b-1 controls MAIT cell development. *Immunol Cell Biol* *97*, 190-202.

Xiao, X., Li, K., Ma, X., Liu, B., He, X., Yang, S., Wang, W., Jiang, B., and Cai, J. (2019). Mucosal-Associated Invariant T Cells Expressing the TRAV1-TRAJ33 Chain Are Present in Pigs. *Front Immunol* *10*, 2070.

Yan, J., Allen, S., McDonald, E., Das, I., Mak, J.Y.W., Liu, L., Fairlie, D.P., Meehan, B.S., Chen, Z., Corbett, A.J., *et al.* (2020). MAIT Cells Promote Tumor Initiation, Growth, and Metastases via Tumor MR1. *Cancer discovery* *10*, 124-141.

Zhao, F., Cannons, J.L., Dutta, M., Griffiths, G.M., and Schwartzberg, P.L. (2012). Positive and negative signaling through SLAM receptors regulate synapse organization and thresholds of cytotoxicity. *Immunity* *36*, 1003-1016.


Figure legends.


Figure 1: Model of MAIT antigen circulation. In gut bacteria, the unstable riboflavin precursor 5-A-RU reacts with methylglyoxal or glyoxal from the intermediary metabolism to generate 5-OP-RU or 5-OE-RU, respectively. Bacteria may also produce weak agonists or antagonists that may also bind to MR1 and prevent MAIT activation. The unstable 5-OP/E-RU rapidly crosses the epithelial layer to reach the portal blood and the liver before going through the lung and follow the general blood circulation. This pathway leads to a progressive dilution of 5-OP/E-RU by the afferent blood arriving at each step from other organs. The concentration of 5-OP/E-RU in the arterial blood reaching the thymus is still high enough to generate MR1:5-OP-RU complexes on DP thymocytes able to select MAIT cells, in a process also related to the high expression of MR1 by DP thymocytes. How 5-OP-RU crosses the epithelial layers and whether it is bound to a carrier in the blood are unknown.

Figure 2: Development of 5-OP-RU-specific T cells in the thymus. Upon rearrangement of a TCR with specificity for the 5-OP-RU:MR1 complex, immature thymocytes undergo positive selection by recognizing MR1 at the surface of either TECs (A) or DP thymocytes (B). Positive selection on TECs results in differentiation into naïve conventional-like CD4⁺ T cells. Positive selection on DP thymocytes results in co-stimulation with Slam and engagement of the SAP signaling pathway, which drives expression of PLZF (*Zbtb16*) and effector differentiation into either MAIT1 or MAIT17 cells. Vitamin B2 metabolites such as 5-OP-RU can be captured by several cells in thymus, including DP thymocytes and TECs. Microbiota-derived antigens mediate positive selection of effector MAIT cells and intra-thymic proliferation of MAIT17 cells. Upon thymic egress, MAIT17 cells preferentially localize in mucosal tissues.

Figure 3: MAIT subset positioning in tissues of B6 or B6-MAIT^{Cast} mice. Percentages of MAIT1 (in red) and MAIT17 (in blue) cells among total MAIT cells are indicated.


Figure 4: MAIT17 cell potential effector functions in mucosal barriers including skin, lamina propria and lungs. At steady state, whether MAIT cell are stimulated or not and whether they exert effector functions remain uncertain. In an inflammatory context, depending on the stimulation mode, MAIT cells secrete various effector mediators. How these mediators vary according to the tissue and whether they are secreted by various subpopulations remains unknown. Another hypothesis is that different functions are sequentially triggered. MAIT1 cell functions are less well described as they represent a small fraction of the total population in mucosal barriers






The mouse picture and the organs were retrieved from a published paper (the drawings, not the data)

■ ROR γ t⁺ MAIT cells
■ ROR γ t⁻ MAIT cells





Inflammatory context (infection, tumor...)

