



HAL
open science

Advanced nanomedicines for the treatment of inflammatory diseases

Romain Brusini, Mariana Varna, Patrick Couvreur

► **To cite this version:**

Romain Brusini, Mariana Varna, Patrick Couvreur. Advanced nanomedicines for the treatment of inflammatory diseases. *Advanced Drug Delivery Reviews*, 2020, 157, pp.161 - 178. 10.1016/j.addr.2020.07.010 . hal-03493936

HAL Id: hal-03493936

<https://hal.science/hal-03493936>

Submitted on 15 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Advanced nanomedicines for the treatment of inflammatory diseases

Romain Brusini¹, Mariana Varna¹, Patrick Couvreur^{1#}

¹Institut Galien Paris-Sud, UMR 8612, CNRS, Université Paris-Sud, Université Paris-Saclay,
Faculté de Pharmacie, F-92296 Châtenay-Malabry Cedex, France

#Corresponding author: Prof. Patrick COUVREUR

Email: patrick.couvreur@universite-paris-saclay.fr

Tél: +33 146 835 582

+33 146 835 396

Fax: +33 146 619 334

Abstract

Inflammation, a common feature of many diseases, is an essential immune response that enables survival and maintains tissue homeostasis. However, in some conditions, the inflammatory process becomes detrimental, contributing to the pathogenesis of a disease. Targeting inflammation by using nanomedicines (*i.e.* nanoparticles loaded with a [therapeutic active principle](#)), either through the recognition of molecules overexpressed onto the surface of activated macrophages or endothelial cells, or through enhanced [vasculature permeability](#), or even through biomimicry, offers a promising solution for the treatment of inflammatory diseases. After providing a brief insight on the pathophysiology of inflammation and current therapeutic strategies, the review will discuss, at a pre-clinical stage, the main innovative nanomedicine approaches that have been proposed in the past five years for the resolution of inflammatory disorders, finally focusing on those currently in clinical trials.

Keywords: nanoparticles, drug delivery systems, inflammation, biomimetic nanoparticles, pre-clinical assessment, clinical application

List of abbreviations

5-ASA: 5-AminoSalicylic Acid (also called Mesalazine)

APCs: antigen-presenting cells

AMPS: 2-Acrylamido-2-MethylPropane-Sulfonic acid

ATP: Adenosine Triphosphate

bFGF: basic fibroblast growth factor

COVID-19: coronavirus disease 2019

DC: dendritic cells

Dex: Dexamethasone

EAE: experimental autoimmune encephalomyelitis

ELVIS: Extravasation through Leaky Vasculature and subsequent Inflammatory cell-mediated Sequestration

EPR: Enhanced Permeability and Retention

FDA: Food and Drug Administration

HA: Hyaluronic Acid

HIF: Hypoxia-Inducible transcription Factor

IBD: Inflammatory Bowel Diseases

ICAM: Inter-Cellular Adhesion Molecule

IFN- γ : Interferon gamma

IL-10: Interleukin 10

LMWH: Low Molecular Weight Heparin

LPS: LipoPolySaccharide

LXR: Liver X Receptor

MAPK: Mitogen Activated Protein Kinase

MPO: Myeloperoxidase

MPSS: MethylPrednisolone Sodium Succinate

NF- κ B: Nuclear Factor-kappa B

NIPAM: N-Isopropyl acrylamide

NPs: Nanoparticles

NSAID: Non-Steroidal Anti-inflammatory Drug

pDNA: plasmid DeoxyriboNucleic Acid

PEG: Polyethylene Glycol

PEI: Polyethyleneimine

PLGA: Poly(Lactic-co-Glycolic Acid)

PRR: Pattern-Recognition Receptor

PSL: PhosphatidylSerine-containing Liposome

ROS: Radical Oxygen Species

RA: Rheumatoid arthritis

TCR: T cell receptors

TLR: Toll-Like Receptor

TNF- α : Tumor Necrosis Factor alpha

VCAM: Vascular Cell Adhesion Molecule

VEGF: Vascular Endothelial Growth Factor

1. Introduction

Inflammation represents a mechanism of the body's defence, originating from multiple causes such as infectious agents (viruses, bacteria), physical agents, radical oxygen species (ROS), metabolic stress (hypoxia), to cite only few examples[1]. In some situations, the inflammatory process may, however, become detrimental. The COVID-19 infection, associated in certain cases with a strong cytokine storm, a viral sepsis and an uncontrolled systemic inflammation, leading to acute respiratory distress represents an example of such adverse reaction[2, 3]. Historically, the first to define the clinical symptoms of inflammation was Celsus in the first century AD, who described four cardinal signs of inflammation as follows: "*rubor et tumor cum calore et dolore*" (redness and swelling with heat and pain). In 1858, Virchow added a fifth cardinal sign: "*functio laesa*" (disturbance of function)[1].

Depending on the origin of the agent, inflammation can be classified by whether it is caused by an external agent or an endogenous abnormal response. Based on its duration, inflammation can either be acute or chronic. Acute inflammation has been considered a defence of innate immune response induced by infection or injury[4], while chronic inflammation can accompany some pathological states without any infection or injury, like in obesity[5].

Recently, Antonelli and Kushner redefined inflammation as the innate immune response to potentially harmful stimuli such as pathogens, injury, and metabolic stress[6]. [Indeed, proper inflammatory responses initiate a broad spectrum of innate immune protection and orchestrate long-term adaptive immunity toward specific pathogens. As such, the host immune system brings out layers of regulatory mechanisms that in return modulate the initiation, progression and resolution of inflammation\[6, 7\].](#) Generally, the inflammatory response (Figure 1) [is perceived as a reaction of recognition of the infectious agent \(inducers\), a step which involves specialized \(sensors\) body cells as well as circulating proteins. After recognition, there is a production of mediators of inflammation which target the diseased tissue\[6, 8\].](#) [But both, innate and adaptive immune system, trigger the inflammatory process. Indeed, the components of the innate immunity include phagocytes \(i.e., neutrophils, monocytes, macrophages\), inflammation-related serum proteins \(e.g., complement, C-reactive protein\), cell receptors that signal a defensive response \(e.g., Toll-like receptors\) and cells that release cytokines and inflammatory mediators \(i.e., macrophages, mast cells, natural-killer cells\)\[7\].](#) The complement system not only acts as the first line of defense for the body, but also links innate and adaptive immunity and plays an important role in peripheral lymph nodes to

enhance B and T cell responses. Activation of adaptive immunity occurs after innate immunity signaling and/or antigen presentation by specialized cells known as antigen-presenting cells (APCs), which include dendritic cells (DC) and macrophages. In contrast to the innate immune system, the adaptive immune system is highly specific against one or more antigens after their recognition by specialized receptors at the surface of B and T lymphocytes[9]. B cells are the main producers of antibodies which recognize and bind antigens. T cells are lymphocytes that express T cell receptors (TCR) on their surface and serve essential roles in cell-mediated immunity. However, more and more data show that this distinction between innate immunity and adaptive immunity, does not correspond to reality since enormous interdependence between the two immune responses exists. For example, nitric oxide and reactive oxygen species (ROS) produced by macrophages, dendritic cells or other components of the innate immunity can modulate T cell function and survival[10].

In most cases, the acute inflammatory response is controlled over time, and is normally finished once the insult is eliminated and the damaged tissue repaired. However, in some situations, when the inflammatory inducer persists, a chronic inflammatory response may establish. And when the inflammatory processes last for a long period of time, inflammation may become harmful and damaging for the tissues, which can in turn, contribute to the development and pathogenesis of chronic diseases of altered homeostasis. For example, lymphocyte T cells are thought to be the main triggers of autoimmune disease processes. In addition to innate immunity, adaptive mechanisms, through antigen recognition by the lymphocytes, production of antibodies, and their interaction with complement and phagocytic cells of the innate immunity such as macrophages, control both the nature and shape of the inflammatory responses. Thus, through this retro-activation of innate mechanisms, adaptive responses frequently provoke a persistent inflammation, which contributes to the process of chronic inflammation[7].

Inflammation is a very complex mechanism with, however, new advances in the understanding of its molecular mechanisms each year. Excellent reviews exist in this area describing the pathophysiological mechanisms[1, 6, 8]. It is now known that a large spectrum of diseases are accompanied by an uncontrolled inflammatory response, among these being sepsis, asthma, obesity and type 2 diabetes, neurodegenerative diseases, cardiovascular diseases (e.g. atherosclerosis, cardiac ischemia/ reperfusion), cancer, bowel disease, Crohn disease, rheumatoid polyarthritis, to cite only a few[11]. The understanding of the pathophysiology of the inflammation has allowed to identify proteins expressed by certain cells during inflammation (e.g. ICAM, VCAM, Selectins, ...). Additional phenomena such as

the EPR (enhanced permeability and retention) effect have been identified. This effect, which is an interesting feature of inflamed tissues, leads to an increased vascular permeability coupled to a deregulated neovascularization[12]. EPR effect was first described in tumors where discontinuous endothelium in newly formed and immature vessels are formed. In inflammatory diseases, a similar effect was described with, however, some specificity. Thus, loss of endothelium integrity can result in abnormal neoangiogenesis observed in diseased tissues. The neovascularization mechanisms is described to be induced by hypoxia and inflammation which are critical factors. Hypoxia promotes activation of hypoxia-inducible transcription factors (*e.g.* HIF-1, HIF-2) and subsequently expression of growth factor (*e.g.* endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF)), finally leading to a loss of intercellular junctions between endothelial cells[12, 13]. On the other hand, neutrophils, which are predominate in acute inflammation, promote vascular permeability by releasing neutrophil elastase, myeloperoxidase and MMP-9, which contribute to enhanced vascular permeability. Wang *et al*, referred this phenomenon as ELVIS effect (Extravasation through Leaky Vasculature and subsequent Inflammatory cell-mediated Sequestration) and use it to deliver polymer prodrugs[14, 15]. Other studies have exploited the leaky vasculature during inflammation to deliver nanomedicines like PEGylated polycyanoacrylate [16] or liposomes [17] to the spinal cord of rats with experimental autoimmune encephalomyelitis (EAE).

Today, the therapeutic strategies include the use of steroidal or non-steroidal anti-inflammatory drugs. But two additional therapeutic approaches have recently emerged: the use of anti-leukotrienes that inhibit the recruitment of inflammatory cells and of pro-inflammatory cytokine inhibitors, like anti-TNF alpha or anti-IL-1 monoclonal antibodies. However, these treatments are sometimes not enough to obtain an optimal pharmacological activity. Drug limitations include non-specific biodistribution, low bioavailability and/or short half-life into the body. Moreover, high dosages are needed to be administered to patients, which cause off-target side effects with only modest success in controlling inflammatory disease symptoms[18].

To avoid some of these limitations, the use of nanoparticles (NPs), with a size from a few tenths to a few hundred of nanometers, has gained increasing interest. It is nowadays possible to obtain biocompatible nanomedicines (*i.e.* NPs loaded with active principle) with a highly controlled shape, size and surface charge[19-22]. Moreover, a passive exploitation of **this characteristic leaky vasculature** has allowed an increased delivery and accumulation of nanomedicines through sub-endothelial space. **Additionally**, targeting moieties on

nanoparticles' surface may allow an active accumulation and a controlled drug release into the diseased cells and tissues, reducing toxicity and side-effects[23-25]. Therefore, many nanomedicines were developed with the aim to treat diseases with an inflammatory background, including cancer[26, 27], cardiovascular pathologies[28-30], autoimmune diseases[31], metabolic syndrome[32], neurodegenerative diseases[22, 33, 34].

We have excluded in this review nanomedicines dedicated to cancer therapy where inflammatory processes occurs, because there are already excellent reviews on this subject[35-37]. Nevertheless, to the best of our knowledge, there is no recent review making the state of the art on the use of nanomedicines when inflammation in general becomes detrimental.

This review will focus on the new nanomedicine concepts that have emerged during the last five years concerning the management of inflammatory diseases at the pre-clinical stage. The last part deals with the very low number of nanomedicines in clinical trials.

2. Recent nanomedicines for the treatment of inflammation

A better understanding of the molecular and cellular events underlying the inflammatory response has opened some new perspectives in the treatment of inflammation, particularly through the development of well-designed nanomedicines. Indeed, nanoparticles (NPs) can be specifically engineered to go preferentially to the target tissue from the site of administration, thus addressing issues of conventional therapies such as off-target organ side effects and systemic toxicity, exacerbated by frequent and long-term dosing. The formulation of nanoparticles regulating the expression of pro- and anti-inflammatory molecules and targeting inflammatory sensors or macrophages through phagocytosis, holds great promise for the treatment of inflammatory diseases (Figure 2, Table 1). In addition, using nanocarriers to specifically target effector cells, particularly antigen-presenting cells, could be of great value to promote cellular response or immune tolerance thanks to their modulability which allows them to passively (optimizing the size and/or surface charge of nanoparticles) or actively (decorating nanoparticles with specific antibodies) target these cells[38]. Considerable attention to develop more effective anti-inflammatory nanomedicines, in order to overcome the side effects observed with conventional therapy, has resulted in the development on anti-inflammatory nanomedicines.

2.1. Steroid-based nanomedicines

Corticosteroids are a class of steroid hormones involved in a wide range of physiological processes, including stress response, immune response and regulation of inflammation. Particularly, glucocorticoids are potent anti-inflammatory corticosteroids, regardless of the inflammation's cause, whose effect is mainly through a lipocortin-1 synthesis mechanism[39, 40]. Some examples of the mostly used glucocorticoids for the treatment of inflammation are hydrocortisone and derivatives, prednisolone, dexamethasone or budesonide. Their therapeutic strength is extensively recognized, but they inexorably exhibit severe side effects after few weeks of treatment (such as weight gain, high blood pressure, high blood sugar, or even cataracts, bone loss, and muscle weakness)[41, 42] limiting their clinical use. Thus, various nano-formulations of corticosteroids[43] were proposed to overcome those limitations (Figure 3, Table 1).

Prednisolone is one of the most popular corticosteroids used for encapsulation into nanocarriers. For example, prednisolone nanocapsules, composed of a polymer wall surrounding an oily core, and coated with a pH-responsive enteric polymer (Eudragit S100), were designed and tested in healthy rats. This oral system allowed a high encapsulation efficiency of ca. 90% and a drug release triggered into the colon, as a consequence of neutral pH. This should increase the selectivity and efficacy of prednisolone for the treatment of inflammatory bowel diseases (IBD) or other colonic diseases[44].

Similarly, for oral administration, spherical polymer microsponges with a size of around 30 μ m, but with nano-porous microstructures, were also loaded with prednisolone and coated with Eudragit S100. The microsponges structure facilitated plastic deformation which was needed for the formation of mechanically strong tablets applicable for clinics. In addition, these microsponges displayed a high drug loading (around 86%), allowing dose and dose frequency reduction which should decrease prednisolone side effects and have some interest for the treatment of IBD[45]. However, the proof of concept was only validated *in vitro*, while *in vivo* studies are still lacking to validate this approach.

Another interesting strategy, based on [orally or/and rectally administered](#) lectin-decorated glucocorticoid loaded nanoparticles, was developed for an active targeting and selective adhesion to the inflamed tissue *in vitro* in experimental colitis. This strategy seems to be a promising tool for treatment of inflammatory bowel diseases[46]. Other recent pre-clinical studies have further demonstrated the therapeutic added value of prednisolone nano-formulations. For example, prednisolone loaded liposomes were assessed in a 8-week-old male LEWIS rat model of renal ischaemia-reperfusion injury, and liposomes were found to

accumulate in the inflamed kidney after systemic delivery. The treatment increased the presence of anti-inflammatory macrophages and reduced monocyte chemoattractant protein-1 mRNA production, concomitant to a reduced pro-inflammatory profile in the kidney[47]. Although leading to very interesting data, it is important to note that the experiments were performed with young age animals and the final point of analysis remained quite short (*i.e.* 96 h). And young animals respond more efficiently than older ones, whereas, in the daily medical practice elderly people represent the population of patients the more affected by inflammatory diseases.

Rheumatoid arthritis (RA) is a chronic, systemic inflammatory disease treated with high doses of glucocorticoids. To avoid non-specific body biodistribution, solid-lipid nanoparticles were coated with hyaluronic acid in order to target hyaluronic receptor CD44 over-expressed on the surface of synovial lymphocytes, macrophages and fibroblasts in inflamed joints in RA. After intravenous injection into 6-week-old mice with collagen-induced arthritis, a significant improvement of the disease has been obtained, likely due to the specific accumulation of NPs into the inflamed tissue[48]. Folate-targeted double liposomes (*i.e.* small unilamellar vesicles encapsulated into a big liposome) containing two therapeutic agents (Prednisolone and Methotrexate) were also developed and assessed by systemic administration in collagen-induced arthritis in rats. These liposomes displayed high stability, enhanced loading capability and further enhanced the site-specific drug delivery to activated macrophages at inflamed joints in RA[49]. Gaafar *et al.* designed elastic niosome (“ethoniosomes”) nanoparticles loaded also with prednisolone for ocular delivery. The ethoniosomes composed of Span60 and cholesterol and using ethanol 20% v/v as a hydrating solution to the surfactant/lipid materials were tested in rabbits subjected to a clove oil-induced severe ocular inflammation. A reduction to half of the time for complete healing was observed, in addition to a reduced intraocular pressure, one of the main side effects observed with conventional anti-inflammatory treatments. Besides, this system demonstrated good physical stability for at least 2 months at the fridge temperature with good ocular tolerability and minimal ocular irritation, suggesting good potential for translation into the clinic[50].

Methylprednisolone, already used in clinics for spinal cord injury treatment, is unfortunately accompanied by dose-related side effects due to systemic injection. The loading of methylprednisolone sodium succinate (MPSS) into polycaprolactone NPs dispersed in a fibrin glue-based gel system was tested in a rat model of induced spinal cord injury either by topical or intraperitoneal administration. The localized delivery of MPSS onto the lesion site enabled

a reduced damage on spinal cord[51]. This therapeutic effect was explained by a dramatic decrease of caspase-3 and a moderate decrease in pro-inflammatory cytokines.

Dexamethasone (Dex) is another glucocorticoid drug with anti-inflammatory activity. A recent study demonstrates that dendrimer-Dex gel attenuated corneal inflammation more efficiently than free-Dex, by decreasing macrophage infiltration and pro-inflammatory cytokines expression, in a rat corneal inflammation model. The designed system is a biocompatible hydrogel conjugated to Dex and based on hydroxyl-terminated polyamidoamine dendrimers and hyaluronic acid, cross-linked *via* thiolene click chemistry for subconjunctival injection[52]. Noteworthy, a single subconjunctival injection of this nano-formulation led to a prolonged efficacy for a period of 2 weeks with a reduced central corneal thickness and an improved corneal clarity, without elevation of the intraocular pressure. Moreover, the subconjunctival route is clinically accessible and can allow the administration of a relatively large volume of nanomedicine. [Dexamethasone nanomedicines could also be very efficient for the treatment of rheumatoid arthritis or ulcerative colitis, bypassing the serious adverse effects associated with important doses of drug free.](#) Thus, Wang *et al.* developed self-assembled poly(ethylene glycol) / poly(ϵ -caprolactone) polymer micelles loaded with Dex for intravenous injection at low dose of Dex (0.8 mg/kg body weight) in a rat model of arthritis. Although displaying moderate adverse effects, these polymeric micelles demonstrated substantial anti-inflammatory potential, including suppression of paw swelling and erythema, down-regulation of pro-inflammatory cytokines expression and protection of articular cartilage and bone from degradation and erosion[53].

Spherical polymer nanoconstructs loaded with Dex were also intravenously injected to a mouse model of colitis. A strong systemic anti-inflammatory effect was observed through a reduced macrophage infiltration and the expression of pro-inflammatory cytokines. These cellular effects were accompanied by a better animal recovery manifested by a reduced weight loss and a decreased rectal bleeding. These benefits were less evident in animals treated with the drug free[54]. Furthermore, using the same model, Dianzani *et al.* assessed the efficacy of [orally administered](#) multi-drug nanoparticles co-loaded with Dexamethasone and Butyrate. Both *in vitro* and *in vivo*, the treatment induced a significant decrease of pro-inflammatory cytokine release, which was more effective at doses 10-fold lower than the dose required to achieve the similar pharmacological efficacy with the single free drugs treatments[55]. If delivering simultaneously multiple drugs may be interesting, further research is, however, necessary to better understand the exact mechanism underlying the enhanced anti-inflammatory effect of this multidrug nano-formulation. [Very recently, by using an interesting](#)

approach, Wang's team optimized a prodrug composed of N-(2-hydroxypropyl) methacrylamide (HPMA) copolymer linked to dexamethasone for the treatment of inflammatory arthritis. By intravenous injection in a rat model of arthritis, they observed very effective and safe therapeutic anti-inflammatory effects compared to other prodrug formulations. This attractive study promoted the importance of identifying the most potent therapeutic nanomedicines by evaluating the design and screening of activation mechanism[56].

Budesonide is a topical anti-inflammatory synthetic steroid, which was formulated for oral administration as first-line therapy in mild to moderate ileocecal Crohn's disease patients[57]. In order to minimize early drug release and activity in the stomach and small intestine, Naeem *et al.* encapsulated Budesonide into dual pH- and time-dependent polymer nanoparticles composed of Eudragit FS30D, as a pH-sensitive polymer, and Eudragit RS100, as a controlled-release polymer. It was shown that in a mouse model of colitis, dual pH/time-dependent systems had greater therapeutic potential than single Eudragit FS30D or single Eudragit RS100 system. This beneficial effect was explained by the release of Budesonide at the right place (*i.e.* colon) during the right time (*i.e.* 24h)[58]. The use of Eudragit with various functionalities, either pH-sensitive or for controlled release, opens interesting perspective for clinical applications, particularly for colon-targeting treatments such as in IBD. Thus, nanotechnology has been used in oral dosage formulation design as strategies to further enhance uptake into diseased tissue within the colon [59]

Other nano-formulations to target and decrease inflammation are in development. For example, Budesonide loaded into Hyaluronic Acid (HA) or poly(lactic-co-glycolic acid) (PLGA) nanoparticles exhibited higher anti-inflammatory effects than the free drug *in vitro* on human colon carcinoma Caco-2 cell line and on inflamed intestinal mucosa in mice respectively[60, 61].

Together these preclinical results underline the feasibility to improve the delivery of anti-inflammatory drugs *in vivo*. However, to be closer to the human diseases, the choice of the animal model, the duration of the treatment and the end points deserve to be better thoughtful, as will be discussed below (see part 4).

2.2. Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) nanomedicines

Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) act through the inhibition of cyclooxygenase proteins that metabolize arachidonic acid into prostaglandins[62]. Due to

short half-life and high percentage of protein binding, high doses of NSAIDs are needed to be efficient, which in turn causes undesirable side effects, such as an increased risk of gastrointestinal and cardiovascular complications[62, 63]. Some NSAIDs NPs have recently been in development in order to counteract these side effects but, for the moment, they are only investigated at an *in vitro* pre-clinical stage.

Gonçalves *et al.* have loaded Diclofenac, a NSAID that inhibits prostaglandin synthesis by reducing cyclooxygenase-1 and cyclooxygenase-2 with relative equivalence, into chitosan-poly(γ -glutamic acid) NPs. This approach did not show any cytotoxicity for human macrophages. NPs were rapidly phagocytosed and activated these cells, also inhibiting prostaglandin E2 and IL-6 production[64]. More recently, Shang *et al.* designed a novel intestinal targeted drug delivery system for oral administration; this system that combines PEG₉₅₀ and methacrylic acid (MAA)) hydrogel and Ibuprofen-loaded PLGA NPs, could site-specifically deliver the drug into alkaline environment at a sustained-release manner. Such process improved the bioavailability and decreased the side effects of Ibuprofen[65].

Celecoxib is a COX-2 inhibitor indicated for long-term treatment of rheumatoid arthritis. It is administered orally, but unfortunately, this molecule shows adverse systemic complications in patients. In rats with experimental arthritis, the topical application of a gel containing this anti-inflammatory drug into solid lipid nanoparticles was found more efficient in inhibiting the inflammatory process, compared to untreated or conventional gel treated animals[66].

2.3. Nano-encapsulated anti-inflammatory mediators

Anti-inflammatory mediators represent another approach for the treatment of inflammatory diseases. For instance, cytokine IL-10, one of the main anti-inflammatory cytokines possesses multiple effects in immuno-regulation and inflammation[67]. It mainly downregulates the expression of other pro- or anti-inflammatory cytokines, enhances immune B cell survival, proliferation, as well as, antibody production. Besides, IL-10 is implied in blocking NF- κ B activity and is involved in the regulation of the JAK-STAT signalling pathway. But its clinical use is complicated by important cytokine-related side effects [68]. To overcome these limitations, new IL-10 anti-inflammatory nano-formulations were developed (Table 1).

A typical example is the development of nano-sized phosphatidylserine-containing liposomes (PSL) conjugated to IL-10. Toita *et al.* showed in a mouse model of obesity that intraperitoneal injection of IL-10-conjugated PSL had a better macrophage targeting ability and an enhanced anti-inflammatory effect than PSL alone, due to the synergistic anti-

inflammatory effects of IL-10 and PSL. It was suggested that this association could be used as a macrophage-targeted therapy for inflammation-related diseases[69].

Another example is the development of biodegradable polyester NPs incorporating IL-10 for the targeted delivery of atherosclerotic plaques. Intravenous injection of these NPs significantly reduced acute inflammation in an atherosclerotic murine model and were shown to be more potent than IL-10 free. In addition, it was observed that plaque rupture was prevented by increasing the fibrous cap thickness, simultaneously to the decrease of the necrotic cores[70]. More recently, Baganizi *et al.* have bioconjugated recombinant IL-10 to poly(vinylpyrrolidone)-coated silver NPs. These nanoparticles showed an improvement of the cytokine stability and a better anti-inflammatory effectiveness in mouse macrophages as manifested by a decrease in IL-6 and TNF α production[71]. However, this study was limited by the fact that silver NPs could not be delivered *in vivo* for inflammation treatment. Indeed, at the cellular level, toxicities have been reported, including reactive oxygen species generation, DNA damages and cytokine induction during *in vitro* studies[72, 73]. *In vivo*, studies also indicated adverse effects on many organs of the circulatory, respiratory, central nervous, hepatic and dermal systems[74, 75].

Other studies focused on the nano-formulation of antagonists of the pro-inflammatory cytokine IL-1 receptor. Particularly, self-assembled IL-1 receptor antagonist (IL-1Ra)-presenting nanoparticles were developed. Two examples of this kind of NPs are: IL-1Ra-poly(2-hydroxyethyl methacrylate)-pyridine NPs and IL-1Ra core/satellite NPs, based on methyl viologen-functionalized polymeric NPs as the core, and a trans-isomer of azobenzene modified IL-1Ra as the corona. The pharmacological efficacy of these nanodevices were investigated after local injection in a rat model of inflamed joints[76,77] and both IL-1Ra-presenting nanoparticles demonstrated interesting anti-inflammatory effects, mainly inhibiting the IL-1 β -stimulated pro-inflammatory production.

2.4. Anti-inflammatory peptides nanomedicines

Anti-inflammatory peptides, which are small bioactive molecules containing up to 50 amino acids, are raising increasing interest[78]. For example, the anti-inflammatory peptide KFAK, an inhibitor of mitogen-activated protein kinase-2, was assessed in a cartilage explant inflammation model. To increase the peptide's therapeutic lifetime, KFAK was first loaded into co-polymerized 2-acrylamido-2-methylpropane sulfonic acid (AMPS) and N-isopropylacrylamide (NIPAM) monomers, to form pH-dependant poly(NIPAM-co-AMPS) nanoparticles[79, 80]. These NPs demonstrated the ability to selectively diffuse through the

degraded cartilage explants before to release the peptide, allowing a significant reduction of pro-inflammatory cytokine production. However, the authors noted that the high affinity of KFAK towards the nanoparticles core and the lack of core degradation resulted in the release of less than half of the loaded peptides, thus limiting the therapeutic applications of this approach[80]. Then, the same authors developed a more complex delivery system, based on poly(NIPAM) nanoparticles with degradable disulfide crosslinks (abbreviated as NGPEGSS), loaded with KFAK peptides. The system relies on the co-polymerization of NIPAM and AMPS monomers, coated with polyethylene glycol (PEG) and using the degradable crosslinker bis(acryloyl)cystamine (BAC) for more effective release into the intracellular compartment[80, 81]. *Ex vivo*, a preferential accumulation of the NPs was observed throughout the inflamed aggrecan-depleted cartilage explants compared to their healthy counterparts and, simultaneously, a significant reduction of IL-6 pro-inflammatory cytokine was measured. Moreover, these NPs were stable and released less than 10% of the loaded KFAK over 96h at pH=7.4 in a non-reducing environment. Thus, their ability to diffuse into the damaged cartilage, maintaining the majority of their payload prior to cell uptake by chondrocytes and macrophages, combined with a release of the peptide into these cells, represented an improvement comparatively to the treatment with the free peptide or with the previously described poly(NIPAM-co-AMPS) NPs[81].

2.5. Nanomedicines for gene therapy

Another strategy to control the inflammation is based on the manipulation of genetic material. In recent years, significant progresses have been made to design gene delivery systems for obtaining enhanced therapeutic efficacy with less off-target side effects. It is expected that targeted gene therapy could reduce the above-mentioned limitation by localizing the targeted gene at the site of the inflammation. Small interfering RNAs (siRNAs) were found to be a promising approach for silencing specific genes involved in inflammation. However, siRNA delivery into the cytosol of immune cells remains a challenge.

Many different mediators were considered for the development of gene nano-therapies (Table 1). For example, siRNA silencing proinflammatory cytokine tumor necrosis factor (TNF α) expression was entrapped into mannose-modified trimethyl chitosan-cysteine nanoparticles. *In vitro*, the NPs were internalized by rat peritoneal exudate cells, through endocytosis and micropinocytosis pathways. But more interestingly, in a rat model of acute hepatic injury, TNF α siRNA-loaded NPs orally administered induced an effective TNF α knockdown at a very low dose (*i.e.* 50 μ g/kg)[82]. Along the same line, cationic lipid nanoemulsions were also

proposed for the delivery of anti-TNF α siRNA to the brain. It was shown that after intranasal administration in a rat model of LPS-induced neuroinflammation, the nanoemulsion concentrated preferentially into the brain, inducing a local down-regulation of TNF α [83]. Similarly, poly(ethylene glycol)-poly(caprolactone) polymeric micelles conjugated with anti-TNF α siRNA were administered intranasally in a rat model of cerebral ischemia-reperfusion injury associated with neuroinflammation. The rats subjected to the micelles treatment exerted a decrease of TNF α production and a significant improvement of their neurological score, as compared with the treatment using naked anti-TNF α siRNA[84].

Very recently, intranasal administration of cationic phosphorus dendriplexes loaded with anti-TNF α siRNA were shown very effective in a murine acute lung inflammatory model. Indeed, the authors observed a specific targeting toward macrophages, resulting in a major inhibition of TNF α , as well as, a better anti-inflammatory regulation of other cytokines[85].

Since in inflammation sites there is an overexpression of folic acid receptors by macrophages, a modified chitosan nanocarrier has been constructed, deploying folic acid, diethylethylamine and PEG groups to deliver TNF α siRNA. A significant decrease of inflammation in a murine arthritis model, close to human disease, was observed in animal receiving intraperitoneal injections of these NPs[86]. Nevertheless, the observed results should be discussed since the effects were observed only after multiple days of treatment, whereas TNF α levels usually tend to decrease rapidly (within a day) after siRNA treatment.

In order to downregulate TNF α and reduce RA progression, Aldayel *et al* designed PEGylated solid-lipid nanoparticles composed of lecithin and cholesterol and loaded with lyophilized TNF- α -siRNA. When this formulation was intravenously injected in a mouse model of collagen antibody-induced arthritis refractory to methotrexate therapy, the results obtained showed a good treatment efficacy, opening the hope for patients refractory to methotrexate[87]. Similar results have been obtained with lipid-polymer hybrid nanoparticles loaded with siRNA directed against TNF α . Intra-articular injection, in an experimental arthritis mouse model, showed that this nanotherapy could efficiently suppress the inflammatory process, this effect being obtained with a low TNF α -siRNA dose (*i.e* 1 μ g)[88].

Gene therapy was also evaluated on a different aspect of inflammation, relying on the control of macrophage polarization. Indeed, M1 macrophages are pro-inflammatory and have a central role in host defence against infection, while M2 macrophages are associated with responses to anti-inflammatory reactions and tissue remodelling[89]. Therefore, re-polarization of macrophages from the M1 phenotypes to M2 phenotypes by plasmid DNA

(pDNA), siRNA or miRNA, represents an exciting prospect for the treatment of inflammatory diseases. In this context, IL-10 pDNA was encapsulated into alginate-based nanoparticles decorated with tuftsin, a tetrapeptide located in the heavy chain of immunoglobulin G which actively targets macrophages[90]. After **intraperitoneal** administration, these NPs were found to concentrate in inflamed paws of arthritic rats, due to a continuous recruitment of macrophages in the inflamed tissues, and a sustained local and systemic IL-10 expression was observed. Another nano-formulation of IL-10 pDNA was designed, using hyaluronic acid-poly(ethyleneimine) (HA-PEI) NPs. In this study, IL-10 pDNA and IL-4 pDNA, who are both supposed to play a key role in the polarization of macrophages, were encapsulated into these NPs. It was found that the *in vitro* transfection occurred in peritoneal macrophages over-expressing CD44 receptor and increased M2 phenotype surface markers due to the expression of IL-4 and IL-10. Besides, the intraperitoneal administration of these NPs to LPS-induced inflamed mice also significantly increased the M2 markers expression, showed elevated levels of serum IL-10 and reduced inflammation[91]. When microRNA-223, which is a miRNA that has been proved to alter macrophage polarization and activation, was encapsulated in the same nanocarrier[92], similar re-polarization of macrophages and anti-inflammatory effects were observed *in vitro*[93].

Other similar approaches are described in Table 1. But the crucial message here is that if gene nanotherapy could be effective in pre-clinical rodent models, it remains urgent to validate the proof-of-concept in bigger animal models, closer to human diseases.

2.6. Biomimetic nanoparticles

When injected intravenously, most of the nanomedicines are rapidly captured and eliminated by the immune system if they are not PEGylated. Biomimetic nanoparticles are hybrid nanostructures in which the uppermost layer is similar to a cell membrane (*e.g.* erythrocytes, immune cells, cancer cells and platelets)[94, 95]. In this category are included cell membrane-coated nanoparticles and liposomes engineered with cell membrane proteins. This strategy brings the biological characteristics of the source cells and leads to an exhibited prolonged circulation of the nanoparticles and an active targeting capability; moreover, these biomimetic nanoparticles are less likely to be recognized by the immune system since the membrane coating has the inherent characteristics of the mother cells. Excellent reviews have been published in this field [96-99] and we cover here only the very recent publications.

Current clinical treatment of rheumatoid arthritis primarily targets the inflammatory response by using anti-cytokine biologics such as those inhibiting tumour necrosis factor alpha (TNF-

α) and interleukin (IL-1). However, the response rate still remains unsatisfactory[100]. Since neutrophils play an important role in resolving inflammation and repairing tissues damage[101], in a very interesting approach Zhang *et al.*, fused neutrophil membrane onto polymeric cores and injected these hybrid biomimetic nanoparticles [into the knee joint](#) in a mouse model of collagen-induced arthritis and in a human transgenic mouse model of arthritis. The results showed a significant therapeutic efficacy by reducing joint damage and suppressing overall arthritis severity[102]. Of note, the authors used a biodegradable polymer which is FDA approved (*i.e* PLGA), to prepare the nanoparticles core. In another strategy, macrophage membrane-coated polymeric nanoparticles have shown to bind and neutralize endotoxins and to inhibit systemic inflammatory response in a mice lipopolysaccharide (LPS) model of inflammation [after intravenous injection](#)[103].

In another study, biomimetic nanoparticles were synthesized using membrane proteins purified from activated J774 macrophages and loaded with rapamycin. In a model of ApoE^{-/-} mice with atherosclerotic plaques, it was observed that [systemic injection](#) of these hybrid NPs was able to suppress cell proliferation within the aorta[104].

Other strategies have been developed such as leukocyte-mimetic liposomes with the ability to translocate through inflamed endothelial cell layers[105]. Macrophage-derived microvesicles-coated PLGA nanoparticles were able to target rheumatoid arthritis (RA) [after intravenous injection](#), displaying significant enhancement of the therapeutic efficacy in a collagen-induced arthritis mouse model. Similarly, macrophage-coated nanoparticles encapsulating tacrolimus significantly suppressed the progression of RA in mice[106].

Together these studies suggest that hybrid biomimetic nanomedicines represent a new and promising approach, since they possess unique biodistribution characteristics exhibited by different cell types used and, flexible designs derived from adaptable nanoparticle cores. However, some important concerns remain including possible immune reactions and complement activation after their administration. Besides, the scaling-up and the sterilisation of these biomimetic nanodevices also require specific developments.

2.7. Other applications of nanoparticles for immune system modulation

Strategies to activate or inhibit receptors implied in the inflammatory response are also potential ways to reduce inflammation (Table 1, Figure 3). One example is the liver X receptors (LXRs), whose activation in macrophages is atheroprotective and suppresses inflammation[107]. Thus, some recent studies focused on the formulation of LXR agonists in macrophage-targeted nanoparticles, such as mannose functionalized dendrimeric NPs[108],

biodegradable diblock PLGA-b-PEG copolymer NPs[109], or synthetic high density lipoprotein NPs[110]. **Intravenous injection** of all these different nano-formulations were found to increase the expression of LXR target genes, to exert anti-inflammatory effects and to inhibit the development and necrosis of plaques in LDLr-deficient and ApoE-deficient mouse models of atherosclerosis. Particularly, these nano-formulations allowed an enhanced cholesterol efflux[108-110], a reduction of CD68-positive macrophage content of plaques by 50%[109] and an up-regulation of expression of ATP-binding cassette transporters[110].

Nanomedicines with immunosuppressants are also in development for improving the pharmacological efficacy and reducing the adverse events of these molecules. **Topical administration** of tacrolimus (FK506) loaded onto monolein nanoparticles were tested in a psoriasis-like skin inflammation in mice, and appeared to be more efficient than tacrolimus free dissolved in propylene glycol[111]. **Positively charged Tacrolimus-loaded nanomicelles** were found to prolong eye surface retention, to enhance corneal permeability, to exert anti-apoptotic effects on the corneal epithelium and overall to suppress expression of inflammatory related factors in a dry eye disease rabbit model.[112]. Besides, PEGylated liposomes loaded with the immunosuppressive drug FK506, were found to inhibit the expression of cytokines, such as interferon- γ and tumor necrosis factor- α , and reduced inflammation and fibrosis of the myocardium in a rat experimental autoimmune myocarditis after systemic administration[113]. Liposomal nano-formulations were also used to encapsulate cyclosporine A, another immunosuppressive drug. For example, **intravenous administration** of liposomal-cyclosporine A could improve cerebral ischemia and reperfusion injuries like neuroinflammation in a rat model, by inhibiting the inflammatory responses including MPO activity and TNF α level[114]. A further study on neuroinflammation in rats also proved the anti-inflammatory efficacy of cyclosporine-A using a cationic oil-in-water nanoemulsion upon intranasal administration[115].

Studies on mesalazine (5-ASA), an aminosalicylate anti-inflammatory drug mainly used in inflammatory bowel disease, focused in ameliorating the contact between the drug and the inflamed tissues by increasing adhesion and retention time. Practically, 5-ASA loaded onto silicon dioxide nanoparticles were tested in a dextran sodium sulfate-induced mouse model of ulcerative colitis. A significant decrease of IL-6, TNF- α and MPO was noted after **oral** administration of these NPs[116]. Along the same line, a very recent study designed an inclusion complex of 5-ASA in hydroxypropyl- β -cyclodextrin loaded onto chitosan NPs. This delivery system exhibited a sustained-release profile, and the NPs were more effective than the free drug in inhibiting the secretion of pro-inflammatory cytokine-stimulated colon cancer

cells[117]. In another study, polymeric nanoparticles were [rectally administered](#) to protect low molecular weight heparin (LMWH) from intestinal degradation and to provide targeted delivery to inflamed tissue in experimental colitis mice. Such a combination shows a highly specific therapy by its protection against degradation in luminal environment and selective drug delivery inflamed intestinal tissue[118].

An unusual approach consisted in the design of prodrugs that will spontaneously self-assemble as nanoparticles in solution. This is what our group has developed through the construction of nanoparticles made of squalene, a natural and biocompatible lipid precursor of the cholesterol's biosynthesis. Of note, this lipid is already used in clinic as vaccine adjuvant. Squalene has been employed as a building block for the bio-conjugation with different hydrophilic drug molecules, and demonstrated the ability to self-assemble in aqueous medium in the form of nanoparticles[119, 120]. In general, these squalene-based nanoparticles were found to be non-cytotoxic, exhibited a high drug payload, an enhanced therapeutic response, and they were quite simple to prepare. Based on this concept, very recently, we discovered a new painkiller nanomedicine consisting in leu-enkephalin conjugated to squalene. And, the nanoparticles showed prolonged anti-pain effects on inflammatory pain in rats. Since they were found to target the inflamed area after intravenous administration and to act only on the peripheral opioid receptors, this nanomedicine opens the door to the treatment of intense pain without inducing tolerance and addiction, as observed with morphine and related synthetic opioids. The targeting of the inflamed area was found to occur by the EPR-like effect[121].

Squalene has also been conjugated to adenosine, a crucial regulatory autocrine and paracrine factor with anti-inflammatory properties[122, 123]. Adenosine is physiologically presents at low levels in unstressed tissues, and is rapidly released from the intracellular space to the extracellular space in response to pathophysiological conditions, such as hypoxia, ischemia, inflammation or trauma, to generate various cellular responses that aim to restore tissue homeostasis[122, 123]. Nevertheless, the use of adenosine in clinic is limited because of a very short plasma half-life and the need to administer high dosages which induce important side effects. Squalenoyl-adenosine nanoparticles may therefore represent a promising approach in the treatment of inflammatory disorders surpassing adenosine limitations. [Intravenous administration](#) of these nanoparticles already demonstrated a significant neuroprotective effect in experimental models of stroke in mice and spinal cord injury in a rats, where inflammation is part of the diseases[124].

Since inflammation is accompanied by the overproduction of radical oxygen species[125], molecules with antioxidant capacities have been formulated as nanoparticles in order to

increase the targeting of inflamed tissues and improve drug bioavailability. For example lycopene, a carotenoid drug with antioxidant and anti-inflammatory activity[126] and glutathione[127] were both tested in animals with experimental rheumatoid arthritis. *Wu et al.* also developed andrographolide-loaded poly(ethylene glycol) and poly(propylene sulphide) micelles which showed to synchronically alleviate inflammation and oxidative stress in an ApoE^{-/-} atherosclerotic mice model[128]. In addition, our group recently developed a novel antioxidant/anti-inflammatory multidrug squalene-based nanoparticle formulation, allowing simultaneous delivery of adenosine and tocopherol in tandem to targeted sites by exploiting dysfunctions of the endothelial barrier at the sites of acute inflammation. These nanoparticles demonstrated promising anti-inflammatory and protective effects in mice models of endotoxemia and lethal systemic shock, with potential to treat uncontrolled inflammation associated with many diseases, including severe COVID-19[129].

Another interesting example is represented by metallic nanoparticles, the archetype being gold nanoparticles. The use of gold for therapeutic purposes dates back to Antiquity in China and Egypt, but only recently the scientific community developed a strong interest in gold nanoparticles for biomedical applications, especially because they can be easily functionalised and possess imaging and therapeutic capabilities[130]. The anti-inflammatory property of nanogold was recently explained by the down regulation of inflammatory mediators like TNF- α , IL-1 β , COX-2 and transcription factor NF-kB (Nuclear factor-kB)[131]. Although systemic administration of these nanoparticles produced a significant anti-inflammatory activity *in vivo*, on different mouse models[132, 133], it has to be noted that gold, even if well tolerated by the body, is not a biodegradable material and may remain for a long time into the body, as previously demonstrated in a mouse model[134].

Apart from being used as drug delivery platforms, NPs have also been engineered to modulate the immune system. The induction of antigen-specific adaptive immunity exclusively occurs in lymphoid tissues, where T and B cells meet antigen or antigen-loaded APC (dendritic cells, macrophages). Since dendritic cells can activate T cells, vaccination protocols targeting DCs are promising approaches to enhance immune responses. When injected into the blood, nanoparticles interact most frequently with antigen-presenting cells (APCs). Among the APCs, dendritic cells (DCs), which are the most specialized, are capturing and processing antigens before migration to lymphoid tissues, allowing T or B cells activation. By this way, nanoparticles may allow modulating the immune response. For example, De Koker *et al.* engineered hybrid nanoparticles composed of silica- poly(methacrylic acid) polymer. When

injected *in vivo*, these NPs target and accumulate in lymph nodes, suggesting a possible application for the lymphatic delivery of antigens and immune-modulation[135]. And Muller *et al.* designed a versatile nanoparticle vaccine platform to promote an antigen-specific humoral response. These NPs were able to sustain a prolonged antigen presentation to APCs and to produce a strong immune response, as compared to the free antigen. Moreover, the size, charge, and surface characteristics of the NPs were found to be key parameters for improving the lymphatic trafficking of the NPs and their subsequent uptake by APCs[136].

To overcome resistance or production of antibodies which may occur after treatment with Rituximab (anti-CD20), another interesting approach was developed by Kopeček *et al.* who developed polymer prodrugs to target B cells. This approach, initially tested for B-cell malignancy treatment[137-140], was recently evaluated in a pre-clinical model of inflammatory rheumatoid arthritis (RA)[141]. The treatment, tested in a collagen-induced rheumatoid arthritis in mice, was composed of two conjugates: (i) a bispecific engager, Fab'-MORF1 (obtained by linking Fab' fragment of CD20 mAb to the 3' end of a morpholino oligonucleotide (MORF1) via a thioether bond) and (ii) a receptor crosslinking mediated by HPMA-based P-(MORF2). Compared to untreated mice, a delayed progress of RA was observed. And this new approach could open the way for the treatment of other autoimmune diseases.

However, this research area based on the modulation of the immune system has mainly been developed for cancer immunotherapy (see reviews[142-144]), while the studies reporting the modulation of immune system in other inflammatory diseases are less. In this sense, Gao *et al.*, designed an innovative antibacterial vaccine. The authors used *Escherichia coli* as a model pathogen membrane and coated them onto small gold nanoparticles (AuNPs) with a diameter of 30 nm. When injected subcutaneously, the NPs induced rapid activation and maturation of dendritic cells in the lymph nodes of the vaccinated mice, with in addition, the generation of durable antibody responses. These results demonstrated that using natural bacterial membranes to coat synthetic nanoparticles holds great promise for antibacterial vaccines[145]. Autoimmune diseases are caused by antigenically complex immune responses of the adaptive and innate immune system against specific cells, tissues or organs. Type 1 diabetes is a T cell-dependent autoimmune disease that is characterized by the destruction of insulin-producing β cells in the pancreas. Yeste *et al.*, engineered nanoparticles to deliver a tolerogenic molecule, for inducing a tolerogenic phenotype in DCs. A significant delay of diabetes onset was observed in nonobese diabetic mice receiving this immunomodulatory

nanoformulation[146]. For the future, a deeper understanding on the NP-immune cell crosstalk could guide the development of nano-immunotherapeutic approaches with an increased therapeutic efficacy.

3. Clinical trials and commercial markets

Many nanomedicines, mainly liposomes, have entered clinical practice, and even more are being investigated in clinical trials for a wide variety of therapeutic and imaging indications. To date, only few tenths of nanomedicines are FDA approved and few hundreds of clinical trials are investigating new nanomedicines[147], cancer therapy being the main application. Recently, three other nanomedicines have been approved: Patisiran/ONPATTRO, VYXEOS, AND NBTXR3/Hensify[148]. However, to the best of our knowledge, very few anti-inflammatory nanoparticles have succeeded to reach clinical trials (Table 2).

Anti-inflammatory liposomal formulations are one of the most promising nanomedicines for the treatment of acute and chronic inflammation. For example, after intravenous administration, long-circulating prednisolone-containing liposomes were found to accumulate into macrophages of patients with symptomatic iliofemoral atherosclerosis, as a result of prolonged circulation half-life. Nevertheless, in contrast to the pre-clinical efficacy data obtained in rabbits[149], short-term treatment did not reduce arterial wall permeability or arterial wall inflammation in humans[150-152]. This may have resulted from an insufficient dose of prednisolone reaching the plaque; and although increasing the dose will likely resolve this issue, it would also introduce the risk of adverse effects. Another possible explanation may be a timing issue since, based on pre-clinical results, the authors evaluated the clinical results only after 10 days of treatment. But most studies evaluating the impact of anti-inflammatory agents in human cardiovascular diseases showed therapeutic efficacy only after several weeks of treatment [150]. Despite the lack of anti-inflammatory activity, the study proved for the first time that targeted delivery of nanomedicines to atherosclerotic lesions is feasible in humans. This could provide guidance for the development of future nano-formulations for the imaging or the treatment of atherosclerosis.

Interestingly, such prednisolone-encapsulated liposome formulations were also tested in phase II clinical trials for the treatment of arteriovenous fistula[153], graves ophthalmopathy, and ulcerative colitis[154]. Recently, the clinical study for the treatment of ulcerative colitis exhibited strong positive results in 70% of the patients, with 4 patients out of 13 in remission, and 7 out of 13 showing a reduction of the endoscopy sub-score[155]. In addition, a phase III

clinical trial for the treatment of rheumatoid arthritis has started but no results are available for the moment.

A liposomal formulation of cyclosporine A in a lipogel was also developed for a topical application to treat mild to moderate stable plaque psoriasis and tested in an early-phase clinical trial[156]. The clinical performance of this liposomal gel was superior to that of cyclosporine incorporated in a conventional dermatological formulation at equal drug concentration. Pharmacokinetic studies and other safety evaluation protocols indicated the good tolerance of the liposomal formulation which has induced uniform reduction of plaque elevation[156]. Other liposomal formulations are currently in development, such as liposomal corticosteroids or liposomal dexamethasone, whose clinical translation has started very recently in patients with progressive multiple myeloma[157].

Pitavastatin is a statin known to have vasodilating properties and to modulate inflammatory response. To evaluate the safety, tolerability, and pharmacokinetics for intravenous administration of Pitavastatin-loaded PLGA nanoparticles (NK-104-NP), a phase I clinical trial was recently conducted in healthy Japanese male subjects. The study proved that NK-104-NPs exhibited dose-dependent pharmacokinetics and was well tolerated with no significant adverse effects in healthy volunteers[158].

In a very recent pilot clinical study, an oil-in-water nanoemulsion containing all trans- retinoic acid (Tretinoin), showed a good efficiency against acne after topical application[159]. Similarly, when subjected to focal thermolysis of sebaceous gland by near infrared laser, silica-gold nanoshells (150 nm) coated with PEG, were effective to provide a significant reduction in occurrence and appearance of inflammatory lesions[160].

In conclusion, very few nanomedicines specifically designed for the treatment of inflammation are nowadays assessed in clinical trials. In order to operate translation from the bench to the bedside, several experimental challenges need to be addressed. Key issues related to the clinical development of anti-inflammatory nanoparticles include biological challenges such as understanding the biology of specific diseases and the biological interaction of nanoparticles in patients, commercialization hurdles related to large-scale manufacturing, to cost-effectiveness in comparison to current therapies and to regulatory standards, biocompatibility and safety[161]. Reducing complexity in nanoparticle design and synthesis should, therefore, be one of the main goals to generate clinically translatable nano-sized therapeutics.

4. General discussion

In recent years, cellular and molecular advances have improved the comprehension of the inflammatory response, leading to the discovery of multiple targets for biological therapies against inflammatory diseases. In addition to a variety of safe and quite effective anti-inflammatory agents already available, such as aspirin and other nonsteroidal anti-inflammatory drugs, many biological agents have also been developed including antibodies and nucleic acids for cytokine inhibition or to reduce the extravasation of immune cells into the inflamed tissues[162]. In spite of promising progresses made in developing such treatments, the widespread distribution of the inflammatory signaling pathways in multiple cell types and tissues leads to important off-target side effects and to systemic toxicities. Combined with a relatively low bioavailability or/and limited half-lives, necessitating repeated and expensive high dosages responsible for severe side effects, current therapies still remain with clinical hurdles to overcome.

This has prompted the academic research to design nanocarriers in order to improve the delivery of anti-inflammatory compounds. After administration, such nanomedicines follow a kinetic process of absorption, distribution, metabolization and elimination. In addition, they distribute in a different way in the body depending if their surface is naked, decorated with PEG, or loaded with a ligand for an active targeting. In this way, numerous nanoparticles loaded with various anti-inflammatory molecules were developed and have shown real therapeutic improvements in pre-clinical models (Figure 3).

Other than being drug delivery carriers, nanoparticles have also been used as vaccine adjuvant and for modulation of the immune system. Different data suggest that the shape of the NPs can affect the interactions with innate immune system. Once injected into the body the NPs immediately encounter the innate immune system and generate specific response as foreign substances. Apart that the coating with poly (ethylene glycol) delays the protein corona formation at their surface, the physicochemical properties of NPs (size, charge, hydrophobicity and stiffness) will also determine their interaction with the innate system.

For example, it was shown that the uptake of nanorods by macrophages was more efficient than the uptake of nanospheres[163]. And a comparison of the complement activation using spherical, rods and disks polystyrene NPs evidenced that rods and disks generated more important complement activation than spheres at later time points[164].

But today it remains still unclear which physical properties or chemical composition of nanoparticles can drive to immune modulation.

There is also an urgent need to develop new *in vitro* and *in vivo* pre-clinical models, better mimicking the pathophysiology of the disease in humans. For instance, the EPR effect has become during decades a general condition to allow the targeting of nanoparticles or liposomes through leaky vasculatures towards tumor or inflamed areas, in most of the experimental pre-clinical rodent models. But in the clinical reality, the occurrence of the EPR effect is far to be systematic. Various other parameters need to be taken into consideration, like the distribution and diffusion deep into the inflamed tissue (which is bigger in humans than in mice), the extend of drug release, the blood perfusion of the inflamed area, the nature of the extracellular matrix, *etc.* Thus, concerning the inflammatory processes, are the pre-clinical models really reliable? [And to what extent the observed vascular permeability in pre-clinical models is transposable to human?](#) Answering those questions and developing clinically relevant animal models is a key to allow a better selection of disease-driven nano-formulations[165].

Different administration ways, oral, local, topical or intravenous have been tested with anti-inflammatory nanomedicines. While intravenous delivery allows a direct access to the whole body, the local delivery proved to be effective, allowing to directly concentrate the drug into the diseased tissue. If the choice of administration route depends evidently on the localization of the inflammation site, it should also customize the choice of the materials and the design of the nanomedicines (presence or not of PEG or other ligands).

[The route of administration such as intravenous or the direct injection into the inflamed site can influence the induction of the immune response.](#) Thus, when NPs are administered intravenously, the protein corona formed at the surface of the NPs and the complement eventually activated can alter biodistribution and therapeutic efficacy. But when the NPs are administered locally, the immune response is mainly dependent on the tissue residential cells. Chemical and physical properties of the NPs, including size, surface charge, and surface chemistry, are other important factors in the context of systemic inflammation. For example, it was observed that LPS-induced inflammation resulted in splenic macrophage polarization and altered leukocyte uptake of polystyrene nanoparticles depending on their size[166]. Concerning local administration, poly(lactic acid) particles need a size of at least 3 μm to guarantee the retention in inflamed joints[167].

Of note, as detailed in this review, different types of nanocarriers have been employed for the delivery of anti-inflammatory drugs, including liposomes, polymer nanoparticles, micelles or prodrugs. To compare their delivery abilities is quite hazardous because the experimental

conditions are diverse. Liposomes are evidently the most advanced in terms of clinical translation and some liposomal formulations are even FDA approved. However, they generally display a lower stability in biological fluids than polymer nanoparticles or prodrugs, sometimes leading to a fast drug leakage. And their drug loading capacities are often limited. On the other hand, in case of intravenous administration, polymer micelles may disassemble due to a high critical micellar concentration which doesn't occur with polymer nanoparticles which may be considered as frozen micelles.

But as shown in this review, the clinical translation of the nanoparticles remains still limited. This is due, in part, to the complexity related to the pharmaceutical development of nanomedicines, including their biocompatibility and safety. We also note that chronic inflammatory diseases bring together confounding factors such as age, overweight and obesity and sometimes genetic predisposition to cite only a few. However, most of the preclinical studies to assess anti-inflammatory nanomedicines are performed on young and healthy animals, with an induced accelerated inflammatory disease and subject to normal fed. These factors should therefore be carefully considered in order to improve the translation from preclinical models to the clinic. In addition, the difference between rodent and human immune systems may represent an important obstacle for translating studies performed in mice into human clinical trials. This is the reason why there is an urgent need to develop pre-clinical experimental models with pathophysiological similarities closer to human inflammatory disease.

Bio-inspired nanodevices represent a new generation of innovative nanomedicines with the aim to mimic natural circulatory cells. They are able to increase the blood circulation time and to improve the distribution of the loaded drug towards the inflammatory cells and tissues. However, their immunogenicity, scale-up and characterization remain important hurdles before starting clinical trials.

Besides, problems related to the scaling-up, the government regulations and the overall cost-effectiveness if compared to the current available anti-inflammatory therapies are other important limitations in the success of the anti-inflammatory nanomedicines in general. The often complex architectural design of many nanomedicines described in this review, likely results in difficulties for performing reproducible sample preparations, safe and in good quantities[168]. And this is particularly important since a slight modification of the size, the shape and/or the nanoparticle surface chemistry may dramatically influence the stability, the

interaction with biological media, as well as, the biodistribution and clearance of the nanocarriers. Thus, reliable and standardized methodologies to obtain reproducible nanoparticles are absolutely required[169]. There is often a gap between the academic environment where innovative nanocarriers are designed and the industrial context, where reproducible preparation and manufacturing processes supported by analytical methodologies are needed.

5. Conclusion

As shown in this review, there is an increasing interest to improve the delivery of small drug molecules or biotherapeutics (*i.e.* nucleic acids, peptides, proteins) for the resolution of inflammatory diseases. This represents a unique opportunity for pharmaceutically developable nanoscaled drug delivery systems, especially to optimize drug safety and efficacy through improved biodistribution and pharmacokinetics and to resolve the issues related to the difficulty to administer fragile biomolecules.

Acknowledgments

The authors gratefully acknowledge the financial support from the 7th EuroNanoMed-II call for proposals, project NanoHeart n°ANR-16-ENM2-0005-01.

Conflict of interest

The authors declare no competing interest.

Author Contributions

The manuscript was written with the contribution of all the authors.

Figures and Tables Captions

Figure 1: Schematic representation of the inflammatory response.

Briefly, inducer signals (1) (*e.g.* wound, pathogens) trigger sensors (2) on inflammatory cells present in the damaged area, which start to induce the production and release of multiple mediators (3). Mediators, including plasma proteases, chemokines, cytokines, and vascular modification factors lead in turn to vascular modifications, recruitment of blood platelets and other inflammatory cells such as phagocytic leukocytes (*e.g.* neutrophils) and act on effector cells (4) and tissues to resolve inflammation. [Adaptive immunity is also implicated in inflammation response, through lymphocytes B and T able to recognize and respond to antigens presented by antigen-presenting cells such as dendritic cells. Adaptive mechanisms may function either by direct cytotoxic effects or by secretion of antibodies interacting with elements of the innate inflammatory response \(complement proteins, phagocytic cells...\).](#)

Figure 2: Anti-inflammatory nanomedicines: strategies and targets

Different types of nanoparticles were developed or are still in pre-clinical development for the management of inflammation. Among them liposomes, polymer nanoparticles, [micelles](#), dendrimers, or hydrogel-based formulations. These nanoparticles could be naked or functionalized with targeting moieties, such as coating groups and antibodies or other affinity ligands. This allows them to target passively, through the leaky vasculature or actively the main actors of inflammation, including macrophages, endothelial cells, membrane receptors on inflammatory cells, anti-inflammatory genes and cytokines.

Figure 3: Examples of anti-inflammatory diseases and their innovative nanomedicines.

A large spectrum of materials and different strategies are developed for the treatment of inflammatory diseases.

Table 1: Overview of new innovative nanomedicines for pre-clinical therapies against inflammatory disorders.

Lack of information in a table cell will be noted N/A (not available).

Abbreviations: AMPS: 2-Acrylamido-2-MethylPropane-Sulfonic acid, ATP: Adenosine Triphosphate, DOTAP: 1,2-Dioleoyloxy-3-(trimethylammonium)propane, IFN- γ : Interferon gamma, IL-10 (or IL-1Ra): Interleukin 10 (or Interleukin 1 receptor antagonist), LPS:

LipoPolySaccharide, MCP-1: monocyte chemoattractant protein 1, MPO: Myeloperoxidase, NF- κ B: Nuclear Factor-kappa B, NIPAM: N-Isopropyl acrylamide, NPs: Nanoparticles, pDNA: plasmid DeoxyriboNucleic Acid, PEG: Polyethylene Glycol, PEI: Polyethyleneimine, PLGA: Poly(Lactic-co-Glycolic Acid), TNF- α : Tumor Necrosis Factor alpha

Table 2: Some examples of anti-inflammatory nanomedicines under clinical trials

Bibliography

- [1] R. Chovatiya, R. Medzhitov, Stress, inflammation, and defense of homeostasis, *Mol Cell*, 54 (2014) 281-288.
- [2] W.F.E. Hossam M. Ashour, Md. Masudur Rahman , Hatem A. Elshabrawy Insights into the Recent 2019 Novel Coronavirus (SARS-CoV-2) in Light of Past Human Coronavirus Outbreaks, *Pathogens* 9(2020).
- [3] A.A. Meo SA, Al-Khlaiwi T, Meo IM, Halepoto DM, Iqbal M, Usmani AM, Hajjar W, Ahmed N., Novel coronavirus 2019-nCoV: prevalence, biological and clinical characteristics comparison with SARS-CoV and MERS-CoV, *Eur Rev Med Pharmacol Sci*, 24 (2020) 2012-2019.
- [4] T.S. Xiao, Innate immunity and inflammation, *Cell Mol Immunol*, 14 (2017) 1-3.
- [5] G.S. Hotamisligil, Endoplasmic reticulum stress and the inflammatory basis of metabolic disease, *Cell*, 140 (2010) 900-917.
- [6] M. Antonelli, I. Kushner, It's time to redefine inflammation, *FASEB J*, 31 (2017) 1787-1791.
- [7] D.A. Cronkite, T.M. Strutt, The Regulation of Inflammation by Innate and Adaptive Lymphocytes, *J Immunol Res*, 2018 (2018) 1467538.
- [8] R. Medzhitov, Inflammation 2010: new adventures of an old flame, *Cell*, 140 (2010) 771-776.
- [9] D.D. Chaplin, Overview of the immune response, *J Allergy Clin Immunol*, 125 (2010) S3-23.
- [10] D.G. Harrison, T.J. Guzik, H.E. Lob, M.S. Madhur, P.J. Marvar, S.R. Thabet, A. Vinh, C.M. Weyand, Inflammation, immunity, and hypertension, *Hypertension*, 57 (2011) 132-140.
- [11] R. Molinaro, C. Boada, G.M. Del Rosal, K.A. Hartman, C. Corbo, E.D. Andrews, N.E. Toledano-Furman, J.P. Cooke, E. Tasciotti, Vascular Inflammation: A Novel Access Route for Nanomedicine, *Methodist Deakey Cardiovasc J*, 12 (2016) 169-174.
- [12] M. Durymanov, T. Kamaletdinova, S.E. Lehmann, J. Reineke, Exploiting passive nanomedicine accumulation at sites of enhanced vascular permeability for non-cancerous applications, *J Control Release*, 261 (2017) 10-22.
- [13] E.A. Azzopardi, E.L. Ferguson, D.W. Thomas, The enhanced permeability retention effect: a new paradigm for drug targeting in infection, *J Antimicrob Chemother*, 68 (2013) 257-274.
- [14] D. Wang, S.R. Goldring, The bone, the joints and the Balm of Gilead, *Mol Pharm*, 8 (2011) 991-993.
- [15] F. Yuan, L.D. Quan, L. Cui, S.R. Goldring, D. Wang, Development of macromolecular prodrug for rheumatoid arthritis, *Adv Drug Deliv Rev*, 64 (2012) 1205-1219.
- [16] P. Calvo, B. Gouritin, H. Villarroya, F. Eclancher, C. Giannavola, C. Klein, J.P. Andreux, P. Couvreur, Quantification and localization of PEGylated polycyanoacrylate nanoparticles in brain and spinal cord during experimental allergic encephalomyelitis in the rat, *Eur J Neurosci*, 15 (2002) 1317-1326.
- [17] G. Cavaletti, A. Casetti, A. Canta, S. Galbiati, A. Gilardini, N. Oggioni, V. Rodriguez-Menendez, A. Fasano, G.M. Liuzzi, U. Fattler, S. Ries, J. Nieland, P. Riccio, H. Haas, Cationic liposomes target sites of acute neuroinflammation in experimental autoimmune encephalomyelitis, *Mol Pharm*, 6 (2009) 1363-1370.
- [18] I. Tabas, C.K. Glass, Anti-inflammatory therapy in chronic disease: challenges and opportunities, *Science*, 339 (2013) 166-172.

- [19] A. Banerjee, J. Qi, R. Gogoi, J. Wong, S. Mitragotri, Role of nanoparticle size, shape and surface chemistry in oral drug delivery, *J Control Release*, 238 (2016) 176-185.
- [20] A.B. Jindal, The effect of particle shape on cellular interaction and drug delivery applications of micro- and nanoparticles, *Int J Pharm*, 532 (2017) 450-465.
- [21] D.H. Jo, J.H. Kim, T.G. Lee, Size, surface charge, and shape determine therapeutic effects of nanoparticles on brain and retinal diseases, *Nanomedicine*, 11 (2015) 1603-1611.
- [22] V. Agrahari, P.A. Burnouf, T. Burnouf, Nanoformulation properties, characterization, and behavior in complex biological matrices: Challenges and opportunities for brain-targeted drug delivery applications and enhanced translational potential, *Adv Drug Deliv Rev*, 148 (2019) 146-180.
- [23] G.A. Duncan, M.A. Bevan, Computational design of nanoparticle drug delivery systems for selective targeting, *Nanoscale*, 7 (2015) 15332-15340.
- [24] H. Li, X. Zhang, K. Wang, H. Liu, Y. Wei, Facile preparation of biocompatible and robust fluorescent polymeric nanoparticles via PEGylation and cross-linking, *ACS Appl Mater Interfaces*, 7 (2015) 4241-4246.
- [25] M.J. Santos-Martinez, K. Rahme, J.J. Corbalan, C. Faulkner, J.D. Holmes, L. Tajber, C. Medina, M.W. Radomski, Pegylation increases platelet biocompatibility of gold nanoparticles, *J Biomed Nanotechnol*, 10 (2014) 1004-1015.
- [26] J. Iqbal, B.A. Abbasi, R. Ahmad, T. Mahmood, B. Ali, A.T. Khalil, S. Kanwal, S.A. Shah, M.M. Alam, H. Badshah, A. Munir, Nanomedicines for developing cancer nanotherapeutics: from benchtop to bedside and beyond, *Appl Microbiol Biotechnol*, 102 (2018) 9449-9470.
- [27] S. Tran, P.J. DeGiovanni, B. Piel, P. Rai, Cancer nanomedicine: a review of recent success in drug delivery, *Clin Transl Med*, 6 (2017) 44.
- [28] K.S. Kim, G. Khang, D. Lee, Application of nanomedicine in cardiovascular diseases and stroke, *Curr Pharm Des*, 17 (2011) 1825-1833.
- [29] V.M. Martin Gimenez, D.E. Kassuha, W. Manucha, Nanomedicine applied to cardiovascular diseases: latest developments, *Ther Adv Cardiovasc Dis*, 11 (2017) 133-142.
- [30] M.V. F. Dormont, P. Couvreur, Nanoplumbers: biomaterials to fight cardiovascular diseases, *Materials Today*, 21 (2018) 122-143.
- [31] M. Gharagozloo, S. Majewski, M. Foldvari, Therapeutic applications of nanomedicine in autoimmune diseases: from immunosuppression to tolerance induction, *Nanomedicine*, 11 (2015) 1003-1018.
- [32] O. Veisoh, B.C. Tang, K.A. Whitehead, D.G. Anderson, R. Langer, Managing diabetes with nanomedicine: challenges and opportunities, *Nat Rev Drug Discov*, 14 (2015) 45-57.
- [33] M. Goldsmith, L. Abramovitz, D. Peer, Precision nanomedicine in neurodegenerative diseases, *ACS Nano*, 8 (2014) 1958-1965.
- [34] A.V. Kabanov, H.E. Gendelman, Nanomedicine in the diagnosis and therapy of neurodegenerative disorders, *Prog Polym Sci*, 32 (2007) 1054-1082.
- [35] R. Molinaro, C. Corbo, M. Livingston, M. Evangelopoulos, A. Parodi, C. Boada, M. Agostini, E. Tasciotti, Inflammation and Cancer: In Medio Stat Nano, *Curr Med Chem*, 25 (2018) 4208-4223.
- [36] L. Salvioni, M.A. Rizzuto, J.A. Bertolini, L. Pandolfi, M. Colombo, D. Prospero, Thirty Years of Cancer Nanomedicine: Success, Frustration, and Hope, *Cancers (Basel)*, 11 (2019).
- [37] D.J. Irvine, E.L. Dane, Enhancing cancer immunotherapy with nanomedicine, *Nat Rev Immunol*, (2020).

- [38] T.G. Dacoba, A. Olivera, D. Torres, J. Crecente-Campo, M.J. Alonso, Modulating the immune system through nanotechnology, *Semin Immunol*, 34 (2017) 78-102.
- [39] P.J. Barnes, Glucocorticoids, *Chem Immunol Allergy*, 100 (2014) 311-316.
- [40] E. Oppong, A.C. Cato, Effects of Glucocorticoids in the Immune System, *Adv Exp Med Biol*, 872 (2015) 217-233.
- [41] E.-M.C. Hussain Ali, Maïke Windbergs, Claus-Michael Lehr, Nanomedicines for the treatment of inflammatory bowel diseases, *European Journal of Nanomedicine*, 5 (2013) 23-38.
- [42] M. Oray, K. Abu Samra, N. Ebrahimiadib, H. Meese, C.S. Foster, Long-term side effects of glucocorticoids, *Expert Opin Drug Saf*, 15 (2016) 457-465.
- [43] J. Youshia, A. Lamprecht, Size-dependent nanoparticulate drug delivery in inflammatory bowel diseases, *Expert Opin Drug Deliv*, 13 (2016) 281-294.
- [44] S.J. Kshirsagar, M.R. Bhalekar, J.N. Patel, S.K. Mohapatra, N.S. Shewale, Preparation and characterization of nanocapsules for colon-targeted drug delivery system, *Pharm Dev Technol*, 17 (2012) 607-613.
- [45] A. Kumari, A. Jain, P. Hurkat, A. Tiwari, S.K. Jain, Eudragit S100 coated microsponges for Colon targeting of prednisolone, *Drug Dev Ind Pharm*, 44 (2018) 902-913.
- [46] B. Moulari, A. Beduneau, Y. Pellequer, A. Lamprecht, Lectin-decorated nanoparticles enhance binding to the inflamed tissue in experimental colitis, *J Control Release*, 188 (2014) 9-17.
- [47] C.M.A. van Alem, M. Boonstra, J. Prins, T. Bezhaeva, M.F. van Essen, J.M. Ruben, A.L. Vahrmeijer, E.P. van der Veer, J.W. de Fijter, M.E. Reinders, O. Meijer, J.M. Metselaar, C. van Kooten, J.I. Rotmans, Local delivery of liposomal prednisolone leads to an anti-inflammatory profile in renal ischaemia-reperfusion injury in the rat, *Nephrol Dial Transplant*, 33 (2018) 44-53.
- [48] P.H. Zhou, B. Qiu, R.H. Deng, H.J. Li, X.F. Xu, X.F. Shang, Chondroprotective Effects of Hyaluronic Acid-Chitosan Nanoparticles Containing Plasmid DNA Encoding Cytokine Response Modifier A in a Rat Knee Osteoarthritis Model, *Cell Physiol Biochem*, 47 (2018) 1207-1216.
- [49] A. Verma, A. Jain, A. Tiwari, S. Saraf, P.K. Panda, G.P. Agrawal, S.K. Jain, Folate Conjugated Double Liposomes Bearing Prednisolone and Methotrexate for Targeting Rheumatoid Arthritis, *Pharm Res*, 36 (2019) 123.
- [50] P.M. Gaafar, O.Y. Abdallah, R.M. Farid, H. Abdelkader, Preparation, characterization and evaluation of novel elastic nano-sized niosomes (ethoniosomes) for ocular delivery of prednisolone, *J Liposome Res*, 24 (2014) 204-215.
- [51] Y. Karabey-Akyurek, A.G. Gurcay, O. Gurcan, O.F. Turkoglu, S. Yabanoglu-Ciftci, H. Eroglu, M.F. Sargon, E. Bilensoy, L. Oner, Localized delivery of methylprednisolone sodium succinate with polymeric nanoparticles in experimental injured spinal cord model, *Pharm Dev Technol*, 22 (2017) 972-981.
- [52] U. Soiberman, S.P. Kambhampati, T. Wu, M.K. Mishra, Y. Oh, R. Sharma, J. Wang, A.E. Al Towerki, S. Yiu, W.J. Stark, R.M. Kannan, Subconjunctival injectable dendrimer-dexamethasone gel for the treatment of corneal inflammation, *Biomaterials*, 125 (2017) 38-53.
- [53] Q. Wang, J. Jiang, W. Chen, H. Jiang, Z. Zhang, X. Sun, Targeted delivery of low-dose dexamethasone using PCL-PEG micelles for effective treatment of rheumatoid arthritis, *J Control Release*, 230 (2016) 64-72.
- [54] A. Lee, C. De Mei, M. Ferreira, R. Marotta, H.Y. Yoon, K. Kim, I.C. Kwon, P. Decuzzi, Dexamethasone-loaded Polymeric Nanoconstructs for Monitoring and Treating Inflammatory Bowel Disease, *Theranostics*, 7 (2017) 3653-3666.

- [55] C. Dianza, F. Foglietta, B. Ferrara, A.C. Rosa, E. Muntoni, P. Gasco, C. Della Pepa, R. Canaparo, L. Serpe, Solid lipid nanoparticles delivering anti-inflammatory drugs to treat inflammatory bowel disease: Effects in an in vivo model, *World J Gastroenterol*, 23 (2017) 4200-4210.
- [56] Z. Jia, G. Zhao, X. Wei, D. Kong, Y. Sun, Y. Zhou, S.M. Lele, E.V. Fehringer, K.L. Garvin, S.R. Goldring, D. Wang, Structural optimization of HPMA copolymer-based dexamethasone prodrug for improved treatment of inflammatory arthritis, *J Control Release*, (2020).
- [57] G.R. Greenberg, B.G. Feagan, F. Martin, L.R. Sutherland, A.B. Thomson, C.N. Williams, L.G. Nilsson, T. Persson, Oral budesonide for active Crohn's disease. Canadian Inflammatory Bowel Disease Study Group, *N Engl J Med*, 331 (1994) 836-841.
- [58] M. Naeem, M. Choi, J. Cao, Y. Lee, M. Ikram, S. Yoon, J. Lee, H.R. Moon, M.S. Kim, Y. Jung, J.W. Yoo, Colon-targeted delivery of budesonide using dual pH- and time-dependent polymeric nanoparticles for colitis therapy, *Drug Des Devel Ther*, 9 (2015) 3789-3799.
- [59] S. Hua, E. Marks, J.J. Schneider, S. Keely, Advances in oral nano-delivery systems for colon targeted drug delivery in inflammatory bowel disease: selective targeting to diseased versus healthy tissue, *Nanomedicine*, 11 (2015) 1117-1132.
- [60] H. Ali, B. Weigmann, E.M. Collnot, S.A. Khan, M. Windbergs, C.M. Lehr, Budesonide Loaded PLGA Nanoparticles for Targeting the Inflamed Intestinal Mucosa--Pharmaceutical Characterization and Fluorescence Imaging, *Pharm Res*, 33 (2016) 1085-1092.
- [61] S.Y. Vafaei, M. Esmaeili, M. Amini, F. Atyabi, S.N. Ostad, R. Dinarvand, Self assembled hyaluronic acid nanoparticles as a potential carrier for targeting the inflamed intestinal mucosa, *Carbohydr Polym*, 144 (2016) 371-381.
- [62] S. Wongrakpanich, A. Wongrakpanich, K. Melhado, J. Rangaswami, A Comprehensive Review of Non-Steroidal Anti-Inflammatory Drug Use in The Elderly, *Aging Dis*, 9 (2018) 143-150.
- [63] C. Sostres, C.J. Gargallo, M.T. Arroyo, A. Lanas, Adverse effects of non-steroidal anti-inflammatory drugs (NSAIDs, aspirin and coxibs) on upper gastrointestinal tract, *Best Pract Res Clin Gastroenterol*, 24 (2010) 121-132.
- [64] R.M. Goncalves, A.C. Pereira, I.O. Pereira, M.J. Oliveira, M.A. Barbosa, Macrophage response to chitosan/poly-(gamma-glutamic acid) nanoparticles carrying an anti-inflammatory drug, *J Mater Sci Mater Med*, 26 (2015) 167.
- [65] Q. Shang, S. Huang, A. Zhang, J. Feng, S. Yang, The binary complex of poly(PEGMA-co-MAA) hydrogel and PLGA nanoparticles as a novel oral drug delivery system for ibuprofen delivery, *J Biomater Sci Polym Ed*, 28 (2017) 1874-1887.
- [66] P. Nirbhavane, G. Sharma, B. Singh, G.K. Khuller, V.G. Goni, A.B. Patil, O.P. Katare, Preclinical Explorative Assessment of Celecoxib-Based Biocompatible Lipidic Nanocarriers for the Management of CFA-Induced Rheumatoid Arthritis in Wistar Rats, *AAPS PharmSciTech*, 19 (2018) 3187-3198.
- [67] W.K.E. Ip, N. Hoshi, D.S. Shouval, S. Snapper, R. Medzhitov, Anti-inflammatory effect of IL-10 mediated by metabolic reprogramming of macrophages, *Science*, 356 (2017) 513-519.
- [68] A. Viscido, A. Capannolo, G. Latella, R. Caprilli, G. Frieri, Nanotechnology in the treatment of inflammatory bowel diseases, *J Crohns Colitis*, 8 (2014) 903-918.
- [69] R. Toita, T. Kawano, M. Murata, J.H. Kang, Anti-obesity and anti-inflammatory effects of macrophage-targeted interleukin-10-conjugated liposomes in obese mice, *Biomaterials*, 110 (2016) 81-88.

- [70] N. Kamaly, G. Fredman, J.J. Fojas, M. Subramanian, W.I. Choi, K. Zepeda, C. Vilos, M. Yu, S. Gadde, J. Wu, J. Milton, R. Carvalho Leitao, L. Rosa Fernandes, M. Hasan, H. Gao, V. Nguyen, J. Harris, I. Tabas, O.C. Farokhzad, Targeted Interleukin-10 Nanotherapeutics Developed with a Microfluidic Chip Enhance Resolution of Inflammation in Advanced Atherosclerosis, *ACS Nano*, 10 (2016) 5280-5292.
- [71] D.R. Baganizi, E. Nyairo, S.A. Duncan, S.R. Singh, V.A. Dennis, Interleukin-10 Conjugation to Carboxylated PVP-Coated Silver Nanoparticles for Improved Stability and Therapeutic Efficacy, *Nanomaterials (Basel)*, 7 (2017).
- [72] M.C. Stensberg, Q. Wei, E.S. McLamore, D.M. Porterfield, A. Wei, M.S. Sepulveda, Toxicological studies on silver nanoparticles: challenges and opportunities in assessment, monitoring and imaging, *Nanomedicine (Lond)*, 6 (2011) 879-898.
- [73] L. Yildirimer, N.T. Thanh, M. Loizidou, A.M. Seifalian, Toxicology and clinical potential of nanoparticles, *Nano Today*, 6 (2011) 585-607.
- [74] J.H. Sung, J.H. Ji, J.U. Yoon, D.S. Kim, M.Y. Song, J. Jeong, B.S. Han, J.H. Han, Y.H. Chung, J. Kim, T.S. Kim, H.K. Chang, E.J. Lee, J.H. Lee, I.J. Yu, Lung function changes in Sprague-Dawley rats after prolonged inhalation exposure to silver nanoparticles, *Inhal Toxicol*, 20 (2008) 567-574.
- [75] K. Bilberg, M.B. Hovgaard, F. Besenbacher, E. Baatrup, In Vivo Toxicity of Silver Nanoparticles and Silver Ions in Zebrafish (*Danio rerio*), *J Toxicol*, 2012 (2012) 293784.
- [76] R. Agarwal, T.M. Volkmer, P. Wang, L.A. Lee, Q. Wang, A.J. Garcia, Synthesis of self-assembled IL-1Ra-presenting nanoparticles for the treatment of osteoarthritis, *J Biomed Mater Res A*, 104 (2016) 595-599.
- [77] T. Wang, Y. Tang, X. He, J. Yan, C. Wang, X. Feng, Self-Assembled Raspberry-Like Core/Satellite Nanoparticles for Anti-Inflammatory Protein Delivery, *ACS Appl Mater Interfaces*, 9 (2017) 6902-6907.
- [78] S. Gupta, A.K. Sharma, V. Shastri, M.K. Madhu, V.K. Sharma, Prediction of anti-inflammatory proteins/peptides: an insilico approach, *J Transl Med*, 15 (2017) 7.
- [79] R.L. Bartlett, 2nd, S. Sharma, A. Panitch, Cell-penetrating peptides released from thermosensitive nanoparticles suppress pro-inflammatory cytokine response by specifically targeting inflamed cartilage explants, *Nanomedicine*, 9 (2013) 419-427.
- [80] S. Poh, J.B. Lin, A. Panitch, Release of anti-inflammatory peptides from thermosensitive nanoparticles with degradable cross-links suppresses pro-inflammatory cytokine production, *Biomacromolecules*, 16 (2015) 1191-1200.
- [81] J.B. Lin, S. Poh, A. Panitch, Controlled release of anti-inflammatory peptides from reducible thermosensitive nanoparticles suppresses cartilage inflammation, *Nanomedicine*, 12 (2016) 2095-2100.
- [82] C. He, L. Yin, Y. Song, C. Tang, C. Yin, Optimization of multifunctional chitosan-siRNA nanoparticles for oral delivery applications, targeting TNF-alpha silencing in rats, *Acta Biomater*, 17 (2015) 98-106.
- [83] S. Yadav, S.K. Gandham, R. Panicucci, M.M. Amiji, Intranasal brain delivery of cationic nanoemulsion-encapsulated TNFalpha siRNA in prevention of experimental neuroinflammation, *Nanomedicine*, 12 (2016) 987-1002.
- [84] T. Kanazawa, T. Kurano, H. Ibaraki, Y. Takashima, T. Suzuki, Y. Seta, Therapeutic Effects in a Transient Middle Cerebral Artery Occlusion Rat Model by Nose-To-Brain Delivery of Anti-TNF-Alpha siRNA with Cell-Penetrating Peptide-Modified Polymer Micelles, *Pharmaceutics*, 11 (2019).
- [85] A. Bohr, N. Tsapis, I. Andreana, A. Chamarat, C. Foged, C. Delomenie, M. Noiray, N. El Brahmi, J.P. Majoral, S. Mignani, E. Fattal, Anti-Inflammatory Effect of Anti-TNF-alpha SiRNA Cationic Phosphorus Dendrimer Nanocomplexes Administered Intranasally in a Murine Acute Lung Injury Model, *Biomacromolecules*, 18 (2017) 2379-2388.

- [86] Q. Shi, E.P. Rondon-Cavanzo, I.P. Dalla Picola, M.J. Tiera, X. Zhang, K. Dai, H.A. Benabdoune, M. Benderdour, J.C. Fernandes, In vivo therapeutic efficacy of TNFalpha silencing by folate-PEG-chitosan-DEAE/siRNA nanoparticles in arthritic mice, *Int J Nanomedicine*, 13 (2018) 387-402.
- [87] A.M. Aldayel, H.L. O'Mary, S.A. Valdes, X. Li, S.G. Thakkar, B.E. Mustafa, Z. Cui, Lipid nanoparticles with minimum burst release of TNF-alpha siRNA show strong activity against rheumatoid arthritis unresponsive to methotrexate, *J Control Release*, 283 (2018) 280-289.
- [88] M.A.A. Jansen, L.H. Klausen, K. Thanki, J. Lyngso, J. Skov Pedersen, H. Franzyk, H.M. Nielsen, W. van Eden, M. Dong, F. Broere, C. Foged, X. Zeng, Lipidoid-polymer hybrid nanoparticles loaded with TNF siRNA suppress inflammation after intra-articular administration in a murine experimental arthritis model, *Eur J Pharm Biopharm*, 142 (2019) 38-48.
- [89] Y.C. Liu, X.B. Zou, Y.F. Chai, Y.M. Yao, Macrophage polarization in inflammatory diseases, *Int J Biol Sci*, 10 (2014) 520-529.
- [90] S. Jain, T.H. Tran, M. Amiji, Macrophage repolarization with targeted alginate nanoparticles containing IL-10 plasmid DNA for the treatment of experimental arthritis, *Biomaterials*, 61 (2015) 162-177.
- [91] T.H. Tran, R. Rastogi, J. Shelke, M.M. Amiji, Modulation of Macrophage Functional Polarity towards Anti-Inflammatory Phenotype with Plasmid DNA Delivery in CD44 Targeting Hyaluronic Acid Nanoparticles, *Sci Rep*, 5 (2015) 16632.
- [92] M. Haneklaus, M. Gerlic, L.A. O'Neill, S.L. Masters, miR-223: infection, inflammation and cancer, *J Intern Med*, 274 (2013) 215-226.
- [93] T.H. Tran, S. Krishnan, M.M. Amiji, MicroRNA-223 Induced Repolarization of Peritoneal Macrophages Using CD44 Targeting Hyaluronic Acid Nanoparticles for Anti-Inflammatory Effects, *PLoS One*, 11 (2016) e0152024.
- [94] A.M. Carmona-Ribeiro, Preparation and characterization of biomimetic nanoparticles for drug delivery, *Methods Mol Biol*, 906 (2012) 283-294.
- [95] C.M. Hu, R.H. Fang, K.C. Wang, B.T. Luk, S. Thamphiwatana, D. Dehaini, P. Nguyen, P. Angsantikul, C.H. Wen, A.V. Kroll, C. Carpenter, M. Ramesh, V. Qu, S.H. Patel, J. Zhu, W. Shi, F.M. Hofman, T.C. Chen, W. Gao, K. Zhang, S. Chien, L. Zhang, Nanoparticle biointerfacing by platelet membrane cloaking, *Nature*, 526 (2015) 118-121.
- [96] H. Yan, D. Shao, Y.H. Lao, M. Li, H. Hu, K.W. Leong, Engineering Cell Membrane-Based Nanotherapeutics to Target Inflammation, *Adv Sci (Weinh)*, 6 (2019) 1900605.
- [97] K. Jin, Z. Luo, B. Zhang, Z. Pang, Biomimetic nanoparticles for inflammation targeting, *Acta Pharm Sin B*, 8 (2018) 23-33.
- [98] V. Vijayan, S. Uthaman, I.K. Park, Cell Membrane-Camouflaged Nanoparticles: A Promising Biomimetic Strategy for Cancer Theragnostics, *Polymers (Basel)*, 10 (2018).
- [99] H. Wang, Y. Liu, R. He, D. Xu, J. Zang, N. Weeranoppanant, H. Dong, Y. Li, Cell membrane biomimetic nanoparticles for inflammation and cancer targeting in drug delivery, *Biomater Sci*, 8 (2020) 552-568.
- [100] G.S. Firestein, I.B. McInnes, Immunopathogenesis of Rheumatoid Arthritis, *Immunity*, 46 (2017) 183-196.
- [101] M. Horckmans, L. Ring, J. Duchene, D. Santovito, M.J. Schloss, M. Drechsler, C. Weber, O. Soehnlein, S. Steffens, Neutrophils orchestrate post-myocardial infarction healing by polarizing macrophages towards a reparative phenotype, *Eur Heart J*, 38 (2017) 187-197.
- [102] Q. Zhang, D. Dehaini, Y. Zhang, J. Zhou, X. Chen, L. Zhang, R.H. Fang, W. Gao, Neutrophil membrane-coated nanoparticles inhibit synovial inflammation and alleviate joint damage in inflammatory arthritis, *Nat Nanotechnol*, 13 (2018) 1182-1190.

- [103] S. Thamphiwatana, P. Angsantikul, T. Escajadillo, Q. Zhang, J. Olson, B.T. Luk, S. Zhang, R.H. Fang, W. Gao, V. Nizet, L. Zhang, Macrophage-like nanoparticles concurrently absorbing endotoxins and proinflammatory cytokines for sepsis management, *Proc Natl Acad Sci U S A*, 114 (2017) 11488-11493.
- [104] C. Boada, A. Zinger, C. Tsao, P. Zhao, J.O. Martinez, K. Hartman, T. Naoi, R. Sukhoveshin, M. Sushnitha, R. Molinaro, B. Trachtenberg, J.P. Cooke, E. Tasciotti, Rapamycin-Loaded Biomimetic Nanoparticles Reverse Vascular Inflammation, *Circ Res*, 126 (2020) 25-37.
- [105] T. Fukuta, S. Yoshimi, T. Tanaka, K. Kogure, Leukocyte-mimetic liposomes possessing leukocyte membrane proteins pass through inflamed endothelial cell layer by regulating intercellular junctions, *Int J Pharm*, 563 (2019) 314-323.
- [106] R. Li, Y. He, Y. Zhu, L. Jiang, S. Zhang, J. Qin, Q. Wu, W. Dai, S. Shen, Z. Pang, J. Wang, Route to Rheumatoid Arthritis by Macrophage-Derived Microvesicle-Coated Nanoparticles, *Nano Lett*, 19 (2019) 124-134.
- [107] N. Zelcer, P. Tontonoz, Liver X receptors as integrators of metabolic and inflammatory signaling, *J Clin Invest*, 116 (2006) 607-614.
- [108] H. He, Q. Yuan, J. Bie, R.L. Wallace, P.J. Yannie, J. Wang, M.G. Lancina, 3rd, O.Y. Zolotarskaya, W. Korzun, H. Yang, S. Ghosh, Development of mannose functionalized dendrimeric nanoparticles for targeted delivery to macrophages: use of this platform to modulate atherosclerosis, *Transl Res*, 193 (2018) 13-30.
- [109] X.Q. Zhang, O. Even-Or, X. Xu, M. van Rosmalen, L. Lim, S. Gadde, O.C. Farokhzad, E.A. Fisher, Nanoparticles containing a liver X receptor agonist inhibit inflammation and atherosclerosis, *Adv Healthc Mater*, 4 (2015) 228-236.
- [110] Y. Guo, W. Yuan, B. Yu, R. Kuai, W. Hu, E.E. Morin, M.T. Garcia-Barrio, J. Zhang, J.J. Moon, A. Schwendeman, Y. Eugene Chen, Synthetic High-Density Lipoprotein-Mediated Targeted Delivery of Liver X Receptors Agonist Promotes Atherosclerosis Regression, *EBioMedicine*, 28 (2018) 225-233.
- [111] R.K. Thapa, B.K. Yoo, Evaluation of the effect of tacrolimus-loaded liquid crystalline nanoparticles on psoriasis-like skin inflammation, *J Dermatolog Treat*, 25 (2014) 22-25.
- [112] C.G. Sen Lin, Doudou Wang, Qi Xie, Biao Wu, Jingjie Wang, Kaihui Nan*, Qinxiang Zheng*, and Wei Chen*, Overcoming the Anatomical and Physiological Barriers in Topical Eye Surface Medication Using a Peptide-Decorated Polymeric Micelle, *ACS Appl. Mater. Interfaces*, 11 (2019).
- [113] K. Okuda, H.Y. Fu, T. Matsuzaki, R. Araki, S. Tsuchida, P.V. Thanikachalam, T. Fukuta, T. Asai, M. Yamato, S. Sanada, H. Asanuma, Y. Asano, M. Asakura, H. Hanawa, H. Hao, N. Oku, S. Takashima, M. Kitakaze, Y. Sakata, T. Minamino, Targeted Therapy for Acute Autoimmune Myocarditis with Nano-Sized Liposomal FK506 in Rats, *PLoS One*, 11 (2016) e0160944.
- [114] A. Partoazar, S. Nasoohi, S.M. Rezayat, K. Gilani, S.E. Mehr, A. Amani, N. Rahimi, A.R. Dehpour, Nanoliposome containing cyclosporine A reduced neuroinflammation responses and improved neurological activities in cerebral ischemia/reperfusion in rat, *Fundam Clin Pharmacol*, 31 (2017) 185-193.
- [115] S. Yadav, G. Pawar, P. Kulkarni, C. Ferris, M. Amiji, CNS Delivery and Anti-Inflammatory Effects of Intranasally Administered Cyclosporine-A in Cationic Nanoformulations, *J Pharmacol Exp Ther*, 370 (2019) 843-854.
- [116] H. Tang, D. Xiang, F. Wang, J. Mao, X. Tan, Y. Wang, 5-ASA-loaded SiO₂ nanoparticles-a novel drug delivery system targeting therapy on ulcerative colitis in mice, *Mol Med Rep*, 15 (2017) 1117-1122.

- [117] P. Tang, Q. Sun, L. Zhao, H. Pu, H. Yang, S. Zhang, R. Gan, N. Gan, H. Li, Mesalazine/hydroxypropyl-beta-cyclodextrin/chitosan nanoparticles with sustained release and enhanced anti-inflammation activity, *Carbohydr Polym*, 198 (2018) 418-425.
- [118] T. Yazeji, B. Moulari, A. Beduneau, V. Stein, D. Dietrich, Y. Pellequer, A. Lamprecht, Nanoparticle-based delivery enhances anti-inflammatory effect of low molecular weight heparin in experimental ulcerative colitis, *Drug Deliv*, 24 (2017) 811-817.
- [119] P. Couvreur, B. Stella, L.H. Reddy, H. Hillaireau, C. Dubernet, D. Desmaele, S. Lepetre-Mouelhi, F. Rocco, N. Dereuddre-Bosquet, P. Clayette, V. Rosilio, V. Marsaud, J.M. Renoir, L. Cattel, Squalenoyl nanomedicines as potential therapeutics, *Nano Lett*, 6 (2006) 2544-2548.
- [120] D. Desmaele, R. Gref, P. Couvreur, Squalenoylation: a generic platform for nanoparticulate drug delivery, *J Control Release*, 161 (2012) 609-618.
- [121] J. Feng, S. Lepetre-Mouelhi, A. Gautier, S. Mura, C. Cailleau, F. Coudore, M. Hamon, P. Couvreur, A new painkiller nanomedicine to bypass the blood-brain barrier and the use of morphine, *Sci Adv*, 5 (2019) eaau5148.
- [122] L. Antonioli, C. Blandizzi, P. Pacher, G. Hasko, Immunity, inflammation and cancer: a leading role for adenosine, *Nat Rev Cancer*, 13 (2013) 842-857.
- [123] G. Hasko, J. Linden, B. Cronstein, P. Pacher, Adenosine receptors: therapeutic aspects for inflammatory and immune diseases, *Nat Rev Drug Discov*, 7 (2008) 759-770.
- [124] A. Gaudin, M. Yemisci, H. Eroglu, S. Lepetre-Mouelhi, O.F. Turkoglu, B. Donmez-Demir, S. Caban, M.F. Sargon, S. Garcia-Argote, G. Pieters, O. Loreau, B. Rousseau, O. Tagit, N. Hildebrandt, Y. Le Dantec, J. Mougin, S. Valetti, H. Chacun, V. Nicolas, D. Desmaele, K. Andrieux, Y. Capan, T. Dalkara, P. Couvreur, Squalenoyl adenosine nanoparticles provide neuroprotection after stroke and spinal cord injury, *Nat Nanotechnol*, 9 (2014) 1054-1062.
- [125] J. Egea, I. Fabregat, Y.M. Frapart, P. Ghezzi, A. Gorchach, T. Kietzmann, K. Kubaichuk, U.G. Knaus, M.G. Lopez, G. Olaso-Gonzalez, A. Petry, R. Schulz, J. Vina, P. Winyard, K. Abbas, O.S. Ademowo, C.B. Afonso, I. Andreadou, H. Antelmann, F. Antunes, M. Aslan, M.M. Bachschmid, R.M. Barbosa, V. Belousov, C. Berndt, D. Bernlohr, E. Bertran, A. Bindoli, S.P. Bottari, P.M. Brito, G. Carrara, A.I. Casas, A. Chatzi, N. Chondrogianni, M. Conrad, M.S. Cooke, J.G. Costa, A. Cuadrado, P. My-Chan Dang, B. De Smet, B. Debelec-Butuner, I.H.K. Dias, J.D. Dunn, A.J. Edson, M. El Assar, J. El-Benna, P. Ferdinandy, A.S. Fernandes, K.E. Fladmark, U. Forstermann, R. Giniatullin, Z. Giricz, A. Gorbe, H. Griffiths, V. Hampl, A. Hanf, J. Herget, P. Hernansanz-Agustin, M. Hillion, J. Huang, S. Ilikay, P. Jansen-Durr, V. Jaquet, J.A. Joles, B. Kalyanaraman, D. Kaminsky, M. Karbaschi, M. Kleanthous, L.O. Klotz, B. Korac, K.S. Korkmaz, R. Koziel, D. Kracun, K.H. Krause, V. Kren, T. Krieg, J. Laranjinha, A. Lazou, H. Li, A. Martinez-Ruiz, R. Matsui, G.J. McBean, S.P. Meredith, J. Messens, V. Miguel, Y. Mikhed, I. Milisav, L. Milkovic, A. Miranda-Vizuet, M. Mojovic, M. Monsalve, P.A. Mouthuy, J. Mulvey, T. Munzel, V. Muzykantov, I.T.N. Nguyen, M. Oelze, N.G. Oliveira, C.M. Palmeira, N. Papaevgeniou, A. Pavicevic, B. Pedre, F. Peyrot, M. Phylactides, G.G. Pircalabioru, A.R. Pitt, H.E. Poulsen, I. Prieto, M.P. Rigobello, N. Robledinos-Anton, L. Rodriguez-Manas, A.P. Rolo, F. Rousset, T. Ruskovska, N. Saraiva, S. Sasson, K. Schroder, K. Semen, T. Seredenina, A. Shakirzyanova, G.L. Smith, T. Soldati, B.C. Sousa, C.M. Spickett, A. Stancic, M.J. Stasia, H. Steinbrenner, V. Stepanic, S. Steven, K. Tokatlidis, E. Tuncay, B. Turan, F. Ursini, J. Vacek, O. Vajnerova, K. Valentova, F. Van Breusegem, L. Varisli, E.A. Veal, A.S. Yalcin, O. Yelisyeveva, N. Zarkovic, M. Zatloukalova, J. Zielonka, R.M. Touyz, A. Papapetropoulos, T. Grune, S. Lamas, H. Schmidt, F. Di Lisa, A. Daiber, *European*

- contribution to the study of ROS: A summary of the findings and prospects for the future from the COST action BM1203 (EU-ROS), *Redox Biol*, 13 (2017) 94-162.
- [126] V.M. Moia, F. Leal Portilho, T. Almeida Padua, L. Barbosa Correa, E. Ricci-Junior, E. Cruz Rosas, L. Magalhaes Rebelo Alencar, F. Savio Mendes Sinfronio, A. Sampson, S. Hussain Iram, F. Alexis, M.D. de OliveiraHenriques, R. Santos-Oliveira, Lycopene used as Anti-inflammatory Nanodrug for the Treatment of Rheumatoid Arthritis: Animal assay, Pharmacokinetics, ABC Transporter and Tissue Deposition, *Colloids Surf B Biointerfaces*, 188 (2020) 110814.
- [127] M.O. Kadry, Liposomal glutathione as a promising candidate for immunological rheumatoid arthritis therapy, *Heliyon*, 5 (2019) e02162.
- [128] T. Wu, X. Chen, Y. Wang, H. Xiao, Y. Peng, L. Lin, W. Xia, M. Long, J. Tao, X. Shuai, Aortic plaque-targeted andrographolide delivery with oxidation-sensitive micelle effectively treats atherosclerosis via simultaneous ROS capture and anti-inflammation, *Nanomedicine*, 14 (2018) 2215-2226.
- [129] R.B. Flavio Dormont, Catherine Cailleau, Franceline Reynaud, Arnaud Peramo, Amandine Gendron, Julie Mouglin, Françoise Gaudin, Mariana Varna & Patrick Couvreur, Squalene-based multidrug nanoparticles for improved mitigation of uncontrolled inflammation, *Science Advances*, (2020).
- [130] M. Varna, H.V. Xuan, E. Fort, Gold nanoparticles in cardiovascular imaging, *Wiley Interdiscip Rev Nanomed Nanobiotechnol*, 10 (2018).
- [131] M.A. Khan, M.J. Khan, Nano-gold displayed anti-inflammatory property via NF- κ B pathways by suppressing COX-2 activity, *Artif Cells Nanomed Biotechnol*, 46 (2018) 1149-1158.
- [132] Y. Zhao, Z. He, R. Wang, P. Cai, X. Zhang, Q. Yuan, J. Zhang, F. Gao, X. Gao, Comparison of the Therapeutic Effects of Gold Nanoclusters and Gold Nanoparticles on Rheumatoid Arthritis, *J Biomed Nanotechnol*, 15 (2019) 2281-2290.
- [133] F. Gao, Q. Yuan, P. Cai, L. Gao, L. Zhao, M. Liu, Y. Yao, Z. Chai, X. Gao, Au Clusters Treat Rheumatoid Arthritis with Uniquely Reversing Cartilage/Bone Destruction, *Adv Sci (Weinh)*, 6 (2019) 1801671.
- [134] R.P. Pannerec-Varna M, Bousquet G, Ferreira I, Leboeuf C, Boisgard R, Gapihan G, Verine J, Palpant B, Bossy E, et al., In vivo uptake and cellular distribution of gold nanoshells in a preclinical model of xenografted human renal cancer, *Gold Bull* 46 (2013) 257-265.
- [135] S. De Koker, J. Cui, N. Vanparijs, L. Albertazzi, J. Grooten, F. Caruso, B.G. De Geest, Engineering Polymer Hydrogel Nanoparticles for Lymph Node-Targeted Delivery, *Angew Chem Int Ed Engl*, 55 (2016) 1334-1339.
- [136] S.N. Mueller, S. Tian, J.M. DeSimone, Rapid and Persistent Delivery of Antigen by Lymph Node Targeting PRINT Nanoparticle Vaccine Carrier To Promote Humoral Immunity, *Mol Pharm*, 12 (2015) 1356-1365.
- [137] T.W. Chu, J. Yang, R. Zhang, M. Sima, J. Kopecek, Cell surface self-assembly of hybrid nanoconjugates via oligonucleotide hybridization induces apoptosis, *ACS Nano*, 8 (2014) 719-730.
- [138] T.W. Chu, R. Zhang, J. Yang, M.P. Chao, P.J. Shami, J. Kopecek, A Two-Step Pretargeted Nanotherapy for CD20 Crosslinking May Achieve Superior Anti-Lymphoma Efficacy to Rituximab, *Theranostics*, 5 (2015) 834-846.
- [139] L. Li, J. Yang, J. Wang, J. Kopecek, Amplification of CD20 Cross-Linking in Rituximab-Resistant B-Lymphoma Cells Enhances Apoptosis Induction by Drug-Free Macromolecular Therapeutics, *ACS Nano*, 12 (2018) 3658-3670.
- [140] J. Wang, L. Li, J. Yang, P.M. Clair, M.J. Glenn, D.M. Stephens, D.C. Radford, K.M. Kosak, M.W. Deininger, P.J. Shami, J. Kopecek, Drug-free macromolecular therapeutics

- induce apoptosis in cells isolated from patients with B cell malignancies with enhanced apoptosis induction by pretreatment with gemcitabine, *Nanomedicine*, 16 (2019) 217-225.
- [141] J. Wang, Y. Li, L. Li, J. Yang, J. Kopecek, Exploration and Evaluation of Therapeutic Efficacy of Drug-Free Macromolecular Therapeutics in Collagen-Induced Rheumatoid Arthritis Mouse Model, *Macromol Biosci*, 20 (2020) e1900445.
- [142] J. Du, Y.S. Zhang, D. Hobson, P. Hydring, Nanoparticles for immune system targeting, *Drug Discov Today*, 22 (2017) 1295-1301.
- [143] M.S. Goldberg, Improving cancer immunotherapy through nanotechnology, *Nat Rev Cancer*, 19 (2019) 587-602.
- [144] E. Zupancic, C. Curato, M. Paisana, C. Rodrigues, Z. Porat, A.S. Viana, C.A.M. Afonso, J. Pinto, R. Gaspar, J.N. Moreira, R. Satchi-Fainaro, S. Jung, H.F. Florindo, Rational design of nanoparticles towards targeting antigen-presenting cells and improved T cell priming, *J Control Release*, 258 (2017) 182-195.
- [145] W. Gao, R.H. Fang, S. Thamphiwatana, B.T. Luk, J. Li, P. Angsantikul, Q. Zhang, C.M. Hu, L. Zhang, Modulating antibacterial immunity via bacterial membrane-coated nanoparticles, *Nano Lett*, 15 (2015) 1403-1409.
- [146] A. Yeste, M.C. Takenaka, I.D. Mascanfroni, M. Nadeau, J.E. Kenison, B. Patel, A.M. Tukpah, J.A. Babon, M. DeNicola, S.C. Kent, D. Pozo, F.J. Quintana, Tolerogenic nanoparticles inhibit T cell-mediated autoimmunity through SOCS2, *Sci Signal*, 9 (2016) ra61.
- [147] A.C. Anselmo, S. Mitragotri, Nanoparticles in the clinic, *Bioeng Transl Med*, 1 (2016) 10-29.
- [148] A.C. Anselmo, S. Mitragotri, Nanoparticles in the clinic: An update, *Bioeng Transl Med*, 4 (2019) e10143.
- [149] M.E. Lobatto, Z.A. Fayad, S. Silvera, E. Vucic, C. Calcagno, V. Mani, S.D. Dickson, K. Nicolay, M. Banciu, R.M. Schiffelers, J.M. Metselaar, L. van Bloois, H.S. Wu, J.T. Fallon, J.H. Rudd, V. Fuster, E.A. Fisher, G. Storm, W.J. Mulder, Multimodal clinical imaging to longitudinally assess a nanomedical anti-inflammatory treatment in experimental atherosclerosis, *Mol Pharm*, 7 (2010) 2020-2029.
- [150] F.M. van der Valk, D.F. van Wijk, M.E. Lobatto, H.J. Verberne, G. Storm, M.C. Willems, D.A. Legemate, A.J. Nederveen, C. Calcagno, V. Mani, S. Ramachandran, M.P. Paridaans, M.J. Otten, G.M. Dallinga-Thie, Z.A. Fayad, M. Nieuwdorp, D.M. Schulte, J.M. Metselaar, W.J. Mulder, E.S. Stroes, Prednisolone-containing liposomes accumulate in human atherosclerotic macrophages upon intravenous administration, *Nanomedicine*, 11 (2015) 1039-1046.
- [151] A Proof of Concept Study to Determine the Local Delivery and Efficacy of Nanocort - Full Text View - ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT01647685>.
- [152] Silencing Inflammatory Activity by Injecting Nanocort in Patients at Risk for Atherosclerotic Disease - Full Text View - ClinicalTrials.gov, <https://clinicaltrials.gov/ct2/show/NCT01601106>.
- [153] B.M. Voorzaat, J. van Schaik, K.E. van der Bogt, L. Vogt, L. Huisman, B.A. Gabreels, I.M. van der Meer, R.G. van Eps, D. Eefting, M.C. Weijmer, H.O. Groeneveld, R.C. van Nieuwenhuizen, H. Boom, C.A. Verburgh, K. van der Putten, J.I. Rotmans, Improvement of radiocephalic fistula maturation: rationale and design of the Liposomal Prednisolone to Improve Hemodialysis Fistula Maturation (LIPMAT) study - a randomized controlled trial, *J Vasc Access*, 18 (2017) 114-117.
- [154] n.d. Liposome encapsulated prednisolone - AdisInsight, <https://adisinsight.springer.com/drugs/800028576>.

- [155] n.d.h.e.n. Enceladus Pharmaceuticals BV announces positive results from a phase 2a study in patients with active ulcerative colitis with Nanocort® | Enceladus Pharmaceuticals.
- [156] R. Kumar, S. Dogra, B. Amarji, B. Singh, S. Kumar, Sharma, K. Vinay, R. Mahajan, O.P. Katare, Efficacy of Novel Topical Liposomal Formulation of Cyclosporine in Mild to Moderate Stable Plaque Psoriasis: A Randomized Clinical Trial, *JAMA Dermatol*, 152 (2016) 807-815.
- [157] n.d.h. A Evaluation of the Safety of Oncocort IV Pegylated Liposomal Dexamethasone Phosphate in Patients With Progressive Multiple Myeloma - Full Text View - ClinicalTrials.gov.
- [158] K. Nakano, T. Matoba, J.I. Koga, Y. Kashihara, M. Fukae, I. Ieiri, M. Shiramoto, S. Irie, J. Kishimoto, K. Todaka, K. Egashira, Safety, Tolerability, and Pharmacokinetics of NK-104-NP, *Int Heart J*, 59 (2018) 1015-1025.
- [159] M. Sabouri, A. Samadi, S. Ahmad Nasrollahi, E.S. Farboud, B. Mirrahimi, H. Hassanzadeh, M. Nassiri Kashani, R. Dinarvand, A. Firooz, Tretinoin Loaded Nanoemulsion for Acne Vulgaris: Fabrication, Physicochemical and Clinical Efficacy Assessments, *Skin Pharmacol Physiol*, 31 (2018) 316-323.
- [160] S.A.P.R.o.T.I.C.S.i.A. S. Inc, (n.d.). <https://www.prnewswire.com/news-releases/sebacia-announces-positive-results-of-two-independent-clinical-studies-in-acne-300026759.html>.
- [161] S. Hua, M.B.C. de Matos, J.M. Metselaar, G. Storm, Current Trends and Challenges in the Clinical Translation of Nanoparticulate Nanomedicines: Pathways for Translational Development and Commercialization, *Front Pharmacol*, 9 (2018) 790.
- [162] C.A. Dinarello, Anti-inflammatory Agents: Present and Future, *Cell*, 140 (2010) 935-950.
- [163] N. Doshi, S. Mitragotri, Macrophages recognize size and shape of their targets, *PLoS One*, 5 (2010) e10051.
- [164] P.P. Wibroe, A.C. Anselmo, P.H. Nilsson, A. Sarode, V. Gupta, R. Urbanics, J. Szebeni, A.C. Hunter, S. Mitragotri, T.E. Mollnes, S.M. Moghimi, Bypassing adverse injection reactions to nanoparticles through shape modification and attachment to erythrocytes, *Nat Nanotechnol*, 12 (2017) 589-594.
- [165] H.S. Leong, K.S. Butler, C.J. Brinker, M. Azzawi, S. Conlan, C. Dufes, A. Owen, S. Rannard, C. Scott, C. Chen, M.A. Dobrovolskaia, S.V. Kozlov, A. Prina-Mello, R. Schmid, P. Wick, F. Caputo, P. Boisseau, R.M. Crist, S.E. McNeil, B. Fadeel, L. Tran, S.F. Hansen, N.B. Hartmann, L.P.W. Clausen, L.M. Skjolding, A. Baun, M. Agerstrand, Z. Gu, D.A. Lamprou, C. Hoskins, L. Huang, W. Song, H. Cao, X. Liu, K.D. Jandt, W. Jiang, B.Y.S. Kim, K.E. Wheeler, A.J. Chetwynd, I. Lynch, S.M. Moghimi, A. Nel, T. Xia, P.S. Weiss, B. Sarmiento, J. das Neves, H.A. Santos, L. Santos, S. Mitragotri, S. Little, D. Peer, M.M. Amiji, M.J. Alonso, A. Petri-Fink, S. Balog, A. Lee, B. Drasler, B. Rothen-Rutishauser, S. Wilhelm, H. Acar, R.G. Harrison, C. Mao, P. Mukherjee, R. Ramesh, L.R. McNally, S. Busatto, J. Wolfram, P. Bergese, M. Ferrari, R.H. Fang, L. Zhang, J. Zheng, C. Peng, B. Du, M. Yu, D.M. Charron, G. Zheng, C. Pastore, On the issue of transparency and reproducibility in nanomedicine, *Nat Nanotechnol*, 14 (2019) 629-635.
- [166] K.H. Chen, D.J. Lundy, E.K. Toh, C.H. Chen, C. Shih, P. Chen, H.C. Chang, J.J. Lai, P.S. Stayton, A.S. Hoffman, P.C. Hsieh, Nanoparticle distribution during systemic inflammation is size-dependent and organ-specific, *Nanoscale*, 7 (2015) 15863-15872.
- [167] J. Pradal, P. Maudens, C. Gabay, C.A. Seemayer, O. Jordan, E. Allemann, Effect of particle size on the biodistribution of nano- and microparticles following intra-articular injection in mice, *Int J Pharm*, 498 (2016) 119-129.

- [168] F. Dormont, M. Rouquette, C. Mahatsekake, F. Gobeaux, A. Peramo, R. Brusini, S. Calet, F. Testard, S. Lepetre-Mouelhi, D. Desmaele, M. Varna, P. Couvreur, Translation of nanomedicines from lab to industrial scale synthesis: The case of squalene-adenosine nanoparticles, *J Control Release*, 307 (2019) 302-314.
- [169] A. Lamprecht, Nanomedicines in gastroenterology and hepatology, *Nat Rev Gastroenterol Hepatol*, 12 (2015) 195-204.
- [170] M. Zhou, J. Hou, Z. Zhong, N. Hao, Y. Lin, C. Li, Targeted delivery of hyaluronic acid-coated solid lipid nanoparticles for rheumatoid arthritis therapy, *Drug Deliv*, 25 (2018) 716-722.
- [171] S. Kanhai, I. Stavrakaki, W. Gladdines, P. J. Gaillard, E. S. Klaassen, G. J. Groeneveld, Glutathione - PEGylated liposomal methylprednisolone in comparison to free methylprednisolone: slow release characteristics and prolonged lymphocyte depression in a first - in - human study, *British Journal of Clinical Pharmacology*, 84 (2018) 1020-1028.
- [172] D. Paithankar, Girish Munavalli, Arielle Kauvar, Jenifer Lloyd, Richard Blomgren, Linda Faupel, Todd Meyer, Samir Mitragotri, Ultrasonic delivery of silica-gold nanoshells for photothermolysis of sebaceous glands in humans: Nanotechnology from the bench to clinic, *Journal of Controlled Release*, 206 (2015) 30-36.


Bacteria Injury

1 INDUCERS

2 SENSORS

3 MEDIATORS

4 EFFECTORS




Anti-inflammatory nanomedicines


Drug delivery strategy

Main therapeutic targets

Types of nanoparticles



Targeting moieties




PASSIVE TARGETING




ACTIVE TARGETING



Targeted receptors
(e.g. folate receptor, ICAM, VCAM)



Macrophage



Endothelial cell



Nucleic acids
(e.g. mRNA, genes)



Circulating proteins
(e.g. cytokines)




Table 1: Some examples of new innovative nanomedicine systems for pre-clinical therapies against inflammatory disorders

Nanocarrier	Active principle	Inflammatory lesion	Animal model	Treatment time	Tested doses and route of administration	Main results	Ref
Eudragit S100 polymeric NPs	Prednisolone	Healthy	Wistar Albino rats	Every 30mins, from 1h to 6h	5 mg/kg of NPs, oral administration	<ul style="list-style-type: none"> • Specific colon targeting with a lag time 	[45]
Pegylated liposomes	Prednisolone	Renal ischemia and reperfusion injury	8-week-old male LEW/HanHsd rats	96h	10 mg/kg of NPs, intravenous (i.v.) injection	<ul style="list-style-type: none"> • Specific accumulation in inflamed kidney • Increase of anti-inflammatory macrophages recruitment • Decrease of MCP-1 mRNA production 	[47]
Hyaluronic acid-coated solid lipid NPs	Prednisolone	Collagen-antibody induced model of arthritis in Freud's adjuvant	6-week-old male DBA/1 mice	4 injections, once every three days	15 mg/kg of equivalent prednisolone each time, intra-articular injections	<ul style="list-style-type: none"> • Selective accumulation in inflamed tissue • Good therapeutic efficacy based on analysis of joints, pannus formation, bone preservation and reduction of pro-inflammatory cytokines 	[170]
Nano-sized elastic niosome	Prednisolone	Clove oil-induced severe ocular inflammation	Albino rabbits	Every 4h, 3 times a day, for 6 days	Equivalent to 500 µg of prednisolone (A or P) each time, ocular delivery	<ul style="list-style-type: none"> • Time for complete healing reduced to half • Significant decrease of intraocular pressure elevation 	[50]
Polycaprolactone NPs dispersed in fibrin glue-based gel system	Methyl-prednisolone sodium succinate	Acute spinal cord injury	Male Wistar Albino rats	24h	Doses = N/A, topical and intraperitoneal administration	<ul style="list-style-type: none"> • Reduced damage on spinal cord • Decrease in caspase 3 and pro-inflammatory cytokines levels 	[51]
Hyaluronic acid hydrogel-based hydroxyl-terminated polyamidoamine dendrimers	Dexamethasone	Corneal alkali burn model	8-week-old Lewis rats	24h, 72h, 7 days or 14 days	Equivalent to 1.76 mg /kg of Dex, subconjunctival injection	<ul style="list-style-type: none"> • 1.6-fold better anti-inflammatory activity at 10-fold lower concentration than free drug in vitro • <i>In vivo</i>: corneal features improvements, and decrease in macrophage infiltration 	[52]

Poly(ethylene glycol)-poly(ϵ -caprolactone) polymer micelles	Dexamethasone	Adjuvant arthritis model by mycobacteria inoculation in Freund's adjuvant	6-8 weeks Wistar rats	4 injections of Dex, on days 14, 16, 18, 20 after arthritis induction	0.8 mg/kg of Dex, intravenous injection	<ul style="list-style-type: none"> • Paw swelling and erythema suppression • Down-regulation of pro-inflammatory cytokines • Protection of articular cartilage and bone from degradation and erosion 	[53]
Spherical polymeric nanoconstructs	Dexamethasone	Dextran sodium sulfate induced model of arthritis	8-week-old female C57Bl/6J mice	At day 2 after lesion: 6 consecutive treatments distributed until 16 days	5 mg/kg each time, i.v. injections	<ul style="list-style-type: none"> • Reduced weight loss and rectal bleeding • Reduced macrophage infiltration and pro-inflammatory cytokines production 	[54]
Solid lipid NPs	Dexamethasone and Butyrate	Dextran sodium sulfate induced model of arthritis	8-week-old male BALB/c mice	Daily treatment started from day 6 after lesion, for 3 days	0.1 mg/kg of dexamethasone and 4 mg/kg of Butyrate, oral administration	<ul style="list-style-type: none"> • Significant decrease of pro-inflammatory cytokines, more effectively at doses 10-fold lower than with free drugs 	[55]
N-(2-hydroxypropyl)methacrylamide (HPMA) copolymer NPs	Dexamethasone	Adjuvant-induced arthritis	Male Lewis rats	One injection, at day 14, 15, 16 or 17 after arthritis induction	10 mg/kg of dexamethasone, intravenous injection	<ul style="list-style-type: none"> • Better systemic bone quality and bone protection • Amelioration of joint inflammation 	[56]
Dual pH- and time-dependant polymeric NPs composed of Eudragit FS30D and Eudragit RS100	Budesonide	Dextran sodium sulfate-induced model of colitis	7-week-old mice	2h, 6h, or 10h	0.5 mg/kg of drug (0.168 mg/kg of budesonide), oral administration	<ul style="list-style-type: none"> • Improved colon-specific drug targeting • Body characteristics improvement 	[58]
PLGA NPs	Budesonide	Oxazolone-mediated experimental colitis	8 to 12-week-old BALB/c mice	18h	42 μ g of NPs, oral administration	<ul style="list-style-type: none"> • Selective targeting at the site of inflammation 	[60]
Phosphatidyl serine:phosphatidyl choline (3:7 mol/mol) liposomes	Recombinant mouse IL-10	12 weeks High Fat Diet-induced obesity and atherosclerosis	C57Bl/6 mice	One treatment after 12 weeks of diet	1 μ g/mouse of equivalent rIL-10, intraperitoneal injection	<ul style="list-style-type: none"> • Significant decrease in visceral fat weight • Reduced size of adipocytes • Reduced IL-6 and TNF-α secretion, and level of 	[69]

Biodegradable polyester polymeric NPs made with core polymer I, II and PLGA-PEG-CoIV	Recombinant mouse IL-10	Zymosan A (0.2 mg/mouse)-induced peritonitis (1) and 12 weeks of Western Diet-induced atherosclerosis (2)	8 to 10-week-old female C57Bl/6J (1) and 8 to 10-week-old male Ldlr ^{-/-} (2)	4h treatment (1) and 1/week for 4 weeks (2)	100 or 500 ng/mouse of equivalent rIL-10 (1) and 5 µg of rIL-10/ i.v. injection (2)	alanine amino transferase (decreased liver injury) <ul style="list-style-type: none"> •Improvement of fibrous cap thickness •Reduction of necrotic core area •Tempered inflammation in peritonitis with decrease in polymorphonuclear neutrophils infiltration •Better macrophage targeting ability 	[70]
Poly(NIPAm-co-AMPS) NPs	Anti-inflammatory cell penetrating peptide KAFAK	Osteoarthritis model induced by removal of native aggrecan of cartilage explants and IL1β treatment	3-month-old bovine knee joints explants	One treatment after 2 days of culture	50 µg NPs, treatment on cartilage explants	<ul style="list-style-type: none"> •Specific targeting of inflamed joint tissues •Reduction of pro-inflammatory cytokines 	[79]
NGPEGSS NPs system incorporating NIPAm, AMPS, BAC, and PEG polymers	Anti-inflammatory cell penetrating peptide KAFAK	Osteoarthritis model by removal of native aggrecan cartilage explants and IL1β treatment	3-month-old bovine knee joints explants	One treatment after 2 days of culture	50 µg NPs, treatment on cartilage explants	<ul style="list-style-type: none"> •Specific targeting of inflamed joint tissues Significant suppression of IL-6 production on days 4,6 and 8 	[81]
Chitosan/poly(γ-glutamic acid) NPs	Diclofenac	LPS-induced inflammation	<i>In vitro</i> human macrophages	24h, 48h	0.7 mg/mL, <i>in vitro</i> treatment	<ul style="list-style-type: none"> •Rapid internalization (within 3 hours) without toxicity below 0.7 mg/mL •Stimulation of production of prostaglandin E2 Reduction of IL-6 production 	[64]
Mannose-modified trimethyl chitosan-cysteine NPs	TNF-α siRNA	Acute hepatic injury induced by LPS and D-GaIN	Male Sprague-Dawley rats	2h	20 µg or 50 µg of equivalent TNF-α siRNA/kg, oral administration	<ul style="list-style-type: none"> •Effective TNF-α knockdown at low doses •Dramatic alleviation of inflammatory conditions in the liver (congested central vein, swollen, disarranged hepatocytes, and broken cytolemma) 	[82]

Nanoemulsion formulation of cationic lipid DOTAP	TNF- α siRNA	LPS intranigral injection-induced Parkinson's disease model	Male Sprague-Dawley rats	6h, 24h	1.5 mg/kg of NPs, intranasal administration	<ul style="list-style-type: none"> • Preferential concentration into the brain • Brain downregulation of TNF-α 	[115]
Poly(ethylene glycol)-poly(caprolactone) polymer micelles conjugated with Tat cell-penetrating peptide and siRNA targeting TNF- α	TNF- α siRNA	transient middle cerebral artery occlusion model of cerebral ischemia-reperfusion injury	10-week-old Sprague-Dawley male rats	At 30 min after middle cerebral artery occlusion	Equivalent of 30 μ g of siRNA, intranasal administration	<ul style="list-style-type: none"> • Transport of the drug to the brain directly across the nasal mucosa, without going through the BBB • Reduced infarcted area and better neurological scores • Significant suppression of TNF-α production 	[84]
Dendrimers functionalized with cationic pyrrolidinium or morpholinium surface groups	TNF- α siRNA	LPS-induced acute lung inflammation model by intranasal administration of LPS	6 to 8-week-old female CD-1 mice	28h, 96h	Pre-treatment 24h before inflammation, with 2 mg/kg of eq. siRNA concentration, intranasal administration	<ul style="list-style-type: none"> • Specific targeting in macrophages • Major inhibition of TNF-α • Modulation of all other cytokines (strong anti-inflammatory effect) 	[85]
Modified chitosan nano-carrier deploying folic acid, diethylethylamine and PEG groups	TNF- α siRNA	Collagen-antibody induced model of arthritis	8 to 12-week-old female DBA/1 mice	Treatment at day 1, 3, 5, 7 after induction of inflammation and sacrifice at day 10	4 intraperitoneal injections at 50 μ g of siRNA each	<ul style="list-style-type: none"> • Significant decrease of inflammation observed by improved clinical scores and lower TNF-α concentration in inflamed tissues • Decrease in articular cartilage destruction and bone loss 	[86]
Tuftsin/alginate NPs	IL-10 pDNA	Adjuvant arthritis model by inoculation with <i>Mycobacterium butyricum</i> in Freund's adjuvant	Male Lewis rats	From 1h to 24h	Intraperitoneal injection at day 19 post-lesion of 3 mg of NPs	<ul style="list-style-type: none"> • Specific targeting of macrophages in inflamed paws • Continuous recruitment of M2 macrophages to inflamed tissues • Sustained local and systemic IL-10 expression 	[90]

Hyaluronic acid/polyethyleneimine NPs	IL-10 and IL-4 pDNAs	Intraperitoneal injection of Brewer-thioglycollate medium to recruit peritoneal macrophages	6 to 8-week-old C57Bl/6 mice	48h	2h post inflammation, intraperitoneal injection of 100 µg of equivalent pDNA	<ul style="list-style-type: none"> • Effective targeting of peritoneal macrophages over-expressing CD44 receptors • Significant increase of Arg/iNOS ratio and CD163 expression • Reduced level of TNF-α and IL-1β in peritoneal macrophages and fluid • Increased IL-10 and IL-4 expression 	[91]
Hyaluronic acid/chitosan NPS	Cytokine response modifier A pDNA	Anterior cruciate ligament transection model of osteoarthritis	6 to 8-week-old Male Sprague-Dawley rats	3 injections, every 4 weeks starting 4 weeks after lesion	4 µg of NPs/rat each time, i.v. injections	<ul style="list-style-type: none"> • Significant inhibition of cartilage damage, synovial inflammation, and loss of type II collagen • Down regulation of IL-1β, MMP-3 and MMP-13 in synovial tissues 	[48]
Mannose-functionalized dendrimeric NPs using polyamidoamine dendrimer	Liver X receptor (LXR) ligand T0901317	12 weeks of Western Diet to induce atherosclerosis	Ldlr ^{-/-} mice	Once per week for 4 weeks after the 12 weeks of diet	200 µg of NPs by i.v. injection	<ul style="list-style-type: none"> • Specific uptake by macrophages and not hepatocytes in atherosclerotic plaques • Increased expression of LXR target genes (ABCA1/ABCG1) • Enhanced cholesterol efflux • Significant reduction in atherosclerotic plaque progression, plaque necrosis, and plaque inflammation 	[108]
Biodegradable diblock PLGA-b-PEG copolymer NPs	Liver X receptor (LXR) ligand GW3965	Zymosan A-induced peritonitis	Ldlr ^{-/-} mice	5h treatment (1) or over 2 weeks (2)	10 mg/kg of equivalent agonist 1h prior to Zymosan injection (1) or 6 injections of 10 mg/kg (2), i.v. injections	<ul style="list-style-type: none"> • Significantly more efficient than free drug for inducing LXR target gene expression and suppressing inflammatory factors • Reduction of CD68-positive macrophage content of plaques (by 50%) 	[109]

Phospholipid-reconstituted ApoA-I peptide-derived synthetic HDL	Liver X receptor (LXR) ligand T0901317	Atherogenic diet for 14 weeks	8-week-old male ApoE-deficient mice	3 times a week for 6 weeks	30 mg/kg of NPs, which is equal to 1.5 mg/kg of ligands, i.v. injections	<ul style="list-style-type: none"> • Specific accumulation in plaque and plaque regression • Up-regulation of expression of ATP-binding cassette transporters and increased cholesterol efflux in macrophages 	[110]
Raspberry-like core/satellite NPs with methyl viologen-functionalized polymeric NPs (core) and azobenzene (corona)	IL-1Ra antagonist protein	Healthy	8 to 10-week-old male Sprague-Dawley rats	One treatment	5 µg of cy7-labelled NPs, intra-articular injection	<ul style="list-style-type: none"> • Improved retention time due to prolong half-life time of IL-1Ra in the joint compared to soluble form 	[77]
Oleic acid-incorporated liquid crystalline monoolein-based NPs	Tacrolimus (FK506)	Daily topical dose (1.5 mg) of Imiquimod on shaved back for 5 days to induce a skin inflammation (psoriasis)	8 to 11-week-old BALB/c mice	Once a day, for 7 days after inflammation induction	30 µg of equivalent Tacrolimus each time, topical administration	<ul style="list-style-type: none"> • Significant increase in the amount of drug permeated and retained • More efficient in the treatment of skin inflammation than Tacrolimus in propylene glycol 	[111]
Positively charged PEP-PEG-PBG polymer micelles	FK506	Dry eye disease animal model by subcutaneous injection of scopolamine	6-week-old C57BL/6 mice	4 times a day for 5 days	Topical application of 30 µg of the formulation each time	<ul style="list-style-type: none"> • Significantly reduce apoptosis of corneal epithelium • Suppression of inflammatory-related factors (TNF-α, IL-6, MMP-9...) 	[112]
PEGylated liposomes	FK506	Experimental autoimmune myocarditis model induced by immunization with porcine myosin	7-week-old male Lewis rats	Treatment on day 14 and 17, sacrificed on day 21	0.035 mg/kg, 0.17 mg/kg or 0.35 mg/kg of equivalent drug per treatment, i.v. injection	<ul style="list-style-type: none"> • Increased level of FK506 in both plasma and hearts • Suppression of pro-inflammatory cytokine expression such as TNF-α and IFNγ • Reduced inflammation and fibrosis in the myocardium 	[113]
PEGylated liposomes	Cyclosporin A	Transient middle cerebral artery	Male Wistar rats	5min after ischemia, for 48h	2.5 mg/kg of equivalent	<ul style="list-style-type: none"> • Significant recovery of the infarct size, brain oedema, 	[114]

		occlusion by 90 min brain ischemia induction, followed by 48h reperfusion			cyclosporin A, i.v. injection	and neurological activities compared to free drug	
Omega-3 fatty acid rich flaxseed oil-based nanoemulsion system	Cyclosporin A	LPS model of neuroinflammation	Sprague-Dawley rats	9h	Pre-treatment 3h prior to inflammation, with 5 mg/kg of NPs, intranasal administration	<ul style="list-style-type: none"> • Inhibition of inflammatory responses including MPO activity and TNF-α levels • Higher uptake in major regions of the brain • Inhibition of pro-inflammatory cytokines 	[115]
Silicon dioxide NPs	Mesalazine	5% dextran sodium sulphate-induced ulcerative colitis model	8 to 9-week-old male BALB/c mice	Once a day, for 7 days, sacrifice on day 8	100 mg/kg per day of NPs, oral administration	<ul style="list-style-type: none"> • Significant decrease of IL-6 and TNF-α cytokine production and MPO activity 	[116]
Squalene-adenosine NPs	Tocopherol (VitE)	Endotoxemia model of inflammation induced by intraperitoneal injection of LPS	(8-12)-week-old Male C57BL/6 and female BALB/c mice	One treatment, 30 min after LPS injection	15 mg/kg SQAd NPs (5.5 mg/kg of equivalent adenosine) and 15 mg/kg VitE, intravenous injection	<ul style="list-style-type: none"> • Significant decrease in TNF-α and increase in IL-10 production • Significant reduction of MCP-1 and IL-6 in the lung and kidney • Mitigation of MDA levels • Improved survival rate and clinical scores in lethal LPS model 	[129]

Table 2: Some examples of nanomedicines under clinical trials

Nanocarrier	Active principle	Disease	Patients and clinical phase	Main Results	Ref.
PEGylated liposomes	Prednisolone	Ilio-femoral atherosclerosis	14 patients, mean age 70 years (\pm 7 years)	<ul style="list-style-type: none"> the treatment did not reduce arterial wall permeability or inflammation prolonged circulation half-life of prednisolone 	[150]
PEGylated liposomes	Prednisolone	Moderate to severe active ulcerative colitis	18 patients (iv administration)	<ul style="list-style-type: none"> 54 weeks after initiation of treatment only 4 patients out of 13 in remission 	[155]
Liposome	Cyclosporine (lipogel or cream)	Chronic plaque psoriasis	38 patients (receiving cyclosporine lipogel or conventional cyclosporine cream or placebo)	<ul style="list-style-type: none"> 14 weeks after initiation of treatment, complete clearance was observed in lesion sites in 41% patients treated with cyclosporine lipogel and none for patients treated with cyclosporine cream or placebo gel 	[156]
Glutathione-PEGylated liposomes	Methyl prednisolone	--	Assessment of safety, pharmacokinetics and pharmacodynamics	<ul style="list-style-type: none"> slow release formulation with reduced toxicity and prolonged decrease in the lymphocyte count 	[171]
PLGA	Pitavastatine	Healthy volunteers	40 patients, Phase I, iv administration	<ul style="list-style-type: none"> well tolerated no significant adverse response 	[158]
Nanoemulsion (oil in water emulsion)	Tretionin	Acnee vulgaris	10 patients with mild to moderate acnee vulgaris lesions	<ul style="list-style-type: none"> clinical safety of the formulation in humans 	[159]
PEGylated silica - gold nanoshells	Photothermolysis (ultrasound 10 sec to 10 W/cm ²)	Acnee vulgaris	37 patients	<ul style="list-style-type: none"> efficacy in inducing photothermal disruption of acne vulgaris 	[172]

PRE-CLINICAL MODELS

CLINICAL ASSESSMENTS

