



HAL
open science

Impact on mental health of the COVID-19 outbreak among community pharmacists during the sanitary lockdown period

M. Lange, S. Joo, P.-A. Couette, S. de Jaegher, F. Joly, X. Humbert

► **To cite this version:**

M. Lange, S. Joo, P.-A. Couette, S. de Jaegher, F. Joly, et al.. Impact on mental health of the COVID-19 outbreak among community pharmacists during the sanitary lockdown period. *Annales Pharmaceutiques Françaises*, 2020, 78, pp.459 - 463. 10.1016/j.pharma.2020.09.002 . hal-03494055

HAL Id: hal-03494055

<https://hal.science/hal-03494055v1>

Submitted on 7 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Impact on mental health of the COVID-19 outbreak among community pharmacists during the sanitary lockdown period

Impact de l'épidémie de COVID-19 sur la santé mentale des pharmaciens d'officine pendant la période de confinement

Running head: COVID-19 psychological impact in community pharmacists

Marie LANGE^{1,2}, Sarah JOO³, Pierre-André COUETTE³, Sophie de JAEGHER³, Florence JOLY^{1,2}, Xavier HUMBERT³

1 Clinical Research Department, Centre François Baclesse, 14000 Caen, France

2 Normandie Univ, UNICAEN, INSERM, ANTICIPE, 14000 Caen, France

3 Normandie Univ, UNICAEN, UFR Santé, Family Medicine Department, 14000 Caen, France

Corresponding author:

Dr Xavier HUMBERT

EA 4650 Signalisation, électrophysiologie et imagerie des lésions d'ischémie-reperfusion myocardique

Département de médecine générale

Pôle de formation et de recherche en santé, 2, rue des Rochambelles, 14000 Caen, France.

Tel : +332.31.56.82.21; Fax : +332.31.56.82.15 ; e-mail: xavier.humbert@unicaen.fr

Abstract

Objectives: COVID-19 outbreak can impact mental health including health care workers. The aim of this study was to assess the psychological impact of COVID-19 in French community pharmacists.

Material and methods: We carried out a postal-based survey to assess the psychological impact of COVID-19 in French owner community pharmacists based on three validated self-report questionnaires: Perceived Stress scale, Impact of Event Scale-revised and Maslach Burnout Inventory.

Results: The sample consists of 135 community pharmacists. Twenty-three pharmacists reported significant post-traumatic stress symptoms (17%). High burnout symptoms were found in 33 (25%), 46 (34.9%) and 4 (3%) participants. Females scored higher than males for all questionnaires ($p=0.01$).

Conclusions: This study is the first study which showed the psychological impact of COVID-19 in community pharmacists. Based on validated self-report questionnaires, up to 35% of pharmacists reported psychological disturbances. Interventions to promote psychological well-being of healthcare workers need to be developing.

Keywords: COVID-19, mental health, pharmacists, burnout

Résumé

Objectif : L'épidémie de COVID-19 a pu avoir un impact sur la santé mentale y compris celle des professionnels de santé. L'objectif de cette étude était d'évaluer l'impact psychologique du COVID-19 de pharmaciens d'officine français.

Matériels et méthodes : Nous avons réalisé une enquête postale pour évaluer l'impact psychologique du COVID-19 chez des pharmaciens d'officine français titulaires à partir de 3 auto-questionnaires validés : l'échelle de stress perçu, l'échelle révisée *Impact of Event* et l'inventaire de Burnout de Maslach.

Résultats : L'échantillon consistait en 135 pharmaciens d'officine. Ving-trois pharmaciens rapportaient des symptômes significatifs de stress post-traumatique (17%). Un burnout élevé était observé chez 33 (25%), 46 (34, 9%) et 4 (3%) participants. Les femmes avaient des scores plus élevés que les hommes à l'ensemble des questionnaires.

Conclusion : Cette étude est la première montrant l'impact psychologique du COVID-19 chez les pharmaciens d'officine. Basée sur des questionnaires validés, jusqu'à 35% des pharmaciens rapportent des difficultés psychologiques. Des interventions visant à promouvoir le bien-être psychologique des professionnels de santé devraient être développées.

Mots clés : COVID-19, santé mentale, pharmaciens, burnout

1. Introduction

In December 2019, a novel coronavirus (named SARS-CoV-2, COVID-19) was identified as causing a cluster of pneumonia cases in Wuhan in China [1]. On March 2020, the COVID-19 outbreak was declared as a pandemic by the World Health Organization.

As healthcare workers, pharmacists were played a role in hindering the spread of coronavirus by increasing patient awareness, especially by advising them on precautionary measures and providing appropriate information [2,3]. Furthermore, they were sometimes the unique source to provide protection material such as surgical masks or hydroalcoholic solution. Community pharmacists appeared like one of the most accessible healthcare professionals during this public health crisis [3].

COVID-19 outbreak can impact mental health including health care workers. Several studies had been start to be published which reported psychological symptoms during COVID-19 outbreak in general population [4,5] or in health care workers [6–10], including stress, anxiety and sleep disturbances.

The main objective of this study was to assess the psychological impact of COVID-19 in French community pharmacists. Secondary objectives were to assess the relation between psychological scores and other variables such as epidemic location status, change in pharmacy organization, geographic areas and gender. This study was conducted in the French region of Normandy, an area less affected by the COVID-19 pandemic than some other regions in the country.

2. Material and Methods

We carried out a postal-based survey the 15th April 2020 to assess the impact of COVID-19 on the mental health of all owner community pharmacists who worked in the French region of

Normandy (departments of Calvados, Orne and Manche), one month after COVID-19 sanitary lockdown. No recall of the sending was made. This survey was approved by Ethic Department of University of Caen Normandy (Authorization n° TG_COMPO_PEDAGO_SANTE_14-20180529-01R1, 6th April 2020).

It consisted of sociodemographic, geographic areas (rural or urban [11]), changes in work organization and three psychological validated self-report questionnaires. High epidemic location has been defined by areas where localized clusters have been identified before COVID-19 lockdown (4 clusters identified at the time of the study).

The self-report questionnaires used were: the Perceived Stress scale (PSS) [12] (higher score represents high level of stress), the Impact of Event Scale-revised (IES-R) [13] (higher score represents high level of stress and significant post-traumatic stress symptoms were defined by a score ≥ 33) and the Maslach Burnout Inventory (MBI ; emotional exhaustion (EE), depersonalisation (DP) and personal accomplishment (PA)) (Jackson et al., 1996[14]. Higher scores for EE and DP represent higher burnout symptoms. Higher score of PA represents lower burnout symptoms. High burnout symptoms were defined by scores ≥ 30 and ≥ 12 , respectively for EE and DP and ≤ 33 for PA.

Statistical analysis

Data were expressed as means and standard deviations (SD) or percentage. Comparisons between groups (gender or exposition) were done by *t* Student's test or chi-2 test, where appropriate, using NCSS version 9 (Hintze J, Kaysville, Utah, USA: www.ncss.com).

4. Results

The response rate was 31.1% ($n=135$; 434 questionnaires were sent). The sample consists of 135 community pharmacists (59.1% females (3 missing responses), mean age= 47.9 ± 11.4 ; Table 1). Forty six worked in urban areas (34.1%). The pharmacies localized in high epidemic location represents 24.4% of the sample ($n=33$).

Among them 119 (88.1%) had change work organization since the epidemic situation (including dedicated waiting area, sanitized of area, hydroalcoholic solution for patients...).

The mean scores of the PSS was 15.9 (SD=7.4).

The mean scores of the IES-R was 20.6 (SD=15.1). Twenty-three pharmacists reported significant post-traumatic stress symptoms (17%).

The mean scores of the EE, DP and PA were respectively 23.0 (SD=11.4), 10.9 (SD=5.5) and 48.1 (SD=7.2). High burnout symptoms were found in 33 (25%), 46 (34.9%) and 4 (3%) participants according respectively, EE, DP and PA scores.

Epidemic location, geographic areas, change in pharmacy organization were not significantly associated with psychological scores.

Females scored higher than males for PSS, IES-R and DP ($p=0.01$, 0.01 and 0.001 respectively, Table 2).

5. Discussion

This study is the first study which showed the psychological impact of COVID-19 in community pharmacists during the sanitary lockdown period. Based on self-report validated questionnaires,

up to 35% of pharmacists reported psychological disturbances. Females reported more stress and burnout symptoms than males.

First results on the mental health impact of the COVID-19 outbreak showed significant psychological disturbances in healthcare workers. In a sample of more than 1200 nurses and physicians who worked in China, 50% reported symptoms of depression, 45% anxiety, 34% insomnia and 71% distress [8]. In medical staff members in China, 36% had insomnia symptoms during the COVID-19 outbreak [10]. Outside China, mental health impact of COVID-19 had also showed in Oman physicians [6] and in French urologists [15]. Our findings are in line with these first results.

Beyond maintaining a stable supply of pharmaceuticals and hygiene products, community pharmacists are charged of major responsibilities including advising, informing, and educating the community [16]. They were sometimes the unique source to provide protection material to patients such as masks. Thus, the disruption in surgical mask supplies in the beginning of the sanitary lockdown period was able to induce stress to reply to the high demand of patients.

If pharmacists are including at the frontline beating the COVID-19 pandemic [17], no study focuses on psychological impact of the situation on these professionals and no study assesses burnout in healthcare workers during this epidemic crisis. We showed that 3 to 35% of community pharmacists reported psychological disturbances including high burnout symptoms and post-traumatic stress symptoms. Previous study, outside COVID-19 context, in community pharmacists in all France showed that until 37% of community pharmacists presented high burnout symptoms [18]. These results are not directly comparable because the present study was realized in the context of COVID-19 pandemic, the other not and the present one included only owner community pharmacists and was performed in only one region in France.

In our study, women reported more psychological difficulties than men. Similar effect of gender have been found in healthcare workers in China [7,8], in Oman physicians [6] and or in general population [4,5] during COVID-19 outbreak. Nevertheless, previous researches in French community pharmacies showed the surprising opposite pattern [18,19]. At the opposition of these studies, only owner pharmacists were included in our study which could be explained these differences.

Epidemic location was no significantly associated with psychological scores. A previous study also did not shown relationship between psychological scores and working with COVID-19 patients [6]. In our study, this result could be explained by the fact that Normandy region was not a high exposed COVID-19 region in France with only 4 localized clusters at the time of the study.

Others limitations of the study include small sample size and low response rate. Furthermore, we did not have reference of psychological symptoms before the sanitary lockdown period of our participants, nor control group and longitudinal data would be interesting.

6. Conclusion

This study is the first study which showed the psychological impact of COVID-19 in community pharmacists during the sanitary lockdown period. Based on self-report validated questionnaires, up to 35% of pharmacists reported psychological disturbances. Interventions to promote psychological well-being of healthcare workers need to be developing.

Acknowledgements: The authors would like to thank all participants.

Funding: no funding.

Disclosure of interest: The authors declare that they have no competing interest.

REFERENCES

- [1] Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med.* 2020;382:727–733.
- [2] Al-Quteimat OM, Amer AM. SARS-CoV-2 outbreak: How can pharmacists help? *Res Social Adm Pharm.* 2020;
- [3] Cadogan CA, Hughes CM. On the frontline against COVID-19: Community pharmacists' contribution during a public health crisis. *Res Social Adm Pharm.* 2020;
- [4] Liu S, Luo P, Tang M, et al. Providing pharmacy services during the coronavirus pandemic. *Int J Clin Pharm.* 2020;42:299–304.
- [5] Wang C, Pan R, Wan X, et al. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health.* 2020;17.
- [6] Badahdah AM, Khamis F, Mahyijari NA. The psychological well-being of physicians during COVID-19 outbreak in Oman. *Psychiatry Res.* 2020;289:113053.
- [7] Du J, Dong L, Wang T, et al. Psychological symptoms among frontline healthcare workers during COVID-19 outbreak in Wuhan. *Gen Hosp Psychiatry.* 2020;
- [8] Lai J, Ma S, Wang Y, et al. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw Open.* 2020;3:e203976.
- [9] Kang L, Ma S, Chen M, et al. Impact on mental health and perceptions of psychological care among medical and nursing staff in Wuhan during the 2019 novel coronavirus disease outbreak: A cross-sectional study. *Brain Behav Immun.* 2020;
- [10] Zhang C, Yang L, Liu S, et al. Survey of Insomnia and Related Social Psychological Factors Among Medical Staff Involved in the 2019 Novel Coronavirus Disease Outbreak. *Front Psychiatry.* 2020;11:306.
- [11] Insee - Statistiques locales [Internet]. [cited 2020 Sep 11]. Available from: <https://statistiques-locales.insee.fr/#c=home>.

- [12] Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav.* 1983;24:385–396.
- [13] Creamer M, Bell R, Failla S. Psychometric properties of the Impact of Event Scale - Revised. *Behav Res Ther.* 2003;41:1489–1496.
- [14] Jackson SE, Leiter MP, Maslach, C. MBI: Maslach burnout inventory. Palo Alto, CA: Consulting Psychologists Press; 1996.
- [15] Abdessater M, Rouprêt M, Misrai V, et al. COVID19 pandemic impacts on anxiety of French urologist in training: Outcomes from a national survey. *Prog Urol.* 2020;
- [16] Ung COL. Community pharmacist in public health emergencies: Quick to action against the coronavirus 2019-nCoV outbreak. *Research in Social and Administrative Pharmacy.* 2020;16:583–586.
- [17] Bukhari N, Rasheed H, Nayyer B, et al. Pharmacists at the frontline beating the COVID-19 pandemic. *J Pharm Policy Pract.* 2020;13:8.
- [18] Balaýssac D, Pereira B, Virof J, et al. Burnout, associated comorbidities and coping strategies in French community pharmacies-BOP study: A nationwide cross-sectional study. *PLoS ONE.* 2017;12:e0182956.
- [19] Balaýssac D, Pereira B, Virof J, et al. Work-related stress, associated comorbidities and stress causes in French community pharmacies: a nationwide cross-sectional study. *PeerJ.* 2017;5:e3973.

Table 1: Demographic, organization and psychological characteristics of pharmacists

	All sample (n=135)
Female (n, %) [missing 3]	78 (59.1)
Age (y) mean (SD)	47.9 (11.4)
Urban areas (n, %)	46 (34.1)
Change in pharmacy organization (yes; n, %)	119 (88.1)
Perceived stress scale; mean (SD)	15.9 (7.4)
Impact of Event scale; mean (SD)	20.6 (15.1)
Post-traumatic stress symptoms (n, %)	23 (17.0)
Emotional exhaustion*; mean (SD) [missing 3]	23 (11.4)
Low (n, %)	56 (42.4)
Middle (n, %) burnout symptoms	43 (32.6)
High (n, %)	33 (25.0)
Depersonalisation*; mean (SD) [missing 3]	10.9 (5.5)
Low (n, %)	18 (13.6)
Middle (n, %) burnout symptoms	68 (51.5)
High (n, %)	46 (34.9)
Personal accomplishment*; mean (SD) [missing 3]	48.1 (7.2)
Low (n, %)	117 (88.7)
Middle (n, %) burnout symptoms	11 (8.3)
High (n, %)	4 (3.0)

* Maslach Burnout Inventory

Table 2: Demographic, exposition and psychological characteristics of pharmacists according to gender

	Females (n=78)	Males (n=54)	p value
Age (y) mean (SD)	47.1 (11.0)	49.20 (11.9)	0.31
Epidemic location Covid-19			
High (n, %)	63 (80.8)	37 (68.5)	0.26
Low (n, %)	15 (19.2)	17 (31.5)	
Perceived stress scale; mean (SD)	17.4 (7.6)	14.2 (6.6)	0.01
Impact of Event scale; mean (SD)	23.7 (16.2)	16.7 (12.32)	0.01
Emotional exhaustion*; mean (SD)	24.1 (12.2)	22.0 (10.1)	0.29
Depersonalisation*; mean (SD)	9.5 (4.3)	12.9 (6.4)	0.001
Personal accomplishment*; mean (SD)	48.3 (7.0)	47.5 (7.7)	0.66

* Maslach Burnout Inventory