



HAL
open science

Environmental influence on canopy phenology in the dry tropics

Frédéric Dô, Venceslas A Goudiaby, Olivier Gimenez, Amadou L Diagne, Mayecor Diouf, Alain Rocheteau, Leonard E Akpo

► **To cite this version:**

Frédéric Dô, Venceslas A Goudiaby, Olivier Gimenez, Amadou L Diagne, Mayecor Diouf, et al.. Environmental influence on canopy phenology in the dry tropics. *Forest Ecology and Management*, 2005, 215 (1-3), pp.319-328. <10.1016/j.foreco.2005.05.022>. <hal-03498945>

HAL Id: hal-03498945

<https://hal.science/hal-03498945v1>

Submitted on 25 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

**ENVIRONMENTAL INFLUENCE ON CANOPY PHENOLOGY IN THE
2 DRY TROPICS**

FREDERIC C. DO¹, VENCESLAS A. GOUDIABY², OLIVIER GIMENEZ³, AMADOU L. DIAGNE⁴, MAYECOR
4 DIOUF², ALAIN ROCHETEAU¹, LEONARD E. AKPO²

¹ *IRD-UR060, Centre d'Ecologie Fonctionnelle et Evolutive, 1919 Rte de Mende, 34293 Montpellier France*

6 ² *Faculté des Sciences et techniques (Biologie Végétale), Université Cheikh Anta Diop, BP 5005 Dakar, Sénégal*

³ *CNRS, Centre d'Ecologie Fonctionnelle et Evolutive, 1919 Rte de Mende, 34293 Montpellier France*

8 ⁴ *Institut des Sciences de l'Environnement, Université Cheikh Anta Diop, BP 5005 Dakar, Sénégal*

10

Abstract

12 In the dry tropics, leaf activity of dominant woody species in the dry season is fundamental for
ecosystem vitality. Prediction of the potential impact of climate change on leaf phenology of these
14 species is a major challenge in Tropical Ecology. However, the environmental controls of inter-annual
variation of leaf flushing and leaf fall are poorly understood. We focused on a comparison between the
16 two major drivers considered in the dry tropics: soil water availability and atmospheric conditions. We
performed a 5 year phenological study of 39 individual trees of *Acacia tortilis* ssp. *raddiana* in the
18 northern Sahelian zone. Our data show that a significant inter-annual variation, up to 2.0 months,
affects both the timing of the peaks of leaf flush and leaf fall. The full canopy stage lasted between 5
20 and 8 months, broadly including the rainy season and the “cool” dry season, the timing of leaf flush
peak varying from one month before the first rains to one month after. The trees have access to a deep
22 water table (31 m) which had no significant inter-annual variation. Predictions of canopy fullness from
upper soil water availability (SWA), temperature, relative humidity and vapor pressure deficit (VPD),
24 were compared applying polynomial logistic regressions pooling all the years. We found that
atmospheric conditions and particularly VPDmax (midday value) were far better predictors than SWA.
26 Simulations based on VPDmax, significantly predicted the inter-annual variations both in the timing of
leaf flush peak (RMSE = 0.5 month) and in the timing of leaf fall peak (RMSE = 0.6 month). We
28 conclude that the inter-annual variation of canopy phenology is mainly a fine tuning to atmospheric
conditions.

30

*Key words: Sahelian Africa; Acacia tortilis; canopy fullness; leaf flushing; leaf fall; vapour pressure
32 deficit; soil water availability.*

34

INTRODUCTION

36 In dry tropical regions such as Sahelian Africa, leaf activity of dominant woody species in the dry
 season is a keystone resource for domestic herbivores, wildlife and ecosystems as a whole (Le
 38 Houérou 1989). Prediction of the potential impact of climate change on the timing of leaf flushing and
 leaf fall, i.e. leaf phenology, is a new and major challenge in Tropical Ecology.

40 Authors are in agreement on a primary genetic control and an influence of day length on leaf
 phenology (Halevy and Orshan 1973; Reich and Borchert 1984; Borchert 1994 a&b; Reich 1995;
 42 Williams et al. 1997, Morellato et al. 2000). The controversy is over precisely which environmental
 control can best explain inter-annual and inter-site variations in the timings of leaf flush and leaf fall.
 44 The hypothesis that has gained most support is that of control by soil water availability through tree
 water status (Borchert 1994 a,b,c; Eamus and Prior 2001). However, several irrigation experiments in
 46 the field have shown little impact of soil water availability on leaf phenology and a number of authors
 have suggested that evaporative demand may also be an important driver (Wright and Cornejo 1990;
 48 Duff et al. 1997; Myers et al. 1998).

In conducting a five and a half years study on 39 individuals of *Acacia tortilis* in the Sahelian zone, we
 50 have chosen to investigate the issue by using an inter-annual analysis, which is rare in the bibliography
 (Poupon 1979, Corlett and Lafrankie 1998). The Sahelian zone *sensu stricto* represents one of the
 52 driest savannas with 200-400 mm y⁻¹ of rainfall and with high inter-annual variability (C.V. = 37%),
 potential evaporation is in excess of 2000 mm y⁻¹ and groundwater is usually located deeper than 30 m
 54 (Le Houerou 1989). *Acacia tortilis* (Forsk.) Hayne *ssp. raddiana* (Savi) Brenan, is a dominant tree of
 the area which is classified as semi-evergreen or semi-deciduous (Breman and Kessler 1995, Le
 56 Floc'h and Grouzis 2003), i.e. the loss of leaves which takes place in the late dry season is and its rate
 varies with years, sites and individuals. The tree regularly exhibits leaf flush before the onset of rains.
 58 It is known that its roots can reach the deep water table (Lhote 1961; Deans et al. 2004) in common
 with numerous woody species of low-rainfall areas. However, high seasonal fluctuations of predawn
 60 leaf water potential attest to the influence of soil water availability near the soil surface on tree water
 status (Berger et al. 1996). In this study, we focused on the annual attainment and maintenance of the

62 full canopy stage in the tree population. The full canopy stage of a tree is assessed when the proportion
of the branches supporting green expanded leaves represents more than 50 % of those in the canopy.
64 Such simple phenological assessment at a monthly interval is a good indicator of the seasonal
transpiration and activity of the tree (Do and Rocheteau 2003). The timings when foliation rose above
66 or fell beneath the 50 % value defined the timings of leaf flushing and leaf fall peaks, respectively.
These timings allow the calculation of full canopy stage duration. Using concurrent measurements of
68 climatic conditions and soil water availability, the study had two objectives. The first objective was to
assess the inter-annual variation in the timings of leaf flushing and leaf fall and by consequence the
70 changes in the full canopy stage duration and period in the year. The second objective was to
determine which environmental variable can best explain inter-annual change in the timings of leaf
72 flushing and leaf fall peaks. In this water-controlled biome, we focused on a comparison between soil
water availability and the following atmospheric variables: temperature, relative humidity and vapour
74 pressure deficit which is a combination of the former variables and the driving force of water loss.

METHODS

76 **Location, climate and vegetation**

The study was carried out in northern Senegal, near Souilene Village (16°20' N, 15°25' W) located 20
78 km south of Dagana in the grazing land ecosystem of the sandy Ferlo. Annual rainfall averages 280
mm, mean annual temperature is 28.7°C and average maximum and minimum temperatures are,
80 respectively: 41 and 22 °C for the hottest month (May) and 31 and 14 °C for the coldest month
(January). The sparse woody community is dominated by three species: *Acacia tortilis* and *Balanites*
82 *aegyptiaca* (L.) Del, for the tree layer (90 individuals ha⁻¹) and *Boscia senegalensis* (Pers.) Lam for the
shrub layer (50 individuals ha⁻¹). The soil is very deep and sandy (90 % sand).

84 **Population and measurements**

The studied population included 39 mature individuals located in a 1 ha area that has been protected
86 since 1989. Trunk diameter, which was measured 30 cm above the soil, varied from 5 to 45 cm, with
70 % of the population having diameters between 15 and 30 cm. Tree height ranged from 5 to 10 m.
88 Local microclimate was automatically monitored by a datalogger (21X, Campbell Scientific Ltd,

Leics, UK), which recorded hourly values of air temperature, relative humidity, incoming short wave
 90 radiation and rainfall. The water Vapour Pressure Deficit (VPD) was calculated from air temperature
 (T) and relative humidity (RH):

$$92 \quad \text{VPD} = e_s(T) - e \quad \text{and,} \quad e = \text{RH} * e_s(T)$$

where $e_s(T)$ is the saturation vapour pressure at air temperature T and e, the actual vapour pressure.

94 Soil moisture was measured to a depth of 4 meters by neutron probe (Solo 25, Nardeux S.A., Les Ulis,
 France) using six access tubes. The soil wetting front rarely exceeded one meter depth. Soil water
 96 availability (SWA) is expressed for the layer 0-100 cm deep, calculated between moisture contents at
 field capacity and the driest moisture contents which were less than those prevailing at a water
 98 potential of -1.5 MPa. The soil profile down to the water table was augered by the Geological
 Department of Dakar University (Dr Abdoulaye Faye). Samples were taken every 0.5 m. Actual
 100 moisture content and moisture equivalent to -1.5 MPa were determined in the laboratory. Over the
 period June 1995- August 2000, phenological observations and soil measurements were carried out
 102 every month in the dry season and twice a month in the rainy season, except in August 1995 and 1997.
 Climatic data were not available between 13 and 18 April 1996 and between 29 September and 10
 104 October 1996.

Assessment of phenological stages

106 For each tree, a phenological stage for the foliage was allocated according to the 6 following
 categories taking into account the proportion of branches bearing green expanded leaves in the canopy
 108 (Grouzis and Sicot 1980): 0, leafless, 0%; 1, leaf buds opening, <10 %; 2, start of leaf expansion, 10-
 50 % of the branches; 3, full canopy, > 50 %; 4, start of leaf senescence, changing colour, <50 %; 5,
 110 full leaf senescence or fall, <10 %.

Data analysis

112 To express canopy fullness, the phenological stages were converted to their midrange
 percentage value of branches bearing green expanded leaves in the canopy: 0, 0 %; 1, 5 %; 2, 30 %; 3,
 114 75 %; 4, 30 %; 5, 5%. Thus the maximum value of canopy fullness is 75 by definition.

The timings of leaf flush or leaf fall peaks were calculated for each tree by linear interpolation such as
 116 the time when the canopy fullness rose above or fell below the 50 % value in the long term (> 1

month). Time is expressed in numerical values of months (mo.), e.g. 6.5 for the middle of June.

118 Because phenological cycles are displayed across two years, we use for simplicity's sake, the year of
the rainy season to name the phenological cycle, e.g. 1998 for 1998/99. Timings of leaf fall peak
120 located at the beginning of the second year are expressed as greater than month 12, e.g. 13.5 for the
middle of January. The duration of full canopy was calculated for each tree by difference between the
122 timings of leaf fall peak and leaf flush peak.

Year effects on the duration of full canopy and the timings of leaf flush peak and leaf fall peak
124 were tested using a repeated-measures analysis of variance (SAS proc. Mixed, SAS Institute Inc.
1999). Potential influence of environmental variables (edaphic or atmospheric) on changes in canopy
126 fullness, were investigated by regression analysis between values of canopy fullness and each variable,
considering all trees and pooling data for all of the years. For atmospheric variables (temperature,
128 relative humidity and vapour pressure deficit), the average value for the 15 days preceding
phenological observations was considered. For soil water availability, the actual SWA and the amount
130 of rainfall were used. Since at the tree level, canopy fullness values were qualitative and ordinal
response variables, classified as 0, 5, 30 and 75, we specifically performed ordered polynomial logistic
132 regressions with a cumulative logit link (SAS proc. Genmod, SAS Institute Inc. 1999). This model
relates the cumulative log of odds of the response variable to the explanatory variable in a linear form
134 (Agresti 1984). We assessed the validity of the regression by computing the significance of each
variable using Wald tests (Agresti 1984). Because we wanted to compare the predictive power of
136 environmental variables, the c statistic was computed (Hosmer and Lemeshow 2000). It measures the
degree to which predicted probabilities agree with actual outcomes. The c statistic ranges from 0.5 to
138 1. A 0.5 value means that the model is no better than assigning observations randomly into outcome
categories. Variables with c statistic value greater than 80 % were therefore selected as having high
140 predictive power. The analysis gives the values of the explanatory variable (environmental variable)
corresponding to the selected probability of each response (0, 5, 30, 75). For environmental variables
142 having good predictive power, regressions were used to predict for each year, the times of leaf flush
and leaf fall peaks such as the day when the probability of full canopy (response 75) rose above or fell
144 below the 50 % value in the long term (> 1 month). Inputs were climatic data for the 15 days

preceding each day. Quality of prediction was evaluated by the root mean square error:

146
$$RMSE = \sqrt{\sum_1^n (s_i - m_i)^2 / n}$$
, s_i is the model prediction, m_i the corresponding
 observation, i the year and n the total number of years.

148

RESULTS

150 During the study period (1995-2000), annual precipitation averaged 230 mm with minima of 146 and
 153 mm in 1997 and 1996, and a maximum of 367 mm in 1999 (Table 1, Fig. 1a). It is noteworthy that
 152 inter-annual variability over the six year period (C.V. = 37%) was similar to the long-term variability
 for the area. Rains occurred mainly between July and September. The dates for onset of the rains were
 154 variable, *i.e.* about the start of June in 1996 and 1997, but the end of July in 1998. August and
 September were the wettest months. On average, 70 % of the total annual precipitation fell during
 156 these 2 months. SWA in the upper layer (0-100 cm) was closely correlated with annual rainfall (Fig.
 1b). In the wet season, SWA increased proportionally with rainfall and reached 80% for the wettest
 158 year 1999. However, by January, SWA decreased to similar values between years, (from 7 to 9 %).
 Subsequently, SWA declined to small values of 3 to 5%, before the start of the rains.

160 Atmospheric conditions in the rainy season were characterized by highest relative humidities
 (RH), above 50 %, with mean temperatures around 30°C (Fig. 1a). This was the time of the lowest
 162 evaporative demand: daily means of vapor pressure deficit (VPD) were less than 2 kPa (Fig. 1b). RH
 increased between May and June, from one to two months before the start of the rains. This is a
 164 common phenomenon induced by the northerly movement of the Inter-Tropical Convergence Zone but
 there were notable differences between years (Fig. 1a). RH decreased sharply after the rainy season,
 166 then it fluctuated, from 10 to 50 %, with notable differences between years. In the dry season, two
 climatic seasons are usually distinguished according to air temperature: the “cool” dry season lasting
 168 from December until February and the hot dry season lasting from March until June (Fig.1b, month 15
 to 17). The hot dry season with low RH and increasing temperatures is characterized by the highest
 170 evaporative demand: mean VPD generally exceeds 2 kPa and maximum VPD regularly exceeds 5 kPa.

Evolution of Canopy fullness showed an annual cycle with both similarities and differences
172 between years (Fig. 1b, Table 1). The defoliation phase was centred around the hot dry season.
Defoliation was never complete at population scale as shown by Canopy fullness which never reached
174 a zero value (Fig. 1b). The point closest to zero occurred in April 2000 (Fig. 1a: point at 16 mo. in
“1999” or 4 mo. in “2000”), which corresponded to 63 % of fully defoliated trees. The full canopy
176 stage lasted an average 7 months (Table 1), from mid-July to the start of February, broadly including
the rainy and cool dry seasons (Fig. 1). The inter-annual variation of full canopy stage duration was
178 large, e.g. more than two months between the extreme years: 5.2 mo. in 1996 and 7.5 mo. in 1998 or
1999 (Table 1). The repeated-measures analysis of variance revealed a highly significant ($P < 0.001$)
180 effect of year on canopy duration. The observation interval which was at the best biweekly and more
often monthly, precludes discussion of differences or equalities at less than 0.5 months. The
182 magnitude of differences clearly distinguishes two groups, the group of the years 1996 and 1997
where duration was shortened by 25 % in comparison with the group of the years 1995, 1998 and
184 1999. The duration differences resulted from variation in a two month range both in the timings of
peaks for leaf flush and leaf fall (Table 1). The timing of leaf flush peak varied from the start of June
186 (years 2000, 1999), more than one month before the first rains (Fig. 1), to the end of July in the rainy
season (years 1995, 1996, 1997). The situation in 1998 was intermediate: the start of leaf flushing was
188 early, in May as in 1999 and 2000, but was followed by a defoliation before leaf flush peaked in July
(Fig. 1). The timing of leaf fall peak varied from the start of January (year 1996) to the start of March
190 (year 1995). The years 1996 and 1997 had intermediate defoliation in December. The inter-individual
variation was large and variable depending upon year (standard deviation in Table 1). For example, for
192 the same average timing of leaf flush peak, the inter-individual variation was fourfold larger in 1999
than in 2000. Similar results occurred between 1995 and 1996 for the timing of leaf fall peak. At the
194 individual level, the time between the onset of flushing and full foliation was approximately 1 to 1.5
mo (data not shown). Figure 1b shows the average of canopy fullness in the population, so durations
196 were increased by an averaging effect in the case of large inter-individual variation.

The influence of SWA on the inter-annual variation of canopy phenology could involve
198 ground water and/or upper soil water. The top of the water table was found at 31 m depth, consistent

with the water level found in village wells in the study area. Between the upper layer of soil and the
200 water table, SWA for plants was very limited because the actual moisture contents were equivalent to
a water potential of -1.5 MPa. Living root fragments of *A. tortilis*, which can be identified from their
202 very special smell, were collected at 25 m depth. These confirmed that the trees have access to the
water table. However no fluctuations of the deep water table which could explain inter-seasonal
204 variability of leaf phenology, are known in the area (Gaye and Edmunds 1996).

Results of regression analysis between canopy fullness and environmental variables over all
206 years are summarized in Table 2. For upper SWA, two variables were considered: the actual SWA and
the amount of rainfall in the 15 days preceding phenological observations. In the completely dry
208 season (timing of leaf fall), changes of SWA were slow and actual SWA, measured at monthly
intervals, was a good integrator of soil conditions for the 15 day period of time. By contrast, during the
210 transition between dry and rainy seasons (timing of leaf flush), due to the effect of rainfall, measured
SWA was a poor predictor and rainfall amount (hourly recording) was used as a surrogate (see Fig. 1).
212 Whatever the analysis, there was a significant relationship ($P < 0.05$) with all environmental variables.
This was due to the fact that during the considered periods, soil water and atmospheric variables were
214 all moving broadly at the same seasonal time as canopy fullness (See Fig. 1). The key point is the
comparison between predictive powers of variables (c values in Table 2). For the whole time of
216 deciduousness (mainly from January to August), VPDmax is from far a higher predictive variable than
SWA or rainfall (Table 2). VPDmax is the value measured around midday, which defines the
218 maximum evaporative demand during the day. The value of VPDmax corresponding to a prediction of
50% of canopy fullness was 4.8 kPa.

220 When the time of deciduousness was divided between times of leaf flush and leaf fall, there
was a remarkable switch between the predictive powers of relative humidity (RH) and temperature
222 (T_a), which are combined in VPD (Table 2). During the time of leaf flush, RH and VPD are the
variables which best predicted canopy fullness. During the time of leaf fall, temperature and VPD are
224 the variables which best predicted canopy fullness. The same analysis considering a period of 30 days
before phenological observations gave similar results (data not shown).

226 Regression analyses were performed pooling all of the years. For environmental variables
having a high predictive power, regressions were used to predict for each year, the timings of leaf
228 flush and leaf fall peaks. Inputs were climatic data for the 15 days preceding each date. The day when
the probability of full canopy rose above (leaf flush) or fell below (leaf fall) the 50 % value in the long
230 term (> 1 month) was retained. The precision of “long term” is particularly important for the
prediction of the date of leaf fall peak because the dynamics of climatic variables fluctuated strongly at
232 this time (see Fig. 1). Root mean square errors of predictions are given in Table 2 and detailed results
for VPDmax are shown in Figure 2. Atmospheric conditions explained a large part of interannual
234 variation in both the timing of leaf flush peak (RMSE = 0.4- 0.5 month) and the timing of leaf fall
peak (RMSE = 0.6 – 0.7 month, Table 2). Finally, combination of Figures 2a and 2b illustrates that
236 VPDmax can explain a part of the duration and period of canopy fullness for the 5 complete years of
the study (RMSE = 0.8). The period containing October and November was not considered in the
238 regression-prediction analysis, however, simulations based on atmospheric conditions also predict
defoliation for this period, which was not systematically observed (Fig. 1).

240

DISCUSSION

242 The full canopy stage of *A. tortilis* sp. *raddiana* lasts between 5 and 8 months, broadly including the
rainy season and the “cool” dry season. The seasonal period and duration of full canopy stage is in
244 agreement with general description of its leaf phenology in Sahelian Africa (Grouzis and Sicot 1980,
Diouf and Zaafour 2003). As with the majority of species in the seasonally dry tropics, the defoliation
246 phase was centered around the hot dry season.

The first main result of this study was large inter-annual variation in the start and end of the
248 full canopy stage on the same site, more than 2 months in each case. One consequence was a 25 %
shortfall of the stage duration for two of the years (1996 and 1997). Such variation confirms that
250 genetics and day length cannot be the unique controlling factors of canopy phenology and that other
factors have complementary influence on the timings of leaf flush and leaf fall peaks. We should insist
252 that here we deal with the major timing of leaf flush and not initial leaf flush which is often discussed
in the literature (Borchert 1994c). Initial leaf flush was generally observed between April and May.

254 Finally we discuss which factors influence the timing of the massive leaf flush which follows the
initial leaf flush.

256 The second main result was that atmospheric conditions and particularly VPDmax could
explain inter-annual change of canopy fullness better than upper or deep SWA. The influence of
258 annual rainfall on canopy phenology through upper SWA is a current hypothesis in the Sahelian zone :
higher rainfall would induce a larger residual upper soil water availability in the dry season which
260 could delay the timing of defoliation and would advance the time of leaf flushing peak. Our results do
not support this hypothesis despite a large inter-annual variation in rainfall (from 146 mm in 1997 to
262 367 mm in 1999). It was particularly notable that three month after rainy season, upper SWA reached
similar and very small values in all years. How was this possible? The phenological stage or canopy
264 fullness percentage does not describe leaf number or area which were two times higher for the wettest
years, 1998 and 1999 than for 1997 (Goudiaby 2003). Biomass of grass was also greater at the end of
266 the wettest rainy seasons. These two factors involve greater water uptake, which can explain how
SWA reached similar small values.

268 The third main result was that atmospheric conditions and particularly VPDmax predicted a
significant part of the inter-annual variations in both the timing of leaf flush peak and of leaf fall peak.
270 We conclude that these inter-annual variations of canopy phenology were mainly controlled by
atmospheric conditions. Comparable data on this point are scarce in the literature although the possible
272 influence of relative humidity on leaf flush was often hypothesized. One reason may be that
phenological studies on the same individuals, which last for more than one or two years, are rare in the
274 tropics. Another reason could be that until now, VPD was not always considered as a climatic variable
influencing savanna tree phenology (De Bie *et al.* 1998, Chidumayo 2001). Moreover, the conditions
276 of water constraint (rainfall amount, evaporative demand and depth to the water table) in the
ecosystem studied here were more extreme than in savanna situations detailed elsewhere in the
278 literature (Eamus and Prior 2001). However, an inter-annual analysis with “only” 5 to 6 years cannot
definitively demonstrate that VPD is a major proximal cue of canopy phenology in the dry tropics.
280 Nevertheless, the results give a new and clear insight on the influence of atmospheric conditions.

What would be the adaptative significance of such response to atmospheric conditions? This
282 behaviour leads to maximization of the duration of canopy fullness below a VPD threshold between
4.5 and 4.9 kPa. Such behaviour maximizes the duration of photosynthetic activity under a threshold
284 of evaporative demand and does not reduce water loss. Field observation at the terminal branch scale
in 1997 and 1998 showed, that the annual phenological pattern corresponds to three main cohorts of
286 leaves (Goudiaby, 2003). The first cohort begins between April and May, the second cohort begins in
July and the last begins between December and January. The cohort of July expressed the longest leaf
288 life span around 6-7 months. The life span of the initial and final cohorts was very short and variable
depending upon the year. Leaf stomata are known to be very sensitive to VPD. We hypothesize that
290 the response of the two extreme cohorts of leaves to climatic conditions could explain inter-annual
variation of canopy phenology.

292 Do these results mean that soil water availability has no influence on leaf phenology of *Acacia tortilis*?
We do not support this. Firstly, leaf activity of trees during large periods of the dry season is only
294 possible because trees have access to extra-annual water in the deep soil. Secondly, the existence of
large spatial variation in canopy deciduousness at the landscape scale suggests a strong effect of SWA.
296 It was particularly notable at the border of the Senegal river, near Dagana city, where the water table is
close to the top soil that the tree population behaves like an evergreen species. Finally, we believe that
298 the average phenological pattern fits the prevailing local SWA's, annual rainfall amount and depth to
the water table. While the inter-annual variation of canopy phenology is a fine tuning to atmospheric
300 conditions.

In conclusion, if such large inter-annual variation of canopy phenology is common to other dominant
302 species of the seasonally dry tropics, which is fairly possible, the consequences for the potential
impact of climate change are important and not only rainfall should be considered.

304 **ACKNOWLEDGEMENTS**

We thank Youm Mbaye, Hyacinthe Tendeng and Baïdi Li for technical assistance. We are grateful to
306 Michel Grouzis for initiating the research program. The research was partly funded by the European

Commission contract INCO-DC « Minimizing competition in dryland agroforestry Contract No.
 308 ERBIC 18CT 98 0322». We specially thank J.D. Deans for English revision.

LITERATURE CITED

- 310 Agresti, A. (1984). *Analysis of ordinal categorical data*. New York: Wiley.
- Berger A., Grouzis M., Fournier C., 1996. The water status of six woody species coexisting in the
 312 Sahel (Ferlo, Senegal). *J. Trop. Ecol.* 12: 607-627.
- Borchert, R. 1994a. Soil and stem water storage determine phenology and distribution of tropical dry
 314 forest trees. *Ecology* **75**:1437-1449.
- Borchert, R. 1994b. Water status and development of tropical trees during seasonal drought. *Trees*
 316 **8**:115-125.
- Borchert, R. 1994c. Induction of rehydration and bud break by irrigation or rain in deciduous trees of a
 318 tropical dry forest in Costa Rica. *Trees* **8**:115-125.
- Borchert, R. 1998. Responses of tropical trees to rainfall seasonality and its long-term changes.
 320 *Climatic Change* **39**:381-393.
- Breman, H. and J-J. Kessler. 1995. *Woody plants in Agro-Ecosystems of Semi-Arid Regions, with*
 322 *Emphasis on the Sahelian Countries*. Springer-Verlag Berlin Heidelberg, Germany.
- Chidumayo, E. N. 2001. Climate and phenology of savanna vegetation in southern Africa. *Journal of*
 324 *Vegetation Science* **12**:347-354.
- Corlett, R. T. and JR. J. Lafrankie. 1998. Potential impacts of climate change on tropical asian forest
 326 through influence on phenology. *Climatic. Change* **39**:439-453.
- De Bie, S., P. Ketner, M. Paase and C. Geerling. 1998. Woody plant phenology in the West Africa
 328 savanna. *Journal of Biogeography* **25**:883-900.
- Deans, J. D., W. M. Edmunds, D. K. Lindley, C. B. Gaye, B. Dreyfus, J. Nizinski, M. Neyra, R.C.
 330 Munro, K. Ingleby. 2004. Nitrogen in interstitial waters in the Sahel: Natural Baseline, Pollutant
 or resource ? *Plant and Soil*, *in press*.

- 332 Do F. & A. Rocheteau 2003. Cycle annuel de transpiration d'*Acacia raddiana* par la mesure des
flux de sève brute (Nord-Sénégal). In "Un arbre au désert: *Acacia raddiana*", Editeurs
334 scientifiques M. Grouzis et E. Le Floch, IRD éditions, Paris, pp. 119-142.
- Duff, G. A., B. A. Myers, R. J. Williams, D. Eamus, A. O'Grady and I. R. Fordyce 1997. Seasonal
336 patterns in soil moisture, vapour pressure deficit, tree canopy cover and predawn water potential
in a northern Australian savanna. *Australian Journal of Botany* **45**:211-224.
- 338 Diouf M. & M. S. Zaafouri, 2003. Phénologie comparée d'*Acacia raddiana* au nord et au sud du
Sahara. In "Un arbre au désert: *Acacia raddiana*", Editeurs scientifiques M. Grouzis et E. Le
340 Floch, IRD éditions, Paris, pp. 103-118.
- Eamus, D. and L. Prior. 2001. Ecophysiology of trees of seasonally dry tropics: Comparisons among
342 phenologies. *Advances in Ecological Research*. **32**:113-197.
- Gaye, C. B. and W. M. Edmunds. 1996. Groundwater recharge estimation using chloride, stable
344 isotopes and tritium profiles in the sands of north-western Senegal. *Environmental Geology*
27:246-251.
- 346 Goudiaby V.C.A. 2003. Impact d'un déficit pluviométrique sur la feuillaison d'*Acacia tortilis* (Forsk.)
Hayne subsp. raddiana (Savi) Brenan var. *raddiana* dans le Nord-Ferlo au Sénégal. Thèse de
348 Doctorat, Faculté des Sciences et Techniques, Université Cheikh Anta Diop, DAKAR, 82 p.
- Grouzis, M. & M. Sicot. 1980. A method for the phenological study of browse populations in the
350 Sahel: the influence of some ecological factors. Pages 233-240 in H. N. Le Houérou, editor.
Browse in Africa : The current state of knowledge. ILCA edition, Addis Abeba, Ethiopia.
- 352 Halevy, G. and G. Orshan. 1973. Ecological studies on *Acacia* species in the Negev and Sinai: II.
Phenology of *Acacia raddiana*, *A. tortilis* and *A. Gerrardii negevensis*. *Israelian Journal of*
354 *Botany* **22**:120-138.
- Hosmer, D. W., Jr., & S. Lemeshow (2000). *Applied logistic regression* (2nd ed.). New York: Wiley.
- 356 IPCC. Climate change 2001: Impacts, Adaptation, and Vulnerability. Cambridge University Press,
New York, USA.

- 358 Le Floch E. & M. Grouzis, 2003. *Acacia raddiana*, un arbre des zones arides à usages
multiples. In “Un arbre au désert: *Acacia raddiana*”, Editeurs scientifiques M. Grouzis et E. Le
360 Floch, IRD éditions, Paris, pp. 21-58.
- Le Houérou, H. N. 1989. The grazing land ecosystems of the African Sahel. Ecological studies,
362 Volume 75, Springer-Verlag, Berlin.
- Lhote, H. 1961. Au sujet de l’arbre du Ténére. Bulletin de liaison Saharienne 12:49-54.
- 364 Morellato, L.P.C., D.C. Talora, A Takahasi, C.C. Bencke, E.C. Romera, and V.B. Zipparo 2000.
Phenology of atlantic forest trees: A comparative study. Biotropica 32(4b):811-823.
- 366 Myers, B. A., R. J. Williams, I. Fordyce, G. A. Duff & D. Eamus 1998. Does irrigation affect leaf
phenology in deciduous and evergreen trees of the savannas of northern Australia ? Australian
368 Journal of Ecology 23:329-339.
- Poupon H. 1979. Etude de la phénologie de la strate ligneuse à Fété Olé (Sénégal septentrional) de
370 1971 à 1977. Bulletin de l’IFAN, Tome 41, série A, n° 1:44-85.
- Reich, P. B. 1995. Phenology of tropical forest: patterns, causes, and consequences. Canadian Journal
372 of Botany 73:164-174.
- Reich, P. B. & , R. Borchert 1984. Water stress and tree phenology in tropical dry forest in the
374 lowlands of Costa Rica. Journal of Ecology 72:61-74.
- SAS Institute Inc. (1999). *SAS/STAT® user’s guide* (Version 8, Vol. 2). Cary, NC: SAS Institute Inc.
- 376 Williams, R. J., B. A. Myers, W. J. Muller, G. A. Duff, & D. Eamus. 1997. Leaf phenology of woody
species in a north Australian tropical savanna. Ecology 78(8):2542-2558.
- 378 Wright, S. J. & Cornejo, F. H. 1990. Seasonal drought and leaf fall in a tropical forest. Ecology
71:1165-1175.
- 380

LIST OF FIGURES

- 382 Figure 1: Inter-annual variation of environmental conditions and leaf phenology for *Acacia tortilis*
 growing at Souilène in North Senegal. **a)** daily rainfall, 15 days moving average of mean air
 384 temperature and relative humidity (RH) **b)** average Canopy Fullness of 39 individuals, Soil
 Water Availability (SWA) in the upper layer (0-1 m), 15 days moving average of mean air
 386 vapour pressure deficit (VPD). Time is expressed in numerical values of months, e.g. 6.5 for the
 middle of June. The study ended in August 2000.
- 388 Figure 2: Timings of leaf flush peak (a) and leaf fall peak (b) predicted by models based on the
 dynamics of maximum vapour pressure deficit (VPDmax) versus observed timings. Input was
 390 the average value of VPDmax for the preceding 15 days. Point labels specify annual
 phenological cycles. Horizontal bars indicate standard deviation of observed data (means of 39
 392 individuals). RMSE is the root mean square error (see methods).

LIST OF TABLES

- 394 Table 1: Mean values of timing of leaf flush peak, timing of leaf fall peak, and duration of full canopy
 stage expressed in months for 39 individuals of *Acacia tortilis* growing at Souilène in North
 396 Senegal. Repeated-measures ANOVA revealed highly significant effects of year ($P < 0.001$) on
 the three variables. The study ended in August 2000 before observation of the timing of leaf fall
 398 for the phenological cycle 2000-01.
- Table 2. Parameters of regression-prediction analysis testing which environmental variable can best
 400 explains inter-annual changes in canopy fullness (CF) and the timings of leaf fall and leaf flush
 peaks (50% of CF). Variables were upper soil water availability (SWA), rainfall and daily
 402 mean, minimum and maximum air temperatures (T), relative humidity (RH) and vapour
 pressure deficit (VPD). Polynomial logistic regressions were performed (see methods). C is the
 404 predictive power, which is high when (*) is added. 50% CF indicates the value of variables

406 corresponding to 50 % of Canopy Fullness. RMSE is root mean square error of predicted timings of leaf fall and leaf flush peaks against observed timings.