



HAL
open science

Black hole mass measurement using ALMA observations of [CI] and CO emissions in the Seyfert 1 galaxy NGC 7469

Dieu D. Nguyen, Takuma Izumi, Sabine Thater, Masatoshi Imanishi, Taiki Kawamuro, Shunsuke Baba, Suzuka Nakano, Jean L. Turner, Kotaro Kohno, Satoki Matsushita, et al.

► To cite this version:

Dieu D. Nguyen, Takuma Izumi, Sabine Thater, Masatoshi Imanishi, Taiki Kawamuro, et al.. Black hole mass measurement using ALMA observations of [CI] and CO emissions in the Seyfert 1 galaxy NGC 7469. *Monthly Notices of the Royal Astronomical Society*, 2021, 504, pp.4123-4142. 10.1093/mnras/stab1002 . hal-03502254

HAL Id: hal-03502254

<https://hal.science/hal-03502254>

Submitted on 20 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Black hole mass measurement using ALMA observations of [CI] and CO emissions in the Seyfert 1 galaxy NGC 7469

Dieu D. Nguyen¹,^{1,2} Takuma Izumi^{3,4}, Sabine Thater⁵, Masatoshi Imanishi^{3,4}, Taiki Kawamuro^{3,6}, Shunsuke Baba³, Suzuka Nakano³, Jean L. Turner⁷, Kotaro Kohno^{8,9}, Satoki Matsushita¹⁰, Sergio Martín^{11,12}, David S. Meier¹³, Phuong M. Nguyen¹⁴ and Lam T. Nguyen¹⁵

¹Faculty of Fundamental Sciences, Phenikaa University, Yen Nghia, Hadong, Hanoi 12116, Vietnam

²Phenikaa Institute for Advanced Study (PIAS), Phenikaa University, Yen Nghia, Hadong, Hanoi 12116, Vietnam

³National Astronomical Observatory of Japan (NAOJ), National Institute of Natural Sciences (NINS), 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

⁴Department of Astronomical Science, The Graduate University for Advanced Studies (SOKENDAI), 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

⁵Department of Astrophysics, University of Vienna, Türkenschanzstrasse 17, 1180 Wien, Austria

⁶Núcleo de Astronomía de la Facultad de Ingeniería, Universidad Diego Portales, Av. Ejército Libertador 441, Santiago, Chile

⁷Department of Physics and Astronomy, University of California at Los Angeles (UCLA), Los Angeles, CA 90095-1547, USA

⁸Institute of Astronomy, Graduate School of Science, The University of Tokyo, 2-21-1 Osawa, Mitaka, Tokyo 181-0015, Japan

⁹Research centre for the Early Universe, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo, Tokyo 113-0033, Japan

¹⁰Academia Sinica, Institute of Astronomy & Astrophysics (ASIAA), PO Box 23-141, Taipei 10617, Taiwan

¹¹European Southern Observatory, Alonso de Córdova, 3107, Vitacura, Santiago 763-0355, Chile

¹²Joint ALMA Observatory, Alonso de Córdova, 3107, Vitacura, Santiago 763-0355, Chile

¹³Department of Physics, New Mexico Institute of Mining and Technology, Socorro, NM 87801, USA

¹⁴Department of Physics, Quy Nhon University, 170 An Duong Vuong, Quy Nhon, Vietnam

¹⁵LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Univ. Paris Diderot, Sorbonne Paris Cité, 5 place Jules Janssen, F-92195 Meudon, France

Accepted 2021 April 5. Received 2021 March 5; in original form 2021 January 29

ABSTRACT

We present a supermassive black hole (SMBH) mass measurement in the Seyfert 1 galaxy NGC 7469 using Atacama Large Millimeter/submillimeter Array (ALMA) observations of the atomic-[CI](1–0) and molecular-¹²CO(1–0) emission lines at the spatial resolution of ≈ 0.3 (or ≈ 100 pc). These emissions reveal that NGC 7469 hosts a circumnuclear gas disc (CND) with a ring-like structure and a two-arm/bi-symmetric spiral pattern within it, surrounded by a starbursting ring. The CND has a relatively low $\sigma_{\text{gas}}/V \approx 0.35$ ($r \lesssim 0.5$) and ≈ 0.19 ($r > 0.5$), suggesting that the gas is dynamically settled and suitable for dynamically deriving the mass of its central source. As is expected from X-ray dominated region (XDR) effects that dramatically increase an atomic carbon abundance by dissociating CO molecules, we suggest that the atomic [CI](1–0) emission is a better probe of SMBH masses than CO emission in active galactic nuclei (AGNs). Our dynamical model using the [CI](1–0) kinematics yields a $M_{\text{BH}} = 1.78_{-1.10}^{+2.69} \times 10^7 M_{\odot}$ and $M/L_{\text{F547M}} = 2.25_{-0.43}^{+0.40} (M_{\odot}/L_{\odot})$. The model using the ¹²CO(1–0) kinematics also gives a consistent M_{BH} with a larger uncertainty, up to an order of magnitude, i.e. $M_{\text{BH}} = 1.60_{-1.45}^{+11.52} \times 10^7 M_{\odot}$. This newly dynamical M_{BH} is ≈ 2 times higher than the mass determined from the reverberation mapped (RM) method using emissions arising in the unresolved broad-line region (BLR). Given this new M_{BH} , we are able to constrain the specific RM dimensionless scaling factor of $f = 7.2_{-3.4}^{+4.2}$ for the AGN BLR in NGC 7469. The gas within the unresolved BLR thus has a Keplerian virial velocity component and the inclination of $i \approx 11.0_{-2.5}^{+2.2}$, confirming its face-on orientation in a Seyfert 1 AGN by assuming a geometrically thin BLR model.

Key words: galaxies: nuclei – galaxies: ISM – galaxies: kinematics and dynamics – galaxies: Seyfert – galaxies: spiral – (galaxies:) quasars: supermassive black holes.

1 INTRODUCTION

Supermassive black holes (SMBHs) appear to be ubiquitous at the centres of massive galaxies. They have a strong influence on their bulge environments recorded in the scaling relations between the black hole mass (M_{BH}) and the galaxy’s macroscopic properties, e.g. stellar-velocity dispersion (σ_{*} ; Ferrarese & Merritt 2000; Gebhardt

et al. 2000) and the bulge-stellar mass (M_{bulge} ; Kormendy & Richstone 1995; Häring & Rix 2004; McConnell et al. 2013), despite the bulge extending well beyond the black hole’s sphere of influence (SOI; Kormendy & Ho 2013). These M_{BH} –galaxy scaling relations suggest SMBHs and galaxies grow and evolve together through a series of accretion and merger events with feedback that regulates star formation (e.g. Schawinski et al. 2007).

The current data at the low- M_{BH} regime ($< 5 \times 10^7 M_{\odot}$) hint a steeper $M_{\text{BH}}-M_{\text{bulge}}$ slope for the lower mass galaxies ($M_{*} < 10^{10} M_{\odot}$;

* E-mail: dieu.nguyenduc@phenikaa-uni.edu.vn

Nguyen et al. 2018, 2019; Nguyen et al. 2020) despite a large scatter. One possible explanation is that black hole growth follows a different evolutionary track of the growth of bimodality-black hole seeds (e.g. Pacucci et al. 2017, 2018). However, this black hole census remains incomplete, and typical scatter is about more than two orders of magnitude (e.g. Nguyen et al. 2019). Additionally, another important but often unexplored point is the systematics in different methods that may bias M_{BH} determinations from masers (e.g. Greene et al. 2010; Kuo et al. 2011), stellar- and gas-dynamics (e.g. Krajnović et al. 2018; Thater et al. 2019), reverberation mapping (RM, e.g. Peterson et al. 2004), velocity widths of broad optical emission lines (e.g. Baldassare et al. 2015). Often the different methods also do not give consistent results. Recently, there was a few direct comparisons but crucial to determine such bias, including (1) M_{BH} in M87 inferred from stellar- (Gebhardt et al. 2011) versus ionized-gas kinematics (Walsh et al. 2013) using the observations from the Event Horizon Telescope (Event Horizon Telescope Collaboration 2019a, b, c) and (2) M_{BH} in NGC 404 inferred from stellar- versus molecular-gas kinematics (Davis et al. 2020). Any confident interpretation of low-mass black hole growth will require both larger numbers of M_{BH} measurements and greater measurement precision.

The development of the cold-gas-dynamical method relies on high-spatial-resolution interferometric data at mm/(sub)mm wavelengths observed with Atacama Large Millimeter/submillimeter Array (ALMA), which has introduced a new powerful method to determine central M_{BH} in various types of galaxy morphologies and masses (Davis 2014; Onishi et al. 2015). Most of these works utilized $^{12}\text{CO}(2-1)$, which is bright and the best trade-off between sensitivity and resolution. Using the circumnuclear molecular gas disc's (CND, a massive gaseous disc with a few 100 pc in size) kinematics at the observational scale \approx the black hole's SOI reduces the error associated with the galaxy-stellar-mass uncertainty (Boizelle et al. 2019, 2021; North et al. 2019). Recent efforts of Davis et al. (2020) and Nguyen et al. submitted have even pushed the possible of the method down to the lower M_{BH} regime of $\approx 10^{5-6} M_{\odot}$.

In this paper, we applied the cold-gas-dynamical method to estimate the central M_{BH} in the Seyfert 1 active galactic nucleus (AGN) NGC 7469 using ALMA observations. NGC 7469 is one of the brightest type-1 AGNs, and therefore, its M_{BH} was measured by various methods. However, the results range largely between 10^5 and $10^8 M_{\odot}$. The M_{BH} of NGC 7469 was determined via the direct signatures of radio (VLA 8.4 GHz; Pérez-Torres et al. 2009) and X-ray (*XMM-Newton* and *Swift*; Liu et al. 2014) radiation, providing a $M_{\text{BH}} = 1.6_{-1.5}^{+24.7} \times 10^7$ and $M_{\text{BH}} = 6_{-3}^{+27} \times 10^5 M_{\odot}$ using the Fundamental Plane of Plotkin et al. (2012) and Gültekin et al. (2019), respectively. Whereas, the RM method used the indirect signature from the broad-line region (BLR) variability using H β and He II $\lambda 4686$ emissions (Peterson et al. 2004, 2014; Wang et al. 2014), giving a $M_{\text{BH}} = (1.0 \pm 0.1) \times 10^7 M_{\odot}$. Our simple but powerful method that uses cold gas dynamics will provide another direct constraint to its central black hole mass. Accurate M_{BH} estimate in AGNs, in turn, will shed light on the geometry/orientation of the ionizing photon-emitted BLR that is close to the central SMBH, which is unresolved with the current facilities but critical for the unified AGN model.

As a follow-up to the ALMA emission-lines survey of the AGNs in NGC 7469 (Izumi et al. 2020, hereafter I20), we utilized the bright, high signal-to-noise (S/N) and high-spatial-resolution data of the molecular- $^{12}\text{CO}(1-0)$ ($\theta_{\text{beam}} \approx 0''.29$) and the atomic-[CI](1-0) lines ($\theta_{\text{beam}} \approx 0''.31$) to perform cold-gas-dynamical models and weigh the central SMBH. We emphasize that the atomic-[CI](1-0) line is a potentially powerful tool to probe the nuclear gas dynamics

in AGNs. It is very bright at the CND-scale due to X-ray Dominated Region (XDR) processes around the AGNs (i.e. a mass accreting SMBH) that include a highly efficient dissociation of CO molecules into C atoms (e.g. Maloney, Hollenbach & Tielens 1996; Meijerink & Spaans 2005). This process is recently confirmed in NGC 7469 as an elevated C/CO flux ratio and/or abundance ratio (I20). Hence, [CI](1-0) can potentially be a better probe of nuclear gas dynamics than $^{12}\text{CO}(2-1)$ around AGNs and in determining M_{BH} . The atomic-[CI](1-0) (rest frequency of ≈ 492 GHz) line has a higher rest-frame frequency than $^{12}\text{CO}(2-1)$ (rest frequency of ≈ 230 GHz) or $^{12}\text{CO}(1-0)$ (rest frequency of ≈ 115 GHz) and thus can be a suitable tracer to perform similar observations and dynamical modellings towards high-redshift AGNs.

The paper is organized into six Sections. We present all crucial properties of NGC 7469 in Section 2 and *HST* and ALMA observations in Section 3. The mass modelling of NGC 7469 and the kinematics molecular simulation (KinMS; Davis 2014) tool, used to estimate the central M_{BH} , and the results are described in Section 4. We then discuss the atomic-[CI](1-0) emission is probably a new gas-tracer (also better than the frequent use of low-J CO lines) to estimate M_{BH} in the AGNs and compare this new M_{BH} with its RM-based estimates, as well as the usage this new M_{BH} to constrain the inclination angle of the unresolved BLR in Section 5. Finally, we conclude our findings in Section 6.

Throughout the article, all the maps are plotted with the orientation of north up and east to the left. We adopt an angular-size distance to NGC 7469 of $D_A = 68.4 \pm 18.8$ Mpc, where the error is 1σ scatter among 17 distances from NASA/IPAC Extragalactic Galaxies (NED¹), giving a physical scale of $330 \text{ pc arcsec}^{-1}$ in the standard flat Universe with $H_0 \approx 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M \approx 0.3$, and $\Omega_\Lambda \approx 0.7$ (corresponding to a luminosity distance of $D_L = 70.8$ Mpc, I20). The derived M_{BH} scales linearly with the assumed distance, so any change to the distance will result in a compensated shift in M_{BH} . We should emphasize that this is the lowest spatial-resolution limit at which we still can perform an accurate- M_{BH} measurement using of the atomic-[CI](1-0) emission observed with ALMA. All quantities are quoted with the Galactic extinction correction to recover their intrinsic values assuming $A_V = 0.184$ (Schlafly & Finkbeiner 2011) and the interstellar extinction law of Cardelli, Clayton & Mathis (1989).

2 NGC 7469

NGC 7469 (Mrk 1514, Arp 298) is identified as a stellar-bar spiral, with a Hubble type (R')SAB(rs)a at near-infrared (NIR) wavelength (Knapen, Shlosman & Peletier 2000). It is also classified as a luminous infrared galaxy (LIRG) based on its high infrared (IR) luminosity ($L_{8-1000 \mu\text{m}} = 10^{11.7} L_{\odot}$; Sanders et al. 2003).

Analysis from the molecular $^{12}\text{CO}(1-0)$ and HCN(1-0) emissions observed with IRAM Plateau de Bure Interferometer (PdBI) interferometer (Guilloteau et al. 1992) yields a systemic velocity $V_{\text{LSR}} = 4925 \text{ km s}^{-1}$, kinematic inclination $i = 45^\circ$, and position angle PA = 128° (Davies, Tacconi & Genzel 2004, henceforth D04).

Optical spectroscopy confirmed the nucleus of NGC 7469 hosts a luminous Seyfert 1 AGN traced by (i) strong emission at K band $2.2 \mu\text{m}$ (Genzel et al. 1995; Lonsdale et al. 2003; Imanishi & Wada 2004), (ii) a core jet-like structure (e.g. Lonsdale et al. 2003; Alberdi et al. 2006) and ionized gas outflows (Scott et al. 2005; Blustin et al. 2007), and (iii) UV and X-ray variability (e.g. Kriss et al. 2000; Nandra et al.

¹<https://ned.ipac.caltech.edu/>

2000; Petrucci et al. 2004; Scott et al. 2005). This variability is also seen in optical broad Balmer lines ($\text{FWHM}_{\text{H}\beta\lambda 4861} = 4369 \text{ km s}^{-1}$; Bonatto & Pastoriza 1990; Collier et al. 1998; Peterson et al. 2014). Davies et al. (2007) find a young stellar population ($\approx 110\text{--}190 \text{ Myr}$) in the NGC 7469's CNL. The average star formation rate there is also high ($\langle \text{SFR} \rangle = 50\text{--}100 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$), indicating a composite core of a type 1 AGN and a starburst ring distributed in the annulus of $1''.5\text{--}2''.5$ ($\approx 500\text{--}833 \text{ pc}$) at the central kpc region of this galaxy (Soifer et al. 2003; Díaz-Santos et al. 2007). Genzel et al. (1995) argue that this nucleus region accounts for two-thirds of the galaxy bolometric luminosity ($L_{\text{bol.}} = 10^{45.3} \text{ erg s}^{-1}$; Kaspi et al. 2000). This starburst ring was also clearly detected by ALMA high resolution ($< 0''.6$) (sub)mm dust continuum data (Izumi et al. 2015; Imanishi, Nakanishi & Izumi 2016).

Liu et al. (2014) investigated the nuclear X-ray spectra observed by *XMM-Newton* and *Swift* of the NGC 7469 AGN, which originates from inverse Compton scattering excited by hot and compact corona near the SMBH. They measured hard/soft X-ray luminosities of $\log L_{2\text{--}10 \text{ keV}} = 43.170 \pm 0.009 \text{ erg s}^{-1}$ and $\log L_{14\text{--}195 \text{ keV}} = 43.602^{+0.315}_{-1.184} \text{ erg s}^{-1}$. Additionally, Pérez-Torres et al. (2009) use VLA 8.4-GHz observations at $\approx 0''.3$ resolution to estimate the radio emission of the nucleus of $\log L_{8.4 \text{ GHz}} = 36.959 \pm 0.009 \text{ erg s}^{-1}$. The radiations suggest that the AGN is shining at the Eddington ratio of ≈ 0.3 (Petrucci et al. 2004).

The nuclear $^{12}\text{CO}(1\text{--}0)$ gas forms a CNL at the centre and a ring-like morphology located at $\approx 1''.5\text{--}2''.5$. Additionally, there is a bar or a pair of spiral arms between the ring and the CNL with weak NIR emission (D04; Izumi et al. 2015). The total molecular gas mass inferred from $^{12}\text{CO}(1\text{--}0)$ within the radius of $r \lesssim 3''.5$ ($\approx 1.2 \text{ kpc}$) is $2.7 \times 10^9 \text{ M}_{\odot}$ (D04, I20), while this mass of the entire galaxy is $M_{\text{H}_2} \approx 10^{10} \text{ M}_{\odot}$ (Meixner et al. 1990).

NGC 7469 has a bulge mass of $M_{\text{bulge}} = (1.1 \pm 0.3) \times 10^{11} \text{ M}_{\odot}$ (and $M_{\text{B}} = -20.9 \pm 0.2 \text{ mag}$) based on the black hole-to-bulge mass relation in AGN (Wandel 2002), while its bulge–disc luminosity decomposition from *HST/R*-band image gives $M_{\text{R}} = -22.08 \pm 0.75 \text{ mag}$ (McLure, Dunlop & Kukula 2000; McLure & Dunlop 2001). Also, Onken et al. (2004) used slit-spectroscopy to estimate its bulge/spheroid velocity dispersion of $\sigma_{*} = 152 \pm 16 \text{ km s}^{-1}$ using the stellar absorption lines of Ca II triplet (CaT) in the NIR regime excited by the AGNs.

We summarized these properties of NGC 7469 in Table 1.

3 DATA

3.1 Hubble Space Telescope (HST) images

We used the optical *HST* observations in the WFC3/UVIS-FIX *F547M* and ACS/WFC *F814W* filters to create a mass-follows-light model (Section 4.2). The *F814W* image suffers from central saturation due to the bright AGN, which will be masked out in the subsequent analysis. Also, there is a 20 per cent difference in the pixel sizes between the two instruments, WFC3 versus ACS. We thus downloaded the raw `flt` frames of the ACS *F814W* image from *HST*/Mikulski Archive for Space Telescopes (MAST) and re-reduced these images using the `drizzlepac`/`Astrodizzle` package² (Avila et al. 2012) to a final pixel scale of $0''.04$. More details of these images are shown in Table 2.

Table 1. Properties of NGC 7469.

Parameter (Unit) (1)	Value (2)	References (3)
Morphology	(R')SAB(rs)a	(1)
Nuclear activity	Seyfert 1	(2)
R.A. (ICRS)	23 ^h 03 ^m 15 ^s .617	(3)
Dec. (ICRS)	+08°52'26".00	(3)
Position angle (°)	128 (or 308*)	(4)
Inclination angle (°)	45	(4)
Angular-size distance (Mpc)	68.4	(3, 5, 6, 7)
Luminosity distance (Mpc)	70.8	(7)
Comoving radial distance (Mpc)	69.6	(7)
Redshift	0.0163	(7)
Linear scale (pc arcsec ⁻¹)	330	(7)
$\log L_{2\text{--}10 \text{ keV}}$ (erg s ⁻¹)	43.17	(8)
$\log L_{14\text{--}195 \text{ keV}}$ (erg s ⁻¹)	43.60	(8)
$\log L_{8.4 \text{ GHz}}$ (erg s ⁻¹)	36.96	(9)
$\log L_{8\text{--}1000 \mu\text{m}}$ (erg s ⁻¹)	44.58	(10)
$\log L_{\text{bol.}}$ (erg s ⁻¹)	45.30	(11)
Bulge stellar mass (M_{\odot})	$(1.1 \pm 0.3) \times 10^{11}$	(12)
Total gas mass (M_{\odot})	1.0×10^{10}	(6)
$M_{\text{CND, total}}$ (M_{\odot})	2.7×10^9	(4, 5)
M_{B} (mag)	-20.9 ± 0.2	(12)
M_{R} (mag)	-22.08 ± 0.75	(13, 14)
Velocity dispersion (km s ⁻¹)	152 ± 16	(15)
$\langle \text{SFR} \rangle$ ($M_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$)	50–100	(4)
Stellar age (Myr)	110–190	(4)
RM-based M_{BH} (M_{\odot})	$(1.0 \pm 0.1) \times 10^7$	(16, 17, 18)
[CI](1–0)-based M_{BH} (M_{\odot})	$1.8^{+2.7}_{-1.1} \times 10^7$	(19)

Note. (1): Knapen et al. (2000); (2): Osterbrock & Martel (1993); (3): I20; (4): D04; (5): Izumi et al. (2015); (6): Meixner et al. (1990); (7) <http://www.astro.ucla.edu/~wright/CosmoCalc.html>; (8): Liu et al. (2014); (9): Pérez-Torres et al. (2009); (10): Sanders et al. (2003); (11): Kaspi et al. (2000); (12): Wandel (2002); (13): McLure et al. (2000); (14): McLure & Dunlop (2001); (15): Onken et al. (2004); (16): Peterson et al. (2014); (17): Wang et al. (2014); (18): Peterson et al. (2004); (19): this paper. *: The difference in 180° of the PA will flip the CNL's velocity-position diagram but does not change the dynamical results.

We aligned these *HST* images to the galaxy centre determined from I20. Next, we used the `Tiny Tim`³ routine (Krist 1995; Krist, Hook & Stoehr 2011) to create point spread function (PSF) models for individual *HST* exposure frames of the involved filters. Then, we inserted them into the corresponding `flt` images at the galaxy centre in each frame to simulate the observations. The final PSF model is the combination of these PSF models created by `Astrodizzle` (Rusli et al. 2013; den Brok et al. 2015; Nguyen et al. 2017, 2018; Thater et al. 2019), which has an FWHM of $0''.08$ ($\approx 27 \text{ pc}$) and will be used to decompose the mass model (Section 4.2).

Fig. 1 shows the *F547M* image. There is a starburst ring with bright and resolved super-star clusters distributed around the bright nucleus within the annulus of $1''.5\text{--}2''.5$ ($\approx 500\text{--}833 \text{ pc}$), as well as prominent dust lanes on the north and north-east sides, extending to a radius of at least 10 arcsec ($\approx 3.3 \text{ kpc}$) from the centre.

3.2 ALMA observations

Izumi et al. (2020) reported a variety of bright gas emissions in the nucleus of NGC 7469, including molecules (e.g. $^{12}\text{CO}(1\text{--}0)$, $^{12}\text{CO}(2\text{--}1)$, $^{12}\text{CO}(3\text{--}2)$) and atomic [CI](1–0). An

²<https://www.stsci.edu/scientific-community/software/drizzlepac>

³<http://tinytim.stsci.edu/sourcecode.php>.

Table 2. *HST*/WFC3 UVIS-FIX and ACS WFC images.

Filter	Camera	Aperture	UT date	PID	PI	Pixel scale (" /pix)	Exposure time (s)	Zero-point ^a (Vega mag)	A_{λ}^b (mag)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
<i>F547M</i>	WFC3	UVIS2-FIX	2009 November 11	GO-11661	Bentz	0.04	3×370	24.748	0.075
<i>F814W</i> (saturation)	ACS	WFC	2006 June 12	GO-10592	Evans	0.05	2×200	25.943	0.105




Figure 1. The *HST*/WFC3 *F547M* image of NGC 7649 within the FOV of 50 arcsec \times 50 arcsec ($\approx 16.5 \times 16.5$ kpc²) overlaid with the contours of the integrated intensities of molecular-¹²CO(1-0) (panel a) and atomic-[CII](1-0) (panel b) emissions, respectively. These maps show the large-scale image and the co-spatial distributions of the dust lanes seen in optical and the molecular/atomic gas seen in mm/sub-mm wavelengths. Here, emission contours are spaced at $0.1 \times n \times I_{0,\max}$ of ¹²CO(1-0) and [CII](1-0) integrated intensities shown in the colour bars of the panel (a) of Figs 2 and 3, $n = 1, 9$. In panels (c) and (d), we zoom to the FOV of 15 arcsec \times 15 arcsec ($\approx 5 \times 5$ kpc²) for ¹²CO(1-0) and 6 arcsec \times 6 arcsec ($\approx 2 \times 2$ kpc²) for [CII](1-0) to show the detailed distributions of the gas emissions at the centre of NGC 7469.

Table 3. ALMA observations of the molecular- $^{12}\text{CO}(1-0)$ and atomic-[CI](1-0) emission lines.

Band	Emission	Frequency (GHz)	BMAJ (arcsec/pc)	BMIN (arcsec/pc)	BPA ($^\circ$)	RMS (mJy beam $^{-1}$ km s $^{-1}$)	V_{sys} (km s $^{-1}$)	Velocity width (km s $^{-1}$)	LAS (arcsec)	Velocity res. (km s $^{-1}$)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)
3	$^{12}\text{CO}(1-0)$	115.271	0.37/123	0.27/90	-72.5	34	4850	400	3.05	5.2
8	[CI](1-0)	492.161	0.34/113	0.31/103	75.7	227	4831	390	4.53	2.5

Note. (1)–(3): ALMA observed bands, emission lines, and their rest frequencies. (4)–(6): The synthesized beam sizes (both in arcsec and parsec) and orientations. BMAJ, BMIN, and BPA are the major axis, minor axis, and position angle of the beam, respectively. (7): The root-mean-squares (rms) noise. (8)–(10): the systemic velocity, total velocity width of the whole CNB and the largest angular size of each emission line. (11): The velocity resolution of the emissions.

accurate weighing of the central SMBH requires that these emissions are observed at high spatial resolution, and show bright, compact distributions towards the galaxy centre and regular kinematics in the entire CNB. Owing to the XDR effect (i.e. dissociation of CO into C), [CI](1-0) surely probes the close vicinity of this AGN as evidenced by the position-velocity diagram (PVD). We thus can extend this analysis towards higher redshift thanks to the higher rest frequency of [CI], which is much easier than the CO lines observed here. On the other hand, an extended CNB traces the total enclosed mass out to the radius covered by the gas, which helps to separate the M_{BH} from the masses of the remaining galaxy components (i.e. stars + ISM + non-luminous stellar remnants) better. Regarding this aspect, the ≈ 5 per cent higher in spatial resolution and more extended of $^{12}\text{CO}(1-0)$ make it a better transition for dynamical modelling than other CO lines. We thus chose the atomic-[CI](1-0) and molecular- $^{12}\text{CO}(1-0)$ emissions among these as the best bets to perform M_{BH} measurement that compromises both these requirements.

The nuclear molecular- $^{12}\text{CO}(1-0)$ and atomic-[CI](1-0) emissions of NGC 7469 were observed in bands 3 and 8 of ALMA cycle 5 (PID: 2017.1.00078.S, PI: T. Izumi) at the spatial-resolution scales of $\approx 0''.3$ (or ≈ 100 pc) that are ≈ 55 times larger than the SOI of the black holes predicted from RM ($\approx 0''.006$ or ≈ 1.9 pc; Peterson et al. 2004, 2014; Wang et al. 2014). These bright and high-quality observations reveal that their kinematics are suitable for gas-dynamical modelling to measure the central M_{BH} . Details of data reduction and imaging of these visibilities are presented in I20. In Table 3, we show the observational properties of these two emissions.

We created a 3D cubes of (RA, Dec., and velocity) for these $^{12}\text{CO}(1-0)$ and [CI](1-0) emissions using the Common Astronomy Software Applications (CASA; McMullin et al. 2007, package version 5.4). The $^{12}\text{CO}(1-0)$ and [CI](1-0) integrated intensities are shown in contours in Fig. 1, overlaid on the $F547M$ image. It is clear that the atomic-[CI](1-0) emission is more compact and strongly peaked towards the centre (< 5 arcsec or 1.7 kpc) than the molecular- $^{12}\text{CO}(1-0)$ emission when we solely look at the central kpc regions (i.e. inside the starburst ring). However, the molecular- $^{12}\text{CO}(1-0)$ emission shows a more extended distribution (≈ 7 arcsec or 2.3 kpc) with some fainter features distributed further out at ≈ 15 –20 arcsec (or ≈ 5 –6.6 kpc).

We also created the integrated intensity (moment 0), intensity-weighted mean LOS velocity field (moment 1), and intensity-weighted LOS velocity dispersion (moment 2) maps for $^{12}\text{CO}(1-0)$ and [CI](1-0) using the moments masking technique (Dame, Hartmann & Thaddeus 2001; Dame 2011) and showed them in the panels (a), (b), and (c) of Figs 2 and 3, respectively. To do this, we spatially smoothed each channel by a factor of α then varied α and gauged the spatial and velocity coherence of the signal. The spatial smoothing increases the sensitivity but decreases the angular resolution, helping to suppress noise peaks. Next, we performed ‘ $\beta\sigma$ -clipping’, where β is a positive factor and σ is the RMS noise

(recorded in column 7 of Table 3). Thus, we decided all channels with intensities $< \beta\sigma$ were set to zero at any given spatial position. This mask created from the smoothed cube was only used to identify and mask out emission-free regions of the original cube, while the moment maps with full spatial and velocity resolutions were created from the original cube still. We varied α and β for their appropriate choices to obtain the best moment maps, which yields $\alpha = 3$ and $\beta = 0.5$.

The gas have ring-like structures with diameters of ≈ 7 arcsec for $^{12}\text{CO}(1-0)$ and < 5 arcsec for [CI](1-0), indicating by features of two-arm/bi-symmetric spiral structures (showing as the two cyan arcs in the panel a of Figs 2 and 3). In the molecular- $^{12}\text{CO}(1-0)$ gas integrated intensity map [panel (a) of Fig. 2], the emission is resolved with the three high-surface-brightness regions within $0''.5$ and other high-surface-brightness areas associated with the star-forming ring (black circle). Whereas, the atomic-[CI](1-0) gas emission is centrally peaked within central $0''.5$, and the two-arm spiral structure of the high-surface-density regions opens a bit narrower than that of $^{12}\text{CO}(1-0)$ (panel a of Fig. 3).

The intensity-weighted mean LOS velocity fields, which are beam-convolved, of both emission lines reveal a rotating disc with a total velocity width (ΔV) of ≈ 400 km s $^{-1}$ for $^{12}\text{CO}(1-0)$ and ≈ 390 km s $^{-1}$ for [CI](1-0) and small warps seen along the minor axis (panel b of Figs 2 and 3). These warps would make our dynamical modellings become complicated and hard to measure the M_{BH} . Panel (c) of Figs 2 and 3 shows the intensity-weighted mean LOS velocity dispersion (σ_{gas}) maps of these emissions. The σ_{gas} shows spaxel variations within the ranges of $\approx (3\text{--}51)$ km s $^{-1}$ for $^{12}\text{CO}(1-0)$ and $\approx (3\text{--}67)$ km s $^{-1}$ for [CI](1-0). Generally, the high intensity-weighted mean LOS velocity dispersion regions are co-spatial with the high-surface-brightness density regions, i.e. the centrally peaked region, star-forming ring, and regions associated with the two-arm spiral structure. It is notable that at large radii ($r \gtrsim 2$ arcsec or 660 pc), these high-velocity dispersion regions are highly dominant in the west and north-west sides of the $^{12}\text{CO}(1-0)$ map, while in the [CI](1-0) map they are mostly located in the south and south-east sides. The origin of these inverse distributions of the intensity-weighted mean LOS velocity dispersions between these two emission lines is unknown. However, they do not affect the dynamical modelling of M_{BH} .

In panel (d) of Figs 2 and 3, we show the integrated spectrum of the $^{12}\text{CO}(1-0)$ and [CI](1-0) CNBs created by extracting the fluxes within the nuclear regions of 10 arcsec \times 10 arcsec ($\approx 3.3 \times 3.3$ kpc 2) and $7''.2 \times 7''.2$ ($\approx 2.4 \times 2.4$ kpc 2) of the corresponding datacubes. These profiles show a classical double-horn shape of a rotating disc but asymmetric with some missing blueshifted velocity channels.

Similarly, Fig. 4 shows the PVDs of $^{12}\text{CO}(1-0)$ and [CI](1-0) extracted from a cut of three pixels in width ($0''.81 \approx 270$ pc) through the major axis (PA = 308°). There is sharp increase of the rotations towards the galaxy centre in both emissions, interpreted as the Keplerian motions (i.e. $V_{\text{circ}}(r) \propto r^{-0.5}$) caused by either an




Figure 2. The moment maps of $^{12}\text{CO}(1-0)$ show in the order of the integrated intensity (panel a), the intensity-weighted mean LOS velocity field (panel b), and the intensity-weighted LOS velocity dispersion (panel c) within the FOV of $10 \text{ arcsec} \times 10 \text{ arcsec}$ ($\approx 3.3 \times 3.3 \text{ kpc}^2$). In the moment 0 map, the black circle shows the position of the starburst ring (Soifer et al. 2003; Díaz-Santos et al. 2007), while the cyan arcs highlight the two-arm spiral structure of the CND. In the moment 1 map, there is a small warp elongated along the minor axis, creating a twist seen in this direction. In the moment 2 map, the high-intensity-weighted LOS velocity dispersion regions are co-spatial with the high-surface-brightness density regions in the moment 0 map within the region of 2 arcsec ($\approx 660 \text{ pc}$). Further west of this region where the gas distributes regularly in the CND, the LOS velocity dispersion is also high and peaks at the region where the gas is faint (mom 0) and the LOS velocity field drops suddenly in the redshifted side (mom 1) due to low S/N. The synthesized beam size listed in Table 3 is shown as an ellipse at the bottom left of each panel. The $(\Delta\text{RA}, \Delta\text{Dec.}) = (0, 0)$ position on these maps indicates the kinematic/galaxy centre. In panel (d), we show the integrated spectrum extracted within the same FOV, showing an asymmetric double-horn shape of a rotating disc.

SMBH or a centrally massive and compact cluster of baryonic matter residing at the heart of NGC 7469. The asymmetry in the CND is only seen in the blueshifted Keplerian motion of the $^{12}\text{CO}(1-0)$ PVD with $\approx 20 \text{ km s}^{-1}$ less than the redshifted Keplerian motion. This asymmetric emission distribution in the PVD may be caused by a smaller fraction of molecular gas mass on the blueshifted component

than that of the redshifted component. Other possibility is that they have different excitation condition (Imanishi et al. 2018; Izumi et al. 2018). Here, we rule out the possibility of dust extinction because there is less extinction in this blueshifted component (Section 4.4). However, this rotating-asymmetric feature is not present in the $[\text{C}]\text{I}(1-0)$ PVD, suggesting this emission line is actually a more




Figure 3. Same as Fig. 2 but for [CII](1–0) in a smaller FOV of $7.2 \text{ arcsec} \times 7.2 \text{ arcsec}$ (or $2.4 \times 2.4 \text{ kpc}^2$), but some differences are described here. In the integrated intensity (panel a) map, the black circle and blue spiral arms illustrate the same region and the arcs of the two-arm spiral structure of the CND as similar as those in the panel (a) of Fig. 2, although we show them in a smaller FOV of the map (i.e. their illustrations are bigger in sizes). In the moment 2 map (panel c), there is an opposite distribution of the atomic-[CII](1–0) gas compared to the molecular- $^{12}\text{CO}(1–0)$ gas in Fig. 2. The high-velocity-dispersion regions are mostly located in the south and south-east sides of the nucleus not in the north and north-west. The reason for these inverse velocity-dispersion distributions is currently unknown. Future works of both atomic and molecular gas surveys in a sample of AGN will disentangle this issue better.

appropriate line to measure the M_{BH} than the molecular- $^{12}\text{CO}(1–0)$ line for this AGN NGC 7469. This argument is strengthened when we examine these emissions in the galaxy-minor-axis ($\text{PA} = 218^\circ$) PVDs as shown in Fig. 5. The [CII](1–0) emission is symmetric, while the $^{12}\text{CO}(1–0)$ emission is asymmetric with missing the blueshifted motions towards the galaxy centre. On the other hand, the latter is more extended towards larger radii than the former. These features again tell us that [CII](1–0) is a best bet to estimate the centrally

compact mass (i.e. black hole mass), while the $^{12}\text{CO}(1–0)$ line is preferable in tracing the enclosed mass (i.e. mass-to-light ratio, M/L) of NGC 7469 out to $\approx 7 \text{ arcsec}$ [see panel a of Fig. 1; also see the usage of $^{12}\text{CO}(2–1)$ in Izumi et al. (2020)]. Importantly, at the spatial scales measured in this work, there are somewhat insignificant non-circular motions (i.e. outflows or inflows $\approx 20 \pm 10$ (kinematic uncertainty) km s^{-1} ; will be discussed later in Section 4.3) are found in both lines.




Figure 4. The PVDs extract along the major axis (PA = 308°) in the FOVs of 10 arcsec \times 10 arcsec ($\approx 3.3 \times 3.3$ kpc²) for ¹²CO(1–0) (panel a) and 7.2 arcsec \times 7.2 arcsec ($\approx 2.4 \times 2.4$ kpc²) for [CI](1–0) (panel b) with a slit of three pixels in width (0 $'$.81 or 270 pc). Keplerian motions towards the galaxy centre present in both diagrams. However, that motion of [CI](1–0) is more symmetric than ¹²CO(1–0). Small pluses at the bottom left corners show the errors of our measurements. Contour levels are plotted at $n \times 0.1 \times I_{0,\max}$ of ¹²CO(1–0) and [CI](1–0) shown in the colour bars (also Figs 2 and 3), $n = 1, 9$. It is clear that the atomic-[CI](1–0) emission is much more centrally concentrated than the molecular-¹²CO(1–0) emission.




Figure 5. Same as Fig. 4 for the PVDs extract along the minor axis (PA = 218°). There are no circular motions in both emissions. However, the PVD of [CI](1–0) (red) is more symmetric than the PVD of ¹²CO(1–0) (blue). The horizontal lines with corresponding colours indicate the systemic velocities of [CI](1–0) and ¹²CO(1–0), while the vertical black line anchors the emission centre.

4 DYNAMICAL MODELS

In this section, we present the KinMS tool (Section 4.1; Davis 2014) and the *default-mass model* (Section 4.2) of NGC 7469 used to estimate the central M_{BH} . Next, we report the results in Section 4.3. Alternatively, we test our dynamical modellings with various galaxy mass models based on different assumptions of colour variability and colour– M/L correlations (Nguyen et al. 2019; Bell & de Jong 2001; hereafter N19; B01) in Section 4.4. A simpler version of this

analysis is presented in I20 as well to dynamical measure the CO-to- or [CI]-to- H_2 mass conversion factors.

4.1 KinMS model

The KinMS tool optimizes a datacube in two steps. First, it creates a simulated cube with a given set of model parameters for comparison to observables. Secondly, it determines the best-fitting model by using an efficient method to explore the parameter space.

To create a simulated cube, KinMS adopts a parametric function describing the distribution and kinematics of molecular gas. The gas is assumed to move on circular orbits governed by a circular velocity curve, calculated from the `mge_circular_velocity` procedure within the Interactive Data Language (IDL) Jeans Anisotropic Modelling (JAM⁴; Cappellari 2008) package. This circular velocity curve is then used as the corresponding input of the axisymmetric-mass model specified via the multiGaussian expansion (MGE; Emsellem, Monnet & Bacon 1994; Cappellari 2002) parametrization, including all mass components: stars, the interstellar medium (ISM; gas + dust), and a putative SMBH (Davis et al. 2013).

The KinMS tool then uses this simulated cube for optimizing the data cube. Specifically, it utilizes the EMCEE technique (Foreman-Mackey et al. 2013), the affine-invariant ensemble sampler based on the Markov Chain Monte Carlo (MCMC) method, and the Bayesian analysis framework to walk through the parameter space (Goodman & Weare 2010). The model calculates χ^2 at every step and uses it to determine the next move until reaching the minimum- χ^2 . In practice, this is all accomplished by using the PYTHON code `KINMSpy_MCMC`.⁵ To find the best-fitting parameters and errors, we determine the

⁴<https://purl.org/cappellari/software>

⁵https://github.com/TimothyADavis/KinMSpy_MCMC

probability distribution function (PDF) likelihood calculated from the full calculation pool for the posterior distribution.

Dynamical pressure distributed by the highly turbulent σ_{gas} of a thick disc could potentially cause bias in the M_{BH} because the gas pressure supports the gas against gravity (Binney & Tremaine 1987). Observations with a high ratio of $\sigma_{\text{gas}}/V_{\text{rot}}$ will spoil the thin disc assumption of gas moving on purely circular orbits (Coccatto et al. 2006; Barth et al. 2016a; Boizelle et al. 2019). Using the dynamics of [CI](1–0) and $^{12}\text{CO}(2-1)$, I20 found $\sigma_{\text{gas}}/V_{\text{rot}} \approx 0.35$ ($r \lesssim 0''.5$) and $\sigma_{\text{gas}}/V_{\text{rot}} \approx 0.19$ ($r > 0''.5$) [see panels (a) and (b) of their Fig. 6], which indicates a dynamically thin ($d_t \approx 0$) and cold ($V_{\text{rot}} \approx V_{\text{circ}}$) CND for NGC 7469. That means the dynamically asymmetric drift from turbulent velocity dispersion of the gas can be ignored (Davis et al. 2020, Nguyen et al., submitted).

The KinMS tool also uses a radial parametric function to describe gas surface brightness distribution (SB). Here, we assumed the axisymmetric SBs for [CI](1–0) and $^{12}\text{CO}(1-0)$ and modelled them using a Gaussian. We identified the Gaussian centre to be the kinematic centre (x_c, y_c) and incorporated their normalization factors into the total flux parameters (F). Thus, we described the $^{12}\text{CO}(1-0)$ and [CI](1–0) SBs with only one free parameter: the dispersion of the Gaussian (G_σ).

Additionally, to account for the small warps seen in both intensity-weighted mean LOS velocity fields of $^{12}\text{CO}(1-0)$ and [CI](1–0) [panels (c) of Figs 2 and 3], we extracted their radial PA profiles along the major axis via the `kinometry`⁶ code (Krajinović et al. 2006), then used them as an additional input in the same manner of using the circular velocity curve. These PA profiles vary in the range of (295–325)° for $^{12}\text{CO}(1-0)$ and (300–320)° for [CI](1–0) across 4 arcsec as shown in Fig. 7. Accounting for PA variations in our dynamical models help to mimic the warps seen along the galaxy-minor axis of the velocity-field maps of $^{12}\text{CO}(1-0)$ [panel (b) of Fig. 2] and [CI](1–0) [panel b of Fig. 3]. Dynamical models with constant PAs will create axisymmetric kinematic maps that cannot well reproduce the data, which are asymmetric along the minor axis. This mismatch between the data and the model biases the M_{BH} estimates and produces some large residuals or artificial non-circular motions in the `Data-Model` maps of the velocity fields.

Overall, the KinMS model matches the observations with nine free parameters, including G_σ , i , F , x_c , y_c , M_{BH} , M/L , σ_{gas} , and a velocity offset ($v_{\text{off}} = V - V_{\text{sys}} - V_{\text{rot}}$) of the kinematic centre relative to the rotation and the systemic velocity of the whole galaxy.

4.2 The default-galaxy-mass model

4.2.1 Stellar mass

We applied the IDL `mge_fit_sectors` routine, version 4.14⁷ (Cappellari 2002) of the MGE model, to parametrize the stellar-mass distribution. We first modelled the $F547M$ PSF as circular MGE model and tabulated it in Table 4. Then, we included this PSF MGE in the second axisymmetric-MGE fit for the photometric MGE. We also masked out pixels contaminated by the AGN (unresolved point source at the centre, $r < 0''.06$ or ≈ 20 pc), the prominent starburst ring and bright stars. This MGE can be de-projected analytically with a specific axial ratio to reconstruct a 3D distribution of the entire galaxy. This best-fitting MGE model is tabulated in Table 5

and shown in Fig. 8, which illustrates the agreement between the data and the model in the form of 2D contours at the same radii and contour levels.

4.2.2 ISM mass

Izumi et al. (2020) estimated the total molecular mass of $M_{\text{H}_2} \approx 2.7 \times 10^9 M_\odot$ within the region of $r \leq 3$ arcsec ($\lesssim 1$ kpc) using the dynamics of $^{12}\text{CO}(2-1)$ and [CI](1–0). In this work, we estimated this mass using the $^{12}\text{CO}(1-0)$ flux calculated in circular apertures. We converted these aperture fluxes into M_{H_2} by assuming the CO-to- H_2 conversion factor for starburst galaxies: $X_{\text{CO}} = 1.0 \pm 0.3 \times 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$ (Kuno et al. 2000, 2007; Bolatto et al. 2013). We found the same amount of M_{H_2} within 3 arcsec and $M_{\text{H}_2}(r \lesssim 7'') \approx 4 \times 10^9 M_\odot$.

Papadopoulos & Allen (2000) also estimated the dust mass of $\approx 2.5 \times 10^7 M_\odot$ in the same region of $r \leq 7$ arcsec ($\lesssim 2.3$ kpc). These values give a gas-to-dust ratio of ≈ 160 , which is close to the Galactic value. We then accounted for these ISM mass (molecular gas + dust) in the mass model by turning on the `gasgrav` mechanism in the KinMS model. Here, we assumed the molecular gas and dust are distributed accordingly to the surface brightness densities of $^{12}\text{CO}(1-0)$ and [CI](1–0).

The stars and ISM masses directly predict the circular velocity of the $^{12}\text{CO}(1-0)$ and [CI](1–0) CNDs using a constant M/L_{F547M} for the stellar-mass component as a free parameter in our dynamical models. We define the total mass model with this stellar-mass model as the *default-mass model* and quote its results throughout the paper.

4.3 Results of the default-mass model

We performed the KinMS fit in an area of 60 pixels \times 60 pixels (6 arcsec \times 6 arcsec, $\approx 2 \times 2$ kpc²). The model started with an initial guess of parameters that are flat priors in linear scales, except for M_{BH} was in log-scale (to ensure efficient sampling of the posterior). The search ranges of these parameters are shown in column 2 of Table 6. We limit the velocity channels ($-300, 300$) km s⁻¹ related to the systemic velocity of ≈ 4850 and ≈ 4831 km s⁻¹ for $^{12}\text{CO}(1-0)$ and [CI](1–0), respectively. To ensure our fit converges, we ran the model with 3×10^5 iterations, then exclude the first 20 per cent of the calculations as the burn-in phase and used the rest 80 per cent to produce the final posterior PDF of all nine free parameters.

Previous works showed that the ALMA noise covariance of the uncertainties, meaning the contribution of systematic errors dominates over the statistical errors (Mitzkus, Cappellari & Walcher 2017). van den Bosch & van de Ven (2009) suggested that both errors give the same contribution and increased the $\Delta\chi^2$ required to define a given confidence level (CL) by the standard deviation of the χ^2 , namely $\sqrt{2(N-P)} \approx \sqrt{2N}$ (see section 15.1 of Press 2007), where $N = 60 \times 60 \times 60$ is the number of constraints and $P = 9$ is the number of model parameters or degree of freedom. Accordingly, in the Bayesian framework, we rescaled the model uncertainties by dividing the model log likelihood by $\sqrt{2N}$ or equivalently multiplying the `rms` by $(2N)^{1/4}$ (Table 3). This rescaling approach was adopted in recent papers using KinMS and ALMA data (Onishi et al. 2017; Nagai et al. 2019; North et al. 2019; Smith et al. 2019; Davis et al. 2020; Nguyen et al. 2020; Nguyen et al., submitted; Smith et al. 2021a) to yield more realistic uncertainties.

We interpreted our KinMS models' results with an SMBH at the centre of NGC 7469 to match the Keplerian motions of the $^{12}\text{CO}(1-0)$ and [CI](1–0) kinematics. The best-fitting model is identified

⁶<http://davor.krajinovic.org/idl/#kinometry>

⁷<http://purl.org/cappellari/software>




Figure 6. The $4.8 \text{ arcsec} \times 4.8 \text{ arcsec}$ ($\approx 1.7 \times 1.7 \text{ kpc}^2$) FOV of the intensity-weighted mean LOS velocity field map of $^{12}\text{CO}(1-0)$ (panel a), the map derived from the best-fitting KinMS model shown in panel (c) of Fig. 10 (panel b), and the residual (Data - Model) LOS map (panel c).




Figure 7. The radial position angle (PA) profiles of the $^{12}\text{CO}(1-0)$ (cyan) and $[\text{CI}](1-0)$ (red) emission lines extracted along the galaxy major axis using the Kinemetry code (Krajinović et al. 2006).

Table 4. MGE Parameters of the *HST*/WFC3 *F547M* PSF.

j	Total count of Gaussian $_j$	σ_j (arcsec)	q_j
(1)	(2)	(3)	(4)
1	0.18	0.02	0.99
2	0.64	0.06	1.00
3	0.07	0.18	0.99
4	0.06	0.37	0.99
5	0.02	0.92	0.98
6	0.03	1.55	0.98

Note. Total count, width (σ_j), and axial ratio (q_j) of each Gaussian j .

via the Bayesian analysis of the post-burn-in PDFs generated by MCMC. Here, the best-fitting value of each parameter is the median of the marginalized parameter posterior PDFs of all other parameters, and its uncertainty are all models within (31–69 per cent), (16–84 per cent), (2.3–97.7 per cent), and (0.14–99.86 per cent) of the PDFs or 0.5σ , 1σ , 2σ , and 3σ CLs, respectively.

Table 5. The *HST*/WFC3 *F547M* stellar-light-MGE model.

j	$\log(\Sigma_{*,j}/L_{\odot} \text{ pc}^2)$	σ_j (arcsec)	q_j
(1)	(2)	(3)	(4)
1	5.52	0.02	1.00
2	4.79	0.04	1.00
3	3.44	0.27	1.00
4	2.82	0.96	1.00
5	1.73	1.94	0.86
6	1.45	7.60	0.62
7	0.34	18.1	0.82

Note. Stellar-light-surface density ($\Sigma_{*,j}$), width (σ_j), and axial ratio (q_j) of each deconvolved Gaussian component j .

4.3.1 M_{BH} derived with the atomic- $[\text{CI}](1-0)$ kinematics

At 3σ CL, the best-fitting model gives $M_{\text{BH}} = 1.78_{-1.10}^{+2.69} \times 10^7 M_{\odot}$ and $M/L_{\text{F547M}} = 2.20_{-0.40}^{+0.43} (M_{\odot}/L_{\odot})$ at $\chi^2_{\text{min, red}} = 1.94$. Other best-fitting parameters and their likelihoods show in Table 6.

4.3.2 M_{BH} derived with the molecular- $^{12}\text{CO}(1-0)$ kinematics

Similarly, at 3σ CL, This best-fitting model has $M_{\text{BH}} = 1.60_{-1.45}^{+11.52} \times 10^7 M_{\odot}$ and $M/L_{\text{F547M}} = 2.22_{-0.22}^{+0.20} (M_{\odot}/L_{\odot})$ at $\chi^2_{\text{min, red}} = 1.91$. Also, see Table 6 for the full model's description.

To demonstrate how well the models describe the observables, we show the observed PVDs overlaid with the best-fitting PVDs in Fig. 9 for $[\text{CI}](1-0)$ and Fig. 10 for $^{12}\text{CO}(1-0)$. The models without, with lower mass, and with overly massive SMBHs also add for comparisons. All these models do not fit the CND's central Keplerian motions. The intensity-weighted mean LOS velocity field of the data, the best-fitting models, and their residual fields (*Data* - *Model*) within the FOV of $5''.2 \times 5''.2$ ($\approx 1.7 \times 1.7 \text{ kpc}^2$) show in Fig. 11 for $[\text{CI}](1-0)$ and Fig. 6 for $^{12}\text{CO}(1-0)$. Within this FOV, most of the residual map has the amplitude of $\approx 10 \text{ km s}^{-1}$, the limit in which we bin the spectrum together (the channel width or the uncertainty of our kinematic measurements). However, there is a large residual ($\approx 30 \text{ km s}^{-1}$) on the north-west and south-east sides of the galaxy centre for $[\text{CI}](1-0)$ and $^{12}\text{CO}(1-0)$, respectively, which might be interpreted as real non-circular motions. Nevertheless, these high-velocity residuals in both emissions are perhaps the results of the asymmetric rotation seen in panel (d) of Figs 2 and 3 (Section 3.2),




Figure 8. The comparison between the *HST/F547M* photometry of NGC 7469 (black) and its corresponding best-fitting MGE model (red) at the same radii, which illustrate the whole galactic scale 60 arcsec \times 60 arcsec ($\approx 20 \times 20$ kpc², panel a) and the zoom-in of 14 arcsec \times 14 arcsec ($\approx 4.6 \times 4.6$ kpc², panel b) FOV. Yellow regions are the pixel excluded during the MGE fit.

Table 6. Best-fitting model parameters and associated statistical uncertainties for the *default-mass model* and the kinematics of [CI](1–0) and ¹²CO(1–0).

Parameters (Units)	Search range (Uniform)	Best fit	1 σ error (1–84%)	3 σ error (0.14–99.86%)	Best fit	1 σ error (1–84%)	3 σ error (0.14–99.86%)
<i>Emission line</i>							
<i>Black hole:</i>							
$\log(M_{\text{BH}}/M_{\odot})$	(1, 9)	7.25	−0.14, +0.13	−0.42, +0.40	7.22	−0.30, +0.31	−0.90, +0.93
$M/L_{F547M} (M_{\odot}/L_{\odot})$	(0.1, 5.0)	2.20	−0.14, +0.13	−0.43, +0.40	2.22	−0.07, +0.06	−0.22, +0.20
<i>Gas CNDS:</i>							
$\sigma_{\text{gas}} (\text{km s}^{-1})$	(1, 50)	26.53	−0.12, +0.17	−0.36, +0.40	22.70	−1.14, +1.34	−4.00, +4.02
G_{σ} (arcsec)	(0.1, 1.5)	0.80	−0.01, +0.02	−0.03, +0.06	0.60	−0.04, +0.04	−0.12, +0.12
$F (\text{Jy km s}^{-1})$	(1, 100)	76.98	−0.41, +0.43	−1.23, +1.29	37.30	−1.54, +1.39	−4.51, +4.32
$i (^{\circ})$	(45, 89)	53.59	−0.42, +0.41	−1.26, +1.25	52.90	−1.40, +1.17	−3.20, +3.10
<i>Nuisance:</i>							
x_{c} (arcsec)	(−1.5, +1.5)	−0.02	−0.07, +0.06	−0.21, +0.20	+0.07	−0.13, +0.12	−0.38, +0.36
y_{c} (arcsec)	(−1.5, +1.5)	−0.01	−0.08, +0.07	−0.24, +0.21	−0.16	−0.14, +0.14	−0.42, +0.42
$v_{\text{off}} (\text{km s}^{-1})$	(−5, +5)	0.01	−0.07, +0.07	−0.21, +0.21	0.48	−0.11, +0.13	−0.33, +0.37

Note. The table columns list, respectively, each parameter name, search range, best fit and uncertainty at 1 σ (1–84%) and 3 σ (0.14–99.86%) CLs of the PDF. The parameters x_{c} , y_{c} , and v_{off} are parameters defined relative to the adopted galaxy centre (23^h03^m15^s.617, +08^o52′26″.00) and $V_{\text{sys}} = 4831$ and 4834 km s^{−1} for [CI](1–0) and ¹²CO(1–0), respectively.

or perhaps due to the beam smearing effects because of the limited-spatial resolutions of our data, and consequently, the black hole could be a bit more massive. Future higher resolution observations will better estimate these non-circular motions, which will deliver a more robust M_{BH} .

We show the PDF and 2D marginalizations of each free parameters of KinMS models of [CI](1–0) and ¹²CO(1–0) in Appendix, respectively. There is no covariance found among parameters of the [CI](1–0) kinematics. For ¹²CO(1–0), covariance of M_{BH} versus M/L_{F547M} is present as the result of the degeneracy between the potentials of the SMBH and galaxy itself, happening when the observational scales have not resolved the SMBH’s SOI. Also, other covariances of G_{σ} and σ_{gas} versus other parameters in the two bottom row panels of its posterior cause by no clear central peak in the ¹²CO(1–0) integrated intensity map (panel a of Fig. 2). Three bright clumps of high-brightness-density at the ¹²CO(1–0) CNDS

centre increases the explored ranges for x_{c} and y_{c} , then resulted in such covariances among the parameters during a simultaneous constraint.

As we mentioned that the atomic-[CI](1–0) line might be a better transition to do dynamical modelling for the central M_{BH} than the molecular-¹²CO(1–0) line in Section 1. *We proved here that the KinMS model using the [CI](1–0) kinematics produces a significantly smaller M_{BH} -uncertainty than that of the KinMS model using the ¹²CO(1–0) kinematics.* Also, there is no correlation between the constrained- M_{BH} and M/L_{F547M} , meaning an excellent constraint. We thus quote the parameter values inferred from the KinMS model using the kinematics measured from the atomic-[CI](1–0) line throughout this work. However, it should be noticed that although the M/L_{F547M} -uncertainty inferred from the KinMS model using the ¹²CO(1–0) kinematics is a factor of ≈ 2 smaller (in log scale) than that of the KinMS model using the [CI](1–0) kinematics, the correlation




Figure 9. Comparisons between data and a few KinMS models using the *default-mass model* and the [CI](1–0) emission. The PVD was extracted along the major axis (orange scale and grey contours) in the same manner in Fig. 4. The model PVDs are extracted identically from models that are different in M_{BH} (cyan contours), including the model without an SMBH (panel a) and the models with a small (panel b), the best fit (panel c, $M_{\text{BH}} = 1.78 \times 10^7 M_{\odot}$ and $M/L_{F547M} = 2.20 (M_{\odot}/L_{\odot})$), and an overly large (panel d) SMBH. The models in panels a, b, and d are not good fits for the data in the central part of the atomic-[CI](1–0) CND’s kinematics as they fail to produce the raising rotation of the Keplerian motion towards the galaxy centre.

between its constrained- M_{BH} and M/L_{F547M} is present. This M/L_{F547M} uncertainty mismatch is probably caused by (1) the more extended distribution of $^{12}\text{CO}(1-0)$ kinematics (≈ 7 arcsec) comparing to that of [CI](1–0) (< 5 arcsec) as seen in panels (a) and (b) of Fig. 1, meaning it is the better probe of the extended mass or M/L_{F547M} . Also, (2) the somewhat higher symmetry of the extended kinematics of $^{12}\text{CO}(1-0)$ than that of [CI](1–0) seen in their extracted PVDs along the galaxy-major axis (the lower velocity wings, Fig. 4) help to reduce its M/L_{F547M} uncertainty [see the consistencies between the data and the best-fitting models in panels (c) of Figs 9 and 10].

Boizelle et al. (2019) discussed the angular-resolution limit at which one can constrain a reliable M_{BH} is $\theta_{\text{beam}} \lesssim \theta_{\text{rSOI}}$, where θ_{rSOI} is the angle subtended by r_{SOI} . Measurements using data with larger synthesized beams are more susceptible to systematic biases from stellar-mass uncertainties. Given $M_{\text{BH}} \approx 1.8 \times 10^7 M_{\odot}$ and $\sigma_{*} \approx 150 \text{ km s}^{-1}$ (Onken et al. 2004) gives $r_{\text{SOI}} \equiv GM_{\text{BH}}/\sigma_{*}^2 \approx 3 \text{ pc}$ ($\approx 0''.01$), a factor of ≈ 30 times smaller than what our ALMA observations can resolve. This work thus belongs to the majority of M_{BH} measurements that have $\theta_{\text{beam}} > \theta_{\text{rSOI}}$ (Davis et al. 2013, 2017,

2018; Onishi et al. 2015, 2017; Smith et al. 2019; Nguyen et al. 2020, Nguyen et al. submitted) and the lowest angular resolution with M_{BH} ever measured. Despite such a large θ_{beam} , our dynamical models using the atomic-[CI] kinematics can constrain a robust M_{BH} within a robust systemic/statistic uncertainty ($\gtrsim 35$ per cent when $\theta_{\text{beam}} > \theta_{\text{rSOI}}$; Nguyen et al. 2020). While the same constraints using the molecular- $^{12}\text{CO}(1-0)$ emission were subject to much larger errors with possible estimated- M_{BH} that are all within the mass range of 10^6 – $10^8 M_{\odot}$ (Table 6). The main reason is that the atomic-[CI] emission is highly concentrated at the galaxy centre (close to the SMBH) due to XDR effects around the AGN (120), while the molecular- $^{12}\text{CO}(1-0)$ emission is much more extended (Fig. 1). However, observations at higher angular resolutions would benefit our M_{BH} estimate by further reducing the uncertainties arising from the stellar mass.

It is also worth mentioning that the dynamical modelling of the other CO lines observed in this program did not improve the inferred- M_{BH} uncertainty significantly comparing to the chosen $^{12}\text{CO}(1-0)$. These molecular emissions distributions are a bit more compact towards the centre of NGC 7469, but their spatial resolutions are




Figure 10. Same as Fig. 9 but for $^{12}\text{CO}(1-0)$. The best-fitting model has $M_{\text{BH}} = 1.6 \times 10^7 M_{\odot}$ and $M/L_{F547M} = 2.22 (M_{\odot}/L_{\odot})$.




Figure 11. Same as Fig. 6 but for $[\text{CO}(1-0)]$. The best-fitting KinMS model is the one shown in panel (c) of Fig. 10.

a little lower than $^{12}\text{CO}(1-0)$. Additionally, the appearance of the $^{12}\text{CO}(2-1)$ and $^{12}\text{CO}(3-2)$ PVDs are similar to that of $^{12}\text{CO}(1-0)$ PVD. i.e. there would be a ‘hole’ at the AGN position (or CO-deficit; Smith et al. 2019; Nguyen et al. 2020), hinted by three resolved clumps in the integrated intensity map of the $^{12}\text{CO}(1-0)$ emission in

panel (a) of Fig. 2; also see fig. 2 of Izumi et al. (2020). Furthermore, the compactness of the molecular emission has an advantage in reducing the M_{BH} uncertainty but increasing the M/L_{F547M} uncertainty in the same manner as seen in the case of $[\text{CI}](1-0)$, which is clearly centrally peaked [panel (a) of Fig. 3].




Figure 12. The *HST* $F547M-F814W$ ($\approx V - I$) colour map within $7''.5$ (≈ 2.5 kpc) nuclear region. The map centres at the ALMA continuum peak. Redder areas are resulted from older populations and dust extinction, while bluer areas are resulted from very young star-forming regions. The southern and western half of the nucleus appears to have little extinction. The black trail at the centre includes the excluded spaxels of saturation in the $F814W$ image. The black contours show the *HST*/ $F547M$ continuum emission at the surface brightness density of 17.2, 18.3, and 19.1 mag arcsec $^{-2}$ (also see the photometry in V band of Doroshenko, Lyutyi & Shenavrin 1998).

4.4 The stellar-mass models account for spatial variations of populations and extinction

There is a colour variation due to complex stellar populations and dust extinction in the nucleus region of NGC 7469 shown in Fig. 12, implying that our assumption of a constant M/L is too simple. Here, we used an additional *HST*/ $F814W$ ($\approx I$) image (Table 2) in combination with the $F547M$ ($\approx V$) image to create a $F547M-F814W$ colour map (Fig. 12). Next, we convolved each astrometrically aligned image to the other PSF image (e.g. the $F547M$ image convolved with the $F814W$ PSF) to mitigate spurious gradients near the centre of the galaxy due to the difference of PSF widths. Then, we subtracted off the sky background level on each image calculated in an annulus located at (20–22 arcsec) away from the galaxy centre.

There is a heavy dust-extinction on the northern and north-east sides, while the southern and south-west sides of the nucleus have little extinction. Also, the starburst ring appears to host very young star clusters, suggesting the M/L variation. Such variation had a significant impact on the dynamical models of $\lesssim 10^7 M_{\odot}$ SMBHs (McConnell & Ma 2013; Thater et al. 2019; Nguyen et al. 2018, N19). In principle, we could apply the colour- M/L scaling relations for the young nucleus (< 1 Gyr) compiled by B01 and N19 to convert this colour map into their corresponding M/L maps (panels a and b of Fig. 13) and mass-surface-density maps (panels c and d of Fig. 13) that account for the variation of stellar populations pixel by pixel.

However, the central saturation found in $F814W$ prevents us from creating an accurate mass-surface density at the galaxy centre, where the SMBH’s gravitational potential dominates.

The N19-mass map predicts less mass at redder regions of contaminated dust and more mass at the bluer regions of young stars in the starburst ring than the B01-mass map’s prediction. It is also much more symmetric and thus is a better description of true mass distribution in the nucleus of NGC 7469 because the N19 colour- M/L correlation specifically applied for young populations (< 1 Gyr) and accounted for the extinction by dust. On the other hand, the B01 relation only relies on the spectrophotometric evolution models of spiral galaxies and the colours of the integrated stellar populations.

We converted the B01 and N19 mass-surface-density maps into their MGE forms with the central stellar mass in the centrally saturated spaxels interpolated from the outer pixels. Then, in their KinMS models, we first replaced the M/L_{F547M} parameter by the mass-scaling factor $\Gamma = (M/L_{\text{dyn}})/(M/L_{\text{pop}})$ to scale the mass-surface-density profiles directly, where (M/L_{dyn}) is the mass-to-light ratio constrained from dynamical model and (M/L_{pop}) is that determined from stellar populations. Next, we allowed M_{BH} and Γ to vary as free parameters, fixed other well-constrained parameters at their best fits shown in Table 6, and ran the model with the kinematics of the atomic-[CI](1–0) emission only. These resulted in $M_{\text{BH}} = 2.2^{+3.3}_{-1.3} \times 10^7 M_{\odot}$ and $\Gamma = 1.09^{+0.21}_{-0.23}$ for the B01-mass map and $M_{\text{BH}} = 2.9^{+1.8}_{-1.7} \times 10^7 M_{\odot}$ and $\Gamma = 0.93^{+0.20}_{-0.20}$ for the N19-mass map. Each of these masses is a factor of ≈ 1.3 and 1.6 higher than our best-fitting model in Section 4.3, indicating that the specific stellar populations and dust extinction in the nucleus of NGC 7469 have somewhat of an impact on our dynamical modelling.

4.5 Other uncertainty sources on the M_{BH} estimate

We test here the robustness of our dynamical model under the influence of sources of error other than the uncertainties in ALMA kinematics and stellar-mass errors. To this end, we tested dynamical models with the atomic-[CI](1–0) kinematics and allowed only the M_{BH} and M/L_{F547M} changing as free parameters while fixed other parameters at their best-fitting values in Table 6. The results of these tests include the best-fitting parameters associated with 1σ (16–84 per cent) and 3σ (0.14–99.86 per cent) CLs recorded in Table 7.

4.5.1 Izumi et al. (2020) CO-to- versus [CI]-to- H_2 conversion factor

Our choice of the starbursting galaxy conversion factor to estimate the molecular gas mass in Section 4.2 is uncertain because it could vary on small scales of kinematics and could also be lower in the very centre due to stellar processes. Meier, Turner & Hurt (2008) have found that X_{CO} varies in the range of $(0.5-1) \times 10^{20}$ cm $^{-2}$ (K km s $^{-1}$) $^{-1}$ in local normal spiral galaxies due to spiral arm/bar streaming. It is also too high in clouds with substantial stellar content. Based on dynamical modellings, I20 found the CO-to- and [CI]-to- H_2 conversion factor of $X_{\text{CO}} = 1.9 \times 10^{20}$ cm $^{-2}$ (K km s $^{-1}$) $^{-1}$ and $X_{[\text{CI}](1-0)} = 2.1 \times 10^{20}$ cm $^{-2}$ (K km s $^{-1}$) $^{-1}$, respectively, at the innermost ≈ 100 pc region of NGC 7469. The CO-to- H_2 conversion factor is, coincidentally equal to, while the [CI]-to- H_2 conversion factor is smaller than, those derived for Galactic star-forming regions due to the elevated C^0 abundance in the XDR. These suggest that our original adoption of the starburst CO-to- H_2 conversion factor may be inappropriate and may underestimate the total molecular gas mass by a factor of ≈ 2 . We, therefore,




Figure 13. The M/L_V maps constructed from the $F547M$ – $F814W$ colour map using the B01 (panel a) and the N19 (panel b) colour– M/L correlation. Panel (c): The nuclear surface-density-mass maps of NGC 7469 constructed by multiplying the $F547M$ surface-density-luminosity map to the B01 (panel c) and the N19 (panel d) M/L_V map pixel by pixel. Other illustrations (e.g. contours and saturated spaxels) are similar to Fig. 12.

used instead these newly derived conversion factors of NGC 7469 to test our modellings. The best-fitting KinMS models then yield $M_{\text{BH}} = 9.3^{+9.2}_{-4.6} \times 10^6 M_\odot$ and $M/L_{F547M} = 2.27^{+0.46}_{-0.45} (M_\odot/L_\odot)$ for CO-to- H_2 and $M_{\text{BH}} = 9.8^{+12.6}_{-5.5} \times 10^6 M_\odot$ and $M/L_{F547M} =$

$2.30^{+0.43}_{-0.43} (M_\odot/L_\odot)$ for [CI]-to- H_2 , suggesting these adopted conversion factors have a significant impact on the derived M_{BH} but within 3σ uncertainties of our best-fitting models in Section 4.3 and Table 6.

Table 7. Best-fitting model parameters and associated statistical uncertainties of the atomic-[CI](1–0) kinematics.

Parameter	Best fit	1 σ uncertainty (16–84%)	3 σ uncertainty (0.14–99.86%)
(1)	(2)	(3)	(4)
Bell & de Jong (2001) mass model			
$\log(M_{\text{BH}}/M_{\odot})$	7.35	−0.13, +0.13	−0.40, +0.38
Γ	1.09	−0.08, +0.07	−0.23, +0.21
Nguyen et al. (2019) mass model			
$\log(M_{\text{BH}}/M_{\odot})$	7.46	−0.14, +0.13	−0.42, +0.40
Γ	0.93	−0.07, +0.07	−0.21, +0.21
Izumi et al. (2020) CO-to-H₂ conversion factor			
$\log(M_{\text{BH}}/M_{\odot})$	6.97	−0.10, +0.10	−0.30, +0.30
$M/L_{F547M} (M_{\odot}/L_{\odot})$	2.27	−0.15, +0.15	−0.45, +0.46
Izumi et al. (2020) [CI]-to-H₂ conversion factor			
$\log(M_{\text{BH}}/M_{\odot})$	6.99	−0.12, +0.12	−0.36, +0.36
$M/L_{F547M} (M_{\odot}/L_{\odot})$	2.30	−0.14, +0.14	−0.43, +0.43
Axisymmetric ISM distribution			
$\log(M_{\text{BH}}/M_{\odot})$	7.26	−0.14, +0.14	−0.42, +0.42
$M/L_{F547M} (M_{\odot}/L_{\odot})$	2.19	−0.13, +0.13	−0.40, +0.39
AGN light contamination			
$\log(M_{\text{BH}}/M_{\odot})$	6.93	−0.09, +0.10	−0.28, +0.30
$M/L_{F547M} (M_{\odot}/L_{\odot})$	2.05	−0.07, +0.07	−0.21, +0.21

Note. The search range for Γ was set within (0.1–2.0), while the search ranges of $\log(M_{\text{BH}}/M_{\odot})$ and M/L_{F547M} were kept the same as in Table 6.

4.5.2 Axisymmetric ISM distribution

A different distribution of the molecular gas and dust within the nucleus of NGC 7469 may affect our dynamic results. Following Davis et al. (2020), we tested the impact of the ISM distribution using the axisymmetric assumption by converting the atomic-[CI](1–0) integrated intensity map into the M_{H_2} (+dust) map (120), then parametrizing this ISM-mass map into another MGE form. The dynamical model uses this axisymmetric-ISM-MGE mass (we turned off the gasGrav mechanism during the KinMS fit) gives $M_{\text{BH}} = 1.8^{+3.0}_{-1.1} \times 10^7 M_{\odot}$ and $M/L_{F547M} = 2.19^{+0.40}_{-0.39} (M_{\odot}/L_{\odot})$, suggesting our derived M_{BH} is less sensitive to different assumptions of ISM distributions as long as its total mass is determined accurately.

4.5.3 AGN light contamination

The bright AGN at the centre of NGC 7469 may contaminate a few central pixels of the $F547M$ image. The excess emission here increases the stellar mass density and decreases both the best fits of M_{BH} and M/L_{F547M} but still within 2σ CL (see Tables 6 and 7). To have this conclusion, we re-ran the $F547M$ -photometric-MGE model without masking those central pixels and used it in the KinMS model. We thus conclude that the AGN-light distribution indeed impacts our dynamical result but not significant.

5 DISCUSSION

5.1 Constraining the BLR inclination angle

RM is the commonly use method to reveal SMBHs and constrain their masses in Seyfert 1 AGN using the emission of the BLRs in the time domain rather than resolving it in spatial scale (Blandford & McKee 1982; Peterson & Caldwell 1993; Kaspi 2001; Kaspi et al. 2005; Bentz et al. 2006a, b, 2013). It uses the intrinsic time variability

of AGN emission to measure the time delay between continuum variations in the accretion discs and the broad emission-line response (e.g. Woo & Urry 2002a, b; McLure & Dunlop 2004; Shen et al. 2011; Woo et al. 2019). The reverberation time lag for a broad emission line, multiplied by the speed of light c , gives a measure of the response-weighted radius of the BLR. When combined with the velocity width of the broad emission line (ΔV), the M_{BH} can be estimated using the virial equation: $M_{\text{BH}} = \langle f \rangle \times R_{\text{BLR}}(\Delta V)^2/G$, where $\langle f \rangle = 4.31 \pm 1.05$ is an average dimensionless scale factor incorporating information of the geometry, kinematics, and orientation of the BLR (Grier et al. 2013). However, RM provides a rough- M_{BH} estimate because (1) the normalization factor ($\langle f \rangle$) changes significantly in the range of (3.2–7.8) depending on samples and calibration methods (Graham et al. 2011; Woo et al. 2010, 2013; Park et al. 2012; Onken et al. 2014) and (2) its uncertainty has a scatter of >0.5 dex (Onken et al. 2004) generally unknown for individual objects due to our inability to directly resolve the BLR. These give a concern that non-virial motions in the BLR clouds or radiation pressure might cause large errors in RM-derived M_{BH} (Krolik 2001).

The RM-based MBH of NGC 7469 was determined with the canonical normalization factor, and thus there still remains systematic uncertainty given the possibility that the BLR of NGC 7469 may have different geometry and kinematics (Peterson et al. 2014; Wang et al. 2014) into account. These particular physical properties could include non-virial velocity components (Denney et al. 2009, 2010), radiation pressure perturbations (Marconi et al. 2008; Netzer & Marziani 2010), the relative thickness (h/R_{BLR}) of the Keplerian BLR orbital plane (Gaskell 2009), and the LOS inclination angle (i) of this plane (Wills & Browne 1986; Runnoe et al. 2014; Shen & Ho 2014). All of these properties were subsumed into the average dimensionless scaling factor for RM AGNs: $\langle f \rangle = 4.31 \pm 1.05$ (Grier et al. 2013). Given our dynamical M_{BH} constraint of $1.78 \times 10^7 M_{\odot}$ and its 1σ uncertainty (Table 6) and account for the consistency with 1σ errors of R_{BLR} and ΔV (Peterson et al. 2014), we find $f = 7.2^{+4.2}_{-3.4}$ for the BLR in NGC 7469 (replacing $\langle f \rangle$ by f).

We use this specific f to infer the geometry of the BLR based on its analytical expression of a planar BLR of a thick disc, which is given by the expression: $f = \left[4 \left(\sin^2 i + \left(\frac{h}{R_{\text{BLR}}} \right)^2 \right) \right]^{-1}$, where i is the inclination angle of the system relative to the projected LOS of the Keplerian velocity on the BLR orbital plane (Collin et al. 2006; Decarli et al. 2008). In this scenario, the BLR thickness may result from radiation pressure of an accretion disc, which creates turbulent motions, disc outflowing winds, and non-coplanar orbits (Collin et al. 2006; Czerny et al. 2016). Theoretically, active SMBHs are thought to accrete material in the form of accretion discs and powered by accretion flows. The accretion discs convert gravitational energy into intense radiation (Shakura & Sunyaev 1973), making gas in the BLR move with velocities of thousands of kilometers per second. Under virial equilibrium, the M_{BH} can be determined from the intrinsic FWHM_{int} of observed lines by using the following equation: $M_{\text{BH}} = f G^{-1} R_{\text{BLR}} \text{FWHM}_{\text{int}}^2$. In practice, however, the f factor derived by comparing single epoch M_{BH} with that masses obtained from the $M_{\text{BH}}-L$ (Decarli et al. 2008) or $M_{\text{BH}}-\sigma$ (Shen & Ho 2014) scaling relations of spheroidal galaxies, or from amplitude of the excess X-ray variability variance (Nikolajuk et al. 2006) reveal the anticorrelation between f and FWHM_{obs} of the broad emission lines ($f \propto \text{FWHM}_{\text{obs}}^{-1}$), implying that $\text{FWHM}_{\text{int}} \propto \text{FWHM}_{\text{obs}}^{1/2}$.

Recently, Mejía-Restrepo et al. (2018) used a sample of 39 $z \sim 1.55$, high S/N spectroscopic type-1 AGN observed with VLT/X-Shooter spectrograph to perform Monte Carlo simulations of the LOS inclination dependence of FWHM_{int} of the virialised velocity

component of the BLR. They found that only thin BLRs ($h/R_{\text{BLR}} \lesssim 0.1$) can reproduce the observed bi-dimensional distribution of $f \propto \text{FWHM}_{\text{obs}}^{-1}$ and the predicted $\text{FWHM}_{\text{int}} \propto \text{FWHM}_{\text{obs}}^{1/2}$. We thus consider the thin disc BLR model for the NGC 7469 AGN in which $(h/R_{\text{BLR}}) \approx 0$, giving the BLR's LOS inclination of $i \approx 11.0^{+2.2}_{-2.5}$, which tightly constrains the face-on orientation of the BLR. However, the physics of a compact BLR is likely to remain constant over time (i.e. consistent in terms of the Eddington ratio), making it is possible to study active SMBHs at low and high redshifts (Mejía-Restrepo et al. 2018).

5.2 A new game changer for constraining M_{BH} in AGN

This work shows that the M_{BH} in the AGN type 1 NGC 7469 can be constrained using the atomic-[CI](1–0) kinematics. The atomic-[CI](1–0) emission line is bright, strongly peaked towards the galaxy centre at the CND-scale due to the XDR around the AGN (120). These unique features suggest that the atomic-[CI](1–0) line must be more appropriate to measure central M_{BH} in active galaxies than the low-J CO-molecular lines. This newly precise measurement of M_{BH} using ALMA atomic-gas dynamics can also be used to constrain the virial geometric factor f of the BLRs in Seyfert 1 AGN NGC 7469, then provide the first insights into the unresolved structure of BLRs (except the BLR of 3C 273 seems to be spatially resolved by using GRAVITY; Gravity Collaboration 2018) as the effect of line-of-sight (LOS) inclination (i) in a planar distribution of the BLR emitting gas in which $f(i)$ seems following an anticorrelation (Mejía-Restrepo et al. 2018).

Constraining a functional form for such an important correlation based on a large sample of nearby AGN will provide a better understanding on the correlation between the M_{BH} and AGN properties (e.g. FWHM and the velocity dispersion of the emitting gas in the BLR) that will help to derive more accurate RM-based M_{BH} in more distant AGN beyond the local Universe ($z \gtrsim 0.1$). Besides, these precise measurements of M_{BH} will ultimately be a key to understand the M_{BH} function and the mean radiative efficiency of SMBH accretion (Shankar, Weinberg & Miralda-Escudé 2009; Shankar et al. 2016). Fortunately, high-spatial-resolution and high sensitivity of ALMA observations now allow us to probe the X-ray dissociated atomic-gas kinematics close to the SMBHs' SOI and infer their masses accurately through the resolved manner [i.e. it can achieve the synthesized beam of $\theta_{[\text{CI}](1-0)} \approx 0''.009$ (or 9 mas) with the most extended baseline of 16 km], which was impossible in previous studies (i.e. both RM and dynamics with low-J CO molecular tracers). This work pioneers the usage of the atomic-[CI](1–0) line as a new kinematics tracer and can be a game-changer in the M_{BH} measurement field, at least for the active galaxies where other methods cannot work well. More importantly, the atomic-[CI](1–0) emission has a higher rest-frequency than low-J CO-molecular lines, suitable for performing similar dynamical observations/measurements towards higher redshift objects (i.e. the synthesis beam of $\theta_{[\text{CI}](1-0)} \approx 9$ mas is good enough for constraining any black holes with $M_{\text{BH}} > 9 \times 10^8 M_{\odot}$ at all redshift⁸). Thus,

⁸Note that the curve r_{SOI} varies as a function of both redshift (z) and black hole mass (M_{BH}), $r_{\text{SOI}} \propto f(z, M_{\text{BH}})$, which has the minimum at $z \approx 1$. Given the current majority measurements in the local Universe with ALMA observations that have $\theta_{\text{beam}} > 2 \times r_{\text{SOI}}$ (references in the text), the speculated M_{BH} values could be smaller by a factor of a few. For example, Nguyen et al. (2020) constrain the M_{BH} in NGC 3504 at the observational scale of six times larger than its r_{SOI} . We thus argue that depending on the case, M_{BH} could be

we can extend this atomic-[CI](1–0) line survey and analysis for high-redshift AGN in the future.

5.3 Comparison to RM-based M_{BH} and scaling relations

As discussed in Section 1, the central M_{BH} of NGC 7469 was determined via both the direct signatures of radio and X-ray radiation on the Fundamental Plane and the indirect signature of the BLR variability. However, these M_{BH} occupy a wide range of mass from 10^5 to $10^8 M_{\odot}$. Our gas-dynamical models using ALMA observation in this work give an ≈ 40 per cent higher value for the M_{BH} than the estimate of Peterson et al. (2014) and Wang et al. (2014) using RM of the H β and He II $\lambda 4686$ emission lines, respectively.

Our derived M_{BH} of NGC 7469 is consistent with the empirical $M_{\text{BH}}-M_{\text{bulge}}$ correlation of both ETGs and late-type galaxies (LTGs; Sahu et al. 2019a; Sahu, Graham & Davis 2019b) within 3σ uncertainty. However, it is about one order of magnitude below the same correlations compiled mainly for higher mass galaxies that have Sérsic profiles without central cores of Kormendy & Ho (2013), McConnell et al. (2013), and Saglia et al. (2016), as well as 1.5 mag below the same correlation of Scott, Graham & Schombert (2013), compiled for the core-Sérsic and lower mass galaxies ($M_{\star} \lesssim 2 \times 10^{10} M_{\odot}$). It is also well below the Greene, Strader & Ho (2020) correlation with the same amount of magnitude compiled for all types and masses of galaxies. In the context of $M_{\text{BH}}-\sigma_{\star}$, the NGC 7469 M_{BH} is consistent with the McConnell et al. (2013), Kormendy & Ho (2013), Saglia et al. (2016), and Sahu et al. (2019a, b) correlations rather than the correlation of Greene et al. (2020). We show our measurement of NGC 7469 in the contexts of the Kormendy & Ho (2013) and Sahu et al. (2019a) $M_{\text{BH}}-M_{\text{bulge}}$ and $M_{\text{BH}}-\sigma_{\star}$ correlations in Fig. 14 only. Here, the former is the most high-cited scaling relation, while the latter is the recent most completed compilation correlation with adding new measurements and revisions from 2013.

Smith et al. (2021b) recently investigated the empirical correlations between M_{BH} and the flat rotation velocities measured either at a large radius in their rotation curves or via spatially integrated emission line widths of a sample of both spatially resolved and unresolved galaxies (so-called $\Delta V_{\text{CO}}-M_{\text{BH}}$ correlation) that have M_{BH} constrained via molecular-gas-dynamical modellings. Their spatially resolved sub-sample includes 27 galaxies with CO-CND detected, and nine of them have dynamical- M_{BH} constrained via ALMA in the same manner we did here for NGC 7469. The unresolved sub-sample, on the other hand, includes 24 same targets with six ALMA dynamical- M_{BH} . Here, they assumed the rotation velocity traced by the de-projected integrated CO emission line width and found a tight correlation of 24 spatially resolved CO discs (three targets were omitted): $\log(M_{\text{BH}}/M_{\odot}) = (7.5 \pm 0.1) + (8.5 \pm 0.9)[\log(W_{50}/\sin i \text{ km s}^{-1}) - 2.7]$, where W_{50} is the FWHM of a double-horned emission line profile and i the inclination of the CO disc. Another tight correlation between this de-projected CO line widths (flat rotation velocity) and the stellar-velocity dispersion averaged within one effective radius of $\log(\sigma_{\star}/\text{km s}^{-1}) = (2.20 \pm 0.02) + (1.1 \pm 0.1)[\log(W_{50}/\sin i \text{ km s}^{-1}) - 2.7]$ was also found, which is so-called the $\Delta V_{\text{CO}}-\sigma_{\star}$ correlation. In the context of the $\Delta V_{\text{CO}}-M_{\text{BH}}$ correlation and with $\Delta V_{\text{CO}} \approx 400 \text{ km s}^{-1}$ (Section 3.2), NGC 7469 is about ≈ 0.2 dex of out 1σ intrinsic scatter (0.5 dex). This outlier may be caused by the nearly face-on orientation of the CO/[CI] disc, implying that the inclination uncertainties are very large. However,

lower by an appropriate factor in comparison to the value inferred from the function $r_{\text{SOI}} \propto f(z, M_{\text{BH}})$.




Figure 14. Our NGC 7469 M_{BH} (red encircle) in the context of the $M_{\text{BH}}-M_{\text{bulge}}$ (left) and $M_{\text{BH}}-\sigma_*$ scaling relations (right). 24 molecular-gas- M_{BH} measurements using both ALMA (Onishi et al. 2015; Barth et al. 2016a, b; Davis et al. 2017, 2018, 2020; Boizelle et al. 2019, 2020; Combes et al. 2019; Smith et al. 2019; Nagai et al. 2019; North et al. 2019; Thater 2019; Nguyen et al. 2020, Nguyen et al., submitted, Smith et al. 2021a) and CARMA (Davis et al. 2013; Onishi et al. 2017) observations are plotted in cyan. The scaling relations of Sahu, Graham & Davis (2019a) and Kormendy & Ho (2013) are plotted in the specific-solid-colour lines shown in the legend. M_{bulge} are taken from McConnell et al. (2013), Krajnović et al. (2013), Salo et al. (2015), Huang et al. (2016), Savorgnan et al. (2016), and Sani et al. (2018), while σ_* are taken from Ho et al. (2009), Seth et al. (2010), Coccato et al. (2013), Krajnović et al. (2013), Kormendy & Ho (2013), Savorgnan et al. (2016), and Sani et al. (2018). Error bars of our measurement is 3σ uncertainty.

in terms of the $\Delta V_{\text{CO}}-\sigma_*$ correlation, the M_{BH} of NGC 7469 is well predicted (scatter $\lesssim 0.1$ dex). As companions to other M_{BH} -galaxy’s properties scaling relations, these two correlations can also be used to estimate the local- M_{BH} function.

6 CONCLUSIONS

We present dynamical-mass measurements for the central SMBH in the AGN type 1 NGC 7469 using *HST* imaging and the bright atomic-[CI](1–0) and molecular- $^{12}\text{CO}(1-0)$ emissions observed by ALMA. We highlight our main results below:

(i) NGC 7469 hosts multiple transitions of molecular-CO and atomic-[CI] lines within the CNB and starburst ring. Detailed analysis of the $^{12}\text{CO}(1-0)$ and [CI](1–0) emissions find a dynamically settled CNB in the inner 1 arcsec of the galaxy centre.

(ii) Our dynamical modelling for the atomic-[CI](1–0) data suggests the presence of a central SMBH of $M_{\text{BH}} = 1.78^{+2.69}_{-1.10} \times 10^7 M_{\odot}$, while that of the molecular- $^{12}\text{CO}(1-0)$ data provides a mass of $M_{\text{BH}} = 1.60^{+11.52}_{-1.45} \times 10^7 M_{\odot}$, a factor of $\approx (1.5-2)$ higher than the Peterson et al. (2014) RM-based- M_{BH} . These two models also find a consistent $M/L_{F547M} \approx 2.20 (M_{\odot}/L_{\odot})$.

(iii) The atomic-[CI](1–0) emission may be the best line for performing the gas-dynamical modelling and constraining central M_{BH} in active galaxies as [CI](1–0) morphological distribution is closer to the SMBH and brighter than the low-J CO lines. Thus, it is an excellent tracer to do cross-checks between RM-based and gas-dynamical methods in near and far AGN. However, we should caution that users need to test this point further with other AGN and higher spatial resolution data.

(iv) Our new M_{BH} estimate for NGC 7469 provides a specific value for the RM AGN dimensionless scaling factor of the BLR of

$f = 7.2^{+4.2}_{-3.4}$ for the first time. This value is ≈ 40 per cent higher than the average value of $\langle f \rangle$ applied for all previous indirect RM-based M_{BH} estimates in Seyfert 1 AGN. It also reveals a thin accretion disc for the unresolved BLR, which is oriented face-on with the LOS inclination of $i \approx 11.0^{+2.2}_{-2.5}$.

(v) Our M_{BH} is consistent with the empirical $M_{\text{BH}}-M_{\text{bulge}}$ correlations of the recent compilation of Sahu et al. (2019b) for all types of galaxies, which have direct- M_{BH} measurements. However, it is almost one order of magnitude below the correlations of Kormendy & Ho (2013), McConnell et al. (2013), and Saglia et al. (2016) of the more massive SMBHs and galaxies without the Sérsic bulges in their surface brightness profiles. This negative offset is even larger (roughly 1.5 mag) in the context of the Greene et al. (2020) scaling relation compiled for all types and masses of galaxies and the Scott et al. (2013) correlation compiled for lower mass galaxies with Sérsic cores. In the context of the $M_{\text{BH}}-\sigma_*$ correlation, the M_{BH} seems consistent with the Kormendy & Ho (2013), McConnell et al. (2013), Saglia et al. (2016), and Sahu et al. (2019a, b) scaling relations but inconsistent with the compilation of Greene et al. (2020), which is above the locations of the SMBH about one order of magnitude. Our M_{BH} is also consistent with the $\Delta V_{\text{CO}}-\sigma_*$ correlation but offset 0.2 dex outside 1σ (0.5 dex) scatter of the $\Delta V_{\text{CO}}-M_{\text{BH}}$ scaling relation, which are recently compiled by Smith et al. (2021b).

(vi) The NGC 7469’s M_{BH} has been constrained consistently using molecular/atomic gas of ALMA, although the observational scale of the data is a factor of ≈ 30 times larger than the angle subtended by the SMBH’s r_{SOI} . Because of the poorly resolvable scale of our data, we argue this consistency applies to the case of NGC 7469 only.

(vii) This work pioneers the user of the atomic-[CI] line in constraining M_{BH} in AGN accurately, which then allows estimating the BLR’s inclination angle, a crucial ingredient to understand the unified AGN model.

Facilities: ALMA and HST/UVIS WFC3.

Software: IDL, CASA, ASTROPY, EMCEE, KINMS, KINOMETRY, and MGEFIT.

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

ACKNOWLEDGEMENTS

The authors would like to thank the anonymous referee for their careful reading and useful comments, that helped to improve the paper greatly. The authors also thank National Astronomical Observatory of Japan (NAOJ), National Institute of Natural Sciences (NINS), and the PHENIKAA Institute for Advanced Study (PIAS of PHENIKAA University) for supporting this work. TI, SB, and KK gratitude to the support from the Japan Society for the Promotion of Science (JSPS) KAKENHI grant #JP20K14531, 19J00892, and JP17H06130 (and the NAOJ ALMA Scientific Research Grant Number 2017-06B), respectively. ST acknowledges funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement no 724857 (Consolidator Grant ArcheoDyn).

This paper makes use of the following ALMA data: ADS/JAO.ALMA #2017.1.00078.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada) and NSC and ASIAA (Taiwan) and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, AUI/NRAO and NAOJ. We also thank the ALMA operators and staff and the ALMA help desk for diligent feedback and invaluable assistance in processing these data.

REFERENCES

- Alberdi A., Colina L., Torrelles J. M., Panagia N., Wilson A. S., Garrington S. T., 2006, *ApJ*, 638, 938
- Avila R. J., Hack W. J., STSci AstroDrizzle Team, 2012, *American Astronomical Society Meeting Abstracts* 220. p. #135.13
- Baldassare V. F., Reines A. E., Gallo E., Greene J. E., 2015, *ApJ*, 809, L14
- Barth A. J., Boizelle B. D., Darling J., Baker A. J., Buote D. A., Ho L. C., Walsh J. L., 2016a, *ApJ*, 822, L28
- Barth A. J., Darling J., Baker A. J., Boizelle B. D., Buote D. A., Ho L. C., Walsh J. L., 2016b, *ApJ*, 823, 51
- Bell E. F., de Jong R. S., 2001, *ApJ*, 550, 212
- Bentz M. C., Peterson B. M., Pogge R. W., Vestergaard M., Onken C. A., 2006a, *ApJ*, 644, 133
- Bentz M. C. et al., 2006b, *ApJ*, 651, 775
- Bentz M. C. et al., 2013, *ApJ*, 767, 149
- Binney J., Tremaine S., 1987, *Galactic Dynamics*. Princeton Univ. Press, Princeton
- Blandford R. D., McKee C. F., 1982, *ApJ*, 255, 419
- Blustein A. J. et al., 2007, *A&A*, 466, 107
- Boizelle B. D., Barth A. J., Walsh J. L., Buote D. A., Baker A. J., Darling J., Ho L. C., 2019, *ApJ*, 881, 10
- Boizelle B. D. et al., 2021, *ApJ*, 908, 1
- Bonatto C. J., Pastoriza M. G., 1990, *ApJ*, 353, 445
- Bolatto A. D. et al., 2013, *Nature*, 499, 450
- Cappellari M., 2002, *MNRAS*, 333, 400
- Cappellari M., 2008, *MNRAS*, 390, 71
- Cardelli J. A., Clayton G. C., Mathis J. S., 1989, *ApJ*, 345, 245
- Coccato L., Sarzi M., Pizzella A., Corsini E. M., Dalla Bontà E., Bertola F., 2006, *MNRAS*, 366, 1050
- Coccato L., Morelli L., Pizzella A., Corsini E. M., Buson L. M., Dalla Bontà E., 2013, *A&A*, 549, A3
- Collier S. J. et al., 1998, *ApJ*, 500, 162
- Collin S., Kawaguchi T., Peterson B. M., Vestergaard M., 2006, *A&A*, 456, 75
- Combes F. et al., 2019, *A&A*, 623, A79
- Czerny B., Du P., Wang J.-M., Karas V., 2016, *ApJ*, 832, 15
- Dame T. M., 2011, preprint([arXiv:1101.1499](https://arxiv.org/abs/1101.1499))
- Dame T. M., Hartmann D., Thaddeus P., 2001, *ApJ*, 547, 792
- Davies R. I., Tacconi L. J., Genzel R., 2004, *ApJ*, 602, 148
- Davies R. I., Müller Sánchez F., Genzel R., Tacconi L. J., Hicks E. K. S., Friedrich S., Sternberg A., 2007, *ApJ*, 671, 1388
- Davis T. A., 2014, *MNRAS*, 443, 911
- Davis T. A., Bureau M., Cappellari M., Sarzi M., Blitz L., 2013, *Nature*, 494, 328
- Davis T. A., Bureau M., Onishi K., Cappellari M., Iguchi S., Sarzi M., 2017, *MNRAS*, 468, 4675
- Davis T. A. et al., 2018, *MNRAS*, 473, 3818
- Davis T. A. et al., 2020, *MNRAS*, 496, 4061
- Decarli R., Labita M., Treves A., Falomo R., 2008, *MNRAS*, 387, 1237
- den Brok M. et al., 2015, *ApJ*, 809, 101
- Denney K. D. et al., 2009, *ApJ*, 704, L80
- Denney K. D. et al., 2010, *ApJ*, 721, 715
- Díaz-Santos T., Alonso-Herrero A., Colina L., Ryder S. D., Knapen J. H., 2007, *ApJ*, 661, 149
- Doroshenko V. T., Lyutyi V. M., Shenavrin V. I., 1998, *Astron. Lett.*, 24, 160
- Emsellem E., Monnet G., Bacon R., 1994, *A&A*, 285, 723
- Event Horizon Telescope Collaboration, 2019a, *ApJ*, 875, L1
- Event Horizon Telescope Collaboration, 2019b, *ApJ*, 875, L4
- Event Horizon Telescope Collaboration, 2019c, *ApJ*, 875, L6
- Ferrarese L., Merritt D., 2000, *ApJ*, 539, L9
- Foreman-Mackey D., Hogg D. W., Lang D., Goodman J., 2013, *PASP*, 125, 306
- Gaskell C. M., 2009, *New Astron. Rev.*, 53, 140
- Gebhardt K. et al., 2000, *ApJ*, 539, L13
- Gebhardt K., Adams J., Richstone D., Lauer T. R., Faber S. M., Gültekin K., Murphy J., Tremaine S., 2011, *ApJ*, 729, 119
- Genzel R., Weitzel L., Tacconi-Garman L. E., Blietz M., Cameron M., Krabbe A., Lutz D., Sternberg A., 1995, *ApJ*, 444, 129
- Goodman J., Weare J., 2010, *Commun. Appl. Math. Computat. Sci.*, 5, 65
- Graham A. W., Onken C. A., Athanassoula E., Combes F., 2011, *MNRAS*, 412, 2211
- Gravity Collaboration, 2018, *Nature*, 563, 657
- Greene J. E. et al., 2010, *ApJ*, 721, 26
- Greene J. E., Strader J., Ho L. C., 2020, *ARA&A*, 58, 257
- Grier C. J. et al., 2013, *ApJ*, 764, 47
- Guilloteau S. et al., 1992, *A&A*, 262, 624
- Gültekin K., King A. L., Cackett E. M., Nyland K., Miller J. M., Di Matteo T., Markoff S., Rupen M. P., 2019, *ApJ*, 871, 80
- Häring N., Rix H.-W., 2004, *ApJ*, 604, L89
- Ho L. C., Greene J. E., Filippenko A. V., Sargent W. L. W., 2009, *ApJS*, 183, 1
- Huang S., Ho L. C., Peng C. Y., Li Z.-Y., Barth A. J., 2016, *ApJ*, 821, 114
- Imanishi M., Wada K., 2004, *ApJ*, 617, 214
- Imanishi M., Nakanishi K., Izumi T., 2016, *AJ*, 152, 218
- Imanishi M., Nakanishi K., Izumi T., Wada K., 2018, *ApJ*, 853, L25
- Izumi T. et al., 2015, *ApJ*, 811, 39
- Izumi T., Wada K., Fukushige R., Hamamura S., Kohno K., 2018, *ApJ*, 867, 48
- Izumi T. et al., 2020, *ApJ*, 898, 75
- Kaspi S., 2001, *ASP Conf. Ser. Vol. 224, Reverberation Measurements of Quasars and the Size-Mass-Luminosity Relationships in Active Galactic Nuclei*. Astron. Soc. Pac., San Francisco, p. 347
- Kaspi S., Smith P. S., Netzer H., Maoz D., Jannuzi B. T., Giveon U., 2000, *ApJ*, 533, 631
- Kaspi S., Maoz D., Netzer H., Peterson B. M., Vestergaard M., Jannuzi B. T., 2005, *ApJ*, 629, 61

- Knapen J. H., Shlosman I., Peletier R. F., 2000, *ApJ*, 529, 93
- Kormendy J., Ho L. C., 2013, *ARA&A*, 51, 511
- Kormendy J., Richstone D., 1995, *ARA&A*, 33, 581
- Krajnović D., Cappellari M., de Zeeuw P. T., Copin Y., 2006, *MNRAS*, 366, 787
- Krajnović D. et al., 2013, *MNRAS*, 432, 1768
- Krajnović D. et al., 2018, *MNRAS*, 477, 3030
- Kriss G. A., Peterson B. M., Crenshaw D. M., Zheng W., 2000, *ApJ*, 535, 58
- Krist J., 1995, in Shaw R. A., Payne H. E., Hayes J. J. E., eds, *ASP Conf. Ser. Vol. 77, Astronomical Data Analysis Software and Systems IV*. Astron. Soc. Pac., San Francisco, p. 349
- Krist J. E., Hook R. N., Stoehr F., 2011, in Kahan M. A., ed., *Proc. SPIE Conf. Ser. Vol. 8127, Optical Modeling and Performance Predictions V*. SPIE, Bellingham
- Krolik J. H., 2001, *ApJ*, 551, 72
- Kuno N., Nishiyama K., Nakai N., Sorai K., Vila-Vilaró B., Handa T., 2000, *PASJ*, 52, 775
- Kuno N. et al., 2007, *PASJ*, 59, 117
- Kuo C. Y. et al., 2011, *ApJ*, 727, 20
- Liu T., Wang J.-X., Yang H., Zhu F.-F., Zhou Y.-Y., 2014, *ApJ*, 783, 106
- Lonsdale C. J., Lonsdale C. J., Smith H. E., Diamond P. J., 2003, *ApJ*, 592, 804
- Maloney P. R., Hollenbach D. J., Tielens A. G. G. M., 1996, *ApJ*, 466, 561
- Marconi A., Axon D. J., Maiolino R., Nagao T., Pastorini G., Pietrini P., Robinson A., Torricelli G., 2008, *ApJ*, 678, 693
- McConnell N. J., Ma C.-P., 2013, *ApJ*, 764, 184
- McConnell N. J., Chen S.-F. S., Ma C.-P., Greene J. E., Lauer T. R., Gebhardt K., 2013, *ApJ*, 768, L21
- McLure R. J., Dunlop J. S., 2001, *MNRAS*, 327, 199
- McLure R. J., Dunlop J. S., 2004, *MNRAS*, 352, 1390
- McLure R. J., Dunlop J. S., Kukula M. J., 2000, *MNRAS*, 318, 693
- McMullin J. P., Waters B., Schiebel D., Young W., Golap K., 2007, in Shaw R. A., Hill F., Bell D. J., eds, *ASP Conf. Ser. Vol. 376, Astronomical Data Analysis Software and Systems XVI*. Astron. Soc. Pac., San Francisco, p. 127
- Meier D. S., Turner J. L., Hurt R. L., 2008, *ApJ*, 675, 281
- Meijerink R., Spaans M., 2005, *A&A*, 436, 397
- Meixner M., Puchalsky R., Blitz L., Wright M., Heckman T., 1990, *ApJ*, 354, 158
- Mejía-Restrepo J. E., Lira P., Netzer H., Trakhtenbrot B., Capellupo D. M., 2018, *Nat. Astron.*, 2, 63
- Mitzkus M., Cappellari M., Walcher C. J., 2017, *MNRAS*, 464, 4789
- Nagai H. et al., 2019, *ApJ*, 883, 193
- Nandra K., Le T., George I. M., Edelson R. A., Mushotzky R. F., Peterson B. M., Turner T. J., 2000, *ApJ*, 544, 734
- Netzer H., Marziani P., 2010, *ApJ*, 724, 318
- Nguyen D. D. et al., 2017, *ApJ*, 836, 237
- Nguyen D. D. et al., 2018, *ApJ*, 858, 118
- Nguyen D. D. et al., 2019, *ApJ*, 872, 104
- Nguyen D. D. et al., 2020, *ApJ*, 892, 68
- Nikołajuk M., Czerny B., Ziółkowski J., Gierliński M., 2006, *MNRAS*, 370, 1534
- North E. V. et al., 2019, *MNRAS*, 490, 319
- Onishi K., Iguchi S., Sheth K., Kohno K., 2015, *ApJ*, 806, 39
- Onishi K., Iguchi S., Davis T. A., Bureau M., Cappellari M., Sarzi M., Blitz L., 2017, *MNRAS*, 468, 4663
- Onken C. A., Ferrarese L., Merritt D., Peterson B. M., Pogge R. W., Vestergaard M., Wandel A., 2004, *ApJ*, 615, 645
- Onken et al., 2014, *ApJ*, 791, 37
- Osterbrock D. E., Martel A., 1993, *ApJ*, 414, 552
- Pacucci F., Natarajan P., Volonteri M., Cappelluti N., Urry C. M., 2017, *ApJ*, 850, L42
- Pacucci F., Loeb A., Mezcua M., Martín-Navarro I., 2018, *ApJ*, 864, L6
- Papadopoulos P. P., Allen M. L., 2000, *ApJ*, 537, 631
- Park D. et al., 2012, *ApJ*, 747, 30
- Pérez-Torres M. A., Alberdi A., Colina L., Torrelles J. M., Panagia N., Wilson A., Kankare E., Mattila S., 2009, *MNRAS*, 399, 1641
- Peterson R. C., Caldwell N., 1993, *AJ*, 105, 1411
- Peterson B. M. et al., 2004, *ApJ*, 613, 682
- Peterson B. M. et al., 2014, *ApJ*, 795, 149
- Petrucci P. O., Maraschi L., Haardt F., Nandra K., 2004, *A&A*, 413, 477
- Plotkin R. M., Markoff S., Kelly B. C., Kording E., Anderson S. F., 2012, *MNRAS*, 419, 267
- Press M., 2007, *Sci. Am.*, 297, 100
- Runnoe J. C., Brotherton M. S., DiPompeo M. A., Shang Z., 2014, *MNRAS*, 438, 3263
- Rusli S. P., Erwin P., Saglia R. P., Thomas J., Fabricius M., Bender R., Nowak N., 2013, *AJ*, 146, 160
- Saglia R. P. et al., 2016, *ApJ*, 818, 47
- Sahu N., Graham A. W., Davis B. L., 2019a, *ApJ*, 876, 155
- Sahu N., Graham A. W., Davis B. L., 2019b, *ApJ*, 887, 10
- Salo H. et al., 2015, *ApJS*, 219, 4
- Sanders D. B., Mazzarella J. M., Kim D.-C., Surace J. A., Soifer B. T., 2003, *AJ*, 126, 1607
- Sani E. et al., 2018, *Front. Astron Space Sci.*, 5, 2
- Savognan G. A. D., Graham A. W., Marconi A., Sani E., 2016, *ApJ*, 817, 21
- Schawinski K., Thomas D., Sarzi M., Maraston C., Kaviraj S., Joo S.-J., Yi S. K., Silk J., 2007, *MNRAS*, 382, 1415
- Schlafly E. F., Finkbeiner D. P., 2011, *ApJ*, 737, 103
- Scott J. E. et al., 2005, *ApJ*, 634, 193
- Scott N., Graham A. W., Schombert J., 2013, *ApJ*, 768, 76
- Seth A. C. et al., 2010, *ApJ*, 714, 713
- Shakura N. I., Sunyaev R. A., 1973, in Bradt H., Giacconi R., eds, *IAU Symp. 55, X- and Gamma-Ray Astronomy*. Kluwer, Dordrecht, p. 155
- Shankar F., Weinberg D. H., Miralda-Escudé J., 2009, *ApJ*, 690, 20
- Shankar F. et al., 2016, *MNRAS*, 460, 3119
- Shen Y., Ho L. C., 2014, *Nature*, 513, 210
- Shen Y., Liu X., Greene J. E., Strauss M. A., 2011, *ApJ*, 735, 48
- Smith M. D. et al., 2019, *MNRAS*, 485, 4359
- Smith M. D. et al., 2021a, *MNRAS*, 503, 4
- Smith M. D. et al., 2021b, *MNRAS*, 500, 1933
- Soifer B. T., Bock J. J., Marsh K., Neugebauer G., Matthews K., Egami E., Armus L., 2003, *AJ*, 126, 143
- Thater S., 2019, preprint ([arXiv:1911.11491](https://arxiv.org/abs/1911.11491))
- Thater S., Krajnović D., Cappellari M., Davis T. A., de Zeeuw P. T., McDermid R. M., Sarzi M., 2019, *A&A*, 625, A62
- van den Bosch R. C. E., van de Ven G., 2009, *MNRAS*, 398, 1117
- Walsh J. L., Barth A. J., Ho L. C., Sarzi M., 2013, *ApJ*, 770, 86
- Wandel A., 2002, *ApJ*, 565, 762
- Wang J.-M. et al., 2014, *ApJ*, 793, 108
- Wills B. J., Browne I. W. A., 1986, *ApJ*, 302, 56
- Woo J.-H., Urry C. M., 2002a, *ApJ*, 579, 530
- Woo J.-H., Urry C. M., 2002b, *ApJ*, 581, L5
- Woo J.-H. et al., 2010, *ApJ*, 716, 269
- Woo J.-H., Schulze A., Park D., Kang W.-R., Kim S. C., Riechers D. A., 2013, *ApJ*, 772, 49
- Woo J.-H., Cho H., Gallo E., Hodges-Kluck E., Le H. A. N., Shin J., Son D., Horst J. C., 2019, *Nat. Astron.*, 3, 755

SUPPORTING INFORMATION

Supplementary data are available at *MNRAS* online.

Fig12.pdf

Fig13.pdf

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.