

# **End of the Century pCO(2) Levels Do Not Impact Calcification in Mediterranean Cold-Water Corals**

Cornelia Maier, Alexander Schubert, Maria M. Berzunza Sanchez, Markus G

Weinbauer, Pierre Watremez, Jean-Pierre Gattuso

## **To cite this version:**

Cornelia Maier, Alexander Schubert, Maria M. Berzunza Sanchez, Markus G Weinbauer, Pierre Watremez, et al.. End of the Century pCO(2) Levels Do Not Impact Calcification in Mediterranean Cold-Water Corals. PLoS ONE, 2013, 8 (4), 10.1371/journal.pone.0062655 . hal-03502634

## **HAL Id: hal-03502634 <https://hal.science/hal-03502634>**

Submitted on 26 Dec 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



[Distributed under a Creative Commons Attribution 4.0 International License](http://creativecommons.org/licenses/by/4.0/)

## End of the Century  $pCO<sub>2</sub>$  Levels Do Not Impact Calcification in Mediterranean Cold-Water Corals

## Cornelia Maier<sup>1,2\*</sup>, Alexander Schubert<sup>1,2</sup>, Maria M. Berzunza Sànchez<sup>1,2</sup>, Markus G. Weinbauer<sup>1,2</sup>, Pierre Watremez<sup>3</sup>, Jean-Pierre Gattuso<sup>1,2</sup>

1 Centre National de la Recherche Scientifique-Institut National des Sciences de l'Univers, Laboratoire d'Océanographie de Villefranche, Villefranche-sur-mer, France, 2 Université Pierre et Marie Curie-Paris 6, Observatoire Océanologique de Villefranche, Villefranche-sur-mer, France, 3 Agence des Aires Marines Protegées, Brest, France

## Abstract

Ocean acidification caused by anthropogenic uptake of  $CO<sub>2</sub>$  is perceived to be a major threat to calcifying organisms. Coldwater corals were thought to be strongly affected by a decrease in ocean pH due to their abundance in deep and cold waters which, in contrast to tropical coral reef waters, will soon become corrosive to calcium carbonate. Calcification rates of two Mediterranean cold-water coral species, Lophelia pertusa and Madrepora oculata, were measured under variable partial pressure of  $CO_2$  (pCO<sub>2</sub>) that ranged between 380 uatm for present-day conditions and 930 uatm for the end of the century. The present study addressed both short- and long-term responses by repeatedly determining calcification rates on the same specimens over a period of 9 months. Besides studying the direct, short-term response to elevated  $pCO<sub>2</sub>$  levels, the study aimed to elucidate the potential for acclimation of calcification of cold-water corals to ocean acidification. Net calcification of both species was unaffected by the levels of  $pCO<sub>2</sub>$  investigated and revealed no short-term shock and, therefore, no long-term acclimation in calcification to changes in the carbonate chemistry. There was an effect of time during repeated experiments with increasing net calcification rates for both species, however, as this pattern was found in all treatments, there is no indication that acclimation of calcification to ocean acidification occurred. The use of controls (initial and ambient net calcification rates) indicated that this increase was not caused by acclimation in calcification response to higher pCO2. An extrapolation of these data suggests that calcification of these two cold-water corals will not be affected by the  $pCO<sub>2</sub>$  level projected at the end of the century.

Citation: Maier C, Schubert A, Berzunza Sànchez MM, Weinbauer MG, Watremez P, et al. (2013) End of the Century pCO<sub>2</sub> Levels Do Not Impact Calcification in Mediterranean Cold-Water Corals. PLoS ONE 8(4): e62655. doi:10.1371/journal.pone.0062655

Editor: John Murray Roberts, Heriot-Watt University, United Kingdom

Received October 3, 2012; Accepted March 24, 2013; Published April 30, 2013

Copyright: © 2013 Maier et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Financial support was provided by the European Commission through a Marie-Curie Fellowship to CM (MECCA, project no 220299) and the project COMP via the Prince Albert II of Monaco Foundation, the German Academic Exchange Service (DAAD). This work is a contribution to the 'European Project on Ocean Acidification' (EPOCA) which received funding from the European Community's Seventh Framework Programme FP7/2007–2013) under grant agreement no. 211384. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

\* E-mail: maier.conny@gmail.com

#### Introduction

Ocean acidification is one of the major threats to the marine environment and has become a central research focus in marine science during the last decade. The ocean and atmosphere exchange carbon dioxide  $(CO<sub>2</sub>)$  [1], and the net uptake of  $CO<sub>2</sub>$  by the ocean causes the pH to decline. Due to anthropogenic activity, the oocean pH has already declined by 0.1 units since preindustrial times and will further decline by about 0.4 units until the end of the century [2]. The concentration of  $CO<sub>2</sub>$  in seawater is steadily increasing at an unprecedented rate of change and it is anticipated that atmospheric  $CO<sub>2</sub>$  concentrations increase fourfold between 1750 and 2100, reaching values above 1000 ppm [3]. Many calcifying organisms will be affected by ocean acidification with a decrease of the growth of their shells or skeletons [4–7]. Also, reproduction and larval growth are thought to be impeded [8–10]. For tropical coral reefs, the predictions as to the detrimental effects of ocean acidification are almost unison and foresee a decline in reef growth [11–15] and a shift in species composition with a decrease in diversity [16,17]. On the other hand, it has been shown, that the response to ocean acidification can be highly variable for different taxonomic groups [18]. Investigations on cold-water corals or deep-sea corals are scarce. So far only three experimental studies on the effects of ocean acidification have been published. The rate of net calcification of Lophelia pertusa from Norwegian waters was shown to decline under future  $pCO_2$  conditions [19,20], while no or even a positive longterm (6 months) response of net calcification was observed [20]. A third study investigated the short-term response of net calcification of the Mediterranean coral *Madrepora oculata* under  $pCO<sub>2</sub>$  values reflecting past, pre-industrial and future conditions. No effect on net calcification was found between ambient and elevated  $pCO<sub>2</sub>$ , while calcification rates were twice as high under pre-industrial  $pCO<sub>2</sub>$  than under ambient or elevated  $pCO<sub>2</sub>$ . This suggests that present-day calcification rates of M. oculata have already drastically declined since pre-industrial times [21]. Cold-water corals were initially thought to become affected by ocean acidification before their tropical relatives because they inhabit deeper and colder waters where the aragonite saturation state  $(\Omega_a)$  is lower than in shallower, warmer regions [22]. More than 70% of cold-water coral communities are found in regions that will be undersaturated with respect to aragonite by the end of the century [23]. However, experimental as well as observational evidence has shown that some cold-water corals are able to cope and maintain positive

skeletal growth even in waters undersaturated in  $\Omega_a$  [19,20,24]. Furthermore, the study of Form & Riebesell [20] found a decline of net calcification at elevated  $pCO<sub>2</sub>$  in a short-term experiment and a stimulating effect in a long-term experiment from which they concluded that the cold-water coral L. pertusa is able to acclimate to higher  $pCO<sub>2</sub>$ . In the present study, the question of acclimation is addressed using a thorough experimental design in which net calcification of individual coral samples is repeatedly measured as a function of exposure time to respective  $pCO<sub>2</sub>$  levels. The two branching cold-water coral species present in the Mediterranean Sea, L. pertusa and M. oculata are investigated to discern whether one of the two species is more resistant than the other or has a higher potential for acclimation. In the Mediterranean Sea, M. oculata is more widespread than  $L$ . pertusa [25]. This could indicate that the prevailing conditions in the Mediterranean Sea are more favourable for M. oculata than for L. pertusa and this might result in a divergent species response to changing ocean pH. It is well known, that the distribution of cold-water corals is controlled by temperature within a range of 4 to  $12^{\circ}$ C [26]. In the Mediterranean Sea, the temperature at depths where cold-water corals occur ranges between 12.5 to almost  $14^{\circ}$ C [25,27] which is, therefore, above the common temperature range. Additionally, total alkalinity is higher in the Mediterranean Sea than open oceans and consequently absorbs more atmospheric  $CO<sub>2</sub>$ . It has been shown, that anthropogenic  $CO<sub>2</sub>$  has already affected the Mediterranean Sea and that the pH has decreased by 0.05 to 0.14 pH units, depending on the depth considered, since preindustrial times [28].

#### Materials and Methods

The present study aimed at elucidating (1) the effect of an increase in  $CO<sub>2</sub>$  (ocean acidification) on rates of net calcification on the two Mediterranean cold-water coral species, M. oculata and L. pertusa, (2) to find out whether there is a species effect and (3) to evaluate short- (shock) and longer-term (acclimation) responses in calcification. This has been addressed by measuring net calcification before, immediately after and 1, 2, 3 and 9 months after  $pCO<sub>2</sub>$  has been adjusted.

### Sampling and Experimental Set-up

During the MedSeaCan cruise in June 2009, the cold-water corals L. pertusa and M. oculata were sampled in the canyon of Lacaze Duthiers using a remotely operated vehicle (ROV) and the vessel MINIBEX (COMEX, France). Corals were sampled at water depths of 500 (42°32.98'N, 03°25.21'E), 267 (42°34.98'N, 03°24.15'E) and 260 m (42°35.07'N, 03°24.14'E). On board, corals were maintained in a plastic container  $(1040\times640\times515$  mm) at a controlled temperature of  $12.5\pm0.5^{\circ}$ C using a chilling unit attached to a water pump  $(1000 l h<sup>-1</sup>)$ . Corals were transported back to the laboratory and maintained in a climate room at  $13^{\circ}$ C until the experiments started in August 2009. Coral branches were sub-divided into smaller fragments (Table 1) and placed into vials of 4.5 or 8 cm inner diameter and a volume of ca 300 and 1000 ml. The vials were placed in 4 aquaria that served as water baths  $(13^{\circ}C)$  as well as overflow basins for seawater from the vials. Each vial received running seawater (Mediterranean surface water with a salinity of 38) and air using silicone tubings of 0.5/2.5 and 1.0/3.0 mm inner/outer diameter, respectively. The seawater was filtered through 2 layers of micron bags  $(5 \text{ and } 1 \text{ }\mu\text{m})$  into two 100 l storage tanks in a climate room which was set to  $11^{\circ}$ C. Temperature in the water baths containing the vials was maintained at  $\pm 0.1^{\circ}$ C using electronic temperature controllers

(Corema) and heaters (Tetratec® HT75). Temperature homogeneity was obtained by circulation pumps (JBL Pro Flow 500, 500 l  $h^{-1}$ ). Seawater was distributed by gravity from the storage tanks to the vials at a flow rate of  $32 \pm 14$  ml h<sup>-1</sup>. Air was supplied through a small tube (ca. 8 cm height $\times0.7/1.0$  cm inner/outer diameter) inserted vertically in the vials, preventing the bubbles to be in direct contact with the coral fragments and providing an airlift and mixing. The air was pre-mixed using mass flow controllers (MFCs, ANALYT MC-GFC17, 0–10 l for air and 0–10 ml for pure  $CO<sub>2</sub>$ ) and an air compressor (Jun-Air OF302-25B) at a flow rate of  $4 \times 11$  $min^{-1}$  distributed to  $4\times21$  vials. Corals were fed 3 times a week with freshly hatched Artemia larvae and 1 time a week with frozen krill. The water bath containing the vials and overflowing seawater was also adjusted to the target pCO<sub>2</sub> by bubbling with an air stone (HOBBY ceramic air diffuser, 150 mm). Prior to each feeding the seawater of the water bath with respective  $pCO<sub>2</sub>$  was filtered (Tetratec EX 1200, 1200 l  $h^{-1}$ ) and the strong water flow generated by the filtration unit was used to clean vials and remove old prey and detritus.

#### Determination of Rates of Net Calcification

Before changing the  $pCO<sub>2</sub>$  levels in the experimental set-up, net calcification for each coral fragments was determined using the alkalinity anomaly technique [29] in order to provide an initial control  $(T_0)$  at ambient pCO<sub>2</sub>. Subsequently, the CO<sub>2</sub> concentration of the air used to bubble the vials was adjusted to  $4 \text{ pCO}_2$  treatment levels (A–D) with treatment A at 280 (low), B at 400 (ambient), C at 700 (elevated) and D at 1000 ppm (elevated) using the MFCs. To adjust to lower pCO2 than ambient (treatment A, 280 ppm), soda lime was used to generate low-CO2-air (5–10 ppm) which was mixed with pure CO2. For treatment B ambient air was used and for treatment C (700 ppm) and D (1000 ppm) ambient air was mixed with pure  $CO<sub>2</sub>$ . The exact mixing of air or  $CO<sub>2</sub>$ -free air with pure CO2 was adjusted by a LI-COR CO2-analyser (LI-6252). The pH in the different treatments during coral maintenance was monitored on a weekly base using a commercial pH module and electrode (IKS aquastar) which was calibrated to the NBS standard (buffers 4 and 7). This was done to control that treatment levels remained relatively constant, however, these measurements were not used to assess the carbonate chemistry, which was determined by analysis of  $A_T$  and  $C_T$  (see below for details) 1) during 2-day incubation to determine calcification rates via the alkalinity anomaly technique and 2) in maintenance vials containing corals and in blanks (vials without corals) after 9 months when net calcification rates were established using the buoyant weight technique.

To discriminate between short- and longer-term effects, net calcification was determined immediately after changing the  $pCO<sub>2</sub>$  by transferring coral fragments directly from ambient into incubation vials adjusted to target air- $CO_2$  mix  $(T_1)$  and at about monthly intervals during the first 3 months  $(T_2, T_3, T_4, \text{ with})$ 29, 57 and 89 days of exposure, respectively) using the alkalinity anomaly technique. Additionally, net calcification was again determined after approximately 9 months  $(267 \pm 14$  days) with the buoyant weight technique [30] using a balance (Mettler Toledo) with a precision of 1 mg.

For determination of net calcification rates using the alkalinity anomaly technique, corals and blanks (seawater without corals) were placed for 2 days in the same type of vials than those in which they were maintained but with a constant volume of 200 or 700 ml. To maintain  $pCO<sub>2</sub>$  levels constant vials were aerated with the same air- $CO<sub>2</sub>$  mix that was also used during maintenance periods for treatments A–D. At the end of the incubation, seawater **Table 1.** Initial size (polyp number and skeletal weight) and initial net calcification rates (G) at  $T_0$  (before manipulating pCO<sub>2</sub>) of the corals L. pertusa (LP) and M. oculata (MO) used in repeated incubations of  $pCO<sub>2</sub>$  treatment A–D.



Means  $\pm$  S.D.

doi:10.1371/journal.pone.0062655.t001

was sub-sampled to determine inorganic nutrients, dissolved inorganic carbon  $(C_T)$  and total alkalinity  $(A_T)$  as described in Maier et al. [21]. Other parameters of the carbonate chemistry (pCO<sub>2</sub>, pH on the total scale, pH<sub>T</sub>, and  $\Omega_a$ ) were determined from  $C_T$ ,  $A_T$ , temperature (13<sup>o</sup>C), salinity (38) and hydrostatic pressure (0) using the software package seacarb [32] (Table 2 and S1). Rates of net calcification were calculated from differences in  $A_T$  from blanks (seawater without corals incubated in parallel to coral samples) and coral incubations and were corrected for changes in the concentration of inorganic nutrients. For the incubation at  $T_3$ , no samples for inorganic nutrient concentrations were taken. Therefore, the regression functions between calcification rates corrected (or uncorrected) for inorganic nutrient release (Figure S1) were used to make the nutrient correction for  $T_3$ . Data were normalized to the initial skeletal dry weight of coral fragments and reported in %  $d^{-1}$  [21,31] using the exponential growth function  $G [% d^{-1}] = ((W_n/W_0)^{1/n} - 1) * 100$ ; with  $G =$  net calcification rate,  $W_n =$  Weight after n days,  $W_0 =$  initial weight and d time interval (days) for growth increment (alkalinity anomaly meth $od = 2$  days, buoyant weight method  $= 267$  days).

### Statistical Analysis

Statistical analyses were conducted using the software package Statistica 7.0. A repeated measures analysis of variance (ANOVA) was used. Only coral fragments for which net calcification rates were established at all time steps  $(T_0-T_4)$  and buoyant weight after 9 months) were considered for analysis as this is a pre-requisite for repeated measures testing. Therefore N is the same for all repeated measurements (Table S1). For independent samples (non-repeated measures analyses) a one-way ANOVA comparison of carbonate chemistry between  $pCO<sub>2</sub>$  treatments or initial size and initial net calcification rates  $(T_0)$  was used. If applicable, a post-hoc test was performed using the Honest Significance Difference (HSD) test for either equal or unequal N. The respective statistical tests used are also given in the text, tables and figure legends. Values are given as mean  $\pm$  S.D. unless stated otherwise.

## Results and Discussion

## Initial Carbonate Chemistry, Concentration of Inorganic Nutrients and Calcification

For the initial incubation at ambient pCO<sub>2</sub> (T<sub>0</sub>), the average  $A_T$ and  $C_T$  of blank controls (incubation without corals) was  $2616\pm27$ and 2374 $\pm$ 21 µmol kg<sup>-1</sup> ( $\pm$  standard deviation, throughout this paper;  $N = 30$ . pH<sub>T</sub>, pCO<sub>2</sub> and  $\Omega_a$  were  $8.05 \pm 0.03$ ,  $447.3 \pm 31.7$  µatm and  $2.68 \pm 0.16$ , respectively. The concentrations of phosphate and ammonium were  $0.03\pm0.07$  and  $1.03\pm1.48$  µmol kg<sup>-1</sup> (N = 16). A<sub>T</sub> and C<sub>T</sub> decreased during the 2-day incubations of both coral species;  $A_T$  was on average  $2546 \pm 47$  and  $2571 \pm 32$  µmol kg<sup>-1</sup> and C<sub>T</sub> was  $2333 \pm 39$  and 2356 $\pm$ 41 µmol kg<sup>-1</sup> (N = 23 and 18), respectively. As a consequence, pH<sub>T</sub> and  $\Omega$ <sub>a</sub> were also lower in coral vials than in blanks while pCO<sub>2</sub> increased (Table 2). For both species,  $pH<sub>T</sub>$  was lower by 0.05 units and  $\Omega$ <sub>a</sub> was lower by 0.03, whereas pCO<sub>2</sub> increased by 52 for  $M$ . *oculata* and 60  $\mu$ atm for  $L$ . *pertusa*. Cold-water corals can release significant amounts of inorganic nutrients [33], and phosphate concentrations increased by  $0.413$  and  $0.483 \mu$ mol  $\text{kg}^{-1}$  to 0.44±0.49 and 0.51±0.34 µmol kg<sup>-1</sup>, while ammonium increased by 6.80 and 3.81  $\mu$ mol kg<sup>-1</sup> to 7.83±4.97 and  $4.84\pm5.17$  µmol kg<sup>-1</sup> for M. *oculata* and L. *pertusa* (N = 18 and 22), respectively.

Before changing the  $pCO<sub>2</sub>$  to respective treatment levels A–D (low-high pCO<sub>2</sub>), mean calcification rates were  $0.006 \pm 0.006\%$  $d^{-1}$  for *L. pertusa* (N = 18) and 0.023±0.022%  $d^{-1}$  for *M. oculata*  $(N = 23)$  (Table 1). For both species, the rates of net calcification were significantly correlated with skeletal weight. Smaller coral fragments exhibit higher net calcification rates following a negative logarithmic trend (Figure S2). This negative dependence of size and age for cold-water coral calcification is in accordance with earlier studies [19,21,36]. However, despite the fact that we worked with a relatively large size range, neither size (weight or polyp number), nor initial calcification rates measured under ambient  $pCO<sub>2</sub>$  (Table 1) differed significantly between the coral fragments used in the four  $pCO_2$  treatments (1-way ANOVA,  $p \ge 0.15$ ) and net calcification rates were well within the range of earlier findings [19–21,34,35].

Table 2. Parameters of the carbonate chemistry for the total anomaly technique at  $T_0$  (ambient air, prior to adjusting to respective  $pCO<sub>2</sub>$  levels) and at T<sub>1</sub>-T<sub>4</sub> (immediately and 1, 2 and 3 months after adjusting  $pCO<sub>2</sub>$ , respectively).



Blank is seawater incubated in parallel without coral (N≥4, see Table S1) and used to subtract from A<sub>T</sub> of coral incubations to determine net calcification rates. LP is L. pertusa, MO is M. oculata. A<sub>T</sub> and C<sub>T</sub> (bold) were measured after 2-day incubations and other parameters were calculated from A<sub>T</sub>, C<sub>T</sub>, temperature of 13<sup>°</sup>C and salinity of 38 using the software package seacarb [31]. Values are means  $\pm$  S.D. (N is given in Table 1). doi:10.1371/journal.pone.0062655.t002

## Carbonate Chemistry of Blanks after Adjusting  $pCO<sub>2</sub>$ Levels

After the  $CO_2$  was adjusted to the 4 pCO<sub>2</sub> levels (A–D), the carbonate chemistry changed accordingly (Table 2). For the blanks,  $A_T$  was relatively uniform around 2600 µmol kg<sup>-1</sup> for treatments A to D. The  $C_T$  of blanks increased from  $2313\pm3$  to 2492 $\pm$ 4 µmol kg<sup>-1</sup>, and pH<sub>T</sub> decreased from 8.14 $\pm$ 0.01 to 7.76 $\pm$ 0.01,  $\Omega_{a}$  from 3.2 $\pm$ 0.04 to 1.5 $\pm$ 0.03 and pCO<sub>2</sub> levels ranged from  $349\pm6$  to  $929\pm25$  µatm for treatments A to D, respectively. For time step  $T_1-T_4$ ,  $A_T$  was not significantly different between treatments A–D (One-way ANOVA,  $p>0.2$ ), while it was significant between maintenance vials buoyant weight, treatment B and D, and buoyant weight, treatment D and T4 treatment D. Other parameters of the carbonate chemistry ( $C_T$ , pCO<sub>2</sub>, pH and  $\Omega_a$ ) were in general significantly different between single pCO<sub>2</sub> treatments (One-way ANOVA, Tukey Honest-Significant Difference (HSD) post-hoc test,  $p<0.05$ ) with exceptions for adjacent treatment levels where p-values between treatments were  $>0.05$ . The actual pCO<sub>2</sub> values of seawater differed from those applied by the MFCs and revealed a reduced range between 349 to 929 instead of 280 to

1000 µatm, respectively. This is probably due to the fact that all  $4 \text{ pCO}_2$  treatments were maintained and incubated in the same climate room and mixing of seawater with the overlying air most likely took place due to the vertical water circulation that was generated by the aeration system.

## Carbonate Chemistry and Calcification Rates in Coral Incubation after Adjusting  $pCO<sub>2</sub>$  Levels

For coral incubation of  $T_1-T_4$ , the  $A_T$  of seawater containing L. pertusa was on average  $2496 \pm 45$ , while A<sub>T</sub> for M. oculata incubation was on average  $2520 \pm 21$  (Table 2 and Table S1). The C<sub>T</sub> of *L. pertusa* increased from treatment A–D from  $2236 \pm 23$ to 2492 $\pm$ 4 µmol kg<sup>-1</sup> while C<sub>T</sub> of *M. oculata* increased from  $2256 \pm 11$  to  $2408 \pm 7$  µmol kg<sup>-1</sup> for treatment A to D, respectively. Similar as for the T<sub>0</sub>, the DIC, pH<sub>T</sub> and  $\Omega$ <sub>a</sub> were slightly lower in coral vials than in corresponding blanks while  $pCO<sub>2</sub>$  was higher (Table 2). This means, that also other parameters of the carbonate system than the  $A_T$  changed during the 2-day incubation as a consequence of coral calcification and metabolism. It is evident, that these shifts cannot be avoided as the determination of calcification rates is based on the fact, that the precipitation of 1 mole  $CaCO<sub>3</sub>$  decreases the A<sub>T</sub> by 2 mole and  $C<sub>T</sub>$  by 1 mole [37]. Also, the  $CO<sub>2</sub>$  released by coral respiration and calcification into the surrounding seawater was apparently not completely equilibrated by aeration with respective air- $CO<sub>2</sub>$  gas mix and the  $pCO<sub>2</sub>$  levels of coral vials increased by an average 20 to 40 µatm during the 2-day incubation (Table 2). The  $A_T$ decreased by about 100 µmole relative to blanks, and there was no significant difference of  $A_T$  between pCO<sub>2</sub> treatments or repeated incubation  $T_1$  to  $T_4$  at the end of incubation (Table S2). The excretion of inorganic nutrients by cold-water corals increases the  $A_T$ , i.e. it counteracts the decrease caused by calcification. Thus, calcification rates determined by the alkalinity anomaly technique are an underestimate of actual calcification rates if not corrected for. In a previous study using freshly collected cold-water corals it has been demonstrated that calcification rates of both species would be underestimated by 10% if inorganic nutrients were not taken into account [21]. In the present study, inorganic nutrient concentrations were used to correct calcification rates using the following linear equations:  $G_{L,persusa} = 1.025 * G_{uncorrected} +000019$  $(R = 0.997, N = 81, p << 0.001)$  and  $G_{M.\text{oculata}} = 1.027*G_{\text{uncorrected}}$ +0.00093 ( $R = 0.993$ ,  $N = 102$ ,  $p \ll 0.001$ ) (Figure S1). This means that the ratio of inorganic nutrient release to net calcification was 4-times higher for freshly collected corals than for corals maintained in aquaria.

### Calcification Rates and  $pCO<sub>2</sub>$  Treatment Effects

After the  $pCO<sub>2</sub>$  had been adjusted to the intended treatment levels, calcification rates were measured immediately  $(T_1)$  and 1, 2 and 3 months  $(T_2-T_4)$  after adjusting the pCO<sub>2</sub> levels using the alkalinity anomaly technique and additionally after 9 months using the buoyant weighing technique. Pooled data from repeated measurements (average  $T_1 - T_4$ ) of alkalinity anomaly corresponding to approximately 3-months of coral growth and the buoyant weight after 9-months provided similar results and average calcification rates of L. pertusa slightly varied between treatments A and D from  $0.011 \pm 0.008$  to  $0.017 \pm 0.012\%$  d<sup>-1</sup> for the alkalinity anomaly method and between  $0.010 \pm 0.008$  and  $0.021\pm0.037\%$  d<sup>-1</sup> for the buoyant weight technique (Figure 1, Table S3). For M. oculata average values ranged between  $0.023 \pm 0.012$  and  $0.035 \pm 0.025\%$  d<sup>-1</sup> and between  $0.017\pm0.014$  and  $0.038\pm0.057\%$  d<sup>-1</sup> for the alkalinity anomaly and buoyant weight technique, respectively. For both coral species, there was neither a significant effect between methods used to measure calcification rates (time span of maintenance at respective pCO<sub>2</sub> levels) nor a significant pCO<sub>2</sub> treatment effect or a combined effect of methods (exposure time) and  $pCO<sub>2</sub>$  levels (Repeated measures ANOVA,  $p > 0.05$ , Table S4A). As we used two different methods to establish net calcification rates over midterm and longer-term calcification, we provide a methods comparison for a similar time interval (mid-term) between alkalinity anomaly and buoyant weight showing that the 2 methods provide similar results (SI 1). These data were not included in the manuscript, as they do not comprise the same N, which is mandatory for the repeated measures design used here.

## Short-term (Shock) and Longer-term (Acclimation) Response to Variable  $pCO<sub>2</sub>$

The experimental approach used in the present study was designed to discriminate between short-term ''shock'' effects due to fast changes in  $pCO<sub>2</sub>$  levels and longer-term acclimation response of net calcification. The results from repeated determination of calcification rates over time revealed no short-term, or long-term response in calcification to higher  $pCO<sub>2</sub>$  for either of the two species investigated (Figure 2). There was a continuous increase in



Figure 1. Net calcification rates of a) M. oculata and b) L. pertusa at various pCO<sub>2</sub> levels pooled over  $T_1-T_4$  (total anomaly technique) and from buoyant weight technique spanning 4 or 9 months, respectively. Values are mean  $\pm$  SD, N $\geq$ 4 (Table 1). doi:10.1371/journal.pone.0062655.g001

calcification rates for the coral  $L$ . pertusa in treatment  $D$  which had highest  $pCO<sub>2</sub>$  levels (Figure 1), however this increase was statistically not significant (repeated measures ANOVA, Table S4B). Thus, our data do not support an earlier suggestion for a potential short-term shock response with long-term acclimation of net calcification rates as proposed by Form & Riebesell [20]. Yet, there was an effect of time on net calcification rates for both coral species. In general, average calcification rates (pooled for  $pCO<sub>2</sub>$ treatments) increased until  $T_3$  and then decreased at  $T_4$  again (repeated measures ANOVA, unequal N HSD, Table S4C). This pattern was independent of  $pCO<sub>2</sub>$  treatment and similar for  $L$ . pertusa and M. oculata. This indicates, that factors other than  $pCO<sub>2</sub>$ must have been responsible for the changes in calcification rates with time, that were either driven by intrinsic controls (e.g. reproductive cycles) or a general acclimation to the maintenance conditions in the vials or aquarium (independent of  $pCO<sub>2</sub>$ ).

The lack of  $pCO<sub>2</sub>$  treatment effects seems to contradict other studies reporting on a short-term calcification response of L. pertusa



Figure 2. Net calcification rates (G) of a) M. oculata and b) L. pertusa at  $T_0$ –T4 at the 4 pCO<sub>2</sub> treatment levels (pCO<sub>2</sub> values at different time steps and for the two species are given in Table S1). The pCO<sub>2</sub> at T<sub>0</sub> was at ambient pCO<sub>2</sub> and served as initial control, while T<sub>1</sub>-T<sub>4</sub> was after pCO<sub>2</sub> has been changed to respective treatment levels. Values are mean  $\pm$  SE and SD. doi:10.1371/journal.pone.0062655.g002

to elevated  $pCO<sub>2</sub>$  levels [19,20]. In the following, an attempt is made to reconcile these contradictory findings. Form & Riebesell found a negative correlation between calcification rates and  $CO<sub>2</sub>$ concentration in their short-term experiment. However, this negative correlation might have been due to one very high value for calcification rates at the lowest  $pCO<sub>2</sub>$  which forced the slope to a negative trend while all other values were within a certain range independent of  $pCO_2$  ([20]; Figure 2) similar to the findings of the present study. In the short-term study by Maier et al. [19] there was also a clear negative response to increasing  $pCO<sub>2</sub>$  on net calcification rates. In that study, the experimental set up was different as very small incubation vials (50 ml) and a closed system approach was used. Thus, the initial values for carbonate chemistry were comparable with the present study and that of Form & Riebesell [20], however, the carbonate chemistry changed drastically during incubation and mean  $pCO<sub>2</sub>$  values as high as 2160 µatm were reached ([19]; Table S4). The conclusion with respect to an expected 50% decrease in calcification rates by the

end of the century might thus have been a misinterpretation with respect to relating net calcification rates to the initial  $pCO<sub>2</sub>$  range and not to the much higher  $pCO<sub>2</sub>$  values that were actually reached during incubation due to the closed system approach and small incubation volume.

For M. oculata, so far only one study exists with respect to ocean acidification and it addresses the short-term  $pCO<sub>2</sub>$  effect on net calcification rates [21]. In that study, the  $pCO<sub>2</sub>$  was changed in two ways: 1) to that of pre-industrial concentrations (285 ppm) and 2) to values projected for the end of the century (865 ppm). Similar to the present study, there was no change in net calcification rates between ambient and future  $pCO<sub>2</sub>$  levels, while calcification rates doubled when  $pCO<sub>2</sub>$  was set to values of pre-industrial times indicating that the increase in  $pCO<sub>2</sub>$  that took place since preindustrial times had already a negative impact on M. oculata calcification.

The results of the present study are in contrast to the acclimation hypothesis postulated by Form & Riebesell [20]. Their conclusion with respect to acclimation was based on the fact, that there was a negative short-term response, while in the longterm study higher  $pCO<sub>2</sub>$  had even higher calcification rates than the two lower  $pCO<sub>2</sub>$  levels. However, there is indication that their data can be interpreted differently. First, the long-term experiment of Form & Riebesell [20] lacked the ambient  $pCO<sub>2</sub>$  control and did thus not cover the same  $pCO<sub>2</sub>$  range than their short-term study. However, an ambient  $pCO<sub>2</sub>$  treatment in the long-term study would have been pivotal, specifically because the negative response in the short-term experiment was caused by a higher calcification rate at ambient pCO2. Second, the study by Form & Riebesell [20] also lacked initial controls. It is therefore possible, that the coral fragments used in the high  $pCO<sub>2</sub>$  treatment had already higher initial calcification rates under ambient  $pCO<sub>2</sub>$ conditions. In contrast, the experimental design of the present study comprised both initial and ambient controls and a timeseries for calcification rates which allowed a better evaluation of shock or short-term responses versus long-term acclimation in calcification rates. Due to a lack in short-term response of calcification for the range in  $pCO<sub>2</sub>$  levels studied, our data do not provide evidence that acclimation had played a role in the longterm calcification response to increasing  $pCO<sub>2</sub>$ . This does not mean that no acclimation took place, it rather means that the mechanism(s) enabling cold-water corals to maintain calcification rates constant over a large  $pCO<sub>2</sub>$  range (independent of the duration of the exposure) remain to be identified.

## Summary View of pCO2 Effects on Cold-water Coral Calcification

For a pCO<sub>2</sub> range between 350 and 1000  $\mu$ atm, no effect on net calcification rates, neither for short- nor long-term exposure could be distinguished, and there is evidence that the recently postulated cold-water coral acclimation hypothesis [20] does not hold as such. The present study revealed significant changes observed as function of time for all  $pCO<sub>2</sub>$  treatments, but this must be attributed to other causes than  $pCO<sub>2</sub>$  either related to aquarium conditions or coral biology. Including the previous 3 studies on cold-water coral response to ocean acidification and the results of the present study a certain pattern emerges: the response of coldwater corals L. pertusa and M. oculata to increasing  $pCO<sub>2</sub>$  is nonlinear and net calcification rates remain constant for a  $pCO<sub>2</sub>$  range between ambient and somewhere above 1000  $\mu$ atm where  $\Omega$ <sub>a</sub> is already close to or even below 1. The negative short-term response in the study by Maier et al. (2009) indicated, that once a threshold at high  $pCO<sub>2</sub>$  has been reached, a significant decline in net calcification rates can be expected with increasing  $pCO<sub>2</sub>$  it can

further be assumed that this threshold lies somewhere below 2000  $\mu$ atm. For the Mediterranean coral M. oculata it appears that a 1st threshold had already been surpassed since pre-industrial as indicated by the increase in calcification rates of M. oculata at reduced  $pCO<sub>2</sub>$  [21]. In this respect, the exceptionally high calcification value of L. pertusa at lowest  $pCO_2$  in the short-term experiment of Form & Riebesell [20] might be indicative of such a threshold between present-day and pre-industrial  $pCO<sub>2</sub>$ , however, this needs further investigation.

A non-linear response is contrary to findings for tropical, zooxanthellate corals, that generally reveal a linear, negative calcification response to increasing  $pCO<sub>2</sub>$  [13–15]. However, a non-linear response had already been revealed for temperate zooxanthellate corals which, similar to the Mediterranean coldwater corals, remained unaffected within a large range of presentday to end of the century projections of  $pCO<sub>2</sub>$  [18,38,39] and a drastic reduction in calcification at a  $pCO<sub>2</sub>$  of 2850  $\mu$ atm [40]. The responses of the organisms to increasing ocean acidification are variable and complex [18] and even enhanced calcification at higher  $pCO<sub>2</sub>$  had been proposed for some taxa [41–43].

Up to now it is unclear how the corals are able to resist increasing  $pCO<sub>2</sub>$  levels and how they maintain calcification rates constant over such a large  $pCO<sub>2</sub>$  gradient. It has been proposed that cold-water corals are able to resist increasing ocean acidification by their ability to maintain a high pH within their calicoblastic, calcifying fluid [44]. The way the calcifying fluid is sheltered and replenished with cations from ambient seawater is crucial in how a coral responds to increasing ocean acidification [45–46]. Also, an explanation why temperate corals can resist to higher  $pCO<sub>2</sub>$  levels are their lower growth rates [38] as less carbonate ions will be required in the same time for calcification. This could also explain why cold-water corals grow in waters with an  $\Omega$ <sub>a</sub> around 1, while fast growing tropical corals are found in waters with an  $\Omega_a$  above 3.5. Overall, cold-water corals seem well adapted to low  $\Omega$ <sub>a</sub> which may explain their resistance to increasing  $pCO<sub>2</sub>$  levels up to a certain threshold. Tropical corals can experience drastic short-term changes of  $pCO<sub>2</sub>$  in the close environment due to diurnal changes in  $CO<sub>2</sub>$  uptake and release driven by light-dependent changes in the metabolic functioning of reef organisms [47]. Nothing is known on naturally occurring short-term changes in the seawater carbonate chemistry close to cold-water corals but due to the lack of photosynthetic activity in these depths they are likely less pronounced than in tropical reef systems. Nevertheless the question remains if any naturally occurring short-term changes render cold-water corals resistant to fast changes in  $pCO_2$  and a large range of  $pCO_2$  with values reaching more than 1000 µatm. Finally, it is still questionable if acclimation of calcification to increasing  $pCO<sub>2</sub>$  is a likely scenario in the natural environment. The resistance to increasing  $pCO<sub>2</sub>$ levels and the maintenance of constant calcification requires energy and might thus be sustainable during short-term exposure while energy requirements might not be sustained over longer time of exposure to higher  $pCO<sub>2</sub>$ . This might especially be the case in the natural cold-water coral environment, where regular food supply as usually provided during aquarium maintenance is not always guaranteed and where other stressors such as predation, disease and temperature abnormalities may further impede coral growth.

#### Supporting Information

Figure S1 Calcification rates (G) of M. oculata and L. pertusa corrected and uncorrected for inorganic nutrient reslease during incubation. Correlation analysis were significant with  $R = 0.993$ ,  $N = 102$ ,  $p \lt 0.001$  and  $R = 0.997$ ,  $N = 81$ , p $<< 0.001$  for M. oculata and L. pertusa, respectively. (PDF)

Figure S2 Calcification rates (G) versus skeleletal weight (SW) of M. oculata and L. pertusa under ambient  $pCO<sub>2</sub>$  conditions (T<sub>0</sub>). Logarithmic regressions are significant with  $p = 0.01$  for both species;  $N = 23$  and 18 for M. oculata and L. pertusa, respectively.

(PDF)

Table S1 Parameters of the carbonate chemistry (CC) and the inorganic nutrients  $(IN)$  phosphate  $(PO<sub>4</sub>)$  and ammonium (NH<sub>4</sub>) for the incubation times  $T_0-T_4$  and  $pCO<sub>2</sub>$  treatments A–D after 2-day incubation using the alkalinity anomaly technique. Additionally, CC was established after 9 months (267 days) when net calcification rates were measured using the buoyant weight (BW) technique.  $T_0$  was established prior to adjusting  $pCO<sub>2</sub>$  at ambient for all treatments, while  $T_1 - T_4$  and BW are values derived from 2-day incubations immediately and 1, 2 and 3 months after adjusting  $pCO<sub>2</sub>$  and after 9 months, respectively. LP (Lophelia pertusa), MO (Madrepora oculata) and blank (incubated in parallel without coral) at respective  $pCO<sub>2</sub>$ treatment levels. Values are given as mean  $\pm$  S.D. (PDF)

Table S2 Post-hoc results (Tukey honest-significance difference test for unequal N) of breakdown ANOVA for parameters of the carbonate chemistry  $(A_T, C_T, pCO_2)$ , pH<sub>T</sub>,  $\Omega_a$ ) for the coral incubations with T<sub>1</sub>–T<sub>4</sub> (directly and 1, 2, 3 months after  $pCO<sub>2</sub>$  was changed to respective treatment levels) established during 2-day incubation using alkalinity anomaly and after 9 months (267 days) using buoyant weight (BW) technique to determine net calcification rates. Matrix of p-values for L. pertusa at upper right and for *M. oculata* at lower left part of the table with  $p<0.05$ (italic). Bold values are corresponding treatments at different incubation times.

## (PDF)

#### References

- 1. IPPC (2007) Climate change 2007: the physical science basis. Cambridge University Press, Cambridge, UK.
- 2. Orr JC (2011) Recent and future changes in ocean carbonate chemistry. In Gattuso, J-P & Hansson, L, editors. Ocean Acidification. Oxford University Press, Oxford. 41–66.
- 3. Plattner G-K, Joos F, Stocker TF, Marchal O (2001) Feedback mechanisms and sensitivities of ocean carbon uptake under global warming. Tellus 53B: 564–592.
- 4. Riebesell U, Zondervan I, Rost B, Tortell PD, Zeebe RE, et al. (2000) Reduced calcification of marine plankton in response to increased atmospheric CO2. Nature 407: 364–367.
- 5. Fabry VJ (2008) Marine calcifiers in a high-CO<sub>2</sub> ocean. Science 320: 1020, doi: 10.1126/science.1157130.
- 6. Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M, et al. (2008) Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. Nature 454: 96–99.
- 7. Martin S, Gattuso J-P (2009) Response of Mediterranean coralline algae to ocean acidification and elevated temperature. Global Change Biol 15: 2089– 2100.
- 8. Albright R (2011) Reviewing the effects of ocean acidification on sexual reproduction and early life history stages of reef-building corals. J Mar Biol doi:10.1155/2011/473615, 14pp.
- 9. Gazeau F, Gattuso J-P, Dawber C, Pronker AE, Peene F, et al. (2010) Effect of ocean acidification on the early life stages of the blue mussel Mytilus edulis. Biogeosciences 7: 2051–2060.
- 10. Doropoulos C, Ward S, Diaz-Pulido G, Hoegh-Guldberg O, Mumby PJ (2012) Ocean acidification reduces coral recruitment by disrupting intimate larval-algal settlement interactions. Ecol Lett 15: 338–346, doi: 10.1111/j.1461- 0248.2012.01743.x.
- 11. Kleypas JA, Buddemeier RW, Gattuso J-P (2001) The future of coral reefs in an age of global change. Int J Earth Sciences 90: 426–437.

Table S3 Average calcification rates (G) of the coldwater corals Lophelia pertusa (LP) and Madrepora oculata (MO) at  $T_0$  to  $T_4$  determined by the total anomaly technique in monthly time intervals and 2 days of incubation, and calcification rates determined by buoyant weight (G(BW)) after maintenance of corals for 9–10 months under respective  $pCO<sub>2</sub>$  treatment levels. (PDF)

Table S4 Statistical results for repeated measures ANOVA of calcification rates (G). A Comparison between total alkalinity (TA) method (average G, pooled T1–T4) and Buoyant Weighting (BW) for  $pCO_2$  treatments A–D; and **B** for comparison of repeated measurements  $T_0-T_4$  for the 4 pCO<sub>2</sub> treatments. Table C gives the matrix for p-values of the Tukey-Honest-Significance post-hoc comparison for unequal N of the variable R1 ( $T_0-T_4$ ) for M. oculata (lower left) and L. pertusa (upper right). Significant p are marked in bold, italic (PDF)

File S1 Method comparison between total alkalinity anomaly (TAA) and buoyant weight (BW) technique to establish net calcification rates  $(G)$  of  $M$ . oculata and  $L$ . pertusa.

#### (PDF)

#### Acknowledgments

We thank captain, crew and scientific shipboard staff of R/V Minibex (COMEX). Cruises on RV Minibex were part of a project on canyons of the Mediterranean Sea (MedSeaCan) lead by the French Marine Protected Areas Agency (AAMP), France. Thanks to F. Bils for help with coral maintenance and experiments and S. Comeau for assistance in analysis of  $A_T$ .

### Author Contributions

Conceived and designed the experiments: CM MW JPG PW. Performed the experiments: CM AS MMBS. Analyzed the data: CM AS MMBS. Contributed reagents/materials/analysis tools: CM JPG MW PW. Wrote the paper: CM.

- 12. Kleypas JA, Yates KK (2009) Coral reefs and ocean acidification. Oceanography 22: 108–117.
- 13. Langdon C, Atkinson MJ (2005) Effect of elevated  $pCO_2$  on photosynthesis and calcification of corals and interactions with seasonal change in temperature/ irradiance and nutrient enrichment. J Geophys Res 110: C09S07, doi:10.1029/ 2004JC002576.
- 14. De'ath G, Lough JM, Fabricius KE (2009) Declining coral calcification on the Great Barrier Reef. Science 323: 116–119.
- 15. Gattuso J-P, Allemand D, Frankignoulle M (1999) Photosynthesis and calcification at cellular, organismal and community levels in coral reefs: A review on interactions and control by carbonate chemistry. Amer Zool 39: 160– 183.
- 16. Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, et al. (2007) Coral reefs under rapid climate change and ocean acidification. Science 318: 1737, doi: 10.1126/science.1152509.
- 17. Fabricius KE, Langdon C, Uthicke S, Humphrey C, Noonan S, et al. (2011) Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations. Nature Climate Change 1: 165–169, doi:10.1038/nclimate1122.
- 18. Ries JB, Cohen AL, McCorkle DC (2009) A nonlinear calcification response to  $CO<sub>2</sub>$ -induced ocean acidification by the coral *Oculina arbuscula*. Geology 37: 1057–1152.
- 19. Maier C, Hegeman J, Weinbauer MG, Gattuso J-P (2009) Calcification of the cold-water coral Lophelia pertusa under ambient and reduced pH. Biogeosciences 6: 1671–1680.
- 20. Form AU, Riebesell U (2012) Acclimation to ocean acidification during longterm CO2 exposure in the cold-water coral Lophelia pertusa. Global Change Biol 18: 843–853, doi: 10.1111/j.1365-2486.2011.02583.x.
- 21. Maier C, Watremez P, Taviani M, Weinbauer MG, Gattuso J-P (2012) Calcification rates and the effect of ocean acidification on Mediterranean coldwater corals. Proc. R. Soc. Lond., B 279: 1713–1723, doi: 10.1098/ rspb.2011.1763.
- 22. Orr JC, Fabry VJ, Aumont O, Bopp L, Doney SC, et al. (2005) Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. Nature 437: 681–686.
- 23. Guinotte JM, Orr J, Cairns S, Freiwald A, Morgan L, et al. (2006) Will humaninduced changes in seawater chemistry alter the distribution of deep-sea scleractinian corals? Front Ecol Environ 4: 141–146.
- 24. Thresher RE, Tilbrook B, Fallon S, Wilson NC, Adkins J (2011) Effects of chronic low carbonate saturation levels on the distribution, growth and skeletal chemistry of deep-sea corals and other seamount megabenthos. Mar Ecol Prog Ser 442: 87–99.
- 25. Freiwald A, Beuck L, Rüggeberg A, Taviani M, Hebbeln D (2009) The white coral community in the central Mediterranean Sea revealed by ROV surveys. Oceanography 22: 36–52.
- 26. Roberts JM, Wheeler AJ, Freiwald A (2006) Reefs of the deep: The biology and geology of cold-water coral ecosystems. Science 312: 543–547.
- 27. Taviani M, Freiwald A, Zibrowius H (2005) Deep coral growth in the Mediterranean Sea: an overview. In: Freiwald A & Roberts JM editors. Coldwater corals and ecosystems. Springer-Verlag, Berlin Heidelberg. 137–156.
- 28. Touratier F, Goyet C (2011) Impact of the Eastern Mediterranean Transient on the distribution of anthropogenic  $CO<sub>2</sub>$  and first estimate of acidification for the Mediterranean Sea. Deep Sea Res I 58: 1–15.
- 29. Chisholm JRM, Gattuso J-P (1991) Validation of the alkalinity anomaly technique for investigating calcification and photosynthesis in coral reef communities. Limnol Oceanogr 36: 1232–1239.
- 30. Davies PS (1989) Short-term growth measurements of corals using an accurate buoyant weighing technique. Mar Biol 101: 389–395.
- 31. Reynaud S, Ferrier-Page`s C, Meibom A, Mostefaoui S, Mortlock R, et al. (2007) Light and temperature effects on Sr/Ca and Mg/Ca ratios in the scleractinian coral Acropora sp. Geochim. Cosmochim Acta 71 (2): 354–362.
- 32. Lavigne H, Gattuso J-P (2011) seacarb: seawater carbonate chemistry with R. R package version 2.4. Available: http://CRAN.R-project.org/package = seacarb. Accessed 9 September 2012.
- 33. Maier C, Kluijver AD, Agis M, Brussaard CPD, van Duyl FC, et al. (2011) Dynamics of nutrients, total organic carbon, prokaryotes and viruses in onboard incubations of cold-water corals. Biogeosciences 8: 2609–2620.
- 34. Orejas C, Ferrier-Pagès C, Reynaud S, Gori A, Beraud E, et al. (2011) Longterm growth rates of four Mediterranean cold-water coral species maintained in aquaria. Mar Ecol Prog Ser 429: 57–65.
- 35. Orejas C, Gori A, Gili JM (2008) Growth rates of live Lophelia pertusa and Madrepora oculata from the Mediterranean Sea maintained in aquaria. Coral Reefs 27: 255.
- 36. Mortensen PB (2001) Aquarium observations on the deep-water coral Lophelia pertusa (L., 1758) (Scleractinia) and selected associated invertebrates. Ophelia 54: 83–104.
- 37. Wolf-Gladrow DA, Zeebe RE, Klaas C, Körtzinger A, Dickson AG (2007) Total alkalinity: The explicit conservative expression and its application to biogeochemical processes. Deep Sea Res I 106: 287–300.
- 38. Rodolfo-Metalpa R, Martin S, Ferrier-Pagès C, Gattuso J-P (2010) Response of the temperate coral *Cladocora caespitosa* to mid-and long-term exposure to  $pCO<sub>2</sub>$ and temperature levels projected for the year 2100 AD. Biogeosciences 7: 289– 300.
- 39. Rodolfo-Metalpa R, Houlbrèque F, Tambutté É, Boisson F, Baggini C, et al. (2011) Coral and mollusc resistance to ocean acidification adversely affected by warming. Nature Climate Change 1: 308–311.
- 40. Ries JB, Cohen AL, McCorkle DC (2010) A nonlinear calcification response to  $CO<sub>2</sub>$ -induced ocean acidification by the coral Oculina arbuscula. Coral Reefs 29: 661–674.
- 41. Fiorini S, Middelburg JJ, Gattuso J-P (2011) Testing the effects of elevated pCO<sub>2</sub> on coccolithophores (Prymnesiophyceae): Comparison between haploid and diploid life stages. J Phycol: doi:10.1111/j.1529-8817.2011.01080.x.
- 42. Iglesias-Rodriguez MD, Halloran PR, Rickaby REM, Hall IR, Colmenero-Hidalgo E, et al. (2008) Phytoplankton calcification in a high-CO<sub>2</sub> world. Science 320: 336–340.
- 43. Wood HL, Spicer JI, Widdicombe S (2008) Ocean acidification may increase calcification rates, but at a cost. Proc R Soc Lond B 275: 1767–1773.
- 44. McCulloch M, Trotter J, Montagna P, Falter J, Dunbar R, et al. (2012) Resilience of cold-water scleractinian corals to ocean acidification: Boron isotopic systematics of pH and saturation state up-regulation. Geochim Cosmochim Acta 87: 21–34; doi:10.1016/j.gca.2012.03.027.
- 45. Tambutté S, Holcomb M, Ferrier-Pagès C, Reynaud S, Tambutté E, et al. (2011) Coral biomineralization: From the gene to the environment. J exp mar Biol Ecol 408: 58–78.
- 46. Gagnon AC, Adkins JF, Erez J (2012) Seawater transport during coral biomineralization. Earth Planet Sci 329–330: 150–161.
- 47. Gattuso J-P, Frankignoulle M, Smith SV (1999) Measurement of community metabolism and significance in the coral reef  $CO_2$  source-sink debate. Proc Natl Acad Sci 96: 13017–13022.