



HAL
open science

Semiparametric reference curves for EDEN cohort

Sandie Ferrigno

► **To cite this version:**

Sandie Ferrigno. Semiparametric reference curves for EDEN cohort. CMStatistics 2021, Dec 2021, Londres, United Kingdom. hal-03507796

HAL Id: hal-03507796

<https://hal.science/hal-03507796>

Submitted on 3 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Semiparametric reference curves for EDEN cohort

SANDIE FERRIGNO

Inria team BIGS Nancy - Grand-Est, Lab IECL, Université de Lorraine, France

OBJECTIVES

In medicine, **reference curves** [1] [2] [6] are an important tool for clinical practice to screen for fetuses that may be of abnormal size for gestational age.

Data are from the French **EDEN** mother-child cohort [3]. As measures of child growth, we are studying fetal weight and height that depends on the gestational age, in the third trimester of mother's pregnancy.

Semi-parametric methods as **LMS methods** [4][1] and Generalized Additive Model for Location, Scale and Shape (**GAMLSS**) [5] R package are used to construct and implement these curves.

LMS METHODS

The LMS method was developed by Cole [2] and Cole and Green [1]. Let the positive random variable Y (weight or height) be defined through the transformed random variable Z given by

$$Z = \begin{cases} \frac{1}{\sigma\nu} \left[\left(\frac{Y}{\mu} \right)^\nu - 1 \right], \nu \neq 0 \\ \frac{1}{\sigma} \log \left(\frac{Y}{\mu} \right), \nu = 0 \end{cases},$$

with $Y > 0$, $\mu > 0$, $\sigma > 0$ and $-\infty < \nu < \infty$. Z is assumed to follow a truncated standard normal distribution. The three parameters μ (location parameter), σ (scale parameter) and ν (skewness parameter) depend on the weeks of amenorrhea (GW) through smoothing functions $\mu = h_1(GW)$, $\log \sigma = h_2(GW)$, $\nu = h_3(GW)$ with $h_1(\cdot)$, $h_2(\cdot)$ and $h_3(\cdot)$ nonparametric smoothing functions as cubic splines. The distribution of Y is given by

$$Y \sim D(\mu, \sigma, \nu)$$

where the distribution D represent the Box-Cox Cole and Green (BCCG) distribution. Extensions of this method [5], as Box-Cox power exponential (BCPE) and Box-Cox t (BCT) are also investigated.

EDEN COHORT

EDEN [3] is a mother-child cohort study investigating the prenatal and early postnatal determinants of child health and development. 2002 pregnant women were recruited before 24 weeks of amenorrhoea in two maternity clinics from France and from may 2003 to september 2006, 1899 newborns were then included. The main outcomes of interest are fetal and postnatal growth, adiposity development, respiratory health, atopy, behaviour and bone, cognitive and motor development.

Our interest focuses on the study of the variables weight (in grammes) and height (in centimeters)

of fetuses according to the weeks of amenorrhea (GW). The data are not exact for weeks of amenorrhea less than 28 because we have few births before this date. The calculation of statistics on the data also requires to start working from 32 weeks of amenorrhea so that the data are in sufficient number. We observe that the majority of these fetuses are born for an age between 38 and 41 GW.

We acknowledge the commitment of the EDEN mother-child cohort study group and more particularly Barbara Heude which allowed us to use the data from the cohort.

DATA AND CENTILES




Figure 1: Weight for male and female fetuses from 28 to 42 gestational weeks.




Figure 2: Height for male and female fetuses from 28 to 42 gestational weeks.

Centiles are usually chosen from a symmetric subset of the 3rd, 10th, 50th, 90th and 97th.

REFERENCE CURVES FOR WEIGHT




Figure 3: Centiles curves using the Box-Cox and Green (BCCG) distribution for the male and female fetuses weight data and cubic splines smoothing.




Figure 4: Centiles curves using the Box-Cox power exponential (BCPE) distribution for the male and female fetuses weight data and cubic splines smoothing.




Figure 5: Centiles curves using the Box-Cox t (BCT) distribution for the male and female fetuses weight data and cubic splines smoothing.

REFERENCE CURVES FOR HEIGHT




Figure 6: Centiles curves using the Box-Cox and Green (BCCG) distribution for the male and female fetuses height data and cubic splines smoothing.




Figure 7: Centiles curves using the Box-Cox power exponential (BCPE) distribution for the male and female fetuses height data and cubic splines smoothing.




Figure 8: Centiles curves using the Box-Cox t (BCT) distribution for the male and female fetuses height data and cubic splines smoothing.

REFERENCES

- [1] T. J. Cole and P. J. Green. Smoothing reference centile curves : the lms method and penalized likelihood. *Journal of the Royal Statistical Society*, pages 1305–1319, 1992.
- [2] T. J. Cole. Fitting smoothed centile curves to reference data. *Journal of the Royal Statistical Society*, pages 385–418, 1988.
- [3] B. Heude *et al.* Cohort profile: The eden mother-child cohort on the prenatal and early postnatal determinants of child health and development. *International Journal of Epidemiology*, pages 353–363, 2016.
- [4] E. O. Ohuma and D. G. Altman. Statistical methodology for constructing gestational age-related charts using cross-sectional and longitudinal data : The intergrowth-21 project as a case study. *Statistics in Medicine*, pages 1–20, 2018.
- [5] M D. Stasinopoulos *et al.* *Flexible regression and smoothing*. Chapman & Hall Book, 2017.
- [6] P. Royston and E. M. Wright. How to construct 'normal ranges' for fetal variables. *Ultrasound Obstet Gynecol*, pages 30–38, 1998.