



HAL
open science

Unbounded-energy solutions to the fluid+disk system and long-time behavior for large initial data

Guillaume Ferriere, Matthieu Hillairet

► **To cite this version:**

Guillaume Ferriere, Matthieu Hillairet. Unbounded-energy solutions to the fluid+disk system and long-time behavior for large initial data. 2022. hal-03511566v1

HAL Id: hal-03511566

<https://hal.science/hal-03511566v1>

Preprint submitted on 5 Jan 2022 (v1), last revised 8 Aug 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Unbounded-energy solutions to the fluid+disk system and long-time behavior for large initial data.

GUILLAUME FERRIERE AND MATTHIEU HILLAIRET

Institut de Recherche Mathématique Avancée, UMR 7501 Université de Strasbourg et CNRS, France,
guillaume.ferriere@math.unistra.fr

IMAG, Univ Montpellier, CNRS, Montpellier, France, matthieu.hillairet@umontpellier.fr

Abstract

In this paper, we analyse the long-time behavior of solutions to a coupled system describing the motion of a rigid disk in a 2D viscous incompressible fluid. Following previous approaches in [4, 13, 14] we look at the problem in the system of coordinates associated with the center of mass of the disk. Doing so, we introduce a further nonlinearity to the classical Navier Stokes equations. In comparison with the classical nonlinearities, this new term lacks time and space integrability, thus complicating strongly the analysis of the long-time behavior of solutions.

We provide herein two refined tools : a refined analysis of the Gagliardo-Nirenberg inequalities and a thorough description of fractional powers of the so-called fluid-structure operator [2]. On the basis of these two tools we extend decay estimates obtained in [4] to arbitrary initial data and show local stability of the Lamb-Oseen vortex in the spirit of [7, 8].

1. INTRODUCTION

In this paper, we pursue the studies on the long-time behavior of solutions to the following model for the motion of a rigid disk inside a viscous incompressible fluid:

$$\frac{\partial u}{\partial t} + (u \cdot \nabla)u - \nu \Delta u + \nabla p = 0 \quad \text{for } x \in \mathcal{F}(t), \quad (1.1)$$

$$\operatorname{div} u = 0 \quad \text{for } x \in \mathcal{F}(t), \quad (1.2)$$

$$u(t, x) = h'(t) + \omega(t)(x - h(t))^\perp \quad \text{for } x \in \partial B(t), \quad (1.3)$$

$$mh''(t) = - \int_{\partial B(t)} \Sigma(u, p)n \, d\sigma(x) \quad (1.4)$$

$$\mathcal{J}\omega'(t) = - \int_{\partial B(t)} (x - h(t))^\perp \cdot \Sigma(u, p)n \, d\sigma(x). \quad (1.5)$$

Here $u \in \mathbb{R}^2$ and $p \in \mathbb{R}$ stand for the velocity-field/pressure unknowns describing the behavior of a homogeneous incompressible viscous fluid. The rigid solid disk occupies the domain $B(t) := B(h(t), 1)$ and its motion is described by a translation velocity $\ell = h'(t)$ and a rotation velocity ω . Doing so, we prescribe the evolution of the fluid+disk system by integrating the incompressible Navier-Stokes equations (1.1)-(1.2) in the fluid domain $\mathcal{F}(t) := \mathbb{R}^2 \setminus \overline{B(t)}$ and the Newton equation of solid dynamics (1.4)-(1.5). We emphasize that the motion of the fluid and the solid are both unknowns. The system is complemented with no-slip interface conditions (1.3) and transmission of normal stress. The stress tensor $\Sigma(u, p)$ appearing then in the Newton laws is the fluid stress tensor

$$\Sigma(v, p) = -p\operatorname{Id} + 2\nu D(v),$$

with

$$D(v)_{i,j} = \frac{1}{2} \left(\frac{\partial v_i}{\partial x_j} + \frac{\partial v_j}{\partial x_i} \right), \quad 1 \leq i, j \leq 2.$$

We remind that $\nu > 0$ stands for the fluid viscosity and that, due to the incompressibility condition, the viscous operator appearing in (1.1) reads:

$$\nu \Delta u - \nabla p = \operatorname{div} \Sigma(u, p)$$

where, by convention, the divergence operator of a matrix is computed row-wise. By scaling arguments, we prescribed that the density of the fluid is constant equal to 1 and that the solid has radius 1. Below, it appears also that the viscosity ν has only an influence through a time-scaling so we fix $\nu = 1$ for simplicity. The quantity m and \mathcal{J} appearing in the Newton laws represent respectively the mass and inertia of the solid disk. In the 2D case under consideration here, the inertia \mathcal{J} is time-independent. The symbol n appearing in the integrals of (1.4)-(1.5) stands for the normal to $\partial B(t)$ inward $B(t)$. We keep the convention that the normal is directed outward the fluid domain throughout the paper. Like in [4], our motivation for studying this system is to analyse the energy exchange between the solid body and the rigid disk, we do not include any forcing term such as gravity in the system.

Systems like (1.1)–(1.5) coupling ODEs and PDEs and describing the motion of solid bodies inside a viscous fluid have been the subject of numerous studies in the past years. Regarding the specific case of one rigid disk in an unbounded viscous fluid, the Cauchy theory for finite-energy initial data is studied in [13]. The authors remark therein that solutions to (1.1)–(1.5) satisfy the *a priori* estimate:

$$\frac{1}{2} \left[m|\ell(t)|^2 + \mathcal{J}\omega(t)^2 + \int_{\mathcal{F}(t)} |u(t, \cdot)|^2 \right] + \int_0^t \int_{\mathcal{F}(s)} |D(u)|^2 = \frac{1}{2} \left[m|\ell(0)|^2 + \mathcal{J}\omega(0)^2 + \int_{\mathcal{F}(0)} |u(0, \cdot)|^2 \right]$$

This opens the way to the construction of global-in-time finite-energy solution for arbitrary data. To this purpose, the authors operate the change of unknowns:

$$v(t, x) = u(t, x - h(t)), \quad \tilde{p} = p(t, x - h(t)), \quad \ell(t) = h'(t), \quad \omega(t) = \omega(t). \quad (1.6)$$

and obtain the new system:

$$\frac{\partial v}{\partial t} + ((v - \ell(t)) \cdot \nabla)v - \Delta v + \nabla \tilde{p} = 0 \quad \text{for } x \in \mathcal{F}_0, \quad (1.7)$$

$$\operatorname{div} v = 0 \quad \text{for } x \in \mathcal{F}_0, \quad (1.8)$$

$$v(t, x) = \ell(t) + \omega(t) x^\perp \quad \text{for } x \in \partial B_0, \quad (1.9)$$

$$m\ell'(t) = - \int_{\partial B_0} \Sigma(v, \tilde{p})n \, d\sigma(x) \quad (1.10)$$

$$\mathcal{J}\omega'(t) = - \int_{\partial B(t)} x^\perp \cdot \Sigma(v, \tilde{p})n \, d\sigma(x), \quad (1.11)$$

where $B_0 = B(0, 1)$ and $\mathcal{F}_0 = \mathbb{R}^2 \setminus \overline{B_0}$. With this change of unknowns, we have now a problem in a fixed geometry that we can complete prescribing an initial condition. Setting an initial time $t_0 \geq 0$ that can be strictly positive, this condition reads:

$$v|_{t=t_0} = v_0 \quad \text{for } x \in \mathcal{F}_0, \quad \ell(t_0) = \ell_0, \quad \omega(t_0) = \omega_0. \quad (1.12)$$

Despite (1.7)-(1.11) is an autonomous problem, we introduce here a generalized Cauchy-problem with arbitrary initial time. This will have an influence below because of our choice for initial data. We recall here also that the pressure \tilde{p} can be seen as the Lagrange-multiplier of the divergence-free condition involved in the system above. For this reason, there is no initial condition on \tilde{p} . In our formalism, the pressure will also be a secondary unknown that is taken rid *via* a projector argument and that can be recovered *a posteriori*. For all these reasons, we state our results in terms of (v, ℓ, ω) only. For instance, in [13], the authors consider the case $t_0 = 0$. They consider initial data $(v_0, \ell_0, \omega_0) \in L^2(\mathcal{F}_0) \times \mathbb{R}^2 \times \mathbb{R}$ such that :

$$\operatorname{div} v_0 = 0 \text{ in } \mathcal{F}_0 \quad v_0 \cdot n = (\ell_0 + \omega_0 x^\perp) \cdot n \text{ on } \partial B_0 \quad (1.13)$$

and construct global-in-time finite-energy solutions in the sense that:

- $v \in C([0, \infty); L^2(\mathcal{F}_0) - w)$ with $\nabla v \in L^2((0, \infty); L^2(\mathcal{F}_0))$

- $(\ell, \omega) \in C([0, \infty); \mathbb{R}^3)$
- (v, ℓ, ω) solve (1.7)-(1.12) with the associated *a priori* estimate.

$$\frac{1}{2} \left[m|\ell(t)|^2 + \mathcal{J}\omega(t)^2 + \int_{\mathcal{F}_0} |v(t, \cdot)|^2 \right] + \int_{\mathcal{F}_0} |D(v)|^2 = \frac{1}{2} \left[m|\ell(0)|^2 + \mathcal{J}\omega(0)^2 + \int_{\mathcal{F}_0} |v(0, \cdot)|^2 \right]$$

The results are extended to L^q -initial data in [4].

Solutions to the Cauchy-problem are constructed via a perturbative approach. First the authors consider the linearized system:

$$\frac{\partial v}{\partial t} - \Delta v + \nabla \tilde{p} = 0 \quad \text{for } t \in (0, \infty), x \in \mathcal{F}_0, \quad (1.14)$$

$$\operatorname{div} v = 0 \quad \text{for } t \in (0, \infty), x \in \mathcal{F}_0, \quad (1.15)$$

$$v(t, x) = \ell(t) + \omega(t) x^\perp \quad \text{for } t \in (0, \infty), x \in \partial B_0, \quad (1.16)$$

$$m\ell'(t) = - \int_{\partial B_0} \Sigma(v, \tilde{p}) n \, d\sigma(x) \quad \text{for } t \in (0, \infty), \quad (1.17)$$

$$\mathcal{J}\omega'(t) = - \int_{\partial B(t)} x^\perp \cdot \Sigma(v, \tilde{p}) n \, d\sigma(x) \quad \text{for } t \in (0, \infty), \quad (1.18)$$

They show that this system can be rewritten into an infinite-dimensional differential system

$$\partial_t V - AV = 0 \quad (1.19)$$

by constructing an unknown V encoding simultaneously (v, ℓ, ω) and a specific unbounded operator A (that we call fluid-structure operator following [2]). We give more details on these constructions in the next section. Finite-energy solutions to the linearized system are obtained by remarking that A is an accretive positive self-adjoint operator which implies the existence of a contraction semigroup $(S(t))_{t>0}$ solving (1.19). The nonlinear system can be then interpreted in the form of a nonlinear infinite-differential system:

$$\partial_t V - AV = F(V)$$

and mild-solutions are constructed via a Kato-type argument. Since these mild-solutions are finite-energy solutions and finite-energy solutions are unique, this yields "the" finite-energy solution. Actually, this argument is performed on regularized H^1 initial data in [13] (and finite-energy solutions are obtained then by a compactness argument). But, as we shall see below (see **Theorem 3.4**), the reasoning extends to L^2 initial data.

The long-time behavior of solutions to (1.7)-(1.12) is tackled in [4] by the second author in collaboration with S. Ervedoza and C. Lacave. Firstly the properties of the fluid-structure semigroup $(S(t))_{t \geq 0}$ are studied in a non-Hilbert setting which yields explicit bounds for the large-time decay of L^q -initial data and an explicit first order term for sufficiently localized initial data. *Via* perturbative arguments, these decay rates are extended to the finite-energy solutions to the full nonlinear problem (1.7)–(1.12) for initial data such that $v_0 \in L^q(\mathcal{F}_0) \cap L^2(\mathcal{F}_0)$ for some $q \in (1, 2)$ with $\|v_0\|_{L^2(\mathcal{F}_0)} + |\ell_0| + |\omega_0|$ sufficiently small (depending only on q).

In this paper we pursue the computations of [4] in two directions. Firstly, we extend the decay-rate computation of finite-energy solutions to (1.7)–(1.12) for arbitrary data in $L^q \cap L^2$. Namely, our first result reads:

Theorem 1.1. *Let $q \in (1, 2)$ and assume that $t_0 = 0$ and that the initial data $(v_0, \ell_0, \omega_0) \in L^2(\mathcal{F}_0) \times \mathbb{R}^2 \times \mathbb{R}$ satisfy the compatibility condition (1.13) and the further condition $v_0 \in L^q(\mathcal{F}_0)$. Then, the unique finite-energy solution (v, ℓ, ω) of (1.7)-(1.12) satisfies:*

$$\sup_{t>0} t^{\frac{1}{p} - \frac{1}{q}} \|v(t)\|_{L^p(\mathcal{F}_0)} < \infty \quad \forall p \in (2, \infty) \quad (1.20)$$

$$\sup_{t>0} t^{\frac{1}{q}} |\ell(t)| < \infty \quad (1.21)$$

This result must be compared with [4, Theorem 1.3] where a further smallness is required. We point out that, like in [4], our result states that the decay of solutions to the nonlinear problem (1.7)-(1.12) is the same as the decay of solutions of the linearized system (1.14)-(1.18). However, we are still not able to extract a leading term for the nonlinear system.

The proof of this first result is based on adapting the global stability argument in [7]. Namely, we use that the fluid-structure operator A underlying the resolution of the linearized problem (1.14)-(1.18) is self-adjoint and positive. We can then construct the fractional powers A^μ for $\mu \in (-1, 1)$ and analyze their ranges and domains. To extract a decay of any solution to (1.7)-(1.12), we first compute an energy estimates on $U = A^{-\mu}V$ for a μ adapted to the integrability of the initial data v_0 . One key new difficulty is that the nonlinearities in (1.7)-(1.12) involve the term $\ell \cdot \nabla v$. It turns out that handling this term requires to prove a similar time-integrability of ℓ as the one of ∇v and in particular that $\ell \in L^2((0, \infty))$. This property is obtained in a first independent step.

In a second direction, we also extend the analysis to infinite energy initial data. Indeed, similarly to the introductory remark of [7] in the case of a still particle, one may observe that the total amount of the fluid vorticity $\omega := \partial_2 v_1 - \partial_1 v_2$ in solutions to (1.7)-(1.11) has to vanish. This property fails however in many contexts. We recall that, in the absence of a disk, a central object is the normalized Lamb-Oseen vortex:

$$\Theta(t, x) = \frac{1}{2\pi} \frac{x^\perp}{|x|^2} \left(1 - e^{-\frac{|x|^2}{4(1+t)}}\right), \quad x \in \mathbb{R}^2 \setminus \{(0, 0)\}, \quad t \geq 0, \quad (1.22)$$

since any solution to the Navier Stokes equations on \mathbb{R}^2 converges to a multiple of this profile given by the initial mass of the vorticity [8]. This result is extended to the Navier Stokes equations outside a still obstacle [7] showing that any bounded-energy perturbation of a small Lamb-Oseen vortex behaves in large-time like the Lamb-Oseen vortex.

We consider herein the local stability of the Lamb-Oseen vortex Θ in the case of the full fluid+disk problem (1.7)-(1.11). For this, we first see that Θ can be written under the form $\Theta(t, x) = g(t, |x|^2) x^\perp$, where

$$g(t, r) = \frac{1 - e^{-\frac{r}{4(1+t)}}}{2\pi r}.$$

Hence, the Lamb-Oseen vortex on ∂B_0 is a pure rotation. We can then assume initial data are of the form

$$v_0 = \alpha \Theta(t_0, \cdot) + w_0 \quad \ell_w^0 = \ell_w^0 \quad \omega_0 = \frac{\alpha}{2\pi} (1 - \exp(-1/4(1 + t_0))) + \omega_w^0 \quad (1.23)$$

where w_0 is localized in space and

$$w_0 = \ell_w^0 + \omega_w^0 x^\perp \text{ on } \partial B_0. \quad (1.24)$$

Furthermore, we remark (or recall) that the Lamb-Oseen vortex yields a solution to the Navier Stokes equations with an explicit pressure:

$$\nabla \Pi = \alpha^2 \frac{x}{|x|^2} |\Theta(t, x)|^2 \quad \forall x \in \mathbb{R}^2 \setminus \{(0, 0)\}.$$

Hence, plugging the ansatz:

$$\begin{aligned} v(t, x) &= \alpha \Theta(t, x) + w(t, x), & \tilde{p}(t, x) &= \alpha^2 \Pi(t, x) + q(t, x), \\ \ell_w(t, x) &= \ell_w(t, x) & \omega_w(t, x) &= \alpha g(t, 1) + \omega_w(t, x). \end{aligned} \quad (1.25)$$

into (1.7)-(1.12), we obtain the perturbed system:

$$\frac{\partial w}{\partial t} + ((w - \ell_w(t)) \cdot \nabla) w - \Delta w + \nabla \varpi = -\alpha \left[(\Theta \cdot \nabla) w + ((w - \ell_w(t)) \cdot \nabla) \Theta \right] \quad \text{in } (0, \infty) \times \mathcal{F}_0, \quad (1.26)$$

$$\operatorname{div} w = 0 \quad \text{in } (0, \infty) \times \mathcal{F}_0, \quad (1.27)$$

$$w(t, x) = \ell_w(t) + \omega_w(t) x^\perp \quad \text{on } (0, \infty) \times \partial B_0, \quad (1.28)$$

$$m \ell_w'(t) = - \int_{\partial B_0} \Sigma(w, \varpi) n \, d\sigma(x) \quad \text{on } (0, \infty), \quad (1.29)$$

$$\mathcal{J}\omega'_w(t) = - \int_{\partial B(t)} x^\perp \cdot \Sigma(w, \varpi) n \, d\sigma(x) + \alpha\zeta(t) \quad \text{on } (0, \infty), \quad (1.30)$$

$$w|_{t=0} = w_0 \quad \text{on } \mathcal{F}_0, \quad (1.31)$$

$$\ell_w(0) = \ell_w^0, \quad \omega_w(0) = \omega_w^0. \quad (1.32)$$

with an explicit source term ζ . We detail this computation in **Section 3**. We can then rely on the study of the fluid-structure semi-group to construct a mild-solution to (1.26)–(1.32):

$$W(t) = S(t - t_0)W_0 + \int_{t_0}^{t-s} S(t - s)F_\alpha(s)ds \quad (1.33)$$

with a source term F_α to be made precise later on.

In this direction, our first result shows that this Duhamel-formula yields a suitable solution to our problem:

Theorem 1.2. *Let $(\alpha, t_0) \in \mathbb{R} \times [0, \infty)$ and $(w_0, \ell_w^0, \omega_w^0) \in L^2(\mathcal{F}_0) \cap \mathbb{R}^2 \cap \mathbb{R}$ such that (1.24). Then, the Duhamel formula (1.33) yields a triplet (w, ℓ_w, ω_w) such that:*

1. $w \in C([t_0, \infty); L^2(\mathcal{F}_0))$, with $\nabla w \in L^2((t_0, \infty); L^2(\mathcal{F}_0))$
2. $(\ell_w, \omega_w) \in C([t_0, \infty); \mathbb{R}^3)$
3. (w, ℓ_w, ω_w) is a solution to (1.26)–(1.32)

By reconstructing (v, ℓ, ω) via (1.25), we recover a global-in-time solution for unbounded-energy initial data of the form (1.23). We can then look at the large-time behavior of these solutions. To state this second result we shall start from a sufficiently developed Lamb-Oseen vortex, meaning that the radius of the vortex is sufficiently large, or that we consider the problem (1.7)–(1.12) starting from a time t_0 sufficiently large with an initial data obtained by perturbing $\alpha\Theta(t_0, \cdot)$ like in (1.23) with a small perturbation in \mathcal{L}^2 . We have then the following theorem:

Theorem 1.3. *Let the assumptions of Theorem 1.2 be in force and assume further that t_0 is sufficiently large, α is sufficiently small, $w_0 \in L^q(\mathcal{F}_0)$ for some $q \in (1, 2)$ and W_0 is small enough in \mathcal{L}^2 . The constructed solution (v, ℓ, ω) to (1.7)–(1.12) with initial condition (1.23) satisfies:*

$$\lim_{t \rightarrow \infty} t^{\frac{1}{2} - \frac{1}{p}} \|v(t) - \alpha\Theta(t, \cdot)\|_{L^p(\mathcal{F}_0)} = 0 \quad \forall p \in (2, \infty) \quad (1.34)$$

$$\sup_{t > t_0} (t - t_0)^{\frac{1}{q}} (|\ell(t)|) < \infty \quad (1.35)$$

Some comments are in order. First, the decay rate prescribed in (1.34) implies that $\alpha\Theta$ is indeed the leading term for large times. However, the explicit formula (1.22) entails that we have $|\Theta(t, x)| \leq 1/t$ on ∂B_0 so that the remainder may be much larger on ∂B_0 and induce a leading translation velocity. The complementary inequality (1.35) fixes then a minimal decay of the translation velocity depending only on the integrability of the initial perturbation.

The proofs of the two latter theorems rely on the $L^p - L^q$ properties of the semigroup $(S(t))_{t \geq 0}$ obtained in [4]. One key-difficulty in both cases is again the term $\ell_w \cdot \nabla w$. This term has limited space integrability (we cannot expect better than $\nabla w \in L^2(\mathcal{F}_0)$) and time-decay ($|\ell_w|$ decays a little less than $\|\nabla w\|_{L^2(\mathcal{F}_0)}$ but strictly less *a priori*). Hence, to handle this term we have to estimate sharply the loss of time-decay between $|\ell_w|$ and $\|\nabla w\|_{L^2(\mathcal{F}_0)}$. This is obtained by applying a sharp version of the Gagliardo-Nirenberg inequality and of the associated constant, following [3].

The outline of the paper is as follows. In the next section we provide preliminary lemmas. We explain the construction of the capital-letter unknowns and fluid-structure operator A . We recall the results of [4] on the decay properties of the semigroup and complement the analysis with a description of the fractional powers of A in the spirit of [7]. Finally, we recall the Gagliardo Nirenberg analysis underlying the stability analysis of the Lamb-Oseen vortex. In Section 3 we detail the proofs of Theorem 1.2 and Theorem 1.3. Section 4 is devoted to the proof of Theorem 1.1. Some further technicalities are presented in an appendix.

2. PRELIMINARY CONSTRUCTIONS AND TECHNICAL LEMMAS

In this section, we first recall the construction of function spaces that enable to handle the fluid unknown v and solid unknowns (ℓ, ω) at once. We also recall the construction of the unbounded operator A underlying the resolution of (1.14)-(1.18). These constructions are reproduced from [4, 13, 14].

The first key-issue we address is related to the problem of controlling the body linear velocity by the fluid velocity-field. In the forthcoming analysis, one would hope to be able to control the linear velocity $|\ell|$ by $\|\nabla v\|_{L^2(\mathcal{F}_0)}$ only. However, in full generality, this is possible in $3D$ but it turns out to be false in $2D$. This can be seen as reminiscent either of the fact that $\dot{H}^1(\mathbb{R}^2)$ embeds in no $L^p(\mathbb{R}^2)$ space or of the Stokes paradox [6, Introduction of Section V]. Here, we exchange such a control for an almost optimal control in the form of a family of Gagliardo-Nirenberg inequalities with an explicit estimate of the embedding constants. The second key-contribution of this section is the analysis of the fractional powers of the operator A .

2.1. Function spaces and Gagliardo-Nirenberg inequality

As classical in fluid+disk systems, we treat (1.7)-(1.12) by encoding all the unknowns (v, ℓ, ω) into one unified unknown with the following construction. From a triplet $(v, \ell, \omega) \in [C_c^\infty(\overline{\mathcal{F}_0})] \times \mathbb{R}^2 \times \mathbb{R}$ verifying

$$\operatorname{div} v = 0 \quad \text{in } \mathcal{F}_0, \quad v = \ell + \omega x^\perp \quad \text{on } \partial B_0,$$

we define a divergence-free vector field denoted V on \mathbb{R}^2 obtained by extending v by $\ell + \omega x^\perp$ in B_0 . Adapted to such V , we introduce the function spaces \mathcal{L}^p ($p \in [1, \infty]$) defined by

$$\mathcal{L}^p := \{V \in [L^p(\mathbb{R}^2)]^2, \operatorname{div} V = 0 \text{ in } \mathbb{R}^2, D(V) = 0 \text{ in } B_0\}.$$

We recall that, since B_0 is connected, the condition $D(V) = 0$ on B_0 implies that $V|_{B_0}$ is a rigid velocity-field. Conversely, we adapt below the convention that for $V \in \mathcal{L}^p$ we denote $v = \mathbb{1}_{\mathcal{F}_0} V$ and $(\ell_v, \omega_v) \in \mathbb{R}^2 \times \mathbb{R}$ the translation/angular velocities characterizing V in B_0 .

We recall now some classical properties of these spaces. When $p \in [1, \infty)$, we endow \mathcal{L}^p with the norm

$$\|V\|_{\mathcal{L}^p}^p = \int_{\mathcal{F}_0} |V|^p + \frac{m}{\pi} \int_{B_0} |V|^p,$$

(and the corresponding definition when $p = \infty$). When $(p, p') \in [1, \infty]$ are conjugate, we equip $(\mathcal{L}^p, \mathcal{L}^{p'})$ with the duality pairing:

$$\langle V, W \rangle_{\mathcal{L}^p, \mathcal{L}^{p'}} = \int_{\mathcal{F}_0} V \cdot W + \frac{m}{\pi} \int_{B_0} V \cdot W.$$

For any $p \in [1, \infty]$, it is straightforward that \mathcal{L}^p is a closed subspace of

$$L_\sigma^p(\mathbb{R}^2) := \{V \in L^p(\mathbb{R}^2) \text{ s.t. } \operatorname{div} V = 0\}$$

which is itself a closed subspace of $[L^p(\mathbb{R}^2)]^2$. In particular, there exists a projector $\mathbb{P}_p : [L^p(\mathbb{R}^2)]^2 \rightarrow \mathcal{L}^p$. When $p \in (1, \infty)$, this projector is analyzed in previous references such as [14]. Since all the \mathbb{P}_p coincide on $C_c^\infty(\mathbb{R}^2)$ we can drop the p -dependency and denote this projector with \mathbb{P} . Our analysis below relies on the following fundamental lemma whose proof can be found in [14, Remark 2.4]:

Lemma 2.1. *Given $p \in (1, \infty)$ the projector $\mathbb{P} : [L^p(\mathbb{R}^2)]^2 \rightarrow \mathcal{L}^p$ is bounded.*

We also define

$$\mathcal{H}^1 := \mathcal{L}^2 \cap [H^1(\mathbb{R}^2)]^2.$$

As a closed subspace of $[H^1(\mathbb{R}^2)]^2$ this is a separable Hilbert space when equipped with the norm

$$\|V\|_{\mathcal{H}^1} = \|V\|_{\mathcal{L}^2} + \|\nabla V\|_{L^2},$$

in which the set of $C_c^\infty(\mathbb{R}^2)$ -solenoidal vector-field is dense. Implicitly in the gradient norm, we use the shortcut L^2 for $L^2(\mathbb{R}^2)$. We keep this convention for norms of Lebesgue and Sobolev spaces in what follows. The \mathcal{H}^1 -norm is associated with a Korn inequality that reads as follows:

Lemma 2.2. For any $V \in \mathcal{H}^1$ there holds:

$$\int_{\mathbb{R}^2} |\nabla V|^2 = 2 \int_{\mathcal{F}_0} |D(V)|^2. \quad (2.1)$$

We refer to [13, Lemma 4.1] for a proof.

We complement this part of the section with a Gagliardo-Nirenberg inequality that will enable to control the linear velocity associated with a fluid velocity-field. This inequality reads as the following lemma:

Lemma 2.3. There exists $C > 0$ such that, for any $p \geq 2$ and any $u \in H^1(\mathbb{R}^2)$, there holds

$$\|u\|_{L^p} \leq C \sqrt{p} \|u\|_{L^2}^{\frac{2}{p}} \|\nabla u\|_{L^2}^{1-\frac{2}{p}}.$$

Proof. We use the following result of [3].

Lemma 2.4 ([3, Theorem 1.1]). Let $d \geq 2$ and $q \geq 1$ such that $q \leq \frac{d}{d-2}$ if $d \geq 3$. Define

$$\mathcal{D}^q(\mathbb{R}^d) = \left\{ u \in L^{q+1}(\mathbb{R}^d) \cap L^{2q}(\mathbb{R}^d) \mid \nabla u \in L^2(\mathbb{R}^d) \right\}$$

Then, for any function $u \in \mathcal{D}^q(\mathbb{R}^d)$, there holds

$$\|u\|_{L^{2q}} \leq A_{q,d} \|\nabla u\|_{L^2}^\theta \|u\|_{L^{q+1}}^{1-\theta},$$

where

$$A_{q,d} := \left(\frac{y(q-1)^2}{2\pi d} \right)^{\frac{\theta}{2}} \left(\frac{2y-d}{2y} \right)^{\frac{1}{2q}} \left(\frac{\Gamma(y)}{\Gamma(y-\frac{d}{2})} \right)^{\frac{\theta}{d}},$$

with

$$\theta = \frac{d(q-1)}{q(d+2-(d-2)q)}, \quad y = \frac{q+1}{q-1}.$$

Let $u \in H^1(\mathbb{R}^2)$. Applying the previous lemma with $d = 2$ and $q = \frac{p}{2}$, we get

$$\|u\|_{L^p} \leq A_{q,2} \|\nabla u\|_{L^2}^\theta \|u\|_{L^{q+1}}^{1-\theta}, \quad (2.2)$$

with

$$\theta = \frac{q-1}{2q} = \frac{1}{2} - \frac{1}{p}, \quad y = \frac{q+1}{q-1} = 1 + \frac{4}{p-2}$$

and

$$A_{q,2} = \left(\frac{y(q-1)^2}{4\pi} \right)^{\frac{\theta}{2}} \left(\frac{y-1}{y} \right)^{\frac{1}{p}} \left(\frac{\Gamma(y)}{\Gamma(y-1)} \right)^{\frac{\theta}{2}}.$$

Using the property of the Gamma function, we have $\Gamma(y) = (y-1)\Gamma(y-1)$, so that

$$A_{q,2} = \left(\frac{(p+2)(p-2)}{16\pi} \right)^{\frac{1}{4}-\frac{1}{2p}} \left(\frac{4}{p+2} \right)^{\frac{1}{p}} \left(\frac{4}{p-2} \right)^{\frac{1}{4}-\frac{1}{2p}} \leq C p^{\frac{1}{4}}. \quad (2.3)$$

Moreover, by interpolation, there holds

$$\|u\|_{L^{q+1}} \leq \|u\|_{L^2}^{\frac{2}{p+2}} \|u\|_{L^p}^{\frac{p}{p+2}}$$

Thus, putting this and (2.3) into (2.2) yields

$$\|u\|_{L^p} \leq C p^{\frac{1}{4}} \|\nabla u\|_{L^2}^{\frac{1}{2}-\frac{1}{p}} \|u\|_{L^2}^{\frac{1}{p}} \|u\|_{L^p}^{\frac{1}{2}},$$

The conclusion follows. \square

The above lemma entails the following control that we shall use without mention below:

Corollary 2.5. Let $p \geq 2$ and $V \in \mathcal{L}^p \cap H^1(\mathbb{R}^2)$. There exists a constant C independent of p and V such that:

$$|\ell_v| \leq C \sqrt{p} \|V\|_{\mathcal{L}^2}^{\frac{2}{p}} \|\nabla V\|_{L^2(\mathbb{R}^2)}^{1-\frac{2}{p}}.$$

2.2. Construction of the unbounded operator A and related properties

With the construction of the previous part in this section, we can now define the fluid-structure operator A which enables to rewrite the system (1.14)-(1.18) into the infinite-dimensional differential system (1.19). Following [4, 13, 14] we set:

$$\mathcal{D}(A) := \left\{ W \in \mathcal{H}^1(\mathbb{R}^2) \text{ s.t. } w = W|_{\mathcal{F}_0} \in [H^2(\mathcal{F}_0)]^2 \right\}.$$

We point out that such vector-fields admit a discontinuity of normal derivative on ∂B_0 . This is a key property that enables a non-trivial solid dynamics. For any $W \in \mathcal{D}(A)$ we set $AW = \mathbb{P}AW$ where (keeping the convention that $w = W|_{\mathcal{F}_0}$)

$$AW = \begin{cases} -\Delta w & \text{in } \mathcal{F}_0 \\ \frac{2}{m} \left(\int_{\partial B_0} D(w)n d\sigma \right) + 2\mathcal{J}^{-1} \left(\int_{\partial B_0} z^\perp D(w)n d\sigma \right) y^\perp & \text{in } B_0. \end{cases}$$

We note that this induces indeed an unbounded operator $\mathcal{D}(A) \rightarrow \mathcal{L}^2(\mathbb{R})$ because for any $W \in \mathcal{D}(A)$ we have $AW \in [L^2(\mathbb{R}^2)]^2$ (so that in particular \mathbb{P} corresponds actually to the L^2 -projection).

2.2.1. Previous analysis of A . In [13] the properties of A are studied in this hilbertian framework. We gather here the main conclusions. First, we have that the unbounded operator $(A, \mathcal{D}(A))$ is an accretive self-adjoint positive operator on \mathcal{L}^2 . Hence, the Cauchy problem

$$\begin{cases} \partial_t V + AV = 0 \\ V|_{t=0} = V_0 \end{cases} \quad (2.4)$$

has a unique solution for any $V_0 \in \mathcal{L}^2$ defining thus a contraction semi-group $(S(t))_{t>0}$. The relations between this semi-group and our linearized system is the content of the following proposition:

Proposition 2.6. *For any $V_0 \in \mathcal{L}^2$, the unique solution*

$$V := S(t)V_0 \in C([0, \infty); \mathcal{L}^2) \cap C^1((0, \infty); \mathcal{L}^2) \cap C((0, \infty); \mathcal{D}(A))$$

to the Cauchy problem (2.4) yields a vector field v and velocities (ℓ_v, ω_v) satisfying

- $v \in C([0, \infty); L^2(\mathcal{F}_0)) \cap C((0, \infty); H^2(\mathcal{F}_0))$,
- $(\ell_v, \omega_v) \in C([0, \infty); \mathbb{R}^2 \times \mathbb{R})$,

and a pressure $p \in C((0, \infty); H_{loc}^1(\mathcal{F}_0))$ such that (1.14)-(1.18) holds true with initial condition:

$$\ell_v(0) = \ell_0, \quad \omega_v(0) = \omega_0, \quad v(0, \cdot) = v_0 \quad \text{in } \mathcal{F}_0.$$

Remarking that the spaces $(\mathcal{L}^p)_{p \in (1, \infty)}$ share $\mathcal{L}^2 \cap C_c^\infty(\mathbb{R}^d)$ as dense subspace the properties of the semi-group $(S(t))_{t>0}$ are extended to the non-hilbertian setting in [4]. This is the content of the following lemma:

Lemma 2.7 ([4, Theorem 1.1]). *For each $q \in (1, \infty)$, the fluid-structure operator A generates a semigroup on \mathcal{L}^q which satisfies:*

- *For all $p \in [q, \infty]$, there exists $K_1 = K_1(p, q) > 0$ such that for every $V_0 \in \mathcal{L}^q$:*

$$\|S(t)V_0\|_{\mathcal{L}^p} \leq K_1 t^{\frac{1}{p} - \frac{1}{q}} \|V_0\|_{\mathcal{L}^q} \quad \text{for all } t > 0.$$

- *If $q \leq 2$, for $p \in [q, 2]$, there exists $K_2 = K_2(p, q) > 0$ such that for every $V_0 \in \mathcal{L}^q$:*

$$\|\nabla S(t)V_0\|_{L^p(\mathcal{F}_0)} \leq K_2 t^{-\frac{1}{2} + \frac{1}{p} - \frac{1}{q}} \|V_0\|_{\mathcal{L}^q} \quad \text{for all } t > 0.$$

- For $p \in [\max(2, q), \infty)$, there exists $K_3 = K_3(p, q) > 0$ such that for every $V_0 \in \mathcal{L}^q$:

$$\|\nabla S(t)V_0\|_{L^p(\mathcal{F}_0)} \leq \begin{cases} K_3 t^{-\frac{1}{2} + \frac{1}{p} - \frac{1}{q}} \|V_0\|_{\mathcal{L}^q} & \text{for all } 0 < t < 1, \\ K_3 t^{-\frac{1}{q}} \|V_0\|_{\mathcal{L}^q} & \text{for all } t \geq 1. \end{cases}$$

The above estimates for the gradient are only on \mathcal{F}_0 . However, when $V_0 \in \mathcal{L}^2$, $V(t) = S(t)V_0$ is in \mathcal{H}^1 (since it is in $\mathcal{D}(A)$) for $t > 0$ so that Lemma 2.2 applies. Thus, the estimates in Lemma 2.7 are sufficient to get a full \mathcal{H}^1 estimate. Last, we also recall duality decay estimates as shown in [4].

Lemma 2.8 ([4, Corollaries 3.10 and 3.11]). *Assume $1 < q \leq p < \infty$ and let $F \in L^q(\mathbb{R}^2; M_2(\mathbb{R}))$ satisfying $F = 0$ on B_0 . The following decay estimates for $V(t) = S(t)\mathbb{P} \operatorname{div} F$ hold true:*

- if $q \geq 2$, there exists $K_4 = K_4(p, q) > 0$ such that for all $t > 0$:

$$\|V(t)\|_{\mathcal{L}^p} \leq K_4 t^{-\frac{1}{2} + \frac{1}{p} - \frac{1}{q}} \|F\|_{L^q(\mathbb{R}^2)}.$$

- if $q \leq 2$, there exists $K_5 = K_5(p, q) > 0$ such that:

$$\|V(t)\|_{\mathcal{L}^p} \leq \begin{cases} K_5 t^{-\frac{1}{2} + \frac{1}{p} - \frac{1}{q}} \|F\|_{L^q(\mathbb{R}^2)} & \text{for all } 0 < t < 1, \\ K_5 t^{-1 + \frac{1}{p}} \|F\|_{L^q(\mathbb{R}^2)} & \text{for all } 1 \leq t. \end{cases}$$

- There exists $K_\ell = K_\ell(q) > 0$ such that for all $t > 0$:

$$|\ell_{V(t)}| \leq K_\ell t^{-\frac{1}{2} - \frac{1}{q}} \|F\|_{L^q(\mathbb{R}^2)}.$$

2.2.2. *Further material on A .* In this part, we complement the analysis of A with more properties of its fractional powers. The fluid-structure operator A being self-adjoint and positive definite, we may define A^μ for $\mu \in (-1, 1)$ through its spectral representation [12, Section II.3.2]. Since A is injective, we have that these fraction powers (either positive or negative) are positive self-adjoint operator with dense domains.

Our first proposition concerns the square-root of A .

Lemma 2.9. 1. *We have $\mathcal{D}(A^{\frac{1}{2}}) = \mathcal{H}^1(\mathbb{R}^2)$ and*

$$\|A^{\frac{1}{2}}V\|_{\mathcal{L}^2} = \sqrt{2}\|D(v)\|_{L^2(\mathcal{F}_0)}. \quad (2.5)$$

2. *Let $F \in L^2(\mathcal{F}_0)^{2 \times 2}$ then,*

$$\mathbb{P} \operatorname{div} F \in \mathcal{D}(A^{-\frac{1}{2}}) \quad \text{with} \quad \|A^{-\frac{1}{2}}\mathbb{P} \operatorname{div} F\|_{\mathcal{L}^2} \leq \|F\|_{L^2(\mathcal{F}_0)}.$$

Proof. We refer to [13, p. 63] for a proof of the first item. As for the second item, we follow [7] and propose a proof based on the approach of [12, Lemma III-2.6.1]. We point out that, the assumption $F \in L^2(\mathcal{F}_0)^{2 \times 2}$ has to be understood as

$$F = (F_{i,j})_{1 \leq i, j \leq 2} \in L^2(\mathbb{R}^2)^{2 \times 2} \quad \text{with} \quad \operatorname{Supp}(F_{i,j}) \subset \mathcal{F}_0 \quad \forall 1 \leq i, j \leq 2.$$

In order to enlighten the need of this further assumption on the support of F , we provide a complete proof.

By a standard approximation argument, we assume that $F \in C_c^\infty(\mathcal{F}_0)^{2 \times 2}$. Since $A^{-\frac{1}{2}}$ is self-adjoint, and because of the identities (2.1) and (2.5), our proof reduces to obtaining the bound:

$$|\langle \mathbb{P} \operatorname{div} F, A^{-\frac{1}{2}}w \rangle| \leq \|F\|_{L^2(\mathcal{F}_0)} \|\nabla A^{-\frac{1}{2}}w\|_{L^2(\mathbb{R}^2)} \quad \forall w \in \mathcal{D}(A^{-\frac{1}{2}})$$

Let $w \in \mathcal{D}(A^{-\frac{1}{2}})$ so that there exists $v \in \mathcal{D}(A^{\frac{1}{2}})$ for which $w = A^{\frac{1}{2}}v$ (and thus $v = A^{-\frac{1}{2}}w$). We have then by definition of projectors \mathbb{P} and integration by parts:

$$\begin{aligned} \langle \mathbb{P}\operatorname{div}F, A^{-\frac{1}{2}}w \rangle &= \frac{m}{\pi} \int_{B_0} \operatorname{div}F \cdot v + \int_{\mathcal{F}_0} \operatorname{div}F \cdot v \\ &= - \int_{\mathcal{F}_0} F : \nabla v. \end{aligned}$$

We conclude with a standard Cauchy-Schwarz inequality □

In the proof above, if we do not make further assumption on the support of F and take $w \in \mathcal{D}(A^{-\frac{1}{2}})$, the last identity yields:

$$\langle \mathbb{P}\operatorname{div}F, A^{-\frac{1}{2}}w \rangle = \left(\frac{m}{\pi} - 1\right) \int_{\partial B_0} Fn \cdot v - \int_{\mathcal{F}_0} F : \nabla v$$

where:

$$\int_{\partial B_0} Fn \cdot v d\sigma = \int_{\partial B_0} Fnd\sigma \cdot \ell_v + \int_{\partial B_0} Fn \cdot n^\perp d\sigma\omega.$$

To relax the assumption on the support of F we should be able to control this further term by $\|\nabla v\|_{L^2(\mathbb{R}^2)}$. This implies to obtain the boundedness of the mapping $v \mapsto \ell_v$ on $\mathcal{D}(A)$ endowed with the $\dot{H}^1(\mathbb{R}^2)$ topology. However the Stokes paradox (see [6, Introduction of section V]) implies in particular that this property does not hold true. Nevertheless, for sufficiently smooth F , we can slightly relax the assumption on the support by requiring some momentum to vanish on ∂B_0 . We remark also that, while in the $L^2(\mathcal{F}_0)$ -density of $C_c^\infty(\mathcal{F}_0)$ -tensor-fields, the divergence operator may create non-trivial distributions on ∂B_0 , the operator $A^{-\frac{1}{2}}\mathbb{P}\operatorname{div}$ does not.

We proceed with the analysis of the range of A^μ for $\mu \in (0, 1/2)$ corresponding to [7, Lemma 5.1]. This is the content of the next lemma:

Lemma 2.10. *Let $q \in (1, 2)$ and $\mu < 1/q - 1/2$. For all $v \in \mathcal{L}^2(\mathbb{R}^2) \cap [L^q(\mathbb{R}^2)]^2$ there exists a unique $w \in \mathcal{D}(A^\mu)$ such that $v = A^\mu w$. Furthermore, there exists a constant $C = C(q, \mu) > 0$ depending only on q and μ for which $\|w\|_{\mathcal{L}^2} \leq C(\|v\|_{L^q(\mathbb{R}^2)} + \|v\|_{L^2(\mathbb{R}^2)})$.*

We point out that, in this statement, the condition $v \in \mathcal{L}^2 \cap [L^q(\mathbb{R}^2)]^2$ reads also $v \in \mathcal{L}^2 \cap \mathcal{L}^q$. What remains of this section is devoted to the proof of this result. We first remark that the proof of [7, Lemma 5.1] yields from [9, Lemma 2.2]. So, our proof reduces mostly to check that the fluid-structure operator A satisfies the key-properties necessary to reproduce the proofs of these latter lemmas (that were concerned initially with the standard Stokes operator with homogeneous boundary conditions). In comparison with these previous results, we have a loss in terms of the correspondence $q \rightarrow \mu$ and also in the control which involves the L^2 -norm. In [9] the authors obtain similar results with $\mu = 1/q - 1/2$ and a control with the L^q -norm only. It seems we might not get such optimal bounds in our case. But this will not depreciate the final result.

In [9], the properties of the Stokes operator are analyzed on \mathcal{F}_0 when complemented with vanishing boundary conditions. The main argument is performed on a Laplace system and divergence-free constraints are then handled *via* abstract Heinz-Kato arguments (see [12, Lemma II.3.2.3, p. 100]). With our setting, this Laplace operator reads as follows. We set:

$$L_0^2[B_0] := \{V \in [L^2(\mathbb{R}^2)]^2 \text{ s.t. } V = 0 \text{ on } B_0\}$$

and $\mathbb{P}_0 : [L^2(\mathbb{R}^2)]^2 \rightarrow L_0^2[B_0]$ the corresponding orthogonal projection. Then, we define the operator $\tilde{\mathcal{A}}_0$ by

$$\mathcal{D}(\tilde{\mathcal{A}}_0) := \{V \in L_0^2[B_0] \text{ s.t. } v \in [H^2(\mathcal{F}_0)]^2\}.$$

with

$$\tilde{\mathcal{A}}_0[V] = \mathbb{P}_0[-\mathbb{1}_{\mathcal{F}_0}\Delta V], \quad \forall V \in \mathcal{D}(\tilde{\mathcal{A}}_0).$$

To take advantage of the analysis of [9] in order to study the fractional powers of A , we propose to use the same Heinz-Kato argument to handle the divergence-free constraint and to focus on the remaining Laplace equation

(completed with non-standard integral boundary conditions) with the help of $\tilde{\mathcal{A}}_0$. The operator $\tilde{\mathcal{A}}_0$ will take hold of the PDE and we shall complement the analysis with a fine study of our non standard boundary conditions. To this end, we first rewrite the integral boundary conditions introduced by A . This is the content of the following lemma:

Proposition 2.11. *Let $V \in \mathcal{D}(A)$ then there holds:*

$$\mathcal{A}V = \frac{1}{m} \left(\int_{\partial B_0} \partial_n v d\sigma \right) + \mathcal{J}^{-1} \left(\int_{\partial B_0} z^\perp \cdot \partial_n v d\sigma + 2\omega_v \right) y^\perp \quad \text{on } B_0.$$

Proof. It is sufficient to prove that, for any $V \in \mathcal{D}(A)$ and any $(\ell, \omega) \in \mathbb{R}^2 \times \mathbb{R}$ there holds:

$$\int_{\partial B_0} 2D(v)n \cdot (\ell + \omega z^\perp) d\sigma = \int_{\partial B_0} \partial_n v d\sigma \cdot \ell + \left(\int_{\partial B_0} z^\perp \cdot \partial_n v d\sigma + 2\omega_v \right) \omega.$$

So, let $V \in \mathcal{D}(A)$. Given $(\ell, \omega) \in \mathbb{R}^2 \times \mathbb{R}$ let:

$$W = \nabla^\perp \left[\chi(y) \left(\ell \cdot y^\perp + \omega |y|^2 \right) \right]$$

where $\chi \in C_c^\infty(\mathbb{R}^2)$ is fixed but arbitrary satisfying $\mathbf{1}_{B_0} \leq \chi \leq 1$. We note that with such conventions, there holds $W \in \mathcal{D}(A)$ with $\ell_W = \ell$ and $\omega_W = \omega$. We have then by integration by parts (using several times that w, W and v, V are divergence free):

$$\begin{aligned} \int_{\partial B_0} 2D(v)n \cdot (\ell + \omega z^\perp) d\sigma &= \int_{\partial B_0} 2D(v)n \cdot w d\sigma \\ &= \int_{\mathcal{F}_0} \operatorname{div}(2D(v)) \cdot w + 2D(v) : D(w) \\ &= \int_{\mathcal{F}_0} \Delta v \cdot w + \int_{\mathbb{R}^2} \nabla V : \nabla W \\ &= \int_{\partial B_0} \partial_n v \cdot w d\sigma + 2\omega_v \omega \\ &= \int_{\partial B_0} \partial_n v \cdot (\ell + \omega z^\perp) d\sigma + 2\omega_v \omega. \end{aligned}$$

The term $\omega_v \omega$ appearing on the fourth line is the contribution of the (skew-symmetric part of the) gradients ∇V and ∇W on B_0 . This ends the proof. \square

Thanks to **Proposition 2.11** we can now rewrite the fluid-structure operator $A = \mathbb{P}\tilde{\mathcal{A}}$ where $\tilde{\mathcal{A}}$ is defined (without the divergence-free constraint) by the formula:

$$\tilde{\mathcal{A}}W = \begin{cases} -\Delta w & \text{in } \mathcal{F}_0 \\ \frac{1}{m} \left(\int_{\partial B_0} \partial_n w d\sigma \right) + \mathcal{J}^{-1} \left(\int_{\partial B_0} z^\perp \cdot \partial_n w d\sigma + 2\omega_w \right) y^\perp & \text{in } B_0, \end{cases}$$

for $W \in \mathcal{D}(\tilde{\mathcal{A}}) = L^2[B_0] \cap [H^1(\mathbb{R}^2)]^2 \cap [H^2(\mathcal{F}_0)]^2$. Here, we denote:

$$L^2[B_0] = \left\{ W \in [H^1(\mathbb{R}^2)]^2 \text{ s.t. } W = \ell_W + \omega_W y^\perp \text{ on } B_0 \right\}.$$

We may reproduce here classical computations to obtain that $\tilde{\mathcal{A}}$ is a selfadjoint positive operator on $L^2[B_0]$ since it is associated with the quadratic form:

$$\langle \tilde{\mathcal{A}}W, V \rangle = \int_{\mathbb{R}^2} \nabla W : \nabla V, \quad \forall (W, V) \in \mathcal{D}(\tilde{\mathcal{A}}).$$

We point out that the duality bracket is still the one associated with the disk density. In particular, we have that (note that ∇W is the skew-symmetric matrix associated with ω_W on B_0):

$$\|\tilde{\mathcal{A}}^{\frac{1}{2}}W\|_{L^2[B_0]}^2 = \int_{\mathbb{R}^2} |\nabla W|^2 \quad \forall W \in \mathcal{D}(\tilde{\mathcal{A}}^{\frac{1}{2}}). \quad (2.6)$$

and, for $\lambda > 0$:

$$\|(\tilde{\mathcal{A}} + \lambda)^{\frac{1}{2}}W\|_{L^2[B_0]}^2 = \int_{\mathbb{R}^2} |\nabla W|^2 + \lambda \langle W, W \rangle \quad \forall W \in \mathcal{D}(\tilde{\mathcal{A}}^{\frac{1}{2}x}). \quad (2.7)$$

We recall that similar identities hold with the operator A . Thanks to these two latter identities, we can reproduce the procedure of [9, Lemma 2.2] and the proof of Lemma 2.10 reduces to obtaining the following proposition:

Proposition 2.12. *Let $q \in (1, 2)$ and $\mu < 1/q - 1/2$. For all $\varepsilon > 0$, there exists a mapping $R_{\mu, \varepsilon} : L^2[B_0] \cap [L^q(\mathbb{R}^2)]^2 \rightarrow L^2[B_0]$ satisfying:*

- for arbitrary $W \in L^2[B_0]$ there holds:

$$(\tilde{\mathcal{A}} + \varepsilon)^{-\mu}W = (\tilde{\mathcal{A}}_0 + \varepsilon)^{-\mu}(\mathbf{1}_{\mathcal{F}_0}W) + R_{\mu, \varepsilon}W$$

- there exists a constant $C := C(\mu) > 0$ depending on μ but independent of $\varepsilon > 0$ and $W \in L^2[B_0] \cap [L^q(\mathbb{R}^2)]^2$ such that :

$$\|R_{\mu, \varepsilon}W\|_{L^2(\mathbb{R}^2)} \leq C\|W\|_{L^q(\mathbb{R}^2)}. \quad (2.8)$$

We postpone the proof of this proposition to Appendix B. For completeness, we provide a proof of **Lemma 2.10** with this proposition at-hand.

Proof of Lemma 2.10. The proof follows a standard regularization-compactness scheme. Let $\mu \in (0, 1/2)$ and $q \in (1, 2)$ such that $\mu < 1/q - 1/2$. Given $W \in \mathcal{L}^2 \cap [L^q(\mathbb{R}^2)]^2$ and $\varepsilon \in (0, \infty)$ we can construct $(A + \varepsilon)^{-\mu}W$. Formula (2.7) with a Heinz-Kato argument imply then that

$$\|(A + \varepsilon)^{-\mu}W\|_{\mathcal{L}^2} \leq \|(\tilde{\mathcal{A}} + \varepsilon)^{-\mu}W\|_{L^2[B_0]}.$$

However, we have that:

$$(\tilde{\mathcal{A}} + \varepsilon)^{-\mu}W = (\tilde{\mathcal{A}}_0 + \varepsilon)^{-\mu}(\mathbf{1}_{\mathcal{F}_0}W) + R_{\mu, \varepsilon}W$$

For the first term, according to [9, Eq. (2.2)] (that holds componentwise in our setting) and a Hardy-Littlewood-Sobolev inequality, there holds:

$$\|(\tilde{\mathcal{A}}_0 + \varepsilon)^{-\mu}(\mathbf{1}_{\mathcal{F}_0}W)\|_{L^2} \leq C\|W\|_{L^{q'}(\mathcal{F}_0)}$$

where $1/q' = \mu + 1/2$. We have then $q' \in (q, 2)$ so that, by interpolation, we derive:

$$\|(\tilde{\mathcal{A}}_0 + \varepsilon)^{-\mu}(\mathbf{1}_{\mathcal{F}_0}W)\|_{L^2} \leq C(\|W\|_{L^q(\mathcal{F}_0)} + \|W\|_{L^2(\mathcal{F}_0)})$$

As for the other part, applying the previous proposition, we conclude that:

$$\|R_{\mu, \varepsilon}W\|_{L^2} \leq C(\mu)\|W\|_{L^q}.$$

Letting $\varepsilon \rightarrow 0$, we have thus that $(A + \varepsilon)^{-\mu}W$ converges to some V (in \mathcal{L}^2) that satisfies $A^\mu V = W$ with the expected control $\|V\|_{\mathcal{L}^2} \leq C(\|W\|_{L^q} + \|W\|_{L^2(\mathcal{F}_0)})$. \square

3. STABILITY OF THE OSEEN VORTEX

In this section, we construct global-in-time solutions to (1.26)-(1.32) for arbitrary w_0 and analyze the long-time behavior for small perturbations of fully-developed Oseen vortex.

To this end, we have first the following useful estimates in the same spirit as Lemma 2.1 of [7] (so that we do not detail the proof):

Lemma 3.1. *1. For any $p \in (2, \infty]$, there exists a constant $a_p > 0$ such that for all $t \geq 0$*

$$\|\Theta(t)\|_{L^p} \leq \frac{a_p}{(1+t)^{\frac{1}{2}-\frac{1}{p}}}. \quad (3.1)$$

2. For any $p \in (1, \infty]$, there exists $b_p > 0$ such that for all $t \geq 0$

$$\|\nabla\Theta(t)\|_{L^p} \leq \frac{b_p}{(1+t)^{1-\frac{1}{p}}}.$$

3. For all $t, s \geq 0$, we have

$$\|\Theta(t) - \Theta(s)\|_{L^2}^2 \leq \frac{1}{4\pi} \left| \log \frac{1+t}{1+s} \right|. \quad (3.2)$$

4. There exists a constant $\kappa_1 > 0$ such that for all $t, s \geq 0$,

$$\|\nabla\Theta(t) - \nabla\Theta(s)\|_{L^2}^2 \leq \kappa_1 \left| \frac{1}{1+t} - \frac{1}{1+s} \right|.$$

We recall then that, contrary to [7], we don't need to use a cut-off function. Indeed, the boundary conditions are here more suitable than the no-slip boundary condition of [7] for the Oseen vortex, since Θ is a pure rotation on $\partial B_0 : \Theta(t, x) = g(t, 1) x^\perp$ on ∂B_0 . From this remark and the construction of the pressure Π in the introduction, we obtain that, when plugging the ansatz (1.25) into (1.7)-(1.11), we may have a remainder term in the Newton laws only. Furthermore, we have the following proposition:

Proposition 3.2. *For all $t \geq 0$, there exists $C > 0$ such that for all $t \geq 0$*

$$\left| \int_{\partial B_0} x^\perp \cdot \Sigma(\Theta(t), \Pi(t)) n \, d\sigma(x) \right| + |\partial_t g(t, 1)| \leq C \frac{1}{(1+t)^2}.$$

There also holds for all $t \geq 0$

$$\int_{\partial B_0} \Sigma(\Theta(t), \Pi(t)) n \, d\sigma(x) = 0.$$

In particular, we see that there is actually no remainder in the Newton law for the linear momentum. But there is one in the Newton law on the angular momentum:

$$\zeta(t) := - \int_{\partial B_0} x^\perp \cdot \Sigma(\Theta(t), \Pi(t)) n \, d\sigma(x) - \mathcal{J} \partial_t g(t, 1).$$

The previous result yields the following estimate for this remainder.

Corollary 3.3. *There exists $C > 0$ such that for all $t \geq 0$,*

$$|\zeta(t)| \leq \frac{C}{(1+t)^2}.$$

Eventually, going to capital-letter unknowns, we obtain with similar arguments as in [13] that we have a solution (w, ℓ_w, ω) to (1.26)-(1.32) if the associated W satisfies (1.33) with

$$F_\alpha(s) = \zeta(s)x^\perp \frac{\mathbb{1}_{B_0}}{\mathcal{J}} - \mathbb{P} \left[((\Theta(s) \cdot \nabla)w(s) + (w(s) \cdot \nabla)\Theta(s) - (\ell_{W(s)} \cdot \nabla)\Theta(s))\mathbb{1}_{\mathcal{F}_0} \right] \\ - \mathbb{P} \left[((w(s) - \ell_{W(s)}) \cdot \nabla)w(s)\mathbb{1}_{\mathcal{F}_0} \right].$$

We proceed with the proof of **Theorem 1.2** and **Theorem 1.3**. We first study in the next subsection the Duhamel formula (1.33) on small timespan or for small data. This shall imply that we have local-in-time solutions to (1.26)-(1.32) as well as **Theorem 1.3**. Thanks to this first study, we have also that non-extendable solutions to (1.26)-(1.32) must blow-up in the \mathcal{L}^2 norm so that **Theorem 1.2** yields from an *a priori* estimate on $\|W\|_{\mathcal{L}^2}$ that we prove in a last subsection.

3.1. Local-in-time Cauchy theory and proof of Theorem 1.3

The main result of this part is the following theorem:

Theorem 3.4. *1. For any $\alpha \in \mathbb{R}$, any $t_0 \geq 0$ and any $W_0 \in \mathcal{L}^2$, there exists $T = T(\alpha, W_0) > 0$ such that (1.33) has a unique solution $W \in \mathcal{C}^0([t_0, t_0 + T], \mathcal{L}^2) \cap \mathcal{C}^0((t_0, t_0 + T], H^1(\mathbb{R}^2))$. Moreover, any upper bound on $|\alpha| + \|W_0\|_{\mathcal{L}^2}$ gives a lower bound on the local existence time T .*

2. There exists positive constants K_0, δ, K_6 and T_0 such that, if $t_0 \geq T_0$, if $|\alpha| \leq \delta$, and if $\|W_0\|_{\mathcal{L}^2} \leq K_6$, then the solution $W(t)$ of (1.33) given by the previous part is global in time and satisfies

$$\sup_{t \geq t_0} \|W(t)\|_{\mathcal{L}^2} + \sup_{t > t_0} (t - t_0)^{\frac{1}{2}} (\|\nabla w(t)\|_{L^2(\mathcal{F}_0)} + |\ell_{W(t)}|) \leq K_0 (\|W_0\|_{\mathcal{L}^2} + |\alpha|(1 + t_0)^{-\frac{5}{4}}).$$

In addition, if

$$M := \sup_{\tau \geq 0} \tau^\mu \|S(\tau)W_0\|_{\mathcal{L}^2} + \sup_{\tau > 0} \tau^{\mu + \frac{1}{2}} \left(\|\nabla S(\tau)W_0\|_{L^2(\mathcal{F}_0)} + |\ell_{S(\tau)W_0}| \right) < \infty, \quad (3.3)$$

for a fixed $\mu \in (0, \frac{1}{2})$, then

$$\sup_{t \geq t_0} (t - t_0)^\mu \|W(t)\|_{\mathcal{L}^2} + \sup_{t > t_0} (t - t_0)^{\mu + \frac{1}{2}} (\|\nabla w(t)\|_{L^2(\mathcal{F}_0)} + |\ell_{W(t)}|) \leq 2M + C|\alpha|$$

for some $C > 0$.

Proof. The proof is very similar to the proof of Proposition 3.2 of [7], who followed the classical approach of Fujita and Kato [5]. Given $t_0 \geq 0$, we introduce the Banach space $X := \mathcal{C}^0([t_0, \infty), \mathcal{L}^2) \cap \mathcal{C}^0((t_0, \infty), H^1(\mathbb{R}^2) \cap L^\infty(B_0))$, equipped with the norm

$$\|W\|_X = \sup_{t \geq t_0} \|W(t)\|_{\mathcal{L}^2} + \sup_{t > t_0} (t - t_0)^{\frac{1}{2}} (\|\nabla w(t)\|_{L^2(\mathcal{F}_0)} + |\ell_{W(t)}|).$$

From Lemma 2.7, we know that $S(t - t_0)W_0 \in X$ and $\|S(t - t_0)W_0\|_X \leq K\|W_0\|_{\mathcal{L}^2}$ for some constant $K > 0$. On the other hand, given any $W \in X$, we denote for $t \geq t_0$:

$$F_0(t) = \int_{t_0}^t S(t - s) \mathbb{P} \left[\zeta(s)x^\perp \frac{\mathbb{1}_{B_0}}{\mathcal{J}} \right] ds, \\ (F_1 W)(t) = \int_{t_0}^t S(t - s) \mathbb{P} \left[((\Theta(s) \cdot \nabla)w(s)) \right] ds, \\ (F_2 W)(t) = \int_{t_0}^t S(t - s) \mathbb{P} \left[(w(s) - \ell_{W(s)}) \cdot \nabla w(s) \right] ds, \\ (F_3 W)(t) = \int_{t_0}^t S(t - s) \mathbb{P} \left[(w(s) - \ell_{W(s)}) \cdot \nabla \Theta(s) \right] ds,$$

$$(FW)(t) = \alpha F_0(t) + \alpha(F_1W)(t) + (F_2W)(t) + \alpha(F_3W)(t).$$

We show that F maps X into X and that, for some constant $K > 0$,

$$\|FW\|_X \leq K(|\alpha|(1+t_0)^{-\frac{1}{4}} + |\alpha|\|W\|_X + \|W\|_X^2), \quad (3.4)$$

$$\|FW_1 - FW_2\|_X \leq K(|\alpha| + \|W_1\|_X + \|W_2\|_X)\|W_1 - W_2\|_X. \quad (3.5)$$

For this, we compute now bounds successively for F_0 , F_1 , F_2 , and F_3 . First, using Corollary 3.3 and Lemma 2.7 (with $q = \frac{4}{3}$), we get for all $t \geq t_0$

$$\|F_0(t)\|_{\mathcal{L}^2} + (t-t_0)^{\frac{1}{2}}(\|\nabla F_0(t)\|_{L^2(\mathcal{F}_0)} + \|F_0(t)\|_{L^\infty}) \leq C \int_{t_0}^t \left(\frac{1}{(t-s)^{\frac{1}{4}}} + \frac{(t-t_0)^{\frac{1}{2}}}{(t-s)^{\frac{3}{4}}} \right) \frac{1}{(1+s)^2} ds \leq \frac{C}{(1+t_0)^{\frac{5}{4}}}.$$

Then, we control F_2 with the help of Lemma A.1 (see Appendix A) which ensures that:

$$\|F_2(t)\|_{\mathcal{L}^2} + (t-t_0)^{\frac{1}{2}}(\|\nabla F_2(t)\|_{L^2(\mathcal{F}_0)} + \|F_2(t)\|_{L^\infty}) \leq C\|W\|_X^2$$

Similarly, there holds

$$\begin{aligned} \|(w - \ell_{W(s)} \cdot \nabla)\Theta(s)\|_{L^{\frac{4}{3}}(\mathcal{F}_0)} &\leq \|w\|_{L^4(\mathcal{F}_0)}\|\nabla\Theta(s)\|_{L^2(\mathcal{F}_0)} + |\ell_{W(s)}|\|\nabla\Theta(s)\|_{L^{\frac{4}{3}}(\mathcal{F}_0)} \\ &\leq C \left(\frac{1}{(s-t_0)^{\frac{1}{4}}(1+s)^{\frac{1}{2}}} + \frac{1}{(s-t_0)^{\frac{1}{2}}(1+s)^{\frac{1}{4}}} \right) \|W\|_X \end{aligned}$$

so that, applying the boundedness of $\mathbb{P} : L^{4/3}(\mathbb{R}^2) \rightarrow \mathcal{L}^{4/3}$ (see [14, Remark 2.4]):

$$\begin{aligned} \|F_3W(t)\|_{\mathcal{L}^2} + (t-t_0)^{\frac{1}{2}}(\|\nabla F_3W(t)\|_{L^2(\mathcal{F}_0)} + \|F_3W(t)\|_{L^\infty}) \\ \leq C \int_{t_0}^t \left(\frac{1}{(t-s)^{\frac{1}{4}}} + \frac{(t-t_0)^{\frac{1}{2}}}{(t-s)^{\frac{3}{4}}} \right) \left(\frac{1}{(s-t_0)^{\frac{1}{4}}(1+s)^{\frac{1}{2}}} + \frac{1}{(s-t_0)^{\frac{1}{2}}(1+s)^{\frac{1}{4}}} \right) ds \|W\|_X \\ \leq C\|W\|_X. \end{aligned}$$

We finally bound F_1W . To this end, the procedure is similar to that of [7]. First, we observe that $\Theta \cdot n = 0$ on ∂B so that we can rewrite:

$$\mathbb{P}[\mathbf{1}_{\mathcal{F}_0}(\Theta \cdot \nabla)w] = A^{\frac{1}{2}}A^{-\frac{1}{2}}\mathbb{P}\operatorname{div}(\mathbf{1}_{\mathcal{F}_0}\Theta \otimes w).$$

Moreover, using Lemma 2.9 and the estimate (3.1), we compute

$$\left\| A^{-\frac{1}{2}}\mathbb{P}\operatorname{div}(\mathbf{1}_{\mathcal{F}_0}\Theta \otimes w) \right\|_{L^2(\mathcal{F}_0)} \leq \|\Theta(s)w(s)\|_{L^2(\mathcal{F}_0)} \leq \frac{C}{(1+s)^{\frac{1}{2}}}\|W\|_X.$$

Therefore, Lemmas 2.7 and (2.5) and the fact that $A^{\frac{1}{2}}$ commutes with the semigroup lead to:

$$\begin{aligned} \|(F_1W)(t)\|_{\mathcal{L}^2} &\leq \int_{t_0}^t (t-s)^{-\frac{1}{2}} \left\| A^{-\frac{1}{2}}\mathbb{P}\operatorname{div}(\mathbf{1}_{\mathcal{F}_0}\Theta \otimes w(s)) \right\|_{\mathcal{L}^2} ds \\ &\leq C \int_{t_0}^t (t-s)^{-\frac{1}{2}}(1+s)^{-\frac{1}{2}}\|W\|_X ds \leq C\|W\|_X, \end{aligned}$$

and

$$\begin{aligned} (t-t_0)^{\frac{1}{2}}(\|\nabla(F_1W)(t)\|_{L^2(\mathcal{F}_0)} + |\ell_{W(t)}|) &\leq \int_{t_0}^{\frac{t+t_0}{2}} \frac{(t-t_0)^{\frac{1}{2}}}{t-s} \left\| A^{-\frac{1}{2}}\mathbb{P}\operatorname{div}(\mathbf{1}_{\mathcal{F}_0}\Theta \otimes w)(s) \right\|_{\mathcal{L}^2} ds \\ &\quad + \int_{\frac{t+t_0}{2}}^t \frac{(t-t_0)^{\frac{1}{2}}}{(t-s)^{\frac{1}{2}}} \|(\Theta(s) \cdot \nabla)w(s)\|_{L^2(\mathcal{F}_0)} ds \end{aligned}$$

$$\begin{aligned}
&\leq C \left[\int_{t_0}^{\frac{t+t_0}{2}} \frac{(t-t_0)^{\frac{1}{2}}}{t-s} (1+s)^{-\frac{1}{2}} ds \|W\|_X \right. \\
&\quad \left. + \int_{\frac{t+t_0}{2}}^t \frac{(t-t_0)^{\frac{1}{2}}}{(t-s)^{\frac{1}{2}}} \frac{1}{(1+s)^{\frac{1}{2}}(s-t_0)^{\frac{1}{2}}} ds \|W\|_X \right] \\
&\leq C \|W\|_X.
\end{aligned}$$

Since $FW = \alpha F_0 + \alpha F_1 W + F_2 W + \alpha F_3 W$, this concludes the proof of (3.4). The Lipschitz bound (3.5) is established in the same way.

Now let $r > 0$ such that $4Kr \leq 1$ and define $B_r := \{W \in X \mid \|W\|_X \leq r\}$. If we assume that $4|\alpha|K \leq 1$, $4K\|W_0\|_{\mathcal{L}^2} \leq r$ and $4K|\alpha|(1+t_0)^{-\frac{5}{4}} \leq r$, then (3.4) and (3.5) imply that the map $W \mapsto S(t-t_0)W_0 + FW$ leaves the closed ball B_r invariant and is a strict contraction in B_r . By construction, the unique fixed point of this map in B_r is the desired solution of (1.33). This proves part 2 of Theorem 3.4 with

$$K_0 = K, \quad \delta = \frac{1}{4K}, \quad K_6 = \frac{1}{16K^2}, \quad T_0 = (4K)^{\frac{4}{5}}.$$

The first part of this proof can be modified in a classical way (see for instance [1, 5]) to yield the local Cauchy theory, i.e. part 1 of Theorem 3.4.

In a second step, we assume that (3.3) holds for some $\mu \in (0, \frac{1}{2})$. Given any $T > t_0$, we denote

$$\mathcal{E}_T = \sup_{t_0 \leq t \leq T} (t-t_0)^\mu \|W(t)\|_{\mathcal{L}^2} + \sup_{t_0 < t \leq T} (t-t_0)^{\mu+\frac{1}{2}} (\|\nabla w(t)\|_{L^2(\mathcal{F}_0)} + |\ell_{W(t)}|),$$

where W (also represented by the triplet (w, ℓ_W, ω_W)) is the solution of (1.33) previously constructed. Since $W(t) = S(t-t_0)W_0 + (FW)(t)$, we have

$$\mathcal{E}_T \leq M + \sup_{t_0 \leq t \leq T} (t-t_0)^\mu \|(FW)(t)\|_{\mathcal{L}^2} + \sup_{t_0 < t \leq T} (t-t_0)^{\mu+\frac{1}{2}} (\|\nabla(FW)(t)\|_{L^2(\mathcal{F}_0)} + |\ell_{(FW)(t)}|),$$

where M is defined in (3.3). Let $p \in (1, 2)$ be such that $\frac{1}{p} > \mu + \frac{1}{2}$ and define $q \in (2, \infty)$ such that $\frac{1}{q} = \frac{1}{p} - \frac{1}{2}$. In particular, $\frac{1}{2} > \frac{1}{q} > \mu$. First, we have in a similar way as previously:

$$\begin{aligned}
&(t-t_0)^\mu \|F_0(t)\|_{\mathcal{L}^2} + (t-t_0)^{\mu+\frac{1}{2}} (\|\nabla F_0(t)\|_{L^2(\mathcal{F}_0)} + \|F_0(t)\|_{L^\infty}) \\
&\leq C \int_{t_0}^t \left(\frac{(t-t_0)^\mu}{(t-s)^{\frac{1}{q}}} + \frac{(t-t_0)^{\mu+\frac{1}{2}}}{(t-s)^{\frac{1}{p}}} \right) \frac{1}{(1+s)^2} ds \\
&\leq \frac{C}{(1+t_0)^{\frac{1}{q}-\mu+1}}.
\end{aligned}$$

The same computations as previously can be done for $F_1 W$, $F_2 W$ and $F_3 W$ introducing the further decay of W induced by \mathcal{E}_T (see [7] for more details), so that we finally get

$$\mathcal{E}_T \leq M + \tilde{K} (|\alpha|(1+t_0)^{\mu-\frac{1}{q}-1} + |\alpha|\mathcal{E}_T + \|W\|_X \mathcal{E}_T), \quad (3.6)$$

for some positive constant \tilde{K} independent of T and t_0 . Taking δ and K_6 smaller and t_0 larger if needed, we can ensure that $2\tilde{K}(|\alpha| + \|W\|_X) \leq 1$, so that (3.6) leads to

$$\mathcal{E}_T \leq 2M + 2\tilde{K}|\alpha|(1+t_0)^{\mu-\frac{1}{q}-1},$$

for all $T > t_0$. This concludes the proof. \square

To conclude this part, we provide two comments regarding the link between **Theorem 3.4** and the claimed results. The local-in-time Cauchy theory entails straightforwardly from item 1 in **Theorem 3.4**. The remark on the time of existence T also entails that non-extendable solution are global if $\|W\|_{\mathcal{L}^2}$ does not blow up. To obtain **Theorem 1.2** we complement the study with an *a priori* estimate in the next subsection. Concerning item 2, we recall that, for the linearized system we have the decay estimates of Lemma 2.7. Hence we infer the content of **Theorem 1.3** by remarking that, if $W_0 \in \mathcal{L}^2 \cap \mathcal{L}^q$ (meaning that $w_0 \in L^q(\mathcal{F}_0)$) is small in \mathcal{L}^2 then the assumption (3.3) is satisfied with $\mu = \frac{1}{q} - \frac{1}{2}$.

3.2. A logarithmic energy estimate

In this section we complement the proof of **Theorem 1.2** by establishing an estimate in the energy space. This is the content of the next lemma:

Lemma 3.5. *There exists a constant $K > 0$ such that, for any $\alpha \in \mathbb{R}$ and any $W_0 \in \mathcal{L}^2$, the solution of (1.33) with initial data W_0 provided by Theorem 3.4 is global in time and satisfies, for all $t \geq 0$,*

$$\|W(t)\|_{\mathcal{L}^2}^2 + \int_0^t \|D(w(s))\|_{L^2(\mathcal{F}_0)} ds \leq K \left(\|W_0\|_{\mathcal{L}^2}^2 + \alpha^2 \log(1+t) + K_\alpha \right),$$

where $K_\alpha = \alpha^2(1 + \log(1 + |\alpha|))$.

Proof. Fix $\alpha \in \mathbb{R}$, $W_0 \in \mathcal{L}^2$, and let $W \in \mathcal{C}^0([0, T], \mathcal{L}^2) \cap \mathcal{C}^0((0, T], \mathcal{H}^1)$ be the solution provided by Theorem 3.4 with initial data $W(0) = W_0$. We recall that we denote $V = W + \alpha\Theta$. Given any $\tau \geq 0$, we define then,

$$\begin{aligned} \tilde{w}_\tau(t, x) &= v(t, x) - \alpha\Theta(t + \tau, x) = w(t, x) + \alpha \left(\Theta(t, x) - \Theta(t + \tau, x) \right), \quad \text{for all } x \in \mathcal{F}_0, \\ \ell_{\tilde{W}_\tau(t)} &= \ell_V(t) = \ell_{W(t)}, \\ \omega_{\tilde{W}_\tau(t)} &= \omega_V - \alpha g(t + \tau, 1) = \omega_{W(t)} + \alpha \left(g(t, 1) - g(t + \tau, 1) \right). \end{aligned}$$

The given \tilde{W}_τ (represented by the triplet $(\tilde{w}_\tau, \ell_{\tilde{W}_\tau(t)}, \omega_{\tilde{W}_\tau(t)})$) satisfy the system of equations (1.26)-(1.32) (or equivalently (1.33)), where $\Theta(t)$ and $\zeta(t)$ are replaced by $\Theta(t + \tau)$ and $\zeta(t + \tau)$. Assume first that the solutions are smooth enough. Multiplying both sides of (1.26) by \tilde{w}_τ and integrating by parts over \mathcal{F}_0 (using the fact that \tilde{w}_τ and Θ are divergence-free), we find

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\tilde{w}_\tau\|_{L^2(\mathcal{F}_0)}^2 + 2 \|D(\tilde{w}_\tau(t))\|_{L^2(\mathcal{F}_0)}^2 &= \int_{\partial B_0} \tilde{w}_\tau(t) \cdot \Sigma(\tilde{w}_\tau(t)) n \, d\sigma(x) - \frac{\alpha}{2} \int_{\partial B_0} \Theta(t + \tau) |\tilde{w}_\tau|^2 \cdot n \, d\sigma(x) \\ &\quad - \alpha \int_{\mathcal{F}_0} \tilde{w}_\tau(t) \cdot \left((\tilde{w}_\tau(t) - \ell_{\tilde{W}_\tau(t)}) \cdot \nabla \right) \Theta(t + \tau) \, dx + \frac{1}{2} \int_{\partial B_0} |\tilde{w}_\tau(t)|^2 (\tilde{w}_\tau(t) - \ell_{\tilde{W}_\tau(t)}) \cdot n \, d\sigma(x). \end{aligned}$$

Since Θ and $\tilde{w}_\tau(t) - \ell_{\tilde{W}_\tau(t)}$ are orthogonal to n on ∂B_0 , the second and fourth terms also vanish. (1.28)-(1.30) then yield

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \left(\|\tilde{w}_\tau(t)\|_{L^2(\mathcal{F}_0)}^2 + m \left| \ell_{\tilde{W}_\tau(t)} \right|^2 + \mathcal{J} \left| \omega_{\tilde{W}_\tau(t)} \right|^2 \right) + 2 \|D(\tilde{w}_\tau(t))\|_{L^2(\mathcal{F}_0)}^2 \\ = -\alpha \int_{\mathcal{F}_0} \tilde{w}_\tau(t) \cdot \left((\tilde{w}_\tau(t) - \ell_{\tilde{W}_\tau(t)}) \cdot \nabla \right) \Theta(t + \tau) \, dx + \alpha \zeta(t + \tau) \omega_{\tilde{W}_\tau(t)}. \end{aligned}$$

The right-hand side can be estimated as usual with Lemma 3.1:

$$\begin{aligned} \left| \int_{\mathcal{F}_0} \tilde{w}_\tau(t) \cdot \left((\tilde{w}_\tau(t) - \ell_{\tilde{W}_\tau(t)}) \cdot \nabla \right) \Theta(t + \tau) \, dx \right| &\leq \|\tilde{w}_\tau(t)\|_{L^2(\mathcal{F}_0)} \frac{C}{1 + \tau + t}, \\ \left| \int_{\mathcal{F}_0} \tilde{w}_\tau(t) \cdot \left(\ell_{\tilde{W}_\tau(t)} \cdot \nabla \right) \Theta(t + \tau) \, dx \right| &\leq \|\tilde{w}_\tau(t)\|_{L^2(\mathcal{F}_0)} \left| \ell_{\tilde{W}_\tau(t)} \right| \frac{C}{(1 + \tau + t)^{\frac{1}{2}}}, \\ \left| \zeta(t + \tau) \omega_{\tilde{W}_\tau(t)} \right| &\leq \frac{C}{(1 + \tau + t)^2} \left| \omega_{\tilde{W}_\tau(t)} \right| \leq \frac{C}{(1 + \tau + t)^2} \left(\left| \omega_{\tilde{W}_\tau(t)} \right|^2 + 1 \right) \end{aligned}$$

Integrating in time from 0 to t for any $t > 0$ leads to

$$\begin{aligned} \frac{1}{2} \left\| \tilde{W}_\tau(t) \right\|_{\mathcal{L}^2}^2 + 2 \int_0^t \|D(\tilde{w}_\tau(s))\|_{L^2(\mathcal{F}_0)}^2 \, ds \\ \leq \frac{1}{2} \left\| \tilde{W}_\tau(0) \right\|_{\mathcal{L}^2}^2 + K |\alpha| \int_0^t \left(\frac{\|\tilde{w}_\tau(s)\|_{L^2(\mathcal{F}_0)}^2}{1 + \tau + s} + \frac{\|\tilde{w}_\tau(s)\|_{L^2(\mathcal{F}_0)} \left| \ell_{\tilde{W}_\tau(s)} \right|}{(1 + \tau + s)^{\frac{1}{2}}} + \frac{\left| \omega_{\tilde{W}_\tau(s)} \right|^2}{(1 + \tau + s)^2} + \frac{1}{(1 + \tau + s)^2} \right) \, ds, \end{aligned} \tag{3.7}$$

for some constant $K > 0$, independent of τ in particular. Such an estimate then also holds for weaker solutions. From this estimate, for $\tau = 0$, the Gronwall lemma shows that $\left\| \tilde{W}_\tau(t) \right\|_{\mathcal{L}^2}^2$ is bounded locally in time. Therefore, the solution W of (1.26)-(1.32) provided by the part 1 of Theorem 3.4 is global in time. Then, for general $\tau \geq 0$, we need to better estimate the second term, in particular $\left| \ell_{\tilde{W}_\tau(s)} \right|$ which should decrease faster than $\|\tilde{w}_\tau(s)\|_{L^2(\mathcal{F}_0)}$ (or $\left\| \tilde{W}_\tau(s) \right\|_{\mathcal{L}^2}$ equivalently). For this, we use Corollary 2.5. Applying it for $p = 2 + \log(1 + \tau + s)$, we get:

$$\begin{aligned} \left| \ell_{\tilde{W}_\tau(s)} \right| &\leq C(2 + \log(1 + \tau + s))^{\frac{1}{2}} \left\| \tilde{W}_\tau(s) \right\|_{L^2(\mathbb{R}^2)}^{\frac{2}{2+\log(1+\tau+s)}} \left\| \nabla \tilde{W}_\tau(s) \right\|_{L^2(\mathbb{R}^2)}^{1-\frac{2}{2+\log(1+\tau+s)}} \\ &\leq C(2 + \log(1 + \tau + s))^{\frac{1}{2}} \left\| \tilde{W}_\tau(s) \right\|_{\mathcal{L}^2}^{\frac{2}{2+\log(1+\tau+s)}} \left\| D(\tilde{W}_\tau(s)) \right\|_{L^2(\mathcal{F}_0)}^{1-\frac{2}{2+\log(1+\tau+s)}}, \end{aligned}$$

where we have used Lemma 2.2 in the last estimate. Then, we obtain:

$$\begin{aligned} \frac{\|\tilde{w}_\tau(s)\|_{L^2(\mathcal{F}_0)} \left| \ell_{\tilde{W}_\tau(s)} \right|}{(1 + \tau + s)^{\frac{1}{2}}} &\leq C \left(\frac{2 + \log(1 + \tau + s)}{1 + \tau + s} \right)^{\frac{1}{2}} \left\| \tilde{W}_\tau(s) \right\|_{\mathcal{L}^2}^{1+\frac{2}{2+\log(1+\tau+s)}} \left\| D(\tilde{W}_\tau(s)) \right\|_{L^2(\mathcal{F}_0)}^{1-\frac{2}{2+\log(1+\tau+s)}} \\ &\leq \left\| D(\tilde{W}_\tau(s)) \right\|_{L^2(\mathcal{F}_0)}^2 + C \left[\frac{2 + \log(1 + \tau + s)}{1 + \tau + s} \right]^{\xi(\tau+s)} \left\| \tilde{W}_\tau(s) \right\|_{\mathcal{L}^2}^2, \end{aligned}$$

where

$$\xi(x) := \frac{1}{1 + \frac{2}{2+\log(1+x)}} = 1 - \frac{1}{2 + \log(1+x)} \frac{2}{1 + \frac{2}{2+\log(1+x)}}.$$

In particular, we can easily compute that

$$\left[\frac{2 + \log(1 + \tau + s)}{1 + \tau + s} \right]^{\xi(\tau+s)} \leq C \frac{2 + \log(1 + \tau + s)}{1 + \tau + s}.$$

Therefore, we obtain

$$\begin{aligned} \frac{1}{2} \left\| \tilde{W}_\tau(t) \right\|_{\mathcal{L}^2}^2 + \int_0^t \left\| D(\tilde{w}_\tau(s)) \right\|_{L^2(\mathcal{F}_0)}^2 ds \\ \leq \frac{1}{2} \left\| \tilde{W}_\tau(0) \right\|_{\mathcal{L}^2}^2 + \frac{K|\alpha|}{1 + \tau} + K|\alpha| \int_0^t \left\| \tilde{W}_\tau(s) \right\|_{L^2(\mathcal{F}_0)}^2 \frac{2 + \log(1 + \tau + s)}{1 + \tau + s} ds, \end{aligned}$$

By applying the Gronwall lemma, we get

$$\begin{aligned} \frac{1}{2} \left\| \tilde{W}_\tau(t) \right\|_{\mathcal{L}^2}^2 + \int_0^t \left\| D(\tilde{w}_\tau(s)) \right\|_{L^2(\mathcal{F}_0)}^2 ds \\ \leq K \left[\left\| \tilde{W}_\tau(0) \right\|_{\mathcal{L}^2}^2 + \frac{|\alpha|}{1 + \tau} \right] \exp \left[K|\alpha| (\log(1 + \tau + t)^2 - \log(1 + \tau)^2 + \log(1 + \tau + t) - \log(1 + \tau)) \right] \end{aligned}$$

Now take $\tau = (\chi t)^2$ where $\chi = 1 + |\alpha|$, we get:

$$\log(1 + \tau + t) - \log(1 + \tau) = \log \left(1 + \frac{t}{1 + (\chi t)^2} \right) \leq C$$

and

$$\begin{aligned} \log(1 + \tau + t)^2 - \log(1 + \tau)^2 &= \log \left(1 + \frac{t}{1 + (\chi t)^2} \right) \left(\log(1 + t + (\chi t)^2) + \log(1 + (\chi t)^2) \right) \\ &\leq C \frac{t \log(1 + (\chi t)^2)}{1 + (\chi t)^2} \leq \frac{C}{\chi}. \end{aligned}$$

Thanks to the estimate (3.2) and the explicit expression of $g(t, r)$, there also holds

$$\begin{aligned} \left\| \tilde{W}_\tau(0) \right\|_{\mathcal{L}^2}^2 &\leq 2\|W_0\|_{\mathcal{L}^2}^2 + 2\alpha^2\|\Theta(0) - \Theta(\tau)\|_{L^2(\mathbb{R}^2)} + 2\alpha^2|g(0, 1) - g(\tau, 1)| \\ &\leq \|W_0\|_{\mathcal{L}^2}^2 + C\alpha^2\left(1 + \log(1 + (\chi t)^2)\right) \\ &\leq \|W_0\|_{\mathcal{L}^2}^2 + C\alpha^2\left(1 + \log(1 + |\alpha|) + \log(1 + t)\right), \end{aligned}$$

but also

$$\begin{aligned} \|W(t)\|_{\mathcal{L}^2}^2 &\leq 2\left\| \tilde{W}_\tau(t) \right\|_{\mathcal{L}^2}^2 + 2\alpha^2\|\Theta(t + \tau) - \Theta(t)\|_{\mathcal{L}^2}^2 \\ &\leq 2\left\| \tilde{W}_\tau(t) \right\|_{\mathcal{L}^2}^2 + \frac{\alpha^2}{2\pi} \log\left(1 + \frac{t}{1 + (\chi t)^2}\right) \\ &\leq 2\left\| \tilde{W}_\tau(t) \right\|_{\mathcal{L}^2}^2 + \frac{\alpha^2}{2\pi} \frac{t}{1 + (\chi t)^2} \\ &\leq 2\left\| \tilde{W}_\tau(t) \right\|_{\mathcal{L}^2}^2 + \frac{\alpha^2}{2\chi\pi}, \end{aligned}$$

and

$$\begin{aligned} \int_0^t \|D(w(s))\|_{L^2(\mathcal{F}_0)}^2 ds &\leq 2 \int_0^t \|D(\tilde{w}_\tau(s))\|_{L^2(\mathcal{F}_0)}^2 ds + 2\alpha^2 \int_0^t \|D(\Theta(\tau + s) - \Theta(s))\|_{L^2(\mathcal{F}_0)}^2 ds \\ &\leq 2 \int_0^t \|D(\tilde{w}_\tau(s))\|_{L^2(\mathcal{F}_0)}^2 ds + 2\alpha^2 \int_0^t \|\nabla(\Theta(\tau + s) - \Theta(s))\|_{L^2(\mathcal{F}_0)}^2 ds \\ &\leq 2 \int_0^t \|D(\tilde{w}_\tau(s))\|_{L^2(\mathcal{F}_0)}^2 ds + 2\kappa_1\alpha^2 \int_0^t \left(\frac{1}{1 + \tau} - \frac{1}{1 + \tau + s}\right) ds \\ &\leq 2 \int_0^t \|D(\tilde{w}_\tau(s))\|_{L^2(\mathcal{F}_0)}^2 ds + 2\kappa_1\alpha^2 \left(\frac{t}{1 + (\chi t)^2} - \log\left(1 + \frac{t}{1 + (\chi t)^2}\right)\right) ds \\ &\leq 2 \int_0^t \|D(\tilde{w}_\tau(s))\|_{L^2(\mathcal{F}_0)}^2 ds + 2\kappa_1 \frac{\alpha^2}{\chi}. \end{aligned}$$

The last five estimates put together (along with $\chi \geq 1$) lead to the result. \square

4. GLOBAL STABILITY FOR FINITE-ENERGY SOLUTIONS

This last section is devoted to the proof of **Theorem 1.1**. For this, we first recall the partial result in [4] on which relies our analysis:

Lemma 4.1 ([4, Theorem 1.3]). *Let $q \in (1, 2)$ and assume that $V_0 \in \mathcal{L}^q \cap \mathcal{L}^2$ with $\|V_0\|_{\mathcal{L}^2}$ sufficiently small. Then the unique finite-energy weak solution V with initial data V_0 satisfies:*

$$\sup_{t>0} t^{\frac{1}{p} - \frac{1}{q}} \|V(t)\|_{\mathcal{L}^p} < \infty \quad \forall p \in (2, \infty) \quad (4.1)$$

$$\sup_{t>0} t^{\frac{1}{q}} |\ell_v(t)| < \infty. \quad (4.2)$$

Theorem 1.1 is then a direct consequence of the two following propositions that we prove in the next subsections:

Proposition 4.2. *Let $q \in (1, 2)$ and assume that $V_0 \in \mathcal{L}^q \cap \mathcal{L}^2$. Then the unique finite-energy solution V starting from V_0 satisfies:*

$$V \in C([0, \infty); \mathcal{L}^q \cap \mathcal{L}^2) \quad (4.3)$$

$$\nabla V \in L^1_{loc}([0, \infty); L^q(\mathcal{F}_0) \cap L^2(\mathcal{F}_0)) \quad (4.4)$$

$$\ell_V \in L^2((0, \infty)). \quad (4.5)$$

Proposition 4.3. *Let $q \in (1, 2)$ and assume that $V_0 \in \mathcal{L}^q \cap \mathcal{L}^2$. Then the unique finite-energy solution V starting from V_0 satisfies:*

$$\liminf_{t \rightarrow \infty} \|V(t)\|_{\mathcal{L}^2} = 0. \quad (4.6)$$

4.1. Proof of Proposition 4.2

Let $q < 2$ and $V_0 \in \mathcal{L}^q \cap \mathcal{L}^2$. We recall that, by the construction of [13], we have $V \in C([0, \infty); \mathcal{L}^2)$ and $\nabla V \in L^2((0, \infty); L^2(\mathbb{R}^2))$. Furthermore, with the proof of **Theorem 3.4** we know that the solution V is computed through the Duhamel formula:

$$V(t) = S(t)V_0 + \int_0^t S(t-s)\mathbb{P}[\mathbf{1}_{\mathcal{F}_0}(V - \ell_V) \cdot \nabla V]ds. \quad (4.7)$$

since it is the only fixed point of the mapping:

$$\mathcal{D} : W \mapsto S(t)V_0 + \int_0^t S(t-s)\mathbb{P}[\mathbf{1}_{\mathcal{F}_0}(W - \ell_W) \cdot \nabla W]ds.$$

in the space $C([0, T]; \mathcal{L}^2) \cap C((0, T); H^1(\mathcal{F}_0))$ endowed with the X -norm:

$$\|W\|_X = \sup_{[0, T]} \|W\|_{\mathcal{L}^2} + \sup_{[0, T]} \sqrt{t} \|\nabla W\|_{L^2(\mathcal{F}_0)}$$

(for T sufficiently small). We show here that the same property holds adding the property $V \in C([0, T]; \mathcal{L}^q) \cap C((0, T); W^{1,q}(\mathcal{F}_0))$. Let fix B_T the subset in $C([0, T]; \mathcal{L}^2 \cap \mathcal{L}^q) \cap C((0, T); H^1(\mathcal{F}_0) \cap W^{1,q}(\mathcal{F}_0))$ containing W satisfying

$$\|W\|_X \leq 2\|V_0\|_{\mathcal{L}^2} \quad \|W\|_{X_q} := \sup_{[0, T]} \left(\|W(t)\|_{\mathcal{L}^q} + \sqrt{t} \|\nabla W(t)\|_{L^q(\mathbb{R}^2)} \right) \leq (1 + K_2)(\|V_0\|_{\mathcal{L}^q} + \|V_0\|_{\mathcal{L}^2}),$$

where K_2 is the constant involved in **Lemma 2.7**. By adapting the computations in the proof of **Theorem 3.4**, we obtain a time T_0 sufficiently small such that for $T < T_0$ the above mapping is a contraction on B_T for the X -norm. Then, given $W \in B_T$, applying the duality estimates in Lemma 2.8 with $p = q$ we obtain that

$$\|\mathcal{D}[W](t)\|_{\mathcal{L}^q} \leq \|V_0\|_{\mathcal{L}^q} + \int_0^t \phi_q(t-s) \|(W - \ell_W) \otimes W\|_{L^q(\mathcal{F}_0)} ds \quad \forall t \in [0, T]$$

where

$$\phi_q(s) = K_5 \begin{cases} s^{-1/2} & \text{if } s < 1 \\ s^{-1+1/q} & \text{if } s > 1 \end{cases}$$

The last integral we denote $I[W]$ is then bounded by applying the Gagliardo Nirenberg inequality:

$$\begin{aligned} I[W] &\leq \int_0^t |\phi_q(t-s)| \left(\|W\|_{\mathcal{L}^2}^{1/q} \|\nabla W\|_{L^2(\mathcal{F}_0)}^{2(1-1/q)} \right) + |\ell_W| \|W\|_{\mathcal{L}^q} ds \\ &\leq \int_0^t |\phi_q(t-s)| \left(s^{-(1-1/q)} \|W\|_X^2 + \|W\|_X \|W\|_{\mathcal{L}^q} \right) ds. \end{aligned}$$

At this point, we realize that, for $T < 1$ there is an absolute constant \tilde{K}_5 for which:

$$\sup_{[0, T]} \int_0^t \phi_q(t-s) s^{-(1-1/q)} \leq \tilde{K}_5 T^{1/q - 1/2} \quad \sup_{[0, T]} \int_0^t \phi_q(t-s) \leq \tilde{K}_5 \sqrt{T}.$$

Since $q < 1/2$ we can take T_0 smaller (but decreasingly in the quantity $\|V_0\|_{\mathcal{L}^2} + \|V_0\|_{\mathcal{L}^q}$) so that for $T < T_0$:

$$\sup_{[0, T]} \|\mathcal{D}[W](t)\|_{\mathcal{L}^q} \leq \|V_0\|_{\mathcal{L}^q} + \tilde{K}_5 T^{1/q - 1/2} \|V_0\|_{\mathcal{L}^2} \left(\|V_0\|_{\mathcal{L}^2} + \sup_{[0, T_1]} \|W\|_{\mathcal{L}^q} \right) \leq \|V_0\|_{\mathcal{L}^q} + \|V_0\|_{\mathcal{L}^2}. \quad (4.8)$$

As for the gradient, we apply semigroup estimates of Lemma 2.7 to yield that

$$\sqrt{t}\|\nabla\mathcal{D}[W](t)\|_{L^q(\mathcal{F}_0)} \leq K_2\|V_0\|_{\mathcal{L}^q} + K_2 \int_0^t \left(\frac{t}{t-s}\right)^{\frac{1}{2}} \|(W - \ell_w) \cdot \nabla W\|_{L^q(\mathcal{F}_0)} ds.$$

Combining then Hölder inequalities (where $1/q^* = 1/q - 1/2$) together with a Gagliardo-Nirenberg inequality (interpolating the L^{q^*} -norm between the L^2 and \dot{H}^1 norms) and the already obtained bound on $\|W\|_X$, we bound:

$$\begin{aligned} \|W \cdot \nabla W\|_{L^q(\mathcal{F}_0)} &\leq \|W\|_{L^{q^*}(\mathcal{F}_0)} \|\nabla W\|_{L^2(\mathcal{F}_0)} \leq s^{\frac{3}{2}-\frac{1}{q}} \|W\|_X^2 \\ \|\ell_w \cdot \nabla W\|_{L^q(\mathcal{F}_0)} &\leq \sqrt{s} \|W\|_X \|W\|_{X_q} \end{aligned}$$

Since $s < 1$, we end up with:

$$\sqrt{t}\|\nabla\mathcal{D}[W](t)\|_{L^q(\mathcal{F}_0)} \leq K_2\|V_0\|_{\mathcal{L}^q} + K_2 \int_0^t \left(\frac{t}{t-s}\right)^{\frac{1}{2}} ds \|V_0\|_{\mathcal{L}^2} (\|V_0\|_{\mathcal{L}^2} + \|V_0\|_{\mathcal{L}^q}).$$

By a homogeneity argument we have:

$$\int_0^t \left(\frac{t}{t-s}\right)^{\frac{1}{2}} ds \leq Ct^{\frac{3}{2}},$$

hence we can choose T_0 smaller if necessary (but decreasing in the quantity $\|V_0\|_{\mathcal{L}^2} + \|V_0\|_{\mathcal{L}^q}$) so that: for $T < T_0$:

$$\sup_{[0,T]} \sqrt{t}\|\nabla\mathcal{D}[W](t)\|_{L^q(\mathcal{F}_0)} \leq K_2(\|V_0\|_{\mathcal{L}^q} + \|V_0\|_{\mathcal{L}^2}).$$

Finally, \mathcal{D} maps B into B . With similar computations, we obtain that it is a contraction up to restrict to a smaller T_0 again and conclude that we propagate the property $V \in \mathcal{L}^q$ and $\nabla V \in L^q(\mathcal{F}_0)$ on a short time-interval. We note that on this time-interval ΔT , we have

$$\|\nabla W\|_{L^1(0,\Delta T);L^2(\mathcal{F}_0)} \leq \|W\|_X \quad \|\nabla W\|_{L^1(0,\Delta T);L^q(\mathcal{F}_0)} \leq \|W\|_{X_q} \quad (4.9)$$

To obtain further that $V \in \mathcal{L}^q$ and $\nabla V \in L^q(\mathcal{F}_0)$ for all times we remark that by a standard blow-up alternative, it is sufficient to obtain local bounds for $\|V(t)\|_{\mathcal{L}^q} + \|V(t)\|_{\mathcal{L}^2}$. Since this is already known for $\|V(t)\|_{\mathcal{L}^2}$ we focus here on $\|V(t)\|_{\mathcal{L}^q}$. To this end, we note that choosing T_1 so that $\tilde{K}_5 T_1^{\frac{1}{q}-\frac{1}{2}} \|V_0\|_{\mathcal{L}^2} \leq 1/2$ and applying (4.8) with V we have

$$\sup_{[0,T_1]} \|V(t)\|_{\mathcal{L}^q} \leq 2\|V_0\|_{\mathcal{L}^q} + \|V_0\|_{\mathcal{L}^2}.$$

Furthermore, since our system of equation is autonomous, we can reproduce this computation starting from any $t_0 > 0$. Finally, since we already have a uniform bounds for $\|V(t_0)\|_{\mathcal{L}^2}$ we obtain that there exists a short time increment T_1 (independent of the initial data) so that for arbitrary $t_0 > 0$:

$$\sup_{[t_0,t_0+T_1]} \|V(t)\|_{\mathcal{L}^q} \leq 2\|V(t_0)\|_{\mathcal{L}^q} + \|V_0\|_{\mathcal{L}^2}.$$

In particular, there can be no blow-up of $\|V(t)\|_{\mathcal{L}^q}$ in finite-time. Then on a time-interval $[0, T]$ since we have an *a priori* bound for $\|V\|_{\mathcal{L}^q} + \|V\|_{\mathcal{L}^2}$, we can see our solution as a concatenation of local-in-time solutions constructed as above on a small-time interval ΔT . By concatenating the remarks (4.9) on the time-intervals $[n\Delta T, (n+1)\Delta T]$ we conclude that

$$\nabla V \in L^1((0, T); L^2(\mathcal{F}_0) \cap L^q(\mathcal{F}_0)).$$

To complete the proof of **Proposition 4.2**, we show now that $\ell_V \in L^2([0, \infty))$. Since $\nabla V \in L^2([0, \infty))$, we first remark that:

$$\forall \varepsilon > 0, \quad \exists T_\varepsilon > 0 \text{ s.t. } \int_{T_\varepsilon}^\infty \|\nabla V\|_{L^2}^2 < \varepsilon. \quad (4.10)$$

Thanks to the representation formula (4.7), we have then that, for arbitrary $t > 0$ we can split $\ell_v(t) = \ell_S(t) + \ell_{NL}(t)$ where:

$$\ell_S(t) = \ell_{S(t)V_0} \quad \ell_{NL}(t) = \ell_{I(t)}$$

where

$$I(t) = \int_0^t S(t-s) \mathbb{P}[\mathbf{1}_{\mathcal{F}_0}(V - \ell_V) \cdot \nabla V] ds.$$

Since $V_0 \in \mathcal{L}^2 \cap \mathcal{L}^q$ we apply **Lemma 2.7** to yield that:

$$|\ell_S(t)| \leq \min\left(1, \frac{1}{t^{\frac{1}{q}}}\right) \|V_0\|_{\mathcal{L}^q} \in L^2((0, \infty)).$$

For the nonlinear term, we apply the duality estimates of **Lemma 2.7** with $r > 2$. We obtain:

$$|\ell_{NL}(t)| \leq \int_0^t \frac{1}{(t-s)^{\frac{1}{2} + \frac{1}{r}}} \|(V - \ell_v) \otimes V\|_{L^r(\mathcal{F}_0)}$$

At this point, let fix $T > 0$ (sufficiently large) and remark that the right-hand side can be seen as a truncated (time-)convolution of $1/s^{\frac{1}{2} + \frac{1}{r}}$ and $\|(V - \ell_v) \otimes V\|_{L^r(\mathcal{F}_0)} \mathbf{1}_{[0, T]}$. By a Hardy-Littlewood-Sobolev inequality, we have then:

$$\begin{aligned} \|\ell_{NL}\|_{L^2(0, T)} &\leq \| |\cdot|^{-\left(\frac{1}{2} + \frac{1}{r}\right)} * \|(V - \ell_v) \otimes V\|_{L^r(\mathcal{F}_0)} \|_{L^2(\mathbb{R})} \\ &\leq C_r \left(\int_0^T \|(V - \ell_v) \otimes V\|_{L^r(\mathcal{F}_0)}^p \right)^{\frac{1}{p}} \leq C_r \left(\int_0^T \|V\|_{L^{2r}(\mathcal{F}_0)}^{2p} \right)^{\frac{1}{p}} + C_r \left(\int_0^T \|\ell_v|V\|_{L^r(\mathcal{F}_0)}^p \right)^{\frac{1}{p}} \end{aligned}$$

where p is the conjugate exponent of r . For the first-integral on the right-hand side, we apply again a Gagliardo-Nirenberg inequality and the fact that p is the conjugate exponent of r to yield that:

$$\int_0^T \|V\|_{L^{2r}(\mathcal{F}_0)}^{2p} \leq C_r \sup_{[0, T]} \|V(t)\|_{\mathcal{L}^2}^{\frac{2p}{r}} \int_0^T \|\nabla V\|_{L^2}^2 \leq C_r \|V_0\|_{\mathcal{L}^2}^{2p}$$

To estimate the last term, we introduce an intermediate time T_{mid} to be fixed later on. We note here that, for arbitrary $0 \leq T_1 < T_2$ combining a standard Hölder inequality and a Gagliardo Nirenberg inequality entails that (since $p < 2$):

$$\begin{aligned} \int_{T_1}^{T_2} \|\ell_v|V\|_{L^r(\mathcal{F}_0)}^p &\leq \left(\int_{T_1}^{T_2} |\ell_v|^2 \right)^{\frac{p}{2}} \left(\int_{T_1}^{T_2} \|V\|_{L^{\frac{2p}{2-p}}}^{2p} \right)^{1 - \frac{p}{2}} \\ &\leq \|\ell_v\|_{L^2(T_1, T_2)}^p \sup_{(T_1, T_2)} \|V\|_{L^2}^{\frac{2p}{r}} \left(\int_{T_1}^{T_2} \|\nabla V\|_{L^2}^{\frac{2p}{2-p}(1 - \frac{2}{r})} \right)^{1 - \frac{p}{2}} \end{aligned}$$

Recalling that p and r are conjugate exponents yield that

$$\frac{2p}{2-p} \left(1 - \frac{2}{r}\right) = 2$$

and we infer that:

$$\int_{T_1}^{T_2} \|\ell_v|V\|_{L^r(\mathcal{F}_0)}^p \leq C \|\ell_v\|_{L^2(T_1, T_2)}^p \|V_0\|_{\mathcal{L}^2}^{\frac{2p}{r}} \left(\int_{T_1}^{T_2} \|\nabla V\|_{L^2}^2 \right)^{1 - \frac{p}{2}}$$

When $T > T_{mid}$, combining the previous computations between $T_1 = 0$ and $T_2 = T_{mid}$ and between $T_1 = T_{mid}$ and $T_2 = T$, we conclude that:

$$\begin{aligned} \|\ell_{NL}\|_{L^2(0, T)} &\leq C_r \|V_0\|_{\mathcal{L}^2}^2 + C_r \|\ell_v\|_{L^2(0, T_{mid})} \|V_0\|_{\mathcal{L}^2}^{\frac{2}{r}} \left(\int_0^{T_{mid}} \|\nabla V\|_{L^2}^2 \right)^{\frac{1}{p} - \frac{1}{2}} \\ &\quad + C_r \|\ell_v\|_{L^2(T_{mid}, T)} \|V_0\|_{\mathcal{L}^2}^{\frac{2}{r}} \left(\int_{T_{mid}}^T \|\nabla V\|_{L^2}^2 \right)^{\frac{1}{p} - \frac{1}{2}}. \end{aligned}$$

At this point, we recall the remark (4.10) and choose T_{mid} so that:

$$C_r \|V_0\|_{\mathcal{L}^2}^{\frac{2p}{r}} \left(\int_{T_{mid}}^{\infty} \|\nabla V\|_{L^2}^2 \right)^{\frac{1}{p} - \frac{1}{2}} < \frac{1}{2}$$

Splitting $\ell = \ell_S + \ell_{NL}$ and arguing that, on compact time-interval, we can always control $|\ell_v|$ by $\|V\|_{\mathcal{L}^2} \leq \|V_0\|_{\mathcal{L}^2}$, we infer that :

$$\|\ell_v\|_{L^2(0,T)} \leq \|\ell_S\|_{L^2(0,\infty)} + C_r(1 + \sqrt{T_{mid}}) \|V_0\|_{\mathcal{L}^2}^2 + \frac{1}{2} \|\ell_v\|_{L^2(0,T)}.$$

Eventually, we conclude that, for arbitrary $T > T_{mid}$ we have:

$$\|\ell_v\|_{L^2(0,T)} \leq C_{q,r} \left(\|V_0\|_{\mathcal{L}^q} + (1 + \sqrt{T_{mid}}) \|V_0\|_{\mathcal{L}^2}^2 \right).$$

This concludes the proof.

4.2. Proof of Proposition 4.3

This proof is inspired of [7, Section 5]. Let $q < 2$ and $V \in \mathcal{L}^2 \cap \mathcal{L}^q$. We recall that we take $\mu < \frac{1}{q} - \frac{1}{2}$ so that $\mathcal{L}^2 \cap \mathcal{L}^q \subset D(A^{-\mu})$. Thanks to **Proposition 4.2**, we have that the unique finite-energy solution satisfies

- $V \in C([0, \infty); \mathcal{L}^q \cap \mathcal{L}^2) \cap C((0, \infty); W^{1,q}(\mathbb{R}^2) \cap H^1(\mathbb{R}^2))$.
- $\mathbf{1}_{\mathcal{F}_0}(V - \ell_v) \cdot \nabla V \in L^1_{loc}((0, \infty); L^2(\mathbb{R}^2) \cap L^q(\mathbb{R}^2))$

In particular, we have:

$$\partial_t V - AV = \mathbb{P}((V - \ell_v) \cdot \nabla V) \quad V|_{t=0} = V_0$$

where $V \in C([0, \infty); D(A^{-\mu}))$ and $\mathbb{P}((V - \ell_v) \cdot \nabla V) \in L^1_{loc}(0, \infty; D(A^{-\mu}))$. Consequently, we can apply the operator $A^{-\mu}$ to this equation and we obtain that $U = A^{-\mu}V$ is a mild solution to:

$$\partial_t U - AU = A^{-\mu} \mathbb{P}((V - \ell_v) \cdot \nabla V) \quad U|_{t=0} = U_0.$$

We have in particular for arbitrary $t > 0$ that

$$\frac{1}{2} \|U(t)\|_{\mathcal{L}^2}^2 + \int_0^t \|\nabla U(s)\|_{L^2}^2 ds \leq \int_0^t \langle A^{-\mu} \mathbb{P}((V - \ell_v) \cdot \nabla V), U \rangle ds.$$

However, for arbitrary $s \in (0, t)$ there holds:

$$\begin{aligned} |\langle A^{-\mu} \mathbb{P}((V - \ell_v) \cdot \nabla V), U \rangle| &= \left| \int_{\mathcal{F}_0} ((V - \ell_v) \cdot \nabla) A^{-\mu} U \cdot V dx \right| \\ &\leq \left(\|V\|_{L^4(\mathcal{F}_0)}^2 + \|V\|_{L^2} |\ell_v| \right) \|A^{\frac{1}{2}-\mu} U\|_{\mathcal{L}^2} \\ &\leq C \left(|\ell_v| + \|\nabla V\|_{L^2(\mathcal{F}_0)} \right) \|A^\mu U\|_{\mathcal{L}^2} \|A^{\frac{1}{2}-\mu} U\|_{\mathcal{L}^2}, \end{aligned}$$

where we applied a Gagliardo-Nirenberg inequality to pass from the second to the last line. At this point, we argue by interpolation that

$$\|A^\mu U\|_{\mathcal{L}^2} \|A^{\frac{1}{2}-\mu} U\|_{\mathcal{L}^2} \leq C \|U\|_{\mathcal{L}^2} \|A^{\frac{1}{2}} U\|_{\mathcal{L}^2},$$

thus

$$C \left(|\ell_v| + \|\nabla V\|_{L^2(\mathcal{F}_0)} \right) \|U\|_{\mathcal{L}^2} \|A^{\frac{1}{2}} U\|_{\mathcal{L}^2} \leq C \left(|\ell_v| + \|\nabla V\|_{L^2(\mathcal{F}_0)} \right)^2 \|U\|_{\mathcal{L}^2}^2 + \frac{1}{2} \|A^{\frac{1}{2}} U\|_{\mathcal{L}^2}^2.$$

This yields finally that, for all $t \geq 0$:

$$\|U(t)\|_{\mathcal{L}^2}^2 + \int_0^t \|\nabla U(s)\|_{L^2}^2 ds \leq C \int_0^t \left(|\ell_v| + \|\nabla V\|_{L^2(\mathcal{F}_0)} \right)^2 \|U(s)\|_{\mathcal{L}^2}^2 ds.$$

Eventually a Gronwall lemma yields that:

$$\|U(t)\|_{\mathcal{L}^2}^2 + \int_0^t \|\nabla U(s)\|_{L^2}^2 ds \leq \|U_0\|_{\mathcal{L}^2}^2 \exp \left[\int_0^t C \left(|\ell_v| + \|\nabla V\|_{L^2(\mathcal{F}_0)} \right)^2 ds \right]$$

Since the integral in the exponential is bounded by Proposition 4.2, we have then a uniform bound

$$\sup_{t>0} \|U(t)\|_{\mathcal{L}^2}^2 + \int_0^\infty \|\nabla U(s)\|_{L^2}^2 ds \leq C_0$$

where the constant C_0 depends on the whole solution V (and *a priori* not only on V_0).

At this point, we argue in the same manner as in [7, Corollary 4.2]. The situation is even more favorable since we have uniform bounds. Indeed, since $\nabla U \in L^2((0, \infty); L^2(\mathbb{R}^2))$, we can construct a sequence of times t_n growing to infinity such that $\|\nabla U(t_n)\|_{L^2} \rightarrow 0$. We have then that $\|A^{1/2}U(t_n)\|_{\mathcal{L}^2}$ goes to 0 while $\|U(t_n)\|_{\mathcal{L}^2}$ remains bounded. By interpolation, $\|V(t_n)\|_{\mathcal{L}^2} = \|A^\mu U(t_n)\|_{\mathcal{L}^2}$ (where $\mu < 1/2$) goes also to 0 as n goes to infinity. This ends the proof.

A. TECHNICAL LEMMAS

We gather in this section technical lemmas used throughout the paper. We start with handling nonlinearities in the Duhamel formula. We recall that, given $t_0 > 0$, we denote:

$$X := C([t_0, \infty); \mathcal{L}^2) \cap C((t_0, \infty); H^1(\mathbb{R}^2) \cap L^\infty(B_0))$$

that we endow with the norm:

$$\|W\|_X := \sup_{t \geq t_0} \|W(t)\|_{\mathcal{L}^2} + \sup_{t > t_0} (t - t_0)^{\frac{1}{2}} \left(\|\nabla w(t)\|_{L^2(\mathcal{F}_0)} + |\ell_W(t)| \right).$$

Other notations are introduced in Section 3.

Lemma A.1. *Let $t_0 > 0$. Given $(W_a, W_b) \in X$ we denote:*

$$F(t) := \int_{t_0}^t S(t-s) \mathbb{P}[\mathbf{1}_{\mathcal{F}_0}(w_a - \ell_a) \cdot \nabla w_b] ds \quad \forall t \geq t_0.$$

Then there holds:

- $F \in X$
- *there exists a constant $C > 0$ for which:*

$$\|F\|_X \leq C \|W_a\|_X \|W_b\|_X$$

We emphasize that, in this lemma, the assumption $W_a \in X$ induces that, for every $s \geq t_0$, $W_a(s)$ is a rigid motion on B_0 . Obviously, we denote ℓ_a the translation velocity (with respect to the origin) associated with this motion.

Proof. We only give a proof of the second item. To this end, we remark that, since w_a is divergence free:

$$(w_a - \ell_a) \cdot \nabla w_b = \operatorname{div}((w_a - \ell_a) \otimes w_b), \quad \text{on } \mathcal{F}_0.$$

Since $(w_a - \ell_a) \cdot n = 0$ on ∂B we can then extend by 0 to create an $L^2(\mathbb{R}^d)$ -source term which fulfills the assumptions of [4, Corollary 3.10]. This yields, for arbitrary $t > t_0$

$$\|F(t)\|_{\mathcal{L}^2} \leq K \int_{t_0}^t \frac{1}{\sqrt{t-s}} \|(w_a - \ell_a) \otimes w_b\|_{L^2(\mathcal{F}_0)} ds$$

$$\begin{aligned}
&\leq K \int_{t_0}^t \frac{1}{\sqrt{t-s}} \left(|\ell_a| \|w_b\|_{L^2} + \|w_a\|_{L^2}^{1/2} \|\nabla w_a\|_{L^2(\mathcal{F}_0)}^{1/2} \|w_b\|_{L^2}^{1/2} \|\nabla w_b\|_{L^2(\mathcal{F}_0)}^{1/2} \right) \\
&\leq K \int_{t_0}^t \frac{1}{\sqrt{t-s}} \frac{1}{\sqrt{s-t_0}} ds \|w_a\|_X \|w_b\|_X.
\end{aligned}$$

Hence, we have $\|F(t)\|_{\mathcal{L}^2} \leq C \|w_a\|_X \|w_b\|_X$ for arbitrary $t \geq t_0$. For the second part, we split $F = F_1 + F_2 + F_3$ (with $t_{mid} = (t + t_0)/2$):

$$\begin{aligned}
F_1(t) &= \int_{t_0}^{t_{mid}} S(t-s) \mathbb{P} \operatorname{div}[\mathbf{1}_{\mathcal{F}_0}(w_a - \ell_a) \otimes w_b] ds \\
F_2(t) &= \int_{t_{mid}}^t S(t-s) \mathbb{P}[w_a \cdot \nabla w_b] ds \\
F_3(t) &= \int_{t_{mid}}^t S(t-s) \mathbb{P}[\ell_a \cdot \nabla w_b] ds.
\end{aligned}$$

□

For the first term, we combine [4, Corollary 3.10] with standard continuity properties of S . Remarking that:

$$F_1(t) = S(t - t_{mid}) \int_{t_0}^{t_{mid}} S(t_{mid} - s) \mathbb{P} \operatorname{div}[\mathbf{1}_{\mathcal{F}_0}(w_a - \ell_a) \otimes w_b] ds,$$

we obtain with obvious notations and similar computations that:

$$\begin{aligned}
|\ell_1(t)| + \|\nabla F_1(t)\|_{L^2(\mathcal{F}_0)} &\leq \frac{1}{\sqrt{t-t_{mid}}} \left\| \int_{t_0}^{t_{mid}} S(t_{mid} - s) \mathbb{P} \operatorname{div}[\mathbf{1}_{\mathcal{F}_0}(w_a - \ell_a) \otimes w_b] ds \right\|_{\mathcal{L}^2} \\
&\leq \frac{C}{\sqrt{t-t_0}} \int_{t_0}^{t_{mid}} \frac{1}{\sqrt{t_{mid}-s}} \|(w_a - \ell_a) \otimes w_b\|_{L^2(\mathcal{F}_0)} ds \\
&\leq \frac{C}{\sqrt{t-t_0}} \|w_a\|_X \|w_b\|_X.
\end{aligned}$$

For the other terms, we apply standard continuity properties of S . First we note that $w_a \cdot \nabla w_b \in L^{4/3}(\mathbb{R}^2)$ with:

$$\begin{aligned}
\|w_a \cdot \nabla w_b\|_{L^{4/3}(\mathbb{R}^2)} &\leq \|w_a\|_{L^2}^{1/2} \|\nabla w_a\|_{L^2(\mathbb{R}^2)}^{1/2} \|\nabla w_b\|_{L^2(\mathbb{R}^2)} \\
&\leq \frac{1}{(t-t_0)^{3/4}} \|w_a\|_X \|w_b\|_X
\end{aligned}$$

Since $\mathbb{P} : L^{4/3}(\mathbb{R}^2) \rightarrow \mathcal{L}^{4/3}$ is bounded (see [14, Remark 2.4]), we infer that:

$$\begin{aligned}
|\ell_2(t)| + \|\nabla F_2(t)\|_{L^2(\mathcal{F}_0)} &\leq K \int_{t_{mid}}^t \frac{1}{(t-s)^{3/4}} \frac{1}{(t-t_0)^{3/4}} ds \|w_a\|_X \|w_b\|_X \\
&\leq \frac{C}{\sqrt{t-t_0}} \|w_a\|_X \|w_b\|_X.
\end{aligned}$$

Finally, we bound (applying the standard continuity of $\mathbb{P} : L^2(\mathbb{R}^2) \rightarrow \mathcal{L}^2$)

$$\begin{aligned}
|\ell_3(t)| + \|\nabla F_3(t)\|_{L^2(\mathcal{F}_0)} &\leq K \int_{t_{mid}}^t \frac{1}{\sqrt{t-s}} |\ell_a| \|\nabla w_b\|_{L^2(\mathbb{R}^2)} ds \\
&\leq K \int_{t_{mid}}^t \frac{1}{\sqrt{t-s}} \frac{ds}{s-t_0} \|w_a\|_X \|w_b\|_X \\
&\leq \frac{C}{\sqrt{t-t_0}} \|w_a\|_X \|w_b\|_X.
\end{aligned}$$

B. PROOF OF PROPOSITION 2.12

We provide here a proof of Proposition 2.12. To estimate $(\tilde{\mathcal{A}} + \varepsilon)^{-\mu}$ as required in the content of Proposition 2.12 we rely on the integral representation (because $\tilde{\mathcal{A}}$ is a positive selfadjoint operator, see [11, Section 2.6]):

$$(\tilde{\mathcal{A}} + \varepsilon)^{-\mu} = \frac{\sin(\pi\mu)}{\pi} \int_0^\infty \frac{1}{(\lambda + \varepsilon)^\mu} (\tilde{\mathcal{A}} + \lambda + \varepsilon)^{-1} d\lambda. \quad (\text{B.1})$$

In order to construct $R_{\mu,\varepsilon}$ we work at first on a construction of $(\tilde{\mathcal{A}} + \lambda)^{-1}$ involving $(\tilde{\mathcal{A}}_0 + \lambda)^{-1}$ for $\lambda > 0$. To this end, we introduce objects that are crucial to the analysis.

We recall here basics on some modified Bessel functions. The following statements are taken from [10, Section 8]. The function $K_0 : (0, \infty) \rightarrow \mathbb{R}$ is the unique smooth solution to:

$$-\frac{1}{r} \frac{d}{dr} \left[r \frac{d}{dr} K_0(r) \right] + K_0(r) = 0 \quad \forall r > 0,$$

that behaves asymptotically like:

$$K_0(r) \sim \begin{cases} \sqrt{\frac{\pi}{2r}} \exp(-r) & \text{when } r \rightarrow \infty \\ -\ln(r) & \text{when } r \rightarrow 0. \end{cases}$$

Furthermore, all derivatives of K_0 enjoy the same decay at infinity as K_0 and $K_0'(r) \sim -1/r$ in 0. We mention also that $K_0 \geq 0$ and $K_0' \leq 0$ on $(0, \infty)$ (see [10, Theorem 8.1]). Similarly, $K_1 : (0, \infty) \rightarrow \mathbb{R}$ is the smooth solution to:

$$-\frac{1}{r} \frac{d}{dr} \left[r \frac{d}{dr} K_1(r) \right] + \left(1 + \frac{1}{r^2} \right) K_1(r) = 0 \quad \forall r > 0,$$

that has the asymptotic expansion:

$$K_1(r) \sim \begin{cases} \sqrt{\frac{\pi}{2r}} \exp(-r) & \text{when } r \rightarrow \infty \\ \frac{1}{r} & \text{when } r \rightarrow 0 \end{cases}$$

We have again that $K_1 \geq 0$ and $K_1' \leq 0$ on $(0, \infty)$, that the derivatives of K_1 enjoy the same decay as K_1 at infinity and $K_1'(r) \sim -1/r^2$ in 0.

Then, for arbitrary $\lambda > 0$, we define $\phi_\lambda : \mathbb{R}^2 \rightarrow \mathbb{R}$ by:

$$\phi_\lambda(x) = \begin{cases} \frac{K_0(\sqrt{\lambda}|x|)}{\lambda K_0(\sqrt{\lambda}) - \frac{2\pi}{m} \sqrt{\lambda} K_0'(\sqrt{\lambda})} & \text{if } |x| > 1 \\ \frac{K_0(\sqrt{\lambda})}{\lambda K_0(\sqrt{\lambda}) - \frac{2\pi}{m} \sqrt{\lambda} K_0'(\sqrt{\lambda})} & \text{if } |x| \leq 1 \end{cases}$$

and $\psi_\mu : \mathbb{R}^2 \rightarrow \mathbb{R}$ by

$$\psi_\lambda(x) = \begin{cases} \frac{K_1(\sqrt{\lambda}|x|)}{(2\mathcal{J}^{-1} + \lambda)K_1(\sqrt{\lambda}) - \pi\mathcal{J}^{-1}\sqrt{\lambda}K_1'(\sqrt{\lambda})} & \text{if } |x| > 1 \\ \frac{K_1(\sqrt{\lambda})|x|}{(2\mathcal{J}^{-1} + \lambda)K_1(\sqrt{\lambda}) - 2\pi\mathcal{J}^{-1}\sqrt{\lambda}K_1'(\sqrt{\lambda})} & \text{if } |x| \leq 1 \end{cases}$$

We recall that the symbols m and \mathcal{J} appearing in these formulas stand respectively for the mass and inertia of the disk. The aim of this construction is the following proposition:

Proposition B.1. Let $\lambda > 0$. Given $(F, \tau) \in \mathbb{R}^2 \times \mathbb{R}$, let define:

$$V_\lambda[F, \tau](x) = \phi_\lambda(x)F + \psi_\lambda(x)\tau \frac{x^\perp}{|x|} \quad \forall x \in \mathbb{R}^2.$$

Then there holds $V_\lambda[F, \tau] \in \mathcal{D}(\tilde{\mathcal{A}})$ with:

$$(\tilde{\mathcal{A}} + \lambda)V_\lambda[F, \tau] = (F + \tau x^\perp)\mathbb{1}_{B_0}.$$

Proof. Let $\lambda > 0$ and $(F, \tau) \in \mathbb{R}^2 \times \mathbb{R}$. For the proof, we denote $V = V_\lambda[F, \tau]$ for legibility. By construction, ψ_λ and ϕ_λ are continuous on \mathbb{R}^2 . Furthermore, since K_0, K_1 are smooth and decay exponentially at infinity, we have that $\phi_\lambda, \psi_\lambda \in H^2(\mathcal{F}_0)$. The explicit values for ϕ_λ and ψ_λ when $r < 1$ yield also that, on $B(0, 1)$, we have:

$$V(x) = \frac{K_0(\sqrt{\lambda})}{\lambda K_0(\sqrt{\lambda}) - \frac{2\pi}{m}\sqrt{\lambda}K_0'(\sqrt{\lambda})}F + \frac{K_1(\sqrt{\lambda})\tau}{(2\mathcal{J}^{-1} + \lambda)K_1(\sqrt{\lambda}) - 2\pi\mathcal{J}^{-1}\sqrt{\lambda}K_1'(\sqrt{\lambda})}x^\perp \text{ on } B_0.$$

Finally, we obtain that $V \in L^2[B_0] \cap [H^1(\mathbb{R}^2)]^2$ and thus that $V \in \mathcal{D}(\tilde{\mathcal{A}})$.

We go now to polar coordinates (r, θ) and exploit the ODE satisfied by K_0, K_1 to obtain that

$$-\Delta V + \lambda V = (\tilde{\mathcal{A}} + \lambda)V = 0 \quad \text{in } \mathcal{F}_0.$$

This is why we introduced Bessel functions. While, in B_0 , we have:

$$(\tilde{\mathcal{A}} + \lambda)V = \ell + \omega y^\perp$$

with

$$\begin{aligned} \ell &= \frac{1}{m} \int_{\partial B_0} \partial_n v d\sigma + \frac{\lambda K_0(\sqrt{\lambda})}{\lambda K_0(\sqrt{\lambda}) - \frac{2\pi}{m}\sqrt{\lambda}K_0'(\sqrt{\lambda})}F \\ \omega &= \mathcal{J}^{-1} \left(\int_{\partial B_0} z^\perp \partial_n v d\sigma + \frac{2K_1(\sqrt{\lambda})\tau}{(2 + \lambda)K_1(\sqrt{\lambda}) - 2\pi\mathcal{J}^{-1}\sqrt{\lambda}K_1'(\sqrt{\lambda})} \right) \\ &\quad + \frac{\lambda K_1(\sqrt{\lambda})\tau}{(2\mathcal{J}^{-1} + \lambda)K_1(\sqrt{\lambda}) - 2\pi\mathcal{J}^{-1}\sqrt{\lambda}K_1'(\sqrt{\lambda})}. \end{aligned}$$

Going again to polar coordinates (r, θ) , we note that $\partial_n = -\partial_r$ and that $z^\perp = (-\sin(\theta), \cos(\theta))$. For symmetry reasons, we thus have that:

$$\begin{aligned} \int_{\partial B_0} \partial_n v d\sigma &= -\frac{2\pi\sqrt{\lambda}K_0'(\sqrt{\lambda})}{\lambda K_0(\sqrt{\lambda}) - \frac{2\pi}{m}\sqrt{\lambda}K_0'(\sqrt{\lambda})}F \\ \int_{\partial B_0} z^\perp \partial_n v d\sigma &= -\frac{2\pi\sqrt{\lambda}K_1'(\sqrt{\lambda})}{(2\mathcal{J}^{-1} + \lambda)K_1(\sqrt{\lambda}) - 2\pi\mathcal{J}^{-1}\sqrt{\lambda}K_1'(\sqrt{\lambda})}\tau \end{aligned}$$

Introducing these identities in the above computations of ℓ and ω , we end up with $\ell = F$ and $\omega = \tau$. This concludes the proof. \square

We combine now this construction with the operator $\tilde{\mathcal{A}}_0$ to compute the resolvent of $\tilde{\mathcal{A}}$. Given $\lambda > 0$ and $W \in L^2[B_0]$, we have:

$$W = (\ell_W + \omega_W y^\perp)\mathbb{1}_{B_0} + w\mathbb{1}_{\mathcal{F}_0}.$$

Consider $V_\lambda^{(0)}[W] = (\tilde{\mathcal{A}}_0 + \lambda)^{-1}(w\mathbb{1}_{\mathcal{F}_0})$. We have $V_\lambda^{(0)} \in \mathcal{D}(\tilde{\mathcal{A}}_0) \subset \mathcal{D}(\tilde{\mathcal{A}})$ so that we can compute $\tilde{\mathcal{A}}V_\lambda^{(0)}$:

$$(\tilde{\mathcal{A}} + \lambda)(V_\lambda^{(0)}[W]) = \left[\frac{1}{m} \left(\int_{\partial B_0} \partial_n v_\lambda^{(0)}[W] d\sigma \right) + \mathcal{J}^{-1} \left(\int_{\partial B_0} z^\perp \cdot \partial_n v_\lambda^{(0)}[W] d\sigma \right) y^\perp \right] \mathbb{1}_{B_0} + w\mathbb{1}_{\mathcal{F}_0}.$$

Consequently, we correct the value on B_0 by setting:

$$F_\lambda^{(0)}[W] := \frac{1}{m} \left(\int_{\partial B_0} \partial_n v_\lambda^{(0)}[W] d\sigma \right), \quad \tau_\lambda^{(0)}[W] := \mathcal{J}^{-1} \left(\int_{\partial B_0} z^\perp \cdot \partial_n v_\lambda^{(0)}[W] d\sigma \right), \quad (\text{B.2})$$

and

$$\tilde{V}_\lambda[W] := V_\lambda^{(0)}[W] + V_\lambda[\ell_W - F_\lambda^{(0)}[W], \omega_W - \tau_\lambda^{(0)}[W]]. \quad (\text{B.3})$$

By linearity, we obtain that $\tilde{V}_\lambda \in \mathcal{D}(\tilde{\mathcal{A}})$ satisfies:

$$(\tilde{\mathcal{A}} + \lambda)\tilde{V}_\lambda[W] = W.$$

and is the unique one by injectivity of $\tilde{\mathcal{A}} + \lambda$.

With this construction at-hand, we are in position to prove Proposition 2.12.

Proof of Proposition 2.12. Fix $q \in (1, 2)$ and $\mu < \mu_{crit} = 1/q - 1/2$. Let $\varepsilon > 0$ and $W \in L^2[B_0]$. Plugging (B.3) into (B.1) we obtain that

$$\begin{aligned} (\tilde{\mathcal{A}} + \varepsilon)^{-\mu} &= \frac{\sin(\pi\mu)}{\pi} \int_0^\infty \frac{1}{(\lambda + \varepsilon)^\mu} (\tilde{\mathcal{A}}_0 + \varepsilon + \lambda)^{-1} (\mathbf{1}_{\mathcal{F}_0} W) d\lambda \\ &\quad + \frac{\sin(\pi\mu)}{\pi} \int_0^\infty \frac{1}{(\lambda + \varepsilon)^\mu} V_{\varepsilon+\lambda}[\ell_W - F_{\lambda+\varepsilon}^{(0)}[W], \omega_W + \tau_{\lambda+\varepsilon}^{(0)}[W]] d\lambda. \end{aligned}$$

Thus, we have the expected representation formula with:

$$R_{\mu,\varepsilon} W = \frac{\sin(\pi\mu)}{\pi} \int_0^\infty \frac{1}{(\lambda + \varepsilon)^\mu} V_{\varepsilon+\lambda}[\ell_W - F_{\lambda+\varepsilon}^{(0)}[W], \omega_W + \tau_{\lambda+\varepsilon}^{(0)}[W]] d\lambda.$$

To complete the proof, it remains to obtain (2.8). For this, we first bound by introducing the explicit value of $V_{\varepsilon+\lambda}$:

$$\begin{aligned} \|R_{\mu,\varepsilon} W\|_{L^2} &\leq C \int_0^\infty \frac{1}{(\lambda + \varepsilon)^\mu} \|\phi_{\lambda+\varepsilon}\|_{L^2} (|\ell_W| + |F_{\lambda+\varepsilon}^{(0)}[W]|) d\lambda \\ &\quad + C \int_0^\infty \frac{1}{(\lambda + \varepsilon)^\mu} \|\psi_{\lambda+\varepsilon}\|_{L^2} (|\omega_W| + |\tau_{\lambda+\varepsilon}^{(0)}[W]|) d\lambda. \end{aligned}$$

We note here that the constant C appearing in the right-hand side depends only on the physical parameters of the system. We denote by C such constants below. They can depend on the physical parameters or on the data q, μ . They can also vary between lines.

We proceed by estimating the two integrals independently. For the first one, let denote:

$$K(s) = \frac{1}{s^\mu} \|\phi_s\|_{L^2(\mathbb{R}^d)} (|\ell_W| + |F_s^{(0)}[W]|)$$

By looking at the explicit value of ϕ_s , we have:

$$\|\phi_s\|_{L^2(\mathbb{R}^d)} \leq \frac{C}{sK_0(\sqrt{s}) - \sqrt{s}K_0'(\sqrt{s})} \left(K_0(\sqrt{s}) + \frac{1}{\sqrt{s}} \left(\int_{\sqrt{s}}^\infty |K_0(\alpha)|^2 \alpha d\alpha \right)^{\frac{1}{2}} \right).$$

and, with q' the conjugate exponent of q :

$$\|F_s^{(0)}[W]\| \leq C \frac{\|w\|_{L^q(\mathcal{F}_0)}}{K_0(\sqrt{s})s^{\frac{1}{q'}}} \left(\int_{\sqrt{s}}^\infty |K_0(\alpha)|^{q'} \alpha d\alpha \right)^{\frac{1}{q'}}. \quad (\text{B.4})$$

We postpone the proof of this latter inequality to the end of the appendix.

When $s \in (0, 1)$ the asymptotics of K_0 and K_0' ensure that $K_0 \in L^p((0, \infty))$ for all $p \geq 1$ and that

$$\begin{aligned} K(s) &\leq \frac{C}{s^\mu} \frac{\|W\|_{L^q}}{\sqrt{s} |K_0'(\sqrt{s})|} \left(1 + \left(s^{\frac{1}{q'}} K_0(\sqrt{s})\right)^{-1}\right) \left(K_0(\sqrt{s}) + \frac{1}{\sqrt{s}}\right) \\ &\leq \frac{C\|W\|_{L^q}}{s^{\mu-\mu_{crit}+1} |\ln(s)|}, \end{aligned}$$

where $1/s^{\mu-\mu_{crit}+1} |\ln(s)| \in L^1((0, 1))$ since $\mu-\mu_{crit} < 0$. While, when $s \in (1, \infty)$, the same asymptotics guarantee that (remember that $q' > 2$ to bound $\alpha^{1-q'/2} \leq s^{1/2-q'/4}$ for $\sqrt{s} < \alpha$):

$$\begin{aligned} K(s) &\leq \frac{C\|W\|_{L^q}}{s^\mu} \frac{\left(1 + \exp(\sqrt{s}) s^{-\frac{1}{2q'}} \left(\int_{\sqrt{s}}^\infty \exp(-q'\alpha) d\alpha\right)^{\frac{1}{q'}}\right)}{s^{\frac{3}{4}} \exp(-\sqrt{s})} \dots \\ &\dots \left(\frac{\exp(-\sqrt{s})}{s^{\frac{1}{4}}} + \frac{1}{\sqrt{s}} \left(\int_{\sqrt{s}}^\infty \exp(-2\alpha) d\alpha\right)^{\frac{1}{2}}\right) \end{aligned}$$

and finally $K(s) \leq C\|W\|_{L^q(\mathbb{R}^2)} s^{-\mu-\frac{1}{2q'}-1} \in L^1(1, \infty)$. Hence, we have a uniform bound C independent of $\varepsilon \in (0, 1)$ such that:

$$\int_0^\infty \frac{1}{(\lambda + \varepsilon)^\mu} \|\phi_{\lambda+\varepsilon}\|_{L^2(\mathbb{R}^d)} (|\ell_W| + |F_{\lambda+\varepsilon}^{(0)}[W]|) d\lambda \leq \int_0^\infty K(s) ds \leq C\|W\|_{L^q}.$$

For the second integral we denote similarly:

$$\tilde{K}(s) = \frac{1}{s^\mu} \|\psi_s\|_{L^2(\mathbb{R}^2)} (|\omega_W| + |\tau_s^{(0)}[W]|)$$

With the explicit form of ψ_s we have:

$$\|\psi_s\|_{L^2(\mathbb{R}^2)} \leq \frac{C}{(1+s)K_1(\sqrt{s}) - \sqrt{s}K_1'(\sqrt{s})} \left(|K_1(\sqrt{s})| + \frac{1}{\sqrt{s}} \left(\int_{\sqrt{s}}^\infty |K_1(\alpha)|^2 \alpha d\alpha\right)^{\frac{1}{2}} \right).$$

and

$$|\tau_s^{(0)}[W]| \leq C \frac{\|w\|_{L^q(\mathcal{F}_0)}}{K_1(\sqrt{s}) s^{\frac{1}{q'}}} \left(\int_{\sqrt{s}}^\infty |K_1(\alpha)|^{q'} \alpha d\alpha\right)^{\frac{1}{q'}} \quad (\text{B.5})$$

When $s \in (1, \infty)$, K_0 and K_1 admit a similar exponential bound, so we obtain with similar arguments as previously that \tilde{K} is dominated by an L^1 -function multiplied by $\|W\|_{L^q(\mathbb{R}^2)}$. When $s \in (0, 1)$ we proceed more carefully but similarly again. We have $|K_1(\alpha)| \leq 1/\alpha$ when $\alpha < 1$. Hence, we compute that:

$$\int_{\sqrt{s}}^\infty |K_1(\alpha)|^2 \alpha d\alpha \leq C(1 + |\ln(s)|), \quad \int_{\sqrt{s}}^\infty |K_1(\alpha)|^{q'} \alpha d\alpha \leq \frac{1}{s^{\frac{q'}{2}-1}}.$$

Consequently:

$$|\tilde{K}(s)| \leq C(1 + |\ln(s)|)^{\frac{1}{2}} \|W\|_{L^q(\mathbb{R}^2)}.$$

We conclude like previously. \square

To end up this section, we provide a proof of identities (B.4)-(B.5). This is the content of the following proposition:

Proposition B.2. *Let $\lambda > 0$ and $q \in (1, 2)$. There exists a constant C depending only on the physical parameters and q such that, given $W \in L^2[B_0] \cap [L^q(\mathbb{R}^2)]^2$ we have:*

$$|F_\lambda^{(0)}[W]| \leq C \frac{\|w\|_{L^q(\mathcal{F}_0)}}{K_0(\sqrt{\lambda}) \lambda^{\frac{1}{q'}}} \left(\int_{\sqrt{\lambda}}^\infty |K_0(s)|^{q'} s ds\right)^{\frac{1}{q'}}$$

$$|\tau_\lambda^{(0)}[W]| \leq C \frac{\|w\|_{L^q(\mathcal{F}_0)}}{K_1(\sqrt{\lambda})\lambda^{\frac{1}{q'}}} \left(\int_{\sqrt{\lambda}}^{\infty} |K_1(s)|^{q'} s ds \right)^{\frac{1}{q'}}$$

where $F_\lambda^{(0)}[W]$ and $\tau_\lambda^{(0)}[W]$ are defined in (B.2) and q' is the conjugate exponent of q .

Proof. We provide a proof of the second inequality. The first one is obtained with a similar construction based on K_0 .

Let $\omega \in \mathbb{R}$ and

$$u(x) = \frac{K_1(\sqrt{\lambda}|x|)}{K_1(\sqrt{\lambda})} \frac{\omega x^\perp}{|x|} \quad \forall x \in \mathcal{F}_0.$$

By construction, we have:

$$-\Delta u + \lambda u = 0 \text{ on } \mathcal{F}_0, \quad u(x) = \omega x^\perp \text{ on } \partial B_0.$$

Introducing the latter identity into the definition of $\tau_\lambda^{(0)}[W]$, we derive:

$$\tau_\lambda^{(0)}[W]\omega = \int_{\partial B_0} \partial_n v_\lambda^{(0)}[W] u d\sigma,$$

so that we can integrate by parts. Recalling that $v_\lambda^{(0)}[W] = V_\lambda^{(0)}[W]\mathbb{1}_{\mathcal{F}_0}$ satisfies a specific PDE and vanishes on ∂B_0 , then using the PDE satisfied by u , we deduce successively that:

$$\begin{aligned} \tau_\lambda^{(0)}[W]\omega &= \int_{\mathcal{F}_0} \Delta v_\lambda^{(0)} \cdot u + \int_{\mathcal{F}_0} \nabla v_\lambda^{(0)} : \nabla u \\ &= - \int_{\mathcal{F}_0} w \cdot u + \int_{\mathcal{F}_0} \lambda v_\lambda^{(0)} \cdot u - \int_{\mathcal{F}_0} v_\lambda^{(0)} \cdot \Delta u \\ &= - \int_{\mathcal{F}_0} w \cdot u. \end{aligned}$$

Via a standard Hölder inequality and homogeneity arguments we thus infer that:

$$|\tau_\lambda^{(0)}[W]\omega| \leq \frac{\|w\|_{L^q(\mathcal{F}_0)}|\omega|}{K_1(\sqrt{\lambda})\lambda^{\frac{1}{q'}}} \left(\int_{\sqrt{\lambda}}^{\infty} |K_1(s)|^{q'} s ds \right)^{\frac{1}{q'}}.$$

Since ω is arbitrary, this concludes the proof. \square

Acknowledgements. The second author acknowledges support of the Institut Universitaire de France and project "SingFlows" ANR-grant number: ANR-18-CE40-0027. The second author would also like to thank Luc Hillairet for pointing reference [10] on special functions and for enlightening discussions about reference [9].

REFERENCES

- [1] H. Brezis. Remarks on the preceding paper by M. Ben-Artzi: "Global solutions of two-dimensional Navier-Stokes and Euler equations" [Arch. Rational Mech. Anal. **128** (1994), no. 4, 329–358; MR1308857 (96h:35148)]. *Arch. Rational Mech. Anal.*, 128(4):359–360, 1994.
- [2] M. Debayan, S. Ervedoza, and M. Tucsnak. Large time behaviour for the motion of a solid in a viscous incompressible fluid. April 2020.
- [3] M. Del Pino and J. Dolbeault. Best constants for Gagliardo-Nirenberg inequalities and applications to nonlinear diffusions. *J. Math. Pures Appl. (9)*, 81(9):847–875, 2002.
- [4] S. Ervedoza, M. Hillairet, and C. Lacave. Long-time behavior for the two-dimensional motion of a disk in a viscous fluid. *Comm. Math. Phys.*, 329(1):325–382, 2014.

- [5] H. Fujita and T. Kato. On the Navier-Stokes initial value problem. I. *Arch. Rational Mech. Anal.*, 16:269–315, 1964.
- [6] G. P. Galdi. *An introduction to the mathematical theory of the Navier-Stokes equations*. Springer Monographs in Mathematics. Springer, New York, second edition, 2011. Steady-state problems.
- [7] T. Gallay and Y. Maekawa. Long-time asymptotics for two-dimensional exterior flows with small circulation at infinity. *Anal. PDE*, 6(4):973–991, 2013.
- [8] T. Gallay and C. E. Wayne. Global stability of vortex solutions of the two-dimensional Navier-Stokes equation. *Comm. Math. Phys.*, 255(1):97–129, 2005.
- [9] H. Kozono and T. Ogawa. Decay properties of strong solutions for the Navier-Stokes equations in two-dimensional unbounded domains. *Arch. Rational Mech. Anal.*, 122(1):1–17, 1993.
- [10] F. W. J. Olver. *Asymptotics and special functions*. Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1974. Computer Science and Applied Mathematics.
- [11] A. Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1983.
- [12] H. Sohr. *The Navier-Stokes equations*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Verlag, Basel, 2001. An elementary functional analytic approach.
- [13] T. Takahashi and M. Tucsnak. Global strong solutions for the two-dimensional motion of an infinite cylinder in a viscous fluid. *J. Math. Fluid Mech.*, 6(1):53–77, 2004.
- [14] Y. Wang and Z. Xin. Analyticity of the semigroup associated with the fluid-rigid body problem and local existence of strong solutions. *J. Funct. Anal.*, 261(9):2587–2616, 2011.