



**HAL**  
open science

## **Polymeric self-assemblies for photodynamic therapy: A critical approach**

Ugo Till, Laure Gibot, Anne-Françoise Mingotaud, Christophe Mingotaud, Marie-Pierre Rols, Mireille Gaucher, Frédéric Violleau, Patricia Vicendo

### ► **To cite this version:**

Ugo Till, Laure Gibot, Anne-Françoise Mingotaud, Christophe Mingotaud, Marie-Pierre Rols, et al.. Polymeric self-assemblies for photodynamic therapy: A critical approach. Photodiagnosis and Photodynamic Therapy, 2017, 17, pp.A23-A24. 10.1016/j.pdpdt.2017.01.053 . hal-03511855

**HAL Id: hal-03511855**

**<https://hal.science/hal-03511855>**

Submitted on 5 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




## Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible

This is an author's version published in: <http://oatao.univ-toulouse.fr/25141>

**Official URL:** <https://doi.org/10.1016/j.pdpdt.2017.01.053>

**To cite this version:**

Till, Ugo and Gibot, Laure and Mingotaud, Anne-Françoise and Mingotaud, Christophe and Rols, Marie-Pierre and Gaucher, Mireille and Violleau, Frédéric  and Vicendo, Patricia *Polymeric self-assemblies for photodynamic therapy: A critical approach*. (2017) *Photodiagnosis and Photodynamic Therapy*, 17. A23-A24. ISSN 1572-1000

Any correspondence concerning this service should be sent to the repository administrator: [tech-oatao@listes-diff.inp-toulouse.fr](mailto:tech-oatao@listes-diff.inp-toulouse.fr)

**Oral OC-047**

**Polymeric self-assemblies for photodynamic therapy: A critical approach**

U. Till<sup>1,2</sup>, L. Gibot<sup>3</sup>, A.F. Mingotaud<sup>1,\*</sup>,  
C. Mingotaud<sup>1</sup>, M.P. Rols<sup>3</sup>, M. Gaucher<sup>2</sup>,  
F. Violleau<sup>2</sup>, P. Vicendo<sup>1</sup>

<sup>1</sup> *Université de Toulouse; UPS/CNRS; IMRCP, Toulouse, France*

<sup>2</sup> *Université de Toulouse, Institut National Polytechnique de Toulouse – Ecole d'Ingénieurs de Purpan, Département Sciences Agronomiques et Agroalimentaires, UPSP/DGER 115, Toulouse, France*

<sup>3</sup> *Equipe de Biophysique Cellulaire, IPBS-CNRS UMR 5089, Toulouse, France*

The work presented here suggests a new approach in the critical development of polymeric nanovectors for photodynamic therapy (PDT) against cancer. Whereas hundreds of studies quickly jump forward from formation of self-assemblies to biological application without having a thorough examination of the vector solution, we suggest having a parallel assessment of formation/characterization of the nanovectors and biological activity [1]. This is possible by first using a careful physical chemistry characterization of the vectors by both batch techniques (light and neutron scattering, electron microscopy, atomic force microscopy) and Asymmetrical Flow Field-Flow Fractionation (AsFFFF) coupled to adequate detectors (refractometry, light scattering) [2]. This enables us to fully characterize the vectors regarding purity, size and zeta potential.

Data on both polymeric micelles and polymersomes are presented here, using poly(ethyleneoxide-*b*- $\epsilon$ -caprolactone), poly(ethyleneoxide-*b*-D,L-lactide) and poly(ethyleneoxide-*b*-styrene). Self-assemblies exhibiting size range of 20–200 nm are presented and reveal the possible presence of different populations of nanovectors in some cases. Controlled mixtures of different nano-objects are also studied, as well as crosslinked systems. For each new vector, its ability to carry a photosensitizer (Pheophorbide a) for PDT is examined. The activity in PDT either in 2D and 3D cell culture is presented and compared on different batches, in link with the purity analysis. The work shows that selected mixtures of different vectors with different morphologies or sizes may lead to synergetic effects. Also, a strong influence of the crosslinking of the vector has been observed and will be presented.

