### SUPPLEMENTARY MATERIAL

# Association between residential proximity to agricultural crops and adaptive behaviors in children with autism spectrum disorder from the French ELENA cohort

Jeanne Sandrine Ongono\*, Cécile Michelon, Remi Béranger, Emmanuelle Cadot, Valentin Simoncic, Julie Loubersac, Marion Mortamais<sup>#</sup>, Amaria Baghdadli<sup>#</sup>

\* Corresponding author

# Joint last authors

### TABLE OF CONTENTS

/ Interaction analyses	. 2
I/ Supplementary descriptive analyses	. 3
II/ Sensitivity analyses	. 5
1. Additional adjustments and analyses by subgroup of crop exposure scores	. 5
2. Associations between cumulative (from pregnancy to the first two years of life) total exposu	re
score to agricultural crops within a 1000m radius around residential addresses and VABS-II	
subdomains scores.	15
3. Associations between total exposure scores to agricultural crops within a 500m radius arour	١d
residential addresses and VABS-II subdomains scores.	16

### I/ Interaction analyses

**Table S1.** *P-value* of interaction between total exposure score to agricultural crops within a 1000m and 500m radius around residential addresses and the intellectual quotient for developmental periods and each VABS-II subdomain score in children with autism spectrum disorder from the French ELENA cohort.

			Buffer 1000m			
		Pregnancy period			First two years of life period	
	Communication	Daily living skills	Socialization	Communication	Daily living skills	Socialization
Pinteraction	0.0006	<.0001	0.0039	0.0008	<.0001	0.0044
			Buffer 500m			
		Pregnancy period			First two years of life period	
	Communication	Daily living skills	Socialization	Communication	Daily living skills	Socialization
Pinteraction	0.0004	<.0001	0.0075	<.0001	<.0001	0.0042

**Abbreviation:** VABS-II=Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition

### II/ Supplementary descriptive analyses

**Table S2.** Comparison of main characteristics between the study population (n=183) and the "strictly urban"<sup>a</sup> subpopulation excluded for the pregnancy period (n=41) of exposure to agricultural crops in children with ASD from the ELENA cohort.

Characteristics	study population (n=183)	"Strictly urban" subpopulation	P value*
		excluded from analyses (n=41)	
Children age at inclusion	6.1 (3.5)	6.3 (3.4)	<.0001
Children's sex			0.950
Males	142 (78%)	32 (78%)	0.550
Females	41 (22%)	9 (22%)	
Children intellectual quotient (IQ)			
IQ	76.9 (28.9)	75.5 (30.2)	<.0001
Intellectual disability (ID) Status			0.979
ID	71 (39%)	16 (39%)	
Non ID	112 (61%)	25 (61%)	
VABS-II subdomains scores			
Communication	73.0 (15.8)	70.9 (14.0)	<.0001
Daily living skills	74.8 (13.6)	73.2 (11.8)	<.0001
Socialization	71.8 (11.3)	69.6 (11.0)	<.0001
Total exposure score to agricultural crops during pregnancy	29% (27%)	0	-
Exposure score to "crops group 1"* during pregnancy	12% (21%)	0	-
Exposure score to "crops group 2"** during pregnancy	17% (21%)	0	-

a: The "strictly urban" status was defined by more than 99% of urbanized area within a 500-m radius around each address.

Abbreviations: ASD=autism spectrum disorder; ID=intellectual disability; VABS-II= Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition

According to the Corine Land Cover classification (CLC; French Ministry of Environment, Statistics and Observation Department, scale 1:100 000), we defined crop groups as:

\* **Crops group 1**=vineyards, fruit trees, and berry plantations

**\*\* Crops group 2**= non-irrigated arable land (cereals, fodder crops, root crops and fallow land, flower and tree cultivation, and vegetables); annual crops associated with permanent crops (woody crops like shrubs, olives in combination with either non-permanent crops like fodder roots, fodder maize, forage kale, etc., or permanent grass surfaces) and land principally occupied by agriculture with significant areas of natural vegetation.

\*\*\* Wilcoxon or Student tests for quantitative variables, and Chi2 test for qualitative variables.

\* Wilcoxon or Student tests for quantitative variables and Chi2 test for qualitative variables.

#### **III/ Sensitivity analyses**

#### 1. Additional adjustments and analyses by subgroup of crop exposure scores

**Table S3**. Associations [β (CI95%)], between each 20% increase in total exposure score to agricultural crops within a 1000m radius around maternal residential address during pregnancy and VABS-II subdomains scores<sup>a</sup> in children with autism spectrum disorder from the ELENA cohort (n=158).

Total exposure score to crops		VABS-II subdomains scores		
		ASD without ID		
Model*	Communication β (Cl95%) -2.90 (-5.06 to -0.74)	Daily living skills β (Cl95%) -1.88 (-3.74 to -0.01)	<b>Socialization</b> <b>β (CI95%)</b> -1.34 (-3.13 to 0.44)	
		ASD with ID		
Model*	Communication β (Cl95%) 1.18 (-1.69 to 4.06)	Daily living skills β (Cl95%) 0.86 (-1.70 to 3.43)	<b>Socialization</b> <b>β (CI95%)</b> 0.04 (-1.84 to 1.92)	

\*Sensitivity analysis, Model\*= final model [adjusted on Child's sex (female, male), age at the inclusion in the ELENA cohort (<median, >median), higher level of parental education in the household (< high school, university)] additionally adjusted on + maternal folic acid intake (yes/no), alcohol consumption (yes/no), smoking during pregnancy (yes/no), child's birth season (Spring, Summer, Autumn, Winter), recruiting center (*Occitanie, Auvergne-Rhône-Alpes, Ile-de-France, Provence-Alpes-Côte d'Azur, Nouvelle Aquitaine, Grand Est*)

**a**: negative Beta indicating alteration in adaptive functioning

**VABS-II**: Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition

**Table S4**. Associations [ $\beta$  (CI95%)] between each 20% increase in total exposure score to agricultural crops within a 1000m radius around residential address during the first two years of life and VABS-II subdomains scores<sup>a</sup> in children with autism spectrum disorder from the ELENA cohort (n=168).

Total exposure score to crops		VABS-II subdomains scores	
		ASD without ID	
	Communication β (Cl95%)	Daily living skills β (Cl95%)	Socialization β (CI95%)
Model*	-2.36 (-4.33 to -0.38)	-1.49 (-3.17 to 0.18)	-1.30 (-2.88 to 0.29)
		ASD with ID	
	Communication β (Cl95%)	Daily living skills β (Cl95%)	Socialization β (Cl95%)
Model*	2.25 (-0.63 to 5.13)	2.03 (-0.32 to 4.39)	0.21 (-1.57 to 1.98)

\*Sensitivity analysis, Model\*= final model [adjusted on Child's sex (female, male), age at the inclusion in the ELENA cohort (<median, >median), higher level of parental education in the household (< high school, university)] additionally adjusted on + maternal folic acid intake (yes/no), alcohol consumption (yes/no), smoking during pregnancy (yes/no), child's birth season (Spring, Summer, Autumn, Winter), recruiting center (*Occitanie, Auvergne-Rhône-Alpes, Ile-de-France, Provence-Alpes-Côte d'Azur, Nouvelle Aquitaine, Grand Est*)

a: negative Beta indicating alteration in adaptive functioning

VABS-II: Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition

**Table S5**. Associations [ $\beta$  (CI95%)] between each 20% increase in exposure score to crops group 1\* within a 1000m radius around maternal residential address during pregnancy and VABS-II subdomains scores<sup>a</sup> in children with autism spectrum disorder from the ELENA cohort (n=183).

Exposure score to crops group 1*		VABS-II subdomains scores	
		ASD without ID	
	Communication β (CI95%)	Daily living skills β (Cl95%)	Socialization β (Cl95%)
Crude model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	-0.60 (-5.99 to 4.78)	-0.22 (-4.77 to 4.33)	-1.14 (-5.12 to 2.84)
Adjusted model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	-0.46 (-5.70 to 4.79)	-0.98 (-5.70 to 3.75)	-0.97 (-5.05 to 3.12)
		ASD with ID	
	Communication β (CI95%)	Daily living skills β (Cl95%)	Socialization β (Cl95%)
Crude model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	4.53 (-1.47 to 10.52)	1.16 (-4.83 to 7.16)	2.15 (-2.99 to 7.29)
Adjusted model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	5.81 (-0.23 to 11.85)	2.03 (-4.07 to 8.12)	3.76 (-1.10 to 8.61)

**Model 2:** Adjusted model on Child's sex (female, male), age at the inclusion in the ELENA cohort ( $\leq$ median, >median), higher level of parental education in the household ( $\leq$  high school, university)

According to the Corine Land Cover classification (CLC; French Ministry of Environment, Statistics and Observation Department, scale 1:100 000), Crops group 1\*=vineyards, fruit trees, and berry plantations

a: negative Beta indicating alteration in adaptive functioning

**VABS-II**: Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition

**Table S6**. Associations [ $\beta$  (CI95%)] between each 20% increase in exposure score to crops group 2\* within a 1000m radius around maternal residential address during pregnancy and VABS-II subdomains scores<sup>a</sup> in children with autism spectrum disorder from the ELENA cohort (n=183).

Exposure score to crops group 2*		VABS-II subdomains scores	
		ASD without ID	
	Communication β (Cl95%)	Daily living skills β (Cl95%)	Socialization β (Cl95%)
Crude model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	-3.08 (-8.29 to 2.13)	-0.21 (-4.64 to 4.22)	0.46 (-3.42 to 4.35)
Adjusted model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	-2.43 (-7.51 to 2.64)	-0.71 (-5.30 to 3.87)	1.27 (-2.68 to 5.23)
		ASD with ID	
	Communication β (Cl95%)	Daily living skills β (Cl95%)	Socialization β (Cl95%)
Crude model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	0.12 (-6.04 to 6.29)	4.51 (-1.47 to 10.49)	0.33 (-4.90 to 5.56)
Adjusted model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	-1.62 (-7.93 to 4.68)	3.98 (-2.16 to 10.13)	-1.26 (-6.29 to 3.76)

**Model 2:** Adjusted model on Child's sex (female, male), age at the inclusion in the ELENA cohort (≤median, >median), higher level of parental education in the household (≤ high school, university)

According to the Corine Land Cover classification (CLC; French Ministry of Environment, Statistics and Observation Department, scale 1:100 000), **crops group 2**\*\*= non-irrigated arable land (cereals, fodder crops, root crops and fallow land, flower and tree cultivation, and vegetables); annual crops associated with permanent crops (woody crops like shrubs, olives in combination with either non-permanent crops like fodder roots, fodder maize, forage kale, etc., or permanent grass surfaces) and land principally occupied by agriculture with significant areas of natural vegetation.

a: negative Beta indicating alteration in adaptive functioning

**VABS-II**: Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition

**Table S7**. Associations [ $\beta$  (CI95%)] between each 20% increase in exposure score to crops group 1\* within a 1000m radius around residential address during the first two years of life and VABS-II subdomains scores<sup>a</sup> in children with autism spectrum disorder from the ELENA cohort (n=193).

Exposure score to crops group 1*		VABS-II subdomains scores	
		ASD without ID	
	Communication β (Cl95%)	Daily living skills β (Cl95%)	Socialization β (Cl95%)
Crude model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	-1.63 (-6.86 to 3.61)	-0.27 (-4.70 to 4.17)	-0.52 (-4.38 to 3.34)
Adjusted model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	-1.68 (-6.78 to 3.42)	-0.68 (-5.22 to 3.86)	-0.39 (-4.29 to 3.50)
		ASD with ID	
	Communication β (CI95%)	Daily living skills β (Cl95%)	Socialization β (CI95%)
Crude model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	6.10 (0.10 to 12.10)	4.13 (-1.57 to 9.83)	1.69 (-3.13 to 6.50)
Adjusted model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	6.86 (0.95 to 12.77)	4.93 (-0.63 to 10.49)	2.57 (-1.91 to 7.04)

**Model 2:** Adjusted model on Child's sex (female, male), age at the inclusion in the ELENA cohort ( $\leq$ median, >median), higher level of parental education in the household ( $\leq$  high school, university)

According to the Corine Land Cover classification (CLC; French Ministry of Environment, Statistics and Observation Department, scale 1:100 000), Crops group 1\*=vineyards, fruit trees, and berry plantations

a: negative Beta indicating alteration in adaptive functioning

**VABS-II**: Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition

**Table S8**. Associations [ $\beta$  (CI95%)] between each 20% increase in exposure score to crops group 2\* within a 1000m radius around residential address during the first two years of life and VABS-II subdomains scores<sup>a</sup> in children with autism spectrum disorder from the ELENA cohort (n=193).

Exposure score to crops group 2*		VABS-II subdomains scores	
		ASD without ID	
	Communication β (Cl95%)	Daily living skills β (Cl95%)	Socialization β (Cl95%)
Crude model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	-2.93 (-8.01 to 2.15)	-1.09 (-5.41 to 3.23)	-0.07 (-3.84 to 3.69)
Adjusted model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	-2.04 (-7.02 to 2.94)	-1.31 (-5.75 to 3.13)	0.68 (-3.12 to 4.49)
		ASD with ID	
	Communication β (Cl95%)	Daily living skills β (Cl95%)	Socialization β (Cl95%)
Crude model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	0.48 (-5.68 to 6.64)	1.51 (-4.25 to 7.28)	-1.01 (-5.84 to 3.82)
Adjusted model			
Exposure score ≤median	Ref	Ref	Ref
Exposure score >median	-0.60 (-6.77 to 5.56)	0.70 (-5.01 to 6.42)	-2.06 (-6.58 to 2.46)

**Model 2:** Adjusted model on Child's sex (female, male), age at the inclusion in the ELENA cohort ( $\leq$ median, >median), higher level of parental education in the household ( $\leq$  high school, university)

According to the Corine Land Cover classification (CLC; French Ministry of Environment, Statistics and Observation Department, scale 1:100 000), **crops group 2\***= non-irrigated arable land (cereals, fodder crops, root crops and fallow land, flower and tree cultivation, and vegetables); annual crops associated with permanent crops (woody crops like shrubs, olives in combination with either non-permanent crops like fodder roots, fodder maize, forage kale, etc., or permanent grass surfaces) and land principally occupied by agriculture with significant areas of natural vegetation.

a: negative Beta indicating alteration in adaptive functioning

VABS-II: Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition

2. Associations between cumulative (from pregnancy to the first two years of life) total exposure score to agricultural crops within a 1000m radius around residential addresses and VABS-II subdomains scores.

**Table S9**. Associations [ $\beta$  (CI95%)], between each 20% increase in the cumulative total exposure score (from pregnancy to the first two years of children's life) to agricultural crops within a 1000m radius around residential address and VABS-II subdomains scores<sup>a</sup> in children with autism spectrum disorder from the ELENA cohort (n=177, 108 without ID and 69 with ID).

Cumulative total exposure score to		VABS-II subdomains scores	
crops			
	ASD without ID		
	Communication	Daily living skills	Socialization
	β (Cl95%)	β (Cl95%)	β (Cl95%)
Crude model	-2.32 (-4.28 to -0.37)	-1.10 (-2.78 to 0.59)	-1.74 (-3.18 to -0.29)
Adjusted model	-2.33 (-4.25 to -0.40)	-1.46 (-3.21 to 0.29)	-1.60 (-3.10 to-0.10)
		ASD with ID	
	Communication β (Cl95%)	Daily living skills β (Cl95%)	Socialization β (Cl95%)
Crude model	0.53 (-1.84 to 2.89)	0.78 (-1.52 to 3.09)	0.58 (-1.43 to 2.58)
Adjusted model	0.42 (-1.95 to 2.79)	0.62 (-1.68 to 2.93)	0.39 (-1.50 to 2.28)

Adjusted model on Child's sex (female, male), age at the inclusion in the ELENA cohort (<median, >median), higher level of parental education in the

household (≤ high school, university)

*a*: negative Beta indicating alteration in adaptive functioning

**VABS-II**: Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition

#### 3. Associations between total exposure scores to agricultural crops within a 500m radius around residential addresses and VABS-II subdomains scores.

#### 3.1. Pregnancy period

**Table S10**. Associations [β (CI95%)] between each 20% increase in total exposure score to agricultural crops within a 500m radius around maternal residential address during pregnancy and VABS-II subdomains scores<sup>a</sup> in children with autism spectrum disorder from the ELENA cohort (n=183).

Total exposure score to crops		VABS-II subdomains scores	
	ASD without ID		
Crude model	Communication β (Cl95%) -2.73 (-4.89 to -0.58)	Daily living skills β (CI95%) -0.81 (-2.69 to 1.07)	Socialization β (CI95%) -1.65 (-3.28 to -0.02)
Adjusted model	-2.38 (-4.47 to -0.29)	-0.97 (-2.90 to 0.97)	-1.41 (-3.07 to 0.26)
		ASD with ID	
Crude model	<b>Communication</b> <b>β (Cl95%)</b> 0.27 (-2.04 to 2.59)	Daily living skills β (Cl95%) 0.83 (-1.45 to 3.10)	<b>Socialization</b> <b>β (CI95%)</b> 0.96 (-0.99 to 2.91)
Adjusted model	0.06 (-2.25 to 2.36)	0.65 (-1.61 to 2.91)	0.67 (-1.16 to 2.49)

Adjusted model on Child's sex (female, male), age at the inclusion in the ELENA cohort (<median, >median), higher level of parental education in the

household ( $\leq$  high school, university)

**a**: negative Beta indicating alteration in adaptive functioning

**VABS-II**: Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition

#### 3.2. First two years of life period

**Table S11**. Associations [ $\beta$  (CI95%)] between each 20% increase in total exposure score to agricultural crops within a 500m radius around residential address during the first two years of life and VABS-II subdomains scores<sup>a</sup> in children with autism spectrum disorder from the ELENA cohort (n=193).

Total exposure score to crops		VABS-II subdomains scores	
		ASD without ID	
Crude model	Communication β (Cl95%) -1.40 (-3.29 to 0.48)	Daily living skills β (Cl95%) -0.47 (-2.07 to 1.14)	Socialization β (Cl95%) -1.55 (-2.92 to -0.18)
Adjusted model	-1.41 (-3.23 to 0.42)	-0.59 (-2.23 to 1.05)	-1.39 (-2.78 to -0.01)
		ASD with ID	
	Communication β (Cl95%)	Daily living skills β (Cl95%)	Socialization β (Cl95%)
Crude model	0.29 (-2.14 to 2.72)	0.98 (-1.29 to 3.25)	0.80 (-1.10 to 2.70)
Adjusted model	-0.04 (-2.47 to 2.39)	0.79 (-1.46 to 3.03)	0.57 (-1.22 to 2.36)

Adjusted model on Child's sex (female, male), age at the inclusion in the ELENA cohort (≤median, >median), higher level of parental education in the

household (≤ high school, university)

**a**: negative Beta indicating alteration in adaptive functioning

**VABS-II**: Vineland Adaptive Behavior Scales 2<sup>nd</sup> edition