



**HAL**  
open science

## On periodic solutions of fractional-order differential systems with a fixed length of sliding memory

Safa Bourafa, René Lozi, Mohammed-Salah Abdelouahab

► **To cite this version:**

Safa Bourafa, René Lozi, Mohammed-Salah Abdelouahab. On periodic solutions of fractional-order differential systems with a fixed length of sliding memory. *Journal of Innovative Applied Mathematics and Computational Sciences*, 2021, 1 (1), pp.64-78. hal-03513190

**HAL Id: hal-03513190**

**<https://hal.science/hal-03513190>**

Submitted on 5 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# On periodic solutions of fractional-order differential systems with a fixed length of sliding memory

Safa Bourafa  <sup>1</sup>, Mohammed Salah Abdelouahab <sup>1</sup> and René Lozi <sup>2</sup>

<sup>1</sup>Laboratory of Mathematics and their interactions, University Center of Mila, Algeria

<sup>2</sup> Université Côte d'Azur, CNRS, LJAD, Nice, France

Received 16 November 2021, Accepted 25 December 2021

---

**Abstract.** The fractional-order derivative of a non-constant periodic function is not periodic with the same period. Consequently, any time-invariant fractional-order systems do not have a non-constant periodic solution. This property limits the applicability of fractional derivatives and makes it unfavorable to model periodic real phenomena. This article introduces a modification to the Caputo and Riemann-Liouville fractional-order operators by fixing their memory length and varying the lower terminal. It is shown that this modified definition of fractional derivative preserves the periodicity. Therefore, periodic solutions can be expected in fractional-order systems in terms of the new fractional derivative operator. To confirm this assertion, one investigates two examples, one linear system for which one gives an exact periodic solution by its analytical expression and another nonlinear system for which one provides exact periodic solutions using qualitative and numerical methods.


**Keywords:** Fractional-order derivative; sliding fixed memory length; periodic solution.

2020 Mathematics Subject Classification: 34A35, 34K28, 34K40, 45L05.

---

## 1 Introduction

The history of fractional calculus goes back to the end of the 17th century when L'Hopital asked Leibniz what meaning could be ascribed to  $D^n f$  if  $n$  were a fraction? Since that, time-fractional calculus has drawn the attention of many famous mathematicians, such as Euler, Laplace, Fourier, Abel, Liouville, Riemann, and Laurent [18]. The advantages of fractional calculus have been described and pointed out in the last few decades by many authors [8, 15–19]. It has been shown that the fractional-order models of realistic systems are regularly more adequate than usually used integer-order models. Applications of these fractional-order models spread in many fields, such as viscoelastic systems, dielectric polarization, electrode-electrolyte polarization, electromagnetic waves, quantum evolution of complex systems, and so on [6, 10, 11, 14, 20]. There are three definitions most frequently used

 Corresponding author. Email: s.bourafa@centre-univ-mila.dz

for the general fractional differential operators. The first one is the Grünwald-Letnikov (GL) fractional differential operator defined by the limit of a fractional-order backward difference and has an advantage for numerical simulations. The second type is the Riemann-Liouville (RL) definition; this operator played a pivotal role in developing the fractional calculus theory. Using these two fractional differential operators in modeling real phenomena leads to mathematical models with initial conditions expressed in terms of fractional derivatives that do not have known physical interpretation. The third type is the Caputo derivative having the advantage of dealing models with initial conditions expressed in terms of the field variables and their integer-order derivatives, having clear physical interpretations [9]. Recently it has been demonstrated that the fractional-order derivative of a non-constant periodic function is not a periodic function with the same period [13,22,23] and in [5] the authors studied quasi-periodic properties of fractional order integrals and derivatives of periodic functions. As a consequence of the non-periodicity of the fractional derivative of a  $T$ -periodic function, the time-invariant fractional-order systems do not have any non-constant exact periodic solution unless the lower terminal of the derivative is  $\pm\infty$  [12,13,23], which is not realistic. This property limits the applicability of the fractional derivative and makes it unfavorable for periodic real phenomena. In [1], the authors have proposed a modification of the Grünwald-Letnikov fractional differential operator, which consists of fixing the memory length and varying the lower terminal of the derivative. They have demonstrated that the modified definition of fractional derivative preserves the periodicity. The present paper extends this modification to the Caputo and Riemann-Liouville fractional-order operators. Two examples are investigated to confirm that periodic solutions arise in fractional-order systems when the new fractional derivative operator is used. One linear system for which one gives an exact periodic solution defined by its analytical expression and another nonlinear system for which one provides an exact periodic solution using both qualitative and numerical methods.

## 2 Fractional-Order Derivatives

As said above, the most usual definitions of fractional-order derivative are the Grünwald-Letnikov, the Riemann-Liouville and the Caputo definitions [17]. For  $0 < \alpha \notin \mathbb{N}$ , the  $\alpha$ -th order derivative of a function  $f(t)$  with respect to  $t$  and a terminal value  $a$  is given in the sense of

- Grünwald-Letnikov by

$${}^GL_a D_t^\alpha f(x) = \lim_{\substack{h \rightarrow 0 \\ nh = x - a}} h^{-\alpha} \sum_{k=0}^n (-1)^k \binom{\alpha}{k} f(x - kh), \quad (2.1)$$

where  $\binom{\alpha}{k} = \frac{\Gamma(\alpha+1)}{k! \Gamma(\alpha-k+1)}$ .

- Riemann-Liouville by

$${}^{RL}_a D_t^\alpha f(t) = \frac{1}{\Gamma(m - \alpha)} \frac{d^m}{dt^m} \left( \int_a^t (t - \tau)^{m-\alpha-1} f(\tau) d\tau \right). \quad (2.2)$$

- Caputo by

$${}^C D_t^\alpha f(t) = \frac{1}{\Gamma(m-\alpha)} \int_a^t (t-\tau)^{m-\alpha-1} f^{(m)}(\tau) d\tau. \quad (2.3)$$

In (2.2) and (2.3),  $m$  is the first integer greater than  $\alpha$ , and  $\Gamma(\cdot)$  is the Gamma function. The following theorems reveal a remarkable property for the fractional derivatives based on Caputo definition, Grünwald-Letnikov definition, Riemann-Liouville definition [22].

**Theorem 2.1.** *Suppose that  $f(t)$  is a non constant periodic function with period  $T$ . If  $f(t)$  is  $m$ -times differentiable, then the functions  ${}^C D_t^\alpha f(t)$ , where  $0 < \alpha \notin \mathbb{N}$  and  $m$  is the first integer greater than  $\alpha$ , cannot be a periodic functions with period  $T$ .*

**Theorem 2.2.** *Suppose that  $f(t)$  is  $(m-1)$ -times continuously differentiable and  $f^{(m)}(t)$  is bounded. If  $f(t)$  is a non-constant periodic function with period  $T$ , then the functions  ${}^{GL} D_t^\alpha f(t)$  and  ${}^{RL} D_t^\alpha f(t)$ , where  $0 < \alpha \notin \mathbb{N}$  and  $m$  is the first integer greater than  $\alpha$ , cannot be periodic functions with period  $T$ .*

**Example 2.3.** Let  $f(t) = \sin(t)$ . One has

$$\sin(t) = \sum_{p=0}^{\infty} (-1)^p \frac{t^{2p+1}}{(2p+1)!}.$$

Hence

$${}^{RL} D_t^\alpha \sin(t) = t^{1-\alpha} E_{2,2-\alpha}(-t^2),$$

where  $0 < \alpha < 1$  and  $E_{\alpha,\beta}(t)$  is the generalized Mittag-Leffler function defined by

$$E_{\alpha,\beta}(t) = \sum_{k=0}^{\infty} \frac{t^k}{\Gamma(\alpha k + \beta)}.$$

Numerical simulations showed that  $t^{1-\alpha} E_{2,2-\alpha}(-t^2)$  is not a periodic function where  $0 < \alpha < 1$ , even if  $\alpha = 1$  this function is the periodic function  $\cos(t)$ .

As a consequence of the above theorems, periodic solution cannot be expected in fractional-order systems, under any circumstances [22, 23].

**Corollary 2.4.** *A differential equation of fractional-order in the form*

$${}_a D_t^\alpha x(t) = f(x(t)),$$

*where  $0 < \alpha \notin \mathbb{N}$ , cannot have any non-constant smooth periodic solution.*

This property highlights one of the basic differences between fractional-order derivative and integer-order one, and it makes fractional-order systems unfavourable for a wide range of real periodic phenomena. Therefore in this paper one overcomes this problem by imposing a simple modification to both Riemann-Liouville and Caputo definitions.

### 3 The Fractional-Order Derivative with Sliding Fixed Memory Length

one first recalls the Grünwald-Letnikov fractional-order derivative with fixed memory length introduced in [1].

**Definition 3.1.** (The Grünwald-Letnikov fractional derivative with fixed memory length) Let  $\alpha \geq 0$ ,  $L > 0$ ,  $m$  an integer such that  $m - 1 \leq \alpha < m$  and  $f$  an integrable function in the interval  $[a - L, b]$ . The operator  ${}^{MG}_L D_t^\alpha$  defined by :

$${}^{MG}_L D_t^\alpha f(t) = \lim_{h \rightarrow 0} \frac{1}{h^\alpha} \sum_{k=0}^{\frac{L}{h}} (-1)^k \frac{\Gamma(\alpha + 1)}{k! \Gamma(\alpha - k + 1)} f(t - kh), \quad t \in [a, b], \quad (3.1)$$

is called the Grünwald-Letnikov fractional derivative with sliding fixed memory length.

The following proposition gives an evaluation of the limit in the definition of Grünwald-Letnikov fractional derivative with sliding fixed memory length.

**Proposition 3.2.** Under the assumptions of definition (3.1), if the function  $f$  is  $m$ -differentiable with  $f^{(m)} \in L^1[a - L, b]$ , then

$${}^{MG}_L D_t^\alpha f(t) = \sum_{k=0}^{m-1} \frac{f^{(k)}(t - L) L^{k-\alpha}}{\Gamma(k - \alpha + 1)} + \frac{1}{\Gamma(m - \alpha)} \int_{t-L}^t (t - \tau)^{m-\alpha-1} f^{(m)}(\tau) d\tau. \quad (3.2)$$

It has been demonstrated that this modified fractional-order derivative possesses two important properties: the preservation of periodicity and the short memory, which considerably reduces the cost of numerical computations. Furthermore, it has been proven that contrarily to fractional autonomous systems defined using classical fractional derivative, the fractional autonomous systems in terms of the modified fractional derivative can generate exact periodic solutions.

In order to generalize this work, one introduces in this section a similar modification to both Caputo fractional-order derivative and Riemann-Liouville fractional-order derivative as follows.

**Definition 3.3.** (The Caputo fractional derivative with sliding fixed memory length) Let  $\alpha > 0$ ,  $L > 0$ ,  $m$  an integer such that  $m = [\alpha] + 1$  and  $f \in C^m[a - L, b]$ . The Caputo fractional derivative with sliding fixed memory length is defined by

$${}^{MC}_L D_t^\alpha f(t) = \frac{1}{\Gamma(m - \alpha)} \int_{t-L}^t (t - \tau)^{m-\alpha-1} f^{(m)}(\tau) d\tau. \quad (3.3)$$

**Definition 3.4.** (The Riemann-Liouville fractional derivative with sliding fixed memory length) Let  $\alpha \geq 0$ ,  $L > 0$ ,  $m$  an integer such that  $m - 1 \leq \alpha < m$  and  $f$  is a continuous function in  $[a - L, b]$ , the Riemann-Liouville fractional derivative with sliding fixed memory length is defined by

$${}^{MRL}_L D_t^\alpha f(t) = \frac{1}{\Gamma(m - \alpha)} \frac{d^m}{dt^m} \int_{t-L}^t (t - \tau)^{m-\alpha-1} f(\tau) d\tau, \quad (3.4)$$

**Remark 3.5.** From (3.2) and (3.3) one gets

$${}^{MC}_L D_t^\alpha f(t) = {}^{MG}_L D_t^\alpha f(t) - \sum_{k=0}^{m-1} \frac{f^{(k)}(t - L) L^{k-\alpha}}{\Gamma(k - \alpha + 1)}. \quad (3.5)$$

**Proposition 3.6.** Under the assumption that the function  $f(t)$  is  $m$ -times continuously differentiable

$${}^{MRL}D_t^\alpha f(t) = {}^{MG}D_t^\alpha f(t) - \sum_{k=0}^{m-1} \frac{f^{(k)}(t-L)L^{k-\alpha}}{\Gamma(k-\alpha+1)}. \quad (3.6)$$

*Proof.* By differentiation and performing repeatedly integration by parts, one has

$$\begin{aligned} {}^{MRL}D_t^\alpha f(t) &= \frac{1}{\Gamma(m-\alpha)} \frac{d^m}{dt^m} \int_{t-L}^t (t-\tau)^{m-\alpha-1} f(\tau) d\tau, \\ &= -\frac{f^{(m-1)}L^{m-\alpha-1}(t-L)}{\Gamma(m-\alpha)} + \frac{1}{\Gamma(m-\alpha-1)} \frac{d^{m-1}}{dt^{m-1}} \int_{t-L}^t (t-\tau)^{m-\alpha-2} f(\tau) d\tau, \\ &\vdots \\ &= -\sum_{k=0}^{m-1} \frac{f^{(k)}(t-L)L^{k-\alpha}}{\Gamma(k-\alpha+1)} + \frac{1}{\Gamma(-\alpha)} \int_{t-L}^t (t-\tau)^{-\alpha-1} f(\tau) d\tau, \end{aligned}$$

setting  $I = \frac{1}{\Gamma(-\alpha)} \int_{t-L}^t (t-\tau)^{-\alpha-1} f(\tau) d\tau$ , and performing successive integrations by parts one obtains

$$\begin{aligned} I &= \frac{f(t-L)L^{-\alpha}}{\Gamma(1-\alpha)} + \frac{1}{\Gamma(1-\alpha)} \int_{t-L}^t (t-\tau)^{-\alpha} f'(\tau) d\tau, \\ &= \frac{f(t-L)L^{-\alpha}}{\Gamma(1-\alpha)} + \frac{f'(t-L)L^{1-\alpha}}{\Gamma(2-\alpha)} + \frac{1}{\Gamma(2-\alpha)} \int_{t-L}^t (t-\tau)^{-\alpha+1} f^{(2)}(\tau) d\tau, \\ &\vdots \\ &= \sum_{k=0}^{m-1} \frac{f^{(k)}(t-L)L^{k-\alpha}}{\Gamma(k-\alpha+1)} + \frac{1}{\Gamma(m-\alpha)} \int_{t-L}^t (t-\tau)^{-\alpha+m-1} f^{(m)}(\tau) d\tau, \\ &= {}^{MG}D_t^\alpha f(t). \end{aligned}$$

Therefore

$${}^{MRL}D_t^\alpha f(t) = {}^{MG}D_t^\alpha f(t) - \sum_{k=0}^{m-1} \frac{f^{(k)}(t-L)L^{k-\alpha}}{\Gamma(k-\alpha+1)}.$$

□

**Remark 3.7.** From (3.5) and (3.6) one has

$${}^{MRL}D_t^\alpha f(t) = {}^{MC}D_t^\alpha f(t) = {}^{MG}D_t^\alpha f(t) - \sum_{k=0}^{m-1} \frac{f^{(k)}(t-L)L^{k-\alpha}}{\Gamma(k-\alpha+1)}. \quad (3.7)$$

In the following parts, one denotes the operators of Caputo and Riemann-Liouville fractional derivative with sliding fixed memory length by  ${}^M_L D_t^\alpha$ .

### 3.1 Fractional derivative of some elementary functions

In order to highlight the amazing properties of the fractional derivative with sliding fixed memory length one consider two elementary functions (the power and exponential functions), for which one computes their new derivatives.

### 3.1.1 New fractional derivative of the power function

Let  $f(t) = t^n$ ,  $n \in \mathbb{N}^*$ ,  $\alpha > 0$ ,  $L > 0$  and  $m$  is an integer such that  $m - 1 < \alpha < m$ .

If  $n < m$ , then  $f^{(m)}(t) = 0$ , substituting in (3.3) yields  ${}^M_L D_t^\alpha(t^n) = 0$ .

If  $n \geq m$  then by repeated integrations by parts of the relation (3.3) one obtains

$${}^M_L D_t^\alpha(t^n) = \sum_{k=0}^{n-m} \frac{n!L^{-\alpha+m+k}(t-L)^{n-m-k}}{(n-m-k)!\Gamma(-\alpha+m+k+1)}. \quad (3.8)$$

**Remark 3.8.** (Fractional derivative of a constant function)

If  $f$  is a constant function (i.e.  $f(t) = C$  for all  $t \in [a-L, b]$ , and  $C$  any constant including zero) then one has

$${}^M_L D_t^\alpha C = 0.$$

### 3.1.2 Fractional derivative of the exponential function

Let  $f(t) = e^t = \sum_{p=0}^{\infty} \frac{t^p}{p!}$ ,  $\alpha > 0$ ,  $L > 0$  and  $m$  is an integer such that  $m - 1 < \alpha < m$ .

One has

$${}^M_L D_t^\alpha e^t = {}^M_L D_t^\alpha \sum_{p=0}^{\infty} \frac{t^p}{p!} = \sum_{p=0}^{\infty} \frac{1}{p!} {}^M_L D_t^\alpha t^p.$$

From (3.8), one obtains that

$$\begin{aligned} {}^M_L D_t^\alpha(e^t) &= \sum_{p=0}^{\infty} \sum_{k=0}^{p-m} \frac{L^{-\alpha+m+k}(t-L)^{p-m-k}}{(p-m-k)!\Gamma(-\alpha+m+1+k)}, \\ &= \sum_{p=0}^{\infty} \sum_{k=0}^{p-m} \frac{L^{-\alpha+m+k}(t-L)^{p-m-k}}{(p-m-k)!\Gamma(k-\alpha+m+1)}, \\ &= \sum_{p=0}^{\infty} \sum_{k=0}^p \frac{L^{-\alpha+m+k}(t-L)^{p-k}}{(p-k)!\Gamma(k-\alpha+m+1)}, \\ &= \sum_{p=0}^{\infty} \frac{L^{-\alpha+m}(t-L)^p}{p!\Gamma(-\alpha+m+1)} + \sum_{p=0}^{\infty} \frac{L^{-\alpha+m+1}(t-L)^p}{p!\Gamma(-\alpha+m+2)} + \dots, \\ &= \left( \sum_{p=0}^{\infty} \frac{(t-L)^p}{p!\Gamma(-\alpha+m+1)} \right) \left( \sum_{k=0}^{\infty} \frac{L^{-\alpha+m+k}}{\Gamma(-\alpha+m+1+k)} \right), \\ &= e^{t-L} L^{-\alpha+m} \sum_{k=0}^{\infty} \frac{L^k}{\Gamma(-\alpha+m+1+k)}, \\ &= e^{t-L} L^{m-\alpha} E_{1, m+1-\alpha}(L). \end{aligned}$$

## 3.2 Derivative of a periodic function

The main result of this paper is stated in the following theorem.

**Theorem 3.9.** Let  $\alpha > 0$ ,  $L > 0$  and  $m$  an integer such that  $m - 1 < \alpha < m$  and  $f \in C^m[a-L, b]$ . If  $f$  is a periodic function with period  $T$ , Then  ${}^M_L D_t^\alpha f$  is a periodic function with the same period  $T$ .

*Proof.* Suppose that  $f$  is a periodic function with a period  $T$ . The aim of this proof is to demonstrate that the function  $g(t) = {}^M_L D_t^\alpha f$  is a periodic function with the same period  $T$  (i.e.

$g(t+T) = g(t)$ .

One has

$$\begin{aligned} g(t+T) &= {}^M_L D_{t+T}^\alpha f(t+T) = \frac{1}{\Gamma(m-\alpha)} \int_{t+T-L}^{t+T} (t+T-\tau)^{m-\alpha-1} f^{(m)}(\tau+T) d\tau, \\ &= \frac{1}{\Gamma(m-\alpha)} \int_{t-L}^t (t-s)^{m-\alpha-1} f^{(m)}(s+2T) ds, \\ &= \frac{1}{\Gamma(m-\alpha)} \int_{t-L}^t (t-s)^{m-\alpha-1} f^{(m)}(s) ds, \\ &= {}^M_L D_t^\alpha f(t) = g(t). \end{aligned}$$

Thus,  ${}^M_L D_t^\alpha f$  is a periodic function with the same period  $T$ .

□

### 3.2.1 Fractional derivative of some fundamental periodic functions

Note first that the functions  ${}^{MG}_L D_t^\alpha \sin(t)$  and  ${}^{MG}_L D_t^\alpha \cos(t)$  have been calculated in [1].

**Example 3.10.** (Fractional derivative with sliding fixed memory length of the sine function)

By definition

$${}^M_L D_t^\alpha f(t) = {}^{MG}_L D_t^\alpha f(t) - \sum_{k=0}^{m-1} \frac{f^{(k)}(t-L)L^{k-\alpha}}{\Gamma(k-\alpha+1)}.$$

Therefore

$$\begin{aligned} {}^M_L D_t^\alpha \sin(t) &= {}^{MG}_L D_t^\alpha \sin(t) - \sum_{k=0}^{m-1} \frac{\frac{d^k}{dt^k}(\sin(t-L))L^{k-\alpha}}{\Gamma(k-\alpha+1)}, \\ &= L^{-\alpha} \sin(t-L)E_{2,1-\alpha}(-L^2) + L^{1-\alpha} \cos(t-L)E_{2,2-\alpha}(-L^2) \\ &\quad - L^{-\alpha} \sum_{k=0}^{\lfloor \frac{m-1}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+1-\alpha)} \sin(t-L) - L^{1-\alpha} \sum_{k=0}^{\lfloor \frac{m-2}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+2-\alpha)} \cos(t-L), \\ &= L^{-\alpha} \sin(t-L) \left( E_{2,1-\alpha}(-L^2) - \sum_{k=0}^{\lfloor \frac{m-1}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+1-\alpha)} \right) \\ &\quad + L^{1-\alpha} \cos(t-L) \left( E_{2,2-\alpha}(-L^2) - \sum_{k=0}^{\lfloor \frac{m-2}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+2-\alpha)} \right), \\ &= a \sin(t-L) + b \cos(t-L), \end{aligned} \tag{3.9}$$

$$\text{where, } a = L^{-\alpha} \left( E_{2,1-\alpha}(-L^2) - \sum_{k=0}^{\lfloor \frac{m-1}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+1-\alpha)} \right), \quad b = L^{1-\alpha} \left( E_{2,2-\alpha}(-L^2) - \sum_{k=0}^{\lfloor \frac{m-2}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+2-\alpha)} \right).$$

One notices that,  ${}^M_L D_t^\alpha \sin(t)$  is a periodic function with the period  $T = 2\pi$ . This analytical result is displayed in figure (3.1), for some values of  $\alpha$  and  $L = 32.1$ .

**Example 3.11.** (Fractional derivative of cosine function)






Figure 3.1: Fractional derivative of the Sine function for  $L = 32.1$  and some values of  $\alpha$ .

By definition

$$\begin{aligned}
 {}^M D_t^\alpha \cos(t) &= {}^{MG} D_t^\alpha \cos(t) - \sum_{k=0}^{m-1} \frac{d^k}{dt^k} (\cos(t-L)) L^{k-\alpha}, \\
 &= L^{-\alpha} \cos(t-L) E_{2,1-\alpha}(-L^2) - L^{1-\alpha} \sin(t-L) E_{2,2-\alpha}(-L^2) \\
 &\quad - L^{-\alpha} \sum_{k=0}^{\lfloor \frac{m-1}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+1-\alpha)} \cos(t-L) + L^{1-\alpha} \sum_{k=0}^{\lfloor \frac{m-2}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+2-\alpha)} \sin(t-L), \\
 &= L^{-\alpha} \cos(t-L) (E_{2,1-\alpha}(-L^2) - \sum_{k=0}^{\lfloor \frac{m-1}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+1-\alpha)}) \\
 &\quad - L^{1-\alpha} \sin(t-L) (E_{2,2-\alpha}(-L^2) - \sum_{k=0}^{\lfloor \frac{m-2}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+2-\alpha)}), \\
 &= a \cos(t-L) - b \sin(t-L), \tag{3.10}
 \end{aligned}$$

where,

$$a = L^{-\alpha} (E_{2,1-\alpha}(-L^2) - \sum_{k=0}^{\lfloor \frac{m-1}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+1-\alpha)}),$$

and

$$b = L^{1-\alpha} (E_{2,2-\alpha}(-L^2) - \sum_{k=0}^{\lfloor \frac{m-2}{2} \rfloor} \frac{(-L^2)^k}{\Gamma(2k+2-\alpha)}).$$

Obviously  ${}^M D_t^\alpha \cos(t)$  is a periodic function with period  $T = 2\pi$ .

### 3.3 An interpolation property

It is known that the operator of Grünwald-Letnikov fractional derivative with sliding fixed memory length is an extension of the integer-order operator  $\frac{d^m}{dt^m}$ , (see [1]).

The following proposition proves that the Caputo and Riemann-Liouville operators of the

fractional derivative with sliding fixed memory length verifies this property for  $\alpha \rightarrow m$ , but not for  $\alpha \rightarrow m - 1$ .

**Proposition 3.12.** Let  $L > 0$  and  $0 \leq m - 1 < \alpha < m$  such that  $m$  is an integer number, and let  $f(t)$  having  $(m + 1)$  continuous bounded derivatives in  $[a - L, b]$ . Then, for all  $t \in [a, b]$ , one has

$$\lim_{\alpha \rightarrow m} {}^M_L D_t^\alpha f(t) = f^{(m)}(t),$$

and

$$\lim_{\alpha \rightarrow m-1} {}^M_L D_t^\alpha f(t) = f^{(m-1)}(t) - f^{(m-1)}(t - L).$$

*Proof.* One has

$$\begin{aligned} \lim_{\alpha \rightarrow m} {}^M_L D_t^\alpha f(t) &= \lim_{\alpha \rightarrow m} \frac{1}{\Gamma(m - \alpha)} \int_{t-L}^t (t - \tau)^{m-\alpha-1} f^{(m)}(\tau) d\tau, \\ &= \lim_{\alpha \rightarrow m} \frac{L^{m-\alpha} f^{(m)}(t - L)}{\Gamma(m - \alpha + 1)} + \lim_{\alpha \rightarrow m} \frac{1}{\Gamma(m - \alpha + 1)} \\ &\quad \int_{t-L}^t (t - \tau)^{m-\alpha} f^{(m+1)}(\tau) d\tau, \\ &= f^{(m)}(t - L) + \int_{t-L}^t f^{(m+1)}(\tau) d\tau, \\ &= f^{(m)}(t). \end{aligned}$$

For  $\alpha \rightarrow m - 1$ , one has

$$\begin{aligned} \lim_{\alpha \rightarrow m-1} {}^{MC}_L D_t^\alpha f(t) &= \lim_{\alpha \rightarrow m-1} \frac{1}{\Gamma(m - \alpha)} \int_{t-L}^t (t - \tau)^{m-\alpha-1} f^{(m)}(\tau) d\tau, \\ &= \int_{t-L}^t f^{(m)}(\tau) d\tau, \\ &= f^{(m-1)}(t) - f^{(m-1)}(t - L). \end{aligned}$$

□

**Example 3.13.**

Let  $f(t) = e^t$ , then

$${}^M_L D_t^\alpha e^t = e^{t-L} L^{m-\alpha} E_{1, m+1-\alpha}(L),$$

Therefore,

$$\lim_{\alpha \rightarrow m} {}^M_L D_t^\alpha e^t = e^{t-L} E_{1,1}(L) = e^t = f^{(m)}(t).$$

However,

$$\begin{aligned} \lim_{\alpha \rightarrow m-1} {}^M_L D_t^\alpha e^t &= e^{t-L} L E_{1,2}(L) = e^{t-L} (e^L - 1), \\ &= e^t - e^{t-L} = f^{(m)}(t) - f^{(m-1)}(t - L). \end{aligned}$$

**Example 3.14.**

Let  $f(t) = t^n$ , one has

$${}^M_L D_t^\alpha (t^n) = \sum_{k=0}^{n-m} \frac{n! L^{-\alpha+m+k} (t-L)^{n-m-k}}{(n-m-k)! \Gamma(-\alpha+m+k+1)}.$$

Putting  $N = n - m$  and  $t - L = a$ , then

$$\begin{aligned} \lim_{\alpha \rightarrow m} {}^M D_t^\alpha (t^n) &= \sum_{k=0}^N \frac{n! L^k a^{N-k}}{(N-k)! k!} \\ &= \frac{n!}{N!} \sum_{k=0}^N \frac{N! L^k a^{N-k}}{(N-k)! k!} \\ &= \frac{n!}{N!} (a+L)^N = \frac{n!}{(n-m)!} t^{n-m}, \\ &= \frac{d^m}{dt} t^n = f^{(m)}(t). \end{aligned}$$

However,

$$\begin{aligned} \lim_{\alpha \rightarrow m-1} {}^M D_t^\alpha (t^n) &= \sum_{k=0}^N \frac{n! L^{k+1} a^{N-k}}{(N-k)! (k+1)!} \\ &= \frac{n!}{(N+1)!} \sum_{k=0}^{N+1} \frac{(N+1)! L^k a^{N+1-k}}{(N+1-k)! k!} - \frac{n!}{(n-m+1)!} (t-L)^{n-m+1}, \\ &= \frac{n!}{(N+1)!} t^{N+1} - \frac{n!}{(n-(m-1))!} (t-L)^{n-(m-1)}, \\ &= \frac{n!}{(n-(m-1))!} t^{n-(m-1)} - \frac{n!}{(n-(m-1))!} (t-L)^{n-(m-1)}, \\ &= \frac{d^{m-1}}{dt} t^n - \frac{d^{m-1}}{dt} (t-L)^n = f^{(m-1)}(t) - f^{(m-1)}(t-L). \end{aligned}$$

### 3.4 Comparison between some results of classical fractional-order derivatives and fractional order derivatives with sliding fixed memory length

The previous results are summarized in the table (3.1), in order to highlight the differences between classical fractional-order derivative and fractional-order derivative with sliding fixed memory length.

### 3.5 Fractional-order autonomous system with exact periodic solution

As previously mentioned, any autonomous fractional-order system expressed in terms of classical fractional derivatives cannot have any exact periodic solutions [13, 22, 23]. Conversely to these results, one presents some examples showing that fractional-order autonomous systems (linear and nonlinear) expressed in terms of fractional derivatives with sliding fixed memory length can have exact periodic solutions.

**Example 3.15.** (Linear fractional-order system)

Let consider the following linear fractional-order autonomous system

$${}^M D_t^\alpha X(t) = AX(t), \tag{3.11}$$

where  $X(t) \in \mathbb{R}$  and  $A = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}$ , with  $a = L^{-\alpha} (E_{2,1-\alpha}(-L^2) - \sum_{p=0}^{[\frac{m-1}{2}]} \frac{(-L^2)^p}{\Gamma(2p+1-\alpha)})$ ,

$b = L^{1-\alpha} (E_{2,2-\alpha}(-L^2) - \sum_{p=0}^{[\frac{m-2}{2}]} \frac{(-L^2)^p}{\Gamma(2p+2-\alpha)})$ .

Classical fractional derivative ${}^C_a D_t^\alpha$ or ${}^{RL}_a D_t^\alpha$	Fractional derivative with sliding fixed memory length ${}^M_L D_t^\alpha$
${}^C_a D_t^\alpha f(t) = {}^{RL}_a D_t^\alpha f(t) - \sum_{k=0}^{m-1} \frac{f^{(k)}(a)(t-a)^{k-\alpha}}{\Gamma(k-\alpha+1)}$	${}^{MC}_L D_t^\alpha f(t) = {}^{MR}_L D_t^\alpha f(t)$
$\lim_{\alpha \rightarrow m} {}^{RL}_a D_t^\alpha f(t) = \lim_{\alpha \rightarrow m} {}^C_a D_t^\alpha f(t) = f^{(m)}(t)$	$\lim_{\alpha \rightarrow m} {}^M_L D_t^\alpha f(t) = f^{(m)}(t)$
$\lim_{\alpha \rightarrow m-1} {}^{RL}_a D_t^\alpha f(t) = f^{(m-1)}(t),$ $\lim_{\alpha \rightarrow m-1} {}^C_a D_t^\alpha f(t) = f^{(m-1)}(t) - f^{(m-1)}(a)$	$\lim_{\alpha \rightarrow m-1} {}^M_L D_t^\alpha f(t) = f^{(m-1)}(t) - f^{(m-1)}(t-L)$
${}^{RL}_0 D_t^\alpha (t^n) = {}^C_0 D_t^\alpha (t^n) = \frac{\Gamma(n+1)}{\Gamma(n-\alpha+1)} t^{n-\alpha}$	${}^M_L D_t^\alpha (t^n) = \sum_{k=0}^{n-m} \frac{n! L^{-\alpha+m+k} (t-L)^{n-m-k}}{(n-m-k)! \Gamma(-\alpha+m+k+1)}$
${}^{RL}_a D_t^\alpha C = \frac{C}{\Gamma(1-\alpha)} (t-a)^\alpha \neq 0,$ ${}^C_a D_t^\alpha C = 0$	${}^M_L D_t^\alpha C = 0$
${}^{RL}_a D_t^\alpha \sin t = t^{1-\alpha} E_{2,2-\alpha}(-t^2)$	${}^M_a D_t^\alpha \sin t = a \sin(t-L) + b \cos(t-L).$

Table 3.1: Comparison between some results of classical fractional-order derivatives and fractional order derivatives with sliding fixed memory length.

- For  $L = 2k\pi$ , where  $k$  is a non-zero integer. The vector function  $X(t) = c \begin{pmatrix} \cos(t) \\ \sin(t) \end{pmatrix}$ ,  $c \in \mathbb{R}$  is an exact  $2\pi$ -periodic solution for the system (3.11).

By definition,

$${}^{M}_{2k\pi} D_t^\alpha X(t) = c \begin{pmatrix} {}^M_{2k\pi} D_t^\alpha \cos(t) \\ {}^M_{2k\pi} D_t^\alpha \sin(t) \end{pmatrix}.$$

Then, from (3.9) and (3.10) one obtains

$$\begin{aligned} {}^{M}_{2k\pi} D_t^\alpha X(t) &= c \begin{pmatrix} a \cos(t - 2k\pi) - b \sin(t - 2k\pi) \\ a \sin(t - 2k\pi) + b \cos(t - 2k\pi) \end{pmatrix}, \\ &= c \begin{pmatrix} a & -b \\ b & a \end{pmatrix} \begin{pmatrix} \cos(t - 2k\pi) \\ \sin(t - 2k\pi) \end{pmatrix}, \\ &= cA \begin{pmatrix} \cos(t - 2k\pi) \\ \sin(t - 2k\pi) \end{pmatrix}, \\ &= AX(t). \end{aligned}$$

Therefore,  $X(t) = c \begin{pmatrix} \cos(t) \\ \sin(t) \end{pmatrix}$  is an exact  $2\pi$ -periodic solution of (3.11) with  $L = 2k\pi$ .

- For  $L = \frac{\pi}{2}$ , one has

$$\begin{aligned} {}^{M}_{\frac{\pi}{2}} D_t^\alpha X(t) &= c \begin{pmatrix} a \cos(t - \frac{\pi}{2}) - b \sin(t - \frac{\pi}{2}) \\ a \sin(t - \frac{\pi}{2}) + b \cos(t - \frac{\pi}{2}) \end{pmatrix}, \\ &= c \begin{pmatrix} a \sin(t) + b \cos(t) \\ -a \cos(t) + b \sin(t) \end{pmatrix}, \\ &= c \begin{pmatrix} b & a \\ -a & b \end{pmatrix} \begin{pmatrix} \cos(t) \\ \sin(t) \end{pmatrix}, \\ &= cB \begin{pmatrix} \cos(t) \\ \sin(t) \end{pmatrix}, \\ &= BX(t), \end{aligned}$$

with  $B = \begin{pmatrix} b & a \\ -a & b \end{pmatrix} \neq A$ . Thus,  $X(t) = c \begin{pmatrix} \cos(t) \\ \sin(t) \end{pmatrix}$  is not solution of (3.11), but it is an exact  $2\pi$ -periodic solution of the system  ${}^M D_t^\alpha X(t) = BX(t)$ .

**Example 3.16.** (The predator-prey model with Holling type II response function)

All population species possess the property of heredity, which means the passing on traits from parents to their offspring, either through asexual reproduction or sexual reproduction. The offspring cells or organisms acquire the genetic information of their parents through heredity. This property makes fractional differential systems models more efficiently regarding some specific problems than ordinary differential ones.

Motivated by this fact, we introduce the fractional version of the Holling-Tanner model [21] as follows

$$\begin{cases} D^\alpha x = r_1 x \left(1 - \frac{x}{K}\right) - \frac{qxy}{m+x}, \\ D^\alpha y = r_2 y \left(1 - \frac{y}{\gamma x}\right). \end{cases} \quad (3.12)$$

Where  $D \cdot$  denotes a standard fractional-order derivative operator and  $\alpha \in [0, 1]$  is the fractional-order related to the hereditary property of the population (a value of  $\alpha$  close to an integer number means that the population has a weak hereditary property),  $x(t) \geq 0$  and  $y(t) \geq 0$  are the density of prey and predator populations at time  $t$  respectively. The parameters  $r_1$  and  $r_2$  are the intrinsic growth rates,  $K$  represents the carrying capacity of the prey,  $q$  is the maximum number of preys that can be eaten per predator per unit of time,  $m$  is the saturation value (it corresponds to the number of preys necessary to achieve one half the maximum rate  $q$ ),  $\gamma$  is a measure of the quality of the prey as a portion of food for the predator.

Since exact analytical resolution of this nonlinear system is unavailable, one resorts to qualitative and numerical study. For this purpose the parameters are set to  $r_1 = 1$ ,  $r_2 = 0.2$ ,  $K = 25$ ,  $q = \frac{6}{7}$ ,  $m = 1$  and  $\gamma = 0.95$ , the system (3.12) has two equilibrium points  $E_0 = (25, 0)$  and  $E_1 \approx (7.1429, 6.7857)$ .

- The characteristic polynomial of the Jacobian matrix evaluated at  $E_0$  is given by

$$P(\lambda) = \lambda^2 + a_1 \lambda + a_2 = \lambda^2 + 0.8\lambda - 0.2.$$

So  $a_2 = -0.2 < 0$ , then according to Proposition 1 in [7]  $E_0$  is unstable for all  $\alpha \in [0, 2)$ .

- The characteristic polynomial of the Jacobian matrix evaluated at  $E_1$  is given by

$$P(\lambda) = \lambda^2 - 0.1409\lambda + 0.0747.$$

So  $a_1 \approx -0.1409$  and  $a_2 \approx 0.0747 > 0$ .

Applying Hopf-Like Bifurcation theory [2–4] and using Proposition 1 in [7], one obtains the Hopf-Like bifurcation value

$$\alpha^* = \frac{2}{\pi} \cos^{-1} \left( \frac{-a_1}{2\sqrt{a_2}} \right) \approx 0.8341.$$

The fixed point  $E_1$  losses its stability, and a periodic motion ( $S$ -asymptotically periodic for the classical fractional derivative and exact periodic for fractional derivative with sliding fixed memory length) appears.




Figure 3.2: Time evolution and phase portrait of system (3.12) for  $\alpha = 0.9$  (a,b)  $S$ -asymptotically  $T$ -periodic solution with  $T \approx 27.2$  for classical fractional operator. (c,d) Exact  $T$ -periodic solution for the fractional derivative operator with sliding fixed memory length. (e,f) Comparison between the two solutions.

To illustrate these results, one solves the system (3.12) numerically by developing a Matlab code using a discretization technique based on the formula (3.7).

Choosing a value for  $\alpha$  greater than  $\alpha^*$ , for example,  $\alpha = 0.9$ , one compares the solution of (3.12) in terms of classical fractional operator and its solution in terms of the fractional operator with sliding fixed memory length  $L = 30$ . The two trajectories are start from the same initial point  $X_0 = (2.64, 4.88)$ , belonging to the attracting limit cycle. The results are shown in Fig. 3.2.

An  $S$ -asymptotically  $T$ -periodic solution with  $T \approx 27.2$  is obtained for classical fractional operator as shown in Fig. 3.2(a,b), and an exact  $T$ -periodic solution is obtained for the fractional derivative operator with sliding fixed memory length as shown in Fig. 3.2(c,d).

## 4 Conclusion

In this article, one modifies the Caputo and Riemann-Liouville fractional-order derivatives by fixing the memory length and varying the lower terminal of the derivative. It is shown that

the modified fractional derivative operator preserves the periodicity. Consequently, periodic solutions can be obtained in fractional-order systems expressed in terms of the new operator. Two examples are investigated to highlight this property for a linear system provides an analytic expression of an exact periodic solution is computed and for another nonlinear system for which exact periodic solutions are obtained using qualitative and numerical methods.

## Acknowledgments

This research was supported by the Algerian General Directorate for Scientific Research and Technological Development (DGRSDT).

## Conflict of Interest

The authors have no conflicts of interest to disclose.

## References

- [1] M-S. ABDELOUAHAB AND N. HAMRI, *The Grünwald-Letnikov fractional-order derivative with fixed memory length*, *Mediterr. J. Math.* **13** (2016), 557–572.
- [2] M-S. ABDELOUAHAB, N. HAMRI AND J. W. WANG, *Hopf bifurcation and chaos in fractional-order modified hybrid optical system*, *Nonlinear Dyn.* **69** (2012), 275–284.
- [3] M-S. ABDELOUAHAB AND R. LOZI, *Hopf Bifurcation and Chaos in Simplest Fractional-Order Memristor-based Electrical Circuit*, *Indian. J. Ind. Appl. Math.* **6**(2) (2015), 105–119.
- [4] M-S. ABDELOUAHAB, R. LOZI AND G. CHEN, *Complex Canard Explosion in a Fractional-Order FitzHugh–Nagumo Model*, *Int. J. Bifurcation Chaos.* **29**(8) (2019), 1950111–1950133.
- [5] I. AREA, J. LOSADA AND J. J. NIETO, *On quasi-periodicity properties of fractional integrals and fractional derivatives of periodic functions*, *Integral Transforms Spec. Funct.* **27** (2016), 1–16.
- [6] R. L. BAGLEY AND R. A. CALICO, *Fractional order state equations for the control of viscoelastically damped structures*, *J. Guid. Control Dyn.* **14** (1991), 304–311.
- [7] S. BOURAFA, M-S. ABDELOUAHAB AND A. MOUSSAOUI, *On some extended Routh-Hurwitz conditions for fractional-order autonomous systems of order  $\alpha \in [0, 2)$  and their applications to some population dynamic models*, *Chaos Solitons Fractals.* **133** (2020), 109623.
- [8] P. L. BUTZER AND U. WESTPHAL, *An introduction to fractional calculus*, In: Hilfer, R. *Applications of Fractional Calculus in Physics*, World Scientific, Singapore, 2000.
- [9] M. CAPUTO, *Linear models of dissipation whose  $Q$  is almost frequency independent-II*, *Geophys. J. R. Astron. Soc.* **13** (1967), 529–539.
- [10] O. HEAVISIDE, H. J. JOSEPHS AND B. A. BEHREND, *Electromagnetic Theory*, Chelsea Publishing Company, New York, 1971.
- [11] M. ICHISE, Y. NAGAYANAGI AND T. KOJIMA, *An analog simulation of noninteger order transfer functions for analysis of electrode process*, *J. Electroanal. Chem.* **33** (1971), 253–265.

- [12] Y-M. KANG, Y. XIE, J-C. LU AND J. JIANG, *On the nonexistence of non-constant exact periodic solutions in a class of the Caputo fractional-order dynamical systems*, *Nonlinear Dyn.* **82** (2015), 1259–1267.
- [13] E. KASLIK AND S. SIVASUNDARAM, *Non-existence of periodic solutions in fractional-order dynamical systems and a remarkable difference between integer and fractional-order derivatives of periodic functions*, *Nonlinear Analysis: Real World Applications.* **13** (2012), 1489–1497.
- [14] D. KUSNEZOV, A. BULGAC AND G. D. DANG, *Quantum levy processes and fractional kinetics*, *Phys. Rev. Lett.* **82** (1999), 1136–1139.
- [15] C. A. MONJE, Y. Q. CHEN, B. M. VINAGRE, D. XUE, AND V. FELIU, *Fractional-Order Systems and Controls Fundamentals and Applications*, Springer-Verlag London Limited, London, 2010.
- [16] K. B. OLDHAM AND J. SPANIER, *The Fractional Calculus Theory and Applications of Differentiation and Integration to Arbitrary Order*, Academic press, inc, USA, 1974.
- [17] I. PODLUBNY, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [18] B. ROSS, *The development of fractional calculus 1695-1900*, *Hist. Math.* **4** (1977), 75–89.
- [19] S. G. SAMKO, A. A. KILBAS AND O. I. MARICHEV, *Fractional Integrals and Derivatives: Theory and Applications*, Gordon and Breach, Amsterdam, 1993.
- [20] H. H. SUN, A. A. ABDELWAHAB AND B. ONARAL, *Linear approximation of transfer function with a pole of fractional order*, *IEEE Trans. Autom. Control.* **29** (1984), 441–444.
- [21] J. T. TANNER, *The stability and the intrinsic growth rates of prey and predator populations*, *Ecology.* **56** (1975), 855–867.
- [22] M. S. TAVAZOEI, *A note on fractional-order derivatives of periodic functions.* *Automatica J. IFAC.* **46** (2010), 945–948.
- [23] M. S. TAVAZOEI AND M. HAERI, *A proof for non existence of periodic solutions in time invariant fractional order systems*, *Automatica J. IFAC.* **45** (2009), 1886–1890.