



**HAL**  
open science

# On the Kernel Curves Associated with Walks in the Quarter Plane

Thomas Dreyfus, Charlotte Hardouin, Julien Roques, Michael F Singer

► **To cite this version:**

Thomas Dreyfus, Charlotte Hardouin, Julien Roques, Michael F Singer. On the Kernel Curves Associated with Walks in the Quarter Plane. Alin Bostan; Kilian Raschel. Transcendence in Algebra, Combinatorics, Geometry and Number Theory. TRANS19 – Transient Transcendence in Transylvania, Braşov, Romania, May 13–17, 2019, Revised and Extended Contributions, 373, Springer International Publishing, pp.61-89, 2021, Springer Proceedings in Mathematics & Statistics, 978-3-030-84303-8. 10.1007/978-3-030-84304-5\_3. hal-03513338

**HAL Id: hal-03513338**

**<https://hal.science/hal-03513338v1>**

Submitted on 5 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON THE KERNEL CURVES ASSOCIATED WITH WALKS IN THE QUARTER PLANE

THOMAS DREYFUS, CHARLOTTE HARDOUIN, JULIEN ROQUES, AND MICHAEL F. SINGER

ABSTRACT. The *kernel method* is an essential tool for the study of generating series of *walks* in the quarter plane. This method involves equating to zero a certain polynomial - the *kernel polynomial* - and using properties of the curve - the *kernel curve* - this defines. In the present paper, we investigate the basic properties of the *kernel curve* (irreducibility, singularities, genus, uniformization, etc).

## CONTENTS

Introduction	1
1. Nondegenerate walks	4
2. Singularities and genus of the kernel curve	8
3. Involution automorphisms of the kernel curve	15
4. Uniformization of the kernel curve	18
References	24

## INTRODUCTION

Consider a *walk* with small steps in the positive quadrant  $\mathbb{Z}_{\geq 0}^2 = \{0, 1, 2, \dots\}^2$  starting from  $P_0 := (0, 0)$ , that is a succession of points

$$P_0, P_1, \dots, P_k,$$

where each  $P_n$  lies in the quarter plane, where the moves (or steps)  $P_{n+1} - P_n$  belong to  $\{0, \pm 1\}^2$ , and the probability to move in the direction  $P_{n+1} - P_n = (i, j)$  may be interpreted as some *weight-parameter*  $d_{i,j} \in [0, 1]$ , with  $\sum_{(i,j) \in \{0, \pm 1\}^2} d_{i,j} = 1$ . The step set or the *model* of the *walk* is the set of directions with nonzero *weights*, that is

$$\mathcal{S} = \{(i, j) \in \{0, \pm 1\}^2 \mid d_{i,j} \neq 0\}.$$

The following picture is an example of such path:

---

*Date:* December 11, 2020.

2010 *Mathematics Subject Classification.* 05A15, 30D05.

*Key words and phrases.* Random walks, Uniformization of algebraic curves.

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under the Grant Agreement No 648132. The authors would like to thank ANR-19-CE40-0018, ANR-13-JS01-0002-01, ANR-11-LABX-0040-CIMI within the program ANR-11-IDEX-0002-0, LabEx PERSYVAL-Lab ANR-11-LABX-0025-01, the Simons Foundation (#349357, Michael Singer).



The second step of the *kernel method* is to exploit certain involutive birational transformations  $\iota_1, \iota_2$  (they are called  $\zeta, \eta$  in [FIM17]) of the *kernel curve*  $E_t$  of the form

$$\iota_1(x, y) = (x, y') \text{ and } \iota_2(x, y) = (x', y)$$

in order to deduce from (1) some functional equations for  $Q(x, 0, t)$  and  $Q(0, y, t)$ . Hence  $\iota_1$  and  $\iota_2$  switch the roots of the degree two polynomials  $y \mapsto K(x, y, t)$  and  $x \mapsto K(x, y, t)$  respectively. Concretely, the birational transformations  $\iota_1, \iota_2$  are induced by restriction to the curve of the involutive birational transformations  $i_1, i_2$  of  $\mathbb{C}^2$  given by

$$i_1(x, y) = \left( x, \frac{A_{-1}(x)}{A_1(x)y} \right) \text{ and } i_2(x, y) = \left( \frac{B_{-1}(y)}{B_1(y)x}, y \right)$$

where the  $A_i(x) \in x^{-1}\mathbb{Q}[x]$  and the  $B_i(y) \in y^{-1}\mathbb{Q}[y]$  are defined by

$$S(x, y) = A_{-1}(x)\frac{1}{y} + A_0(x) + A_1(x)y = B_{-1}(y)\frac{1}{x} + B_0(y) + B_1(y)x,$$

see [BMM10, Section 3], [KY15, Section 3] or [FIM17]. These  $i_1$  and  $i_2$  are the generators of the group of the *walk*; see [BMM10] for details. Note that although  $i_1$  and  $i_2$  do not depend on  $t$ , the group generated by the induced involutions  $\iota_1$  and  $\iota_2$  may depend on  $t$ , since the order of  $\iota_2 \circ \iota_1$  may depend upon  $t$ , see Remark 4.13.

The third step of the *kernel method* is to use the above mentioned functional equations of  $Q(x, 0, t)$  and  $Q(0, y, t)$  to continue these series as multivalued meromorphic functions. To perform this step, we need an explicit uniformization of the *kernel curve*.

The aim of the present paper is to study the *kernel curve*  $E_t$  and the birational transformations  $\iota_1, \iota_2$ . Note that a similar study has been done in the case  $t = 1$  in [FIM17] and in the *unweighted* case in [KR12]. The goal of the present paper is to extend these works to the *weighted* case when  $t \in ]0, 1[$  is transcendental over  $\mathbb{Q}(d_{i,j})$ . Although many results are similar to [FIM17], the proofs are different. The assumptions we make on  $t$  are crucial in many parts of the proof and it is not clear how the proofs of [FIM17] exactly pass to this context.

We could expect to have classification of the geometric properties of  $\overline{E_t}$  involving configurations of *weights* independent of  $t$ . Hopefully, this paper has been followed by [DR19] where the case for general  $t \in ]0, 1[$  has been considered. The proofs of the latter paper use continuity arguments with respect the parameter  $t$  that permit to deduce many results for algebraic values of  $t$ . Such reasoning needs to be very cautious, and it is not trivial to deduce the results for general  $t \in ]0, 1[$  from the  $t = 1$  case. We will mention explicitly every time if the results are correct for arbitrary values of  $t \in ]0, 1[$ .

The paper is organized as follows. In Section 1, we describe the *nondegenerate models of walks*. In Section 2, we determine the singularities and the genus of the *kernel curve*. In Section 3, we establish the basic properties of  $\iota_1$  and  $\iota_2$ . Finally, in Section 4, we give an explicit uniformization of the *kernel curve*.

**Acknowledgment.** The authors want to warmly thank the referees for their detailed and helpful comments.


## 1. NONDEGENERATE WALKS

From now on, we fix  $t \in ]0, 1[$ , that is transcendental over the field  $\mathbb{Q}(d_{i,j})$ . We start by recalling the notion of *degenerate walks introduced in [FIM17]*.

**Definition 1.1.** A *model of walk* is called *degenerate* if one of the following holds:

- $K(x, y, t)$  is reducible as an element of the polynomial ring  $\mathbb{C}[x, y]$ ,
- $K(x, y, t)$  has  $x$ -degree less than or equal to 1,
- $K(x, y, t)$  has  $y$ -degree less than or equal to 1.


In what follows we will sometimes represent a *model of walks* with arrows. We will also use dashed arrows for a family of models. For instance, the family of models represented by




correspond to models with  $d_{1,1}, d_{1,-1}, d_{0,1} \neq 0$ ,  $d_{1,0} = d_{0,-1} = d_{-1,1} = d_{-1,0} = 0$ , and where nothing is assumed on  $d_{-1,-1}$  and  $d_{0,0}$ . In the following results, the behavior of the *kernel curve* never depends on  $d_{0,0}$ . This is the reason why, to reduce the amount of notations, we have decided to not associate an arrow to  $d_{0,0}$ . The following result is the analog of [FIM17, Lemma 2.3.2], that focuses on the case  $t = 1$ . Our proof differs from the proof of [FIM17, Lemma 2.3.2], which only considered factorization over  $\mathbb{R}[x, y]$ , while in this paper, we need to prove the absence of factorization over  $\mathbb{C}[x, y]$ .

**Proposition 1.2.** A *model of walk* is *degenerate* if and only if at least one of the following holds:

- (1) There exists  $i \in \{-1, 1\}$  such that  $d_{i,-1} = d_{i,0} = d_{i,1} = 0$ . This corresponds to the following families of models of walks



- (2) There exists  $j \in \{-1, 1\}$  such that  $d_{-1,j} = d_{0,j} = d_{1,j} = 0$ . This corresponds to the following families of models of walks



- (3) All the weights are 0 except maybe  $\{d_{1,1}, d_{0,0}, d_{-1,-1}\}$  or  $\{d_{-1,1}, d_{0,0}, d_{1,-1}\}$ . This corresponds to the following families of models of walks



*Proof.* This proof is organized as follows. We begin by showing that (1) (resp. (2)) corresponds to  $K(x, y, t)$  having  $x$ -degree  $\leq 1$  or  $x$ -valuation  $\geq 1$  (resp.  $y$ -degree  $\leq 1$  or  $y$ -valuation  $\geq 1$ ). In these cases, the *model of the walk* is clearly *degenerate*. Assuming (1) and (2) do not hold, we then show that (3) holds if and only if  $K(x, y, t)$  is reducible.

Cases (1) and (2). It is clear that  $K(x, y, t)$  has  $x$ -degree  $\leq 1$  if and only if  $d_{1,-1} = d_{1,0} = d_{1,1} = 0$ . Similarly,  $K(x, y, t)$  has  $y$ -degree  $\leq 1$  if and only if we have  $d_{-1,1} = d_{0,1} = d_{1,1} = 0$ . Furthermore,  $d_{-1,-1} = d_{-1,0} = d_{-1,1} = 0$  if and only if  $K(x, y, t)$  has  $x$ -valuation  $\geq 1$ . Similarly,  $d_{-1,-1} = d_{0,-1} = d_{1,-1} = 0$  if and only if  $K(x, y, t)$  has  $y$ -valuation  $\geq 1$ . In these cases, *the model of the walk* is clearly *degenerate*.

Case (3). We now assume that cases (1) and (2) do not hold. This implies that the model belongs to the family of models  $\left\{ \begin{array}{c} \nearrow \\ \dashrightarrow \\ \nwarrow \end{array} \right\}$  if and only if it belongs to the family of models  $\left\{ \begin{array}{c} \nearrow \\ \swarrow \end{array} \right\}$ . The same holds for the anti-diagonal configuration. If the model belongs to the family of models  $\left\{ \begin{array}{c} \nearrow \\ \swarrow \end{array} \right\}$ , then the *kernel*

$$K(x, y, t) = -d_{-1,-1}t + xy - d_{0,0}txy - d_{1,1}tx^2y^2 \in \mathbb{C}[xy]$$

is a degree two polynomial in  $xy$ . Thus it may be factorized in the following form  $K(x, y, t) = -d_{1,1}t(xy - \alpha)(xy - \beta)$  for some  $\alpha, \beta \in \mathbb{C}$ . If the model belongs to the family of models  $\left\{ \begin{array}{c} \nwarrow \\ \searrow \end{array} \right\}$ , then

$$K(x, y, t) = -d_{-1,1}ty^2 + xy - d_{0,0}txy - d_{1,-1}tx^2.$$

In this situation,  $K(x, y, t)y^{-2} \in \mathbb{C}[x/y]$  may be factorized in the ring  $\mathbb{C}[x/y]$ , proving that  $K(x, y, t)$  may be factorized in  $\mathbb{C}[x, y]$  as well.

Conversely, let us assume that *the model of the walk* is *degenerate*. Recall that we have assumed that cases (1) and (2) do not hold, so  $K(x, y, t)$  has  $x$ - and  $y$ -degree two,  $x$ - and  $y$ -valuation 0, and is reducible. We have to prove that the model belongs to one of the family of models  $\left\{ \begin{array}{c} \nearrow \\ \swarrow \end{array} \right\}$  or  $\left\{ \begin{array}{c} \nwarrow \\ \searrow \end{array} \right\}$ . Let us write a factorization

$$K(x, y, t) = -f_1(x, y)f_2(x, y),$$

with  $f_1(x, y), f_2(x, y) \in \mathbb{C}[x, y]$  not constant. Let us now prove several lemmas on the the polynomials  $f_1(x, y), f_2(x, y) \in \mathbb{C}[x, y]$ .

**Lemma 1.3.** *Both  $f_1(x, y)$  and  $f_2(x, y)$  have bidegree (1, 1).*

*Proof of Lemma 1.3.* Suppose to the contrary that  $f_1(x, y)$  or  $f_2(x, y)$  does not have bidegree (1, 1). Since  $K$  is of bidegree at most (2, 2) then at least one of the  $f_i$ 's has degree 0 in  $x$  or  $y$ . Up to interchange of  $x$  and  $y$  and  $f_1$  and  $f_2$ , we may assume that  $f_1(x, y)$  has  $y$ -degree 0 and we denote it by  $f_1(x)$ . Since we are not in Cases (1) and (2) of Proposition 1.2, the polynomials  $d_{-1,-1}t + d_{0,-1}tx + d_{1,-1}tx^2$  and  $d_{-1,0}t + (d_{0,0}t - 1)x + d_{1,0}tx^2$  are nonzero. By  $K(x, y, t) = -f_1(x)f_2(x, y)$ , we find in particular that  $f_1(x)$  is a common factor of the nonzero polynomials  $d_{-1,-1}t + d_{0,-1}tx + d_{1,-1}tx^2$  and  $d_{-1,0}t + (d_{0,0}t - 1)x + d_{1,0}tx^2$ . Since  $t$  is nonzero, we find that the roots of  $d_{-1,-1}t + d_{0,-1}tx + d_{1,-1}tx^2 = 0$  are algebraic over  $\mathbb{Q}(d_{i,j})$ . On the other hand, since  $t$  is transcendental over  $\mathbb{Q}(d_{i,j})$ , if  $x$  is a root of  $d_{-1,0}t + (d_{0,0}t - 1)x + d_{1,0}tx^2 = 0$  that is algebraic over  $\mathbb{Q}(d_{i,j})$ , then the constant term in  $t$  has to be zero, proving that  $x = 0$ . Therefore, they are polynomials with only zero

as a potential common roots. So the only potential root of  $f_1(x)$  is zero. This means that either  $f_1(x)$  has degree 0, i.e.  $f_1(x) \in \mathbb{C}$ , or  $x$  divides  $f_1(x)$ . In the latter case,  $x$  divides  $K(x, y, t)$ , and we are in Case 1. In both cases, this is a contradiction and proves the lemma.  $\square$

**Lemma 1.4.** *The polynomials  $f_1(x, y)$  and  $f_2(x, y)$  are irreducible in the ring  $\mathbb{C}[x, y]$ .*

*Proof of Lemma 1.4.* To the contrary, suppose that we can find a factorization  $f_1(x, y) = (ax - b)(cy - d)$  for some  $a, b, c, d \in \mathbb{C}$ . Since  $f_1(x, y)$  has bidegree  $(1, 1)$ , we have  $ac \neq 0$ . We then have that

$$0 = K(b/a, y, t) = \frac{b}{a}y - t(\tilde{A}_{-1}(\frac{b}{a}) + \tilde{A}_0(\frac{b}{a})y + \tilde{A}_1(\frac{b}{a})y^2)$$

where  $\tilde{A}_i = xA_i \in \mathbb{Q}[x]$ . Note that  $\tilde{A}_1(x)$  is nonzero because  $K(x, y, t)$  has bidegree  $(2, 2)$ . Equating the  $y^2$ -terms we find that  $\tilde{A}_1(\frac{b}{a}) = 0$  so  $\frac{b}{a}$  is algebraic over  $\mathbb{Q}(d_{i,j})$ . Equating the  $y$ -terms, we obtain that  $\frac{b}{a} - t\tilde{A}_0(\frac{b}{a}) = 0$ . Using the fact that  $t$  is transcendental over  $\mathbb{Q}(d_{i,j})$  and  $\frac{b}{a}$  is algebraic over  $\mathbb{Q}(d_{i,j})$ , we deduce  $\frac{b}{a} = 0$ . Therefore  $b = 0$ . This contradicts the fact that  $K$  has  $x$ -valuation 0. A similar argument shows that  $f_2(x, y)$  is irreducible.  $\square$

**Lemma 1.5.** *Let  $\bar{f}_i(x, y)$  denote the polynomial whose coefficients are the complex conjugates of those of  $f_i(x, y)$ . We may reduce to the case where one of the following cases hold:*

- there exists  $\epsilon \in \{\pm 1\}$  such that  $\bar{f}_1(x, y) = \epsilon f_2(x, y)$ ,
- $\bar{f}_1(x, y) = f_1(x, y) \in \mathbb{R}[x, y]$  and  $\bar{f}_2(x, y) = f_2(x, y) \in \mathbb{R}[x, y]$ .

*Proof of Lemma 1.5.* Unique factorization of polynomials implies that since  $-K(x, y, t) = f_1(x, y)f_2(x, y) = \bar{f}_1(x, y)\bar{f}_2(x, y)$ , there exists  $\lambda \in \mathbb{C}^*$  such that

- either  $\bar{f}_1(x, y) = \lambda f_2(x, y)$  and  $\bar{f}_2(x, y) = \lambda^{-1} f_1(x, y)$ ;
- or  $\bar{f}_1(x, y) = \lambda f_1(x, y)$  and  $\bar{f}_2(x, y) = \lambda^{-1} f_2(x, y)$ .

In the former case, we have  $f_1(x, y) = \bar{\lambda} \bar{f}_2(x, y) = \bar{\lambda} \lambda^{-1} f_1(x, y)$  and so  $\bar{\lambda} \lambda^{-1} = 1$ . This implies that  $\lambda$  is real and replacing  $\bar{f}_1(x, y)$  by  $|\lambda|^{-1/2} f_1(x, y)$  and  $f_2(x, y)$  by  $|\lambda|^{1/2} f_2(x, y)$ , we can assume that either  $\bar{f}_1(x, y) = f_2(x, y)$  and  $\bar{f}_2(x, y) = f_1(x, y)$  or  $\bar{f}_1(x, y) = -f_2(x, y)$  and  $\bar{f}_2(x, y) = -f_1(x, y)$ .

A similar computation in the latter case shows that  $|\lambda| = 1$ . Letting  $\mu$  be a square root of  $\lambda$  we have  $\mu^{-1} = \bar{\mu}$  so  $\lambda = \mu/\bar{\mu}$ . Replacing  $\bar{f}_1(x, y)$  by  $\mu f_1(x, y)$  and  $f_2(x, y)$  by  $\bar{\mu} f_2(x, y)$ , we can assume that  $\bar{f}_1(x, y) = f_1(x, y)$  and  $\bar{f}_2(x, y) = f_2(x, y)$ .  $\square$

Let us continue the proof of Proposition 1.2. For  $i = 1, 2$ , let us write

$$f_i(x, y) = (\alpha_{i,4}x + \alpha_{i,3})y + (\alpha_{i,2}x + \alpha_{i,1}),$$

with  $\alpha_{i,j} \in \mathbb{C}$ . Equating the terms in  $x^i y^j$  with  $-1 \leq i, j \leq 1$ , in  $f_1(x, y)f_2(x, y) = -K(x, y, t)$ , we find (recall that  $d_{i,j} \in [0, 1]$ ,  $t \in ]0, 1[$ )

term	coefficient in $f_1(x, y)f_2(x, y)$	coefficient in $-K(x, y, t)$
1	$\alpha_{1,1}\alpha_{2,1}$	$d_{-1,-1}t \geq 0$
$x$	$\alpha_{1,2}\alpha_{2,1} + \alpha_{1,1}\alpha_{2,2}$	$d_{0,-1}t \geq 0$
$x^2$	$\alpha_{1,2}\alpha_{2,2}$	$d_{1,-1}t \geq 0$
$y$	$\alpha_{1,3}\alpha_{2,1} + \alpha_{1,1}\alpha_{2,3}$	$d_{-1,0}t \geq 0$
$xy$	$\alpha_{1,4}\alpha_{2,1} + \alpha_{1,3}\alpha_{2,2} + \alpha_{1,2}\alpha_{2,3} + \alpha_{1,1}\alpha_{2,4}$	$d_{0,0}t - 1 < 0$
$x^2y$	$\alpha_{1,4}\alpha_{2,2} + \alpha_{1,2}\alpha_{2,4}$	$d_{1,0}t \geq 0$
$y^2$	$\alpha_{1,3}\alpha_{2,3}$	$d_{-1,1}t \geq 0$
$xy^2$	$\alpha_{1,4}\alpha_{2,3} + \alpha_{1,3}\alpha_{2,4}$	$d_{0,1}t \geq 0$
$x^2y^2$	$\alpha_{1,4}\alpha_{2,4}$	$d_{1,1}t \geq 0$

Let us treat separately two cases.

**Case 1:**  $f_1(x, y), f_2(x, y) \notin \mathbb{R}[x, y]$ . So, in this case we have either  $\overline{f_1}(x, y) = f_2(x, y)$  or  $\overline{f_1}(x, y) = -f_2(x, y)$ .

Let us first assume that  $\overline{f_1}(x, y) = f_2(x, y)$ . Then, evaluating the equality  $K(x, y, t) = -f_1(x, y)f_2(x, y)$  at  $x = y = 1$ , we get the following equality  $K(1, 1, t) = -f_1(1, 1)f_2(1, 1) = -|f_1(1, 1)|^2$ . But this is impossible because the left-hand term  $K(1, 1, t) = 1 - t \sum_{i,j \in \{-1, 0, 1\}^2} d_{i,j} = 1 - t$  is  $> 0$  whereas the right-hand term  $-|f_1(1, 1)|^2$  is  $\leq 0$ .

Let us now assume that  $\overline{f_1}(x, y) = -f_2(x, y)$ . Equating the constant terms in the equality  $f_1(x, y)f_2(x, y) = -K(x, y, t)$ , we get  $-|\alpha_{1,1}|^2 = d_{-1,-1}t$ , so  $\alpha_{1,1} = \alpha_{2,1} = d_{-1,-1} = 0$ . Equating the coefficients of  $x^2$  in the equality  $f_1(x, y)f_2(x, y) = -K(x, y, t)$ , we get  $-|\alpha_{1,2}|^2 = d_{1,-1}t$ , so  $\alpha_{1,2} = \alpha_{2,2} = d_{1,-1} = 0$ . It follows that the  $y$ -valuation of  $f_1(x, y)f_2(x, y) = -K(x, y, t)$  is  $\geq 2$ , whence a contradiction.

**Case 2:**  $f_1(x, y), f_2(x, y) \in \mathbb{R}[x, y]$ . We first claim that, after possibly replacing  $f_1(x, y)$  by  $-f_1(x, y)$  and  $f_2(x, y)$  by  $-f_2(x, y)$ , we may assume that  $\alpha_{1,4}, \alpha_{2,4}, \alpha_{1,3}, \alpha_{2,3} \geq 0$ .

Let us first assume that  $\alpha_{1,4}\alpha_{2,4} \neq 0$ . Since  $\alpha_{1,4}\alpha_{2,4} = d_{1,1}t \geq 0$ , we find that  $\alpha_{1,4}, \alpha_{2,4}$  belong simultaneously to  $\mathbb{R}_{>0}$  or  $\mathbb{R}_{<0}$ . After possibly replacing  $f_1(x, y)$  by  $-f_1(x, y)$  and  $f_2(x, y)$  by  $-f_2(x, y)$ , we may assume that  $\alpha_{1,4}, \alpha_{2,4} > 0$ . Since  $\alpha_{1,3}\alpha_{2,3} = d_{-1,1}t \geq 0$ , we have that  $\alpha_{1,3}, \alpha_{2,3}$  belong simultaneously to  $\mathbb{R}_{\geq 0}$  or  $\mathbb{R}_{\leq 0}$ . Then, the equality  $\alpha_{1,4}\alpha_{2,3} + \alpha_{1,3}\alpha_{2,4} = d_{0,1}t \geq 0$  implies that  $\alpha_{1,3}, \alpha_{2,3} \geq 0$ .

We can argue similarly in the case  $\alpha_{1,3}\alpha_{2,3} \neq 0$ .

It remains to consider the case  $\alpha_{1,4}\alpha_{2,4} = \alpha_{1,3}\alpha_{2,3} = 0$ . After possibly replacing  $f_1(x, y)$  by  $-f_1(x, y)$  and  $f_2(x, y)$  by  $-f_2(x, y)$ , we may assume that  $\alpha_{1,4}, \alpha_{2,4} \geq 0$ . The case  $\alpha_{1,4} = \alpha_{1,3} = 0$  is impossible because, otherwise, we would have  $d_{1,1} = d_{-1,1} = d_{0,1} = 0$ , which is excluded. Similarly, the case  $\alpha_{2,4} = \alpha_{2,3} = 0$  is impossible. So, we are left with the cases  $\alpha_{1,4} = \alpha_{2,3} = 0$  or  $\alpha_{2,4} = \alpha_{1,3} = 0$ . In both cases, the equality  $\alpha_{1,4}\alpha_{2,3} + \alpha_{1,3}\alpha_{2,4} = d_{0,1}t \geq 0$  with  $\alpha_{1,4}, \alpha_{2,4} \geq 0$ , implies that  $\alpha_{1,4}, \alpha_{2,4}, \alpha_{1,3}, \alpha_{2,3} \geq 0$ .

Arguing as above, we see that  $\alpha_{1,2}, \alpha_{2,2}, \alpha_{1,1}, \alpha_{2,1}$  all belong to  $\mathbb{R}_{\geq 0}$  or all belong to  $\mathbb{R}_{\leq 0}$ . Using the equation of the  $xy$ -coefficients, we find that  $\alpha_{1,2}, \alpha_{2,2}, \alpha_{1,1}, \alpha_{2,1}$  are all in  $\mathbb{R}_{\leq 0}$ .



Now, equating the coefficients of  $x^2y$  in the equality  $f_1(x, y)f_2(x, y) = -K(x, y, t)$  we get  $\alpha_{1,4}\alpha_{2,2} + \alpha_{1,2}\alpha_{2,4} = d_{1,0}t$ . Using the fact that  $\alpha_{1,4}\alpha_{2,2}, \alpha_{1,2}\alpha_{2,4} \leq 0$  and that  $d_{1,0}t \geq 0$ , we get  $\alpha_{1,4}\alpha_{2,2} = \alpha_{1,2}\alpha_{2,4} = d_{1,0}t = 0$ . Similarly, using the coefficients of  $y$ , we get  $\alpha_{1,3}\alpha_{2,1} = \alpha_{1,1}\alpha_{2,3} = d_{-1,0} = 0$ .

So, we have

$$\alpha_{1,4}\alpha_{2,2} = \alpha_{1,2}\alpha_{2,4} = \alpha_{1,3}\alpha_{2,1} = \alpha_{1,1}\alpha_{2,3} = 0.$$

The fact that  $K(x, y, t)$  has  $x$ - and  $y$ -degree 2 and  $x$ - and  $y$ -valuation 0 implies that, for any  $i \in \{1, 2\}$ , none of the vectors  $(\alpha_{i,4}, \alpha_{i,3}), (\alpha_{i,2}, \alpha_{i,1}), (\alpha_{i,4}, \alpha_{i,2})$  and  $(\alpha_{i,3}, \alpha_{i,1})$  is  $(0, 0)$ . Since  $\alpha_{1,4}\alpha_{2,2} = 0$ , we have  $\alpha_{1,4} = 0$  or  $\alpha_{2,2} = 0$ . If  $\alpha_{1,4} = 0$ , from what precedes, we find

$$\alpha_{1,4} = \alpha_{2,4} = \alpha_{2,1} = \alpha_{1,1} = 0.$$

If  $\alpha_{2,2} = 0$  we obtain

$$\alpha_{2,2} = \alpha_{1,2} = \alpha_{1,3} = \alpha_{2,3} = 0.$$

In the first case, the model belongs to the family of models  $\left\{ \begin{array}{l} \swarrow, \searrow \\ \nearrow, \nwarrow \end{array} \right\}$ . In the second case, we find that the model belongs to the family of models  $\left\{ \begin{array}{l} \swarrow, \nwarrow \\ \nearrow, \searrow \end{array} \right\}$ . This completes the proof.  $\square$

*Remark 1.6.* The fact  $d_{i,j} \in [0, 1]$  are probabilities is crucial in the proof of Proposition 1.2. We do not expect this result to be correct for general  $d_{i,j} \in \mathbb{C}$ .

*Remark 1.7.* The “degenerate models of walks” are called “singular” by certain authors, e.g., in [FIM99, FIM17]. Note also that, in [KR12], “singular walks” has a different meaning and refers to *models of walks* such that the associated *kernel* defines a genus zero curve.

*Remark 1.8.* In [DR19, Proposition 3], the authors show that Proposition 1.2 extends mutatis mutandis to the case when  $t \in ]0, 1[$  is algebraic over  $\mathbb{Q}(d_{i,j})$ . Their proof relies on Proposition 1.2 and uses a continuity argument of the parameter  $t$  to deduce that Proposition 1.2 stays correct for general values of  $t \in ]0, 1[$ .

From now on, we will only consider *nondegenerate models of walks*. In terms of *models of walks*, this only discards one dimensional problems and *models of walks* in the half-plane restricted to the quarter plane that are easier to study, as explained in [BMM10, Section 2.1].

## 2. SINGULARITIES AND GENUS OF THE KERNEL CURVE

The *kernel curve*  $E_t$  is the complex affine algebraic curve defined by

$$E_t = \{(x, y) \in \mathbb{C} \times \mathbb{C} \mid K(x, y, t) = 0\}.$$

We shall now consider a compactification of this curve. We let  $\mathbb{P}^1(\mathbb{C})$  be the complex projective line, which is the quotient of  $(\mathbb{C} \times \mathbb{C}) \setminus \{(0, 0)\}$  by the equivalence relation  $\sim$  defined by

$$(x_0, x_1) \sim (x'_0, x'_1) \Leftrightarrow \exists \lambda \in \mathbb{C}^*, (x'_0, x'_1) = \lambda(x_0, x_1).$$

The equivalence class of  $(x_0, x_1) \in (\mathbb{C} \times \mathbb{C}) \setminus \{(0, 0)\}$  is denoted by  $[x_0 : x_1] \in \mathbb{P}^1(\mathbb{C})$ . The map  $x \mapsto [x : 1]$  embeds  $\mathbb{C}$  inside  $\mathbb{P}^1(\mathbb{C})$ . The latter map is not surjective: its image

is  $\mathbb{P}^1(\mathbb{C}) \setminus \{[1 : 0]\}$ ; the missing point  $[1 : 0]$  is usually denoted by  $\infty$ . Now, we embed  $E_t$  inside  $\mathbb{P}^1(\mathbb{C}) \times \mathbb{P}^1(\mathbb{C})$  via  $(x, y) \mapsto ([x : 1], [y : 1])$ . The *kernel curve*  $\overline{E}_t$  is the closure of this embedding of  $E_t$ . In other words, the *kernel curve*  $\overline{E}_t$  is the algebraic curve defined by

$$\overline{E}_t = \{([x_0 : x_1], [y_0 : y_1]) \in \mathbb{P}^1(\mathbb{C}) \times \mathbb{P}^1(\mathbb{C}) \mid \overline{K}(x_0, x_1, y_0, y_1, t) = 0\}$$

where  $\overline{K}(x_0, x_1, y_0, y_1, t)$  is the following bihomogeneous polynomial

$$(2.1) \quad \overline{K}(x_0, x_1, y_0, y_1, t) = x_1^2 y_1^2 K\left(\frac{x_0}{x_1}, \frac{y_0}{y_1}, t\right) = x_0 x_1 y_0 y_1 - t \sum_{i,j=0}^2 d_{i-1,j-1} x_0^i x_1^{2-i} y_0^j y_1^{2-j}.$$

Although it may seem more natural to take the closure of  $\overline{E}_t$  in  $\mathbb{P}^2(\mathbb{C})$ , the above definition allows us to extend the involutions of  $E_t$  of Section 3 in a natural way as well as allowing us to avoid unnecessary singularities.

We shall now study the singularities and compute the genus of  $\overline{E}_t$ . Recall that since the *model of walk* under consideration is *nondegenerate*, the polynomial  $K(x, y, t)$  is irreducible. We recall that by definition  $P = ([a : b], [c : d]) \in \overline{E}_t$  is called a singularity of the irreducible kernel  $\overline{E}_t$  if

$$\frac{\partial \overline{K}(a, b, c, d, t)}{\partial x_0} = \frac{\partial \overline{K}(a, b, c, d, t)}{\partial x_1} = \frac{\partial \overline{K}(a, b, c, d, t)}{\partial y_0} = \frac{\partial \overline{K}(a, b, c, d, t)}{\partial y_1} = 0.$$

Note that one can check this condition in any affine neighborhood of a point. For example, if  $b, d \neq 0$ , the bihomogeneity of  $\overline{K}$  implies

$$\begin{aligned} 0 = 2\overline{K}(a/b, 1, c/d, 1, t) &= \frac{a}{b} \frac{\partial \overline{K}(a/b, 1, c, /d, 1, t)}{\partial x_0} + \frac{\partial \overline{K}(a/b, 1, c, /d, 1, t)}{\partial x_1} \\ &= \frac{c}{d} \frac{\partial \overline{K}(a/b, 1, c, /d, 1, t)}{\partial y_0} + \frac{\partial \overline{K}(a/b, 1, c, /d, 1, t)}{\partial y_1}. \end{aligned}$$

Therefore the point  $P = ([a : b], [c : d]) \in \overline{E}_t$  is a singular point if and only if

$$\frac{\partial \overline{K}(a/b, 1, c/d, 1, t)}{\partial x_0} = \frac{\partial \overline{K}(a/b, 1, c/d, 1, t)}{\partial y_0} = 0.$$

If  $P = ([a : b], [c : d]) \in \overline{E}_t$  is not a singularity of  $\overline{E}_t$ , then it is called a smooth point of  $\overline{E}_t$ .

We also recall that  $\overline{E}_t$  is called singular if it has at least one singular point. Otherwise, we say that  $\overline{E}_t$  is nonsingular or smooth.

The genus of an algebraic curve is a classical notion in algebraic geometry. It is a nonnegative integer that we may attach to a curve, see [Ful84, Section 8.3] for a definition. The study of the genus of  $\overline{E}_t$  has been considered in [FIM17]. Proposition 2.1 below shows that the smoothness of  $\overline{E}_t$  is intimately related to the value of the genus of  $\overline{E}_t$ . We define the genus of the *weighted model of walk*, as the genus of its *kernel curve*  $\overline{E}_t$ .

Remind the following notations from the introduction :

$$\begin{aligned} K(x, y, t) &= xy - txA_{-1}(x) - txA_0(x)y - txA_1(x)y^2, \\ &= xy - tyB_{-1}(y) - tyB_0(y)x - tyB_1(y)x^2, \end{aligned}$$

where  $x A_i(x) \in \mathbb{Q}[x]$  and  $y B_i(y) \in \mathbb{Q}[y]$ . Then we may write

$$\begin{aligned} \overline{K}(x_0, x_1, y_0, y_1, t) &= \overline{C}_1(x_0, x_1, t) y_1^2 + \overline{B}_1(x_0, x_1, t) y_0 y_1 + \overline{A}_1(x_0, x_1, t) y_0^2 \\ &= \overline{C}_2(y_0, y_1, t) x_1^2 + \overline{B}_2(y_0, y_1, t) x_0 x_1 + \overline{A}_2(y_0, y_1, t) x_0^2. \end{aligned}$$

For any  $[x_0 : x_1]$  and  $[y_0 : y_1]$  in  $\mathbb{P}^1(\mathbb{C})$ , we denote by  $\Delta_1([x_0 : x_1])$  and  $\Delta_2([y_0 : y_1])$  the discriminants of the degree two homogeneous polynomials given by  $y \mapsto \overline{K}(x_0, x_1, y, t)$  and  $x \mapsto \overline{K}(x, y_0, y_1, t)$  respectively, i.e.

$$\begin{aligned} \Delta_1([x_0 : x_1]) &= \overline{B}_1(x_0, x_1, t)^2 - 4\overline{A}_1(x_0, x_1, t)\overline{C}_1(x_0, x_1, t) \\ &= (x_0 x_1 - t^2) \left( (d_{-1,0} x_1^2 - \frac{1}{t} x_0 x_1 + d_{0,0} x_0 x_1 + d_{1,0} x_0^2)^2 \right. \\ &\quad \left. - 4(d_{-1,1} x_1^2 + d_{0,1} x_0 x_1 + d_{1,1} x_0^2)(d_{-1,-1} x_1^2 + d_{0,-1} x_0 x_1 + d_{1,-1} x_0^2) \right) \end{aligned}$$

and

$$\begin{aligned} \Delta_2([y_0 : y_1]) &= \overline{B}_2(y_0, y_1, t)^2 - 4\overline{A}_2(y_0, y_1, t)\overline{C}_2(y_0, y_1, t) \\ &= t^2 \left( (d_{0,-1} y_1^2 - \frac{1}{t} y_0 y_1 + d_{0,0} y_0 y_1 + d_{0,1} y_0^2)^2 \right. \\ &\quad \left. - 4(d_{1,-1} y_1^2 + d_{1,0} y_0 y_1 + d_{1,1} y_0^2)(d_{-1,-1} y_1^2 + d_{-1,0} y_0 y_1 + d_{-1,1} y_0^2) \right). \end{aligned}$$

As we will see in the sequel, see Remark 2.3,  $\Delta_1([x_0 : x_1])$  has a double root if and only if  $\Delta_2([y_0 : y_1])$  has a double root.

**Proposition 2.1.** *For nondegenerate models, the following facts are equivalent:*

- (1) the curve  $\overline{E}_t$  is a genus zero curve;
- (2) the curve  $\overline{E}_t$  is singular;
- (3) the curve  $\overline{E}_t$  has exactly one singularity  $\Omega \in \overline{E}_t$ ;
- (4) there exists  $([a : b], [c : d]) \in \overline{E}_t$  such that the discriminants  $\Delta_1([x_0 : x_1])$  and  $\Delta_2([y_0 : y_1])$  have a root  $[a : b] \in \mathbb{P}^1(\mathbb{C})$  and  $[c : d] \in \mathbb{P}^1(\mathbb{C})$  respectively;
- (5) there exists  $([a : b], [c : d]) \in \overline{E}_t$  such that the discriminants  $\Delta_1([x_0 : x_1])$  and  $\Delta_2([y_0 : y_1])$  have a double root  $[a : b] \in \mathbb{P}^1(\mathbb{C})$  and  $[c : d] \in \mathbb{P}^1(\mathbb{C})$  respectively.

If these properties are satisfied, then the singular point is  $\Omega = ([a : b], [c : d])$  where  $[a : b] \in \mathbb{P}^1(\mathbb{C})$  is a double root of  $\Delta_1([x_0 : x_1])$  and  $[c : d] \in \mathbb{P}^1(\mathbb{C})$  is a double root of  $\Delta_2([y_0 : y_1])$ . If the previous properties are not satisfied, then  $\overline{E}_t$  is a smooth curve of genus one.

*Proof.* By [Dui10, Section 3.3.1], the following formula gives the genus of  $\overline{E}_t$ ,

$$(2.2) \quad g(\overline{E}_t) = 1 - \sum_{P \in \text{Sing}} \frac{m(P)(m(P) - 1)}{2},$$

where  $m(P)$  is a positive integer standing for the multiplicity of a point  $P$ , that is, some partial derivative of order  $m(P)$  does not vanish while for every  $\ell < m(P)$ , the partial derivatives of order  $\ell$  vanish at  $P$ . Since the genus is a nonnegative integer, the above formula shows that  $g(\overline{E}_t)$  is equal to 0 or 1. This formula shows more precisely that  $\overline{E}_t$  is smooth if and only if  $g(\overline{E}_t) = 1$ . Moreover (2.2) shows that if  $\overline{E}_t$  is singular, then there is exactly one singular point that is a double point, and the curve has genus

zero. This proves the equivalence between (1), (2) and (3), and the last statement of the proposition.

Let us prove (4)  $\Rightarrow$  (3). Assume that the discriminant  $\Delta_1([x_0 : x_1])$  (resp.  $\Delta_2([y_0 : y_1])$ ) has a root in  $[a : b] \in \mathbb{P}^1(\mathbb{C})$  (resp.  $[c : d] \in \mathbb{P}^1(\mathbb{C})$ ). Let us write

$$\begin{aligned} & \overline{K}(x_0, x_1, y_0, y_1, t) \\ = & e_{-1,1}(dy_0 - cy_1)^2 + e_{0,1}(bx_0 - ax_1)(dy_0 - cy_1)^2 + e_{1,1}(bx_0 - ax_1)^2(dy_0 - cy_1)^2 \\ + & e_{-1,0}(dy_0 - cy_1) + e_{0,0}(bx_0 - ax_1)(dy_0 - cy_1) + e_{1,0}(bx_0 - ax_1)^2(dy_0 - cy_1) \\ + & e_{-1,-1} + e_{0,-1}(bx_0 - ax_1) + e_{1,-1}(bx_0 - ax_1)^2. \end{aligned}$$

Since  $([a : b], [c : d]) \in \overline{E}_t$ , we have by definition that  $\overline{K}(a, b, c, d, t) = 0$ , i.e.  $e_{-1,-1} = 0$ . Since  $\Delta_1([x_0 : x_1])$  has a root in  $[a : b] \in \mathbb{P}^1(\mathbb{C})$ ,  $K(a, b, y_0, y_1)$  has a double root at  $[c, d]$  and so  $e_{-1,0} = 0$ . Similarly, the fact that  $\Delta_2([y_0 : y_1])$  has a root in  $[c : d] \in \mathbb{P}^1(\mathbb{C})$  implies  $e_{0,-1} = 0$ . This shows that

$$\frac{\partial \overline{K}(a, b, c, d, t)}{\partial x_0} = \frac{\partial \overline{K}(a, b, c, d, t)}{\partial x_1} = \frac{\partial \overline{K}(a, b, c, d, t)}{\partial y_0} = \frac{\partial \overline{K}(a, b, c, d, t)}{\partial y_1} = 0,$$


and, hence,  $([a : b], [c : d])$  is the singular point of  $\overline{E}_t$ .

Let us prove (3)  $\Rightarrow$  (5). If  $\Omega = ([a : b], [c : d])$  is the singular point of  $\overline{E}_t$ , then  $e_{-1,-1} = e_{-1,0} = e_{0,-1} = 0$ , and the discriminants  $\Delta_1([x_0 : x_1])$  and  $\Delta_2([y_0 : y_1])$  have a double root in  $[a : b] \in \mathbb{P}^1(\mathbb{C})$  and  $[c : d] \in \mathbb{P}^1(\mathbb{C})$  respectively.

The implication (5)  $\Rightarrow$  (4) is obvious. □

Our next aim is to describe the genus zero *models of walks*.

**Lemma 2.2.** *The discriminant  $\Delta_2([y_0 : y_1])$  has a double zero if and only if the model of the walk is included in a closed half plane whose boundary passes through  $(0, 0)$ . In other word, this correspond to models of the walks that belong to one of the following eight families*



*Remark 2.3.* As the statement of Lemma 2.2 is symmetric with respect to  $x$  and  $y$  we deduce that the same holds for  $\Delta_1([x_0 : x_1])$ . We then deduce that  $\Delta_1([x_0 : x_1])$  has a double root if and only if  $\Delta_2([y_0 : y_1])$  has a double root.

*Remark 2.4.* In the case  $t = 1$ , it is proved in [FIM17, Lemma 2.3.10] that, besides the models listed in Lemma 2.2, any *nondegenerate* model such that the drift is zero, i.e.

$$(\sum_i id_{i,j}, \sum_j jd_{i,j}) = (0, 0),$$

has a curve  $\overline{E}_t$  of genus 0.

*Proof.* The computations seem to be too complicated to be performed by hand, so we used MAPLE<sup>†</sup>. We are going to prove the result with two strategies. This first one is to write the discriminant of the discriminant  $\Delta_2([y_0 : y_1])$  and study when the latter is 0. The second strategy consists in decomposing the radical of an ideal into its prime

<sup>†</sup>The maple worksheet is available at [https://singer.math.ncsu.edu/ms\\_papers.html](https://singer.math.ncsu.edu/ms_papers.html).

components.

Let us first consider the situation where the double root is at  $(a, b)$  where  $b$  is not zero. Let us set  $y_1 = 1$  and  $y_0 = y$  to obtain the specialization  $\Delta_2([y : 1])$  of  $\Delta_2([y_0 : y_1])$ .

The following MAPLE code calculates the discriminant of the discriminant, its degree and order of vanishing in  $t$ , and then sets the coefficients of powers of  $t$  equal to zero. Solving these equations yields the 8 solutions  $\mathbf{S}[i]$ ,  $i = 1, \dots, 8$  corresponding to the 8 stepsets listed in Lemma 2.2.

```
> K := expand(x*y*(1-t*(add(add(d[i, j]*x^i*y^j, i = -1 .. 1), j = -1 .. 1))));
> DX := discrim(K, x);
> DD := discrim(discrim(K,x),y);
> ldegree(DD,t); degree(DD,t);
```

4  
12

```
> S := solve({seq(coeff(DD,t,i), i=4..12)}, [seq(seq(d[i, j], i=-1..1), j=-1..1)]);
> nops(S);
```

8

```
> S[1];S[2];S[3];S[4];S[5];S[6];S[7];S[8];
```

An alternate approach is to use the *PolynomialIdeals* package

```
> with(PolynomialIdeals):
```

and consider the prime decomposition of the radical of the ideal

```
> J := <seq(coeff(DD,t,i), i=4..12)>;
> PrimeDecomposition(J);
```

$$\begin{array}{lll} \langle d_{-1,-1}, d_{-1,0}, d_{-1,1} \rangle, & \langle d_{-1,-1}, d_{-1,0}, d_{0,-1} \rangle, & \langle d_{-1,-1}, d_{0,-1}, d_{1,-1} \rangle, \\ \langle d_{-1,0}, d_{-1,1}, d_{0,1} \rangle, & \langle d_{-1,1}, d_{0,1}, d_{1,1} \rangle, & \langle d_{0,-1}, d_{1,-1}, d_{1,0} \rangle, \\ \langle d_{0,1}, d_{1,0}, d_{1,1} \rangle, & \langle d_{1,-1}, d_{1,0}, d_{1,1} \rangle. & \end{array}$$

The *PrimeDecomposition* command lists a set of prime ideals whose intersection is the radical of the original ideal. In particular, these ideals have the property that any zero of the original ideal is a zero of one of the listed ideals and vice versa, see [CLO97, Chapter 4, Section 6]. These again correspond to the eight step sets listed in Lemma 2.2.

We now consider the case where the double root is at  $(a, b)$  where  $b = 0$ , that is, at  $(1, 0)$ . We will show that this case leads to *models of walks* already mentioned above.

```
> DDX := expand(z^4*subs(y = 1/z, DX));
```

If  $z = 0$  is a double root then the coefficient of 1 and  $z$  must be zero

```
> coeff(DDX, z, 0); coeff(DDX, z, 1);
```

$$-4t^2d_{-1,1}d_{1,1} + t^2d_{0,1}^2$$

$$-4t^2d_{-1,0}d_{1,1} - 4t^2d_{-1,1}d_{1,0} + 2t^2d_{0,0}d_{0,1} - 2td_{0,1}$$

Taking into account that  $t$  is transcendental over  $\mathbb{Q}(d_{i,j})$ , we are led to three cases, corresponding to three of the step sets in Lemma 2.2.

$$[d_{0,1} = 0, d_{-1,1} = 0, d_{-1,0} = 0]$$


$$[d_{0,1} = 0, d_{-1,1} = 0, d_{1,1} = 0]$$

$$[d_{0,1} = 0, d_{1,1} = 0, d_{1,0} = 0] \quad \square$$

*Remark 2.5.* The proof of Proposition 1.2 proceeds by a direct “hand calculation” while the proof of Lemma 2.2 follows from a simple MAPLE calculation. It would be interesting to have a simple MAPLE based proof of Proposition 1.2 and a hand calculation proof of Lemma 2.2.

**Corollary 2.6.** *The following holds:*

- (1) *The nondegenerate genus zero models of walks are the nondegenerate models whose step set is included in an half space whose boundary passes through  $(0, 0)$ . More precisely, they are nondegenerate models belonging to one of the following families*



- (2) *The nondegenerate genus one models of walks are the models whose step set is not included in any half space whose boundary passes through  $(0, 0)$ .*

*Remark 2.7.* See also [DR19, Proposition 9] for an extension of Corollary 2.6 to the case when  $t \in ]0, 1[$  is algebraic over  $\mathbb{Q}(d_{i,j})$ . Their proof relies on the results of the present section and uses a continuity argument with respect to the parameter  $t$  to deduce that Corollary 2.6 stays correct for general values of  $t \in ]0, 1[$ .

*Proof.* We use Proposition 2.1. We have to determine when there exists  $([a : b], [c : d]) \in \overline{E}_t$  such that the discriminants  $\Delta_1([x_0 : x_1])$  and  $\Delta_2([y_0 : y_1])$  have a double root  $[a : b] \in \mathbb{P}^1(\mathbb{C})$  and  $[c : d] \in \mathbb{P}^1(\mathbb{C})$ . Lemma 2.2 provides such configurations. By Proposition 1.2, the configurations number 1, 3, 5, 7 are dismissed since they led to *singular walks*. Then, if we are considering *nondegenerate genus zero models of walks*, we are in the four families of models considered in (G0). Furthermore, if the step set is not included in any half space whose boundary passes through  $(0, 0)$ , the configurations of Proposition 1.2 and Lemma 2.2 are excluded and by Proposition 2.1, we are in the genus 1 situation.

Conversely, it remains to prove that if the *models of walks* are in the four families of *models* considered in (G0), the kernel has genus 0. Thanks to Proposition 2.1 it suffices to prove that the discriminants have a common zero in that case. This is the goal of the following Lemma 2.8 and Remark 2.9. □

Let us write


$$\Delta_1([x_0 : x_1]) = \sum_{i=0}^4 \alpha_i(t)x_0^i x_1^{4-i}, \quad \text{and} \quad \Delta_2([y_0 : y_1]) = \sum_{i=0}^4 \beta_i(t)y_0^i y_1^{4-i}.$$

where

$$\begin{aligned}
\alpha_0(t) &= t^2 d_{-1,0}^2 - 4t^2 d_{-1,1} d_{-1,-1} \\
\alpha_1(t) &= 2t^2 d_{-1,0} d_{0,0} - 2t d_{-1,0} - 4t^2 d_{-1,1} d_{0,-1} - 4t^2 d_{0,1} d_{-1,-1} \\
\alpha_2(t) &= t^2 d_{0,0}^2 - 2t d_{0,0} + 1 + 2t^2 d_{-1,0} d_{1,0} - 4t^2 d_{-1,1} d_{1,-1} - 4t^2 d_{0,1} d_{0,-1} - 4t^2 d_{1,1} d_{-1,-1} \\
\alpha_3(t) &= -2t d_{1,0} + 2t^2 d_{0,0} d_{1,0} - 4t^2 d_{1,1} d_{0,-1} - 4t^2 d_{0,1} d_{1,-1} \\
\alpha_4(t) &= t^2 d_{1,0}^2 - 4t^2 d_{1,1} d_{1,-1} \\
\beta_0(t) &= t^2 d_{0,-1}^2 - 4t^2 d_{1,-1} d_{-1,-1} \\
\beta_1(t) &= 2t^2 d_{0,-1} d_{0,0} - 2t d_{0,-1} - 4t^2 d_{1,-1} d_{-1,0} - 4t^2 d_{1,0} d_{-1,-1} \\
\beta_2(t) &= t^2 d_{0,0}^2 - 2t d_{0,0} + 1 + 2t^2 d_{0,-1} d_{0,1} - 4t^2 d_{1,-1} d_{-1,1} - 4t^2 d_{1,0} d_{-1,0} - 4t^2 d_{1,1} d_{-1,-1} \\
\beta_3(t) &= -2t d_{0,1} + 2t^2 d_{0,0} d_{0,1} - 4t^2 d_{1,1} d_{-1,0} - 4t^2 d_{1,0} d_{-1,1} \\
\beta_4(t) &= t^2 d_{0,1}^2 - 4t^2 d_{1,1} d_{-1,1}.
\end{aligned}$$

Note that  $\Delta_1([x_0 : x_1])$  (resp.  $\Delta_2([y_0 : y_1])$ ) is of degree 4 and so has four roots counted with multiplicities  $a_1, a_2, a_3, a_4$  (resp.  $b_1, b_2, b_3, b_4$ ) in  $\mathbb{P}^1(\mathbb{C})$ . If the discriminant  $\Delta_1([x_0 : x_1])$  (resp.  $\Delta_2([y_0 : y_1])$ ) has a double root; up to renumbering, we can assume that  $a_1 = a_2$  (resp.  $b_1 = b_2$ ).

**Lemma 2.8.** *Assume that the model of the walk is nondegenerate and belongs to the first family of (G0)*



Then, the walk has genus zero and the singular point of  $\overline{E}_t$  is  $\Omega = ([0 : 1], [0 : 1])$ , that is,  $a_1 = a_2 = [0 : 1]$  (resp.  $b_1 = b_2 = [0 : 1]$ ) is a double root of  $\Delta_1([x_0 : x_1])$  (resp.  $\Delta_2([y_0 : y_1])$ ). The other roots are distinct from one another and from the double root and are given by

	$a_1 = a_2$	$b_1 = b_2$
	$[0 : 1]$	$[0 : 1]$
	$a_3$	$a_4$
$\alpha_4(t) \neq 0$	$[-\alpha_3(t) - \sqrt{\alpha_3(t)^2 - 4\alpha_2(t)\alpha_4(t)} : 2\alpha_4(t)]$	$[-\alpha_3(t) + \sqrt{\alpha_3(t)^2 - 4\alpha_2(t)\alpha_4(t)} : 2\alpha_4(t)]$
$\alpha_4(t) = 0$	$[1 : 0]$	$[-\alpha_2(t) : \alpha_3(t)]$
	$b_3$	$b_4$
$\beta_4(t) \neq 0$	$[-\beta_3(t) - \sqrt{\beta_3(t)^2 - 4\beta_2(t)\beta_4(t)} : 2\beta_4(t)]$	$[-\beta_3(t) + \sqrt{\beta_3(t)^2 - 4\beta_2(t)\beta_4(t)} : 2\beta_4(t)]$
$\beta_4(t) = 0$	$[1 : 0]$	$[-\beta_2(t) : \beta_3(t)]$

*Remark 2.9.* We can extend Lemma 2.8 to the other configurations in (G0) by using the following remarks:

- (1) Replacing  $([x_0 : x_1], [y_0 : y_1])$  by  $([x_0 : x_1], [y_1 : y_0])$ , which corresponds to the variable change  $(x, y) \mapsto (x, y^{-1})$ , amounts to consider a *weighted model of walk* with *weights*  $d'_{i,j} := d_{i,-j}$ . This can be used to extend Lemma 2.8 to the second configuration of (G0); for instance, the singular point of  $\overline{E}_t$  is  $\Omega = ([0 : 1], [1 : 0])$  in that case.
- (2) Replacing  $([x_0 : x_1], [y_0 : y_1])$  by  $([x_1 : x_0], [y_1 : y_0])$  amounts to consider a *weighted model of walk* with *weights*  $d'_{i,j} := d_{-i,-j}$ . This can be used to extend Lemma 2.8 to the third configuration of (G0); for instance, the singular point of  $\overline{E}_t$  is  $\Omega = ([1 : 0], [1 : 0])$  in that case.
- (3) Replacing  $([x_0 : x_1], [y_0 : y_1])$  by  $([x_1 : x_0], [y_0 : y_1])$  amounts to consider a *weighted model of walk* with *weights*  $d'_{i,j} := d_{-i,j}$ . This can be used to extend Lemma 2.8 to the fourth configuration of (G0); for instance, the singular point of  $\overline{E}_t$  is  $\Omega = ([1 : 0], [0 : 1])$  in that case.

*Remark 2.10.* Note that if we consider the  $x_3, x_4$  (resp.  $y_3, y_4$ ) defined in [FIM17, Chapter 6], we have the equality of sets  $\{a_3, a_4\} = \{x_3, x_4\}$  and  $\{b_3, b_4\} = \{y_3, y_4\}$ , but do not have necessarily  $a_i = x_i, b_j = y_j$ , with  $3 \leq i, j \leq 4$ .

*Proof of Lemma 2.8.* We shall prove the lemma for  $\Delta_2([y_0 : y_1])$ , the proof for  $\Delta_1([x_0 : x_1])$  being similar. By assumption,  $d_{-1,-1} = d_{-1,0} = d_{0,-1} = 0$ . Then,  $\alpha_0(t) = \alpha_1(t) = 0$ . Therefore, the discriminant  $\Delta_2([y_0 : y_1])$  has a double root at  $[0 : 1]$  and we can write

$$\Delta_2([y : 1]) = \beta_2(t)y^2 + \beta_3(t)y^3 + \beta_4(t)y^4.$$

Since  $t$  is transcendental over  $\mathbb{Q}(d_{i,j})$ , we see that the coefficient of  $y^2$  is nonzero. Therefore  $[0 : 1]$  is precisely a double root of  $\Delta_2([y_0 : y_1])$ . To see that  $b_3$  and  $b_4$  are distinct, we calculate the discriminant of  $\Delta_2([y : 1])/y^2$ , which is almost the same as the one we considered in the proof of Lemma 2.2. This is a polynomial of degree four in  $t$  with the following coefficients:

term	coefficient
$t^4$	$-16(4d_{-1,1}d_{1,-1}d_{1,1} - d_{1,-1}d_{1,0}^2 - d_{0,0}^2d_{1,1} + d_{0,0}d_{0,1}d_{1,0} - d_{0,1}^2d_{1,-1})d_{-1,1}$
$t^3$	$-16(2d_{0,0}d_{1,1} - d_{0,1}d_{1,0})d_{-1,1}$
$t^2$	$16d_{-1,1}d_{1,1}$
$t$	$0$
$1$	$0$

Let us prove that if  $\Delta_2([y_0 : y_1])$  has a double root different from  $[0 : 1]$ , all the above coefficients must be zero. Recalling that  $d_{-1,1}d_{1,-1} \neq 0$ , from the coefficient of  $t^2$ , we must have  $d_{1,1} = 0$ . From the coefficient of  $t^3$ , we have that  $d_{0,1} = 0$  or  $d_{1,0} = 0$ . From the coefficient of  $t^4$ , we get in both cases  $d_{0,1} = d_{1,0} = 0$ . This implies that the model of the *walk* would be *degenerate*, a contradiction. The formulas for  $b_3$  and  $b_4$  follow from the quadratic formula.  $\square$

### 3. INVOLUTIVE AUTOMORPHISMS OF THE KERNEL CURVE

Following [BMM10, Section 3], [KY15, Section 3] or [FIM17], we consider the involutive rational functions

$$i_1, i_2 : \mathbb{C}^2 \dashrightarrow \mathbb{C}^2$$



given by

$$i_1(x, y) = \left( x, \frac{A_{-1}(x)}{A_1(x)y} \right) \text{ and } i_2(x, y) = \left( \frac{B_{-1}(y)}{B_1(y)x}, y \right).$$

Note that  $i_1, i_2$  are “only” rational functions in the sense that they are a priori not defined when the denominators vanish. The dashed arrow notation used above and in the rest of the paper is a classical notation for rational functions.

The rational functions  $i_1, i_2$  induce involutive rational functions

$$\iota_1, \iota_2 : \overline{E}_t \dashrightarrow \overline{E}_t$$

given by

$$\begin{aligned} \iota_1([x_0 : x_1], [y_0 : y_1]) &= \left( [x_0 : x_1], \left[ \frac{A_{-1}\left(\frac{x_0}{x_1}\right)}{A_1\left(\frac{x_0}{x_1}\right)\frac{y_0}{y_1}} : 1 \right] \right), \\ \text{and } \iota_2([x_0 : x_1], [y_0 : y_1]) &= \left( \left[ \frac{B_{-1}\left(\frac{y_0}{y_1}\right)}{B_1\left(\frac{y_0}{y_1}\right)\frac{x_0}{x_1}} : 1 \right], [y_0 : y_1] \right). \end{aligned}$$


Again, these functions are a priori not defined where the denominators vanish. However, the following result shows that, actually, this is only an “apparent problem”:  $\iota_1$  and  $\iota_2$  can be extended into endomorphisms of  $\overline{E}_t$ . We recall that a rational map  $f : \overline{E}_t \dashrightarrow \overline{E}_t$  is an endomorphism if it is regular at any  $P \in \overline{E}_t$ , i.e. if  $f$  can be represented in suitable affine charts containing  $P$  and  $f(P)$  by a rational function with nonvanishing denominator at  $P$ . More generally, given  $X$  and  $Y$  algebraic varieties, we say that  $f : X \dashrightarrow Y$  is a morphism if  $f$  can be represented by two suitable affine charts containing  $P$  and  $f(P)$  respectively, by a rational function with nonvanishing denominator at  $P$ .

**Proposition 3.1.** *The rational maps  $\iota_1, \iota_2 : \overline{E}_t \dashrightarrow \overline{E}_t$  can be extended into involutive automorphisms of  $\overline{E}_t$ .*

*Proof.* Note that  $\iota_1(x, y)$  is well-defined if the  $x_i$  and the  $y_j$  are nonzero and if  $A_1\left(\frac{x_0}{x_1}\right)\frac{y_0}{y_1}$  is nonzero. This excludes at most finitely many  $(x, y) \in \overline{E}_t$  and, hence, there exists a finite set  $\mathcal{S}_0 \subset \overline{E}_t$  such that  $\iota_1$  is well defined on  $\overline{E}_t \setminus \mathcal{S}_0$ . The map  $\iota_1$  induces a bijection from  $\overline{E}_t \setminus \mathcal{S}_0$  to  $\overline{E}_t \setminus \mathcal{S}_1$ , where  $\mathcal{S}_1$  is a finite set. The same holds for  $\iota_2$ . We have to prove that  $\iota_1, \iota_2 : \overline{E}_t \dashrightarrow \overline{E}_t$  can be extended into endomorphisms of  $\overline{E}_t$ . According to Proposition 2.1, if the curve  $\overline{E}_t$  has genus one, then it is smooth and the result follows from [Har77, Proposition 6.8, p. 43].

It remains to study the case when  $\overline{E}_t$  has genus zero. In that case, Proposition 2.1 ensures that  $\overline{E}_t$  has a unique singularity  $\Omega$ . It follows from [Har77, Proposition 6.8, p. 43] that  $\iota_1$  and  $\iota_2$  can be uniquely extended into morphisms  $\overline{E}_t \setminus \{\Omega\} \rightarrow \overline{E}_t$  still denoted by  $\iota_1$  and  $\iota_2$ . It remains to study  $\iota_1$  and  $\iota_2$  at  $\Omega$ . Let us first assume that the walk under consideration belongs to the family of the first configuration of (G0). Lemma 2.8 ensures that  $\Omega = ([0 : 1], [0 : 1])$ . For  $([x : 1], [y : 1]) \in \overline{E}_t$ , the equation  $K(x, y, t) = 0$  implies that

$$(3.1) \quad \frac{A_{-1}(x)}{A_1(x)y} = \frac{1}{tA_1(x)} - \frac{A_0(x)}{A_1(x)} - y = \frac{x}{t\tilde{A}_1(x)} - \frac{\tilde{A}_0(x)}{\tilde{A}_1(x)} - y$$

FIGURE 1. The maps  $\iota_1, \iota_2$  restricted to the kernel curve  $\overline{E}_t$ 

where  $\tilde{A}_0(x) = xA_0(x) = d_{-1,0} + d_{0,0}x + d_{1,0}x^2$  and  $\tilde{A}_1(x) = xA_1(x) = d_{-1,1} + d_{0,1}x + d_{1,1}x^2$ . Since  $d_{-1,1} \neq 0$ ,  $\tilde{A}_1(x)$  does not vanish at  $x = 0$ . Since  $d_{-1,0} = 0$ ,  $\tilde{A}_0(x)$  vanishes at  $x = 0$ . So, (3.1) shows that  $\iota_1$  is regular at  $\Omega$  and that  $\iota_1(\Omega) = \Omega$ . The argument for  $\iota_2$  is similar.

The other cases listed in (G0) can be treated similarly using a reduction argument as in Remark 2.9.  $\square$

We also consider the automorphism of  $\overline{E}_t$  defined by

$$\sigma = \iota_2 \circ \iota_1.$$

It is easily seen that  $\iota_1$  and  $\iota_2$  are the vertical and horizontal switches of  $\overline{E}_t$  (see Figure 1), i.e. for any  $P = (x, y) \in \overline{E}_t$ , we have

$$\{P, \iota_1(P)\} = \overline{E}_t \cap (\{x\} \times \mathbb{P}^1(\mathbb{C})) \text{ and } \{P, \iota_2(P)\} = \overline{E}_t \cap (\mathbb{P}^1(\mathbb{C}) \times \{y\}).$$

We now give a couple of lemmas for later use.

**Lemma 3.2.** *A point  $P = ([x_0 : x_1], [y_0 : y_1]) \in \overline{E}_t$  is fixed by  $\iota_1$  if and only if  $\Delta_1([x_0 : x_1]) = 0$ . A point  $P = ([x_0 : x_1], [y_0 : y_1]) \in \overline{E}_t$  is fixed by  $\iota_2$  if and only if  $\Delta_2([x_0 : x_1]) = 0$ .*

*Proof.* Assume that  $P$  is fixed by  $\iota_1$ . Then, the polynomial  $[y_0 : y_1] \mapsto \overline{K}(x_0, x_1, y_0, y_1, t)$  has a double root, meaning that the discriminant is zero. This is exactly  $\Delta_1([x_0 : x_1]) = 0$ . Conversely,  $\Delta_1([x_0 : x_1]) = 0$  implies that  $[y_0 : y_1] \mapsto \overline{K}(x_0, x_1, y_0, y_1, t)$  has a double root and therefore  $P$  is fixed by  $\iota_1$ . The proof for  $\iota_2$  is similar.  $\square$

The fixed points of  $\iota_1$  have  $y$ -coordinates that are the double roots of  $y \mapsto \overline{K}(x_0, x_1, y, t)$ , i.e. they are the roots of the discriminant. By Lemma 2.8 and Remark 2.9, there are 3 points of  $\overline{E}_t$  that are fixed by  $\iota_1$ . A similar statement holds for  $\iota_2$ . As is shown in the following lemma, one of them plays a particular role.

**Lemma 3.3.** *Let  $P \in \overline{E}_t$ . The following statements are equivalent:*

- (1)  $P$  is fixed by  $\iota_1$  and  $\iota_2$ ;
- (2)  $P$  is a singular point of  $\overline{E}_t$ ;
- (3)  $P$  is fixed by  $\sigma = \iota_2 \circ \iota_1$ .


*Proof.* Let  $P = ([a : b], [c : d]) \in \overline{E}_t$ . From Proposition 2.1, we have that  $P$  is a singular point if and only if  $\Delta_1([x_0 : x_1])$  and  $\Delta_2([y_0 : y_1])$  vanish at  $[a : b]$  and  $[c : d]$  respectively. We conclude with Lemma 3.2, that (1) is equivalent to (2).

Clearly, (1) implies (3). It remains to prove that (3) implies (1). Assume that  $P = (a_1, b_1)$  is fixed by  $\sigma$ . After writing  $\iota_1(P) = (a_1, b'_1)$  and  $\iota_2(\iota_1(P)) = (a'_1, b'_1)$ , it is clear that  $\sigma(P) = P$  implies successively  $\iota_1(P) = P$  and  $\iota_2(P) = P$ .  $\square$

#### 4. UNIFORMIZATION OF THE KERNEL CURVE

We still consider a *weighted model of nondegenerate walk*. The aim of this section is to give an explicit uniformization of  $\overline{E}_t$ . Thanks to Proposition 2.1, the latter may have genus zero or one. Although there are algorithms to compute such uniformizations, see for instance [vH97, SWPD08], our presentation of explicit uniformizations allows us to understand in detail the pull-backs of  $\sigma$ ,  $\iota_1$  and  $\iota_2$  and therefore their effect on the generating series of the models of walks. Let us start with the genus zero case.

**4.1. Genus zero case.** Let us consider a *nondegenerate weighted model of walks* of genus zero. Thank to Corollary 2.6 combined with Remark 2.9, it suffices to consider the situation where the *nondegenerate model of walk* arises from the following family



Genus zero curves may be parametrized with maps  $\phi : \mathbb{P}^1(\mathbb{C}) \rightarrow \overline{E}_t$  which are bijective outside a finite set. The aim of this subsection, achieved with Proposition 4.7, is to find such a parametrization explicitly. Although we could have just written down the formula for this parametrization and verified its properties, we have preferred to explain how the formula arises. This requires a preliminary study of the automorphisms of  $\mathbb{P}^1(\mathbb{C})$  obtained by pulling back the maps  $\sigma$ ,  $\iota_1$  and  $\iota_2$  by  $\phi$ , which is done with a series of lemmas preceding Proposition 4.7.

According to Lemma 2.8,  $\overline{E}_t$  has a unique singular point  $\Omega = (a_1, b_1) = ([0 : 1], [0 : 1])$ . Moreover  $\Delta_1([x_0 : x_1])$  has degree four with a double root at  $a_1 = [0 : 1]$  and the remaining two roots  $a_3, a_4$  are distinct. We let  $S_3$  and  $S_4$  be the points of  $\overline{E}_t$  with first coordinate  $a_3$  and  $a_4$  respectively. Similarly,  $\Delta_2([y_0 : y_1])$  has degree four with a double root at  $b_1 = [0 : 1]$  and the remaining two roots  $b_3, b_4$  are distinct. We let  $S'_3$  and  $S'_4$  be the points of  $\overline{E}_t$  with second coordinates  $b_3$  and  $b_4$  respectively.

Since  $\overline{E}_t$  has genus zero, there is a rational parametrization of  $\overline{E}_t$  [Ful89, Page 198, Ex.1], i.e. there exists a birational map

$$\begin{aligned} \phi = (\check{x}, \check{y}) : \mathbb{P}^1(\mathbb{C}) &\dashrightarrow \overline{E}_t \\ s &\mapsto (\check{x}(s), \check{y}(s)). \end{aligned}$$

To simplify the notation, we will abusively denote  $(\check{x}, \check{y})$  by  $(x, y)$ . We will now follow the ideas contained in [FIM17] to produce an explicit uniformization of  $\overline{E}_t$  in Proposition 4.7. If we set  $t = 1$ , we recover the uniformization of [FIM17, Section 6.4.3]. However, it is not clear if their proof can be simply modified to hold in our context, so we preferred to give proofs here with a slightly different strategy.

**Lemma 4.1.** *The map  $\phi$  is surjective and induces a bijection between  $\mathbb{P}^1(\mathbb{C}) \setminus \phi^{-1}(\Omega)$  and  $\overline{E}_t \setminus \{\Omega\}$ .*

*Proof.* As any nonconstant rational map from  $\mathbb{P}^1(\mathbb{C})$  to a projective curve,  $\phi$  is actually a surjective morphism of curves, see [Ful89, Corollary 1, Page 160]. Since  $\Omega$  is the unique singular point of  $\overline{E}_t$ , the result follows.  $\square$

The maps  $x, y : \mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{P}^1(\mathbb{C})$  are surjective morphisms of curves as well.

We let  $s_3, s_4 \in \mathbb{P}^1(\mathbb{C})$  (resp.  $s'_3, s'_4 \in \mathbb{P}^1(\mathbb{C})$ ) be such that  $S_3 = \phi(s_3)$  and  $S_4 = \phi(s_4)$  (resp.  $S'_3 = \phi(s'_3)$  and  $S'_4 = \phi(s'_4)$ ).

We will need to know the cardinality of  $x^{-1}(P)$  (resp.  $y^{-1}(P)$ ) for  $P \in \mathbb{P}^1(\mathbb{C})$ . This quantity might depend on  $P$  but it is a general fact about morphisms of curves that the cardinality of  $x^{-1}(P)$  (resp.  $y^{-1}(P)$ ) is constant for  $P$  outside a finite subset of  $\mathbb{P}^1(\mathbb{C})$ . This common value is called the degree of  $x$  (resp.  $y$ ). Inside this finite set, the cardinality can only fall, so is less than the degree.

**Lemma 4.2.** *The morphisms  $x, y : \mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{P}^1(\mathbb{C})$  have degree two.*

*Proof.* We will see that this is a consequence of the fact that  $\overline{E}_t$  is a biquadratic curve. Observe that by Lemma 4.1,  $\phi$  induces a bijection between  $\mathbb{P}^1(\mathbb{C}) \setminus \phi^{-1}(\Omega)$  and  $\overline{E}_t \setminus \{\Omega\}$ . Any  $(a, b) \in \overline{E}_t$  with  $a \neq a_1$  cannot be  $\Omega$  and therefore has a unique preimage by  $\phi$ . Additionally, let  $Z$  be the finite set of zeros of the discriminant  $\Delta_1$ . Then, for any  $a \notin Z$ , the cardinality  $(\{a\} \times \mathbb{P}^1(\mathbb{C})) \cap \overline{E}_t$  is two. Since  $x^{-1}(a) = \phi^{-1}((\{a\} \times \mathbb{P}^1(\mathbb{C})) \cap \overline{E}_t)$  and  $a_1 \in Z$ , it follows that if  $a \notin Z$ , the cardinality of  $x^{-1}(a)$  is two. So,  $x$  has degree two. The argument for  $y$  is similar.  $\square$

Since  $\phi$  is a birational map, the involutive automorphisms  $\iota_1, \iota_2$  of  $\overline{E}_t$  induce involutive automorphisms  $\tilde{\iota}_1, \tilde{\iota}_2$  of  $\mathbb{P}^1(\mathbb{C})$  via  $\phi$ . Similarly,  $\sigma$  induces an automorphism  $\tilde{\sigma}$  of  $\mathbb{P}^1(\mathbb{C})$ . In other words, we have the commutative diagrams

$$\begin{array}{ccc} \overline{E}_t & \xrightarrow{\iota_k} & \overline{E}_t & \text{and} & \overline{E}_t & \xrightarrow{\sigma} & \overline{E}_t \\ \phi \uparrow & & \uparrow \phi & & \phi \uparrow & & \uparrow \phi \\ \mathbb{P}^1(\mathbb{C}) & \xrightarrow{\tilde{\iota}_k} & \mathbb{P}^1(\mathbb{C}) & & \mathbb{P}^1(\mathbb{C}) & \xrightarrow{\tilde{\sigma}} & \mathbb{P}^1(\mathbb{C}) \end{array}$$

Note that since by Lemma 4.1  $\phi$  induces a bijection between  $\mathbb{P}^1(\mathbb{C}) \setminus \phi^{-1}(\Omega)$  and  $\overline{E}_t \setminus \{\Omega\}$  and  $\Omega$  is fixed by  $\iota_1, \iota_2$ , see Lemma 3.3, the group generated by  $\iota_1$  and  $\iota_2$  is isomorphic to the group generated by  $\tilde{\iota}_1$  and  $\tilde{\iota}_2$ . Thus we recover the same group as in [BMM10] for instance. Note that although the cardinal of the group may depends upon  $t$ , see Remark 4.13, since the maps  $\iota_1, \iota_2$  are defined on  $\mathbb{Q}(d_{i,j})(t)$ , two distinct values of  $t$  transcendental over  $\mathbb{Q}(d_{i,j})$  lead to isomorphic groups. We summarize some remarks in the following lemmas.

**Lemma 4.3.** *We have  $x = x \circ \tilde{\iota}_1$  and  $y = y \circ \tilde{\iota}_2$ .*

*Proof.* We obtain  $x = x \circ \tilde{\iota}_1$  by equating the first coordinates in the equality  $\phi \circ \tilde{\iota}_1 = \iota_1 \circ \phi$  and we obtain  $y = y \circ \tilde{\iota}_2$  by equating the second coordinates in the equality  $\phi \circ \tilde{\iota}_2 = \iota_2 \circ \phi$ .  $\square$

**Lemma 4.4.** *Let  $P = \phi(s) \in \overline{E}_t$  and let  $k \in \{1, 2\}$ . We have:*

- if  $\tilde{\iota}_k(s) = s$ , then  $\iota_k(P) = P$ ;
- if  $P \neq \Omega$  and  $\iota_k(P) = P$ , then  $\tilde{\iota}_k(s) = s$ .

Furthermore the map  $\tilde{\iota}_1$  (resp.  $\tilde{\iota}_2$ ) has exactly two fixed points, namely  $s_3$  and  $s_4$  (resp.  $s'_3$  and  $s'_4$ ).

*Proof.* We have  $\iota_k(P) = \iota_k(\phi(s)) = \phi(\tilde{\iota}_k(s))$ . The first assertion is now clear, and the second one follows from the fact that  $\phi$  is injective on  $\overline{E_t} \setminus \phi^{-1}(\Omega)$ . Since  $S_3, S_4 \neq \Omega$  are fixed by  $\iota_1$ , this shows that  $s_3$  and  $s_4$  are fixed by  $\tilde{\iota}_1$ . Similar proof holds for  $\tilde{\iota}_2$ .

It remains to prove that there are exactly two points fixed by  $\tilde{\iota}_k$ . To the contrary, assume that there is a third point fixed by  $\tilde{\iota}_k$ . Since  $\tilde{\iota}_k$  is an automorphism of  $\mathbb{P}^1(\mathbb{C})$ , i.e. an homography, with three fixed points, it is the identity. This is a contradiction and concludes the proof of the lemma.  $\square$

**Lemma 4.5.** *The preimage of  $\Omega$  by  $\phi$  has two elements.*

*Proof.* We know that  $x, y : \mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{P}^1(\mathbb{C})$  have degree two, so  $\phi^{-1}(\Omega)$  has one or two elements. Suppose that  $\phi^{-1}(\Omega)$  has exactly one element, say  $s_1$ . In virtue of  $\phi(s_3) = S_3$  and  $\phi(s_4) = S_4$ ,  $s_1$  is different to  $s_3, s_4$ . Since  $\phi(\tilde{\iota}_1(s_1)) = \iota_1(\phi(s_1)) = \iota_1(\Omega) = \Omega$ , we have  $\tilde{\iota}_1(s_1) = s_1$ . This contradicts Lemma 4.4. Hence,  $\phi^{-1}(\Omega)$  has two elements.  $\square$

From now on, we define  $s_1 \neq s_2$  the two preimages of  $\Omega$  by  $\phi$ , that is

$$\{s_1, s_2\} := \phi^{-1}(\Omega).$$

**Lemma 4.6.** *The map  $\tilde{\iota}_1$  (resp.  $\tilde{\iota}_2$ ) interchanges  $s_1$  and  $s_2$ . The map  $\tilde{\sigma}$  has exactly two distinct fixed points:  $s_1$  and  $s_2$ .*

*Proof.* We have  $\phi(s) = \Omega$  if and only if  $s = s_1$  or  $s_2$  and the equality  $\iota_1(\phi(s)) = \phi(\tilde{\iota}_1(s))$  shows that  $\tilde{\iota}_1$  induces a permutation of  $\phi^{-1}(\Omega) = \{s_1, s_2\}$ . By Lemma 4.4,  $s_1$  is not fixed by  $\tilde{\iota}_1$ , showing that the permutation is not the identity, i.e.  $\tilde{\iota}_1$  interchanges  $s_1$  and  $s_2$ .

The proof for  $\tilde{\iota}_2$  is similar.

As any homography which is not the identity,  $\tilde{\sigma}$  has at most two fixed points in  $\mathbb{P}^1(\mathbb{C})$ . It only remains to prove that  $s_1$  and  $s_2$  are fixed by  $\tilde{\sigma}$ , and this is indeed the case because  $\tilde{\sigma} = \tilde{\iota}_2 \circ \tilde{\iota}_1$  and  $\tilde{\iota}_1, \tilde{\iota}_2$  interchange  $s_1$  and  $s_2$ .  $\square$

We are now ready to give an explicit expression of  $\phi$ . The coefficients  $\alpha_i, \beta_i$  of the discriminants in this situation are given by the formulas

$$\begin{aligned} \alpha_0(t) &= \alpha_1(t) = 0 \\ \beta_0(t) &= \beta_1(t) = 0 \\ \alpha_2(t) &= \beta_2(t) = 1 - 2td_{0,0} + t^2d_{0,0}^2 - 4t^2d_{-1,1}d_{1,-1} \\ \alpha_3(t) &= 2t^2d_{1,0}d_{0,0} - 2td_{1,0} - 4t^2d_{0,1}d_{1,-1} \\ \beta_3(t) &= 2t^2d_{0,1}d_{0,0} - 2td_{0,1} - 4t^2d_{1,0}d_{-1,1} \\ \alpha_4(t) &= t^2(d_{1,0}^2 - 4d_{1,1}d_{1,-1}) \\ \beta_4(t) &= t^2(d_{0,1}^2 - 4d_{1,1}d_{-1,1}). \end{aligned}$$

Note that for  $k = 3, 4$ ,  $\beta_k(t)$ , may be obtained from  $\alpha_k(t)$  by interchanging  $d_{1,0}$  with  $d_{0,1}$  and  $d_{1,-1}$  with  $d_{-1,1}$ .




FIGURE 2. An idealized representation of the uniformization map used in Proposition 4.7. The left hand side represents the complex Riemann sphere and the right hand side the curve  $\overline{E}_t$ , seen as an abstract complex algebraic curve.

**Proposition 4.7.** *An explicit parametrization  $\phi : \mathbb{P}^1(\mathbb{C}) \rightarrow \overline{E}_t$  such that*

$$\tilde{t}_1(s) = \frac{1}{s}, \quad \tilde{t}_2(s) = \frac{\lambda^2}{s} = \frac{q}{s} \quad \text{and} \quad \tilde{\sigma}(s) = qs$$

for a certain  $\lambda \in \mathbb{C}^*$  is given by

$$\phi(s) = \left( \frac{4\alpha_2(t)}{\sqrt{\alpha_3(t)^2 - 4\alpha_2(t)\alpha_4(t)}(s + \frac{1}{s}) - 2\alpha_3(t)}, \frac{4\beta_2(t)}{\sqrt{\beta_3(t)^2 - 4\beta_2(t)\beta_4(t)}(\frac{s}{\lambda} + \frac{\lambda}{s}) - 2\beta_3(t)} \right).$$

Moreover, we have, see Figure 2

$$\begin{aligned} x(0) = x(\infty) = a_1, \quad x(1) = a_3, \quad x(-1) = a_4, \\ y(0) = y(\infty) = b_1, \quad y(\lambda) = b_3, \quad y(-\lambda) = b_4. \end{aligned}$$

*Proof.* According to Lemma 4.6,  $\tilde{t}_1$  is an involutive homography with fixed points  $s_3$  and  $s_4$ , so there exists an homography  $h$  such that  $h(s_3) = 1$ ,  $h(s_4) = -1$  and  $h \circ \tilde{t}_1 \circ h^{-1}(s) = 1/s$ . Up to replacing  $\phi$  by  $\phi \circ h$ , we can assume that  $s_3 = 1$ ,  $s_4 = -1$  and  $\tilde{t}_1(s) = \frac{1}{s}$ . Since  $s_1 \neq s_2$ , we can assume up to renumbering that  $s_1 \neq \infty$ . Let us consider the homography  $k(s) = \frac{s-s_1}{-s_1s+1}$ . Note that  $k$  commutes with  $s \mapsto 1/s$ , so changing  $\phi$  by  $\phi \circ k$  does not affect  $\tilde{t}_1$ , and we can also assume that  $s_1 = [0 : 1]$  and  $s_2 = [1 : 0]$ . Lemma 4.2 and Lemma 4.3 ensure that the morphism  $x : \mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{P}^1(\mathbb{C})$  has degree two and satisfies  $x(s) = x(1/s)$  for all  $s \in \mathbb{P}^1(\mathbb{C})$ . Since the morphism  $x : \mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{P}^1(\mathbb{C})$  has degree two, see Lemma 4.2, it follows that

$$x(s) = \frac{a(s + 1/s) + b}{c(s + 1/s) + d}$$

for some  $a, b, c, d \in \mathbb{C}$ . We have  $x(s_1) = x([0 : 1]) = a_1 = 0$ ,  $x(s_2) = x([1 : 0]) = a_1 = 0$ ,  $x(s_3) = x([1 : 1]) = a_3$  and  $x(s_4) = x([-1 : 1]) = a_4$ . The equality  $x([1 : 0]) = 0$  implies  $a = 0$ . The equalities  $x([1 : 1]) = a_3$  and  $x([-1 : 1]) = a_4$  imply

$$x(s) = \frac{4a_3a_4}{(a_4 - a_3)(s + \frac{1}{s}) + 2(a_3 + a_4)}.$$

The known expressions for  $a_3$  and  $a_4$  given in Lemma 2.8 lead to the expected expression for  $x(s)$ .

According to Lemma 4.6,  $\tilde{t}_2$  is an homography interchanging  $[0 : 1]$  and  $[1 : 0]$ , so  $\tilde{t}_2(s) = \frac{\lambda^2}{s}$  for some  $\lambda \in \mathbb{C}^*$ . Up to renumbering, we have  $s'_3 = \lambda$  and  $s'_4 = -\lambda$ . Using the fact that the morphism  $y : \mathbb{P}^1(\mathbb{C}) \rightarrow \mathbb{P}^1(\mathbb{C})$  has degree two and is invariant by  $\tilde{t}_2$ , and arguing as we did above for  $x$ , we see that there exist  $\alpha, \beta, \gamma, \eta \in \mathbb{C}$  such that

$$y(s) = \frac{\alpha(\frac{s}{\lambda} + \frac{\lambda}{s}) + \beta}{\gamma(\frac{s}{\lambda} + \frac{\lambda}{s}) + \eta}.$$

The equality  $y([1 : 0]) = 0$  implies  $\alpha = 0$ . Using the equalities  $y(s'_3) = y(\lambda) = b_3$  and  $y(s'_4) = y(-\lambda) = b_4$ , and arguing as we did above for  $x$ , we obtain the expected expression for  $y(s)$ .  $\square$

*Remark 4.8.* (1) The uniformization is not unique. More precisely, the possible uniformizations are of the form  $\phi \circ h$ , where  $h$  is an homography. However, if one requires that  $h$  fixes setwise  $\{[0 : 1], [1 : 0]\}$  then  $q$  is uniquely defined up to inversion.  
(2) The real  $q$  or  $q^{-1}$  specializes for  $t = 1$  to the real  $\rho^2$  in [FIM17, Page 178].

The following proposition determines  $q$  up to its inverse. We include this for completeness.

**Proposition 4.9** ([DHRS20], Proposition 1.7, Corollary 1.10). *One of the two complex numbers  $q$  or  $q^{-1}$  is equal to*

$$\frac{-1 + d_{0,0}t - \sqrt{(1 - d_{0,0}t)^2 - 4d_{1,-1}d_{-1,1}t^2}}{-1 + d_{0,0}t + \sqrt{(1 - d_{0,0}t)^2 - 4d_{1,-1}d_{-1,1}t^2}}.$$

Furthermore,  $q \in \mathbb{R} \setminus \{\pm 1\}$ .

*Remark 4.10.* This implies that  $\sigma$  and  $\tilde{\sigma}$  have infinite order (see also [BMM10, FR11]). Because  $\sigma$  is induced on  $\overline{E_t}$  by  $i_1 \circ i_2$ , we find that  $i_1 \circ i_2$  has infinite order as well. It follows that the *group of the walk* introduced in [BMM10], which is by definition the group generated by  $i_1$  and  $i_2$ , has infinite order. Note that this was proved in [BMM10] using a valuation argument.

*Remark 4.11.* We stress the fact that since  $\phi(s)$ ,  $q$  and  $\overline{E_t}$  depend continuously on  $t$  and the set of transcendental number over  $\mathbb{Q}(d_{i,j})$  in  $]0, 1[$  is dense in  $]0, 1[$ , we deduce that Proposition 4.7 and Proposition 4.9 stay valid for every  $t \in ]0, 1[$ .

**4.2. Genus one case.** In this section, we consider the uniformization problem in the genus one context. This problem has been solved in [DR19]. We recall below the main result of [DR19], for the sake of completeness. Let us consider a *nondegenerate model of walk* of genus one. By Proposition 2.1,  $\overline{E}_t$  is a smooth curve of genus one and, by Corollary 2.6, this corresponds to the situation where the step set is not included in any half plane whose boundary passes through  $(0,0)$ . By [WW96, Chapter XX], it is biholomorphic to  $\mathbb{C}/(\mathbb{Z}\omega_1 + \mathbb{Z}\omega_2)$  for some lattice  $\mathbb{Z}\omega_1 + \mathbb{Z}\omega_2$  of  $\mathbb{C}$  via some  $(\mathbb{Z}\omega_1 + \mathbb{Z}\omega_2)$ -periodic holomorphic map

$$(4.1) \quad \begin{aligned} \Lambda : \mathbb{C} &\rightarrow \overline{E}_t \\ \omega &\mapsto (\mathfrak{q}_1(\omega), \mathfrak{q}_2(\omega)), \end{aligned}$$

where  $\mathfrak{q}_1, \mathfrak{q}_2$  are rational functions of  $\wp$  and its derivative  $d\wp/d\omega$ , and  $\wp$  is the Weierstrass function associated with the lattice  $\mathbb{Z}\omega_1 + \mathbb{Z}\omega_2$ :

$$(4.2) \quad \wp(\omega) = \wp(\omega; \omega_1, \omega_2) := \frac{1}{\omega^2} + \sum_{(\ell_1, \ell_2) \in \mathbb{Z}^2 \setminus \{(0,0)\}} \left( \frac{1}{(\omega + \ell_1\omega_1 + \ell_2\omega_2)^2} - \frac{1}{(\ell_1\omega_1 + \ell_2\omega_2)^2} \right).$$

Then, the field of meromorphic functions on  $\overline{E}_t$  is isomorphic to the field of meromorphic functions on  $\mathbb{C}/(\mathbb{Z}\omega_1 + \mathbb{Z}\omega_2)$ , which is itself isomorphic to the field of meromorphic functions on  $\mathbb{C}$  that are  $(\omega_1, \omega_2)$ -periodic. The latter field is equal to  $\mathbb{C}(\wp, \wp')$ , see [WW96].

The maps  $\iota_1, \iota_2$  and  $\sigma$  may be lifted to the  $\omega$ -plane  $\mathbb{C}$ . We denote these lifts by  $\tilde{\iota}_1, \tilde{\iota}_2$  and  $\tilde{\sigma}$  respectively. So we have the commutative diagrams

$$\begin{array}{ccc} \overline{E}_t & \xrightarrow{\iota_k} & \overline{E}_t \\ \uparrow \Lambda & & \uparrow \Lambda \\ \mathbb{C} & \xrightarrow{\tilde{\iota}_k} & \mathbb{C} \end{array} \quad \begin{array}{ccc} \overline{E}_t & \xrightarrow{\sigma} & \overline{E}_t \\ \uparrow \Lambda & & \uparrow \Lambda \\ \mathbb{C} & \xrightarrow{\tilde{\sigma}} & \mathbb{C} \end{array}$$

The following result has been proved

- in [FIM17, Section 3.3] when  $t = 1$ ,
- in [Ras12] in the *unweighted* case for general  $0 < t < 1$ , not necessarily transcendental over  $\mathbb{Q}(d_{i,j})$ ,
- in [DR19, Proposition 18] in the *weighted* case, with general  $0 < t < 1$ , not necessarily transcendental over  $\mathbb{Q}(d_{i,j})$ .

In what follows, we set  $D(\omega) = \Delta_1([\omega : 1])$ . Let us introduce  $z = 2A(x)y + B(x)$ , where  $A(x) = t(d_{-1,1} + d_{0,1}x + d_{1,1}x^2)$ , and  $B(x) = t(d_{-1,0} - \frac{1}{t}x + d_{0,0}x + d_{1,0}x^2)$ .

**Proposition 4.12.** *An explicit uniformization  $\Lambda : \mathbb{C} \rightarrow \overline{E}_t$  such that*

$$\tilde{\iota}_1(\omega) = -\omega, \quad \tilde{\iota}_2(\omega) = -\omega + \omega_3 \quad \text{and} \quad \tilde{\sigma}(\omega) = \omega + \omega_3,$$

for a certain  $\omega_3 \in \mathbb{C}^*$  is given by

$$\Lambda(\omega) = (x(\omega), y(\omega))$$



where  $x(\omega)$  and  $y(\omega)$  are given by

	$x(\omega)$	$z(\omega)$
$a_4 \neq [1:0]$	$\left[ a_4 + \frac{D'(a_4)}{\wp(\omega) - \frac{1}{6}D''(a_4)} : 1 \right]$	$\left[ \frac{D'(a_4)\wp'(\omega)}{2(\wp(\omega) - \frac{1}{6}D''(a_4))^2} : 1 \right]$
$a_4 = [1:0]$	$[\wp(\omega) - \alpha_2/3 : \alpha_3]$	$[-\wp'(\omega) : 2\alpha_3]$

A suitable choice for the periods  $(\omega_1, \omega_2)$  is given by the elliptic integrals

$$(4.3) \quad \omega_1 = \mathbf{i} \int_{a_3}^{a_4} \frac{dx}{\sqrt{|D(x)|}} \in \mathbf{i}\mathbb{R}_{>0} \quad \text{and} \quad \omega_2 = \int_{a_4}^{a_1} \frac{dx}{\sqrt{D(x)}} \in \mathbb{R}_{>0}.$$

Note that, according to [DR19, Section 2],

$$\omega_3 = \int_{a_4}^{X_{\pm}(b_4)} \frac{dx}{\sqrt{D(x)}} \in ]0, \omega_2[.$$

*Remark 4.13.* Contrary to the genus zero situation, the map  $\sigma$  may have finite order.

#### REFERENCES

- [BBMR15] Olivier Bernardi, Mireille Bousquet-Mélou, and Kilian Raschel, *Counting quadrant walks via Tutte's invariant method*, An extended abstract to appear in *Proceedings of FPSAC 2016*, Discrete Math. Theor. Comput. Sci. Proc., arXiv:1511.04298, 2015.
- [BBMR17] ———, *Counting quadrant walks via Tutte's invariant method*, Preprint, arXiv:1708.08215, 2017.
- [BMM10] Mireille Bousquet-Mélou and Marni Mishna, *Walks with small steps in the quarter plane*, Algorithmic probability and combinatorics, Contemp. Math., vol. 520, Amer. Math. Soc., Providence, RI, 2010, pp. 1–39.
- [BRS14] Alin Bostan, Kilian Raschel, and Bruno Salvy, *Non-D-finite excursions in the quarter plane*, J. Combin. Theory Ser. A **121** (2014), 45–63. MR 3115331
- [BvHK10] Alin Bostan, Mark van Hoeij, and Manuel Kauers, *The complete generating function for Gessel walks is algebraic*, Proc. Amer. Math. Soc. **138** (2010), no. 9, 3063–3078.
- [CLO97] David Cox, John Little, and Donal O'Shea, *Ideals, varieties, and algorithms*, second ed., Undergraduate Texts in Mathematics, Springer-Verlag, New York, 1997, An introduction to computational algebraic geometry and commutative algebra. MR 1417938
- [DH19] Thomas Dreyfus and Charlotte Hardouin, *Length derivative of the generating series of walks confined in the quarter plane*, arXiv preprint arXiv:1902.10558 (2019).
- [DHRS18] Thomas Dreyfus, Charlotte Hardouin, Julien Roques, and Michael F Singer, *On the nature of the generating series of walks in the quarter plane*, Inventiones mathematicae (2018), 139–203.
- [DHRS20] ———, *Walks in the quarter plane, genus zero case*, Journal of Combinatorial Theory, Series A. (2020).
- [DR19] Thomas Dreyfus and Kilian Raschel, *Differential transcendence & algebraicity criteria for the series counting weighted quadrant walks*, Publications Mathématiques de Besançon (2019), no. 1, 41–80.
- [Dui10] J. Duistermaat, *Discrete integrable systems: QRT maps and elliptic surfaces*, Springer Monographs in Mathematics, vol. 304, Springer-Verlag, New York, 2010.
- [DW15] D. Denisov and V. Wachtel, *Random walks in cones*, Ann. Probab. **43** (2015), no. 3, 992–1044. MR 3342657
- [FIM99] Guy Fayolle, Roudolf Iasnogorodski, and VA Malyshev, *Random walks in the quarter-plane*, vol. 40, Springer, 1999.

- [FIM17] Guy Fayolle, Roudolf Iasnogorodski, and Vadim Malyshev, *Random walks in the quarter plane. Algebraic methods, boundary value problems, applications to queueing systems and analytic combinatorics. 2nd edition, previously published with the subtitle Algebraic methods, boundary value problems and applications.*, vol. 40, Cham: Springer, 2017 (English).
- [FR11] G. Fayolle and K. Raschel, *Random walks in the quarter-plane with zero drift: an explicit criterion for the finiteness of the associated group*, Markov Process. Related Fields **17** (2011), no. 4, 619–636.
- [Ful84] William Fulton, *Introduction to intersection theory in algebraic geometry*, no. 54, American Mathematical Soc., 1984.
- [Ful89] ———, *Algebraic curves*, Advanced Book Classics, Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA, 1989, An introduction to algebraic geometry, Notes written with the collaboration of Richard Weiss, Reprint of 1969 original.
- [Har77] Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York-Heidelberg, 1977, Graduate Texts in Mathematics, No. 52. MR 0463157
- [KR12] Irina Kurkova and Kilian Raschel, *On the functions counting walks with small steps in the quarter plane*, Publ. Math. Inst. Hautes Études Sci. **116** (2012), 69–114. MR 3090255
- [KY15] Manuel Kauers and Rika Yatchak, *Walks in the quarter plane with multiple steps*, Proceedings of FPSAC 2015, Discrete Math. Theor. Comput. Sci. Proc., Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2015, pp. 25–36.
- [Mis09] Marni Mishna, *Classifying lattice walks restricted to the quarter plane*, J. Combin. Theory Ser. A **116** (2009), no. 2, 460–477. MR 2475028
- [MM14] Stephen Melzer and Marni Mishna, *Singularity analysis via the iterated kernel method*, Combin. Probab. Comput. **23** (2014), no. 5, 861–888. MR 3249228
- [MR09] Marni Mishna and Andrew Rechnitzer, *Two non-holonomic lattice walks in the quarter plane*, Theoret. Comput. Sci. **410** (2009), no. 38-40, 3616–3630. MR 2553316
- [Ras12] Kilian Raschel, *Counting walks in a quadrant: a unified approach via boundary value problems*, J. Eur. Math. Soc. (JEMS) **14** (2012), no. 3, 749–777. MR 2911883
- [SWPD08] J Rafael Sendra, Franz Winkler, and Sonia Pérez-Díaz, *Rational algebraic curves*, Algorithms and Computation in Mathematics **22** (2008).
- [vH97] Mark van Hoeij, *Rational parametrizations of algebraic curves using a canonical divisor*, Journal of Symbolic Computation **23** (1997), no. 2-3, 209–227.
- [WW96] E. Whittaker and G. Watson, *A course of modern analysis*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1996, An introduction to the general theory of infinite processes and of analytic functions; with an account of the principal transcendental functions, Reprint of the fourth (1927) edition. MR 1424469

INSTITUT DE RECHERCHE MATHÉMATIQUE AVANCÉE, U.M.R. 7501 UNIVERSITÉ DE STRASBOURG  
ET C.N.R.S. 7, RUE RENÉ DESCARTES 67084 STRASBOURG, FRANCE

*Email address:* `dreyfus@math.unistra.fr`

UNIVERSITÉ PAUL SABATIER - INSTITUT DE MATHÉMATIQUES DE TOULOUSE, 118 ROUTE DE NARBONNE, 31062 TOULOUSE.

*Email address:* `hardouin@math.univ-toulouse.fr`

UNIV LYON, UNIVERSITÉ CLAUDE BERNARD LYON 1, CNRS UMR 5208, INSTITUT CAMILLE JORDAN, 43 BLVD. DU 11 NOVEMBRE 1918, F-69622 VILLEURBANNE CEDEX, FRANCE

*Email address:* `roques@math.univ-lyon1.fr`

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, BOX 8205, RALEIGH, NC 27695-8205, USA

*Email address:* `singer@ncsu.edu`