



**HAL**  
open science

# Dynamical mechanisms for Kaluza-Klein theories

Frédéric Hélein

► **To cite this version:**

| Frédéric Hélein. Dynamical mechanisms for Kaluza-Klein theories. 2022. hal-03513721v2

**HAL Id: hal-03513721**

**<https://hal.science/hal-03513721v2>**

Preprint submitted on 14 May 2022 (v2), last revised 15 Sep 2022 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Dynamical mechanisms for Kaluza–Klein theories

Frédéric HÉLEIN\*

14 mai 2022

*Abstract : We present variational formulations of gauge theories and Einstein–Yang–Mills equations in the spirit of Kaluza–Klein theories. For gauge theories only a topological fibration is assumed. For gravitation coupled with gauge fields no fibration is assumed : fields are defined on a 'space-time'  $\mathcal{Y}$  of dimension  $4 + r$  without any structure a priori, where  $r$  is the dimension of the structure group. If the latter is compact and simply connected, classical solutions allow to construct a manifold  $\mathcal{X}$  of dimension 4 to be the physical space-time, in such a way that  $\mathcal{Y}$  acquires the structure of a principal bundle over  $\mathcal{X}$ , and lead to solutions of the Einstein–Yang–Mills systems. The special case of the Einstein–Maxwell system is also discussed : it suffices that at least one fiber closes in on a circle to deduce that the 5-dimensional space-time has a fiber bundle structure.*

*Keywords :* Kaluza–Klein, Gauge theories, General Relativity, Differential Geometry

## 1 Introduction

Kaluza–Klein theories (see [1] for an account) goes back to the work of T. Kaluza [8] in 1921 and O. Klein [10] in 1926, in an attempt to unify gravity and electromagnetism as a manifestation of a 5-dimensional relativistic gravity theory on a manifold fibered over some 4-dimensional space-time. This theory was abandoned for various reasons before arousing new interest in the context of supergravity and string theory. Some inconsistency were observed and fixed through the introduction of an additional fields (*radion* or *dilaton*) independantly by P. Jordan [7] in 1947 and Y. Thiry [13] in 1948. The introduction of this field, the physical meaning of which is not clear, can be avoided by not imposing the Einstein equation on the higher dimensional space-time and instead by looking for the critical points of the Einstein–Hilbert action when assuming a fiber bundle structure and equivariance constraints. By following this alternative option the theory was extended to Yang–Mills fields by R. Kerner [9] in 1968, leading to the exact Einstein–Yang–Mills system (see also [2]).

Another problem is to explain why the extra dimensions cannot be observed. One of the most common hypothesis, due to Klein, is to suppose that the extra dimension are

---

\*Institut de Mathématiques de Jussieu, UMR CNRS 7586 Université Denis Diderot — Paris 7, UFR de Mathématiques, Bâtiment Sophie Germain 75205 Paris Cedex 13, France, [helein@math.univ-paris-diderot.fr](mailto:helein@math.univ-paris-diderot.fr)

tiny, so that, in the quantized version, higher modes are too energetic to be observable. Another one is simply to assume that the metric is invariant along extra dimensions. But this assumption needs to be justified.

This is the question we address here : we propose theoretical models coupling gravity with electromagnetism or gauge fields such that, at the classical level, *the invariance of fields along extra dimensions follows from dynamical equations*, so that there is no need to assume this condition *a priori*.

Hence in the following models, fields which are not solutions of the dynamical equations depend in general on  $4+r$  variables, where  $r$  is the number of extra dimensions. But some of the dynamical equations, caused by *auxiliary fields* which play the role of Lagrange multipliers, force the apparition of a fibration of the higher dimensional space-time over some 4-dimensional physical space-time.

However these auxiliary fields may spoil the result (in a way similar to *dilatons*) by creating some artificial matter sources in the final equations. This is where an unexpected *cancellation phenomenon* (20) comes into play, leading to the vanishing of terms coming from auxiliary fields in the equations, when there are projected on the space-time.

## 2 Results

In the following we present several models of growing generality, in order to expound the various mechanisms underlying our theory. In Section 3 we present a model leading to Maxwell equations on a *fixed* space-time but by replacing the usual principal  $U(1)$ -bundle by a topological circle bundle. As a consequence the gauge symmetry group is larger than in the standard theory since, in addition to the usual gauge symmetries, the action functional is invariant by fibration preserving diffeomorphisms. This shows how the invariance of some fields along extra dimensions can be achieved and also gives a first glance at the *cancellation mechanism*.

In Section 4 we extend this construction for Yang–Mills fields, still over a fixed space-time, for a structure group which is *compact* and *simply connected* (e.g. any product of  $SU(k)$ 's). It generalizes results in [4] (where we needed to assume a principal bundle structure *a priori*).

In Section 5 we then turn on the general Kaluza–Klein model for a *compact* and *simply connected* structure group. Fibers are obtained by integrating an exterior differential system through Frobenius's theorem. Most of these results were obtained in [5] (without a bare cosmological constant).

The previous model however does not work for  $U(1)$  since this group is not simply connected. This question is addressed in Section 6 where we show that, if there exists *at least* one integral curve of the exterior differential system which closes, then we obtain a 4-dimensional space-time, a fibration and, at the very end, a solution of the Einstein–Maxwell system.

Throughout the paper all fields are assumed to be smooth.

### 3 Model for Maxwell equations

Consider a connected 4-dimensional space-time manifold  $\mathcal{X}^4$  with a fixed Riemannian metric  $\mathbf{g}_{\mu\nu}$ , a 5-dimensional space-time  $\mathcal{Y}^5$  and suppose that there exists a smooth projection map  $\mathcal{Y}^5 \xrightarrow{P} \mathcal{X}^4$  such that  $dP$  has a maximal rank (equal to 4) everywhere and that, for any  $x \in \mathcal{X}^4$  the inverse image  $\mathbf{f}_x := P^{-1}(x)$  is a topological circle. We denote by  $x = (x^\mu)_{0 \leq \mu \leq 3}$  the coordinates on  $\mathcal{X}^4$ , by  $y$  the coordinate on fibers  $\mathbf{f}_x$  and set  $z = (z^I)_{0 \leq I \leq 4} = (x, y)$ . We describe the metric on  $\mathcal{X}^4$  through a fixed co-vierbein  $e^a = e^a_\mu(x) dx^\mu$  for  $0 \leq a \leq 3$ , so that  $\mathbf{g}_{\mu\nu} = \eta_{ab} e^a_\mu e^b_\nu$ , where  $\eta_{ab}$  is the Minkowski metric.

Dynamical fields on  $\mathcal{Y}^5$  are pairs  $(\theta, \pi)$ , where  $\theta = \theta_\mu(x, y) dx^\mu + \theta_4(x, y) dy = \theta_I(z) dz^I$  and  $\pi = \frac{1}{3!} \pi_{IJK}(z) dz^I \wedge dz^J \wedge dz^K$ . We will however privilege another decomposition of  $\pi$ . Indeed we assume that  $\theta_4 \neq 0$  so that  $(e_0, e_1, e_2, e_3, \theta)$  provides us with a co-5-bein on  $\mathcal{Y}^5$ . We define

$$e^{(4)} = e^0 \wedge e^1 \wedge e^2 \wedge e^3, \quad e_a^{(3)} = \frac{1}{3!} \epsilon_{abcd} e^b \wedge e^c \wedge e^d$$

$$\text{and } e_{ab}^{(2)} = \frac{1}{2!} \epsilon_{abcd} e^c \wedge e^d$$

This allows us to decompose

$$\pi = \frac{1}{2} \pi^{ab}(x, y) e_{ab}^{(2)} \wedge \theta - \pi^a(x, y) e_a^{(3)} \quad (1)$$

We set  $\|\pi\|^2 = \frac{1}{2} \pi^{ab} \pi_{ab}$ , where  $\pi_{ab} = \eta_{ac} \eta_{bd} \pi^{cd}$ , and define the action functional

$$\mathcal{A}[\theta, \pi] = \int_{\mathcal{Y}^5} \frac{1}{2} \|\pi\|^2 e^{(4)} \wedge \theta + \pi \wedge d\theta \quad (2)$$

or  $\mathcal{A}[\theta, \pi] = \int_{\mathcal{Y}^5} \left( \frac{1}{4} \pi^{ab} \pi_{ab} + \frac{1}{2} \Theta_{ab} \pi^{ab} + \Theta_a \pi^a \right) e^{(4)} \wedge \theta$ , where we set  $d\theta = \frac{1}{2} \Theta_{ab} e^a \wedge e^b + \Theta_a e^a \wedge \theta$ .

#### 3.1 Euler–Lagrange equations

We denote by  $(e_I)_{0 \leq I \leq 4}$  the 5-bein dual to  $(e^a, \theta)$ . The Euler–Lagrange equations are

$$\begin{aligned} d\theta(e_a, \partial_y) / \theta_4 = \Theta_a &= 0 & \text{(a)} \\ \Theta_{ab} + \pi_{ab} &= 0 & \text{(b)} \\ d\pi &= \frac{1}{2} \|\pi\|^2 e^{(4)} & \text{(c)} \end{aligned} \quad (3)$$

Equation (3a) has the following consequence. Consider two points  $x_1, x_2 \in \mathcal{X}^4$  and a path  $\gamma$  joining  $x_1$  to  $x_2$  in  $\mathcal{X}^4$ . Its inverse image by  $P$  is a surface  $S \subset \mathcal{Y}^5$  with boundary  $\partial S = \mathbf{f}_{x_2} - \mathbf{f}_{x_1}$ . Since  $\partial_y$  is tangent to  $S$  we have by (3a)

$$\int_{\mathbf{f}_{x_2}} \theta - \int_{\mathbf{f}_{x_1}} \theta = \int_S d\theta = 0 \quad (4)$$

Hence  $q := \int_{\mathbf{f}_x} \theta$  is independent of  $x$  in  $\mathcal{X}^4$ . We can thus define the new coordinate

$$s = f(x, y) = \int_0^y \theta_4(x, y') dy' \text{ mod } [q]$$

and replace the coordinates  $(x, y)$  by  $(x, s)$ , where  $s \in \mathbb{R}/q\mathbb{Z}$ . The form  $\theta$  then reads  $\theta = \mathbf{A}_a(x, s)e^a + ds$ , where  $\mathbf{A}_a(x, f(x, y)) = (\theta - df)_{(x, y)}(e_a)$ . Equation (3a) further implies  $\partial_s \mathbf{A}_a = 0$ , so that

$$\theta = \mathbf{A}_a(x)e^a + ds \quad (5)$$

Hence  $d\theta = \mathbf{F} = \frac{1}{2}\mathbf{F}_{ab}e^a \wedge e^b$ , where  $\mathbf{F}_{ab} = \partial_a \mathbf{A}_b - \partial_a \mathbf{A}_b$ . We define  $p^{ab}(x, s)$  and  $p^a(x, s)$  such that

$$\begin{aligned} p^a(x, f(x, y)) &= \pi^a(x, y) - \pi^{ab}(x, y)\mathbf{A}_b(x, f(x, y)) \\ p^{ab}(x, f(x, y)) &= \pi^{ab}(x, y) \end{aligned}$$

Then  $\pi = \frac{1}{2}p^{ab}e_{ab}^{(2)} \wedge ds - p^a e_a^{(3)}$ , where  $p^{ab} = -\mathbf{F}^{ab}$  by (3b).

Let  $\gamma^a{}_b$  be the spin connection on  $\mathcal{X}^4$  for the vierbein  $(e_a)$ . It satisfies the torsion free condition

$$d^\gamma e^a := de^a + \gamma^a{}_b \wedge e^b = 0. \quad (6)$$

By setting  $d^\gamma p^{ab} := dp^{ab} + \gamma^a{}_c p^{cb} + \gamma^b{}_c p^{ac}$  and  $d^\gamma p^a := dp^a + \gamma^a{}_c p^c$  and by using (6) we find that  $d\pi = \frac{1}{2}d^\gamma p^{ab} \wedge e_{ab}^{(2)} \wedge ds - d^\gamma p^a \wedge e_a^{(3)}$ . By decomposing  $d^\gamma p^{ab} = \partial_c^\gamma p^{ab} e^c + \partial_s p^{ab} ds$  and  $d^\gamma p^a = \partial_c^\gamma p^a e^c + \partial_s p^a ds$  we hence get  $d\pi = (\partial_b^\gamma p^{ab} + \partial_s p^a) e_a^{(3)} \wedge ds - \partial_a^\gamma p^a e^{(4)}$ . Thus (3c) is equivalent to

$$\begin{aligned} \partial_b^\gamma \mathbf{F}^{ab}(x) &= \partial_s p^a(x, s) \quad (\text{a}) \\ \partial_a^\gamma p^a(x, s) &= -\frac{1}{2}\|\mathbf{F}\|^2(x) \quad (\text{b}) \end{aligned} \quad (7)$$

The key point to conclude that  $\mathbf{A}$  is a solution of Maxwell equations in vacuum is to observe that the l.h.s of (7a) is independent of  $s$ , thus

$$\partial_b^\gamma \mathbf{F}^{ab} = \frac{\int_0^q \partial_b^\gamma \mathbf{F}^{ab} ds}{\int_0^q ds} = \frac{\int_0^q \partial_s p^a ds}{q} = 0 \quad (8)$$

because  $p^a$  is  $q$ -periodic in  $s$ . Equation (7b) involves fields which are not observable *a priori*.

### 3.2 Gauge symmetries

The action (2) is invariant under several types of gauge symmetries :

- (i) by diffeomorphisms of  $\mathcal{Y}^5$  which preserves the orientation and the map  $\mathcal{Y}^5 \xrightarrow{P} \mathcal{X}^4$ , i.e. of the form  $T(x, y) = (x, f(x, y))$ , through pull-back  $(\theta, \pi) \mapsto (T^*\theta, T^*\pi)$ ;
- (ii) by gauge transformations  $\theta \mapsto \theta + dV$ , where  $V$  is a function of  $x \in \mathcal{X}^4$ ;
- (iii) by transformations  $\pi \mapsto \pi + \psi$ , where  $\psi$  is any closed 3-form on  $\mathcal{X}^4$  which decays at infinity.

## 4 Non Abelian gauge fields

The previous theory can be extended to non linear gauge theories. We let  $\mathfrak{G}$  be a compact simply connected Lie group of dimension  $r$  and we replace  $\mathcal{Y}^5$  by a manifold

$\mathcal{Y}^{N+1}$  of dimension  $N + 1 = 4 + r$ . We still assume the existence of a smooth projection map  $\mathcal{Y}^{N+1} \xrightarrow{P} \mathcal{X}^4$  such that  $dP$  has maximal rank and, for any  $x \in \mathcal{X}^4$ ,  $f_x := P^{-1}(x)$  is a *connected* submanifold of  $\mathcal{Y}^{N+1}$  of dimension  $r$ . Local coordinates on  $\mathcal{Y}^{N+1}$  are  $z = (z^I)_{0 \leq I \leq N} = (x, y) = (x^\mu, y^i)_{0 \leq \mu \leq 3 < i \leq N}$ , where  $(x^\mu)_{0 \leq \mu \leq 3}$  are coordinates on  $\mathcal{X}^4$  as previously.

Let  $\mathfrak{g}$  be the Lie algebra of  $\mathfrak{G}$ ,  $(\mathbf{t}_i)_{3 < i \leq N}$  a basis of  $\mathfrak{g}$  and  $c_{jk}^i$  the structure coefficients such that  $[\mathbf{t}_i, \mathbf{t}_j] = c_{ij}^k \mathbf{t}_k$ . We note  $\mathfrak{g}^*$  the dual space of  $\mathfrak{g}$  and  $(\mathbf{t}^i)_{3 < i \leq N}$  its dual basis. A consequence of the compactness of  $\mathfrak{G}$  is that  $\mathfrak{g}$  is *unimodular*, which reads

$$c_{ij}^i = c_{ji}^i = 0 \quad (9)$$

We let  $\mathbf{k}$  be a scalar product on  $\mathfrak{g}$  which invariant by the adjoint action of  $\mathfrak{G}$  and we set  $\mathbf{k}_{ij} = \mathbf{k}(\mathbf{t}_i, \mathbf{t}_j)$ . Fields are pairs  $(\theta, \pi)$  where

$$\begin{aligned} \theta &= \theta^i \mathbf{t}_i = (\theta^i_\mu(z) dx^\mu + \theta^i_j(z) dy^j) \mathbf{t}_i \\ \pi &= \pi_i \mathbf{t}^i = \frac{1}{(N-1)!} \pi_{iI_1 \dots I_{N-1}}(z) dz^{I_1} \wedge \dots \wedge dz^{I_{N-1}} \mathbf{t}^i \end{aligned}$$

As for Maxwell case we assume that the rank of  $(e^a, \theta^i)_{0 \leq a \leq 3 < i \leq N}$  is maximal, equal to  $N + 1$ . Hence by introducing the notations

$$\bar{\theta}_{i_1 \dots i_\alpha}^{(r-\alpha)} = \frac{1}{(r-\alpha)!} \epsilon_{i_1 \dots i_\alpha j_{\alpha+1} \dots j_r} \theta^{j_{\alpha+1}} \wedge \dots \wedge \theta^{j_r} \quad (10)$$

for  $0 \leq \alpha \leq r$  and where all indices run from 4 to  $N$ , we instead decompose  $\pi$  as

$$\pi_i = \frac{1}{2} \pi_i^{ab} e_{ab}^{(2)} \wedge \bar{\theta}^{(r)} - \pi_i^{ak} e_a^{(3)} \wedge \bar{\theta}_k^{(r-1)} + \frac{1}{2} \pi_i^{jk} e^{(4)} \wedge \bar{\theta}_{jk}^{(r-2)} \quad (11)$$

where coefficients  $\pi_i^{JK}$  are functions of  $z = (x, y)$ .

The action functional is

$$\mathcal{A}[\theta, \pi] = \int_{\mathcal{Y}^{N+1}} \frac{1}{2} \|\pi\|^2 e^{(4)} \wedge \bar{\theta}^{(r)} + \pi_i \wedge \Theta^i \quad (12)$$

where  $\|\pi\|^2 = \frac{1}{2} \pi_i^{ab} \pi^{i}_{ab}$  with  $\pi^{i}_{ab} = \mathbf{k}^{ij} \pi_j^{cd} \eta_{ac} \eta_{bd}$  and  $\Theta^i := d\theta^i + \frac{1}{2} [\theta \wedge \theta]^i = d\theta^i + \frac{1}{2} c_{jk}^i \theta^j \wedge \theta^k$ .

## 4.1 Euler–Lagrange equations

Using the decomposition  $\Theta^i = \frac{1}{2} \Theta^i_{ab} e^a \wedge e^b + \Theta^i_{ak} e^a \wedge \theta^k + \frac{1}{2} \Theta^i_{jk} \theta^j \wedge \theta^k$  and denoting  $d^\theta \pi_i = d\pi_i - c_{ki}^j \theta^k \wedge \pi_j$  the dynamical equations read

$$\begin{aligned} \pi^{i}_{ab} + \Theta^i_{ab} &= 0 & (a) \\ \Theta^i_{ak} &= 0 & (b) \\ \Theta^i_{jk} &= 0 & (c) \\ d^\theta \pi_i &= \frac{1}{2} \|\pi\|^2 e^{(4)} \wedge \bar{\theta}_i^{(r-1)} & (d) \end{aligned} \quad (13)$$

We first use (13c), which implies that, for any  $3 < i \leq N$  and for any  $x \in \mathcal{X}^4$ , the restriction of  $\Theta^i = d\theta^i + \frac{1}{2}c_{jk}^i\theta^j \wedge \theta^k$  on  $\mathfrak{f} = \mathfrak{f}_x$  vanishes. Hence by Frobenius' theorem (see [3]), given a point  $z \in \mathfrak{f}$ , we can construct a unique map  $g$  from a neighbourhood of  $z$  in  $\mathfrak{f}$  to  $\mathfrak{G}$  such that the restriction to  $\mathfrak{f}$  of  $\theta - g^{-1}dg$  vanishes and such that  $g(z) = 1$ . Since the rank of  $(\theta^i)_{3 < i \leq N}$  is  $r$ , this map is a local diffeomorphism. Actually the inverse map can be extended globally as a map  $T_{\mathfrak{f}} : \mathfrak{G} \rightarrow \mathfrak{f}$  which associates to any  $g \in \mathfrak{G}$  the end value  $v(1)$  of the path  $v : [0, 1] \rightarrow \mathfrak{f}$  which is a solution of  $v(0) = z$ ,  $e^a(\frac{dv}{dt}) = 0$  for  $0 \leq a \leq 3$  and  $\theta(\frac{dv}{dt}) = u^{-1}\frac{du}{dt}$  where  $u : [0, 1] \rightarrow \mathfrak{G}$  is a path such that  $u(0) = 1_{\mathfrak{G}}$  and  $u(1) = g$ . Indeed the definition of  $T_{\mathfrak{f}}(g)$  does not depend on the choice of the path  $u$  since  $\mathfrak{G}$  is simply connected.  $T_{\mathfrak{f}}$  is then a covering map of  $\mathfrak{f}$ . Hence since  $\mathfrak{G}$  is compact,  $\mathfrak{f}$  is compact and is diffeomorphic to a quotient  $\mathfrak{G}_{\mathfrak{f}}$  of  $\mathfrak{G}$  by a finite subgroup.

But all fibers are diffeomorphic to the same group  $\mathfrak{G}_0$ . Indeed for any fixed  $\xi = \xi^a \mathfrak{t}_a$  in  $\mathfrak{g}$ , consider the vector field  $\mathbf{X} = \mathbf{X}(\xi)$  on  $\mathcal{Y}^{N+1}$  defined by  $e^a(\mathbf{X}) = \xi^a$  and  $\theta^i(\mathbf{X}) = 0$ . Then by (6) which, by setting  $\gamma^a_b = \gamma^a_{bc}e^c$ , reads  $de^a = \gamma^a_{bc}e^b \wedge e^c$ , the Lie derivative of  $e^a$  by  $\mathbf{X}$  satisfies  $L_{\mathbf{X}}e^a = 0 \bmod [e^b]$ , for any  $0 \leq a, b \leq 3$ . This implies that the image of a fiber  $\mathfrak{f}$  by the flow map of  $\mathbf{X}$  is another fiber. We can thus construct a diffeomorphism between  $\mathfrak{G}_{\mathfrak{f}} \times B^4(0, \varepsilon) \simeq \mathfrak{f} \times B^4(0, \varepsilon)$  (where  $B^4(0, \varepsilon)$  is the ball of a sufficiently small radius  $\varepsilon$  in  $\mathbb{R}^4$ ) and a neighbourhood of a leaf  $\mathfrak{f}$  in  $\mathcal{Y}^{N+1}$  by mapping  $(z, \xi)$  to  $e^{\mathbf{X}(\xi)}(z)$ . Its inverse map provides us with a local trivialization of  $\mathcal{Y}^{N+1}$ . Since this construction can be done everywhere,  $\mathcal{Y}^{N+1} \rightarrow \mathcal{X}^4$  is endowed with a structure of principal bundle with a structure group  $\mathfrak{G}_0$ .

By choosing a local trivialization of this bundle we can improve the use of Equation (13c) to show the existence of a map  $g$  from an open subset of  $\mathcal{Y}^{N+1}$  to  $\mathfrak{G}_0$  such that the restriction of  $\theta - g^{-1}dg$  to  $\mathfrak{f}$  vanishes for any fiber  $\mathfrak{f}$ . This means that, if we set  $\mathbf{A} = g\theta g^{-1} - dg g^{-1}$  and decompose  $\mathbf{A} = \mathbf{A}_a e^a + \mathbf{A}_i \theta^i$ , then  $\mathbf{A}_i = 0$ . Hence  $\theta_z = g^{-1}(z)\mathbf{A}_a(z)g(z)e^a + g^{-1}(z)dg_z$ .

Equation (13b) then reads  $\partial_k \mathbf{A}_a = 0$ , so that actually  $\mathbf{A}_a(z) = \mathbf{A}_a(x)$  and

$$\theta = g^{-1}\mathbf{A}_a(x)g e^a + g^{-1}dg. \quad (14)$$

Thus  $\mathcal{Y}^{N+1} \rightarrow \mathcal{X}^4$  is endowed with the connection form  $\mathbf{A}$ . We define  $\mathbf{F} = d\mathbf{A} + \frac{1}{2}[\mathbf{A} \wedge \mathbf{A}] = \frac{1}{2}\mathbf{F}_{ab}(x)e^a \wedge e^b$ , so that we have  $\Theta := \Theta^i \mathfrak{t}_i = g^{-1}\mathbf{F}g$ .

Let  $(S_j^i)$  be the matrix of the adjoint action of  $g$  on  $\mathfrak{g}$  in the basis  $(\mathfrak{t}_i)_{3 < i \leq N}$ , i.e. such that  $g\mathfrak{t}_j g^{-1} = S_j^i \mathfrak{t}_i$ . We define  $e^i = S_j^i e^j$  and, by using the same conventions as in (10),  $\bar{e}^{(r)}$ ,  $\bar{e}_i^{(r-1)}$  and  $\bar{e}_{ij}^{(r-2)}$ . Then by (9),  $\bar{e}^{(r)} = \bar{\theta}^{(r)}$ ,  $\bar{e}_i^{(r-1)} = (S^{-1})_i^j \bar{\theta}_j^{(r-1)}$  and  $\bar{e}_{ij}^{(r-2)} = (S^{-1})_i^k (S^{-1})_j^{\ell} \bar{\theta}_{k\ell}^{(r-2)}$ . We also set

$$\begin{aligned} p_i &= (S^{-1})_i^j \pi_j, & p_i^{ab} &= (S^{-1})_i^j \pi_j^{ab} \\ p_i^{ak} &= (S^{-1})_i^j S_{\ell}^k \pi_j^{a\ell}, & p_i^{jk} &= (S^{-1})_i^{\ell} S_m^j S_n^k \pi_{\ell}^{mn} \end{aligned}$$

Then

$$p_i = \frac{1}{2} p_i^{ab} e_{ab}^{(2)} \wedge \bar{e}^{(r)} - p_i^{ak} e_a^{(3)} \wedge \bar{e}_k^{(r-1)} + \frac{1}{2} p_i^{jk} e^{(4)} \wedge \bar{e}_{jk}^{(r-2)} \quad (15)$$

Then (13a) reads  $p^i{}_{ab} + \mathbf{F}^i{}_{ab} = 0$ , where  $p^i{}_{ab} = k^{ij}\eta_{ac}\eta_{bd}p_j{}^{cd}$ , or, by setting  $\mathbf{F}_i{}^{ab} = k_{ij}\eta^{ac}\eta^{bd}\mathbf{F}^j{}_{cd}$ ,

$$p_i{}^{ab} + \mathbf{F}_i{}^{ab} = 0 \quad (16)$$

Lastly, by defining  $d^{\mathbf{A}}p_i = dp_i - c_{ki}^j \mathbf{A}^k \wedge p_j$  and  $\|p\|^2 = \frac{1}{2}p^i{}_{ab}p_i{}^{ab}$ , Equation (13d) translates as

$$d^{\mathbf{A}}p_i = \frac{1}{2}\|p\|^2 e^{(4)} \wedge \bar{e}_i^{(r-1)} \quad (17)$$

The computation of  $d^{\mathbf{A}}p_i$  requires some further notations. We denote by  $\partial_a$  and  $\partial_i$  the operators such that, for any function  $f$ ,  $df = (\partial_a f)e^a + (\partial_i f)e^i$  and by  $\mathbf{A}^i{}_a$  the coefficients such that  $\mathbf{A}^i = \mathbf{A}^i{}_a e^a$ . As in Section 3 we let  $\gamma^a{}_b$  be the spin connection coefficients and we set  $\gamma^a{}_c = \gamma^a{}_{cb}e^b$ . Lastly we define

$$\begin{aligned} \partial_b^{\gamma, \mathbf{A}} p_i{}^{ab} &= \partial_b p_i{}^{ab} - \mathbf{A}^k{}_b c_{ki}^\ell p_\ell{}^{ab} + \gamma^a{}_{cb} p_i{}^{cb} + \gamma^b{}_{cb} p_i{}^{ac} \\ \partial_b^{\gamma, \mathbf{A}} p_i{}^{jb} &= \partial_b p_i{}^{jb} - \mathbf{A}^k{}_b c_{ki}^\ell p_\ell{}^{jb} + \mathbf{A}^k{}_b c_{k\ell}^j p_i{}^{\ell b} + \gamma^b{}_{cb} p_i{}^{jc} \end{aligned}$$

Then we obtain by using (6) and (9) that

$$\begin{aligned} d^{\mathbf{A}}p_i &= (\partial_b^{\gamma, \mathbf{A}} p_i{}^{ab} + \partial_k p_i{}^{ak}) e_a^{(3)} \wedge \bar{e}^{(r)} \\ &+ (\partial_b^{\gamma, \mathbf{A}} p_i{}^{jb} + \partial_k p_i{}^{jk} + \frac{1}{2}\mathbf{F}^j{}_{ab} p_i{}^{ab} + \frac{1}{2}c_{k\ell}^j p_i{}^{k\ell}) e^{(4)} \wedge \bar{e}_j^{(r-1)} \end{aligned} \quad (18)$$

By taking into account (16) we deduce that (17) is equivalent to

$$\begin{aligned} \partial_b^{\gamma, \mathbf{A}} \mathbf{F}_i{}^{ab} &= \partial_k p_i{}^{ak} \\ \partial_b^{\gamma, \mathbf{A}} p_i{}^{jb} + \partial_k p_i{}^{jk} + \frac{1}{2}c_{k\ell}^j p_i{}^{k\ell} &= \frac{1}{2}\|\mathbf{F}\|^2 \delta_i{}^j + \frac{1}{2}\mathbf{F}_i{}^{ab} \mathbf{F}^j{}_{ab} \end{aligned} \quad (19)$$

As in Maxwell theory a key point is to observe that the l.h.s. in the first equation in (19) does not depend of  $y$  (or  $g$ ). Hence since the fibers are compact,

$$\partial_b^{\gamma, \mathbf{A}} \mathbf{F}_i{}^{ab} = \frac{\int_{f_x} \partial_b^{\gamma, \mathbf{A}} \mathbf{F}_i{}^{ab} \bar{e}^{(r)}}{\int_{f_x} \bar{e}^{(r)}} = \frac{\int_{f_x} \partial_k p_i{}^{ak} \bar{e}^{(r)}}{\int_{f_x} \bar{e}^{(r)}} = \frac{\int_{f_x} d(p_i{}^{ak} \bar{e}_k^{(r-1)})}{\int_{f_x} \bar{e}^{(r)}} = 0 \quad (20)$$

Hence  $\mathbf{A}$  is a solution of Yang–Mills equations. The second equation in (19) involves fields which are not observable.

## 4.2 Gauge symmetries

The action (12) is invariant under the following gauge symmetries :

(i) by diffeomorphisms of  $\mathcal{Y}^{N+1}$  which preserves the orientation and the map  $\mathcal{Y}^{N+1} \xrightarrow{P} \mathcal{X}^4$ , i.e. of the form  $T(x, y) = (x, f(x, y))$ , through pull-back  $(\theta, \pi) \mapsto (T^*\theta, T^*\pi)$ ;

(ii) by gauge transformations

$$\begin{aligned} \theta &\longmapsto \text{Ad}_g \theta - dg g^{-1} \\ \pi &\longmapsto \text{Ad}_g^* \pi \end{aligned} \quad (21)$$



for any map  $g : \mathcal{X}^4 \rightarrow \mathfrak{G}$  and where  $\text{Ad}_g \theta = g \theta g^{-1}$  and, if  $\text{Ad}_g \mathbf{t}_j = S_j^i \mathbf{t}_i$ ,  $\text{Ad}_g^* \pi = (S^{-1})_i^j \pi_j \mathbf{t}^i$ ;

(iii) by transformations  $\pi \mapsto \pi + \chi$ , where  $\chi$  has the form

$$\chi = \chi_i^{bk} e_b^{(3)} \wedge \bar{\theta}_k^{(r-1)} + \frac{1}{2} \chi_i^{jk} e_b^{(4)} \wedge \bar{\theta}_{jk}^{(r-2)}$$

decays at infinity and satisfies  $\theta^i \wedge d^\theta \chi_i = 0$ .

## 5 Einstein–Yang–Mills model

The previous theory can be modified in order to couple gravitation with gauge fields. We consider the metric  $\mathbf{h}$  on  $\mathbb{R}^4 \times \mathfrak{g}$  such that  $\mathbb{R}^4 \perp \mathfrak{g}$ ,  $\mathbf{h}$  coincides with the Minkowski metric  $\eta_{ab}$  on  $\mathbb{R}^4$  and with  $\mathbf{k}$  on  $\mathfrak{g}$ . We still assume here that  $\mathfrak{G}$  is compact and simply connected and we still work on a higher dimensional 'space-time'  $\mathcal{Y}^{N+1}$ . However :

(i) We replace the fixed co-vierbein  $(e^a)_{0 \leq a \leq 3}$  by a dynamical one  $(\theta^a)_{0 \leq a \leq 3}$  and we assume that the dynamical fields  $(\theta^I)_{0 \leq I \leq N} = (\theta^a, \theta^i)_{0 \leq a \leq 3 < i \leq N}$  forms a co- $(N+1)$ -bein on  $\mathcal{Y}^{N+1}$ .

(ii) No more fibration of  $\mathcal{Y}^{N+1}$  nor the existence of  $\mathcal{X}^4$  is assumed *a priori*. Even better, we do not make assumption on its topology, beside the fact that  $\mathcal{Y}^{N+1}$  is oriented and connected. Instead we introduce extra auxiliary fields  $(\pi_a)_{0 \leq a \leq 3}$  which are  $(N-1)$ -forms and which play the role of Lagrange multipliers for creating a fibration.

(iii) We also introduce the field  $\varphi^I{}_J = \varphi^I{}_{J,K}(z) dz^K$  with coefficients in the Lie algebra  $so(\mathbb{R}^4 \oplus \mathfrak{g}, \mathbf{h})$  of the Lorentz group, i.e. such that  $\varphi^{IJ} + \varphi^{JI} = 0$ , where  $\varphi^I{}_J = \varphi^{IK} \mathbf{h}_{KJ}$ .

(iv) In the action we replace the term  $\frac{1}{2} \|\pi\|^2 e^{(4)} \wedge \bar{\theta}^{(r)}$  by the  $(N+1)$ -dimensional *Einstein–Palatini* density  $\frac{1}{2} \hat{\theta}_{IJ}^{(N-1)} \wedge \Phi^{IJ}$ , where  $\Phi^{IJ} := d\varphi^{IJ} + \varphi^I{}_K \wedge \varphi^{KJ}$  and  $\hat{\theta}^{(N+1)} = \theta^0 \wedge \dots \wedge \theta^N$  and, for  $0 \leq \alpha$ ,

$$\hat{\theta}_{I_0 \dots I_\alpha}^{(N-\alpha)} = \frac{1}{(N-\alpha)!} \epsilon_{I_0 \dots I_\alpha J_{\alpha+1} \dots J_N} \theta^{J_{\alpha+1}} \wedge \dots \wedge \theta^{J_N}$$

(v) Lastly we impose the constraint

$$\theta^a \wedge \theta^b \wedge \pi_c = \theta^a \wedge \theta^b \wedge \pi_i = 0 \quad (22)$$

*To summarize* : dynamical fields are the 1-forms  $(\theta^I)_{0 \leq I \leq N} = (\theta^a, \theta^i)_{0 \leq a \leq 3 < i \leq N}$ , of rank  $N+1$  everywhere,  $(\varphi^{IJ})_{0 \leq I, J \leq N}$  with  $\varphi^{IJ} + \varphi^{JI} = 0$  and the  $(N-1)$ -forms  $(\pi_I)_{0 \leq I \leq N} = (\pi_a, \pi_i)_{0 \leq a \leq 3 < i \leq N}$ . Constraint (22) is assumed, i.e.  $\theta^a \wedge \theta^b \wedge \pi_I = 0$ . The basic action is

$$\mathcal{A}_0[\theta, \varphi, \pi] = \int_{\mathcal{Y}^{N+1}} \frac{1}{2} \hat{\theta}_{IJ}^{(N-1)} \wedge \Phi^{IJ} + \pi_I \wedge \Theta^I \quad (23)$$

where

$$\begin{aligned} \Theta^a &:= d\theta^a && \text{for } 0 \leq a \leq 3 \\ \Theta^i &:= d\theta^i + \frac{1}{2} c_{jk}^i \theta^j \wedge \theta^k && \text{for } 3 < i \leq N \end{aligned}$$

We shall also incorporate a bare cosmological constant  $\Lambda_0$  and consider the action

$$\mathcal{A}_{\Lambda_0}[\theta, \varphi, \pi] = \mathcal{A}_0[\theta, \varphi, \pi] - \int_{\mathcal{Y}^{N+1}} \Lambda_0 \hat{\theta}^{(N+1)} \quad (24)$$

Constraint (22) can actually be rephrased through the decomposition

$$\pi_I = -\pi_I^{ak} \theta_a^{(3)} \wedge \bar{\theta}_k^{(r-1)} + \frac{1}{2} \pi_I^{jk} \theta^{(4)} \wedge \bar{\theta}_{jk}^{(r-2)} \quad (25)$$

(i.e.  $\pi_I^{ab}$  vanishes). We also use the decomposition

$$\Theta^I = \frac{1}{2} \Theta^I_{JK} \theta^J \wedge \theta^K \quad (26)$$

or  $\Theta^I = \frac{1}{2} \Theta^I_{ab} \theta^a \wedge \theta^b + \Theta^I_{ak} \theta^a \wedge \theta^k + \frac{1}{2} \Theta^I_{jk} \theta^j \wedge \theta^k$ .

## 5.1 Euler–Lagrange equations

We set  $d^\varphi \hat{\theta}_{IJ}^{(N-1)} := d\hat{\theta}_{IJ}^{(N-1)} - \varphi^K_I \wedge \hat{\theta}_{KJ}^{(N-1)} - \varphi^K_J \wedge \hat{\theta}_{IK}^{(N-1)}$ ,  $d^\theta \pi_a = d\pi_a$  if  $0 \leq a \leq 3$ ,  $d^\theta \pi_i = d\pi_i - c_{ki}^j \theta^k \wedge \pi_j$  if  $3 < i \leq N$ . Then the Euler–Lagrange equations of  $\mathcal{A}_{\Lambda_0}[\theta, \varphi, \pi]$  reads :

$$\begin{aligned} \Theta^I_{ak} &= 0 & (a) \\ \Theta^I_{jk} &= 0 & (b) \\ d^\varphi \hat{\theta}_{IJ}^{(N-1)} &= 0 & (c) \\ d^\theta \pi_I + \frac{1}{2} \hat{\theta}_{IJK}^{(N-2)} \wedge \Phi^{JK} - \Theta^K_{IJ} \pi_K^{\ell J} \hat{\theta}_\ell^{(N)} &= \Lambda_0 \hat{\theta}_I^{(N)} & (d) \end{aligned} \quad (27)$$

Consider a solution of this system. We endow  $\mathcal{Y}^{N+1}$  with the metric  $\mathbf{h}_{\mu\nu} dz^\mu \otimes dz^\nu := \mathbf{h}_{IJ} \theta^I \otimes \theta^J$  and we assume that  $(\mathcal{Y}^{N+1}, \mathbf{h})$  is *complete*, meaning that any geodesic curve is defined for all 'time'.

Equation (27c) is equivalent to  $d^\varphi \theta^I := d\theta^I + \varphi^I_J \wedge \theta^J = 0$  thanks to the identity  $d^\varphi \hat{\theta}_{IJ}^{(N-1)} = d^\varphi \theta^K \wedge \hat{\theta}_{IJK}^{(N-2)}$ . It means that the spin connection defined by  $\varphi^{IJ}$  is torsionfree and, hence, coincides with the Levi-Civita connection on  $(\mathcal{Y}^{N+1}, \mathbf{h})$ .

Equation (27b) implies first  $d\theta^a = 0 \bmod [\theta^b]$ , for any  $0 \leq a, b \leq 3$ . This allows us to apply Frobenius' theorem (see [3]) to prove that  $\mathcal{Y}^{N+1}$  is foliated by  $r$ -dimensional submanifolds (leaves)  $\mathfrak{f}$  such that the restriction of  $\theta^a$  to  $\mathfrak{f}$  vanishes, for any  $0 \leq a \leq 3$ . We then define  $\mathcal{X}$  to be the set of leaves of this foliation (not yet a manifold!).

By using the same reasoning as in Section 4 (replacing (13c) by (27b), which implies that the restriction to  $\mathfrak{f}$  of  $\Theta^i = d\theta^i + \frac{1}{2} c_{jk}^i \theta^j \wedge \theta^k$  vanishes) we prove that each leaf  $\mathfrak{f}$  is compact and diffeomorphic to a quotient  $\mathfrak{G}_\mathfrak{f}$  of  $\mathfrak{G}$  by a finite subgroup.

We then prove that the foliation forms actually a fibration. For any fixed  $\xi = \xi^a \mathbf{t}_a$ , consider the vector field  $\mathbf{X} = \mathbf{X}(\xi)$  on  $\mathcal{Y}^{N+1}$  defined by  $\theta^a(\mathbf{X}) = \xi^a$  and  $\theta^i(\mathbf{X}) = 0$ . By (27a,b) we have  $d\theta^a = \frac{1}{2} \Theta^a_{bc} \theta^b \wedge \theta^c$  and thus  $L_{\mathbf{X}} \theta^a = \Theta^a_{bc} \xi^b \theta^c$ . Hence  $L_{\mathbf{X}} \theta^a = 0 \bmod [\theta^b]$ , for any  $0 \leq a, b \leq 3$ . By reasoning as in Section 4 we can construct a local diffeomorphism between  $\mathfrak{G}_\mathfrak{f} \times B^4(0, \varepsilon) \simeq \mathfrak{f} \times B^4(0, \varepsilon)$  and a neighbourhood of  $\mathfrak{f}$  in  $\mathcal{Y}^{N+1}$ . The set  $\mathcal{X}^4 = \mathcal{X}$  can

thus be endowed with a structure of 4-dimensional manifold and  $\mathcal{Y}^{N+1}$  with a structure of principal bundle over  $\mathcal{X}$  with structure group  $\mathfrak{G}_0$ .

Again as in Section 4 a further use of (27b) allows to show the existence of a map  $g$  from an open subset of  $\mathcal{Y}^{N+1}$  to  $\mathfrak{G}_0$  and of a map  $\mathbf{A} = \mathbf{A}_a e^a$ , with  $\mathbf{A}_a = \mathbf{A}^i{}_a \mathbf{t}_i$ , such that  $\theta_z = g^{-1}(z) \mathbf{A}_a(z) e^a g(z) + g^{-1}(z) dg_z$  and, by (27a)  $\mathbf{A}_a$  depends only on  $x$  in  $\mathcal{X}^4$ , so that  $\mathbf{A} = g\theta g^{-1} - dg g^{-1}$  is a 1-form on  $\mathcal{X}^4$ . We define  $\mathbf{F} := d\mathbf{A} + \frac{1}{2}[\mathbf{A} \wedge \mathbf{A}] = \frac{1}{2}\mathbf{F}_{ab}\theta^a \wedge \theta^b$ .

We now let  $(S^I_J)_{0 \leq I, J \leq N}$  such that  $S^a_b = \delta^a_b$ ,  $S^a_j = S^i_b = 0$  and  $S^i_j \mathbf{t}_i = \text{Ad}_g(\mathbf{t}_j) = g\mathbf{t}_j g^{-1}$  (note that  $S^K_I S^L_J \mathbf{h}_{KL} = \mathbf{h}_{IJ}$  since  $\mathbf{k}$  is invariant by  $\text{Ad}_g$ ) and we introduce

$$\begin{aligned} e^I &= S^I_J \theta^J, \quad \text{i.e. } e^a := \theta^a \text{ and } e^i = S^i_j \theta^j \\ p_I &= (S^{-1})^J_I \pi_J \quad \text{i.e. } p_a := \pi_a \text{ and } p_i = (S^{-1})^j_i \pi_j \\ \omega^{IJ} &= S^I_K S^J_L \varphi^{KL} - dS^I_K S^J_L \mathbf{h}^{KL} \\ \Omega^{IJ} &= S^I_K S^J_L \Phi^{KL} \end{aligned}$$

We observe that  $\mathbf{F}^i = S^i_j \Theta^j$  and  $\Omega^{IJ} = d\omega^{IJ} + \omega^I{}_K \wedge \omega^{KJ}$ . We define  $\hat{e}^{(N+1)} := e^0 \wedge \dots \wedge e^N$  and  $\hat{e}_I^{(N)}$ ,  $\hat{e}_{IJ}^{(N-1)}$  and  $\hat{e}_{IJK}^{(N-2)}$  in the same way as previously. We note that, by (9),  $\hat{e}^{(N+1)} = \hat{\theta}^{(N+1)}$ ,  $\hat{e}_I^{(N)} = (S^{-1})^J_I \hat{\theta}_J^{(N)}$ , and  $\hat{e}_{IJK}^{(N-1)} = (S^{-1})^{I'}_I (S^{-1})^{J'}_J (S^{-1})^{K'}_K \hat{\theta}_{I'J'K'}^{(N-1)}$ .

By setting  $d^{\mathbf{A}} p_a = dp_a$  and  $d^{\mathbf{A}} p_i = dp_i - c^j_{ki} \mathbf{A}^k \wedge p_j$  (27d) translates as

$$d^{\mathbf{A}} p_I + \frac{1}{2} \hat{e}_{IJK}^{(N-2)} \wedge \Omega^{JK} - \mathbf{F}^J{}_{Ib} p_J{}^{jb} \hat{e}_j^{(N)} = \Lambda_0 \hat{e}_I^{(N)} \quad (28)$$

The computation of  $d^{\mathbf{A}} p_I$  leads to the same result as in the Yang–Mills case (18), by replacing index  $4 \leq i \leq N$  by  $0 \leq I \leq N$  and with the extra simplification that coefficients  $p_I{}^{ab}$  vanish. We observe that  $\frac{1}{2} \hat{e}_{IJK}^{(N-2)} \wedge \Omega^{JK} = -\text{Ein}(\mathbf{h})_I{}^J e_J^{(N)}$ , where  $\text{Ein}(\mathbf{h})_I{}^J = \text{Ric}(\mathbf{h})_I{}^J - \text{R}(\mathbf{h}) \delta_I{}^J$  is the Einstein tensor for the metric  $\mathbf{h}$  on  $\mathcal{Y}^{N+1}$ . Hence by splitting indices (28) thus gives

$$\begin{aligned} \text{Ein}(\mathbf{h})_a{}^b + \Lambda_0 \delta_a{}^b &= \partial_k p_a{}^{bk} & (a) \\ \text{Ein}(\mathbf{h})_i{}^b &= \partial_k p_i{}^{bk} & (b) \\ \text{Ein}(\mathbf{h})_a{}^j &= \partial_c^{\mathbf{A}} p_a{}^{jc} + \partial_k p_a{}^{jk} + \frac{1}{2} p_a{}^{kl} c_{kl}^j - \mathbf{F}^J{}_{ac} p_J{}^{jc} & (c) \\ \text{Ein}(\mathbf{h})_i{}^j + \Lambda_0 \delta_i{}^j &= \partial_c^{\mathbf{A}} p_i{}^{jc} + \partial_k p_i{}^{jk} + \frac{1}{2} p_i{}^{kl} c_{kl}^j & (d) \end{aligned} \quad (29)$$

where

$$\begin{aligned} \partial_c^{\mathbf{A}} p_a{}^{jc} &= \partial_c p_a{}^{jc} + \mathbf{A}^k{}_c c_{kl}^j p_a{}^{\ell c} \\ \partial_c^{\mathbf{A}} p_i{}^{jc} &= \partial_c p_i{}^{jc} - \mathbf{A}^k{}_c c_{ki}^{\ell} p^{\ell jc} + \mathbf{A}^k{}_c c_{kl}^j p_i{}^{\ell c} \end{aligned} \quad (30)$$

The key point now is that, on the one hand,  $\text{Ein}(\mathbf{h})_I{}^J$  is *constant* on any fiber  $\mathbf{f} = P^{-1}(x)$  and, on the other hand, since  $\mathbf{f}$  is compact without boundary,

$$\int_{\mathbf{f}} \partial_k p_I{}^{bk} \hat{e}^{(N+1)} = \int_{\mathbf{f}} d(p_I{}^{bk} \hat{e}_k^{(N)}) = 0$$

Hence by the same reasoning as in (20) we deduce from (29 a,b) that

$$\text{Ein}(\mathbf{h})_a{}^b + \Lambda_0 \delta_a{}^b = \text{Ein}(\mathbf{h})_i{}^b = 0 \quad (31)$$

(We also deduce *a posteriori* that  $\text{Ein}(\mathbf{h})_a{}^j = 0$  because of the symmetry of the Einstein tensor.)

The remaining task is to recognize that (31) is equivalent to the Einstein–Yang–Mills system on  $\mathcal{X}^4$  equipped with the metric  $\mathbf{g} = \eta_{ab}e^a \otimes e^b$  and the connection  $\mathbf{A}$ . This computation is performed (in the tensorial language) in [9] and [2]. We present it here using the vielbein formalism.

The spin connection coefficients on  $\mathcal{X}^4$  for the metric  $\mathbf{g}$  are  $\gamma^a{}_b = \gamma^a{}_{bc}e^c$  with

$$\gamma^a{}_{bc} = \frac{1}{2}(\Theta^a{}_{bc} - \eta^{ad}\eta_{be}\Theta^e{}_{dc} - \eta^{ad}\eta_{ce}\Theta^e{}_{db})$$

The relation between the  $\gamma^a{}_b$ 's and the spin connection  $\omega^{IJ} = \omega^I{}_K \mathbf{h}^{KJ}$  of  $\mathbf{h}$  on  $\mathcal{Y}^{N+1}$  is given by

$$\begin{aligned} \omega^a{}_b &= \gamma^a{}_b - \frac{1}{2}\mathbf{k}_{ij}\eta^{ac}\mathbf{F}^j{}_{cb}e^i \\ \omega^a{}_i &= \frac{1}{2}\mathbf{k}_{ij}\eta^{ac}\mathbf{F}^j{}_{bc}e^b \\ \omega^i{}_a &= \frac{1}{2}\mathbf{F}^i{}_{ab}e^b \\ \omega^i{}_j &= \frac{1}{2}c^i{}_{jk}(e^k - 2\mathbf{A}^k) \end{aligned} \tag{32}$$

This leads to the expression of the Einstein tensor of  $\mathbf{h}$  in terms of the Einstein tensor of  $\mathbf{g}$  :

$$\begin{aligned} \text{Ein}(\mathbf{h})_a{}^b &= \text{Ein}(\mathbf{g})_a{}^b - \frac{1}{2}\mathbf{F}^i{}_{ac}\mathbf{F}^i{}_{bc} + \frac{1}{4}(\|\mathbf{F}\|^2 + |c|^2)\delta_a{}^b \\ \text{Ein}(\mathbf{h})_i{}^a &= \frac{1}{2}\partial_b^{\gamma, \mathbf{A}}\mathbf{F}_i{}^{ab} \\ \text{Ein}(\mathbf{h})_i{}^j &= \frac{1}{4}\mathbf{F}_i{}^{ab}\mathbf{F}^j{}_{ab} - \frac{1}{4}c^k{}_{il}c^j{}_{km}\mathbf{k}^{\ell m} + \frac{1}{4}(\|\mathbf{F}\|^2 + |c|^2)\delta_i{}^j \end{aligned} \tag{33}$$

where  $\|\mathbf{F}\|^2 = \frac{1}{2}\mathbf{F}_i{}^{ab}\mathbf{F}^i{}_{ab}$ ,  $|c|^2 = \frac{1}{2}c^i{}_{\ell k}c^{\ell}{}_{ij}\mathbf{k}^{jk}$  and  $\partial_b^{\gamma, \mathbf{A}}\mathbf{F}_i{}^{ab} = \partial_b\mathbf{F}_i{}^{ab} - c^j{}_{ki}\mathbf{A}^k{}_b\mathbf{F}_j{}^{ab} + \mathbf{F}_i{}^{cb}\gamma^a{}_{cb} + \gamma^c{}_{bc}\mathbf{F}_i{}^{ab}$ . Hence (31) translates as

$$\begin{aligned} \text{Ein}(\mathbf{g})_a{}^b + \Lambda\delta_a{}^b &= \frac{1}{2}(\mathbf{F}^i{}_{ac}\mathbf{F}^i{}_{bc} - \frac{1}{2}\|\mathbf{F}\|^2\delta_a{}^b) \\ \partial_b^{\gamma, \mathbf{A}}\mathbf{F}_i{}^{ab} &= 0 \end{aligned} \tag{34}$$

where  $\Lambda = \Lambda_0 + \frac{1}{4}|c|^2$ . The first equation is the Einstein equation with a cosmological constant  $\Lambda$  and the stress-energy tensor of the gauge fields, the second one is the Yang–Mills equation on  $(\mathcal{X}^4, \mathbf{g})$ .

## 5.2 Gauge symmetries

The action (23) and the constraint (22) are invariant by orientation preserving diffeomorphisms  $T : \mathcal{Y}^{N+1} \rightarrow \mathcal{Y}^{N+1}$  acting on fields through pull-back  $(\theta, \varphi, \pi) \mapsto (T^*\theta, T^*\varphi, T^*\pi)$ . They are also invariant through the transformation  $(\theta^I, \varphi^{IJ}, \pi_J) \mapsto (S^I{}_J\theta^J, S^I{}_K S^J{}_L \varphi^{KL}, (S^{-1})^J{}_I \pi_J)$  where  $(S^I{}_J)$  is the matrix of  $\text{Ad}_g$ , for some  $g \in \mathfrak{G}$  which is constant (if  $g$  is not constant, the curvature 2-form  $\Phi^{IJ}$  does not transform in a tensorial way).

## 6 Einstein–Maxwell model

What changes if we replace the compact simply connected group  $\mathfrak{G}$  by  $U(1)$ ? The action in (23) becomes

$$\mathcal{A}_0[\theta, \varphi, \pi] = \int_{\mathcal{Y}^5} \frac{1}{2} \hat{\theta}_{IJ}^{(3)} \wedge \Phi^{IJ} + \pi_I \wedge d\theta^I \quad (35)$$

where  $\pi_I = -\pi_I^a \theta_a^{(3)}$ . Critical points of the action  $\mathcal{A}_{\Lambda_0}$  in (24) satisfy System (27), excepted that (27b) does not exist. This has no incidence for solving the system obtained by imposing that the restriction of  $\theta^a$  on  $\mathfrak{f}$  vanishes since the integral leaves are just curves. The key point is that  $U(1)$  is not simply connected. Hence we cannot conclude that the integral leaves are compact and form a fibration in general.

However we shall prove that, if we know that *at least one leaf closes*, then *all leaves close* and are diffeomorphic and we hence get the existence of a space-time  $\mathcal{X}^4$  and a fibration of  $\mathcal{Y}^5$  over it.

Let  $Y$  be the vector field such that  $\theta^a(Y) = 0$  for any  $0 \leq a \leq 3$  and  $\theta^4(Y) = 1$ . Assume that some integral leaf  $\mathfrak{f}_0$  closes. It means that there exists a map  $u : \mathbb{R} \rightarrow \mathcal{Y}^5$  which is a solution of  $\frac{du}{ds} = Y(u)$ , the image of which is  $\mathfrak{f}_0$  and which is periodic, i.e. that there exists  $q > 0$  such that  $u(t+q) = u(t)$ . W.l.g. we can assume that  $q$  is minimal. For any  $\xi = (\xi^a)_{0 \leq a \leq 3}$  in  $\mathbb{R}^4$  let  $X$  be the vector field defined by  $\theta^a(X) = \xi^a$  for  $0 \leq a \leq 3$  and  $\theta^4(X) = 0$ . Then  $[X, Y] = 0$  since  $d\theta^I(X, Y) = 0$  because of (27a). Consider its flow map  $e^X$  and, for a fixed  $t \in \mathbb{R}$ , the map  $v : \mathbb{R} \rightarrow \mathcal{Y}^5$  defined by  $v(s) = e^{tX}(u(s))$ . Since  $X$  and  $Y$  commute  $\frac{dv}{ds} = Y(v)$ . This shows that  $e^{tX}(\mathfrak{f}_0)$ , the image of  $\mathfrak{f}_0$  by  $e^{tX}$ , is an integral leaf  $\mathfrak{f}$  of  $X$ . Since  $\mathfrak{f}_0$  is compact and  $e^{tX}$  is continuous  $\mathfrak{f}$  is compact and hence does not intersect from  $\mathfrak{f}_0$  if  $t$  is sufficiently small. Since this works for all values of  $\xi$  and since  $\mathcal{Y}^5$  is connected, this endows  $\mathcal{Y}^5$  with a topological bundle structure. By proceeding as in Section 3 we can achieve the normalization (4) and then obtain a solution of the Einstein–Maxwell system by using arguments from Section 4 and 5.

By replacing  $\mathfrak{G}$  by, e.g.,  $U(1) \times SU(2) \times SU(3)$  the situation is similar : we do not obtain a fibration in general, but *we do* if *at least one leaf closes*.

## Conclusion

Models presented in Section 3 and 4 provides gauge theories which, in addition to the usual gauge symmetries, are invariant by the action of diffeomorphisms which respect a topological fibration, enforcing thus the similarity between gauge theories and General Relativity. Models in Section 5 and 6 unify gravity and gauge theories at the classical level without symmetry hypotheses. All these models are based on the introduction of auxiliary 'ghosts' fields (but not in the Faddeev–Popov sense), which are not observable at the classical level. One may ask whether these 'ghosts' fields could create physically observable phenomena in a quantum version of the theory.

An intriguing fact is that, although invariant by diffeomorphisms our Kaluza–Klein theory is not invariant by standard gauge transformation. This may be connected to the

difficulty to lift the theory on the bundle of orthonormal frames over space-time (or its spin cover) in the spirit of the *group manifold* approach as initiated in [11, 14] and as done in [6] and further extended in the presence of spinors in [12].

## Références

- [1] T. Appelquist, A. Chodos, and P.G.O. Freund. *Modern Kaluza-Klein theories*. Addison-Wesley Pub. Co., Boston, 1987.
- [2] D. Bleeker. *Gauge Theory and Variational Principles*. Addison-Wesley, Mass., 1981.
- [3] R. L. Bryant, S. S. Chern, R. B. Gardner, H. L. Goldschmidt, and P. A. Griffiths. *Exterior Differential Systems*. Springer Verlag, New-York, Paris, 1991.
- [4] F. Hélein. Multisymplectic formulation of Yang-Mills equations and Ehresmann connections. *Advances in Theoretical and Mathematical Physics*, 19(4) :805–835, 2015. arXiv:1406.3641.
- [5] F. Hélein. A variational principle for kaluza–klein types theories. *Advances in Theoretical and Mathematical Physics*, 24(2) :305–326, 2020. arXiv:1809.03375.
- [6] F. Hélein and D. Vey. Curved space-times by crystallization of liquid fiber bundles. *Foundations of Physics*, 47(1) :1–41, sep 2016. arXiv:1508.07765.
- [7] P. Jordan. Erweiterung der projektiven Relativitätstheorie. *Annalen der Physik*, 436(4-5) :219–228, 1947.
- [8] Th. Kaluza. On the unification problem in physics. *Sitzungsberichte Pruss. Acad. Sci.*, page 966, 1921. Reprinted in English in [1] and in arXiv:1803.08616.
- [9] R. Kerner. Generalization of the Kaluza-Klein theory for an arbitrary non-Abelian gauge group. *Ann. Inst. H. Poincaré*, 9(2) :143, 1968. in www.numdam.org.
- [10] O. Klein. Quantentheorie und fünfdimensionale Relativitätstheorie. *Zeitschrift für Physik*, 37(12) :895–906, dec 1926.
- [11] Y. Ne’eman and T. Regge. Gauge theory of gravity and supergravity on a group manifold. *La Rivista Del Nuovo Cimento*, 1(5) :1–43, 1978.
- [12] Jérémie Pierard de Maujouy. Dirac spinors on generalised frame bundles : a frame bundle formulation for einstein-cartan-dirac theory.
- [13] Y. Thiry. Les équations de la théorie unitaire de Kaluza. *Comptes Rendus Acad. Sci. Paris*, 226(216), 1948.
- [14] M. Toller. Classical field theory in the space of reference frames. *Il Nuovo Cimento B Series 11*, 44(1) :67–98, mar 1978.