



**HAL**  
open science

## The relationship between phenotypic variation among offspring and mother body mass in wild boar: evidence of coin-flipping?

Marlène Gamelon, Jean-Michel Gaillard, Eric Baubet, Sébastien Devillard, Ludovic Say, Serge Brandt, Olivier Gimenez

### ► To cite this version:

Marlène Gamelon, Jean-Michel Gaillard, Eric Baubet, Sébastien Devillard, Ludovic Say, et al.. The relationship between phenotypic variation among offspring and mother body mass in wild boar: evidence of coin-flipping?. *Journal of Animal Ecology*, 2013, 82 (5), pp.937-945. 10.1111/1365-2656.12073 . hal-03515135

**HAL Id: hal-03515135**

**<https://hal.science/hal-03515135v1>**

Submitted on 6 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**The relationship between phenotypic variation among offspring and mother body mass in wild boar: Evidence of coin-flipping?**

Journal:	<i>Journal of Animal Ecology</i>
Manuscript ID:	JAE-2012-00706.R1
Manuscript Type:	Standard Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	GAMELON, Marlène; Biometry and Evolutionary Biology Laboratory, Evolutionary Ecology of Populations Group; Office National de la Chasse et de la Faune Sauvage, GAILLARD, Jean-Michel; Biometry and Evolutionary Biology Laboratory, Evolutionary Ecology of Populations Group, BAUBET, Eric; Office National de la Chasse et de la Faune Sauvage, DEVILLARD, Sébastien; Biometry and Evolutionary Biology Laboratory, Evolutionary Ecology of Populations Group, SAY, Ludovic; Biometry and Evolutionary Biology Laboratory, Evolutionary Ecology of Populations Group, BRANDT, Serge; Office National de la Chasse et de la Faune Sauvage, GIMENEZ, Olivier; Centre d'Ecologie Fonctionnelle et Evolutive,
Key-words:	bet-hedging, phenotypic plasticity, repeatability, reproductive tactic, <i>Sus scrofa scrofa</i>



1083x812mm (72 x 72 DPI)

View Only

1 **The relationship between phenotypic variation among offspring and mother**  
2 **body mass in wild boar: Evidence of coin-flipping?**

3 Marlène Gamelon<sup>1,2,3,4</sup>, Jean-Michel Gaillard<sup>1</sup>, Eric Baubet<sup>2</sup>, Sébastien Devillard<sup>1</sup>, Ludovic  
4 Say<sup>1</sup>, Serge Brandt<sup>2</sup> and Olivier Gimenez<sup>3</sup>

5

6 <sup>1</sup>Université de Lyon, F-69000, Lyon ; Université Lyon 1 ; CNRS, UMR5558, Laboratoire de  
7 Biométrie et Biologie Evolutive, F-69622, Villeurbanne, France.

8 <sup>2</sup>Office National de la Chasse et de la Faune Sauvage, 2 Bis Rue des Religieuses, BP 19,  
9 52120 Châteauvillain, France

10 <sup>3</sup>Centre d'Ecologie Fonctionnelle et Evolutive, Unité Mixte de Recherche 5175, campus  
11 CNRS, 1919 route de Mende, 34293 Montpellier Cedex 5, France

12 <sup>4</sup>E-mail: marlene.gamelon@univ-lyon1.fr, corresponding author

13

14 **Running headline:** Coin-flipping in wild boar

15

16

17

18

19

20

21 **Summary**

22 **1.** In highly variable environments, the optimal reproductive tactics of iteroparous organisms  
23 should minimize variance in yearly reproductive success to maximize the long-term average  
24 reproductive success. To minimize among-year variation in reproductive success individuals  
25 can either minimize the variance in the number of offspring produced at each reproductive  
26 attempt (classical bet-hedging) or maximize the phenotypic diversity of offspring produced  
27 within or among reproductive attempts (coin-flipping).

28 **2.** From a long-term detailed study of an intensively exploited population facing a highly  
29 unpredictable environment, we identify a continuum of reproductive tactics in wild boar  
30 females depending on their body mass.

31 **3.** At one end, light females adjusted litter size to their body mass and produced highly  
32 similar-sized offspring within a litter. These females fitted the hypothesis of individual  
33 optimization commonly reported in warm-blooded species, which involves both an optimal  
34 mass and an optimal number of offspring for a given mother. At the other end of the  
35 continuum, heavy females produced litters of variable size including a mixture of heavy and  
36 light offspring within litters.

37 **4.** Prolific heavy wild boar females diversify the phenotype of their offspring, providing a first  
38 evidence for coin-flipping in a warm-blooded species.

39

40

41 **Key-words:** bet-hedging, phenotypic plasticity, repeatability, reproductive tactic, *Sus scrofa*

42 *scrofa*

## 43 Introduction

44 Bet-hedging (Murphy 1968, Slatkin 1974) refers to the way organisms increase their fitness  
45 by decreasing the variance of annual reproductive success (Philippi & Seger 1989). In  
46 unpredictable and variable environments, the life history tactic leading to the highest  
47 individual fitness involves minimizing among-year variation in reproductive success rather  
48 than consistently maximizing reproductive success each year (Slatkin 1974; Philippi & Seger  
49 1989). Bet-hedging includes two non-exclusive mechanisms, risk spreading and risk-  
50 minimizing (Seger & Brockman 1987). For instance, iteroparity corresponds to a risk-  
51 spreading tactic, which has been interpreted as a response to environmental variation  
52 (Charnov & Schaffer 1973, Schaffer 1974, Orzack & Tuljapurkar 1989, Gaillard & Yoccoz  
53 2003). Likewise, producing each year the same limited number of offspring (Olofsson, Ripa  
54 & Jonsén 2009) and adjusting reproductive effort to a low level relative to available resources  
55 to avoid jeopardizing survival (Richard *et al.* 2002, Gaillard & Yoccoz 2003, Hamel *et al.*  
56 2010, Nevoux *et al.* 2010, Jones 2011) correspond to a risk-minimizing life history tactic.  
57 However, besides varying the number of offspring produced over breeding attempts during  
58 their lifetime, mothers in long-lived iteroparous organisms may change the phenotype of these  
59 offspring by allocating differentially among offspring within a given reproductive attempt  
60 (see Köhl *et al.* 2007 for evidence of differential maternal allocation to male offspring  
61 according to litter composition in saiga antelope *Saiga tatarica*). Such a mechanism involving  
62 developmental plasticity among litter mates has been called coin-flipping by Kaplan &  
63 Cooper (1984). Few studies have reported a phenotypic plasticity in offspring traits (Table 1).  
64 Of 208 studies citing the Kaplan & Cooper's study on 12 January 2012, only 24 tested  
65 developmental plasticity of offspring phenotypic traits or diversification of developmental  
66 time. Most of them dealt with cold-blooded organisms as originally targeted by Kaplan &  
67 Cooper (1984). Only two studies dealt with warm-blooded species (house wrens, *Troglodytes*

68 *aedon*, and yellow-bellied marmots, *Marmota flaviventris*) and provided support for the coin-  
69 flipping hypothesis. In these cases, evidence for developmental plasticity of offspring was  
70 reported among litters (Table 1), leaving open the question of the possible existence of  
71 developmental plasticity within a litter.

72 In polytocous mammals, variation among females in phenotypic traits, genotype, and habitat  
73 quality leads offspring produced by different mothers within a cohort to be more dissimilar  
74 than litter mates, generating so-called family effects (*sensu* Gaillard *et al.* 1998, see Pettorelli  
75 & Durant 2007 for a case study on carnivores). The dependence of offspring traits to maternal  
76 condition (*i.e.*, condition-dependent state *sensu* McNamara & Houston 1996) has been  
77 repeatedly reported in mammals (see Clutton-Brock 1991 for a review) and corresponds to an  
78 individual optimization of the trade-off between size and number of offspring (Smith &  
79 Fretwell 1974, Lloyd 1987). While a trade-off between number and size of offspring is  
80 expected (e.g. Smith and Fretwell 1974, Stearns 1992; see Sæther & Heim 1993 in mammals),  
81 it is rarely observed because of heterogeneity in individual quality (van Noordwijk & de Jong  
82 1986). Thus, large females usually produce large litters of large offspring, whereas small  
83 females produce small litters of small offspring, leading to strong family effects. This is in  
84 apparent contradiction with the coin-flipping hypothesis involving a decoupling between  
85 female attributes and early development of offspring so that females should produce a mixture  
86 of large and small offspring. This discrepancy might explain why evidence for coin-flipping  
87 has not been yet investigated in large mammals for which a tight link between mother and  
88 offspring phenotypes is commonly reported (Clutton-Brock 1991).

89 In the present work we aim to fill the gap by using a long-term detailed study on a wild boar  
90 (*Sus scrofa scrofa*) population. Wild boar is an especially relevant model to test for coin-  
91 flipping because females produce large litter sizes (up to 14, see Servanty *et al.* 2007) at each  
92 reproductive attempt and face highly variable and unpredictable environmental conditions as

93 they mostly feed on forest mast, the abundance of which fluctuates among years (Servanty *et*  
94 *al.* 2007).

95 Using the long-term monitoring of the heavily hunted wild boar population at Châteauvillain-  
96 Arc-en-Barrois, we first tested for individual optimization predicting that large females should  
97 produce large litters while small females should produce small litters. We thus assessed the  
98 relationship between fetus mass and female body mass as well as the relationship between  
99 litter size and female body mass to check for the expected dependence between these two  
100 variables and female phenotype (Clutton-Brock 1991). As mass at birth markedly influences  
101 future survival (Stearns 1992, Roff 2002), especially in large mammals (Gaillard *et al.* 2000),  
102 individual optimization should also lead to marked family effects to allow large females to  
103 recruit more than small ones. To test explicitly the coin-flipping hypothesis, we investigated  
104 developmental plasticity by both evaluating intra- and inter-litter variance of offspring mass.  
105 Under the coin-flipping hypothesis, females should produce a mixture of heavy and light  
106 offspring within a litter.

107

108

109

110

111

112

113

114



## 115 **Materials and methods**

### 116 STUDY AREA AND DATA COLLECTION

117 We studied a wild boar population in northeastern France in the 11,000 ha forest of  
118 Châteauvillain-Arc-en-Barrois. The territory is administratively divided into a core area that  
119 covers 8,500 ha of national forest and a 2,500 ha surrounding area of private or communal  
120 forest. These forests are mainly composed of oak (*Quercus petraea*), beechnut (*Fagus*  
121 *sylvatica*) and hornbeam (*Carpinus betulus*). The climate is intermediate between continental  
122 and oceanic types. Wild boars are hunted each year between October and February. Between  
123 1995 and 2009, we recorded the dressed body mass (*i.e.*, body mass without digestive tract,  
124 heart, lungs, liver, reproductive tract and blood) of each harvested female. Data on female  
125 reproductive status were also collected based on the examination of uteri for the presence of  
126 embryos or fetuses. Litter size was recorded and each fetus was weighed, measured (crown-  
127 rump length, in millimeters), and sexed. This information was available for 319 females and  
128 1,743 fetuses collected during 14 hunting seasons.

129

### 130 LINKING MASS OF FETUSES AND GESTATION STAGE

131 Offspring mass is expected to differ markedly among litters. In particular, not all litters were  
132 at the same gestation stage because females were shot at different dates and also because  
133 mating of wild boar occurs throughout most of the year in the study population, with a birth  
134 peak in mid-April (Gamelon *et al.* 2011). We thus needed to standardize fetus mass by  
135 gestation stage. To assess the gestation stage of a given litter, the mean length of fetuses in the  
136 litter was first calculated. Then, we applied the model provided first by Henry (1968) to link  
137 gestation stage to mean fetus length across litters (gestation stage (in days) = 23.43+0.32\*  
138 mean length (in mm)). The strength of this relationship was very high ( $R^2 = 0.86$ ,  $N = 20$ ,

139 Henry 1968). The gestation stage was thus estimated from Henry (1968)'s model for each  
140 litter. We then regressed individual fetus mass of the entire dataset on the gestation stage and  
141 used the best model to standardize fetus mass at a given gestation stage. Because the  
142 relationship was quadratic (see Results), we corrected in all the following analyses the  
143 individual fetus mass by gestation stage and gestation stage<sup>2</sup> by including them as fixed  
144 effects.

145

#### 146 LINKING LITTER SIZE AND FEMALE BODY MASS

147 Under the hypothesis of individual optimization, large females should produce large litters of  
148 large offspring, whereas small females should produce small litters of small offspring. We  
149 thus first assessed whether larger females produced larger litters than smaller females by  
150 linking litter size to female body mass. We thus fitted three models (on a log-scale): a simple  
151 linear regression model, a quadratic regression model, and a segmented regression model  
152 (Toms & Lesperance 2003). AIC was used for model selection (Burnham & Anderson 2002).

153

#### 154 LINKING MASS OF FETUSES WITH FEMALE BODY MASS

155 We assessed whether large females produced larger offspring than small females by assessing  
156 possible effects of maternal mass on fetus mass. To do this, we used linear mixed models with  
157 normal error terms. We included individual fetus mass as the response variable and mother  
158 identity as a random effect. In addition, we accounted for differences in gestation stage among  
159 litters (by including a quadratic regression as fixed effects) and for other potentially  
160 confounding factors, namely hunting season, female body mass, litter size, and fetus sex. AIC  
161 was used for model selection (Burnham & Anderson 2002).

## 162 WITHIN-LITTER VARIABILITY IN FETUS MASS

163 To provide a measure of within-litter variation in mass, we calculated the coefficient of  
164 variation (CV) of fetus mass for each litter. To correct fetus mass by gestation stage, we used  
165 the quadratic relationship linking fetus mass and gestation stage (see the section “Linking  
166 mass of fetuses and gestation stage” for further details) and we standardized all fetuses at the  
167 same gestation stage of 110 days (110 days corresponds to the latest gestation stage observed  
168 in the dataset). To assess the effect of female body mass on the CV of fetus mass corrected for  
169 gestation stage, we fitted a linear regression between CV of fetus mass and female body mass.  
170 Moreover, we explicitly tested whether the variation in fetus body mass increased with female  
171 body mass by including variance functions into the different models tested. We used the  
172 “varPower” variance function (Pinheiro & Bates 2000) because of its flexibility. Models  
173 including variance functions only differ from models without variance function in that the  
174 within-group errors are allowed to be heteroscedastic. More precisely, the residual term of the  
175 regression linking fetus mass and female body mass followed a Gaussian distribution with  
176 mean zero and a variance that equals  $\sigma^2 \cdot \text{female body mass}^{2\delta}$  where  $\sigma$  is the standard  
177 residual standard deviation and  $\delta$  a parameter to be estimated. Including variance functions  
178 thus specifies that the variance in fetus mass changes with increasing female body mass  
179 (Pinheiro & Bates (2000); see Cleasby & Nakagawa (2011) for an application of variance  
180 functions). AIC was used for model selection (Burnham & Anderson 2002).

181 All these analyses were performed with R 2.12.2 (R Development Core Team 2011).

182

183

184

## 185 Results

### 186 LINKING MASS OF FETUSES AND GESTATION STAGE

187 As expected, the mass of fetuses increased with the gestation stage, on a quadratic way  
188 (gestation stage: slope = -17.175, SE = 0.371; p-value  $\ll$  0.01; gestation stage<sup>2</sup>: slope =  
189 0.209, SE = 0.003; p-value  $\ll$  0.01; R<sup>2</sup> = 0.974; Appendix S1).

190

### 191 LINKING LITTER SIZE AND FEMALE BODY MASS

192 The segmented regression linking litter size with maternal mass provided the best fit to data  
193 (AIC = 174.1; intercept = -1.29 (0.31); slope= 0.76 (SE: 0.08) before the threshold; R<sup>2</sup>=  
194 0.219) when compared to the linear (AIC = 178.2; intercept= -0.67 (SE: 0.26); slope= 0.59  
195 (SE: 0.06); R<sup>2</sup>= 0.204) and quadratic (AIC = 176.3; intercept= -6.23 (SE: 2.89); log(Female  
196 Body Mass)= 3.53 (SE: 1.50); log(Female Body Mass)<sup>2</sup>= -0.38 (SE: 0.19); R<sup>2</sup>= 0.214) models.  
197 The higher the female body mass is, the larger is the litter size until a threshold mass (Fig. 1).  
198 Above this threshold mass, litter size did not increase with body mass. This threshold  
199 corresponded to a female mass of 58.3 kg, at which litter size was about 6.2 fetuses.  
200 Nevertheless, the AIC of the three tested models were very close and the differences among  
201 models did not seem to be biological significant for most of the observed range in female  
202 body mass. There was indeed a quite high variation in litter size both above and below the  
203 threshold value. Moreover, there was a positive relationship between litter size and body  
204 mass, but this relationship became weaker (quadratic model) or stopped to become constant  
205 (piecewise model) as female body mass increases. The relationship between litter size and  
206 female body mass thus appears positive but highly variable, and there is a limit on how many  
207 offspring females can produce on average.

## 208 LINKING MASS OF FETUSES WITH FEMALE BODY MASS

209 The baseline model of variation in fetus mass only included the quadratic effects of gestation  
210 stage and the random effect of mother identity. The best model retained (AIC=16962.28)  
211 included in addition the fixed effects of hunting season, mother body mass (0.295 (SE:  
212 0.117)) and fetus sex (-11.538 (SE: 1.394)) (Table 2A; see Appendix S2 for parameter  
213 estimates). Fetus mass thus depended on maternal mass. The heavier the females are, the  
214 heavier the fetuses are (Fig. 2). Nevertheless, from a biological viewpoint, this relationship  
215 between fetus mass and female body mass was weak as shown by the small effect size  
216 (maternal mass effect=0.2952 (SE: 0.1172)). Moreover, we found no effect of litter size on  
217 fetus mass.

218

## 219 WITHIN-LITTER VARIABILITY IN FETUS MASS

220 The heavier the female body mass is, the higher is the CV of fetus mass (intercept: -3.276e-  
221 03; slope = 3.545e-04; SE = 8.478e-05; p-value << 0.01; Fig. 3), meaning that the difference  
222 in terms of fetus mass within a litter produced by a heavy female was higher than the  
223 difference within a litter produced by a light female.

224 By including variance function into the different mixed models tested (Table 2B), we found  
225 that the best model retained among all the models presented in Table 2 included effects of  
226 hunting season, of mother body mass and of fetus mass as fixed effects, of mother identity as  
227 random effect and included the variance function (AIC=16842.07; see Appendix S2 for  
228 parameter estimates). We therefore found a marked increase in the log-likelihood associated  
229 with the inclusion of the variance function. Consequently, in the best model retained the  
230 variance in fetus mass increases with female body mass. More precisely,  $\delta$  was estimated to

231 0.984 (with a 95% confidence interval of 0.812-1.156) meaning that the variance increases  
232 with the square of female body mass.

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

For Review Only

252 **Discussion**

253 We identified a continuum of reproductive tactics in wild boar females depending on their  
254 body mass. At one end, light females adjusted the size of their litters to their body mass. This  
255 was consistent with the individual optimization process involving a strong positive link  
256 between litter size and mother's phenotypic attributes usually reported in mammals (see  
257 Clutton-Brock 1991 and Gaillard *et al.* 2000 for reviews). However, while previous studies of  
258 large mammals have reported that mothers in better than average condition produce heavier  
259 offspring (e.g., Russell *et al.* 1981 and Holst, Killeen & Cullis 1986 for sheep *Ovis aries*;  
260 Blaxter & Hamilton 1980, Moore, Littlejohn & Cowie 1988 for red deer *Cervus elaphus*),  
261 body mass of female wild boar had a little influence on offspring mass. Fetus mass was  
262 mainly influenced by gestation stage, sex, and year. The influence of gestation stage is trivial  
263 and simply corresponds to fetal growth during gestation. Males were heavier than females, as  
264 expected for a polygynous and dimorphic species like wild boar (Glucksmann 1974). The  
265 marked among-year variation in fetus mass was also expected from the highly variable and  
266 unpredictable food resources wild boars face with. Finally, we did not detect any effect of  
267 litter size on fetus mass. This absence of any evidence for a trade-off between fetus mass and  
268 litter size is especially noteworthy. This indicates that these females adjusted their  
269 reproductive effort mainly through variation in litter size, and suggests that it pays more for  
270 females acquiring an energy surplus to produce more offspring than to produce heavier  
271 offspring. At this end of the continuum, we showed that light females produced fetuses that  
272 have very similar mass. Indeed, the CV of fetus mass was lower in litters produced by light  
273 females than in litters produced by heavy females, indicating little differences in fetus mass  
274 within litters produced by light females.

275 At the other end of the continuum, heavy females produced a quite large variation in litter size  
276 (from 3 to 11). Moreover, we showed that after a threshold body mass, additional body

277 reserves are not translated into additional offspring. This stabilization corresponds to 6 fetuses  
278 on average. This could involve diminishing returns with increasing already large litters. Such  
279 diminishing returns have been reported in several studies (e.g., Jordan & Brooks 2010 on  
280 guppies *Poecilia reticulata*). Such a diminishing return can arise either because the marginal  
281 benefits of continued effort decrease or because the marginal costs of further reproductive  
282 effort increase (Jordan & Brooks 2010). In other words, when diminishing returns occur, a  
283 doubling of reproduction provides less than a doubling of relative success (Frank & Slatkin  
284 1990). In our case, the litter size per-unit female body mass decreases as body mass increases,  
285 providing evidence for a marginal diminishing return. Therefore, heavy wild boar females did  
286 not show any evidence for an individual optimization of their reproductive effort. Such a total  
287 independence between female phenotypic attributes and size or number of offspring produced  
288 despite a twofold variation in litter size has not been yet reported in any mammalian species  
289 to our knowledge. Under the Lack model of optimal litter size (Lack 1948), selection should  
290 act on parents to maximize the number of offspring recruited, leading a trade-off to occur  
291 between number and size of offspring produced (Smith & Fretwell 1974, Lloyd 1987,  
292 Winkler & Wallin 1987). Producing a lot of unviable offspring is obviously not a sustainable  
293 reproductive tactic, and producing a lot of large fetuses was not possible. The reproductive  
294 tactic displayed by most females of large mammals in response to these constraints involves  
295 optimizing this number-size trade-off according to their condition (individual optimization,  
296 Pettifor, Perrins & McCleery 1988, McNamara & Houston 1996). In addition, we showed that  
297 CV of fetus mass was higher in litters produced by heavy females than in litters produced by  
298 light females, indicating marked differences in fetus mass within litters produced by heavy  
299 females. Such a pattern of variation in within-litter fetus mass variation according to female  
300 body mass provides support for coin-flipping in litters produced by heavy females.

301



302 We could thus identify a continuum of reproductive tactics in wild boar females. At one end,  
303 light females display an individual optimization tactic by producing a litter size depending on  
304 their body mass with fetus of similar mass. At the opposite end of the continuum, heavy  
305 females show a coin-flipping tactic by producing offspring with a highly diversified  
306 phenotype. The mating system in this wild boar population could provide a pathway for such  
307 a pattern related to coin-flipping. Indeed, recent works in the studied population have shown  
308 that the number of fathers within a litter increased with litter size (Devillard *et al.* unpublished  
309 data). Nevertheless, large litters are generally produced by heavy females (Fig. 1).  
310 Consequently, the high phenotypic diversity of offspring observed in litters produced by  
311 heavy females could thus result from multiple paternities.

312 Wild boar females, depending on their body mass, have thus different reproductive tactics,  
313 maybe facilitated by the fact that wild boar piglets are not dependent on their size to survive  
314 (Baubet *et al.* 1995) contrary to other large mammalian species of herbivores (Gaillard *et al.*  
315 2000). In absence of survival costs in small-sized piglets, wild boar females can produce a  
316 large range of offspring phenotypes. These females diversified the phenotype of their  
317 offspring, likely to minimize variance in reproductive success among years in the highly  
318 variable and unpredictable environment they faced with. Such a developmental plasticity  
319 might indeed allow females to recruit successfully in both mast years and years without any  
320 mast. Theoretical approaches have also shown that life cycle delays could increase fitness  
321 when environments are sufficiently variable (Tuljapurkar 1990; Tuljapurkar & Wiener 2000).  
322 Previous studies of coin-flipping (Table 1) have shown that a large range of organisms adjust  
323 the number, the birth timing, and the phenotype of their offspring to maximize the number of  
324 recruits in fluctuating environments. However, these empirical studies focusing on  
325 development rate, asynchrony of hatchling, or dispersal capacities of offspring most often  
326 dealt with cold-blooded invertebrates, fishes, amphibians and reptiles. Kaplan & Cooper

327 (1984) themselves, in their original paper, focused on cold-blooded organisms certainly  
328 because developmental plasticity of offspring in warm-blooded species is expected to be  
329 constrained by thermoregulation.

330 Wild boars are hunted in most European forests and have become short-lived animals despite  
331 of their large body size. The generation time in heavily hunted wild boar populations is  
332 around 2 years, which is closer to the turn-over of tit populations than to that of a 60 kg  
333 mammal (Servanty *et al.* 2011). Such unusual demographic patterns have led wild boar  
334 females to have only one or two breeding attempts during their lifetime, starting to breed at  
335 one year of age (Gamelon *et al.* 2011) at a low threshold body mass (between 20 and 25 kg,  
336 Servanty *et al.* 2009). Consequently, wild boar females could exhibit different reproductive  
337 tactics during their life according to their body mass. An investigation of coin-flipping in an  
338 un-hunted wild boar population in which individuals start to breed at a higher threshold body  
339 mass would be required to test whether all heavy females display a coin-flipping tactic.

340

341

342

343

344

345

346

347

348

349 **Acknowledgments**

350 We are grateful to all those who helped capturing and marking wild boars, as well as those  
351 who helped collecting harvested wild boars, particularly P. Van den Bulck and G. Corbeau.  
352 We are grateful to the Office National des Forêts and to F. Jehlé, who allowed us to work on  
353 the study area. We warmly thank Tim Coulson, Marco Festa-Bianchet, Mark Hewison, Barry  
354 Sinervo and Shripad Tuljapurkar for their helpful comments on previous drafts. We also thank  
355 two anonymous referees and Atle Mysterud for constructive comments on previous drafts.

356

357

358

359

360

361

362

363

364

365

366

367

368

369 **References**

- 370 Armitage, K.B. (1986) Individuality, social behavior, and reproductive success in yellow-  
371 bellied marmots. *Ecology*, **67**, 1186–1193.
- 372
- 373 Baubet, E., Van Laere, G. & Gaillard, J-M. (1995) Growth and survival in piglets. *Ibex*  
374 *Journal of Mountain Ecology*, **3**, 71.
- 375
- 376 Blaxter, K.L. & Hamilton, W.J. (1980) Reproduction in farmed red deer. 2. calf growth and  
377 mortality. *The Journal of Agricultural Science*, **95**, 275–284.
- 378
- 379 Blouin-Demers, G. & Weatherhead, P.J. (2007) Allocation of offspring size and sex by female  
380 black ratsnakes. *Oikos*, **116**, 1759–1767.
- 381
- 382 Bowers, E.K., Sakaluk, S.K. & Thompson, C.F. (2011) Adaptive sex allocation in relation to  
383 hatching synchrony and offspring quality in house wrens. *The American Naturalist*, **177**, 617–  
384 629.
- 385
- 386 Brockmann, H.J. (2004) Variable life-history and emergence patterns of the Pipe-Organ mud-  
387 daubing wasp, *Trypoxylon politum* (Hymenoptera: *sphecidae*). *Journal of the Kansas*  
388 *Entomological Society*, **77**, 503–527.
- 389
- 390 Brown, C.A. (2004) Life histories of four species of scorpion in three families (*Buthidae*,  
391 *diplocentridae*, *vaejovidae*) from arizona and new mexico. *Journal of Arachnology*, **32**, 193–  
392 207.
- 393

- 394 Burnham, K.P. & Anderson, D.R. (2002) Model selection and multimodel inference: a  
395 practical information-theoretic approach. Springer.  
396
- 397 Charnov, E.L. & Schaffer, W.M. (1973) Life-History consequences of natural selection:  
398 Cole's result revisited. *The American Naturalist*, **107**, 791–793.  
399
- 400 Cleasby, I.R. & Nakagawa, S. (2011) Neglected biological patterns in the residuals a  
401 behavioural ecologist's guide to co-operating with heteroscedasticity. *Behavioral Ecology and*  
402 *Sociobiology*, **65**, 2361–2372.  
403
- 404 Clutton-Brock, T.H. (1991) The evolution of parental care. Princeton University Press.  
405
- 406 Dziminski, M.A. & Alford, R.A. (2005) Patterns and fitness consequences of intraclutch  
407 variation in egg provisioning in tropical australian frogs. *Oecologia*, **146**, 98–109.  
408
- 409 Dziminski, M.A. & Roberts, J.D. (2006) Fitness consequences of variable maternal  
410 provisioning in quacking frogs (*Crinia georgiana*). *Journal of Evolutionary Biology*, **19**, 144–  
411 155.  
412
- 413 Dziminski, M.A., Vercoe, P.E. & Roberts, J.D. (2009) Variable offspring provisioning and  
414 fitness: a direct test in the field. *Functional Ecology*, **23**, 164–171.  
415
- 416 Edmunds, P. J., Cumbo, V. & Fan, T. (2011) Effects of temperature on the respiration of  
417 brooded larvae from tropical reef corals. *The Journal of Experimental Biology*, **214**, 2783 –  
418 2790.

- 419 Frank, S.A. & Slatkin, M. (1990) Evolution in a variable environment. *The American*  
420 *Naturalist*, **136**, 244-260.
- 421
- 422 Gaillard, J.-M., Andersen, R., Delorme, D. & Linnell, J.D.C. (1998) Family effects on growth  
423 and survival of juvenile roe deer. *Ecology*, **79**, 2878–2889.
- 424
- 425 Gaillard, J.-M., Festa-Bianchet, M., Delorme, D. & Jorgenson, J. (2000) Body mass and  
426 individual fitness in female ungulates: bigger is not always better. *Proceedings of the Royal*  
427 *Society of London. Series B: Biological Sciences*, **267**, 471-477.
- 428
- 429 Gaillard, J.-M., & Yoccoz, N. G. (2003) Temporal variation in survival of mammals: A case of  
430 environmental canalization? *Ecology*, **84**, 3294–3306.
- 431
- 432 Gamelon, M., Besnard, A., Gaillard, J.-M., Servanty, S., Baubet, E., Brandt, S. & Gimenez,  
433 O. (2011) High hunting pressure selects for earlier birth date: wild boar as a case study.  
434 *Evolution*, **65**, 3100–3112.
- 435
- 436 Glucksmann, A. (1974) Sexual dimorphism in mammals. *Biological Reviews*, **49**, 423–475.
- 437
- 438 Goode, J.M., & Ewert, M.A. (2006) Reproductive trends in captive *Heosemys grandis*  
439 (*Geoemydidae*). *Chelonian Conservation and Biology*, **5**, 165–169.
- 440
- 441 Hakalahti, T., Häkkinen, H. & Valtonen, E.T. (2004) Ectoparasitic *Argulus coregoni*  
442 (Crustacea: *branchiura*) hedge their bets - studies on egg hatching dynamics. *Oikos*, **107**,  
443 295–302.

444

445 Halkett, F., Harrington, R., Hullé, M., Kindlmann, P., Menu, F., Rispe, C. &amp; Plantegenest, M.

446 (2004) Dynamics of production of sexual forms in aphids: Theoretical and experimental

447 evidence for adaptive “coin-flipping” plasticity. *The American Naturalist*, **163**, 112–125.

448

449 Hamel, S., Gaillard, J-M, Yoccoz, N.G., Loison, A., Bonenfant, C. &amp; Descamps, S. (2010)

450 Fitness costs of reproduction depend on life speed: empirical evidence from mammalian

451 populations. *Ecology Letters*, **13**, 915–935.

452

453 Henrich, S. (1988) Variation in offspring sizes of the poeciliid fish *Heterandria formosa* in454 relation to fitness. *Oikos*, **51**, 13–18.

455

456 Henry, V.G. (1968) Fetal development in european wild hogs. *The Journal of Wildlife*457 *Management*, **32**, 966–970.

458

459 Holst, P.J., Killeen, I.D. &amp; Cullis, B.R. (1986) Nutrition of the pregnant ewe and its effects on

460 gestation length, lamb birth weight and lamb survival. *Australian Journal Agricultural*461 *Research*, **37**, 647–655.

462

463 Jones, J.H. (2011) Primates and the evolution of long, slow life histories. *Current Biology*, **21**,

464 708–717.

465

466 Jordan, L.A. &amp; Brooks, R.C. (2010) The lifetime costs of increased male reproductive effort:

467 courtship, copulation and the Coolidge effect. *Journal of Evolutionary Biology*, **23**, 2403-

468 2409.

469

470 Kaplan, R.H. (1992) Greater maternal investment can decrease offspring survival in the frog  
471 *Bombina Orientalis*. *Ecology*, **73**, 280–288.

472

473 Kaplan, R.H., & Cooper, W.S. (1984) The evolution of developmental plasticity in  
474 reproductive characteristics: An application of the "Adaptive Coin-Flipping" principle. *The*  
475 *American Naturalist*, **123**, 393–410.

476

477 Koops, M.A., Hutchings, J.A. & Adams, B.K. (2003) Environmental predictability and the  
478 cost of imperfect information: influences on offspring size variability. *Evolutionary Ecology*  
479 *Research*, **5**, 29–42.

480

481 Kühl, A., Myrnerud, A., Erdnenov, G.I., Lushchekina, A.A., Grachev, I.A., Bekenov, A.B. &  
482 Milner-Gulland, E.J. (2007) The “big spenders” of the steppe: sex-specific maternal allocation  
483 and twinning in the saiga antelope. *Proceedings of the Royal Society of London. Series B:*  
484 *Biological Sciences*, **274**, 1293-1299.

485

486 Lack, D. (1948) The significance of Litter-Size. *Journal of Animal Ecology*, **17**, 45–50.

487

488 Lloyd, D.G. (1987) Selection of offspring size at independence and other Size-Versus-  
489 Number strategies. *The American Naturalist*, **129**, 800–817.

490

491 McKee, D. (1997) Intra-clutch offspring size variability in the freshwater zooplankter  
492 *Daphnia magna*. *Hydrobiologia*, **354**, 111–117.

493



- 494 McNamara, J.M. & Houston, A.I. (1996) State-dependent life histories. *Nature*, **380**, 215–  
495 221.
- 496
- 497 Meffe, G. K. (1987) Embryo size variation in mosquitofish: Optimality vs plasticity in  
498 propagule size. *Copeia*, **3**, 762–768.
- 499
- 500 Menu, F. & Debouzie, D. (1993) Coin-flipping plasticity and prolonged diapause in insects:  
501 example of the chestnut weevil *Curculio elephas* (Coleoptera: *curculionidae*). *Oecologia*, **93**,  
502 367–373.
- 503
- 504 Menu, F., Ginoux, M., Rajon, E., Lazzari, C.R. & Rabinovich, J.E. (2010) Adaptive  
505 developmental delay in chagas disease vectors: An evolutionary ecology approach. *PLoS*,  
506 *neglected tropical diseases*, **4**, e691.
- 507
- 508 Moore, G.H., Littlejohn, R.P. & Cowie, G.M. (1988) Factors affecting live weight gain in red  
509 deer calves from birth to weaning. *New Zealand journal of agricultural research*, **31**, 279–  
510 283.
- 511
- 512 Murphy, G.I. (1968) Pattern in life history and the environment. *The American Naturalist*,  
513 **102**, 391–403.
- 514
- 515 Nevoux, M., Forcada, J., Barbraud, C., Croxall, J. & Weimerskirch, H. (2010) Bet-hedging  
516 response to environmental variability, an intraspecific comparison. *Ecology*, **91**, 2416–2427.
- 517
- 518 Olofsson, H., Ripa, J. & Jonzén, N. (2009) Bet-hedging as an evolutionary game: the trade-off

- 519 between egg size and number. *Proceedings of the Royal Society B-Biological Sciences*, **277**,  
520 1149–1151.
- 521
- 522 Orzack, H.S., Tuljapurkar, S. (1989) Population dynamics in variable environments. The  
523 demography and evolution of iteroparity. *The American Naturalist*, **133**, 901–923.
- 524
- 525 Pettifor, R. A., Perrins, C.M. & McCleery, R.H. (1988) Individual optimization of clutch size  
526 in great tits. *Nature*, **336**, 160–162.
- 527
- 528 Pettorelli, N. & Durant, S.M. (2007) Family effects on early survival and variance in long-  
529 term reproductive success of female cheetahs. *Journal of Animal Ecology*, **76**, 908–914.
- 530
- 531 Philippi, T. & Seger, J. (1989) Hedging one's evolutionary bets, revisited. *Trends in Ecology*  
532 *& Evolution*, **4**, 41–44.
- 533
- 534 Pinheiro, J.C. & Bates, D.M. (2000) Mixed-effects models in S and S-PLUS. Springer, New  
535 York, USA.
- 536
- 537 R Development Core Team. (2011) R: A language and environment for statistical computing.  
538 R Foundation for Statistical Computing, Vienna, Austria.
- 539
- 540 Reed, D.H., & Nicholas, A.C. (2008) Spatial and temporal variation in a suite of life-history  
541 traits in two species of wolf spider. *Ecological Entomology*, **33**, 488–496.
- 542
- 543 Richard, A.F., Dewar, R.E., Schwartz, M. & Ratsirarson, J. (2002) Life in the slow lane?

- 544 Demography and life histories of male and female sifaka (*Propithecus verreauxi verreauxi*).  
545 *Journal of Zoology*, **256**, 421–436.
- 546
- 547 Roff, D.A. (2002) Life history evolution. Sinauer Associates Sunderland, Massachusetts.
- 548
- 549 Russel, A.J.F., Foot, J.Z., White, I.R. & Davies, G.J. (1981) The effect of weight at mating  
550 and of nutrition during Mid-Pregnancy on the birth weight of lambs from primiparous ewes.  
551 *The Journal of Agricultural Science*, **97**, 723–729.
- 552
- 553 Sæther, B.E. & Heim, M. (1993) Ecological correlates of individual variation in age at  
554 maturity in female moose (*Alces alces*) - The effects of environmental variability. *Journal of*  
555 *Animal Ecology*, **62**, 482-489.
- 556
- 557 Schaffer, W.M. (1974) Optimal reproductive effort in fluctuating environments. *The American*  
558 *Naturalist*, **108**, 783–790.
- 559
- 560 Seger, J. & Brockman, H.J. (1987) What is bet-hedging? *Oxford Surveys in Evolutionary*  
561 *Biology*, **4**, 182–211.
- 562
- 563 Servanty, S., Gaillard, J-M., Allaine, D., Brandt, S. & Baubet, E. (2007) Litter size and fetal  
564 sex ratio adjustment in a highly polytocous species: the wild boar. *Behavioral Ecology*, **18**,  
565 427–432.
- 566
- 567 Servanty, S., Gaillard, J-M., Ronchi, F., Focardi, S., Baubet, E. & Gimenez, O. (2011)  
568 Influence of harvesting pressure on demographic tactics: implications for wildlife

- 569 management. *Journal of Applied Ecology*, **48**, 835–843.
- 570
- 571 Slatkin, M. (1974) Hedging one's evolutionary bets. *Nature*, **250**, 704–705.
- 572
- 573 Smith, C. C. & Fretwell, S.D. (1974) The optimal balance between size and number of  
574 offspring. *The American Naturalist*, **108**, 499–506.
- 575
- 576 Stearns, S.C. (1992) The evolution of life histories, vol. 248. Oxford University Press Oxford.
- 577
- 578 Tanaka, K. (1995) Variation in offspring size within a population of the web-building spider  
579 *Agelena limbata*. *Researches on Population Ecology*, **37**, 197–202.
- 580
- 581 Thumm, K., & Mahony, M. (2002) Hatching dynamics and bet-hedging in a temperate frog,  
582 *Pseudophryne australis* (Anura: myobatrachidae). *Amphibia-Reptilia*, **23**, 433–444.
- 583
- 584 Thumm, K., & Mahony, M. (2005) Is variable egg size the proximate cause of diversified  
585 bet-hedging in the hatching dynamics of the red-crowned toadlet (*Pseudophryne*  
586 *australis*)(anura: *Myobatrachidae*)? *Journal Information*, **61**, 9–19.
- 587
- 588 Thumm, K., & Mahony, M. (2006). The effect of water level reduction on larval duration in  
589 the red-crowned toadlet *Pseudophryne australis* (Anura: *myobatrachidae*): Bet-hedging or  
590 predictive plasticity? *Amphibia-Reptilia*, **27**, 11–18.
- 591
- 592 Toms, J. & Lesperance, M. (2003) Piecewise regression: a tool for identifying ecological  
593 thresholds. *Ecology*, **84**, 2034–2041.

594

595 Tuljapurkar, S. (1990) Delayed reproduction and fitness in variable environments.

596 *Proceedings of the National Academy of Sciences*, **87**, 1139–1143.

597

598 Tuljapurkar, S. & Wiener, P. (2000) Escape in time: stay young or age gracefully? *Ecological*599 *Modelling*, **133**, 143–159.

600

601 Van Noordwijk, A.J. &amp; de Jong, G. (1986) Acquisition and allocation of resources: their

602 influence on variation in life history tactics. *The American Naturalist*, **128**, 137–142.

603

604 Winkler, D.W. &amp; Wallin, K.. (1987) Offspring size and number: A life history model linking

605 effort per offspring and total effort. *The American Naturalist*, **129**, 708–720.

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621 **Table 1.** List of the 24 papers (out of 208 quoting Kaplan & Cooper (1984)'s paper on coin-  
622 flipping on 12 January 2012) including tests of phenotypic polymorphism among offspring  
623 produced by a given individual. The species considered in the study, the taxonomic order, the  
624 offspring trait measured, the test outcome (coin-flipping (yes) or no (no)), and the reference of  
625 the study are provided. The table is divided into two parts. Part **A** brings together studies  
626 showing a developmental plasticity of offspring while part **B** brings together studies showing  
627 diversification of developmental time. Warm-blooded organisms occur in bold.

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

Order	Species	Traits	Coin-flipping validated	Reference
<b>A</b>				
Anura	Quacking frog ( <i>Crinia georgiana</i> )	Eggs size within clutches	Yes	Dziminski, Vercoe & Roberts 2009
Anura	Quacking frog ( <i>Crinia georgiana</i> )	Eggs size within clutches	Yes	Dziminski & Roberts 2006
Anura	15 Australian frog species	Offspring provisioning within clutch	Yes	Dziminski & Alford 2005
Anura	Red-crowned toadlet ( <i>Pseudophryne australis</i> )	Offspring sizes	Yes	Thumm & Mahony 2005
Anura	Fire-bellied toad ( <i>Bombina orientalis</i> )	Eggs size	Yes	Kaplan 1992
Araneae	Wolf spider ( <i>Rabidosa punctulata</i> , <i>R. rabida</i> )	Offspring size among clutches	No	Reed & Nicholas 2008
Araneae	Web-building spider ( <i>Agelena limbata</i> )	Offspring size within and among clutches	Yes	Tanaka 1995
Chelonia	Giant Asian pond turtle ( <i>Heosemys grandis</i> )	Number and size of eggs among clutches	Yes	Goode & Ewert 2006
Cladocera	Freshwater zooplankter ( <i>Daphnia magna</i> )	Offspring sizes within clutches	No	McKee 1997
Cyprinodontiformes	Poeciliid fish ( <i>Heterandria formosa</i> )	Offspring size at birth	Yes	Henrich 1988
Cyprinodontiformes	Mosquitofish ( <i>Gambusia affinis</i> )	Embryo size	Yes	Meffe 1987
Homoptera	Bird cherry-oat aphid ( <i>Rhopalosiphum padi</i> )	Production of sexuals or parthenogenetic females	Yes	Halkett <i>et al.</i> 2004
<b>Rodentia</b>	<b>Yellow-bellied marmot (<i>Marmota flaviventris</i>)</b>	<b>Social behavior</b>	<b>Yes</b>	<b>Armitage 1986</b>

Salmoniformes	Brook trout ( <i>Salvelinus fontinalis</i> )	Eggs size	Yes	Koops, Hutchings & Adams 2003
Scleractinia	Scleractinian corals ( <i>Pocillopora damicornis</i> , <i>Seriatopora hustrix</i> , <i>Stylophora pistillata</i> )	Dispersal capacities of larvae	Yes	Edmunds, Cumbo & Fan 2011
Scorpiones	4 species of scorpion	Offspring size within clutches	No	Brown 2004
Squamata	Black ratsnakes ( <i>Elaphe obsoleta</i> )	Offspring size among and within clutches	No	Blouin-Demers & Weatherhead 2007
<b>B</b>				
Anura	Red-crowned toadlet ( <i>Pseudophryne australis</i> )	Developmental rates	Yes	Thumm & Mahony 2006
Anura	Red crowned toadlet ( <i>Pseudophryne australis</i> )	Embryonic development to the time of hatching	Yes	Thumm & Mahony 2002
Arguloidea	Parasitic crustacean ( <i>Argulus coregoni</i> )	Intra-clutch variability in hatching among eggs	Yes	Hakalahti, Häkkinen & Valtonen 2004
Coleoptera	Chestnut weevil ( <i>Curculio elephas</i> )	Duration of diapause	Yes	Menu & Debouzie 1993
Hemiptera	Triatominae ( <i>Triatominae</i> )	Developmental delays	Yes	Menu <i>et al.</i> 2010
Hymenoptera	Pipe-Organ Mud-daubing Wasp ( <i>Trypoxylon politum</i> )	Synchrony or asynchrony of emergence pattern	Yes	Brockmann 2004
Passeriformes	<b>House wrens (<i>Troglodytes aedon</i>)</b>	<b>Synchrony or asynchrony in hatching eggs and weight</b>	<b>Yes</b>	<b>Bowers, Sakaluk &amp; Thompson 2011</b>



646 **Table 2.** Model selection of linear mixed models fitted with individual fetus mass (*mass*) as a response variable and mother identity (*mother*) as a  
 647 random factor. The fixed effects correspond to gestation stage (*gestation*) and gestation stage<sup>2</sup> (*gestation*<sup>2</sup>), hunting season (*year*), fetus sex (*sex*),  
 648 mother body mass (*mother body mass*) and litter size (*litter size*). Displayed are the AIC of each model and the difference in AIC between each  
 649 candidate model and the best model ( $\Delta AIC$ ) for **(A)** all possible models presented in the column *Models* and for **(B)** the models presented in the  
 650 column *Models* with a variance function specifying that fetus mass was allowed to differ in relation to female body mass. The selected models  
 651 are in bold. Data come from wild boar females collected in the population of Châteauvillain-Arc-en-Barrois, France.

<b>Models</b>	<b>A</b>		<b>B</b>	
	<b>AIC</b>	<b><math>\Delta AIC</math></b>	<b>AIC</b>	<b><math>\Delta AIC</math></b>
mass~gestation+gestation <sup>2</sup> +year+mother body mass+litter size+sex+mother	16963.60	1.32	16843.23	1.16
mass~gestation+gestation <sup>2</sup> +year+mother body mass+litter size+mother	17028.32	66.04	16903.93	61.86
mass~gestation+gestation <sup>2</sup> +year+mother body mass+mother	17027.01	64.73	16902.73	60.66
<b>mass~gestation+gestation<sup>2</sup>+year+mother body mass+sex+mother</b>	<b>16962.28</b>	<b>0</b>	<b>16842.07</b>	<b>0</b>
mass~gestation+gestation <sup>2</sup> +year+sex+litter size+mother	16968.13	5.85	16847.06	4.99
mass~gestation+gestation <sup>2</sup> +year+sex+mother	16966.20	3.92	16845.09	3.02
mass~gestation+gestation <sup>2</sup> +year+ litter size+mother	17033.48	71.2	16908.19	66.12
mass~gestation+gestation <sup>2</sup> +year+mother	17031.58	69.3	16906.25	64.18
mass~gestation+gestation <sup>2</sup> +mother body mass+litter size+mother	17036.55	74.27	16910.19	68.12
mass~gestation+gestation <sup>2</sup> +mother body mass+sex+mother	16972.24	9.96	16850.11	8.04
mass~gestation+gestation <sup>2</sup> +mother body mass+litter size+sex+mother	16972.64	10.36	16850.33	8.26
mass~gestation+gestation <sup>2</sup> +mother body mass+mother	17036.16	73.88	16909.95	67.88
mass~gestation+gestation <sup>2</sup> +litter size+sex +mother	16974.56	12.28	16852.23	10.16
mass~gestation+gestation <sup>2</sup> +litter size+mother	17039.04	76.76	16912.53	70.46
mass~gestation+gestation <sup>2</sup> +sex+mother	16972.78	10.5	16850.47	8.4

mass~gestation+gestation<sup>2</sup>

17037.21

74.93

16910.72

68.65

---

For Review Only

652 **Figure legends**

653 **Figure 1.** Relationship between litter size and female body mass (on a log-scale) in the wild  
654 boar population of Châteauvillain- Arc-en-Barrois, France. The best fitting model (segmented  
655 regression, solid line) and the linear and quadratic relationships (dotted lines) are displayed.  
656 The vertical solid lines correspond to the threshold body mass estimated from the segmented  
657 regression and its associated confidence interval.

658 **Figure 2.** Relationship between standardized fetus mass (*i.e.*, corrected for gestation stage)  
659 and female body mass. The solid line corresponds to the estimates provided by the linear  
660 mixed model linking standardized fetus mass as a response variable, female body mass, fetus  
661 sex and hunting season as fixed effects and mother identity as a random effect (intercept:  
662 1005.166; slope: 0.295 (SE: 0.117)).

663 **Figure 3.** Relationship between the CV of fetus mass and female body mass from females  
664 collected in the wild boar population of Châteauvillain-Arc-en-Barrois, France (intercept: -  
665 3.276e-03; slope = 3.545e-04 (SE: 8.478e-05); p-value  $\ll$  0.01).

666

667

668


669

670

671

672

673 Fig. 1.




674

675

676

677

678 **Fig. 2.**



679

680

681


682

683

684

685 **Fig. 3.**

686



687

688


689

690

1 **Supporting Information**

2 **Appendix S1**

3 Relationship between fetus mass and gestation stage in the wild boar population of  
4 Châteauvillain-Arc-en-Barrois, France (gestation stage: slope = -17.175, SE = 0.371; p-value  
5  $\ll 0.01$ ; gestation stage<sup>2</sup>: slope = 0.209, SE = 0.003; p-value  $\ll 0.01$ ;  $R^2 = 0.974$ ).



6

7

8

## Appendix S2: Model selection of linear mixed models fitted with individual fetus mass (mass) as a response

	Intercept
<b>A)</b>	
mass~gestation+gestation <sup>2</sup> +year+mother body mass+litter size+sex+mother	380.634381
mass~gestation+gestation <sup>2</sup> +year+mother body mass+litter size+mother	374.019765
mass~gestation+gestation <sup>2</sup> +year+mother body mass+mother	372.598006
<b>mass~gestation+gestation<sup>2</sup>+year+mother body mass+sex+mother</b>	<b>379.211825</b>
mass~gestation+gestation <sup>2</sup> +year+sex+litter size+mother	386.581145
mass~gestation+gestation <sup>2</sup> +year+sex+mother	387.333558
mass~gestation+gestation <sup>2</sup> +year+ litter size+mother	380.227457
mass~gestation+gestation <sup>2</sup> +year+mother	381.145422
mass~gestation+gestation <sup>2</sup> +mother body mass+litter size+mother	366.037266
mass~gestation+gestation <sup>2</sup> +mother body mass+sex+mother	369.833861
mass~gestation+gestation <sup>2</sup> +mother body mass+litter size+sex+mother	372.879994
mass~gestation+gestation <sup>2</sup> +mother body mass+mother	362.987829
mass~gestation+gestation <sup>2</sup> +litter size+sex +mother	377.763622
mass~gestation+gestation <sup>2</sup> +litter size+mother	371.22766
mass~gestation+gestation <sup>2</sup> +sex+mother	375.960021
mass~gestation+gestation <sup>2</sup>	369.662583
<b>B)</b>	
mass~gestation+gestation <sup>2</sup> +year+mother body mass+litter size+sex+mother	377.1696
mass~gestation+gestation <sup>2</sup> +year+mother body mass+litter size+mother	371.4549
mass~gestation+gestation <sup>2</sup> +year+mother body mass+mother	370.2911
<b>mass~gestation+gestation<sup>2</sup>+year+mother body mass+sex+mother</b>	<b>375.9730</b>
mass~gestation+gestation <sup>2</sup> +year+sex+litter size+mother	382.5391
mass~gestation+gestation <sup>2</sup> +year+sex+mother	382.9819
mass~gestation+gestation <sup>2</sup> +year+ litter size+mother	376.9891
mass~gestation+gestation <sup>2</sup> +year+mother	377.5949
mass~gestation+gestation <sup>2</sup> +mother body mass+litter size+mother	364.3586
mass~gestation+gestation <sup>2</sup> +mother body mass+sex+mother	367.4411
mass~gestation+gestation <sup>2</sup> +mother body mass+litter size+sex+mother	370.1793
mass~gestation+gestation <sup>2</sup> +mother body mass+mother	361.6615
mass~gestation+gestation <sup>2</sup> +litter size+sex +mother	375.0776
mass~gestation+gestation <sup>2</sup> +litter size+mother	369.4921
mass~gestation+gestation <sup>2</sup> +sex+mother	373.2981
mass~gestation+gestation <sup>2</sup>	367.9610



response variable and mother identity (mother) as a random factor. The fixed effects correspond to ge

SE	Gestation	SE	Gestation <sup>2</sup>	SE	Year 1997	SE
24.298214	-17.460414	0.649493	0.210740	0.004557	1.667687	18.481280
24.203270	-17.449995	0.647200	0.210706	0.004541	1.360027	18.430091
24.121921	-17.458280	0.646658	0.210823	0.004535	1.575622	18.415428
24.217388	-17.468858	0.648948	0.210859	0.004551	1.883398	18.466631
24.410895	-17.348037	0.654100	0.210455	0.004599	1.368698	18.652701
24.183555	-17.339894	0.652306	0.210400	0.004587	1.280038	18.622173
24.337748	-17.333151	0.652400	0.210412	0.004587	1.035246	18.617337
24.111217	-17.323336	0.650661	0.210346	0.004575	0.926916	18.587900
21.871543	-17.363858	0.639170	0.209844	0.004505	NA	NA
21.866588	-17.385594	0.642490	0.210066	0.004527	NA	NA
21.986713	-17.373723	0.642152	0.209896	0.004526	NA	NA
21.751139	-17.375603	0.639511	0.210013	0.004506	NA	NA
21.953383	-17.293472	0.643980	0.209704	0.004547	NA	NA
21.85719	-17.27853	0.64152	0.20964	0.00453	NA	NA
21.585537	-17.307261	0.642423	0.209795	0.004536	NA	NA
21.488799	-17.290402	0.639943	0.209719	0.004519	NA	NA
24.088181	-17.3991	0.647400	0.2103	0.004557	3.1401	18.237696
23.978827	-17.3927	0.644850	0.2102	0.004540	2.8873	18.168100
23.929573	-17.4071	0.644307	0.2104	0.004534	3.1400	18.156731
24.040415	-17.4138	0.646914	0.2104	0.004551	3.3955	18.227937
24.162234	-17.3013	0.650873	0.2101	0.004590	3.4599	18.372177
23.978335	-17.2942	0.648613	0.2100	0.004575	3.4212	18.340809
24.064796	-17.2924	0.648641	0.2100	0.004575	3.2432	18.310563
23.883754	-17.2828	0.646425	0.2100	0.004561	3.1894	18.280356
21.565384	-17.2938	0.633440	0.2093	0.004480	NA	NA
21.616893	-17.3171	0.637273	0.2095	0.004504	NA	NA
21.695588	-17.2992	0.636818	0.2093	0.004503	NA	NA
21.485677	-17.3115	0.633831	0.2095	0.004481	NA	NA
21.656554	-17.2280	0.638784	0.2092	0.004524	NA	NA
21.538639	-17.2195	0.635754	0.2091	0.004503	NA	NA
21.328217	-17.2443	0.637000	0.2093	0.004491	NA	NA
21.210805	-17.2335	0.633926	0.2092	0.004491	NA	NA







nong female body bass. Displayed are the estimates and the associated SE for each fixed effect for all i

SE	Year 2009	SE	Mother body mass	SE	Litter size
15.619808	-26.341507	10.512601	0.336511	0.135132	-0.765974
15.569515	-26.756219	10.478344	0.351154	0.134593	-0.761081
15.557201	-26.979810	10.467356	0.306138	0.122347	NA
15.607521	-26.566817	10.501591	0.291162	0.122808	NA
15.757725	-24.087746	10.573349	NA	NA	0.215009
15.734505	-23.912606	10.533042	NA	NA	NA
15.721065	-24.408989	10.548713	NA	NA	0.261294
15.698815	-24.196305	10.509310	NA	NA	NA
NA	NA	NA	0.282717	0.133937	-1.200203
NA	NA	NA	0.193927	0.122168	NA
NA	NA	NA	NA	0.134642	-1.203652
NA	NA	NA	0.211577	0.121554	NA
NA	NA	NA	NA	NA	-0.409210
NA	NA	NA	NA	NA	-0.35464
NA	NA	NA	NA	NA	NA
NA	NA	NA	NA	NA	NA
15.447044	-23.3331	10.499089	0.3273	0.138936	-0.8522
15.380652	-23.5533	10.460556	0.3385	0.138557	-0.8335
15.372358	-23.8115	10.451039	0.2847	0.124049	NA
15.440075	-23.5974	10.490455	0.2725	0.124459	NA
15.556002	-21.1073	10.533739	NA	NA	0.1489
15.530104	-20.9634	10.485826	NA	NA	NA
15.496929	-21.2527	10.500213	NA	NA	0.2027
15.471946	-21.0580	10.452921	NA	NA	NA
NA	NA	NA	0.2856	0.957919	-1.2620
NA	NA	NA	0.1892	0.123598	NA
NA	NA	NA	0.2716	0.138223 1422	-1.2772
NA	NA	NA	0.2039	0.123032	NA
NA	NA	NA	NA	NA	-0.4280
NA	NA	NA	NA	NA	-0.3685
NA	NA	NA	NA	NA	NA
NA	NA	NA	NA	NA	NA

## the models . The selecter

SE	Sex	SE
0.948527	-11.539036	1.394038
0.944588	NA	NA
NA	NA	NA
NA	-11.538093	1.394039
0.870981	-11.589028	1.394132
NA	-11.590034	1.394016
0.868395	NA	NA
NA	NA	NA
0.951927	NA	NA
NA	-11.468746	1.394665
0.957026	-11.468034	1.394547
NA	NA	NA
0.872107	-11.514745	1.394530
0.86838	NA	NA
NA	-11.508937	1.394466
NA	NA	NA
0.958618	-9.9865	1.245208
0.954961	NA	NA
NA	NA	NA
NA	-9.9847	1.245326
0.865608	-10.0170	1.245028
NA	-10.0175	1.244922
0.862319	NA	NA
NA	NA	NA
0.957919	NA	NA
NA	-9.9093	1.244814
0.962904	-9.9099	1.244577
NA	NA	NA
0.864402	-9.9438	1.244606
0.859931	NA	NA
NA	-9.9381	1.244600
NA	NA	NA