



Estimation of sex-specific survival with imperfect clues about sex

Roger Pradel, Lory Maurin-Bernier, Olivier Gimenez, Meritxell Genovart, Rémi Choquet, Daniel Oro

► To cite this version:

Roger Pradel, Lory Maurin-Bernier, Olivier Gimenez, Meritxell Genovart, Rémi Choquet, et al.. Estimation of sex-specific survival with imperfect clues about sex. Canadian Journal of Statistics, 2007, 36 (1), pp.29-42. <10.1002/CJS.5550360105>. <hal-03515267>

HAL Id: hal-03515267

<https://hal.science/hal-03515267v1>

Submitted on 6 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Estimation of sex-specific survival with imperfect clues about sex

ROGER PRADEL, LORY MAURIN-BERNIER, OLIVIER GIMENEZ,
MERITXELL GENOVART, RÉMI CHOQUET, AND DANIEL ORO

February 5, 2007

Abstract

In monomorphic species, determination of sex from behavior is prone to errors. Ignoring this fact renders problematic the estimation of sex-specific parameters. We develop capture-recapture survival models that account for uncertainty in the assessment of sex. There is a legitimate concern that some parameters may not be identifiable in these models. We examine parameter redundancy for four basic models with survival and encounter probabilities constant or time-dependent. We further develop a more refined and more appropriate model for an Audouin's gull data set where four distinct behavioral clues have been used. We examine how useful it is to incorporate the least reliable of the clues and the genetic determination of sex available for only a handful of individuals. We finally discuss the implications of our findings for the design of field studies as well as further directions of research.

KEY WORDS: capture-recapture; parameter redundancy; multievent models.

Estimation of sex-specific survival with imperfect clues about sex

February 5, 2007

Abstract

In monomorphic species, determination of sex from behavior is prone to errors. Ignoring this fact renders problematic the estimation of sex-specific parameters. We develop capture-recapture survival models that account for uncertainty in the assessment of sex. There is a legitimate concern that some parameters may not be identifiable in these models. We examine parameter redundancy for four basic models with survival and encounter probabilities constant or time-dependent. We further develop a more refined and more appropriate model for an Audouin's gull data set where four distinct behavioral clues have been used. We examine how useful it is to incorporate the least reliable of the clues and the genetic determination of sex available for only a handful of individuals. We finally discuss the implications of our findings for the design of field studies as well as further directions of research.

KEY WORDS: capture-recapture; parameter redundancy; multievent models.

1 Introduction

Sex differences in survival, dispersal or movement may have important consequences in demography, mating and parental investment patterns (Breitwisch, 1989; Gowaty, 1993), even for species where the two sexes have very similar body size and appearance. Thus, a modern study in ecology cannot ignore a priori this factor. However, naming the sex of an individual may be particularly tricky for monomorphic species observed in the field. Often the field biologist will rely on behavior to distinguish males from females. Some behavioral clues like position during copulation are thought to be quite reliable but other clues like the relative body size in a pair not so much and yet the latter type are easier to gather. With this kind of approach, it is thus not rare that an animal which has once been recorded as a male is later referred to as a female. Confidence in the correct determination of sex will progressively increase with repeated and consistent observations and, eventually, it may be decided that there is no reasonable doubt about the true sex of a particular individual. However, this does not hold for animals seen only a few times and for those with a contradictory record of given sex. A natural tendency for studying nonetheless sex-specific parameters is then to consider the sole individuals for which the sex has been determined with reasonable certainty. However, this approach has shortcomings, especially when the aim of the study is survival. First, it may represent a huge waste of data. For instance, in the field study that has motivated this paper, approximately 80% of the individuals had never been sexed and would thus be discarded. Second, to be sexed with reasonable certainty an individual must often have gone through several observations and hence have survived meanwhile. Then, estimation of survival based on the subsample of known-sex individuals will inevitably be biased high. Another possibility is to segregate the animals into three groups: males, females and unknown-sex individuals. The survival of the unknown-sex individuals can then be constrained to be a weighted average of the survivals of males and females (Oro and Pradel, 2000) with the weights reflecting the proportions of males and females among the unknown. However, these weights are not easy to determine as the relative proportions will depend on the sex-ratio but also on the relative catchability of males and females and on the

47 ability to identify the sex, which may differ between males and females. Thus, while this
48 second method is preferable, it is still imperfect and anyway does not make full use of
49 the information available.

50 We know of one example where sex uncertainty was directly incorporated in a survival
51 analysis (Conroy et al., 1999). This study dealt with serins *Serinus serinus*, a species
52 which is monomorphic only at the juvenile stage. A biometrical measure (wing length)
53 was used to predict the sex of the captured juveniles and an initial state 'predicted male'
54 or 'predicted female' was then assigned to each bird on this basis. If they were ever seen
55 again, their true sex was then determined and they were assigned the new state 'known
56 male' or 'known female'. This procedure allowed the estimation of transition probabilities
57 between the 'predicted sexes' and the 'known sexes' and thus the incorporation of the
58 never recaptured juveniles in the analysis. In this way, survival could be estimated without
59 bias. This model was cast within the framework of multistate capture-recapture models
60 (Arnason, 1972, 1973; Schwarz et al., 1993). Unfortunately, this approach does not work
61 for species which remain monomorphic all their life and this is why we had to develop
62 specific models. We present here a general solution to the problem of sex uncertainty
63 by introducing explicitly probabilities of errors. In essence, as compared to previous
64 approaches, instead of summarizing the data previous to the analysis, we model the very
65 process of sex identification.

66 This work was motivated by the demographic study of an Audouin's gull *Larus au-*
67 *douinii* colony at the Ebro Delta (Spain) (e.g. Oro and Pradel, 2000; Cam et al., 2004)
68 where the birds are sexed upon observation according to four criteria (see Oro et al.,
69 1999, for details): position during copulation, begging food during courtship, courtship
70 feeding and relative body size (the males being on average slightly heavier than the fe-
71 males, see Oro, 1998; Genovart et al., 2003a); or not sexed at all. The information can
72 be coded as three basic events: "the animal is judged to be a male", "the animal is
73 judged to be a female", "no judgment is issued". As this scheme should be relevant to
74 many studies, we start (section 2) by developing a set of four general models that extend
75 the Cormack-Jolly-Seber capture-recapture model and constant-parameter versions of it

(Cormack, 1964; Jolly, 1965, 1982; Seber, 1965) to uncertainty in sex assignment. These models incorporate probabilities of judgement and probabilities of error and can be further developed if needed. They are described in subsection 2.1. Because of the additional parameters involved, there is a legitimate concern that not all parameters may be identifiable. In subsection 2.2, we set off to study redundancy by the method of Catchpole, Morgan and Freeman (1997; 1998). The models of subsection 2.1 do not account for likely differences in probability of error with each of the four criteria used in the Audouin’s gull study, nor do they incorporate such peculiarity of this study as a trend over time in the attempt to identify the sex of the birds. In the next section 3, we construct a more refined and more specialized model, suited to our data. At the same time, we assess the effect of dealing or not dealing with different pieces of information: is it useful to gather the least reliable clue? Should genetic sexing be developed? The last section is a discussion of the interest of this kind of models and of the implications of our findings in terms of field work.

2 Four basic models

2.1 Description

The typical data are presented in Fig. 1. There, the true sex of every single individual is unknown. To write the probability of a given encounter history h , we start by conditioning on the first capture event—as is usual in capture-recapture (CR)—and then apply the law of total probabilities:

$$P(h) = P(f)P(h|f) + P(m)P(h|m).$$

Thus, $P(f)$ and $P(m)$ are the sex proportions among the unmarked captured, and $P(h|f)$ and $P(h|m)$ are the conditional probabilities of h for a female and a male respectively. We need the following parameters:

1. Traditional CR parameters

- ϕ_m, ϕ_f sex-specific survival rates

h	$\#\{h\}$
0000033000	41
0000112133	4
0001011000	2
0033000300	8
0330003313	4
0102220203	1
1001101100	1

Figure 1: Excerpt from the Audouin’s gull data coded according to three criteria: 1=judged male; 2=judged female; 3=no judgement made. $\#\{h\}$ is the number of individuals having encounter history h . The data set has 4025 individuals spread among 917 encounter histories over 10 years of study.

- 102 • p_m, p_f sex-specific encounter rates
- 103 2. New parameters
- 104 • μ , proportion of males in the population
- 105 • e_m, e_f sex-specific probabilities to judge the sex of an individual
- 106 • x_m, x_f sex-specific probabilities to issue the correct judgement.

It is now possible to write out $P(m)$ as $\frac{p_m \mu}{p_m \mu + p_f (1 - \mu)}$ ($P(f) = 1 - P(m)$). As for the conditional probabilities, let us write them for $h = 1 \ 3 \ 2$. If this relates to a male, it has been sexed the first time (probability e_m) correctly (probability x_m). It has then survived to occasion 2 (probability ϕ_m), when it was encountered (probability p_m), but not sexed (probability $1 - e_m$). It survived again (ϕ_m), was encountered at occasion 3 (p_m), and sexed (e_m) incorrectly ($1 - x_m$). All put together, we obtain $P(h|m) = e_m x_m \phi_m p_m (1 - e_m) \phi_m p_m e_m (1 - x_m)$. In a similar way, we would get $P(h|f) = e_f (1 -$

$x_f)\phi_f p_f(1 - e_f)\phi_f p_f e_f x_f$. The complete probability for this encounter history is thus

$$\begin{aligned}
P(h) = & \frac{p_m \mu}{p_m \mu + p_f(1 - \mu)} e_m x_m \phi_m p_m (1 - e_m) \phi_m p_m e_m (1 - x_m) \\
& + \frac{p_f(1 - \mu)}{p_m \mu + p_f(1 - \mu)} e_f(1 - x_f) \phi_f p_f (1 - e_f) \phi_f p_f e_f x_f.
\end{aligned} \tag{1}$$

The probability of each individual encounter history will be obtained in this way and hence the likelihood of the entire data set which, assuming independence of fates, is the product of them. With $\#h$ denoting the number of animals with encounter history h , the likelihood can be written in compact form:

$$L = \prod_h P(h)^{\#h}.$$

Maximum Likelihood Estimators (MLE) are derived by maximizing L as a function of the different parameters. These estimators are known to have excellent statistical properties such as being asymptotically unbiased and this is the approach to parameter estimation retained in leading CR software. However, because the current models do not fall in the category of existing models, we could not take advantage of this feature and wrote our own maximizing programs in MATLAB (Hanselman and Littlefield, 2000).

2.2 Parameter redundancy

The aim of this subsection is to study the impact of the introduction of sex uncertainty on parameter identifiability. As one can easily realize by browsing through the extensive capture-recapture literature, the number of traditional CR models in use is virtually unlimited as new models can be created by changing the kind of effects considered on each type of parameters. For instance, parameters may be constant over time or time-dependent; they may vary by sex, age... It is thus impossible to tell in advance which parameters are or are not identifiable in each possible model. However, it is useful to have some general guidance as to where problems are likely to occur. In traditional CR, the fully time-dependent Cormack-Jolly-Seber model (CJS) plays this role. In this model, it is known that survival over the last time interval is confounded with encounter probability at the last occasion and it is a general feature of the fully time-dependent CR models

that the last interval is critical. This problem disappears if survival or encounter parameters are hold constant. The introduction of sex uncertainty may well add new general identifiability problems. The minimal generalization of the CJS model consists in having the survival and encounter parameters, the 'traditional' parameters of subsection 2.1, still time-dependent and in considering that all of the 'new' parameters are constant over time. In this subsection, we examine whether this model is parameter-redundant. We also examine the three models derived by holding survival or encounter probabilities or both constant over time. In the following, we denote the four models considered by their sole variable part, i.e. (ϕ_{t*s}, p_{t*s}) , (ϕ_{t*s}, p_s) , (ϕ_s, p_{t*s}) , (ϕ_s, p_s) where s stands for sex and t for time.

Catchpole, Morgan and Freeman (1997; 1998) have developed a formal method (hereafter the CMF method) for studying parameter redundancy in models belonging to the exponential family of probability distributions. This method can be applied to the multinomial distribution of animals over the observable encounter histories. It indicates which parameters are directly identifiable and provides identifiable functions of the redundant parameters (see Catchpole and Morgan (1997) and Catchpole et al. (1998) for details). All the calculations being formal, we carried them out with MAPLE. Because the number of capture histories $4^k - 1$ increases rapidly with the number of time steps k , we could apply the procedure only for $k = 3$, but this is sufficient as the results obtained are easily demonstrated to hold in general.

A first conclusion (see Table 1) is that improving the realism of the models by adding new constant nuisance parameters to account for sex uncertainty does not restrict the ability to estimate the parameters of interest, i.e. the survival probabilities. Indeed, the only restriction we observe in the estimation of survival relates to the fully time-dependent model and this restriction is the same that was already present in the corresponding fully time-dependent CJS model. Actually, the new model improves over the CJS model applied to each sex separately in allowing the estimation of the ratio of survivals of males and females at the last occasion; we believe that this result is due to the assumption of a constant sex-ratio in the population. The other case of parameter redundancy concerns

Table 1: Identification of estimable quantities in 4 models incorporating probabilities of error. Superscripts are for time steps. Only 3 occasions were considered. ϕ^2 and p^3 are thus the last survival and last encounter parameters respectively.

Model	#par	Model rank	non estimable parameters	estimable functions
(ϕ_s, p_s)	9	9	none	—
(ϕ_{t*s}, p_s)	11	11	none	—
(ϕ_s, p_{t*s})	13	12	p_m^1, p_f^1	$\frac{p_f^1}{p_m^1}$
(ϕ_{t*s}, p_{t*s})	15	13	$p_m^1, p_m^3, p_f^1, p_f^3, \phi_m^2, \phi_f^2$	$\frac{p_f^1}{p_m^1}, \frac{\phi_f^2}{\phi_m^2}, \phi_m^2 p_m^3, \phi_f^2 p_f^3$

the initial encounter probabilities, p_m^1 and p_f^1 (absent from the CJS model). When encounter probabilities are time-dependent (models (ϕ_{t*s}, p_{t*s}) and (ϕ_s, p_{t*s})), only the ratio $\frac{p_f^1}{p_m^1}$ i.e. the relative catchability of males and females at the initial occasion is estimable. This is easily understood from the likelihood. The initial encounter probabilities appear in it only in the terms $P(m)$ and $P(f)$, and there only through their ratio. For instance, $P(m)$ can be rewritten

$$P(m) = \frac{p_m \mu}{p_m \mu + p_f (1 - \mu)} = \frac{\mu}{\mu + \frac{p_f}{p_m} (1 - \mu)}.$$

2.3 Dual solutions

A redundant parameter is also one that can take all of a continuous range of values while the likelihood remains at its maximum. For a given data set, this can be seen by drawing the profile likelihood curve, i.e. the curve of the maximum value that the likelihood can assume for each value of the parameter under scrutiny: for a redundant parameter, the profile likelihood presents a characteristically flat area at its top. The shape of the profile likelihood is interesting more generally. For instance, when the profile likelihood decreases rapidly away from the optimal value of the parameter, this parameter is estimable with great precision. Away from its maximum, the profile likelihood may

176 exhibit local maxima. If one of these happens to be exactly on the same level as the
177 'absolute' maximum, then this is an instance of non-identifiability without redundancy:
178 two entirely different values of the parameter maximize the likelihood equally well but
179 not the values in-between. This situation cannot be detected by the CMF method. Thus,
180 we examined the profile likelihood curves of the model with traditional parameters hold
181 constant (ϕ_s, p_s) applied to the Audouin's gull data already mentioned (Fig. 1). The
182 parameters were maintained within range by logit transformations. Instead of drawing
183 the profile likelihoods, we actually drew the profile deviances ($D = -2 \log L$) and hence
184 observed the minima rather than the maxima (Fig. 2).

185 A striking feature of model (ϕ_s, p_s) is the existence for each scalar parameter of two
186 distinct values where the deviance is minimized. These values organize into two sets which
187 represent two alternative solutions. This result is in fact data-independent and holds as
188 well for the three other models of this section: (ϕ_{t*s}, p_{t*s}) , (ϕ_{t*s}, p_s) and (ϕ_s, p_{t*s}) (see the
189 appendix for a theoretical demonstration). The deep reason for this fundamental duality
190 comes down to this: given that the true sex of any individual is never known, it is equally
191 acceptable, from the model point of view, to consider that an animal estimated many
192 times to be, say, a male is indeed a male and that most judgments were correct, or that
193 it is a female and most judgments were wrong. Yet, one solution produces probabilities
194 of error above 50%, which is probably not acceptable from the practitioner's point of
195 view. More precisely, one solution is derived from the other by exchanging the survival,
196 encounter and judgment probabilities of males and females, replacing x_m with $1 - x_f$ and
197 x_f with $1 - x_m$ and finally reversing the sex-ratio (see Appendix). As a consequence, the
198 profile deviance curves of sex-specific parameters are the same for males and females. In
199 the case of the Audouin's gull data, they present two close minima (Fig. 2). The profile
200 deviance curve of the proportion of males is symmetrical around 0.5 with two minima,
201 which happen to be very distant with our data, at 0.14 and 0.86, and, incidentally, utterly
202 unrealistic from a biological point of view.




Figure 2: Profile-deviance curves of the parameters of model (ϕ_s, p_s) applied to the Audouin's gull data set. For each parameter, the deviance (y-axis) presents two distinct minima. The curves for p , ϕ and e are the same for males and females.

Table 2: Maximum-likelihood estimates for each of the four models. For time-dependent parameters, the median is given. Each model has also a dual solution with probabilities of errors $>50\%$.

Model	encounter		survival		probability		probability		proportion
	probability				of judgment		to be right		of males
	(p)		(ϕ)		(e)		(x)		(μ)
	m	f	m	f	m	f	m	f	(m)
(ϕ_s, p_s)	0.61	0.72	0.88	0.95	0.02	0.79	0.53	0.56	0.86
(ϕ_{t*s}, p_s)	0.60	0.72	0.89	0.97	0.02	0.79	0.53	0.56	0.87
(ϕ_s, p_{t*s})	0.61	0.74	0.87	0.95	0.02	0.79	0.53	0.56	0.88
(ϕ_{t*s}, p_{t*s})	0.61	0.76	0.86	0.97	0.02	0.79	0.53	0.56	0.88

3 A more realistic model for the Audouin's gull data

The four models introduced in the previous section, fitted to the Audouin's gull data, produce very unreasonable if similar estimates: very high proportions of males that are almost never judged and probabilities of error close to 50% (Table 2). This is not entirely surprising given that they do not incorporate several known important features of the study. In this section, we exemplify the flexibility of our approach by building a more realistic model incorporating our knowledge of the biology and of the way the field work had been conducted.

To start, there was no good reason to believe that the error attached to each of the 4 criteria used (copulation, begging food, courtship feeding and relative body size) was the same. Copulation for instance was suspected to be the most reliable and relative body size, the most error-prone. We thus decided to distinguish the different criteria and recoded the data accordingly (see Table 3). All along the study, sex determination has been a secondary activity but it has been conducted on stringent criteria. This is reflected in the very few obvious mistakes (sexing is consistent over time for a given individual), but the roughly 80% of never-sexed individuals in the data set. Yet, during the course of

Table 3: Codes used to take into account the unequal reliability of each of the 4 criteria used to assess the sex of Audouin’s gulls upon observation at the Ebro delta colony (Spain).

judged from	to be		not judged
	m	f	
copulation	1	5	9
begging food	2	6	
courtship feeding	3	7	
body size	4	8	

the study, sexing has gained ground; especially the criterion ‘relative body size’, initially used very sparingly, has become more common by the end of the study. These features were incorporated in a model denoted $(\phi_s, p_t, \mu, e_T, m4_T, m1, m2, x1, x2, x3, x4)$ with the following characteristics:

- p_t , encounter probability: time but not sex-dependent.
- ϕ_s , survival rate: possibly sex-dependent but constant over time.
- μ , proportion of males: held constant.
- e_T : logit-linear trend over time (T) in the attempt to judge the sex (e). Note that because birds are always judged by pair, we put no sex-dependency on e .
- $m4_T$: logit-linear trend over time (T) in the frequency of use of the criterion ‘relative body size’ ($m4$) among the instances when a judgment was issued.
- $m1$: frequency of use of the criterion ‘copulation’ among the behavioural criteria (i.e. excluding ‘body size’): held constant.
- $m2$ and $m3$: same as $m1$ for ‘begging food’ and ‘courtship feeding’ respectively (one of $m1$, $m2$ or $m3$ is redundant).
- xi : probability to be right when using criterion i ($i = 1, \dots, 4$): held constant.

235 This model has 23 parameters including 1 intercept and 1 slope for e and 1 intercept and
236 1 slope for m_4 and 10 capture probabilities.

237 Additionally, a very few birds (24) had been sexed genetically (Genovart et al.,
238 2003*a,b*). We fitted the previous model using (analysis A) and not using (analysis B)
239 this limited information. When we used it, $P(m)$ in the capture history of the genetically
240 sexed individuals was set to 1 or 0 as appropriate. On the other hand, we considered the
241 effects of not using the presumably less reliable criterion, 'body size'. To do that, the
242 corresponding observations were recoded as a 'no judgment' observation. When doing
243 this latter analysis (analysis C), the genetic determination of sex was not used. Thus,
244 we have a gradient of decreasing amount of information from analysis A to analysis C
245 (Table 4).

246 All three analyses yield estimates in agreement with what is known of the biology of
247 the species. For instance, survival estimates are very close to those estimated previously
248 by CR in the study site (e.g. Oro et al., 1999; Cam et al., 2004). We also note that,
249 as anticipated, body size is the least reliable clue but the copulation criterion does not
250 come out as the obvious best method. The results with and without genetical information
251 are very similar (first 2 columns of Table 4). However, the known sex of only 24 birds
252 suffices to break the tie between the dual solutions: while analyses B and C still have
253 two solutions (only the reasonable one is presented in Table 4), analysis A has just one,
254 as can be seen on the profile deviance curves relative to the proportion of males (Fig. 3).
255 In other, not shown models, where sex-ratio is initially not identifiable, the additional
256 information brought about by the 24 birds renders this and other parameters identifiable.
257 As for the criterion 'body size', even though mistakes are made on average once in every
258 10 judgments and only 15.70% of the judgments made use this criterion, dropping it
259 results in a disproportionate loss of precision (compare the standard errors in the last two
260 columns of Table 4).

Table 4: Results of model $(\phi_s, p_t, \mu, e_T, m4_T, m1, m2, x1, x2, x3, x4)$ applied to the Audouin’s gull data set with different amount of information. Analysis A uses all 4 criteria and the genetic determination of the sex of 24 birds; Analyses B and C do not use the genetic determination of sex; Analysis C does not use the judgments based on body size. Standard errors are given between parentheses. With less information (from left to right), precision decreases. For analyses B and C, we have retained the dual solution that yields proportions of error $<50\%$.

Analysis	A	B	C
proportion of males	0.53 (0.027)	0.53 (0.029)	0.55 (0.034)
female survival	0.91 (0.013)	0.91 (0.014)	0.93 (0.024)
male survival	0.86 (0.013)	0.86 (0.014)	0.84 (0.025)
error copulation	0.06 (0.041)	0.06 (0.041)	0.08 (0.055)
error begging food	0.05 (0.029)	0.06 (0.031)	0.09 (0.077)
error courtship feeding	0.00 (0.161)	0.00 (0.155)	0.00 (0.286)
error body size	0.11 (0.064)	0.09 (0.074)	NA




Figure 3: Improvement in the shape of the deviance brought about by a limited amount of external information shown on the profile deviance of model $(\phi_s, p_t, \mu, e_T, m4_T, m1, m2, x1, x2, x3, x4)$ applied to the Audouin's gull data set and relative to the proportion of males μ . The genetic sexing of 24 individuals renders the profile deviance steeper at its bottom where it retains just one minimum (Analysis A; continuous line) as compared to when the genetic information is not used (Analysis B; dotted line). There is a narrow ridge between the 2 dual solutions in Analysis B.

261 4 Discussion

262 A main concern when we started building these models was the risk of parameter redun-
263 dancy. The models would have been useless if no parameter of interest was estimable.
264 This fear had been voiced in particular by Nichols et al. (2004) when they examined
265 the restricted situation where the sex is not always assessed but, if so, without error.
266 Our results show on the contrary that it is quite possible to run models incorporating
267 probabilities of error and nonetheless gain knowledge about such important biological
268 parameters as survival rates. Although these models have in general two mathematical
269 solutions, one is so unreasonable that it should not be difficult to sort out which one is
270 to be retained.

271 A more serious difficulty is numerical. These models, and more generally the mul-
272 tievent models (Pradel, 2005) to which they belong—but also the now classical multi-
273 state models—present local minima to which the optimization algorithm occasionally
274 converges. The profile deviance curves may help diagnose the problem and improved
275 algorithms may be sought (see, in particular, section 6.2 in Choquet et al., 2005), but it
276 is also possible to guard against local minima by augmenting the information available.
277 Often, such information already exists, but could not be exploited in the traditional CR
278 models. One of the major aims of the newly developed models was indeed to alleviate
279 and valorize the field work by comprehending and exploiting a wider range of data.

280 In this paper, we have examined two ways of gathering additional information: us-
281 ing less efficient clues and sexing genetically some animals. Genetic sexing is of course
282 paramount for the determination of sex, but for practical reasons it cannot usually be
283 performed on every single individual. In our example, genetically sexing very few birds
284 proved sufficient to greatly improve the shape of the deviance and hence the efficiency of
285 the optimization algorithm. We also observed that it eliminated the wrong one of the two
286 dual solutions. Finally, genetic sexing occasionally rendered identifiable new parameters
287 in parameter-redundant models. It should also be noticed that genetic sexing is only one
288 approach to specifying the a priori probability to be a male, $P(m)$, for some individu-
289 als. Other, not necessarily 0 – 1 predictors of sex frequently available in old studies, like

290 biometrical measures, can be incorporated in the model in the same way ($P(m)$ can be
 291 set to some intermediate value between 0 and 1). This approach of incorporating exter-
 292 nal information has been used by Fujiwara and Caswell (2002) although in a way that,
 293 past the first encounter, may not be correct: the conditioning in their stage-assignment
 294 matrix, $P(stage|measure)$, is the reverse of what is needed, $P(measure|stage)$, in the
 295 calculation of the likelihood (see Pradel, 2005, for a formal demonstration). As for the
 296 use of less efficient clues, our example proved that they may be valuable beyond what
 297 seems at first sight attainable based on their frequency of use and efficiency (percentage
 298 of error). We thus encourage people to gather such clues especially if they are not time
 299 consuming (e.g. Redman et al., 2002). After three or four visits, it will always be possible
 300 to assess the quality of the prospective clues as well as the improvement brought about
 301 by such and such identification method in terms of gained precision. If different methods
 302 for sex identification are in competition, it is then possible to decide objectively, from the
 303 informed assessment of their relative costs and benefits, which ones to favor and which
 304 ones to discard.

305 References

- 306 Arnason, A. N. 1972. Parameter estimates from mark-recapture experiments on two
 307 populations subject to migration and death. *Researches on Population Ecology* **13**:97–
 308 113.
- 309 Arnason, A. N. 1973. The estimation of population size, migration rates and survival in
 310 a stratified population. *Researches on Population Ecology* **15**:1–8.
- 311 Breitwisch, R. 1989. Mortality patterns, sex ratios, and parental investment in monoga-
 312 mous birds. *Current Ornithology* **6**:1–50.
- 313 Cam, E., D. Oro, R. Pradel, and J. Jimenez. 2004. Assessment of hypotheses about
 314 dispersal in a long-lived seabird using multistate capture-recapture models. *Journal of*
 315 *Animal Ecology* **73**:723–736.

316 Catchpole, E., B. J. Morgan, and S. Freeman. 1998. Estimation in parameter-redundant
317 models. *Biometrika* **85**:462–8.

318 Catchpole, E. A. and B. J. T. Morgan. 1997. Detecting parameter redundancy. *Biometrika*
319 **84**:187–196.

320 Choquet, R., A.-M. Reboulet, R. Pradel, O. Gimenez, and J.-D. Lebreton, 2005. M-
321 SURGE 1.8 User’s manual. Mimeographed document, CEFÉ UMR CNRS 5175.

322 Conroy, M. J., J. C. Senar, J. E. Hines, and J. Domenech. 1999. Development and appli-
323 cation of a mark-recapture model incorporating predicted sex and transitory behaviour.
324 *Bird Study* **46**:62–73.

325 Cormack, R. M. 1964. Estimates of survival from the sighting of marked animals. *Bio-*
326 *metrika* **51**:429–438.

327 Fujiwara, M. and H. Caswell. 2002. Estimating population projection matrices from
328 multi-stage mark-recapture data. *Ecology* **83**:3257–65.

329 Genovart, M., D. Oro, and F. Bonhomme. 2003*a*. Genetic and morphological differenti-
330 ation between the two largest breeding colonies of audouin’s gull *larus audouinii*. *Ibis*
331 **145**:448–456.

332 Genovart, M., D. Oro, X. Ruiz, R. Griffiths, P. Monaghan, and R. G. Nager. 2003*b*.
333 Seasonal changes in brood sex composition in audouin’s gulls. *Condor* **105**:783–790.

334 Gowaty, P. A. 1993. Differential dispersal, local resource competition, and sex ratio
335 variation in birds. *American Naturalist* **141**:263–280.

336 Hanselman, D. and B. R. Littlefield. 2000. Mastering MATLAB 6. Prentice Hall,
337 Englewood Cliffs, New Jersey, USA.

338 Jolly, G. 1982. Mark-recapture models with parameters constant in time. *Biometrics*
339 **38**:301–21.

- 340 Jolly, G. M. 1965. Explicit estimates from capture-recapture data with both death and
341 immigration-stochastic model. *Biometrika* **52**:225–247.
- 342 Nichols, J. D., W. L. Kendall, J. E. Hines, and J. A. Spendelov. 2004. Estimation of sex-
343 specific survival from capture-recapture data when sex is not always known. *Ecology*
344 **85**:3192–3201.
- 345 Oro, D., 1998. Audouin’s gull account. Pages 47–61 *in* M. Ogilvie, editor. The birds of
346 western palearctic. Oxford University Press, Oxford, UK.
- 347 Oro, D. and R. Pradel. 2000. Determinants of local recruitment in a growing colony of
348 audouin’s gull. *Journal of Animal Ecology* **69**:119–32.
- 349 Oro, D., R. Pradel, and J.-D. Lebreton. 1999. Food availability and nest predation
350 influence life history traits in audouin’s gull, *larus audouinii*. *Oecologia* **118**:438–45.
- 351 Pradel, R. 2005. Multievent: an extension of multistate capture-recapture models to
352 uncertain states. *Biometrics* **61**:442–447.
- 353 Redman, K. K., S. Lewis, R. Griffiths, S. Wanless, and K. C. Hamer. 2002. Sexing
354 northern gannets from DNA, morphology and behavior. *Waterbirds* **25**:230–234.
- 355 Schwarz, C. J., J. F. Schweigert, and A. Arnason. 1993. Estimating migration rates using
356 tag-recovery data. *Biometrics* **49**:177–193.
- 357 Seber, G. A. F. 1965. A note on the multiple-recapture census. *Biometrika* **52**:249–259.

358 Appendix

359 The aim of this appendix is to establish the existence of two distinct dual solutions to
360 the maximization of the likelihood of model (ϕ_s, p_s) presented in subsection 2.1, as well
361 as some related properties of its profile deviance curves. Why this happens is rooted in
362 the law of total probabilities applied to an arbitrary encounter history h .

$$363 \quad P(h) = P(f)P(h|f) + P(m)P(h|m)$$

364 In this formula, the conditional probability $P(h|f)$ is a function of the sole parameters
 365 ϕ_f, p_f, e_f, x_f . This function depends on the particular encounter history h . Let write

$$366 \quad P(h|f) = g_h(\phi_f, p_f, e_f, x_f). \quad (2)$$

367 The probability of the same history conditional on the animal being this time a male
 368 is obtained by using the male parameters instead of the female ones. However, when a
 369 judgment was right for a female, it is wrong for a male (and vice-versa) so that x_f should
 370 be replaced with $1 - x_m$ (see the example of section 2.1 leading to equation 1 if this is
 371 not immediately clear). Thus,

$$372 \quad P(h|m) = g_h(\phi_m, p_m, e_m, 1 - x_m). \quad (3)$$

373 The unconditional probabilities $P(m)$ and $P(f)$ are respectively

$$374 \quad \frac{p_m \mu}{p_m \mu + p_f(1 - \mu)} \text{ and } \frac{p_f(1 - \mu)}{p_m \mu + p_f(1 - \mu)}. \quad (4)$$

375 The transformation $\Theta: [0, 1]^9 \longrightarrow [0, 1]^9$

$$376 \quad (\theta_1, \dots, \theta_9) \mapsto (\theta_2, \theta_1, \theta_4, \theta_3, 1 - \theta_5, \theta_7, \theta_6, 1 - \theta_9, 1 - \theta_8)$$

377 plays a key role. We first establish

378 **Lemma 1.** *The probability of any encounter history, seen as a function of the parameter*
 379 *vector $(\phi_m, \phi_f, p_m, p_f, \mu, e_m, e_f, x_m, x_f)$, is invariant under Θ .*

$$380 \quad \forall h, P(h) \circ \Theta = P(h)$$

381 *Proof.* When the parameters are changed in the following way:

$$382 \quad (\phi_m, \phi_f, p_m, p_f, \mu, e_m, e_f, x_m, x_f) \xrightarrow{\Theta} (\phi_f, \phi_m, p_f, p_m, 1 - \mu, e_f, e_m, 1 - x_f, 1 - x_m),$$

$$384 \quad P(f) \text{ becomes } \frac{p_m \mu}{p_m \mu + p_f(1 - \mu)} = P(m); \quad [\text{from (4)}]$$

$$385 \quad P(m) \text{ becomes } \frac{p_f(1 - \mu)}{p_m \mu + p_f(1 - \mu)} = P(f); \quad [\text{from (4)}]$$

$$386 \quad P(h|f) \text{ becomes } g_h(\phi_m, p_m, e_m, 1 - x_m) = P(h|m); \quad [\text{from (2) and (3)}]$$

$$387 \quad P(h|m) \text{ becomes } g_h(\phi_f, p_f, e_f, x_f) = P(h|f); \quad [\text{from (2) and (3)}]$$

389 so that, eventually, $P(h)$ itself is unchanged. □

390 A consequence of this lemma is that the likelihood and the deviance are equally un-
 391 changed under Θ . Consequently, if the parameter vector $(\hat{\phi}_m, \hat{\phi}_f, \hat{p}_m, \hat{p}_f, \hat{\mu}, \hat{e}_m, \hat{e}_f, \hat{x}_m, \hat{x}_f)$
 392 maximizes the likelihood, its transform by Θ , $(\hat{\phi}_f, \hat{\phi}_m, \hat{p}_f, \hat{p}_m, 1 - \hat{\mu}, \hat{e}_f, \hat{e}_m, 1 - \hat{x}_f, 1 - \hat{x}_m)$,
 393 maximizes it too. Hence, the theorem:

394 **Theorem 1.** *The likelihood of model (ϕ_s, p_s) is maximized at two generally distinct points*
 395 *one of which is the transform of the other by the unipotent mapping Θ .*

396 We now examine some ensuing properties of the profile deviance curves. Let D be
 397 the deviance of model (ϕ_s, p_s) . $D: [0, 1]^9 \rightarrow [0, +\infty]$, is a function of the 9 probability
 398 parameters: $\phi_m, \phi_f, p_m, p_f, \mu, e_m, e_f, x_m, x_f$, which has the property that $D \circ \Theta = D$. If

$$399 \quad E_i^a = \{\theta \in [0, 1]^9, \theta_i = a\},$$

400 the profile deviance function for parameter i is

$$401 \quad P_i(a) \stackrel{\text{def}}{=} \min_{E_i^a} D = \min_{E_i^a} (D \circ \Theta) = \min_{\Theta(E_i^a)} D.$$

402 Given that:

$$\begin{aligned} 403 \quad & \Theta(E_1^a) = E_2^a, & \Theta(E_2^a) = E_1^a, \\ 404 \quad & \Theta(E_3^a) = E_4^a, & \Theta(E_4^a) = E_3^a, \\ 405 \quad & \Theta(E_5^a) = E_5^{1-a}, \\ 406 \quad & \Theta(E_6^a) = E_7^a, & \Theta(E_7^a) = E_6^a, \\ 407 \quad & \Theta(E_8^a) = E_9^{1-a}, & \Theta(E_9^a) = E_8^{1-a}, \end{aligned}$$

409 then,

$$\begin{aligned} 410 \quad & P_1(a) = \min_{E_2^a} D = P_2(a), \\ 411 \quad & P_3(a) = \min_{E_4^a} D = P_4(a), \\ 412 \quad & P_5(a) = \min_{E_5^{1-a}} D = P_5(1-a), \\ 413 \quad & P_6(a) = \min_{E_7^a} D = P_7(a), \\ 414 \quad & P_8(a) = \min_{E_9^{1-a}} D = P_9(1-a). \end{aligned}$$

416 Thus, the profile deviances of the survival (parameters 1 and 2), encounter (parameters 3
417 and 4) and judgment probabilities (parameters 6 and 7) are the same for both sexes. The
418 profile deviance of the probability of error for the males (parameter 8) is the symmetrical
419 with respect to $a=0.5$ of that for the females (parameter 9), and the profile deviance
420 curve of the proportion of males (parameter 5) is symmetrical with respect to $a=0.5$.