



## Synthetic analogues of Huwentoxin-IV spider peptide with altered human Nav1.7/Nav1.6 selectivity ratios

Ludivine Lopez, Jérôme Montnach, Barbara Oliveira-Mendes, Kuldip Khakh, Baptiste Thomas, Sophia Lin, Cécile Caumes, Steven Wesolowski, Sébastien Nicolas, Denis Servent, et al.

### ► To cite this version:

Ludivine Lopez, Jérôme Montnach, Barbara Oliveira-Mendes, Kuldip Khakh, Baptiste Thomas, et al.. Synthetic analogues of Huwentoxin-IV spider peptide with altered human Nav1.7/Nav1.6 selectivity ratios. *Frontiers in Cell and Developmental Biology*, 2021, 9, pp.798588. 10.3389/fcell.2021.798588 . hal-03520574

**HAL Id: hal-03520574**

**<https://hal.science/hal-03520574>**

Submitted on 12 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



# Synthetic Analogues of Huwentoxin-IV Spider Peptide With Altered Human Na<sub>v</sub>1.7/Na<sub>v</sub>1.6 Selectivity Ratios

Ludivine Lopez<sup>1</sup>, Jérôme Montnach<sup>1</sup>, Barbara Oliveira-Mendes<sup>1</sup>, Kuldip Khakh<sup>2</sup>, Baptiste Thomas<sup>3</sup>, Sophia Lin<sup>2</sup>, Cécile Caumes<sup>3</sup>, Steven Wesolowski<sup>2</sup>, Sébastien Nicolas<sup>1</sup>, Denis Servent<sup>4</sup>, Charles Cohen<sup>2</sup>, Rémy Bérout<sup>3</sup>, Evelyne Benoit<sup>4</sup> and Michel De Waard<sup>1,3,5\*</sup>

<sup>1</sup>L'institut du Thorax, INSERM, CNRS, UNIV NANTES, Nantes, France, <sup>2</sup>Xenon Pharmaceuticals, Burnaby, BC, Canada, <sup>3</sup>Smartox Biotechnology, Saint-Egrève, France, <sup>4</sup>Département Médicaments et Technologies pour La Santé (DMTS), Service d'Ingénierie Moléculaire pour La Santé (SiMoS), ERL CNRS/CEA, Institut des Sciences du Vivant Frédéric Joliot, CEA, Université Paris Saclay, Gif-sur-Yvette, France, <sup>5</sup>LabEx « Ion Channels, Science and Therapeutics », Valbonne, France

## OPEN ACCESS

### Edited by:

Giuseppe Calamita,  
University of Bari Aldo Moro, Italy

### Reviewed by:

Jean-François Desaphy,  
University of Bari Aldo Moro, Italy  
Shannon Shields,  
Coda Biotherapeutics, Inc.,  
United States

### \*Correspondence:

Michel De Waard  
michel.dewaard@univ-nantes.fr

### Specialty section:

This article was submitted to  
Cellular Biochemistry,  
a section of the journal  
Frontiers in Cell and Developmental  
Biology

**Received:** 20 October 2021

**Accepted:** 22 November 2021

**Published:** 20 December 2021

### Citation:

Lopez L, Montnach J, Oliveira-Mendes B, Khakh K, Thomas B, Lin S, Caumes C, Wesolowski S, Nicolas S, Servent D, Cohen C, Bérout R, Benoit E and De Waard M (2021) Synthetic Analogues of Huwentoxin-IV Spider Peptide With Altered Human Na<sub>v</sub>1.7/Na<sub>v</sub>1.6 Selectivity Ratios. *Front. Cell Dev. Biol.* 9:798588. doi: 10.3389/fcell.2021.798588

Huwentoxin-IV (HwTx-IV), a peptide discovered in the venom of the Chinese bird spider *Cyriopagopus schmidtii*, has been reported to be a potent antinociceptive compound due to its action on the genetically-validated Na<sub>v</sub>1.7 pain target. Using this peptide for antinociceptive applications *in vivo* suffers from one major drawback, namely its negative impact on the neuromuscular system. Although studied only recently, this effect appears to be due to an interaction between the peptide and the Na<sub>v</sub>1.6 channel subtype located at the presynaptic level. The aim of this work was to investigate how HwTx-IV could be modified in order to alter the original human (h) Na<sub>v</sub>1.7/Na<sub>v</sub>1.6 selectivity ratio of 23. Nineteen HwTx-IV analogues were chemically synthesized and tested for their blocking effects on the Na<sup>+</sup> currents flowing through these two channel subtypes stably expressed in cell lines. Dose-response curves for these analogues were generated, thanks to the use of an automated patch-clamp system. Several key amino acid positions were targeted owing to the information provided by earlier structure-activity relationship (SAR) studies. Among the analogues tested, the potency of HwTx-IV E<sup>4</sup>K was significantly improved for hNa<sub>v</sub>1.6, leading to a decreased hNa<sub>v</sub>1.7/hNa<sub>v</sub>1.6 selectivity ratio (close to 1). Similar decreased selectivity ratios, but with increased potency for both subtypes, were observed for HwTx-IV analogues that combine a substitution at position 4 with a modification of amino acid 1 or 26 (HwTx-IV E<sup>1</sup>G/E<sup>4</sup>G and HwTx-IV E<sup>4</sup>K/R<sup>26</sup>Q). In contrast, increased selectivity ratios (>46) were obtained if the E<sup>4</sup>K mutation was combined to an additional double substitution (R<sup>26</sup>A/Y<sup>33</sup>W) or simply by further substituting the C-terminal amidation of the peptide by a carboxylated motif, linked to a marked loss of potency on hNa<sub>v</sub>1.6 in this latter case. These results demonstrate that it is possible to significantly modulate the selectivity ratio for these two channel subtypes in order to improve the potency of a given analogue for hNa<sub>v</sub>1.6 and/or hNa<sub>v</sub>1.7 subtypes. In addition, selective analogues for hNa<sub>v</sub>1.7, possessing better safety profiles, were produced to limit neuromuscular impairments.

**Keywords:** huwentoxin-IV analogues, *Cyriopagopus schmidtii*, development of pain therapeutics, Na<sub>v</sub>1.6 and Na<sub>v</sub>1.7 channel subtypes, automated patch-clamp, structure-function relationship, peptide synthesis

## INTRODUCTION

For over 10 years, the Na<sub>v</sub>1.7 channel subtype is considered as an attractive pain target owing to several compelling genetic evidences. Loss-of-function mutations in *SCN9A*, the gene encoding human (h) Na<sub>v</sub>1.7, lead to congenital insensitivity to pain (referred to as CIP) (Cox et al., 2006). Conversely, gain-of-function mutations of the same gene lead to erythromelalgia and paroxysmal extreme pain disorder (Estacion et al., 2008). Because of these clinical evidences, the hNa<sub>v</sub>1.7 channel has been the target of choice for high-throughput screening campaigns to identify blockers as pain therapeutics. Two types of libraries were used for this purpose: small organic compounds from leading pharmaceutical companies (MacSari et al., 2012; Nguyen et al., 2012; Sun et al., 2014; Focken et al., 2016; Wu et al., 2017; Cuesta and Meneses, 2021) and natural peptides originating from animal venoms (Trim et al., 2021). Indeed, disulfide-bridged peptides, frequently purified from animal venoms, have been considered as interesting lead compounds (Bordon et al., 2020), either as pore blockers or gating modifiers, for the modulation of ion channels in general and the treatment of pain in particular (Cardoso and Lewis, 2018). Owing to chemical spaces larger than those of small molecules, they possess better affinities and selectivity profiles even when the target of interest shares high sequence identity with other ion channel subtypes. This was illustrated by peptides issued from spider venoms, often possessing an Inhibitory Cystine Knot (ICK) fold, that were shown to be able of distinguishing closely related subtypes of Na<sub>v</sub> channels (for reviews, see (Gonçalves et al., 2018a; Cardoso and Lewis, 2019; Dongol et al., 2019)). This ability is facilitated by the interaction of these peptides with the different voltage-sensor domains, known to be the most divergent in sequence in contrast to pore region.

Along with protoxin II (Montnach et al., 2021), one of the best studied inhibitors of hNa<sub>v</sub>1.7 channel is huwentoxin-IV (HwTx-IV), a 35 amino acid peptide isolated from the venom of the Chinese bird-eating tarantula spider *Cyriopagopus schmidtii* (Peng et al., 2002). HwTx-IV belongs to the NaSpTx family 1 (Klint et al., 2012) because of its sequence. In addition, according to the ICK motif, this peptide is folded with the Cys<sup>2</sup>-Cys<sup>17</sup>, Cys<sup>9</sup>-Cys<sup>24</sup>, Cys<sup>16</sup>-Cys<sup>31</sup> disulfide bridge pattern, which favors the formation of a double-stranded antiparallel beta-sheet (Leu<sup>22</sup>-Ser<sup>25</sup>, Trp<sup>30</sup>-Tyr<sup>33</sup>) along with four turns (Glu<sup>4</sup>-Lys<sup>7</sup>, Pro<sup>11</sup>-Asp<sup>14</sup>, Lys<sup>18</sup>-Lys<sup>21</sup>, Arg<sup>26</sup>-Arg<sup>29</sup>). Overall, this peptide possesses a compact and rigid scaffold (Peng et al., 2002) that confers high protease resistance and thus elevated *in vivo* stability. Like most other spider toxins, HwTx-IV has been reported to be a gating modifier by interacting with the voltage-sensor domain II of hNa<sub>v</sub>1.7 and trapping it into the closed configuration (Xiao et al., 2008; Xiao et al., 2011; Shen et al., 2019; Gao et al., 2020). This binding was suggested to imply residue Glu<sup>753</sup> of the S1-S2 loop and four residues, Glu<sup>811</sup>, Leu<sup>814</sup>, Asp<sup>816</sup> and Glu<sup>819</sup>, of the S3-S4 loop of this channel subtype, which defines an EELDE motif for the toxin interaction. Because of the nature of this complex interaction, HwTx-IV owns *de facto* good natural selectivity for hNa<sub>v</sub>1.7. However, hNa<sub>v</sub>1.1,

hNa<sub>v</sub>1.2, hNa<sub>v</sub>1.3 and hNa<sub>v</sub>1.6 channel subtypes are also sensitive to the peptide, while hNa<sub>v</sub>1.4, hNa<sub>v</sub>1.5 and hNa<sub>v</sub>1.8 are resistant (Rahnama et al., 2017; Gonçalves et al., 2018b). Indeed, the EELDE motif for HwTx-IV binding onto hNa<sub>v</sub>1.7 is preserved in hNa<sub>v</sub>1.6, and only slightly modified (towards an EELNE motif) in hNa<sub>v</sub>1.1, hNa<sub>v</sub>1.2 and hNa<sub>v</sub>1.3. Many more alterations in this motif are detected in hNa<sub>v</sub>1.4, hNa<sub>v</sub>1.5 and hNa<sub>v</sub>1.8. These observations are all coherent with the selectivity data (Xiao et al., 2008; Revell et al., 2013; Rahnama et al., 2017; Gonçalves et al., 2018b; Agwa et al., 2020).

Considering the high potency of HwTx-IV for hNa<sub>v</sub>1.7, it was quite logical to test the analgesic potential of the peptide in pain animal models. HwTx-IV was thus reported as an efficient analgesic in rodent models of inflammatory and neuropathic pain (Liu et al., 2014b), as well as of spontaneous pain induced by the Na<sub>v</sub>1.7 activator OD1, the first toxin isolated from the venom of the scorpion *Odontobuthus doriae* (Rahnama et al., 2017). However, these encouraging preclinical results hampered clinical development because the peptide also produced evident side effects *in vivo* resulting from impairment of neuromuscular transmission (Liu et al., 2014a; Deuis et al., 2016; Flinspach et al., 2017; Rahnama et al., 2017). It is only later that this effect was clearly attributed to the activity of HwTx-IV on the Na<sub>v</sub>1.6 channel, a subtype localized at presynaptic nerve terminals innervating the muscles (Gonçalves et al., 2018b).

Because of the analgesic potential of HwTx-IV, mainstream efforts have been dedicated to producing analogues with enhanced potency on hNa<sub>v</sub>1.7 (Revell et al., 2013; Rahnama et al., 2017). However, most of earlier studies, if not all, considered the potential effects of the peptide sequence modifications on its selectivity profile towards other Na<sub>v</sub> subtypes (in general Na<sub>v</sub>1.5 or Na<sub>v</sub>1.2) with the notable exception of the Na<sub>v</sub>1.6 subtype (Minassian et al., 2013; Neff et al., 2020). The present report aims at investigating the relative selectivity of HwTx-IV and 19 synthetic analogues on both hNa<sub>v</sub>1.7 and hNa<sub>v</sub>1.6 channel subtypes. All these analogues were designed on the basis of earlier SAR investigations and mainly by focusing on amino acid residues shown to influence the hNa<sub>v</sub>1.7/hNa<sub>v</sub>1.2 selectivity ratio. Two goals were pursued: 1) identify analogues with improved selectivity for hNa<sub>v</sub>1.7, and 2) identify new analogues that have improved potencies for hNa<sub>v</sub>1.6. For this purpose, similar experimental conditions for chemical syntheses and functional evaluations were employed to get reliable and exploitable results. Under these conditions, 11 analogues with increased apparent affinity for hNa<sub>v</sub>1.6 were identified, among which 4 improved the half maximal inhibitory concentration (IC<sub>50</sub>) values by more than 36-fold without drastically decreasing the peptide potency for hNa<sub>v</sub>1.7. Conversely, 7 analogues increased the apparent affinity for hNa<sub>v</sub>1.7, while only one of them had a decreased potency for hNa<sub>v</sub>1.6. These data illustrate that adequately mutating HwTx-IV contributes to large improvements of the peptide potency for hNa<sub>v</sub>1.6, while in parallel mildly improving the potency for hNa<sub>v</sub>1.7. This process reaches the point at which the potencies of some of the analogues are similar, in the low nanomolar range, for both channel subtypes.

Peptides	Sequences	MM (Da)
<b>HwTx-IV WT</b>	<b>ECLEIFKACNPSNDQCKSSKLVCSRKTRWCKYQI*</b>	<b>4107.20</b>
HwTx-IV E <sup>1</sup> G	GCLEIFKACNPSNDQCKSSKLVCSRKTRWCKYQI*	4031.93
HwTx-IV E <sup>4</sup> G	ECLGIFKACNPSNDQCKSSKLVCSRKTRWCKYQI*	4031.93
HwTx-IV E <sup>4</sup> K	ECLKIFKACNPSNDQCKSSKLVCSRKTRWCKYQI*	4103.01
HwTx-IV E <sup>4</sup> R	ECLRIFKACNPSNDQCKSSKLVCSRKTRWCKYQI*	4131.01
HwTx-IV K <sup>18</sup> A	ECLEIFKACNPSNDQCCASSKLVCSRKTRWCKYQI*	4046.89
HwTx-IV K <sup>32</sup> N	ECLEIFKACNPSNDQCKSSKLVCSRKTRWCNYQI*	4089.89
HwTx-IV R <sup>26</sup> A	ECLEIFKACNPSNDQCKSSKLVCSAKTRWCKYQI*	4018.89
HwTx-IV R <sup>26</sup> Q	ECLEIFKACNPSNDQCKSSKLVCSQKTRWCKYQI*	4075.91
HwTx-IV T <sup>28</sup> W	ECLEIFKACNPSNDQCKSSKLVCSRKWRWCKYQI*	4188.98
HwTx-IV Y <sup>33</sup> W	ECLEIFKACNPSNDQCKSSKLVCSRKTRWCKWQI*	4126.97
HwTx-IV E <sup>1</sup> G/E <sup>4</sup> G	GCLEIFKACNPSNDQCKSSKLVCSRKTRWCKYQI*	3959.91
HwTx-IV E <sup>4</sup> R/K <sup>32</sup> N	ECLRIFKACNPSNDQCKSSKLVCSRKTRWCNYQI*	4116.95
HwTx-IV E <sup>4</sup> K/R <sup>26</sup> Q	ECLKIFKACNPSNDQCKSSKLVCSQKTRWCKYQI*	4074.97
HwTx-IV E <sup>1</sup> G/E <sup>4</sup> G/K <sup>18</sup> A	GCLEIFKACNPSNDQCCASSKLVCSRKTRWCKYQI*	3902.85
HwTx-IV E <sup>4</sup> K/R <sup>26</sup> A/Y <sup>33</sup> W	ECLKIFKACNPSNDQCKSSKLVCSAKTRWCKWQI*	4040.97
HwTx-IV C <sub>ter</sub> COOH	ECLEIFKACNPSNDQCKSSKLVCSRKTRWCKYQI-COOH	4104.93
HwTx-IV K <sup>36</sup> /C <sub>ter</sub> COOH	ECLEIFKACNPSNDQCKSSKLVCSRKTRWCKYQIK-COOH	4233.03
HwTx-IV E <sup>4</sup> K/C <sub>ter</sub> COOH	ECLKIFKACNPSNDQCKSSKLVCSRKTRWCKYQI-COOH	4103.99
HwTx-IV E <sup>4</sup> G/C <sub>ter</sub> COOH	ECLGIFKACNPSNDQCKSSKLVCSRKTRWCKYQI-COOH	4032.91

**FIGURE 1** | Sequence alignment, with the disulfide bond connectivity, and molecular masses of HwTx-IV and its 19 analogues. The residue substitutions in the peptide analogues are indicated in red. WT: Wild-type. MM: Molecular masses after folding/oxidation. \*: C-terminal amidation.

## MATERIALS AND METHODS

### Chemical Synthesis of HwTx-IV Analogues

HwTx-IV analogues were all chemically assembled and provided by Smartox Biotechnology. Briefly, assembly was done stepwise on 2-chlorotrityl chloride polystyrene resin at a 0.05 mmol or 0.1 mmol scale, using Fmoc-based Solid Phase Peptide Synthesis (SPPS) on a PTI Symphony synthesizer. 20% piperidine in DMF was used to remove the Fmoc protecting group and free amine was coupled using tenfold excess of Fmoc amino acids and HCTU/DIEA activation in NMP/DMF (3 × 15 min). Peptide de-protections and cleavages from the resin were done with TFA/H<sub>2</sub>O/1,3-dimethoxybenzene (DMB)/TIS/2,2'-(Ethylenedioxy) diethanethiol (DODT) 85.1/5/2.5/3.7/3.7 (vol.). They were then precipitated with cold diethyl ether. Oxidative folding of the crude linear peptides were successfully conducted at RT in the conditions optimized for HwTx-IV using a peptide concentration of 0.1 mg/ml in a 0.1 M Tris buffer at pH 8.0 containing 10% of DMSO. The amino acid sequences of HwTx-IV and its 19 analogues are shown in **Figure 1**, along with the molecular masses after folding/oxidation.

### Cell Cultures

CHO cells stably expressing the hNa<sub>v</sub>1.7 channel subtype were cultured in Dulbecco's Modified Eagle's Medium F-12 (DMEM/F12) supplemented with 10% fetal bovine serum, 2 mM glutamine, 200 µg/ml hygromycin B, 10 U/ml penicillin and 10 µg/ml streptomycin (Gibco, Grand Island, NY). HEK-293 cells stably expressing the hNa<sub>v</sub>1.6 channel subtype were cultured in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% fetal calf serum, 1 mM pyruvic acid, 4.5 g/L glucose,

2 mM glutamine, 800 µg/ml G418, 10 U/ml penicillin and 10 µg/ml streptomycin (Gibco, Grand Island, NY). All cell lines were incubated at 37°C in a 5% CO<sub>2</sub> atmosphere. For electrophysiological recordings, cells were detached with trypsin, and floating single cells were diluted (~300,000 cells/ml) in an extracellular solution containing (in mM): 140 NaCl, 4 KCl, 2 CaCl<sub>2</sub>, 1 MgCl<sub>2</sub>, 5 glucose and 10 HEPES (pH 7.4, osmolarity 298 mOsm), that was used throughout the experiments.

### Pharmacological Applications Using the Automated Patch-Clamp System

HwTx-IV analogues were investigated on CHO cells expressing hNa<sub>v</sub>1.7 channel and HEK-293 cells expressing hNa<sub>v</sub>1.6 channel using the automated patch-clamp system from Nanion (SyncroPatch 384 PE; München, Germany). Chips with single-hole and high-resistance (~6–7 MΩ) were used for both cell lines. Voltage pulses and whole-cell recordings were achieved using the PatchControl384 v1.5.2 software (Nanion) and the Biomek v1.0 interface (Beckman Coulter). Prior to recordings, dissociated cells were shaken at 200 RPM in a cell hotel reservoir at 10°C. After cell catching, sealing, whole-cell formation, liquid application, recording, and data acquisition were all performed sequentially and automatically. The intracellular solution contained (in mM): 110 CsF, 10 CsCl, 10 NaCl, 1 MgCl<sub>2</sub>, 1 CaCl<sub>2</sub>, 10 EGTA and 10 HEPES (pH 7.2, osmolarity 280 mOsm), and the extracellular solution was as described above. Whole-cell experiments were done at -100 mV holding potential and at room temperature (18–22°C), while currents triggered at either -10 or 0 mV test potential were sampled at 20 kHz. Stimulation frequency was set



at 0.2 Hz. Each HwTx-IV analogue was prepared at various concentrations in the extracellular solution, itself supplemented with 0.3% of bovine serum albumin (BSA), also added in the control solution. The peptides were distributed in 384-well microplates according to the number of compounds to be tested (generally four), the concentration range defined from the IC<sub>50</sub> of wild-type HwTx-IV, and the number of cells desired for each experimental condition. Compound solutions were diluted 3 times in the patch-clamp recording well by adding 30–60 µl external solution, to reach the final reported concentration and the test volume of 90 µl. For establishing dose-response curves, the compounds were tested at a test potential of either –10 or 0 mV for 50 ms with a pulse every 5 s. Percentages of current inhibition were measured at steady-state of blockage or at the end of a 15-min application time.

## Docking of Toxin Analogues Onto hNa<sub>v</sub>1.7 Channel Structure

Based on the hNa<sub>v</sub>1.7 VSD2-Na<sub>v</sub>Ab chimera channel structure, obtained by cryo-electron microscopy (PDB: 6N4R), and the 3D solution structure of HwTx-IV (PDB code 1MB6), obtained by 2D <sup>1</sup>H-NMR, wild-type amidated or carboxylated HwTx-IV was docked using ZDOCK v.3.0.2. Docking was used because the CryoEM structures reported in the literature lack the HwTx-IV structure in the referred PDB code (6J8G or 6J8H) (Shen et al., 2019) or because the region of interaction is not resolved according to the PDB code (6W6O) (Gao et al., 2020). Also, the earlier docking models, reported after crosslinking experiments, were not publicly available (Tzakoniati et al., 2020). Residues previously reported as implicated in HwTx-IV/hNa<sub>v</sub>1.7 VSD2 channel interaction (T<sup>518</sup>-F<sup>761</sup> and S<sup>807</sup>-E<sup>817</sup> on hNa<sub>v</sub>1.7 VSD2 and S<sup>25</sup>-I<sup>35</sup> on HwTx-IV) have been used to guide docking. Only K<sup>32</sup> of HwTx-IV has been selected as a conserved binding site residue for all analogues. Interactions between HwTx-IV and hNa<sub>v</sub>1.7 VSD2 amino acids have been analyzed using Discovery Studio (Dassault System Biovia) and the resulting 3D structures were drawn with the PyMOL software (The PyMOL Molecular Graphics System, v.2.0 Schrödinger, LLC).

## RESULTS

### Chemical Syntheses of HwTx-IV Analogues

Overall, nineteen analogues were successfully chemically synthesized using fmoc solid-phase peptide chemistry. Proper folding and oxidation of the peptides were evaluated by checking the correspondence between theoretical *versus* experimental masses as assessed by mass spectrometry (Figure 1). Ten analogues with single amino acid substitution were produced, followed by three analogues with double amino acid substitutions, two analogues with triple amino acid substitutions, and four analogues with modifications at the C-terminus. It was remarkable that none of the analogues posed problems for the folding of the peptide (RP-HPLC data and yield calculations, not shown) in spite of the fact that this peptide folds according to an ICK motif and that some analogues harbor up to three substitutions.

### Single Amino Acid-Substituted HwTx-IV Analogues Affecting Peptide Potency Onto hNa<sub>v</sub>1.7


Some of the earlier SAR investigations were performed with recombinant HwTx-IV peptides that are lacking C-terminal amidation (Minassian et al., 2013; Sermadiras et al., 2013; Rahnama et al., 2017; Neff et al., 2020), an amidation that is however crucial for potency onto hNa<sub>v</sub>1.7 (Revell et al., 2013). Also, in some of these reports, several recombinant analogues carried additional residues at either the N- or the C-terminus, all susceptible to affect peptide potency (Minassian et al., 2013; Sermadiras et al., 2013; Neff et al., 2020). Therefore, some analogues that were deemed of interest were tested herein on hNa<sub>v</sub>1.7 to confirm or infirm earlier reports in more standardized conditions (normal C-terminal amidation and absence of extra-non-native amino acid residues).

As shown in Figure 2A, we identified 3 analogues of HwTx-IV with mono-substituted amino acid residues (HwTx-IV E<sup>1</sup>G, HwTx-IV E<sup>4</sup>G, and HwTx-IV K<sup>18</sup>A) that possessed better potency towards the hNa<sub>v</sub>1.7 channel subtype. This was best exemplified by current traces illustrating the percentage of inhibition of Na<sup>+</sup> inward current at 10 nM peptide analogue (upper panel). One substitution brings in a remarkable improvement in HwTx-IX potency onto hNa<sub>v</sub>1.7: the E<sup>4</sup>G mutation with an IC<sub>50</sub> value which is 6.2-fold better than wild-type HwTx-IV (Figure 2B). The improvements made by the E<sup>1</sup>G or the K<sup>18</sup>A mutations were milder. All other mono-substituted analogues of HwTx-IV had decreased potencies towards hNa<sub>v</sub>1.7 with most of them producing almost no inhibition at all at 10 nM (Figure 2A). As shown by the average dose-response curves, the E<sup>4</sup>K mutation was most conservative, followed by the Y<sup>33</sup>W one. As expected from earlier reports (see Discussion section), the most disruptive single point mutation was K<sup>32</sup>N, while R<sup>26</sup>A, R<sup>26</sup>Q and T<sup>28</sup>W had milder effects on the HwTx-IV loss of potency (Figure 2C and Table 1).

### Single Amino Acid-Substituted HwTx-IV Analogues Affecting Peptide Potency Onto hNa<sub>v</sub>1.6

We next tested wild-type HwTx-IV along with its nine mono-substituted amino acid analogues onto the Na<sup>+</sup> current carried by hNa<sub>v</sub>1.6 subtype. We used a concentration of 100 nM to evaluate which analogue performed better than the wild-type HwTx-IV on Na<sup>+</sup> current inhibition. As shown in Figure 3A, 100 nM of HwTx-IV inhibits less than 50% of hNa<sub>v</sub>1.6 current, which is coherent with the average IC<sub>50</sub> value of 226.6 nM (Figure 3B). As such, natively, without any sequence alteration, HwTx-IV is 22.9-fold more potent on hNa<sub>v</sub>1.7 than on hNa<sub>v</sub>1.6 channel subtype (starting selectivity ratio as detailed in Figure 4). Five out of nine mono-substituted amino acid analogues displayed better potency on hNa<sub>v</sub>1.6 than wild-type HwTx-IV itself: HwTx-IV E<sup>1</sup>G (3.6-fold better), HwTx-IV E<sup>4</sup>G (36-fold better), HwTx-IV E<sup>4</sup>K (10.8-fold better), HwTx-IV E<sup>4</sup>R (20.4-fold better) and HwTx-IV Y<sup>33</sup>W (1.1-fold better).



In contrast, 4 mono-substituted analogues of HwTx-IV displayed lower potencies for hNa<sub>v</sub>1.6 compared to wild-type HwTx-IV. Two analogues still affected hNa<sub>v</sub>1.6 with



**TABLE 1 |** Mean ± S.E.M. of IC<sub>50</sub> values (n cells) determined from the concentration-response curves of the effects of wild-type (WT) HwTx-IV and analogues on hNa<sub>v</sub>1.7 and hNa<sub>v</sub>1.6 peak currents.


	hNa <sub>v</sub> 1.7				hNa <sub>v</sub> 1.6			
	Mean IC <sub>50</sub> (nM)	S.E.M.	n	versus WT	Mean IC <sub>50</sub> (nM)	S.E.M.	n	versus WT
<b>HwTx-IV WT</b>	<b>9.9</b>	<b>1.1</b>	<b>5-7</b>	<b>1.00</b>	<b>226.6</b>	<b>1.2</b>	<b>6-10</b>	<b>1.00</b>
HwTx-IV E <sup>1</sup> G	6.9	1.1	3-7	0.70	62.4	1.2	4-7	0.28
HwTx-IV E <sup>4</sup> G	1.6	1.1	6-7	0.16	6.3	1.3	3	0.03
HwTx-IV E <sup>4</sup> K	14.2	1.2	2-8	1.43	20.9	1.3	3	0.09
HwTx-IV E <sup>4</sup> R	—	—	—	—	11.1	1.3	3-5	0.05
HwTx-IV K <sup>18</sup> A	6.4	1.1	3-7	0.65	416.2	1.3	3-7	1.84
HwTx-IV K <sup>32</sup> N	5752.0	1.6	4-7	581	>1,000	-	2-7	>4.4
HwTx-IV R <sup>26</sup> A	111.7	1.2	4-7	11.28	614.2	1.2	3-6	2.71
HwTx-IV R <sup>26</sup> Q	141.4	1.1	3-5	14.28	>1,000	-	2-8	>4.4
HwTx-IV T <sup>28</sup> W	114.1	1.2	4-6	11.53	—	—	—	—
HwTx-IV Y <sup>33</sup> W	22.2	1.2	3-7	2.24	207.2	1.3	3-9	0.91
HwTx-IV E <sup>1</sup> G/E <sup>4</sup> G	2.0	1.1	3-7	0.20	3.7	1.2	1-2	0.02
HwTx-IV E <sup>4</sup> R/K <sup>32</sup> N	>1,000	—	2-8	>101	>333	-	2-3	>1.5
HwTx-IV E <sup>4</sup> K/R <sup>26</sup> Q	3.6	1.1	2-8	0.36	3.4	1.5	5	0.02
HwTx-IV E <sup>1</sup> G/E <sup>4</sup> G/K <sup>18</sup> A	0.8	1.1	3-7	0.08	4.9	1.2	4	0.02
HwTx-IV E <sup>4</sup> K/R <sup>26</sup> A/Y <sup>33</sup> W	0.9	1.1	3-8	0.09	86.4	1.4	5	0.38
HwTx-IV C <sub>ter</sub> COOH	176.3	1.1	5-8	17.81	1,230.0	1.3	2-6	5.43
HwTx-IV K <sup>36</sup> /C <sub>ter</sub> COOH	29.0	1.2	2-6	2.93	1,334.0	1.6	2-7	5.89
HwTx-IV E <sup>4</sup> K/C <sub>ter</sub> COOH	11.9	1.1	4-7	1.20	135.7	1.3	4-10	0.60
HwTx-IV E <sup>4</sup> G/C <sub>ter</sub> COOH	14.9	1.1	2-7	1.51	55.1	1.2	5-10	0.24

*Bold value highlight the wild-type peptide for a better comparison with analogues.*



measurable potency: HwTx-IV K<sup>18</sup>A (1.8-fold reduction) and HwTx-IV R<sup>26</sup>A (2.7-fold reduction). The two other analogues have a too significant shift in IC<sub>50</sub> values to be measured

(HwTx-IV K<sup>32</sup>N and HwTx-IV R<sup>26</sup>Q, both >1  $\mu$ M). In conclusion, among mono-substituted HwTx-IV analogues, the most interesting ones are those with a sequence



alteration at position 4 (HwTx-IV E<sup>4</sup>G or HwTx-IV E<sup>4</sup>K) or 18 (HwTx-IV K<sup>18</sup>A) since these analogues led to an improved potency for hNa<sub>v</sub>1.7 (Figure 4 and Table 1). This improved potency of HwTx-IV K<sup>18</sup>A was associated to an improved selectivity *versus* hNa<sub>v</sub>1.6, which is likely to reduce the molecule side effects resulting from impairment of neuromuscular transmission. In contrast, HwTx-IV E<sup>4</sup>G and HwTx-IV E<sup>4</sup>K tended clearly to abolish differences between potencies for hNa<sub>v</sub>1.7 and hNa<sub>v</sub>1.6, allowing to produce pan-Na<sub>v</sub> blockers.


### Combining Several Substitutions Onto HwTx-IV May Also Bring Competitive Advantages in Improving hNa<sub>v</sub>1.7/hNa<sub>v</sub>1.6 Selectivity Ratio

We first attempted to combine two or three substitutions that individually were shown to be beneficial for the potency of HwTx-IV onto hNa<sub>v</sub>1.7 (E<sup>1</sup>G, E<sup>4</sup>G, K<sup>18</sup>A) or hNa<sub>v</sub>1.6 (E<sup>1</sup>G, E<sup>4</sup>G) (Figures 2A, Figure 3A). The double mutant HwTx-IV E<sup>1</sup>G/E<sup>4</sup>G produced

IC<sub>50</sub> values of  $2.0 \pm 1.1$  nM for hNa<sub>v</sub>1.7 (Figure 5A) and  $3.7 \pm 1.2$  nM for hNa<sub>v</sub>1.6 (Figure 5B). These values are more or less similar to those brought by the single E<sup>4</sup>G substitution, but with a slight improvement introduced by the additional E<sup>1</sup>G mutation in the case of the hNa<sub>v</sub>1.6 subtype. Adding the K<sup>18</sup>A substitution onto the HwTx-IV E<sup>1</sup>G/E<sup>4</sup>G mutant (HwTx-IV E<sup>1</sup>G/E<sup>4</sup>G/K<sup>18</sup>A) did not result in a marked benefit of the peptide potency onto both targets (Figure 5). This was expected since the effect of K<sup>18</sup>A substitution was similar to that of the E<sup>1</sup>G substitution and 4-fold lower to that of the E<sup>4</sup>G substitution on the hNa<sub>v</sub>1.7 subtype. In contrast to the E<sup>1</sup>G and E<sup>4</sup>G substitutions, it did not, by itself, drastically change HwTx-IV potency for the hNa<sub>v</sub>1.6 subtype (Figures 2A, 3A and Table 1). In a second attempt, we combined two substitutions that individually were, *a priori*, not favorable for potency improvements, *i.e.* R<sup>26</sup>Q and E<sup>4</sup>K, although the latter was advantageous for hNa<sub>v</sub>1.6 (Figures 2B, 3).

Nevertheless, the combination of these two substitutions had some logic since they introduce compensatory charge alterations in the whole peptide sequence (loss of negative charge and gain of positive charge at position 4, the latter being





**FIGURE 6 |** Effects of HwTx-IV and its four C-terminal-altered analogues on CHO and HEK-293 cells stably expressing the hNav1.7 and hNav1.6 channel subtypes, respectively. **(A)** Representative traces of current flowing through the hNav1.7 (upper traces) or hNav1.6 (lower traces) subtype, recorded before (black traces) and after (coloured traces) exposure to 10 nM (upper traces) or 100 nM (lower traces) of the indicated peptide. Vertical and horizontal scales (applied to all traces) are 0.5 nA and 5 ms (all traces). **(B)** and **(C)** Concentration-response curves of the effects of HwTx-IV (white circles) and analogues (coloured circles) on hNav1.7 (B) and hNav1.6 (C) peak currents, recorded during  $-10$ -mV and  $0$ -mV test-pulse voltages, respectively. Each value, expressed relatively to that obtained before peptide application, represents the mean  $\pm$  S.E.M. of data obtained from 1–8 cells. The theoretical curves correspond to data point fits with the mean  $IC_{50}$  values indicated in **Table 1**.

compensated by the loss of positive charge at position 26). Interestingly, the double mutant HwTx-IV  $E^4K/R^{26}Q$  yielded excellent potency of the peptide for both channel subtypes (**Figure 5**). Noteworthy, this double mutation yielded yet another HwTx-IV analogue, along with HwTx-IV  $E^1G$  and HwTx-IV  $E^1G/E^4G$ , that possesses similar and high potencies (low nanomolar range) for both channel subtypes. Introducing a third substitution at position 33, which individually was quite neutral for both channel types (**Figures 2B, 3A**), added to those at positions 4 and 26, did not markedly modify the potency for hNav1.7 but produced a 25.4-fold decrease of that for hNav1.6 compared to HwTx-IV  $E^4K/R^{26}Q$  (**Figures 4, 5** and **Table 1**).

We also attempted to produce a charge compensation of the  $E^4R$  mutation, similar to  $R^{26}Q$  in the HwTx-IV  $E^4K/R^{26}Q$  analogue, by an analogous  $K^{32}N$  mutation. As observed by the  $IC_{50}$  values ( $>333$  nM in both cases, **Figure 5**), this strategy did not work because of the crucial role of  $K^{32}$  in the pharmacophore of HwTx-IV (Minassian et al., 2013; Revell et al., 2013).


## Examining the Importance of HwTx-IV C-Terminal Amidation on Peptide Potency for Both Nav Subtypes

We first replaced the C-terminal amidated residue by a carboxylated version of the residue. As shown in **Figure 6**, the potency of HwTx-IV  $C_{ter}COOH$  was significantly reduced

on both channel subtypes (by 18-fold on hNav1.7 and by 5-fold on hNav1.6). These results confirm an earlier report on the role of amidation in HwTx-IV activity (Revell et al., 2013). It is likely that the consequence of amidation on HwTx-IV potency may be explained by the presence of a positive charge at the C-terminus. Indeed, the potency of HwTx-IV  $K^{36}/C_{ter}COOH$ , in which the peptide is still carboxylated but integrates an additional K residue, positively-charged on its side-chain, was greatly improved for the hNav1.7 but not hNav1.6 subtype (**Figure 6**). Similar results were obtained when  $E^4$  was mutated in order to remove the negative charge ( $E^4G$ ), or to replace the negative charge by a positive one ( $E^4K$ ) at this position. Hence, these results are reminiscent of those obtained with the deleterious  $R^{26}$  mutation that also could rescue HwTx-IV potency by an additional  $E^4$  substitution (**Figure 5**). Considering the distal positioning of  $E^4$  relatively to  $R^{26}$  or to  $C_{ter}CONH_2$ , the beneficial effect of  $E^4$  substitution is likely due to a rebalance in dipole moment.

## DISCUSSION

In this manuscript, our goal was to produce HwTx-IV analogues with increased potencies for hNav1.7, hNav1.6, or both. Based on previous studies, we examined a number of analogues for which only a few observations were reported. Finding analogues with



potent activity for hNa<sub>v</sub>1.6 was a particular challenge because the original hNa<sub>v</sub>1.7/hNa<sub>v</sub>1.6 selectivity ratio of 23 was largely in favor of hNa<sub>v</sub>1.7. This endeavor was successful, thanks to mono-, double- or triple-amino acid substitutions. Despite an excellent starting potency for hNa<sub>v</sub>1.7 (mean IC<sub>50</sub> value of 9.9 nM, in accordance with earlier reports (Xiao et al., 2008; Revell et al., 2013; Sermadiras et al., 2013; Rahnema et al., 2017; Agwa et al., 2020)), it was still possible to further improve the activity of HwTx-IV on this channel subtype by mutations of specific positions, *i.e.* E<sup>1</sup>, E<sup>4</sup>, K<sup>18</sup>, E<sup>1</sup>/E<sup>4</sup>, E<sup>4</sup>/R<sup>26</sup>, E<sup>1</sup>/E<sup>4</sup>/K<sup>18</sup> and E<sup>4</sup>/R<sup>26</sup>/Y<sup>33</sup>. The maximal optimization factor on potency was 11–12-fold (with the triple mutants HwTx-IV E<sup>4</sup>K/R<sup>26</sup>A/Y<sup>33</sup>W and E<sup>1</sup>G/E<sup>4</sup>G/K<sup>18</sup>A). Concerning hNa<sub>v</sub>1.6, the challenge was easier to overcome since the starting potency of HwTx-IV was quite low for this channel subtype (mean IC<sub>50</sub> value of 227 nM). The best analogues were (ranked by decreasing order of improved potency): HwTx-IV E<sup>4</sup>K/R<sup>26</sup>Q, E<sup>1</sup>G/E<sup>4</sup>G, E<sup>1</sup>G/E<sup>4</sup>G/K<sup>18</sup>A, E<sup>4</sup>G, and E<sup>4</sup>R, all under or close to 10 nM of IC<sub>50</sub> value (improvement by 20- to 67-fold), and HwTx-IV E<sup>4</sup>K, E<sup>1</sup>G, E<sup>4</sup>K/R<sup>26</sup>A/Y<sup>33</sup>W and Y<sup>33</sup>W with less noticeable improvement (by 1.1- to 11-fold). Five analogues had comparable potencies (low nanomolar range) on the two channel subtypes: HwTx-IV E<sup>4</sup>G, E<sup>4</sup>K, E<sup>1</sup>G/E<sup>4</sup>G, E<sup>4</sup>K/R<sup>26</sup>Q and E<sup>1</sup>G/E<sup>4</sup>G/K<sup>18</sup>A. These data highlight the importance of the type of amino acid at position 4 in obliterating the difference between potencies for the two channel subtypes. However, substitutions at position 4 were not all equipotent since E<sup>4</sup>G was more potent than E<sup>4</sup>K on hNa<sub>v</sub>1.7, possibly because G<sup>4</sup> was

more lipid-friendly than the charged K<sup>4</sup>, at least for this channel type. With regard to selectivity ratios, two observations can be made: 1) some analogues display a large increased selectivity ratio in favor of hNa<sub>v</sub>1.7 (K<sup>18</sup>A, E<sup>4</sup>K/R<sup>26</sup>A/Y<sup>33</sup>W and K<sup>36</sup>/C<sub>ter</sub>COOH, all above 46), while keeping excellent apparent affinity on this subtype, and 2) other analogues display a significant reduction in selectivity ratio in favor of hNa<sub>v</sub>1.6 (E<sup>4</sup>K, E<sup>1</sup>G/E<sup>4</sup>G and E<sup>4</sup>K/R<sup>26</sup>Q, less than 2) (Figure 4 and Table 1).

## Fitting Our Data With Earlier Reports

HwTx-IV has been the focus of several SAR studies, mainly with the aim to improve selectivity for hNa<sub>v</sub>1.7 *versus* other channel subtypes. Indeed, HwTx-IV suffers from toxicity issues *in vivo*, most of the symptoms being however related to an action on Na<sub>v</sub>1.6 (Gonçalves et al., 2018b), a subtype that was not the topic of the earlier SAR investigations. In our study, we found HwTx-IV to be quite potent on hNa<sub>v</sub>1.7 with a mean IC<sub>50</sub> value of 9.9 nM, which compares favorably with the literature as far as the peptide was produced chemically (Sermadiras et al., 2013; Neff et al., 2020). Hence, when HwTx-IV was produced according to recombinant approaches, a significant decrease of potency was always observed (Minassian et al., 2013; Sermadiras et al., 2013; Neff et al., 2020). This decrease may be explained by two factors. The first one is simply the absence of C-terminal amidation. Indeed, in our study, the loss of C-terminal amidation results in a 18-fold decline of potency for hNa<sub>v</sub>1.7 of the chemically-synthesized peptide (see Figures 6A,B and Table 1). This

finding is coherent with an earlier report (Revell et al., 2013). According to our docking simulations, this loss of potency should be due to a reduction of the number of interaction points with hNa<sub>V</sub>1.7 (Figure 7). The second one may be due to incomplete or improper folding of the peptide, which is less controlled using recombinant production than chemical synthesis. Indeed, a comparison between the potencies of amidated synthetic and carboxylated recombinant HwTx-IV shows an even more important decline of potency for hNa<sub>V</sub>1.7 in one study (up to 66-fold) (Sermadiras et al., 2013).

Until 2018, to our knowledge, there was no report illustrating the activity of HwTx-IV on Na<sub>V</sub>1.6, a tetrodotoxin (TTX)-sensitive channel subtype present in the Ranvier nodes of  $\alpha$ -motoneurons (Rahnama et al., 2017). Since then, Gonçalves and collaborators showed that the peptide blocks this channel subtype with an IC<sub>50</sub> value slightly below 100 nM (Gonçalves et al., 2018b), which is not markedly different from the value we obtained in standardized conditions. These results allow a better understanding of the mechanisms of *in vivo* motor toxicity side-effects of HwTx-IV (Liu et al., 2014b; Deuis et al., 2016). More recently, a large-scale SAR investigation was performed with the help of recombinant HwTx-IV analogues tested on both hNa<sub>V</sub>1.7 and hNa<sub>V</sub>1.2 and, occasionally, on a larger panel of other hNa<sub>V</sub> subtypes, without extensively reporting on the activity of wild-type HwTx-IV on hNa<sub>V</sub>1.6 (Neff et al., 2020).

## Lessons From SAR Studies to Determine HwTx-IV Critical Residues for hNa<sub>V</sub>1.7 Block

Several objectives have been pursued in HwTx-IV SAR studies, namely: 1) defining the site of binding of the peptide onto the channel, 2) determining the peptide pharmacophore to get a clear picture of the peptide docking on the channel, 3) clarifying the relationships between the peptide and the channel lipid environment, and 4) improving both the peptide potency and selectivity. The two first aims are closely interconnected.

As stated above (see Introduction section), the blocking effect of HwTx-IV on hNa<sub>V</sub>1.7 is due to its binding onto the voltage-sensor of domain II of the channel subtype. As a result, HwTx-IV traps this voltage-sensor in the inward closed configuration (Xiao et al., 2008), which logically inhibits the gating currents conveyed by domain II movement (Xiao et al., 2014). More precisely, the binding occurs onto the S3-S4 linker of domain II. Amino acid residues D<sup>816</sup> and E<sup>818</sup> of hNa<sub>V</sub>1.7 are involved in this binding since mutating these residues reduces the peptide potency by over 60-fold (Xiao et al., 2008; Xiao et al., 2010). Conversely, mutations in hNa<sub>V</sub>1.4 (N<sup>655</sup>D, Q<sup>657</sup>E) or hNa<sub>V</sub>1.5 (R<sup>812</sup>D, S<sup>814</sup>E), aimed at instating hNa<sub>V</sub>1.7 residues, were shown to produce a gain of HwTx-IV potency on these channel subtypes. A larger scale investigation refined the pictures of hNa<sub>V</sub>1.7 channel residues involved in binding HwTx-IV, and identified a total of five amino acids, one belonging to the S1-S2 linker (E<sup>753</sup>) and the four remaining ones being part of the S3-S4 linker (E<sup>811</sup>, L<sup>814</sup>, D<sup>816</sup> and E<sup>818</sup>) (Xiao et al., 2011). This cluster of 5 residues defines an EELDE motif (herein defined as E<sub>1</sub>E<sub>2</sub>L<sub>3</sub>D<sub>4</sub>E<sub>5</sub> for convenience), and thus suggests that the

interaction largely relies on electrostatic interactions. It is worth noting that HwTx-IV acts in an odd manner on the voltage-sensor of domain II since no clear voltage-dependent effect of the toxin has been reported in the literature, except maybe for very strong depolarizing pulses meant to favor toxin dissociation (Gonçalves et al., 2018b). However, SAR studies may potentially reveal new HwTx-IV analogues displaying voltage-dependent effect and, therefore, it would be of interest to check the voltage-dependence or lack thereof of the best toxin analogues.

To get a hint of how the toxin interacts with its binding site on hNa<sub>V</sub>1.7, several initiatives were launched, including 1) complete SAR investigation of the contribution of HwTx-IV residues, 2) cryo-EM analyses and 3) the use of photocrosslinking probes derived from HwTx-IV.

Concerning the SAR studies on HwTx-IV, the emerging picture is that the couple of residues, W<sup>30</sup> and K<sup>32</sup>, are required to preserve peptide activity onto hNa<sub>V</sub>1.7 (Minassian et al., 2013; Revell et al., 2013; Neff et al., 2020). Modest reduction in potencies (<10-fold) are observed for F<sup>6</sup>A, K<sup>18</sup>A, R<sup>26</sup>A, K<sup>27</sup>A and Y<sup>33</sup>A, all other substitutions being neutral (L<sup>3</sup>A, S<sup>9</sup>A, P<sup>11</sup>A, S<sup>12</sup>A, D<sup>14</sup>A, L<sup>22</sup>A, S<sup>25</sup>A) or modestly beneficial (E<sup>1</sup>A, E<sup>4</sup>A, K<sup>7</sup>A, N<sup>10</sup>A, Q<sup>12</sup>A, S<sup>19</sup>A, S<sup>20</sup>A, K<sup>21</sup>A, V<sup>23</sup>A, T<sup>28</sup>A, R<sup>29</sup>A, Q<sup>34</sup>A and I<sup>35</sup>A) (Revell et al., 2013). These conclusions are relatively well supported by the data of Minassian and collaborators, although these authors use mainly recombinant HwTx-IV, lacking C-terminal amidation and possessing some extra residues in the sequence (Minassian et al., 2013).

For the cryo-EM investigation, the authors replaced the S3-S4 linker of the NaChBac sequence by the one of hNa<sub>V</sub>1.7, and illustrated the interaction of HwTx-IV with this chimera channel (Gao et al., 2020). They concluded that K<sup>27</sup> is located close to E<sub>2</sub>, that K<sup>32</sup> interacts with E<sub>2</sub>, and that Y<sup>33</sup> forms hydrogen bonds with the backbone groups of L<sub>3</sub> and an adjacent A residue. Globally, positive residues of HwTx-IV R<sup>26</sup>, K<sup>27</sup> and K<sup>32</sup> are in close vicinity of the channel, whereas the hydrophobic residues I<sup>5</sup>, F<sup>6</sup> and W<sup>30</sup> are immersed in the lipid bilayer. However, chimera approaches may not perfectly reconstitute the wild-type HwTx-IV binding site. For instance, the K<sup>32</sup>A mutation in HwTx-IV barely impacts peptide effect on the chimera channel (Gao et al., 2020) while, in our study, a similar K<sup>32</sup>N mutation prevents the HwTx-IV-induced block of hNa<sub>V</sub>1.7 (see Figures 2A,C).

The model of interaction was refined by an elegant study that used photocrosslinking probes derived from HwTx-IV and the full-length hNa<sub>V</sub>1.7 (Tzakoniati et al., 2020). L-photomethionine was introduced in HwTx-IV sequence to replace one of 6 residues (I<sup>5</sup>, K<sup>7</sup>, K<sup>21</sup>, K<sup>27</sup>, R<sup>29</sup>, I<sup>35</sup>) deemed proximal to other crucial residues involved in toxin interaction. Among these 6 new analogues, only 2 photoprobes, with substitutions at position 27 or 29, crosslinked to hNa<sub>V</sub>1.7 with high efficiency: photoprobe 27 to the peptide <sup>808</sup>SLVE<sup>811</sup> that belongs to the S3 helix, and photoprobe 29 to the motif <sup>758</sup>TEEF<sup>761</sup> of the S1-S2 loop. These data support the concept that K<sup>27</sup> of HwTx-IV interacts with E<sub>2</sub> of the E<sub>1</sub>E<sub>2</sub>L<sub>3</sub>D<sub>4</sub>E<sub>5</sub> motif and R<sup>29</sup> with a negatively charged residue of S1-S2 loop (most likely E<sup>759</sup>) that differs from E<sub>1</sub>. In this way, they contradict the docking model of Minassian and collaborators that suggested an interaction between K<sup>27</sup> and E<sub>5</sub> rather than E<sub>2</sub>

(Minassian et al., 2013). The refined docking model supports the idea that F<sup>6</sup>, W<sup>30</sup> and Y<sup>33</sup> interact with the lipid bilayer. The crucial K<sup>32</sup> residue would also interact with E<sub>2</sub> of the binding motif (**Figure 7**). According to a summary of this docking, HwTx-IV I<sup>35</sup> would also interact with L<sub>3</sub>. The fact that the R<sup>29</sup>A mutation hardly affects HwTx-IV potency for hNa<sub>v</sub>1.7 (Revell et al., 2013) would indicate that HwTx-IV interaction with the S1-S2 loop has little functional implication for the mechanism of channel block.

How do these data fit with what we know about the toxin/lipid interactions? Part of the conclusions drawn about the definition of HwTx-IV pharmacophore was complexified by the fact that the peptide, besides binding to the targeted channel, also requires partitioning into the lipid membrane for the access to the binding site. This has led to the concept of a three-way interaction for the mechanism of action of the toxin (Agwa et al., 2017), which complicates the design of new analogues as exemplified by the case of protoxin II (Montnach et al., 2021). From the various docking models or Cryo-EM structure previously published (Minassian et al., 2013; Gao et al., 2020; Neff et al., 2020; Tzakoniati et al., 2020), several residues of HwTx-IV are susceptible to directly interact with lipids (I<sup>5</sup>, F<sup>6</sup>, R<sup>29</sup>, W<sup>30</sup>, Y<sup>33</sup>). By mutating non-essential residues for the interaction with hNa<sub>v</sub>1.7, it was shown that the combined mutagenesis of E<sup>1</sup>G, E<sup>4</sup>G, F<sup>6</sup>W and Y<sup>33</sup>W produces an HwTx-IV analogue with increased ability to bind membrane lipids, a property that in turns explains the increased potency of this analogue for hNa<sub>v</sub>1.7 (Agwa et al., 2017). In this mutated analogue, two of the residues are suspected to interact directly with lipids (W<sup>30</sup> and Y<sup>33</sup>), whereas the two other residues may hamper toxin partitioning into the lipid bilayer because of the negative charges associated to E residues. In wild-type HwTx-IV, the E<sup>4</sup> residue forms a salt bridge with K<sup>7</sup>, possibly stabilizing the peptide fold. Mutation of E<sup>4</sup> to G<sup>4</sup> appears to increase the potency for hNa<sub>v</sub>1.7 by improving permeability into the membrane (Klint et al., 2015) and the interaction surface of the peptide with the channel (**Figure 7**). If in addition the authors mutate a non-essential positively charged residue, R<sup>26</sup>A, a further improvement in lipid affinity is observed, also accompanied by an additional improvement in toxin potency for hNa<sub>v</sub>1.7 (Agwa et al., 2020). The lesson from this work is that it is possible to enhance lipid partitioning of a toxin either directly by acting on residues that interact with lipids or indirectly by altering the global hydrophobicity profile of the peptide.

It is therefore clear from SAR and docking studies that there are three ways to alter potency and selectivity of HwTx-IV, namely by 1) mutating residues involved in binding onto the channel itself, 2) substituting residues involved in peptide/lipid interactions, and 3) indirectly playing on the surface charge or hydrophobicity profile of the peptide. By intervening on one or several of these factors, it is possible to modify peptide selectivity. Earlier efforts in that direction were performed by altering the relative potency of HwTx-IV for hNa<sub>v</sub>1.7 and hNa<sub>v</sub>1.2. For instance, the R<sup>29</sup>A mutation was shown to increase potency for hNa<sub>v</sub>1.7 but to considerably reduce the one for hNa<sub>v</sub>1.2 (Minassian et al., 2013). This was possibly due to a weakened electrostatic interaction with the EELNE motif of hNa<sub>v</sub>1.2. It was

also found that the triple mutant E<sup>1</sup>G/E<sup>4</sup>G/Y<sup>33</sup>W (but as recombinant protein, *i.e.* without C-terminal amidation) has an improved IC<sub>50</sub> value for hNa<sub>v</sub>1.7, but without increased activity for either hNa<sub>v</sub>1.1, hNa<sub>v</sub>1.2, hNa<sub>v</sub>1.3, hNa<sub>v</sub>1.5 or hNa<sub>v</sub>1.6 (Rahnama et al., 2017). The analogue with 5 mutations (E<sup>1</sup>G/E<sup>4</sup>G/F<sup>6</sup>W/R<sup>26</sup>A/Y<sup>33</sup>W), not only possessed a better potency for hNa<sub>v</sub>1.7, but also an improved selectivity profile over hNa<sub>v</sub>1.2 (27 times more selective) (Agwa et al., 2020). The same analogue produced a hNa<sub>v</sub>1.7/hNa<sub>v</sub>1.6 selectivity ratio of 11.

## The Pros and Cons of SAR Studies Using HwTx-IV as Template for Selectivity Alterations

The first issue to be discussed is without any doubts linked to the strategy employed to perform the SAR investigation. With an economical perspective, it appears logic to search for analogues that are produced by recombinant means as production costs are lower and yields higher. This method was chosen by several private actors (MedImmune, Janssen Research and Development) but not all (Genentech). Clearly, a comparison of our data testing HwTx-IV analogues onto hNa<sub>v</sub>1.7 illustrates that the presence or not of the C-terminal amidation has important consequences on the interpretation of the SAR data. Interesting differences were noted. For instance, a substitution of T<sup>28</sup> was considered as beneficial for HwTx-IV potency towards hNa<sub>v</sub>1.7 in its recombinant form (Revell et al., 2013), while in our hands it was clearly detrimental. Also, while for both hNa<sub>v</sub>1.7 and hNa<sub>v</sub>1.6 channel subtypes, a HwTx-IV analogue with a carboxylated C-terminus is synonym of loss of potency, there are interesting compensatory mutations capable to restore peptide potency. Part of the beneficial effect of C-terminal amidation could be reproduced by the introduction of an extra K<sup>36</sup> residue, even if the C-terminus was still carboxylated (HwTx-IV K<sup>36</sup>/C<sub>ter</sub>COOH), suggesting that the amine function of the lateral chain could substitute favorably to the C-terminal amidation. Gain of potency for hNa<sub>v</sub>1.7 has been observed earlier if an extra Gly residue was incorporated at position 36 along with a C-terminal amidation (Sermadiras et al., 2013). The gain of potency we observed with HwTx-IV K<sup>36</sup>/C<sub>ter</sub>COOH was true for the hNa<sub>v</sub>1.7 target but not for the hNa<sub>v</sub>1.6 channel subtype, suggesting that the C-terminal amidation is a function of interest to create differences in selectivity between the two channel subtypes. These observations also question the “true” selectivity profiles published with recombinant HwTx-IV. Interestingly, mutation of E<sup>4</sup> also largely compensated for the loss of potency induced by the carboxylated C-terminus sequence. This time, the compensation at work was valid for both channel subtypes, suggesting that all efforts to build recombinant analogues with increased potencies for Na<sub>v</sub> channel subtypes should incorporate an E<sup>4</sup> mutation.

The question that arises now is how far one can expect to go with efforts dedicated at improving selectivity for a given Na<sub>v</sub> subtype? In the work we launched, the task was challenging since the beginning of the project because the crucial E<sub>1</sub>E<sub>2</sub>L<sub>3</sub>D<sub>4</sub>E motif is conserved in hNa<sub>v</sub>1.6. As a reminder, this motif becomes



E<sub>1</sub>E<sub>2</sub>L<sub>3</sub>N<sub>4</sub>E<sub>5</sub> in hNa<sub>V</sub>1.2, a channel subtype that was used as template for earlier SAR studies aimed at altering peptide selectivity. We are aware that these earlier efforts did not necessarily constitute the best guidelines for producing analogues differentially affecting hNa<sub>V</sub>1.7 and hNa<sub>V</sub>1.6 for two reasons: 1) the hNa<sub>V</sub>1.6 binding motif differs from the hNa<sub>V</sub>1.2 motif, and 2) the SAR studies were all performed with recombinant peptides that in addition possess several extra residues at their C-terminus (Neff et al., 2020). Nevertheless, positions that differentially affected the relative hNa<sub>V</sub>1.7/hNa<sub>V</sub>1.2 selectivity ratio when mutated were F<sup>6</sup>, K<sup>18</sup>, R<sup>26</sup>, K<sup>27</sup> and I<sup>35</sup> (Minassian et al., 2013). In contrast to two reports (Minassian et al., 2013; Neff et al., 2020), we found that the K<sup>26</sup> mutation preferentially lowers potency for hNa<sub>V</sub>1.7. More coherent was the fact that the K<sup>18</sup>A mutation improves potency for hNa<sub>V</sub>1.7 while slightly reducing it for hNa<sub>V</sub>1.6, although less drastically than for hNa<sub>V</sub>1.2 (Minassian et al., 2013). In our hands, the most interesting positions tested to differentially affect the selectivity between hNa<sub>V</sub>1.7 and hNa<sub>V</sub>1.6 were E<sup>1</sup>, E<sup>4</sup>, R<sup>26</sup> or the combinations E<sup>1</sup>/E<sup>4</sup>, E<sup>4</sup>/R<sup>26</sup> and E<sup>1</sup>/E<sup>4</sup>/K<sup>18</sup>. For obvious economic reasons, this was not an as exhaustive work as wished to test all possible positions of HwTx-IV sequence and perform a full comparison with the work of Neff and collaborators (Neff et al., 2020). Nevertheless, several conclusions could be reached. First, it is possible to further increase the potency of HwTx-IV for hNa<sub>V</sub>1.7 in spite of an excellent starting affinity. However, the optimization remains in the range of 10-fold maximum, suggesting that there are limitations reached maybe because of the complexity in the docking to the channel combined to the need for partitioning into the lipid membrane. Second, it is clearly possible to optimize potency for hNa<sub>V</sub>1.6 in spite of similarities in the target-binding site, and the extent of this optimization is greater for this subtype than for hNa<sub>V</sub>1.7. Thirdly, there must be differences somewhere in the docking onto hNa<sub>V</sub>1.6 compared to hNa<sub>V</sub>1.7 because some analogues clearly affected the selectivity ratios. One in particular looks as a promising hNa<sub>V</sub>1.7 target (HwTx-IV E<sup>4</sup>K/R<sup>26</sup>A/Y<sup>33</sup>W) as it both improves the potency on this channel type and decreases the one on hNa<sub>V</sub>1.6, which should lead to decreased side effects.

## REFERENCES

- Agwa, A. J., Lawrence, N., Deplazes, E., Cheneval, O., Chen, R. M., Craik, D. J., et al. (2017). Spider Peptide Toxin HwTx-IV Engineered to Bind to Lipid Membranes Has an Increased Inhibitory Potency at Human Voltage-Gated Sodium Channel hNa<sub>V</sub> 1.7. *Biochim. Biophys. Acta (Bba) - Biomembranes* 1859, 835–844. doi:10.1016/j.bbame.2017.01.020
- Agwa, A. J., Tran, P., Mueller, A., Tran, H. N. T., Deuis, J. R., Israel, M. R., et al. (2020). Manipulation of a Spider Peptide Toxin Alters its Affinity for Lipid Bilayers and Potency and Selectivity for Voltage-Gated Sodium Channel Subtype 1.7. *J. Biol. Chem.* 295, 5067–5080. doi:10.1074/jbc.ra119.012281
- Bordon, K. d. C. F., Cologna, C. T., Fornari-Baldo, E. C., Pinheiro-Júnior, E. L., Cerni, F. A., Amorim, F. G., et al. (2020). From Animal Poisons and Venoms to Medicines: Achievements, Challenges and Perspectives in Drug Discovery. *Front. Pharmacol.* 11, 1132. doi:10.3389/fphar.2020.01132

## CONCLUSION

Overall, this work demonstrates that it is possible to create a number of new HwTx-IV analogues with increased potencies for both channel subtypes up to the point of equal potencies on hNa<sub>V</sub>1.7 and hNa<sub>V</sub>1.6. However, it was never possible to create an inversion of the selectivity ratio with a loss of potency towards hNa<sub>V</sub>1.7 combined with nanomolar affinity for hNa<sub>V</sub>1.6. For such a result, we suggest that it is better to start SAR investigation on a toxin that initially has better affinity towards Na<sub>V</sub>1.6.

## DATA AVAILABILITY STATEMENT

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation.

## AUTHOR CONTRIBUTIONS

LL, JM, BP, BO-M, KK, SL and SN performed and analysed the experiments; EB and MDW wrote the manuscript; CC, SW, MW and RB designed and supervised the production of HwTx-IV analogues; DS reviewed and edited the paper. RB and MW acquired the funding. All authors read and approved the final manuscript.

## FUNDING

This work was supported by the Agence Nationale de la Recherche (grant number ANR-11-LABX-0015 to MDW), the Fondation Leducq in the frame of its program of “*Équipement de recherche et plateformes technologiques*”, the Région Pays de la Loire (nouvelle équipe, grant number 2016-11092/11,093 to MDW), and the European FEDER (grant number 2017/FEDER/PL0014592 to MDW). LL is supported by a CIFRE fellowship jointly financed by the ANRT and Smartox Biotechnology, while JM is supported by the Agence Nationale de la Recherche (grant number ANR-18-CE19-0024-02 to MDW).

- Cardoso, F. C., and Lewis, R. J. (2018). Sodium Channels and Pain: from Toxins to Therapies. *Br. J. Pharmacol.* 175, 2138–2157. doi:10.1111/bph.13962
- Cardoso, F. C., and Lewis, R. J. (2019). Structure-Function and Therapeutic Potential of Spider Venom-Derived Cysteine Knot Peptides Targeting Sodium Channels. *Front. Pharmacol.* 10, 366. doi:10.3389/fphar.2019.00366
- Cox, J. J., Reimann, F., Nicholas, A. K., Thornton, G., Roberts, E., Springell, K., et al. (2006). An SCN9A Channelopathy Causes Congenital Inability to Experience Pain. *Nature* 444, 894–898. doi:10.1038/nature05413
- Cuesta, S. A., and Meneses, L. (2021). The Role of Organic Small Molecules in Pain Management. *Molecules* 26. doi:10.3390/molecules26134029
- Deuis, J. R., Wingerd, J. S., Winter, Z., Durek, T., Dekan, Z., Sousa, S. R., et al. (2016). Analgesic Effects of GpTx-1, PF-04856264 and CNV1014802 in a Mouse Model of Na<sub>V</sub>1.7-Mediated Pain. *Toxins (Basel)* 8. doi:10.3390/toxins8030078
- Dongol, Y., Cardoso, F. C., and Lewis, R. J. (2019). Spider Knottin Pharmacology at Voltage-Gated Sodium Channels and Their Potential to Modulate Pain Pathways. *Toxins (Basel)* 11. doi:10.3390/toxins11110626

- Estacion, M., Dib-Hajj, S. D., Benke, P. J., Te Morsche, R. H. M., Eastman, E. M., Macala, L. J., et al. (2008). Nav1.7 Gain-Of-Function Mutations as a Continuum: A1632E Displays Physiological Changes Associated with Erythromelalgia and Paroxysmal Extreme Pain Disorder Mutations and Produces Symptoms of Both Disorders. *J. Neurosci.* 28, 11079–11088. doi:10.1523/jneurosci.3443-08.2008
- Flinspach, M., Xu, Q., Piekarz, A. D., Fellows, R., Hagan, R., Gibbs, A., et al. (2017). Insensitivity to Pain Induced by a Potent Selective Closed-State Nav1.7 Inhibitor. *Sci. Rep.* 7, 39662. doi:10.1038/srep39662
- Focken, T., Liu, S., Chahal, N., Dauphinais, M., Grimwood, M. E., Chowdhury, S., et al. (2016). Discovery of Aryl Sulfonamides as Isoform-Selective Inhibitors of Nav1.7 with Efficacy in Rodent Pain Models. *ACS Med. Chem. Lett.* 7, 277–282. doi:10.1021/acsmchemlett.5b00447
- Gao, S., Valinsky, W. C., On, N. C., Houlihan, P. R., Qu, Q., Liu, L., et al. (2020). Employing NaChBac for Cryo-EM Analysis of Toxin Action on Voltage-Gated Na<sup>+</sup>-channels in Nanodisc. *Proc. Natl. Acad. Sci. USA* 117, 14187–14193. doi:10.1073/pnas.1922903117
- Gonçalves, T. C., Benoit, E., Partiseti, M., and Servent, D. (2018a). The Nav1.7 Channel Subtype as an Antinociceptive Target for Spider Toxins in Adult Dorsal Root Ganglia Neurons. *Front. Pharmacol.* 9, 1000. doi:10.3389/fphar.2018.01000
- Gonçalves, T. C., Boukaiba, R., Molgó, J., Amar, M., Partiseti, M., Servent, D., et al. (2018b). Direct Evidence for High Affinity Blockade of Nav1.6 Channel Subtype by Huwentoxin-IV Spider Peptide, Using Multiscale Functional Approaches. *Neuropharmacology* 133, 404–414. doi:10.1016/j.neuropharm.2018.02.016
- Klint, J. K., Chin, Y. K.-Y., and Mobli, M. (2015). Rational Engineering Defines a Molecular Switch that Is Essential for Activity of Spider-Venom Peptides against the Analgesics Target Nav1.7. *Mol. Pharmacol.* 88, 1002–1010. doi:10.1124/mol.115.100784
- Klint, J. K., Senff, S., Rupasinghe, D. B., Er, S. Y., Herzig, V., Nicholson, G. M., et al. (2012). Spider-venom Peptides that Target Voltage-Gated Sodium Channels: Pharmacological Tools and Potential Therapeutic Leads. *Toxicon* 60, 478–491. doi:10.1016/j.toxicon.2012.04.337
- Liu, Y., Wu, Z., Tang, D., Xun, X., Liu, L., Li, X., et al. (2014b). Analgesic Effects of Huwentoxin-IV on Animal Models of Inflammatory and Neuropathic Pain. *Protein Pept. Lett.* 21, 153–158. doi:10.2174/09298665113206660119
- Liu, Y., Tang, J., Zhang, Y., Xun, X., Tang, D., Peng, D., et al. (2014a). Synthesis and Analgesic Effects of  $\mu$ -TRTX-Hhn1b on Models of Inflammatory and Neuropathic Pain. *Toxins* 6, 2363–2378. doi:10.3390/toxins6082363
- Macsari, I., Besidski, Y., Csjernyik, G., Nilsson, L. I., Sandberg, L., Yngve, U., et al. (2012). 3-Oxoisoindoline-1-carboxamides: Potent, State-dependent Blockers of Voltage-Gated Sodium Channel Nav1.7 with Efficacy in Rat Pain Models. *J. Med. Chem.* 55, 6866–6880. doi:10.1021/jm300623u
- Minassian, N. A., Gibbs, A., Shih, A. Y., Liu, Y., Neff, R. A., Sutton, S. W., et al. (2013). Analysis of the Structural and Molecular Basis of Voltage-Sensitive Sodium Channel Inhibition by the Spider Toxin Huwentoxin-IV ( $\mu$ -TRTX-Hh2a). *J. Biol. Chem.* 288, 22707–22720. doi:10.1074/jbc.m113.461392
- Montnach, J., De Waard, S., Nicolas, S., Burel, S., Osorio, N., Zoukian, C., et al. (2021). Fluorescent- and Tagged-prototoxin II Peptides: Potent Markers of the Na V 1.7 Channel Pain Target. *Br. J. Pharmacol.* 178, 2632–2650. doi:10.1111/bph.15453
- Neff, R. A., Flinspach, M., Gibbs, A., Shih, A. Y., Minassian, N. A., Liu, Y., et al. (2020). Comprehensive Engineering of the Tarantula Venom Peptide Huwentoxin-IV to Inhibit the Human Voltage-Gated Sodium Channel hNav1.7. *J. Biol. Chem.* 295, 1315–1327. doi:10.1074/jbc.ra119.011318
- Nguyen, H. N., Bregman, H., Buchanan, J. L., Du, B., Feric, E., Huang, L., et al. (2012). Discovery and Optimization of Aminopyrimidinones as Potent and State-dependent Nav1.7 Antagonists. *Bioorg. Med. Chem. Lett.* 22, 1055–1060. doi:10.1016/j.bmcl.2011.11.111
- Peng, K., Shu, Q., Liu, Z., and Liang, S. (2002). Function and Solution Structure of Huwentoxin-IV, a Potent Neuronal Tetrodotoxin (TTX)-sensitive Sodium Channel Antagonist from Chinese Bird Spider Selenocosmia Huwena. *J. Biol. Chem.* 277, 47564–47571. doi:10.1074/jbc.m204063200
- Rahnama, S., Deuis, J. R., Cardoso, F. C., Ramanujam, V., Lewis, R. J., Rash, L. D., et al. (2017). The Structure, Dynamics and Selectivity Profile of a Nav1.7 Potency-Optimised Huwentoxin-IV Variant. *PLoS One* 12, e0173551. doi:10.1371/journal.pone.0173551
- Revell, J. D., Lund, P.-E., Linley, J. E., Metcalfe, J., Burmeister, N., Sridharan, S., et al. (2013). Potency Optimization of Huwentoxin-IV on hNav1.7: a Neurotoxin TTX-S Sodium-Channel Antagonist from the Venom of the Chinese Bird-Eating Spider Selenocosmia Huwena. *Peptides* 44, 40–46. doi:10.1016/j.peptides.2013.03.011
- Sermadiras, I., Revell, J., Linley, J. E., Sandercock, A., and Ravn, P. (2013). Recombinant Expression and *In Vitro* Characterisation of Active Huwentoxin-IV. *PLoS One* 8, e83202. doi:10.1371/journal.pone.0083202
- Shen, H., Liu, D., Wu, K., Lei, J., and Yan, N. (2019). Structures of Human Na V 1.7 Channel in Complex with Auxiliary Subunits and Animal Toxins. *Science* 363, 1303–1308. doi:10.1126/science.aaw2493
- Sun, S., Jia, Q., Zenova, A. Y., Chafeev, M., Zhang, Z., Lin, S., et al. (2014). The Discovery of Benzenesulfonamide-Based Potent and Selective Inhibitors of Voltage-Gated Sodium Channel Na V 1.7. *Bioorg. Med. Chem. Lett.* 24, 4397–4401. doi:10.1016/j.bmcl.2014.08.017
- Trim, C. M., Byrne, L. J., and Trim, S. A. (2021). Utilisation of Compounds from Venoms in Drug Discovery. *Prog. Med. Chem.* 60, 1–66. doi:10.1016/bs.pmc.2021.01.001
- Tzakoniati, F., Xu, H., Li, T., Garcia, N., Kugel, C., Payandeh, J., et al. (2020). Development of Photocrosslinking Probes Based on Huwentoxin-IV to Map the Site of Interaction on Nav1.7. *Cel Chem. Biol.* 27, 306–313. doi:10.1016/j.chembiol.2019.10.011
- Wu, Y.-J., Guernon, J., McClure, A., Luo, G., Rajamani, R., Ng, A., et al. (2017). Discovery of Non-zwitterionic Aryl Sulfonamides as Nav1.7 Inhibitors with Efficacy in Preclinical Behavioral Models and Translational Measures of Nociceptive Neuron Activation. *Bioorg. Med. Chem.* 25, 5490–5505. doi:10.1016/j.bmc.2017.08.012
- Xiao, Y., Bingham, J.-P., Zhu, W., Moczydlowski, E., Liang, S., and Cummins, T. R. (2008). Tarantula Huwentoxin-IV Inhibits Neuronal Sodium Channels by Binding to Receptor Site 4 and Trapping the Domain II Voltage Sensor in the Closed Configuration. *J. Biol. Chem.* 283, 27300–27313. doi:10.1074/jbc.m708447200
- Xiao, Y., Blumenthal, K., and Cummins, T. R. (2014). Gating-pore Currents Demonstrate Selective and Specific Modulation of Individual Sodium Channel Voltage-Sensors by Biological Toxins. *Mol. Pharmacol.* 86, 159–167. doi:10.1124/mol.114.092338
- Xiao, Y., Blumenthal, K., Jackson, J. O., 2nd, Liang, S., and Cummins, T. R. (2010). The Tarantula Toxins ProTx-II and Huwentoxin-IV Differentially Interact with Human Nav1.7 Voltage Sensors to Inhibit Channel Activation and Inactivation. *Mol. Pharmacol.* 78, 1124–1134. doi:10.1124/mol.110.066332
- Xiao, Y., Jackson, J. O., 2nd, Liang, S., and Cummins, T. R. (2011). Common Molecular Determinants of Tarantula Huwentoxin-IV Inhibition of Na<sup>+</sup> Channel Voltage Sensors in Domains II and IV. *J. Biol. Chem.* 286, 27301–27310. doi:10.1074/jbc.m111.246876

**Conflict of Interest:** Authors KK, SL, SW, and CC were employed by company Xenon Pharmaceuticals. Authors BT, CC, RB, and LL were employed by company Smartox Biotechnology. MDW is a consultant and founder of Smartox Biotechnology.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors, and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Lopez, Montnach, Oliveira-Mendes, Khakh, Thomas, Lin, Caumes, Wesolowski, Nicolas, Servent, Cohen, Bérout, Benoit and De Waard. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.